

Unit 108 - Field Data Review

Mohawk Valley West Slope demonstration. Method **Tractor**, 63.3 ac, BAF 20 variable-radius cruise on a 300 ft grid, 31 plots, 344 trees tallied. Simulated data with planted recording errors; see docs/methods.md.

Table 1. Stand summary from the BAF 20 cruise, per acre

STAND DENSITY		STAND INFO	
Basal area	222 sq ft/ac (1 SE of mean, BA 2.6 %)	Type	Sierran mixed conifer
Trees per acre	68	CWHR size / density	5 / D (cover 91 %)
QMD	24.5 in	SDI	274 (47 % of max 580)
SAWTIMBER (>= 10 in)		BIOMASS (< 10 in)	
BA QMD	222 sq ft/ac 24.5 in	BA QMD	0 sq ft/ac 0.0 in
TPA	68	TPA	0
Volume	39.2 MBF/ac (7810 cu ft)	Volume	0.0 green tons/ac (0 cu ft)
TREATMENT TARGET		CRUISE DESIGN	
Leave	35 % of max SDI = ~164 sq ft/ac BA	Sampling error (volume)	6.4 % at 95 % (CV 18 %, t = 2.04, n = 31)
Species (BA)	WF 68, PP 66, IC 36, DF 29, SP 22	Stratum standard	40 % of volume - MEETS; plots for standard: 1
		Sampling error (basal area)	5.4 % at 95 % (CV 15 %); secondary, not the standard
		Plot count	31 vs 20-plot local practice, not a handbook standard: ok

Cubic volume = total-stem CVTS by species (PNW-FIA California equations, MacLean and Berger 1976, CARB 2011 compendium) net of defect; Scribner board feet at 5.02 bf/cu ft (Keegan et al. 2010, table 2, California 2000-2006); green tons at PP 45, WF 47, DF 38, SP 49, IC 45 lb/cu ft of wood (Miles and Smith 2009, NRS-38 table 1A). SDI in the summation form; maximum is the basal-area-weighted FVS Western Sierra species maximum (table 3.5.1, rev. 2025-09-23). The Keegan ratio is Scribner log scale per cubic foot of delivered sawlog fiber, so applied to total-stem CVTS it overstates sawlog volume; merchantable cubic volume would be the consistent pairing. Sampling error at 95 % confidence with Student's t on plot net cubic volume per acre, the quantity the handbook standards apply to (FSH 2409.12 ch. 40, 41.1(5)(a) and (b)); stratum standard 40 % of volume for tree-measurement sales; the basal-area error is secondary. Stand-summary BA counts every tallied tree (plot count x BAF); the stand and stock tables exclude records flagged by QA, such as an out-of-range DBH, so the two can differ by a few tenths. Sources and URLs: cruise_data.xlsx, Standards and assumptions.

Table 2. Stand table by 4-inch DBH class, per acre

DBH class, in	Trees	TPA	BA	cu ft	MBF
8-11.9	4	3.6	2.6	66	0.3
12-15.9	22	13.2	14.2	345	1.7
16-19.9	33	12.0	21.3	610	3.1
20-23.9	53	12.7	34.2	1063	5.3
24-27.9	60	10.7	38.7	1307	6.6
28-31.9	44	5.9	28.4	1021	5.1
32-35.9	39	4.0	25.2	919	4.6
36-39.9	35	2.9	22.6	896	4.5
40-43.9	26	1.7	16.8	701	3.5
44-47.9	6	0.3	3.9	167	0.8
48-51.9	7	0.3	4.5	212	1.1
52-55.9	6	0.2	3.9	209	1.1
56-59.9	5	0.2	3.2	153	0.8
60-63.9	4	0.1	2.6	140	0.7
All	344	68.1	221.9	7810	39.2

Table 3. Stock table by species, per acre

Species	Trees	TPA	BA	QMD	cu ft	MBF
WF	106	22.5	68.4	23.6	2344	11.8
PP	103	19.0	66.5	25.3	2641	13.3
IC	56	12.9	36.1	22.7	993	5.0
DF	45	7.0	29.0	27.6	1002	5.0
SP	34	6.7	21.9	24.4	830	4.2
All	344	68.1	221.9	24.5	7810	39.2

Trees = trees tallied in the cruise; TPA, BA (sq ft/ac), cubic feet and Scribner MBF per acre by BAF expansion. Species: PP ponderosa pine, WF white fir, DF Douglas-fir, SP sugar pine, IC incense-cedar.

Unit 108: plots on the 300 ft grid (orange = QA flag)




Figure 1. Plot locations with QA status.

QA findings: no issues found; unit accepted

Table 4. Plot list

Plot	Trees in	BA sq ft/ac	Slope %	Status
108-01	10	200	11	ok
108-02	11	220	22	ok
108-03	9	180	23	ok
108-04	10	200	18	ok
108-05	10	200	16	ok
108-06	12	240	21	ok
108-07	11	220	27	ok
108-08	11	220	23	ok
108-09	11	220	27	ok
108-10	12	240	12	ok
108-11	11	220	14	ok
108-12	14	280	25	ok
108-13	6	120	17	ok
108-14	13	260	6	ok
108-15	14	280	15	ok
108-16	10	200	13	ok
108-17	11	220	9	ok
108-18	11	220	8	ok
108-19	9	180	32	ok
108-20	12	240	17	ok
108-21	10	200	22	ok
108-22	12	240	30	ok
108-23	11	220	18	ok
108-24	12	240	11	ok
108-25	10	200	16	ok
108-26	13	260	15	ok
108-27	13	260	19	ok
108-28	11	220	10	ok
108-29	11	220	13	ok
108-30	10	200	20	ok
108-31	13	260	10	ok