

Unit 404 - Field Data Review

Mohawk Valley West Slope demonstration. Method **Cable**, 51.2 ac, BAF 20 variable-radius cruise on a 300 ft grid, 27 plots, 281 trees tallied. Simulated data with planted recording errors; see docs/methods.md.

Table 1. Stand summary from the BAF 20 cruise, per acre

STAND DENSITY		STAND INFO	
Basal area	208 sq ft/ac (1 SE of mean, BA 3.7 %)	Type	Sierran mixed conifer
Trees per acre	61	CWHR size / density	5 / D (cover 87 %)
QMD	25.0 in	SDI	252 (43 % of max 580)
SAWTIMBER (>= 10 in)		BIOMASS (< 10 in)	
BA QMD	207 sq ft/ac 25.7 in	BA QMD	1 sq ft/ac 5.9 in
TPA	57	TPA	4
Volume	39.2 MBF/ac (7803 cu ft)	Volume	0.2 green tons/ac (10 cu ft)
TREATMENT TARGET		CRUISE DESIGN	
Leave	35 % of max SDI = ~168 sq ft/ac BA	Sampling error (volume)	12.0 % at 95 % (CV 30 %, t = 2.06, n = 27)
Species (BA)	WF 67, PP 64, DF 30, IC 27, SP 20	Stratum standard	40 % of volume - MEETS; plots for standard: 3
		Sampling error (basal area)	2.6 % at 95 % (CV 19 %); secondary, not the standard
		Plot count	27 vs 20-plot local practice, not a handbook standard: ok

Cubic volume = total-stem CVTS by species (PNW-FIA California equations, MacLean and Berger 1976, CARB 2011 compendium) net of defect; Scribner board feet at 5.02 bf/cu ft (Keegan et al. 2010, table 2, California 2000-2006); green tons at PP 45, WF 47, DF 38, SP 49, IC 45 lb/cu ft of wood (Miles and Smith 2009, NRS-38 table 1A). SDI in the summation form; maximum is the basal-area-weighted FVS Western Sierra species maximum (table 3.5.1, rev. 2025-09-23). The Keegan ratio is Scribner log scale per cubic foot of delivered sawlog fiber, so applied to total-stem CVTS it overstates sawlog volume; merchantable cubic volume would be the consistent pairing. Sampling error at 95 % confidence with Student's t on plot net cubic volume per acre, the quantity the handbook standards apply to (FSH 2409.12 ch. 40, 41.1(5)(a) and (b)); stratum standard 40 % of volume for tree-measurement sales; the basal-area error is secondary. Stand-summary BA counts every tallied tree (plot count x BAF); the stand and stock tables exclude records flagged by QA, such as an out-of-range DBH, so the two can differ by a few tenths. Sources and URLs: cruise_data.xlsx, Standards and assumptions.

Table 2. Stand table by 4-inch DBH class, per acre

DBH class, in	Trees	TPA	BA	cu ft	MBF
4-7.9	1	3.9	0.7	10	0.1
8-11.9	4	4.4	3.0	50	0.3
12-15.9	10	6.5	7.4	181	0.9
16-19.9	27	11.5	20.0	555	2.8
20-23.9	36	10.2	26.7	802	4.0
24-27.9	43	8.9	31.9	1062	5.3
28-31.9	38	5.8	28.1	1080	5.4
32-35.9	37	4.3	27.4	1059	5.3
36-39.9	22	2.1	16.3	699	3.5
40-43.9	13	1.0	9.6	411	2.1
44-47.9	17	1.1	12.6	599	3.0
48-51.9	17	0.9	12.6	654	3.3
52-55.9	3	0.1	2.2	115	0.6
56-59.9	3	0.1	2.2	110	0.6
60-63.9	10	0.4	7.4	425	2.1
All	281	61.3	208.1	7814	39.2

Table 3. Stock table by species, per acre

Species	Trees	TPA	BA	QMD	cu ft	MBF
WF	90	23.7	66.7	22.7	2425	12.2
PP	87	16.6	64.4	26.6	2704	13.6
DF	41	7.5	30.4	27.2	1099	5.5
IC	36	6.7	26.7	27.0	792	4.0
SP	27	6.7	20.0	23.4	793	4.0
All	281	61.3	208.1	25.0	7814	39.2

Trees = trees tallied in the cruise; TPA, BA (sq ft/ac), cubic feet and Scribner MBF per acre by BAF expansion. Species: PP ponderosa pine, WF white fir, DF Douglas-fir, SP sugar pine, IC incense-cedar.

Unit 404: plots on the 300 ft grid (orange = QA flag)




Figure 1. Plot locations with QA status.

QA findings: no issues found; unit accepted

Table 4. Plot list

Plot	Trees in	BA sq ft/ac	Slope %	Status
404-01	11	220	65	ok
404-02	11	220	50	ok
404-03	13	260	39	ok
404-04	10	200	67	ok
404-05	11	220	57	ok
404-06	11	220	56	ok
404-07	12	240	54	ok
404-08	9	180	48	ok
404-09	13	260	60	ok
404-10	6	120	41	ok
404-11	12	240	52	ok
404-12	11	220	60	ok
404-13	12	240	44	ok
404-14	9	180	48	ok
404-15	8	160	59	ok
404-16	10	200	58	ok
404-17	11	220	36	ok
404-18	10	200	47	ok
404-19	4	80	46	ok
404-20	10	200	71	ok
404-21	11	220	57	ok
404-22	10	200	43	ok
404-23	10	200	40	ok
404-24	10	200	57	ok
404-25	11	220	50	ok
404-26	12	240	50	ok
404-27	13	260	36	ok