

Unit 401 Field Data Review

Mohawk Valley West Slope demonstration. **Cable** unit, 29.5 ac. 14 plots, BAF 20, variable-radius cruise on a 300 ft grid, 155 trees tallied. Simulated data with planted recording errors; see docs/methods.md.

TABLE 1. STAND SUMMARY, BAF 20 CRUISE, PER ACRE

STAND DENSITY		STAND INFO	
Basal area	221 sq ft/ac (1 SE of mean, BA 2.9 %)	Type	Sierran mixed conifer
Trees per acre	101	CWHR size / density	4 / D (cover 94 %)
QMD	20.0 in	SDI	290 (49 % of max 591)
SAWTIMBER (>= 10 in)		BIOMASS (< 10 in)	
BA QMD	217 sq ft/ac 20.8 in	BA QMD	4 sq ft/ac 9.5 in
TPA	92	TPA	9
Volume	34.8 MBF/ac (6933 cu ft)	Volume	1.5 green tons/ac (68 cu ft)
TREATMENT TARGET		CRUISE DESIGN	
Leave	35 % of max SDI, about 158 sq ft/ac	BA Sampling error (volume)	22.9 % at 95 % (CV 40 %, t = 2.16, n = 14)
Species (BA)	WF 79, PP 66, DF 33, IC 24, SP 20	Stratum standard	40 % of volume, MEETS; plots for standard 5
		Sampling error (basal area)	33 % at 95 % (CV 11 %); reference, not the standard
		Plot count	14; 20-plot local practice, not a handbook standard: SHORT by 6

Cubic volume is total-stem CVTS by species (PNW-FIA California equations, MacLean and Berger 1976, CARB 2011 compendium), net of defect. Scribner board feet at 5.02 bf/cu ft (Keegan et al. 2010, table 2, California 2000-2006). Green tons at PP 45, WF 47, DF 38, SP 49, IC 45 lb/cu ft of wood (Miles and Smith 2009, NRS-38 table 1A). SDI in the summation form; maximum is the basal-area-weighted FVS Western Sierra species maximum (table 3.5.1, rev. 2025-09-23). The Keegan ratio is Scribner log scale per cubic foot of delivered sawlog fiber; applied to total-stem CVTS it overstates sawlog volume. Merchantable cubic volume is the consistent pairing. Sampling error at 95 % confidence, Student's t, on plot net cubic volume per acre, the quantity the handbook standards test (FSH 2409.12 ch. 40, 41.1(5)(a) and (b)); stratum standard 40 % of volume for tree-measurement sales. The basal-area error is reference only. Stand-summary BA counts every tallied tree (plot count x BAF); the stand and stock tables drop records flagged by QA, such as an out-of-range DBH, so the two can differ by a few tenths. Sources and URLs: cruise_data.xlsx, Standards and assumptions.

TABLE 2. STAND TABLE BY 4-INCH DBH CLASS, PER ACRE

DBH class, in	Trees	TPA	BA	cu ft	MBF
8-11.9	9	21.0	12.9	194	1.0
12-15.9	27	35.9	38.6	755	3.8
16-19.9	19	15.2	27.1	654	3.3
20-23.9	15	8.4	21.4	552	2.8
24-27.9	19	7.6	27.1	927	4.7
28-31.9	14	4.1	20.0	669	3.4
32-35.9	23	5.2	32.9	1269	6.4
36-39.9	8	1.5	11.4	512	2.6
40-43.9	1	0.1	1.4	50	0.3
44-47.9	5	0.6	7.1	320	1.6
48-51.9	10	1.1	14.3	725	3.6
52-55.9	2	0.2	2.9	178	0.9
56-59.9	1	0.1	1.4	61	0.3
60-63.9	2	0.1	2.9	137	0.7
All	155	101.1	221.4	7001	35.1

TABLE 3. STOCK TABLE BY SPECIES, PER ACRE

Species	Trees	TPA	BA	QMD	cu ft	MBF
WF	55	31.7	78.6	21.3	2481	12.5
PP	46	30.6	65.7	19.8	2347	11.8
DF	23	13.8	32.9	20.9	1043	5.2
IC	17	12.9	24.3	18.6	542	2.7
SP	14	12.0	20.0	17.5	588	3.0
All	155	101.1	221.4	20.0	7001	35.1

Trees: tally trees. TPA, BA (sq ft/ac), cubic feet and Scribner MBF per acre by BAF expansion. Species codes: PP ponderosa pine, WF white fir, DF Douglas-fir, SP sugar pine, IC incense-cedar.




Figure 1. Plot locations with QA status.

QA findings: none; unit accepted; 6 more plots to meet the 20-plot local practice

TABLE 4. PLOT LIST

Plot	Trees in	BA sq ft/ac	Slope %	Status
401-01	12	240	51	ok
401-02	10	200	46	ok
401-03	11	220	47	ok
401-04	13	260	37	ok
401-05	12	240	40	ok
401-06	10	200	38	ok
401-07	11	220	43	ok
401-08	13	260	36	ok
401-09	10	200	38	ok
401-10	11	220	38	ok
401-11	11	220	36	ok
401-12	9	180	49	ok
401-13	10	200	38	ok
401-14	12	240	46	ok