

Question ID 3cdbf026

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	<div><div></div><div></div><div></div></div>

ID: 3cdbf026

The graph of the equation $ax + ky = 6$ is a line in the xy -plane, where a and k are constants. If the line contains the points $(-2, -6)$ and $(0, -3)$, what is the value of k ?

- A. -2
- B. -1
- C. 2
- D. 3

ID: 3cdbf026 Answer

Correct Answer: A

Rationale

Choice A is correct. The value of k can be found using the slope-intercept form of a linear equation, $y = mx + b$, where m is the slope and b is the y -coordinate of the y -intercept. The equation $ax + ky = 6$ can be rewritten in the form $y = -\frac{ax}{k} + \frac{6}{k}$. One of the given points, $(0, -3)$, is the y -intercept. Thus, the y -coordinate of the y -intercept -3 must be equal to $\frac{6}{k}$. Multiplying both sides by k gives $-3k = 6$. Dividing both sides by -3 gives $k = -2$.

Choices B, C, and D are incorrect and may result from errors made rewriting the given equation.

Question Difficulty: Hard

Question ID 9bbce683

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	<div><div></div><div></div><div></div></div>

ID: 9bbce683

x	y
18	130
23	160
26	178

For line h , the table shows three values of x and their corresponding values of y . Line k is the result of translating line h down 5 units in the xy -plane. What is the x -intercept of line k ?

- A. $(-\frac{26}{3}, 0)$
- B. $(-\frac{9}{2}, 0)$
- C. $(-\frac{11}{3}, 0)$
- D. $(-\frac{17}{6}, 0)$

ID: 9bbce683 Answer

Correct Answer: D

Rationale

Choice D is correct. The equation of line h can be written in slope-intercept form $y = mx + b$, where m is the slope of the line and b is the y -intercept of the line. It's given that line h contains the points 18, 130, 23, 160, and 26, 178. Therefore, its slope m can be found as $\frac{160 - 130}{23 - 18}$, or 6. Substituting 6 for m in the equation $y = mx + b$ yields $y = 6x + b$. Substituting 130 for y and 18 for x in this equation yields $130 = 6(18) + b$, or $130 = 108 + b$. Subtracting 108 from both sides of this equation yields $22 = b$. Substituting 22 for b in $y = 6x + b$ yields $y = 6x + 22$. Since line k is the result of translating line h down 5 units, an equation of line k is $y = 6x + 22 - 5$, or $y = 6x + 17$. Substituting 0 for y in this equation yields $0 = 6x + 17$. Solving this equation for x yields $x = -\frac{17}{6}$. Therefore, the x -intercept of line k is $-\frac{17}{6}, 0$.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice B is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Hard

Question ID fdee0fbf

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	<div><div></div><div></div><div></div></div>

ID: fdee0fbf

In the xy -plane, line k intersects the y -axis at the point $(0, -6)$ and passes through the point $(2, 2)$. If the point $(20, w)$ lies on line k , what is the value of w ?

ID: fdee0fbf Answer

Rationale

The correct answer is 74. The y -intercept of a line in the xy -plane is the ordered pair (x,y) of the point of intersection of the line with the y -axis. Since line k intersects the y -axis at the point $(0, -6)$, it follows that $(0, -6)$ is the y -intercept of this line. An equation of any line in the xy -plane can be written in the form $y = mx + b$, where m is the slope of the line and b is the y -coordinate of the y -intercept. Therefore, the equation of line k can be written as $y = mx + (-6)$, or $y = mx - 6$. The value of m can be found by substituting the x - and y -coordinates from a point on the line, such as $(2,2)$, for x and y , respectively. This results in $2 = 2m - 6$. Solving this equation for m gives $m = 4$. Therefore, an equation of line k is $y = 4x - 6$. The value of w can be found by substituting the x -coordinate, 20, for x in the equation of line k and solving this equation for y . This gives $y = 4(20) - 6$, or $y = 74$. Since w is the y -coordinate of this point, $w = 74$.

Question Difficulty: Hard

Question ID 0b46bad5


Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	■ ■ ■

ID: 0b46bad5


$ax + by = b$

In the equation above, a and b are constants and $0 < a < b$. Which of the following could represent the graph of the equation in the xy -plane?


A.




B.



C.



D.



ID: 0b46bad5 Answer

Correct Answer: C

Rationale

Choice C is correct. The given equation $ax + by = b$ can be rewritten in slope-intercept form, $y = mx + k$, where m represents the slope of the line represented by the equation, and k represents the y-coordinate of the y-intercept of the line. Subtracting ax from both sides of the equation yields $by = -ax + b$, and dividing both sides of this equation by b yields $y = -\frac{a}{b}x + \frac{b}{b}$, or $y = -\frac{a}{b}x + 1$. With the equation now in slope-intercept form, it shows that $k = 1$, which means the y-coordinate of the y-intercept is 1. It's given that a and b are both greater than 0 (positive) and that $a < b$. Since $m = -\frac{a}{b}$, the slope of the line must be a value between -1 and 0. Choice C is the only graph of a line that has a y-value of the y-intercept that is 1 and a slope that is between -1 and 0.


Choices A, B, and D are incorrect because the slopes of the lines in these graphs aren't between -1 and 0.

Question Difficulty: Hard

Question ID c362c210

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	■ ■ ■

ID: c362c210



The points plotted in the coordinate plane above represent the possible numbers of wallflowers and cornflowers that someone can buy at the Garden Store in order to spend exactly \$24.00 total on the two types of flowers. The price of each wallflower is the same and the price of each cornflower is the same. What is the price, in dollars, of 1 cornflower?

ID: c362c210 Answer

Rationale

The correct answer is 1.5. The point (16,0) corresponds to the situation where 16 cornflowers and 0 wallflowers are purchased. Since the total spent on the two types of flowers is \$24.00, it follows that the price of 16 cornflowers is \$24.00, and the price of one cornflower is \$1.50. Note that 1.5 and 3/2 are examples of ways to enter a correct answer.

Question Difficulty: Hard

Question ID 98d3393a

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	<div><div></div><div></div><div></div></div>

ID: 98d3393a

Line ℓ in the xy -plane is perpendicular to the line with equation $x = 2$. What is the slope of line ℓ ?

- A. 0
- B. $-\frac{1}{2}$
- C. -2
- D. The slope of line ℓ is undefined.

ID: 98d3393a Answer

Correct Answer: A

Rationale

Choice A is correct. It is given that line ℓ is perpendicular to a line whose equation is $x = 2$. A line whose equation is a constant value of x is vertical, so ℓ must therefore be horizontal. Horizontal lines have a slope of 0, so ℓ has a slope of 0.

Choice B is incorrect. A line with slope $-\frac{1}{2}$ is perpendicular to a line with slope 2. However, the line with equation $x = 2$ is vertical and has undefined slope (not slope of 2). Choice C is incorrect. A line with slope -2 is perpendicular to a line with slope $\frac{1}{2}$. However, the line with equation $x = 2$ has undefined slope (not slope of $\frac{1}{2}$). Choice D is incorrect; this is the slope of the line $x = 2$ itself, not the slope of a line perpendicular to it.

Question Difficulty: Hard

Question ID cc3e9528

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	<div><div></div><div></div><div></div></div>

ID: cc3e9528

The graph of $9x - 10y = 19$ is translated down 4 units in the xy -plane. What is the x -coordinate of the x -intercept of the resulting graph?

ID: cc3e9528 Answer

Correct Answer: 59/9, 6.555, 6.556

Rationale

The correct answer is $\frac{59}{9}$. When the graph of an equation in the form $Ax + By = C$, where A , B , and C are constants, is translated down k units in the xy -plane, the resulting graph can be represented by the equation $Ax + By + k = C$. It's given that the graph of $9x - 10y = 19$ is translated down 4 units in the xy -plane. Therefore, the resulting graph can be represented by the equation $9x - 10y + 4 = 19$, or $9x - 10y - 40 = 19$. Adding 40 to both sides of this equation yields $9x - 10y = 59$. The x -coordinate of the x -intercept of the graph of an equation in the xy -plane is the value of x in the equation when $y = 0$. Substituting 0 for y in the equation $9x - 10y = 59$ yields $9x - 100 = 59$, or $9x = 59$. Dividing both sides of this equation by 9 yields $x = \frac{59}{9}$. Therefore, the x -coordinate of the x -intercept of the resulting graph is $\frac{59}{9}$. Note that 59/9, 6.555, and 6.556 are examples of ways to enter a correct answer.

Question Difficulty: Hard

Question ID 0366d965

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	<div><div></div><div></div><div></div></div>

ID: 0366d965

x	y
3	7
k	11
12	n

The table above shows the coordinates of three points on a line in the xy -plane, where k and n are constants. If the slope of the line is 2, what is the value of $k + n$?

ID: 0366d965 Answer

Rationale

The correct answer is 30. The slope of a line can be found by using the slope formula, $\frac{y_2 - y_1}{x_2 - x_1}$. It's given that the slope of the line is 2; therefore, $\frac{y_2 - y_1}{x_2 - x_1} = 2$. According to the table, the points $(3, 7)$ and $(k, 11)$ lie on the line. Substituting the coordinates of these points into the equation gives $\frac{11 - 7}{k - 3} = 2$. Multiplying both sides of this equation by $k - 3$ gives $11 - 7 = 2(k - 3)$, or $11 - 7 = 2k - 6$. Solving for k gives $k = 5$. According to the table, the points $(3, 7)$ and $(12, n)$ also lie on the line. Substituting the coordinates of these points into $\frac{y_2 - y_1}{x_2 - x_1} = 2$ gives $\frac{n - 7}{12 - 3} = 2$. Solving for n gives $n = 25$. Therefore, $k + n = 5 + 25$, or 30.

Question Difficulty: Hard

Question ID a35c7164

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	<div><div></div><div></div><div></div></div>

ID: a35c7164

$$5x + 7y = 1$$
$$ax + by = 1$$

In the given pair of equations, a and b are constants. The graph of this pair of equations in the xy -plane is a pair of perpendicular lines. Which of the following pairs of equations also represents a pair of perpendicular lines?

- A. $10x + 7y = 1$
 $ax - 2by = 1$
- B. $10x + 7y = 1$
 $ax + 2by = 1$
- C. $10x + 7y = 1$
 $2ax + by = 1$
- D. $5x - 7y = 1$
 $ax + by = 1$

ID: a35c7164 Answer

Correct Answer: B

Rationale

Choice B is correct. Two lines are perpendicular if their slopes are negative reciprocals, meaning that the slope of the first line is equal to -1 divided by the slope of the second line. Each equation in the given pair of equations can be written in slope-intercept form, $y = mx + b$, where m is the slope of the graph of the equation in the xy -plane and $0, b$ is the y -intercept. For the first equation, $5x + 7y = 1$, subtracting $5x$ from both sides gives $7y = -5x + 1$, and dividing both sides of this equation by 7 gives $y = -\frac{5}{7}x + \frac{1}{7}$. Therefore, the slope of the graph of this equation is $-\frac{5}{7}$. For the second equation, $ax + by = 1$, subtracting ax from both sides gives $by = -ax + 1$, and dividing both sides of this equation by b gives $y = -\frac{a}{b}x + \frac{1}{b}$. Therefore, the slope of the graph of this equation is $-\frac{a}{b}$. Since the graph of the given pair of equations is a pair of perpendicular lines, the slope of the graph of the second equation, $-\frac{a}{b}$, must be the negative reciprocal of the slope of the graph of the first equation, $-\frac{5}{7}$. The negative reciprocal of $-\frac{5}{7}$ is $\frac{7}{5}$, or $\frac{a}{b} = \frac{7}{5}$. Similarly, rewriting the equations in choice B in slope-intercept form yields $y = -\frac{10}{7}x + \frac{1}{7}$ and $y = -\frac{a}{2b}x + \frac{1}{2b}$. It follows that the slope of the graph of the first equation in choice B is $-\frac{10}{7}$ and the slope of the graph of the second equation in choice B is $-\frac{a}{2b}$. Since $\frac{a}{b} = \frac{7}{5}$, $-\frac{a}{2b}$ is equal to $-\frac{1}{2} \cdot \frac{7}{5}$, or $\frac{7}{10}$. Since $\frac{7}{10}$ is the negative reciprocal of $-\frac{10}{7}$, the pair of equations in choice B represents a pair of perpendicular lines.

Choice A is incorrect and may result from conceptual or calculation errors.

Choice C is incorrect and may result from conceptual or calculation errors.


Choice D is incorrect and may result from conceptual or calculation errors.

Question Difficulty: Hard

Question ID c4ea43ef

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	■ ■ ■

ID: c4ea43ef



To earn money for college, Avery works two part-time jobs: A and B. She earns \$10 per hour working at job A and \$20 per hour working at job B. In one week, Avery earned a total of s dollars for working at the two part-time jobs. The graph above represents all possible combinations of numbers of hours Avery could have worked at the two jobs to earn s dollars. What is the value of s ?

- A. 128
- B. 160
- C. 200
- D. 320

ID: c4ea43ef Answer

Correct Answer: B

Rationale

Choice B is correct. Avery earns \$10 per hour working at job A. Therefore, if she works a hours at job A, she will earn $10a$ dollars. Avery earns \$20 per hour working at job B. Therefore, if she works b hours at job B, she will earn $20b$ dollars. The graph shown represents all possible combinations of the number of hours Avery could have worked at the two jobs to earn s dollars. Therefore, if she worked a hours at job A, worked b hours at job B, and earned s dollars from both jobs, the following equation represents the graph: $10a + 20b = s$, where s is a constant. Identifying any point (a,b) from the graph and substituting the values of the coordinates for a and b , respectively, in this equation yield the value of s . For example, the point $(16,0)$, where $a = 16$ and $b = 0$, lies on the graph. Substituting 16 for a and 0 for b in the equation $10a + 20b = s$ yields $10(16) + 20(0) = s$, or $160 = s$. Similarly, the point $(0,8)$, where $a = 0$ and $b = 8$, lies on the graph. Substituting 0 for a and 8 for b in the equation $10a + 20b = s$ yields $10(0) + 20(8) = s$, or $160 = s$.

Choices A, C, and D are incorrect. If the value of s is 128, 200, or 320, then no points (a,b) on the graph will satisfy this equation. For example, if the value of s is 128 (choice A), then the equation $10a + 20b = s$ becomes $10a + 20b = 128$. The point $(16,0)$, where $a = 16$ and $b = 0$, lies on the graph. However, substituting 16 for a and 0 for b in $10a + 20b = s$

yields $10(16) + 20(0) = 128$, or $160 = 128$, which is false. Therefore, $(16, 0)$ doesn't satisfy the equation, and so the value of s can't be 128. Similarly, if $s = 200$ (choice C) or $s = 320$ (choice D), then substituting 16 for a and 0 for b yields $160 = 200$ and $160 = 320$, respectively, which are both false.

Question Difficulty: Hard

Question ID cb58833c

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	<div><div></div><div></div><div></div></div>

ID: cb58833c

The line with the equation $\frac{4}{5}x + \frac{1}{3}y = 1$ is graphed in the xy -plane. What is the x -coordinate of the x -intercept of the line?

ID: cb58833c Answer

Rationale

The correct answer is 1.25. The y -coordinate of the x -intercept is 0, so 0 can be substituted for y , giving $\frac{4}{5}x + \frac{1}{3}(0) = 1$.

This simplifies to $\frac{4}{5}x = 1$. Multiplying both sides of $\frac{4}{5}x = 1$ by 5 gives $4x = 5$. Dividing both sides of $4x = 5$ by 4 gives $x = \frac{5}{4}$, which is equivalent to 1.25. Note that 1.25 and 5/4 are examples of ways to enter a correct answer.

Question Difficulty: Hard

Question ID a7a14e87

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	<div><div></div><div></div><div></div></div>

ID: a7a14e87

In the xy -plane, line k is defined by $x + y = 0$. Line j is perpendicular to line k , and the y -intercept of line j is $(0,3)$. Which of the following is an equation of line j ?

- A. $x + y = 3$
- B. $x + y = -3$
- C. $x - y = 3$
- D. $x - y = -3$

ID: a7a14e87 Answer

Correct Answer: D

Rationale

Choice D is correct. It's given that line j is perpendicular to line k and that line k is defined by the equation $x + y = 0$. This equation can be rewritten in slope-intercept form, $y = mx + b$, where m represents the slope of the line and b represents the y -coordinate of the y -intercept of the line, by subtracting x from both sides of the equation, which yields $y = -x$. Thus, the slope of line k is -1 . Since line j and line k are perpendicular, their slopes are opposite reciprocals of each other. Thus, the slope of line j is 1 . It's given that the y -intercept of line j is $(0,3)$. Therefore, the equation for line j in slope-intercept form is $y = x + 3$, which can be rewritten as $x - y = -3$.

Choices A, B, and C are incorrect and may result from conceptual or calculation errors.

Question Difficulty: Hard

Question ID a1fd2304

Assessment	Test	Domain	Skill	Difficulty
SAT	Math	Algebra	Linear equations in two variables	<div><div></div><div></div><div></div></div>

ID: a1fd2304

How many liters of a 25% saline solution must be added to 3 liters of a 10% saline solution to obtain a 15% saline solution?

ID: a1fd2304 Answer

Rationale

The correct answer is 1.5. The total amount, in liters, of a saline solution can be expressed as the liters of each type of saline solution multiplied by the percent concentration of the saline solution. This gives $3(0.10)$, $x(0.25)$, and $(x + 3)(0.15)$, where x is the amount, in liters, of 25% saline solution and 10%, 15%, and 25% are represented as 0.10, 0.15, and 0.25, respectively. Thus, the equation $3(0.10) + 0.25x = 0.15(x + 3)$ must be true. Multiplying 3 by 0.10 and distributing 0.15 to $(x + 3)$ yields $0.30 + 0.25x = 0.15x + 0.45$. Subtracting $0.15x$ and 0.30 from each side of the equation gives $0.10x = 0.15$. Dividing each side of the equation by 0.10 yields $x = 1.5$. Note that 1.5 and $3/2$ are examples of ways to enter a correct answer.

Question Difficulty: Hard