



TPD<sup>2</sup> Approach

Dual-precision Targeted Protein  
Degradation

2026 Q3


## Forward-looking Statements

This presentation contains forward-looking statements that are based on the current expectations and beliefs of Orum Therapeutics, Inc. (“Orum”). Statements in this presentation regarding matters that are not historical facts, including, but not limited to, statements relating to the potential range of usable antibodies, targetable proteins, degrader payloads, and indications, and the potential market for Orum’s products and technologies; the successful development of a multiple-platform portfolio; the modularity of the PROTAb platform; the selection of targets; the translation of preclinical studies to successful clinical trials; the commercial and therapeutic potential of Orum’s technologies and product candidates; Orum’s expectations with respect to its pipeline, including, but not limited to, timelines, and the potential for the filing of Investigational New Drug applications; the potential to receive milestone and royalty payments from technology sales and licensing transactions and to enter into additional transactions in the future; and the adequacy of past, and the potential for future, fundraising are forward-looking statements. These forward-looking statements are based on management’s expectations and assumptions as of the date of this presentation and are subject to numerous risks and uncertainties, which could cause actual results to differ materially from those expressed or implied by such statements. These risks and uncertainties include, without limitation, the uncertainty of success in research and development activities; risks related to clinical trials, including potential delays, safety issues, or negative results; competition from alternative therapies; reliance on third-party decisions in regard to the technologies we license to them; the risk that we may not be able to maintain and enforce our intellectual property; the risk that our product candidates may not be successfully commercialized or adopted; the impact of any pandemic or other global event; the availability of financing; and risks related to the recruitment and retention of key employees, fluctuating markets and economic conditions, health care reform, prices, and reimbursement rates. The forward-looking statements in this presentation speak only as of the date of this presentation, and we undertake no obligation to update or revise any of the statements. We caution investors not to place considerable reliance on the forward-looking statements contained in this presentation. The contents of this presentation are the proprietary information of Orum and may not be reproduced without its prior consent.

# Orum is a Leader in Degradable-Antibody Conjugates

Daejeon, KR

Boston, US



**First company** to develop a clinical stage **DAC**



**Three FDA IND packages**



**Innovative and proprietary** platform technologies



**Two validating transactions:**  
BMS and Vertex



**Upcoming key events:**  
ORM-1153 – Phase 1 initiation  
ORM-1023 – DC nomination




**Our team:**  
54 employees  
>60% R&D | Daejeon & Boston

# Robust CAGR Expected in ADC Market Amid Increasing Demand for Novel MoA Payload

Market expected to reach \$28B by 2028 (23% CAGR), with ADCs accounting for 30% of global biopharma licensing agreements

Success of maytansinoids, auristatins, and camptothecin payload classes has ushered in a golden era of ADCs, but has also intensified competition as dearth of payload MOA remains



Raffaele Colombo et al. Cancer Discovery (2024)

## ADC payload types

### Tubulin inhibitors

e.g. auristatin, maytansinoid

#### MoA

Inhibit microtubule polymerization, causing mitotic arrest

#### Known Limitation

Peripheral neuropathy, Efflux resistance

### DNA damaging agents

e.g. calicheamicin, PBD

#### MoA

Induce DNA strand crosslinking or cleavage, halting replication

#### Known Limitation

Severe toxicity, Poor stability

### Topoisomerase1

e.g. DXd, SN-38

#### MoA

Inhibit Topo1, leading to accumulation of DNA breaks during replication

#### Known Limitation

Off-target toxicity from free payload, GI toxicity, Interstitial lung disease




## Next-generation payloads are emerging

Novel payloads with improved stability, lower toxicity, new MoAs and the ability to overcome resistance are critical to advance next wave of ADC innovation

# What is a Degradator-Antibody Conjugate (DAC)?

Next generation antibody drug conjugate (ADC) with potential for broad intracellular target diversity



Conventional ADC

VS

Orum TPD<sup>2</sup> DAC

Antibody

Confers selectivity and binding against cell surface antigen, allowing **targeted delivery to target cell**

Linker

Connects payloads to antibody with **stability during circulation**, and **effective release of payload** in presence of cleavage triggers

Payload


- Cytotoxic payload
- Limited MoA

- Targeted protein degrader
- Differentiated MoA



# TPD<sup>2</sup> Dual Precision Mechanism of Action: Targeting for Enhanced Safety & Efficacy

Selective tumor targeting combined with targeted protein degradation drives tumor cell death



✓ *Potentially Improved Efficacy*

✓ *Potential Improvement in PK*

✓ *Potentially Improved Safety & Therapeutic Index*


# Orum's TPD<sup>2</sup> Presents Potential Improved Efficacy, PK and Safety in Pre-Clinical Studies

In pre-clinical studies, Orum's TPD<sup>2</sup> approach presented the potential for each of the following:

A

## Improved Efficacy

*Strong activity in vitro and in vivo*




*TPD<sup>2</sup> shows greater in vivo tumor suppression*

B

## Significant Improvement in PK

*Better and optimal pharmacokinetics*




*Exposure of degraders increased by >100x*

C

## Improved Therapeutic Index (TI)


*Better selectivity and safety profile*



*Enables selective tumor activation, improving the TI by lowering MED and raising MTD*

# Dual-Precision Approach Uniquely Positions Orum Amongst Competitors

TPD<sup>2</sup> uniquely combines the specificity of antibody-drug conjugation with the robust efficacy potential of targeted protein degradation





# Two Deals Completed to Highlight the Value of TPD<sup>2</sup> Technology

## Strategic partnerships enabling clinical development and broader application of TPD<sup>2</sup>

### Asset Sale



#### Partner overview

- Global biopharma leader in oncology, hematology, and molecular glue degraders

#### Deal summary

- Acquired ORM-6151 (BMS-986497)
- **\$100M upfront**; up to **\$80M** in milestones
- BMS assumed full responsibility for all aspects of the program

#### Program highlights

- Phase 1 commenced in May 2024 (NCT06419634)
- Target Indications: AML and MDS

### Platform Deal



#### Partner overview

- Global biotech leader with approved medicines for genetic diseases and a robust pipeline across modalities targeting serious diseases grounded in human biology

#### Deal overview

- Platform deal focused on TPD<sup>2</sup> DACs as **targeted conditioning agents** for use with **gene editing**
- License & option agreement: **up to 3 target-exclusive licenses**

#### Deal economics

- **\$15M upfront payment**
- **Up to \$310M per target** in development and commercialization
- Tiered royalties on future net sales

# Multiplatform Pipeline to Address High Unmet Medical Need

## Dual-precision targeted protein degradation (TPD<sup>2</sup>) approach technologies

| Platform                      | Program               | Target      | Indication                | Discovery                                                                                                      | IND-enabling | Phase 1 |
|-------------------------------|-----------------------|-------------|---------------------------|----------------------------------------------------------------------------------------------------------------|--------------|---------|
| TPD <sup>2</sup> GSPT1        | ORM-1153              | CD123       | AML, Heme-onc             |                                                                                                                |              |         |
|                               | ORM-1023              | Undisclosed | SCLC, NET                 |                                                                                                                |              |         |
|                               | BMS-986497 (ORM-6151) | CD33        | AML, MDS                  |                                                                                                                |              |         |
| TPD <sup>2</sup> New Payloads | Multiple              | Undisclosed | Solid Tumors              |                                                                                                                |              |         |
|                               | Multiple              | Undisclosed | Immunology & Inflammation |                                                                                                                |              |         |
| TPD <sup>2</sup> Undisclosed  | Vertex Partnership    | Undisclosed | Pre-conditioning          |  Up to 3 exclusive licenses |              |         |

SCLC: Small Cell Lung Cancer. NET: Neuroendocrine Tumor. AML: Acute Myeloid Leukemia. MDS: Myelodysplastic Syndromes

## Adaptor linker platform

| Platform                            | Description                                                            |
|-------------------------------------|------------------------------------------------------------------------|
| TPD <sup>2</sup> PROTA <sup>b</sup> | E3 ligase-DAC adaptor linker platform enabling novel payload discovery |



## TPD<sup>2</sup> GSPT1 Platform

### Antibody-enabled Molecular Glues


- ORM-1153
- ORM-1023

# GSPT1 Degraders Are Highly Potent Against a Broad Set of Tumor Types


## GSPT1 is a housekeeping protein involved in translation termination

Degradation of GSPT1 → Integrated stress response → Prolonged response  
→ Caspase activation and apoptosis

Orum's SMol006 (GSPT1 degrader) binds GSPT1 and recruits E3 ligase complex



Ubiquitination and proteosomal degradation of GSPT1




Integrated stress response & apoptosis



Disrupted protein synthesis


Disrupt translation termination



GSPT1 binds eRF1 to mediate stop codon recognition and nascent protein release

## *In vitro* screen of GSPT1 degrader against 250+ cell lines across 20+ tumor types

Hematologic malignancies and NETs, including SCLC, show particular sensitivity to GSPT1 degradation



Note: Each dot represents a cell line.

# ORM-1153: CD123-Targeting GSPT1 DAC

## Combining CD123-targeted delivery with GSPT1 degradation

### Antigen target:

CD123

- Highly expressed in AML, >97% at diagnosis | >98% at relapse
- Low expression in normal HSPCs

### Indication:

Acute myeloid leukemia (AML),  
CD123-expressing hematologic malignancies

### MoA:

CD123-directed GSPT1 protein degradation

### Status:

Phase 1 study

### Antibody


Orum's proprietary Fc-silenced CD123 antibody with enhanced internalization and cross-reactivity to cynomolgus monkey

### Payload

SMol006: proprietary CRBN-based molecular glue degrader targeting GSPT1. Released in the cytosol with limited intrinsic cell permeability

### Linker

$\beta$ -glucuronide linker.  
Interchain cysteines for conjugation with DAR ~4<sup>1</sup>



1. Cartoon location of conjugation only for ease of representation

# TPD<sup>2</sup> GSPT1: Opportunity in Acute Myeloid Leukemia (AML)

## Unmet Needs

### Improving efficacy for R/R patients

Significantly lower efficacy in relapsed/refractory patients than in newly diagnosed AML patients (mOS ~11 mo.)

### Demand for Durable & Less Toxic Therapies


Despite initial CR with venetoclax+HMA, long-term remission is difficult to achieve in unfit patients

### No SoC for TP53m-mutated AML

TP53m patients, accounting for ~5-10% of all cases, show poor response to chemo and face high-risk prognosis, with no established SoC

## AML Market Size (global) & Patient Age Trends

Market projected to grow from \$3.5B in '24 to **\$9.6B** by '34 (10.6% CAGR), with **North America** making up 37% in '24




Source : Towards Healthcare; American Cancer Society



AML is uncommon in people under the age of 45

**68**  
Median diagnosis age

## AML Patient Segmentation (as of 2025; U.S.)



## Standard of Care

| Front Line           | Unfit                                                                                                       | Fit                                                                                                         |
|----------------------|-------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|
|                      | Venetoclax + HMA <sup>1)</sup>                                                                              | Intensive Chemotherapy And / or Targeted Therapy <sup>2)</sup><br>± Hematopoietic stem cell transplantation |
| Recurrent Refractory | Intensive Chemotherapy And / or Targeted Therapy <sup>3)</sup><br>± Hematopoietic stem cell transplantation |                                                                                                             |

<sup>1)</sup> HMA (Hypomethylating Agent): azacitidine or decitabine

<sup>2)</sup> FLT3i (midostaurin, quizartinib), IDHi (ivosidenib), CD33-ADC (gemtuzumab ozogamicin)


<sup>3)</sup> FLT3i (gilteritinib), IDH1i (ivosidenib, olutasidenib), IDH2i (enasidenib), Menin-i (revumenib, ziftomenib)




# ORM-1153's Antibody Demonstrates Enhanced Internalization Efficiency

Sub-nanomolar CD123 binding enables rapid and efficient internalization, and ORM-1153 shows markedly greater in vivo efficacy vs. talacotuzumab-GSPT1

## Internalization efficiency in MV-4-11 (24 hrs)



## Disseminated MV-4-11-Luc xenograft mouse model




| Binding kinetics  | ka (1/Ms) | kd (1/s) | KD (M)   |
|-------------------|-----------|----------|----------|
| ORM-1153 antibody | 8.33E+06  | 5.38E-03 | 6.46E-10 |
| Talacotuzumab     | 8.24E+05  | 5.20E-06 | 6.30E-12 |
| Pivekimab         | 2.90E+06  | 7.35E-05 | 2.53E-11 |
| HT12-GL           | 6.01E+05  | 4.89E-05 | 8.13E-11 |

\*Talacotuzumab: Anti-CD123 humanized monoclonal antibody (formerly CSL362)


\*\*HT12-GL: antibody portion of AZD9829 (CD123-Topo1 ADC)

\*\*\*pHAb Reactive dye




# ORM-1153 Exhibits CD123-Dependent Activity and Strong Potency in AML Cell Lines

## CD123-dependent cytotoxicity in CD123+/- AML cells



## Comparative potency against SoC agents in MV-4-11



- Robust CD123-dependent cytotoxicity is observed, with strong activity in CD123<sup>+</sup> AML cells and negligible activity in CD123<sup>-</sup> cells. ORM-1153 demonstrates ~1,000-fold higher activity than the non-conjugated GSPT1 degrader.
- In MV-4-11(CD123<sup>+</sup>/CD33<sup>+</sup>) cells, ORM-1153 shows ~1,000-fold greater potency than venetoclax and is comparable to gemtuzumab ozogamicin (Mylotarg)


\*Gemtuzumab ozogamicin (Mylotarg): a CD33-targeting ADC approved for the treatment of AML

# ORM-1153 Potently Inhibits Growth in AML TP53 Isogenic Cell Lines


Cytotoxicity retained in TP53 KO cells with ORM-1153 indicating potential activity in TP53 mutant AML

## MV-4-11 TP53 WT/KO cells


### ORM-1153




### Venetoclax



### Pivekimab sunirine (Decnupaz™)



### Pivekimab-SMol249\*




| Test article       | DAR | Fold change (KO / WT) | TP53 WT IC <sub>50</sub> (M) | TP53 KO IC <sub>50</sub> (M) |
|--------------------|-----|-----------------------|------------------------------|------------------------------|
| ORM-1153           | 4   | 0.90                  | 1.66E-12                     | 1.50E-12                     |
| Venetoclax         | -   | 16.26                 | 2.09E-09                     | 3.40E-08                     |
| Pivekimab sunirine | 2   | 57.95                 | 2.64E-12                     | 1.53E-10                     |
| Pivekimab-SMol249* | 2   | 1.23                  | 6.93E-12                     | 8.58E-12                     |

\*Pivekimab-SMol249 is a control generated by conjugating SMol249, the linker-payload used in 1153, to the 442C mutation site of pivekimab, the antibody used in IMGN632

# Robust ORM-1153 Activity in AML PDX Models Regardless of Gene Alterations

Consistent efficacy maintained across high-risk profiles including TP53 and FLT3-ITD

## Ex vivo cytotoxicity and patient sample characteristics (#1 - #7)



| Sample | Gender | Age | Diagnosis | Treatment  | Gene Alteration             | CD123 (ABC) |
|--------|--------|-----|-----------|------------|-----------------------------|-------------|
| #1     | F      | 59  | Relapsed  | Pretreated | IDH1, NPM1, FLT3-ITD        | 11131       |
| #2     | F      | 54  | De novo   | Naïve      | TP53, NPM1, FLT3-ITD, KMT2A | 4775        |
| #3     | M      | NA  | De novo   | Naïve      | WT*                         | 4503        |
| #4     | F      | 68  | De novo   | Naïve      | TP53, FLT3-ITD              | 3566        |
| #5     | M      | 80  | De novo   | Pretreated | FLT3-ITD                    | 3485        |
| #6     | M      | 60  | Relapsed  | Pretreated | TP53, IDH2, NPM1            | 3197        |
| #7     | F      | 21  | De novo   | NA         | WT*                         | 2453        |


\*Not associated with TP53, FLT3, IDH1/2, NPM1, or KMT2A gene alterations  
NA – not available, ABC – antibodies bound per cell

# Robust Tumor Control and Survival Benefit of ORM-1153 in Disseminated AML Model


ORM-1153 drives greater reduction in tumor burden and robust survival compared to SoC and benchmark ADC

## Disseminated MV4-11-Luc xenograft mouse model



### Greater reduction in AML burden over time



### 93% tumor reduction at Day 14



### Robust overall survival at MED




▲ = Dosing day

\*Aza/Ven = Standard of care, azacitidine (2.5 mg/kg, IP QDx5) + venetoclax (100mg/kg, PO 5days/week x 3)

\*\*AZD9829 is a discontinued clinical anti-CD123 ADC with a TOP1 inhibitor payload at DAR8

# Broad Therapeutic Window and Favorable Safety Profile of ORM-1153

## Exposure-based preliminary therapeutic window



- ✓ **Wide therapeutic margin established** between the minimal efficacious dose (MED; 0.1 mg/kg in mice) and the highest non-severely toxic dose (HNSTD; 6.0 mg/kg in NHPs).
- ✓ **Safety profile** in repeat-dose NHP studies, with off-target effects including mild, transient, and reversible reductions in platelets and increases in liver enzymes

# Phase 1 First-in-Human Study of ORM-1153

## Phase 1 Multi-region, First-in-Human Study of ORM-1153 in Patients with R/R AML or Other Hematological Malignancies with CD123 Expression



### Methodology

Open-label, multiregional, Phase 1 dose escalation



### Estimated Enrollment

42 participants



### Sites

4 sites in the US



### Primary Objective

Determine the optimal biological dose (OBD); safety & tolerability



### Secondary Objective

PK, PD, immunogenicity and preliminary anti-leukemic activity




### Duration

24 months (from FPD) to primary completion

### Phase 1 Dose Escalation

CD123 expressing R/R AML

Dose escalation decisions will be governed by Bayesian Optimal Interval Design (BOIN), with dose escalation following a modified Fibonacci sequence.



\*MTD: maximum tolerated dose



# ORM-1023: GSPT1-DAC for Small Cell Lung Cancer and Neuroendocrine Tumors

## Combining targeted delivery with GSPT1 degradation

### Antigen target:

Undisclosed

### Indication:


Small cell lung cancer (SCLC),  
Neuroendocrine tumors (NET)

### MoA:

Target-directed GSPT1 protein degradation

### Status:


DC nomination expected in Q4 2026



# Limited Diversity of Payloads for SCLC ADCs in Development

Antibody targets vary widely, yet the payload heavily relies on TOPO1

## Comparative efficacy landscape of ADCs and TCEs in relapsed small cell lung cancer



- 8 of 9 ADCs with reported Phase I/II efficacy use a TOPO1 payload
- Antibody targets vary widely across B7-H3, DLL3, TROP2, SEZ6 and EGFR/HER3, yet the payload has converged on TOPO1 inhibition


A promising emerging therapy landscape, but concentrated in a single payload class

# Potent Activity of ORM-1023 in SCLC PDX-Derived Models


ORM-1023 showed sub-nanomolar potency across three PDX-derived lines, including the LU5264 model, where lurbinectedin was inactive (>200 nM)

## Ex vivo potency of ORM-1023 in SCLC PDX-derived models\*


LU5217



LU5264



LU5242



| Test Article  | Mechanism                    | IC50 (nM) |        |        |
|---------------|------------------------------|-----------|--------|--------|
|               |                              | LU5217    | LU5264 | LU5242 |
| ORM-1023      | GSPT1 degradation            | 0.279     | 0.186  | 0.017  |
| Lurbinectedin | RNA polymerase II inhibition | 1.846     | >200   | 0.22   |


\* Target antigen expression confirmed by IHC.

# Robust In Vivo Activity of ORM-1023 in SCLC PDX Model


Single-dose ORM-1023 demonstrated potent anti-tumor activity with stable body weight

## In vivo activity of ORM-1023 in LU5242 SCLC PDX model\*

Tumor Volume



Body Weight



\* Target antigen expression confirmed by IHC.




## TPD<sup>2</sup> Novel Degradable Payload Discovery

Enabled by Orum Know-How and  
PROTA Platform

# Orum DAC Discovery Strategy for 2026 and Beyond


## GSPT1 and next-generation payload : Unlocking platform potential across diverse targets

- Address limited activity or resistance associated with conventional payloads




## Indication-tailored payloads: Matching payload-antigen pairs to disease contexts

- "Ultra-specific" dual-targeting approach, with both antibody and payload aligned to disease-specific biology to maximize therapeutic index




# Structure-based Discovery Workflow at Orum: From Target to Lead Candidate




# TPD<sup>2</sup> PROTA<sup>b</sup>: Linker Platform to Facilitate the Development of DACs

## Adaptor for E3 ligase-based degrader conjugation



### 1 CRBN-based degrader 2 Other E3 ligase degrader



Extended to other E3 ligases

Note: Structure of SMol006 molecular glue binding to CRBN E3 ligase

## PROTA<sup>b</sup>-enabled traceless conjugation process


Step 1. PROTA<sup>b</sup> attachment to ADC linker



Step 2. Conjugation to E3-ligase degrader



Step 3. Intracellular traceless release



### Efficient, time-saving development path

- ✓ Reduces DAC development time by 6 to 9 months by eliminating trial-and-error chemistry through moiety-driven molecular adaptation for efficient linking and conjugation

### Compatibility with diverse degrader payloads

- ✓ Offers a universal conjugation handle for both MGDs and heterobifunctional degraders

### Broad linker compatibility

- ✓ Compatible with diverse linker designs, including protease-cleavable, thiol-sensitive, and  $\beta$ -glucuronide-cleavable linkers, etc.




# PROTAAb Enables Targeted Degradation with Diverse Hetero-Bifunctional Degraders

Demonstrates effective use of PROTAAb with non-GSPT1 targets and hetero-bifunctional degraders


## BCMA-BRD4

Anti-BCMA mAb



BRD4 Degradar

## Targeted BRD4 degradation drives cytotoxicity




POC molecules PROTAAb-DACs are highly effective *in vitro*

- ✓ BCMA-BRD4 improved potency by >3 logs compared to a small molecule alone
- ✓ CD79B-IRAK4 induced IRAK4 degradation at low nanomolar range, while maintaining antibody-level target binding affinity

→ Supports platform scalability across diverse intracellular targets


## CD79B-IRAK4

Anti-CD79B mAb




IRAK4 Degradar

## Maintains antibody-level target binding affinity



Dose-dependent degradation of IRAK4 at low nanomolar range


Target-mediated degradation via CD79B binding




# TPD<sup>2</sup> : Significant Potential to Unlock a Vast Number of Indications

## Modular platform technology allows for creation of a multitude of targeted therapies

- Orum's strength lies in its ability to **develop** the antibody and degrader payload separately, tailoring these components to **enhance therapeutic benefit** in each indication




# Expected Timeline for 2026 and Beyond



## Current Partnering Focus

- **Out-license** Orum's programs and platform technology
- Access to complementary external technologies through **collaborations** or **in-licensing**

A dramatic landscape photograph of a snow-capped mountain peak, likely Mount Everest, under a sunset sky with orange and purple clouds. The foreground shows dark, rocky terrain with patches of snow.

# Thank you

A world where no patients suffer from diseases caused by  
undruggable targets