

....


SYMPTOMS & SIGNS – MANIFESTATIONS & NATURAL HISTORY


Outcome of acute HBV infection

90 percent for a perinatally acquired infection would go on to have chronic HBV infection

20 - 50% of those who acquired the infection between 1 and 5yrs develop chronic HBV

Adults - 5%






SYMPTOMS & SIGNS – MANIFESTATIONS & NATURAL HISTORY

••••

Treatment for acute HBV

Treatment for acute HBV is mainly supportive




SYMPTOMS & SIGNS – MANIFESTATIONS & NATURAL HISTORY

•••

Hospital management is based on

- Coagulopathy, deep jaundiced or encephalopathy
- · elderly age at first infection
- have significant comorbidities
- cannot tolerate oral intake
- have poor social support systems



SYMPTOMS & SIGNS – MANIFESTATIONS & NATURAL HISTORY

•••

Antiviral therapy is not indicated, except in a few cases of fulminant hepatic failure, but which should be stopped once the patient has become negative to HBsAg



What we have learned

- HBV is a viral infection that causes acute and chronic diseases
- Transmission of HBV in developing countries is most often horizontal, whilst it is via sexual routes in the West
- Acute HBV presents more dramatically than chronic HBV
- HBV treatment is supportive and rarely needs antiviral agents

