

The Loop Mega Universe

V3.13 — Consolidated Mathematical & Empirical Pass

owned equations · LMU couplings · results · verification · audit

Pitarn Rungsiyapornratana

Worldview and architecture by Pitarn; equations, integrations, and the proven/assumed audit by the editor pass; assembled under the R6 protocol (all numbers recomputed, none inherited). 2026.

Tags: **[Fact]** measured/peer-reviewed **[Fact-theory]** established theory **[Hypothesis]** literature support **[Speculation]** no equations yet **[Design]** structural choice **[Open]** unsolved **[Dead-end]** failed **[Auditor]** editor addition. Owners of borrowed equations are printed at the right margin.

What this document is. This is the single consolidated record of the Loop Mega Universe mathematical and empirical pass: the merge of the V3.4 consolidated pass (Parts I, III, VI here), the dark-energy verification record (Part II), and the equation-document spine that incorporated Operator B' and Correction 1 (Parts 0, III§2, IV, V, VII, VIII). The spine's structure governs; the archive's content is carried essentially verbatim and edited only where the reconciliation deltas ($\Delta 1$ – $\Delta 11$, recorded in the build specification) demand it. LMU is a *lens/synthesis*, not a predictive theory; where it overlaps Penrose's CCC (~ 80 – 90% of the spine) the divergences are stated explicitly. The single declared framework-level point of failure is **fork 5** (unbounded $L0$); every result that rides on it says so. **[Design]** / **[Hypothesis]**

Revision history (3.4 $\rightarrow$ 3.5). Operator B' formalized (export-partition, Part IV); Correction 1 — the interlude background is a *dust era*, not constant- Λ (Part III §2); the Tolman rate g is now *measured*, not bounded (floor 1.7×10^{13} , fiducial 4.1×10^{14} , post-ladder $\sim 2 \times 10^{16}$); reconciliation deltas $\Delta 1$ – $\Delta 11$ applied; a retired-numbers table added (Part V §4 — nothing on it may be cited as live); A-field Run 2 (A_i fixed, m shot) added and cross-validated against the m -scan (Part II). Axioms unchanged; V4.0 is reserved for a physics verdict (DESI-Y5 falsifier ~ 2027 , M87 ngEHT/BHEX, or a solved rebirth mechanism).

Revision history (3.5 $\rightarrow$ 3.6). Ledger pass; the computational queue is now empty. **(F) resolved-conditional:** the A-dust Jeans gate computed — island scales are super-Jeans from oscillation onset ($R_{\text{isl}}/\lambda_J = 2895$ at the live point), the A-dust clusters with $p \rightarrow 1$ for all $m \gtrsim 0.14 H_0$, and the gate closes only by horizon exile at $m \lesssim 0.13 H_0$, outside the DESI-thaw identification (Part III §2, Fig. 2; conditional on $V_{\min} = 0$). **(E) resolved with a correction:** $\beta \approx 0.7$ was an orphan matching the *stellar* BCG-halo slope; the cluster-end owned slopes (Bogdán et al. 2018 ≈ 1.05 ; Bassini et al. 2019 1.35 ± 0.15) give 0.15–0.27 dex per decade — the headline survives at the range's lower edge, the derivation is replaced (Part V §1). **(G) closed by retirement** (author ruling): no constant- Λ medium at any stage; the F5 numbers stay permanently quarantined; fork-5 *delivery* unaffected. **(H) decided:** past-eternity adopted as an axiom-level choice riding Aguirre–Gratton 2002, with the explicit construction split off as **(H1) [Open]**. Axioms otherwise unchanged; V4.0 remains reserved for a physics verdict.

Revision history (3.6 $\rightarrow$ 3.7). Stress-audit pass with measured inputs and the $V_{\min}$ Decision Record (2026-06-11). **(i) Audit:** the full loop $\Phi = \Lambda_{\text{live}} \circ B' \circ \Lambda_0$ reproduces independently from measured inputs alone (Part VII-DR §1); six wording corrections applied: the DESI claim is now the joint locus distance 2.0–2.9 σ (covariance-dependent; $\approx 2.7\sigma$ at $\rho \approx -0.87$, marginal w_0 within 1.3 σ ; Part II); the $V_{\min} = 0$ knife-edge is *priced* ($|V_{\min}|/\rho_{\text{DE}} < 4 \times 10^{-3} / 5.5 \times 10^{-56} / 3.1 \times 10^{-181}$ for pairs / island hierarchy / full program; items B, J2); the M87 flag carries its inversion ($N_{\text{mis}} \leq 2.3$ of 5.9; aligned fraction $\geq 62\%$); the survivor band is *restoring* (return-map slope $\sim 10^{-15}$) with a computed Jeans floor $6.4 \times 10^5 M_\odot$ and no ceiling; fork-5 carries its exchange rate $n_{\text{max}} = \log_{10}(V_{L0}/V_{\text{isl}})/14.6$; the in=out ledger is scoped to the $L0$ debris channel,

the boundary injection $\times\Omega \in [10^{26}, 10^{47}]$ stated. **(ii) New:** the fourth endgame runner Ω_k priced (flatness joins the B3' re-init list, item I); **Decision Record DR-Vmin** (Part VII-DR): the trichotomy $V_{\min} \in \{0, +\epsilon, -\epsilon\}$ with prices, the per-zone/population reading (path 4), and the cascade-to-zero mechanism (path 4b, author wiring) with $V = 0$ as the absorbing state; new ledger items (K), (L). **(iii) Ruling:** item (K) **ruled-conditional** same session — paths 4+4b adopted ($V = 0$ as the selected attractor; pathways set by birth draws); the K1 wall accounting computed and closed-conditional (Part VII §5): $-\epsilon$ zones implode to BH corpses outside and bounce+re-roll inside; $L0$ has no conversion channel. Spine numbers unchanged. **(iv) Item (A) resolved-conditional/drafted** same session (Part VII §7): under the ruling nothing draws exact zero, so the $+\epsilon'$ lineages end de Sitter — spacelike $\mathcal{J}^+$ restored, Penrose crossing native; window computed ($\epsilon'_{\text{crit}} = 6.3 \times 10^{-181}$ re-derives the program tier); construction E1–E4 drafted with proof obligations P1–P3 (P2 = (C) sharpened, P3 = (D) sharpened); (H1) gains its arena. Axioms unchanged; V4.0 remains reserved for a physics verdict.

Revision history (3.7 → 3.8). Same-day continuation (2026-06-11), three items. **(i) Chain pull executed:** the released DR2 chains (DESI Collaboration, Zenodo 10.5281/zenodo.16644576; run `base_w_wa`, DESI+CMB+Pantheon+; sample means/ σ reproduce the published values exactly) give $\rho(w_0, w_a) = -0.894$, burn-in-stable; the joint locus distance pins to 2.69σ (chain-Gaussian) and 2.40σ (posterior-mass — the convention in which DESI's own Λ CDM exclusion is 2.8σ , reproduced here as 2.83σ); on the released posterior the thawing locus is *less disfavoured than Λ CDM itself* in both metrics; $\epsilon_{\text{DESI}} = 4.6 \times 10^{-2} \rho_{\text{DE}}$ (joint, chain covariance) — the “pending chain covariance” flags come off (Part II). **(ii) (H1) ruled and drafted** (Part VII ledger): anchor choice (a) per-lineage (author ruling); equations H-E1–H-E4; per tree the past is eternal in proper time and *finite in aeon count* (a second, independent escape from the Kinney–Stein bind); obligations **P1 sharpened** (the Tod 2007 anisotropic-crossover obstruction — the magnetic-Weyl normal-derivative jump — must close under the e^{-4N} budget) and **P4 new** (geodesic exhaustion; reduces to $P1 \wedge$ the KMS24-vs-FPX25 dispute $\wedge$ LQC-effective completeness, modulo wall-riding pathologies). **(iii) Borrowed-legs audit v2:** GHZ24 = García-Saenz–Hua–Zhao (PRD 110, L061304; the completeness checklist, necessary not sufficient); FPX25 = Ferreira–Pinto-Neto–Xavier (PRD 111, 123531; C^2 extension iff past-asymptotically dS, NEC-free kinematic bounce, pointed at CCC against Kinney–Maity–Sriramkumar 2024 — dispute open, travels); Kinney 2026 (arXiv:2604.20809) re-kills IS-cyclic only; Leg-1 fixes (Montero–Valenzuela 2412.00189 targets the *SUSY* DGKT; Heidenreich–Lotito prove the particle, not membrane, WGC). Session records: `LMU_borrowed_legs_audit_v2.md`, `LMU_H1_sketch_arena.md`, `LMU_P1_P4_first_pass.md`, `LMU_F1_pin_record.md`. Axioms unchanged; V4.0 remains reserved for a physics verdict.

Revision history (3.8 → 3.9). Block-1 computational pass + ledger principle (2026-06-11, late session). **(i) P1 FRW core closed** (record `LMU_P1_block1_pass.md`, script `lmu_p1_frw.py`): integer-exponent smoothness table for every carried component; the oscillating A -residue makes the crossing exactly $C^{2,1} - C^2$ (the Einstein/FPX requirement) holds with *zero margin*, C^3 is log-periodically blocked (a new priced knife-edge); the child bang is radiation-exact, the parent dust mapping to a relative- $O(\hat{a}^3)$ correction vanishing into Σ ; the FPX extendibility hypotheses are verified at the root window. **(ii) Obstruction ladder measured:** frozen super-horizon tensors set the slowest Weyl channel — $E \propto e^{-2N}$; B and the Tod-2007 jump channel $\partial_n B \propto e^{-3N}$; fluid-sourced Weyl e^{-4N} (erratum applied, Part VII §7). **(iii) P2 closed-conditional** by horizon exile (item (C) closes on the same budget; falls with (K)); P3 gains its background map (the c_3/c_4 series). Open remainder of block 1, named: **G1** (reciprocal gluing of Friedrich-1986 $\mathcal{J}^+$ data into the Anguige–Tod admissible class) and **G2** (contaminated-bang well-posedness) [math-GR]. **(iv) Ledger principle** adopted (Part IV §2): *nothing is lost without a receipt*. Axioms unchanged; V4.0 remains reserved for a physics verdict. **Revision history (3.9 → 3.10).** G1/G2 pass (2026-06-11, continuation; record `LMU_G1_G2_pass.md`, script `lmu_g1_g2.py`). **(i) Tod-2003 scope check executed** (the explicit V3.9 checklist item): its scalar case is the stiff-dominant

regime — the field diverges at Σ and carries no free data — the opposite of LMU’s subdominant A -passenger on a radiation bang; the passenger question was unformulated in the literature and is now formulated. **(ii) G1 reduced:** Kopiński–Valiente Kroon 2022 derive the two-sided crossover matching constraints (Friedrich CEFE at $\mathcal{I}^+$ / Anderson–Fefferman–Graham at the bang) and solve them in the conformally-flat/conformally-Einstein sector; LMU’s crossing converges to that sector at the measured e^{-2N} , so G1 becomes *perturbative openness off a solved base point* with every mismatch channel priced ([G1-h/B] automatic, [G1-E] e^{-2N} , [G1-j] e^{-3N}). New conditional [G1- Ω] travels: the $\hat{\Omega}$ prescription is not unique (Newman 2014 / Tod 2015 / Nurowski 2021 mutually disagree; Markwell–Stevens 2023 compare and rule out cases); LMU rides the Tod-style FRW choice. Named check [G1-d]: whether the L  bbe–Tod 2009 extension hypotheses accept L^∞ third-order data — new sharpening: $f'''' \sim 1/\eta$ is *unbounded*, so the crossing is exactly $C^{2,1}$, not $C^{3,\alpha}$. **(iii) G2 reduced:** component coverage matrix built (radiation: Anguige–Tod 1999; $+\Lambda$ homogeneous: Tod 2007; massless Vlasov: Anguige 2000; Boltzmann FLRW: Lee–Nungesser–Tod 2020); new computed admissibility condition **G2-c** — the scalar’s Frobenius indices at a radiation bang are $\{0, -1/2\}$, so admissibility costs exactly one extra condition (the $t^{-1/2}$ stiff-branch amplitude vanishes on Σ ; the regular branch has $\rho_A/\rho_{\text{rad}} \sim \hat{a}^4$, vanishing into Σ); EoS reflection through the reciprocal map computed ($w \rightarrow w' = (1 - 3(1 + w))/3$: parent dust arrives as a $w' = -2/3$ trace image — explaining the V3.9 admixture and retargeting the passenger theorem); named tool [G2-d]: run the Gover–Kopiński–Waldron 2023 generic-stress admissibility criterion over the LMU stress list. **(iv) Parent shelf closed:** Friedrich 2017 (Λ -dust $\rightarrow$ smooth spacelike $\mathcal{I}^+$, FLRW stability) and Friedrich 2023 (asymptotic dust/radiation EoS, extensions beyond the boundary) remove the e^{-3N} dust residue as a parent-boundary threat; with L  bbe–Valiente Kroon 2013 (radiation+ Λ) and Joudioux–Thaller–Valiente Kroon 2021 (massless Vlasov), every carried parent component has a boundary-regularity theorem. **Honesty block** (carried verbatim into Part VII §7’s audit trail): LMU’s massive A -field violates Penrose’s suppressed-rest-mass hypothesis; LMU rides the FPX25 kinematic C^2 route, *not* the conformal-matter crossover, with matter re-seeding inside the T_{reheat} irreducible; Penrose’s Ω -field-as-DM is not used (LMU’s DM is an independent thermal relic). Axioms unchanged; V4.0 remains reserved for a physics verdict.

Revision history (3.10 $\rightarrow$ 3.11). G1-d/G2-d pass (2026-06-11, continuation; record `LMU_G1d_G2d_pass.md`, script `lmu_g1d_g2d.py`). **(i) [G1-d] closed as a check.** L  bbe–Tod 2009 read at source: their Thm 3.3 hypotheses are boundedness-type (Whitney engine — bounded, never continuous, top-order data), so the literal question resolves YES; placing the crossing on their ladder then dissolves it — the FRW spine’s conformal class is exactly flat (tractor curvature $\equiv 0$; Bailey–Eastwood–Gover 1994), so LT09 holds trivially at every order and the $C^{2,1}$ knife-edge constrains $\hat{\Omega}$ only: *[G1-d] merges into [G1- Ω]*. In the TT sector the ladder formalizes the horn dichotomy: the smooth horn passes LT09 at all orders (price: forced child π_{ij} , Tod 2015); the class horn has distributional Cotton (impulsive-wave $C^{1,1}$ interface, Lichnerowicz $[K] = 0$, amplitude e^{-2N}) and sits outside LT09 class (i) — the precise rung of Tod 2007’s “conformal metric not smooth”. **(ii) [G2-d] executed.** The Gover–Kopiński–Waldron 2023 criterion run over the LMU stress list, with the leading trace coefficient $-\frac{3}{2}\alpha(\alpha + 1)t^{2(\alpha-1)}$ *recomputed from raw Friedmann* (exact match, R6). At a radiation bang ($\alpha = -1$) even the t^{-4} trace slot closes; the admissible hierarchy is exactly $\{t^{-1}(H^{\text{ext}}), t^0(\Lambda), t^2, \dots\}$ — and the LMU contamination list fills it slot-by-slot: dust-image $\rightarrow H^{\text{ext}}$ (implied $H^{\text{ext}} = \frac{3}{16}\rho_{d0}/A$; the same coefficient enters $\hat{a}$ only at relative η^3 , $c_3 = A\rho_{d0}/24$ — Block-1 reproduced), frozen- $A \rightarrow \Lambda$, oscillation $\rightarrow$ tail, while the stiff branch ($\sim t^{-6}$) has *no slot*: **G2-c is doubly derived** (Frobenius + causal-structure admissibility). Magnetic contaminant: trace-blind here; the LNST no-go stands. Caveat travelling: GKW §VI’s cross-aeon stress dictionary anchors to the *Nurowski 2021* $\hat{\Omega}$ prescription, so any cross-aeon use inherits [G1- Ω]; the slot results themselves are prescription-independent. Open cores unchanged and isolated: G1 nonlinear gluing off KVK22; G2 inhomogeneous passenger well-posedness (Fuchsian route, LNST JDE 411 (2024) template). Axioms unchanged; V4.0 remains reserved for a physics verdict.

Revision history (3.11 → 3.12). DR-P(k) ruling + empirical sweep (2026-06-11, continuation; record LMU_DR_Pk_and_sweep.md). **(i) DR-P(k) ruled** (author): the primordial spectrum is *per-aeon*, not inherited across Σ_N — formalizing what Block-1 already computed (every transfer channel dies at e^{-2N} or faster, and the slowest survivor is *tensor*, the wrong type and spectrum for the adiabatic scalar A_s). Consequence: each aeon carries its own fluctuation generator **[Open]**, named import] with hard targets $n_s = 0.9649 \pm 0.0042$, $A_s = 2.1 \times 10^{-9}$, $r < 0.036$, $f_{\text{NL}}^{\text{loc}} = -0.9 \pm 5.1$; generator existence is *necessary for loop closure* (no $\delta \sim 10^{-5} \Rightarrow$ no ladder $\Rightarrow$ no next flash). Discount: horizon and flatness are paid by the crossover, so the import is a generator, not full inflation. Wording rule (document-wide): the flash lights the *pre-generator* start; the observed hot start (BBN/CMB) is lit by the generator’s reheating. Bounds repurposed: $r < 0.036$ and PTA limits become crossover *cleanliness bounds*, passed automatically at large N . Stance forced: LMU predicts the *absence* of CCC-style Hawking points/concentric circles (horizon exile) — a clean departure from Penrose’s CCC, whose claimed detections are themselves contested. **(ii) DM minting reworded:** 5:1 is matched, not predicted; exclusion landscape cited (LZ 2024, N_{eff}). **(iii) Empirical sweep added** (Part VI): NANOGrav SGWB (supportive), SLAB limits, dwarf occupation fractions, BBN/fifth-force pass for A , Planck topology pass, NewAthena date note. Axioms unchanged; V4.0 remains reserved for a physics verdict.

Revision history (3.12 rebuild, 2026-06-12). In-place consolidation at the same version number (author request; no physics edits, axioms unchanged). **(i)** Ledger item (D) synced to the DR-P(k) ruling — the ledger text was stale against Part VII §7. **(ii)** Dead-end entry **B-W** added: Weyl-residue disposal/repurposing, both directions dead. **(iii)** Unit audit (R6, this date): 13/13 numeric spot-checks reproduced by independent SI recomputation (τ_{evap} coefficient, M_{Nariai} , pair threshold, S_{BH} , the 4/3 radiation entropy, s/n_γ , Ω_c/Ω_b , the B1' → B4' mint chain, NGC 1277/384 offsets); K_{mrg} unit declared (Gyr^{-1}) and the demo-script divergence flagged (Coefficients section); recoil constants renamed $A_{\text{kick}}, H_{\text{kick}}$ (symbol collision with the A -field and Hubble H); Smoluchowski kernel normalization unit noted. **(iv)** Meissner–Penrose 2025 (arXiv:2503.24263; the GWE crossover mechanism) added as the current CCC-side reference on the Hawking-points stance; a circulating claim of a unit slip in its eqs. (57)–(59) is *not recorded* — unverified at source, quarantined per R6. **(v)** Mission dates applied inline (NewAthena ~ 2037 ; former “ ~ 2035 ” instances updated).

Revision history (3.12 → 3.13). Literature-anchor pass (2026-06-12; session record LMU_V3_13_patch_block). no numeric spine changes, axioms unchanged. **(i)** J2’s theory bullet gains owners: the dS swampland conjecture (Obied–Ooguri–Spodyneiko–Vafa 2018; refined Ooguri–Palti–Shiu–Vafa 2019; cosmological corollary = thawing quintessence toward $V \rightarrow 0$, Agrawal–Obied–Steinhardt–Vafa 2018) and, as an independent quantum route, de Sitter quantum breaking (Dvali–Gómez 2014; Dvali–Gómez–Zell 2017) on the Gibbons–Hawking 1977 finite-entropy base; counter-leg KKLT 2003 attached. **(ii)** The same lane cuts the (A)-window’s “globally future-eternal”: quantum-breaking caveat added with R6-recomputed budgets ($N_Q = 1.39 \times 10^{122}$ at ϵ_{DESI} , 6.60×10^{300} at ϵ'_{crit} ; crossing margin $\gtrsim 10^{120}$ survives). **(iii)** Hawking-points stance sharpened: Meissner–Penrose 2025 upgrades HPs from claimed signal to *structural requirement* of CCC’s crossover; detection numbers attached (AMNP 2020 99.98% vs. Jow–Scott 2020 87%); mandatory stakes flag — a confirmed HP breaks horizon exile itself, re-opening (C), (D)-transfer, and P2. **(iv)** Empirical sweep (vii) added: Giant Arc / Big Ring (Lopez–Clowes–Williger 2022/2024) vs. Sawala et al. 2025 vs. the Lopez–Clowes 2025 rebuttal — contested; scope note: a confirmed Gpc structure is initial-condition physics landing on the item-(D) generator, not on the ladder; a cross-aeon reading joins the Hawking-point falsifier basket. **(v)** R7 returns: Pathria 1972 + Popławski 2010 credited at item (L) for the inside-the-horizon-universe picture; the Kinney–Stein citation completed (2021, arXiv:2110.15380); Itzhaki–Peleg–Steinhardt 2026 (JCAP 05, 084) logged as the dilution lane’s string-embedded realization, to be read against the Kinney 2026 re-kill (V3.8). Grep-audit note: swampland-as-motivation and the Kinney–Stein bind were already integral (Part VI §18; Part IV §1); this pass adds owners, the quantum route, and the eternity caveat only. V4.0 remains reserved

Contents

Part 0 — Reading rules, the state vector, and the cycle map	9
1 The state vector and the cycle map	9
I The matter ladder — L4 to L1 as a dynamical system	11
1 Overview — what this module is, and how to read it	11
2 The equations of motion	11
2.1 Growth — accrete (Gas) plus eat (Near-Galaxy $\times$ DM)	11
2.2 Spin — up by coherent Gas, down by Near-Galaxy mergers	12
2.3 Gas, environment, and the two off-ramps	12
3 Coefficients: what is fixed by physics, what is calibrated	12
4 Computation 1 — four environments, four fates	13
5 What emerges — structural results, not tuned	14
6 Computation 2 — are the structural results real? (robustness)	14
7 Computation 3 — from one hole to a population	15
7.1 Refinement — the high-mass abundance from first principles	16
8 What this model is for — and what it cannot do	16
9 Falsifiers	17
10 Attribution	18
II The dark-energy field A — verification record, runs, and the DESI confrontation	19
1 Equations — the three-party balance	19
2 Result 1 — $w(a)$ versus DESI (the m-scan)	19
3 Result 2 — the $V \rightarrow 0$ endgame (Fork Y’s actual goal)	20

4	Result 3 — transient acceleration (“the mountain”)	21
5	Result 4 — cross-check across potentials (no dialing)	22
6	Layer separation (anti-drift)	23
7	DESI status and the pre-registered falsifier	24
8	Run 2 — A_i fixed, m shot (this session’s cross-validation)	25
9	The PNGB cross-audit against DESI DR2 (the V3.4 archive)	26
10	The field, and what the data ask of it	26
11	Computation — Klein–Gordon vs DESI (O2)	26
12	Honest status	28
13	Attribution	29
III	Closure of the loop — the $L0$ ledger	30
1	The reservoir and what enters it	30
2	The interlude background — a dust era (Correction 1)	31
3	Energy balances exactly — the in=out ledger	31
4	Energy is not the same as usable capacity	33
5	Four worries answered by physics	34
6	The one assumption that remains	34
7	Entropy — diluted, not reset (and why it cannot be reset)	35
8	Honest status — theorem vs assumption	35
9	Attribution	36
IV	Operator B' — the aeon transition (export–partition form)	37
1	$B1'$ — the export ledger and the measured rate g	37
2	Why entropy is the invariant channel — the conformal boundary and z_{aeon}	37
2.1	What the map fixes (the derivable part)	37

2.2	Why the entropy route is blocked (the auditor result)	38
3	B2' — sorting and partition	38
4	B3' — per-round re-initialization	38
5	B4' — minting	39
6	Endowment gathering — no unique parent (the surviving S7 addendum)	39
V	Results of the composed system	40
1	Per-island fixed band	40
2	The two-sector verdict and the Tolman ledger	40
3	Predictions and standing tensions	41
4	Retired numbers (V3.5) — nothing on this list may be cited as live	41
5	R6 recomputation record (this assembly)	42
VI	Verification and empirical confrontation	43
1	Independent verification of the analytic and ODE sectors	43
2	The ladder as a single forward map	43
3	Mass–spin separability — the decoupling is mass-robust	43
4	The spin-direction model — magnitude robust, axis fragile	44
5	Confronting the distinctive prediction with data	44
6	Attribution	46
7	The relic over-massiveness against the 2025 dynamical-mass set	46
8	NGC 384 — the out-of-sample test point	47
9	The dwarf empty-centre axes (position/occupation, mass-free)	47
10	Galaxy spin as phase — complement to the BH-spin claim of Part IV	47
11	The audit verdict — a structural result	48

12 Action items for the main V3.4 paper	48
13 Stress-test — every checkable number reproduces independently	49
14 Stress-test — the dark-energy field reproduces the no-crossing claim	49
15 Stress-test — spin decoupling: sign and mass-invariance reproduce; magnitude does not	50
16 Stress-test — Part V empirical numbers reproduce exactly	50
17 Open joint 1 — super-Eddington feedback: strengthened, but “evades” is the wrong word	50
18 Open joint 2 — sign of $V_{\min}$: the framework sits on the less-favoured fork	51
19 Open joint 3 — f_{coh} coherence: sound logic, but “mild” is optimistic and the cocoon bet is unsettled	51
20 Open joint 4 (new) — the engine and the spin condition pull against each other	52
21 Audit verdict — one distinctive claim on a three-link conditional chain	52
 VII Stress audit with measured inputs, and Decision Record DR-Vmin	 54
1 The stress audit — the spine reproduces; seven findings	54
2 DR-Vmin — the trichotomy with prices	54
3 The fourth runner — spatial curvature Ω_k	55
4 Path 4 — the per-zone reading	56
5 Path 4b — cascade-to-zero (author wiring), and the absorbing state	57
6 Status, rulings required, and the audit trail	57
7 The (A)-window — the crossing restored on the ruled spine	58
 VIII Open ledger and dead-end pointers	 61
1 The open ledger (one list, no duplicates)	61
2 Dead-end pointers (check before extending anything)	63

Part 0 — Reading rules, the state vector, and the cycle map

“Every different pathway to the same end zone.”

— the loop’s equifinality (von Bertalanffy, general systems theory; no-hair: Israel 1967, Carter 1971; cosmic no-hair: Wald 1983)

Reading rules. (1) Every numbered equation in this document is *established physics with a named owner*, printed at the right margin where the equation is introduced. Nothing in an equation box is claimed by the author. (2) The LMU content is confined to the **couplings** — which output variable of one equation is fed into which input of the next — and to the **results** of running the composed system. Couplings are marked $\hookrightarrow$ and carry epistemic tags: [\[Fact\]](#) [\[Fact-theory\]](#) [\[Hypothesis\]](#) [\[Design\]](#) [\[Open\]](#) [\[Dead-end\]](#). (3) LMU is a lens/synthesis, not a predictive theory; where it overlaps Penrose’s CCC (~80–90% of the spine) the divergences are stated explicitly. (4) The single declared framework-level point of failure is **fork 5** (unbounded L0); every result that rides on it says so.

1. The state vector and the cycle map

One island (zone) is tracked by

$$x = (M_{L1}, S, V, \{\text{per-round data: } A_i, \eta, P(k), \hat{n}\}), \quad x_{N+1} = \Phi(x_N) = (\Lambda_{\text{live}} \circ B' \circ \Lambda_0)(x_N).$$

The five **bridge variables** that carry the whole loop: the scale factor/Hubble rate (a, H) (the clock, set by the A-field); the fluctuation field $\sigma(M)$ (structure); black-hole mass M (the ladder); entropy S (the only cross-aeon invariant); and the field value A (re-initialized each round). Figure 1 is the complete wiring.

How the parts map onto the cycle. Part I is the live-phase matter sector (Λ_{live} , the ladder L4→L1); Part II is the live-phase clock (the A-field) and its verification record; Part III is the interlude Λ_0 (the corrected dust background) and the closure ledger; Part IV is the transition operator B' (export–partition); Part V is the result of composing them (Φ , the fixed band and the Tolman ledger); Part VI is the verification and empirical confrontation archive; Part VII the measured-input stress audit and the V_{min} Decision Record; Part VIII the single open ledger; Part IX the master attribution. [\[Design\]](#)




Figure 1: The master loop. Boxes are owned equations (colour = phase); red labels are the LMU coupling variables. The architecture — the choice of arrows — is the only LMU claim in this figure. [Design]

Part I

The matter ladder — L4 to L1 as a dynamical system

1. Overview — what this module is, and how to read it

The body of the framework (the main V3.4 document) argues in words that ordinary matter climbs a two-way ladder of black holes, L4→L1, ranked by an object’s *position in the eating process* rather than by its mass. Module L turned that picture into *equations of motion*. **This module does three things with those equations and then explains what the results mean:**

1. **Integrates** the equations forward in cosmic time for four different environments, and asks whether one set of equations reproduces the four observed kinds of object (§4, Fig. 1).
2. **Sweeps** the free coefficients to ask whether the structural results are real physics or artifacts of the particular numbers chosen (§6, Fig. 2).
3. **Scales up** from one hole to a whole population, via the coagulation (Smoluchowski) equation, to ask what mass function the ladder predicts (§7, Fig. 3).

How the model is used, in one sentence. It is a *forward map from environment to fate*: hand it the conditions a hole is born into — the depth of its host (σ), how much gas it is fed (f_{Edd} , $\dot{M}_{\text{cool}}$), and how many companions it can eat (n_{gal}) — and it returns where that hole ends up: its final mass, its spin, the rung it occupies, and whether it is ejected or quenched on the way. One does not tell it the answer; one tells it the environment and reads off the outcome.

What to keep in mind while reading (stated once, up front). These equations are standard galaxy-evolution physics — accretion, dynamical friction, spin evolution, feedback — assembled in a particular way. **Nothing here distinguishes the framework from Λ CDM**, and the absolute masses are calibration, not predictions. What is genuine is *structure*: which correlations fall out of the dynamics without being inserted, and whether they survive when the numbers are varied. The honest accounting is gathered in §9. **[Design]** (the assembly) / **[Hypothesis]** (the physics).

2. The equations of motion

The state of one climbing system is Pitarn’s four channels plus the running mass and a merger counter:

$$\left[\underbrace{M_{\bullet}}_{\text{mass}}; \underbrace{a_{*}}_{\text{Spin}}, \underbrace{M_g}_{\text{Gas}}, \underbrace{n_{\text{gal}}}_{\text{Near-Galaxy}}, \underbrace{N}_{\text{merger count}} \right], \quad \text{environment fixed by } (\sigma, \rho_{\text{DM}}). \quad (1)$$

Two reductions from Module L are made explicit: the gas channel is carried as a reservoir mass M_g , and the dark-matter density enters through σ (which sets both the escape speed and the dynamical-friction rate). Throughout, q is a merger mass ratio and $\eta = q/(1+q)^2$.

2.1 Growth — accrete (Gas) plus eat (Near-Galaxy $\times$ DM)

$$\boxed{\dot{M}_{\bullet} = \dot{M}_{\bullet}^{\text{acc}} + \dot{M}_{\bullet}^{\text{mrg}}} \quad (2)$$

$$\dot{M}_\bullet^{\text{acc}} = f_{\text{Edd}} \underbrace{\frac{M_g}{M_g + M_{g, \frac{1}{2}}}}_{\text{supply throttle}} \underbrace{\mathcal{C}(M_\bullet, \sigma)}_{\text{feedback cap}} \frac{M_\bullet}{\tau_{\text{Sal}}}. \quad (3)$$

$$\mathcal{C}(M_\bullet, \sigma) = \begin{cases} 1, & f_{\text{Edd}} > 1 \text{ (super-Eddington, photon-trapped)} \\ \max(0, 1 - (M_\bullet/M_\sigma)^\beta), & f_{\text{Edd}} \leq 1 \text{ (King regulation)} \end{cases} \quad (4)$$

Cold dense gas feeds the hole at the Salpeter e -fold $\tau_{\text{Sal}} \approx 45$ Myr; a thinning reservoir starves it (the throttle is the smaller of the Bondi and Eddington rates). The cap $\mathcal{C}$ is the King (2003) self-regulation: *sub-Eddington* growth shuts off as $M_\bullet \rightarrow M_\sigma$ (McConnell–Ma 2013). **The one structural assumption** is the upper branch — *super-Eddington* accretion in a dense cocoon is radiatively trapped, so it evades the M_σ wind and can overshoot (the only way an isolated hole becomes over-massive, §5). **[Fact]** (Bondi, Salpeter, King) / **[Hypothesis]** (super-Edd evades feedback; Inayoshi, Haiman & Ostriker 2016).

$$\dot{M}_\bullet^{\text{mrg}} = \Gamma_{\text{mrg}} q_{\text{typ}} M_\bullet, \quad \Gamma_{\text{mrg}} = K_{\text{mrg}} n_{\text{gal}} \left(\frac{\sigma}{\sigma_0} \right)^{-1} [\text{Chandrasekhar } \rho_{\text{DM}}/\sigma^3, \text{ folded}]. \quad (5)$$

Eating needs a companion (n_{gal}) and a sinking rate set by dynamical friction. **[Fact]** (drag) / **[Hypothesis]** (read as the L3→L2 climb).

2.2 Spin — up by coherent Gas, down by Near-Galaxy mergers

$$\dot{a}_* = + \underbrace{C_{\text{acc}} \frac{\dot{M}_\bullet^{\text{acc}}}{M_\bullet} (a_{\text{eq}} - a_*)}_{\text{coherent accretion spins up}} - \underbrace{C_{\text{mrg}} \frac{\Gamma_{\text{mrg}}}{1+N} a_*}_{\text{random mergers spin down}} \quad (6)$$

Prograde accretion drives $a_* \rightarrow a_{\text{eq}} = 0.998$ (Thorne 1974); many misaligned mergers random-walk it down as $a_* \approx a_0/\sqrt{N}$, whose differential form fixes $C_{\text{mrg}} = \frac{1}{2}$ (*not* free; Hughes & Blandford 2003). This is the *merger-phase* spin-down that protects the survivor at the gate (§5); it is distinct from the terminal Page spin-down at the very end.

2.3 Gas, environment, and the two off-ramps

$$\dot{M}_g = \dot{M}_{\text{cool}}(t) - \dot{M}_\bullet^{\text{acc}}, \quad \dot{n}_{\text{gal}} = -\Gamma_{\text{mrg}}, \quad \dot{N} = +\Gamma_{\text{mrg}}, \quad (7)$$

with cooling inflow set by the halo and the companion supply $n_{\text{gal}} \rightarrow 0$ as the centre eats its surroundings — which is what turns an L2 into the lone L1. The escape speed is the DM-set depth, $v_{\text{esc}} = \xi\sigma$ ($\xi \simeq 2$). The two ways off the climb:

$$\textbf{fall:} \quad \text{eject} \iff v_{\text{recoil}}(a_*, q, \theta) > v_{\text{esc}}(\sigma), \quad v_{\text{recoil}} = \sqrt{v_m^2 + v_s^2}, \quad (8)$$

$$v_m = A_{\text{kick}} \eta^2 \frac{1-q}{1+q}, \quad v_s = H_{\text{kick}} \eta^2 (a_2 - q a_1) \cos \theta,$$

$$\textbf{die:} \quad \text{quench} \iff M_g \rightarrow 0 \Rightarrow \dot{M}_\bullet^{\text{acc}} \rightarrow 0 \text{ (starvation; Bullock et al. 2000)}, \quad (9)$$

with $A_{\text{kick}} = 1.2 \times 10^4$, $H_{\text{kick}} = 6.9 \times 10^3 \text{ km s}^{-1}$ (González 2007; Campanelli 2007); the non-spinning recoil peaks at $\sim 175 \text{ km s}^{-1}$, superkicks of $2000\text{--}5000 \text{ km s}^{-1}$ need high in-plane spin. **[Fact]**

3. Coefficients: what is fixed by physics, what is calibrated

Unit declarations (2026-06-12 audit). K_{mrg} carries Gyr^{-1} (previously unitless in the table; the unit lived only in the scripts). **Flag:** the demo script `forward_map_demo.py` implements $\Gamma_{\text{mrg}} \propto$

$(\sigma/\sigma_0)^{+1}$ with $K = 0.08 \text{ Gyr}^{-1}$ — the *opposite* σ -power to eq. (5); the two laws coincide only at $\sigma \simeq 474 \text{ km s}^{-1}$. Eq. (5) (Chandrasekhar-paced, decreasing in σ) is authoritative; a demo rerun on the corrected law is queued, and until then the four-galaxy classification figures carry this flag. The Smoluchowski kernel normalization likewise carries $[\text{Gyr}^{-1} \text{ per unit comoving number density}]$; units previously lived only in code. [\[Auditor\]](#)

Physics-fixed (not free)	Calibrated-free (5)	Per-archetype inputs
$\tau_{\text{Sal}} = 45 \text{ Myr}$	$C_{\text{acc}} = 1.5$ (spin-up rate)	σ (host depth)
$a_{\text{eq}} = 0.998$ (Thorne)	$K_{\text{mrg}} = 0.45 \text{ Gyr}^{-1}$ (merger norm.)	$f_{\text{Edd}}(t)$ (Eddington ratio)
$C_{\text{mrg}} = \frac{1}{2}$ (from $a \propto N^{-1/2}$)	$\beta = 2$ (feedback sharpness)	$M_{\text{cool}}(t)$ (gas supply)
$\xi = 2$ ($v_{\text{esc}} = \xi\sigma$)	$M_{g, \frac{1}{2}} = 10^7 M_{\odot}$	$M_g(0)$ (initial reservoir)
A, H recoil; M_{σ} (McConnell–Ma)	$q_{\text{typ}} = 0.25$	$n_{\text{gal}}(0)$ (companions)

The decisive choice: σ and n_{gal} are set *independently* — a hole can sit in a deep well ($\sigma = 300$) with *zero* companions (the relic). Tying environment to mass was a logged error; keeping them independent is exactly what lets the ranking be by eating-position, not size. [\[Design\]](#)

4. Computation 1 — four environments, four fates

The same equations (3)–(7) are integrated from a $10^4 M_{\odot}$ seed to 13.5 Gyr with four input sets (field, cluster, isolated-massive, dwarf).

LMU V3.4 -- Ladder dynamical system: one set of EoM (Module L), four environments, four fates




Figure 1: (a) The climb $M_{\bullet}(t)$: one set of equations, four environments, four fates; stars are observed anchors (Sgr A*, M87, NGC 1277, dwarf IMBHs; the NGC 1277 star is drawn at the superseded 2012 anchor — the corrected Walsh 2016 mass is $4.9 \times 10^9 M_{\odot}$, see table footnote). (b) State space a_* vs $M_{\bullet}$: at fixed $\sigma = 300$ the cluster (merger-grown) lands at $a_* \approx 0.55$ while the relic (accretion-grown) freezes at $a_* \approx 1.0$.

archetype	$M_{\bullet}$ (model)	anchor	a_*	dex/ M_{σ} (obs)	N_{mrg}	f_{eject}
FIELD $\rightarrow$ L3 (Sgr A*)	4.6×10^6	4.3×10^6	0.84	+0.05 (+0.01)	0.6	41%
CLUSTER $\rightarrow$ L2 (M87)	6.6×10^9	6.5×10^9	0.55	+0.50 (+0.50)	5.9	0%
ISOLATED-MASS $\rightarrow$ relic	2.0×10^{10}	4.9×10^9 [†]	1.00	+0.99 (+0.24) [†]	0.0	0%
DWARF $\rightarrow$ stuck-L4	4.7×10^4	$\sim 2 \times 10^5$	0.86	+0.01	0.1	83%

[†] Anchor updated (V3.5): the NGC 1277 anchor is the Walsh et al. (2016) adaptive-optics dynamical mass ($4.9 \pm 1.6 \times 10^9 M_{\odot}$ (confirmed by Krajnović et al. 2018), superseding the van den Bosch et al. (2012) value in use

when this run was calibrated (now retired; Part V §4). The archetype’s 2.0×10^{10} output therefore *overshoots* the corrected anchor by ~ 0.6 dex, and NGC 1277 is *lighter* than M87 — the relic claim rides the $n = 6$ sample offset $+0.22 \pm 0.09$ dex (Part VI), not this single anchor; the extreme-overmass flagship is TON 618. The parenthesised observed offset ($+0.24$ dex) is the corrected anchor’s M - σ residual. [\[Auditor\]](#)

What it means. The four rungs of the ladder are not four separate stories that each need their own explanation; they are *one mechanism* — the same five equations — run from four starting environments. A field galaxy fed a steady trickle of gas regulates itself onto the M_σ relation and stops (L3). Put the same seed in a company-rich cluster and mergers carry it past M_σ into a cluster-dominant hole (L2). Give it a dense early gas cocoon but no companions and it overshoots by super-Eddington accretion, then freezes — an over-massive but un-climbed relic (L3-relic). Starve it in a shallow dwarf and it stalls near its seed (stuck-L4). The model’s job is done when the *environment alone* selects the fate; that it lands on the four real anchors is the consistency check that the assembled physics is at least self-consistent.

5. What emerges — structural results, not tuned

The spin coefficients are the *same* for all four runs; the following were not fitted per object.

- **Spin remembers how the hole grew.** At fixed $\sigma = 300$, cluster and relic differ by $\Delta a_* = +0.45$ purely because one grew by mergers (spun down) and the other by accretion (spun up). Spin is therefore a *fossil of assembly history* that is independent of mass — the framework’s one genuinely non-circular structural expectation (its carry-through/C5 signature). [\[Hypothesis\]](#)
- **The survivor cannot be thrown out.** The cluster candidate ($a_* = 0.55$, $v_{\text{esc}} = 600$ km/s) has $f_{\text{eject}} = 0\%$ — protected on *both* coordinates at once, spun down and deep-welled — while the spun-up shallow-welled bottom has $f_{\text{eject}} = 41\text{--}83\%$ per merger. Ejection recycles only the light bottom, never the winner. This supplies the mechanism the survivor argument otherwise only asserts. [\[Hypothesis\]](#)
- **Over-massiveness is forced, not chosen.** An isolated hole ($n_{\text{gal}} = 0$) has $\dot{M}_\bullet^{\text{mrg}} = 0$, so it *cannot* be built over-massive by eating; the only route left is super-Eddington accretion. The relic’s $+0.99$ dex offset is thus forced to be an accretion-overshoot signature — which makes it a testable prediction (§10), not a fudge.

6. Computation 2 — are the structural results real? (robustness)

A worry about §5 is that those three results might be artifacts of the particular coefficients chosen. To test this, all five free coefficients were varied together over factor-2–3 ranges (Monte Carlo, $N = 400$, stiff Radau integrator), re-running cluster, relic, and field for each set.

What it means. The decoupling and the protection are not tuned coincidences — they survive the entire free-parameter space, because they are forced by the *topology* of the two situations, not by the numbers. An isolated hole has no companions, hence no spin-down term in Eq. (6), so it must reach $a_* \rightarrow a_{\text{eq}}$; a merger-rich hole always carries the spin-down term, so it must end lower. The *sign* of Δa_* is therefore guaranteed; only its *size* (how far the cluster spins down) depends on the coefficients, and that stays observably large (median 0.45). Likewise the survivor sits in a deep well and is spun down, so its recoil never reaches escape, whatever the coefficients. In short: §5’s structural claims are promoted from “happens at the fiducial point” to “happens generically.” The absolute masses, by contrast, do move with the coefficients — they remain calibration.




Figure 2: (a) The spin decoupling Δa_* across 400 random coefficient sets: it is positive in 100% of them (median 0.45, minimum 0.30), never vanishing. (b) Ejection probability: the cluster survivor stays at $f_{\text{eject}} = 0\%$ in 100% of the space, while the field bottom remains ejectable (median 40%).

7. Computation 3 — from one hole to a population

A single trajectory describes one hole. The whole ladder is a *population*, governed by the coagulation (Smoluchowski 1916) equation that Module L opened,

$$\begin{aligned} \partial_t n(M, t) = & \underbrace{\frac{1}{2} \int_0^M K n(M') n(M-M') dM'}_{\text{mergers (climb-by-eating)}} - n(M) \int_0^\infty K n(M') dM' \\ & - \underbrace{\partial_M [\dot{M}_{\text{acc}}(M) n]}_{\text{gas climb}} - \underbrace{\Gamma_{\text{ej}}(M) n}_{\text{recoil sink}} + \underbrace{S(M, t)}_{\text{seeds}} \end{aligned} \quad (10)$$

with the friction-motivated kernel $K \propto (M_i + M_j)$, gas advection up to the M_σ ceiling, the recoil sink acting only on low-mass merger products (the §5.2 protection, applied at the population level), and a declining seed source. The scheme conserves mass to 0.1% (a coagulation-solver check), and the recoil branching, not a continuous decay, is what makes it physical.




Figure 3: (a) Build-up of the black-hole mass function over cosmic time: gas accretion lifts seeds to the L3 scale, mergers build the high-mass tail, recoil recycles the bottom. (b) The end-state mass function is steeply declining — many low-mass holes, very few high-mass ones — so the very top of the ladder is reached by almost nothing.

What it means. Read as a population, the ladder predicts exactly the shape one sees: a black-

hole mass function dominated by ordinary galaxy-centre (L3) holes, falling steeply toward the cluster-dominant (L2) scale, with the survivor (L1) end essentially empty. **The rarity of the top is structural** — it is the natural outcome of a cascade in which most objects settle or are recycled long before reaching the summit. This is the population-level reason the main text gives for why L1 has no observed example: not a gap in the data, but a consequence of the dynamics. The recoil sink shows its hand at the bottom, where low-mass spun-up merger products are kicked out and recycled rather than retained.

7.1 Refinement — the high-mass abundance from first principles

The single-population run fixes the mass-function *shape* but calibrates the merger efficiency, so the *number* of L2-scale holes is not yet first-principles. The fix is to resolve the population by environment: most holes live in shallow field hosts (low σ , few companions $\rightarrow$ L3 on $M-\sigma$), a minority in deep cluster hosts (high σ , many companions $\rightarrow$ L2), and these environments occur with the abundance set by the observed velocity-dispersion function (VDF; Sheth et al. 2003). Running the dynamics across a σ grid and weighting by the VDF gives the black-hole mass function as a convolution, $\text{BHMF} = \text{VDF} \circ M_{\bullet}(\sigma)$.




Figure 4: (a) The BH mass function from the environment-weighted dynamics: declining and observed-like, with the L2 end rare *because deep hosts are rare* ($n_{\text{L3}}/n_{\text{L2}} \approx 90$, set by the VDF, not tuned). (b) How environments map to masses: the dynamics produces $M-\sigma$ at low σ (field) and rises above it with merger spin-down at high σ (cluster), plus the over-massive isolated-relic branch.

What it means. The L2 abundance is now first-principles *given the VDF*: the top of the ladder is rare because the deep environments that build it are rare, a number that comes from the dispersion function rather than from a tuned merger rate. This remains the standard $\text{BHMF} = \text{VDF} \times M-\sigma$ construction — non-discriminating — the framework’s additions being the over-massive relic branch and the spin structure; the VDF parameters and the environment model are approximate, and the low-mass (dwarf-IMBH) end is over-produced (observationally uncertain anyway). [\[Hypothesis\]](#)

8. What this model is for — and what it cannot do

What it is for. (i) A *translator*: it converts the verbal ladder into a quantitative forward map, so the picture can be checked for internal consistency rather than only asserted. (ii) A *fingerprinting tool*: because spin records assembly history (§5), the model says that a direct spin measurement reads off whether a hole was merger-built or accretion-built — a use with a concrete observational

target (§10). (iii) A *population predictor of shape*: it says the mass function must be top-sparse, which is why no L1 is seen.

What it cannot do (and does not claim).

- **It does not beat Λ CDM.** Every equation is standard galaxy-evolution physics; the model is a consistency statement of the framework’s *organising idea* (eating-position ranking, survivor, recycling), not a new prediction that discriminates between cosmologies.
- **It does not predict absolute masses.** Five free coefficients plus per-archetype inputs mean matching four anchors is a consistency demonstration, not a fit-test. The weight is on the un-tuned structure of §5–6, which survives the sweep.
- **The M_σ landing of the field is circular** — the King mechanism is built to yield M_σ , so that is calibration, not a prediction. [Open]
- **The population’s high-mass *abundance* is not yet first-principles.** The single-population run gets the mass-function *shape* (declining, top-sparse) but the *number* of L2-scale holes needs the spread of environments (σ , n_{gal} distributions) folded in — the next refinement. That pure light-seed mergers struggle to reach the top is itself the seeding bottleneck the main text acknowledges, resolved there by accretion and carry-through seeds.

9. Falsifiers

Mandatory flag (V3.5; travels with every reuse of the ladder): the cluster spin-down expectation ($a_\star \sim 0.55$) is in tension with the measured M87 spin $a_\star \gtrsim 0.8$ (Drew et al. 2025) — status *leaning-against*, not falsified; the relic-side spin decoupling is unmeasurable in principle (Part VI §21). **V3.7 inversion:** the spin walk $a = a_0/\sqrt{1+N}$ (Hughes & Blandford 2003) turns $a_\star \geq 0.8$ into $N_{\text{misaligned}} \leq 0.56\text{--}2.26$ (lower: pure walk from $a_0 = 0.998$; upper: walk allowing accretion re-spin, lift $\times 1.45$) against the modeled 5.9 — tension factor 2.6–10.6, i.e. an aligned-merger fraction $\geq 62\%$ where the model assumed isotropy.

- **Relic spin.** The model forces isolated over-massive relics (NGC 1277, Mrk 1216) to spin *high* (accretion-frozen) while merger-built cluster BCGs spin *low*. A local relic measured to spin low would break the decoupling. Testable by X-ray reflection (accreting relics) and, for the carry-through version, NewAthena + LISA (late 2030s).
- **A clear major merger in a relic** would break the “grew fast, never climbed” reading.
- **Super-Eddington regulation.** If dense-cocoon super-Eddington accretion is shown to drive the same M_σ wind as sub-Eddington, the isolated-relic over-mass mechanism (§5) fails.

10. Attribution

Element	Owner / source
Four state variables; eating-position ranking; assembly	Pitarn ([Design])
Accretion: Bondi rate; Eddington/Salpeter e -fold	Bondi (1952); Salpeter (1964)
Super-Eddington in dense cocoon (evades feedback)	Inayoshi, Haiman & Ostriker (2016)
Mergers: dynamical-friction sinking	Chandrasekhar (1943); White & Rees (1978)
Spin-up to $a_{\text{eq}} = 0.998$ / merger spin-down $a \propto N^{-1/2}$	Thorne (1974); Hughes & Blandford (2003); Berti & Volonteri (2008)
AGN feedback / M_σ self-regulation; M_σ normalisation	Silk & Rees (1998); King (2003); McConnell & Ma (2013)
Escape speed; recoil / superkick	NFW (1996); González (2007); Campanelli (2007)
Starvation quenching	Bullock et al. (2000)
Population / coagulation equation	Smoluchowski (1916); Press & Schechter (1974)
Integration, sweep, population solver, honest-status audit	editor pass ([Auditor])

Module L-3 (consolidated) integrates, stress-tests, and scales up the equations of motion of Module L. It is a worldview's math pass, not a peer-reviewed result: every coefficient is classified, the calibration is named, the one load-bearing hypothesis (super-Eddington evades feedback) is flagged, and the structural results (spin decoupling, survivor protection, top-sparse mass function) are shown to survive a full free-parameter sweep. The ladder is the best-anchored part of the framework; this module makes its dynamics explicit, its results interpretable, and its freedoms visible.

Part II

The dark-energy field A — verification record, runs, and the DESI confrontation

What this part verifies. Sections 1–7 are the integrated verification record (formerly the separate companion document *Verification of the LMU Dark-Energy Field A*, June 2026; carried here in full — units $M_{\text{pl}} = 1$, $H_0 = 1$, $\Omega_{m0} = 0.31$, $\Omega_{\text{DE}0} = 0.69$). They verify, by direct numerical integration, the dark-energy equation of state $w(a)$ of the scalar field A — the pair (w_0, w_a) — and test it against the value measured by DESI; they also establish the condition $V_{\text{min}} = 0 \Rightarrow H \rightarrow 0 \Rightarrow$ asymptotic Minkowski that defines the Fork-Y endgame, and cross-validate the integration engine against published thawing-quintessence work. **Headline:** canonical thawing gives $w \geq -1$ always; it cannot reach the DESI central $(w_0, w_a) = (-0.84, -0.62)$, which requires past phantom crossing — but it is *not excluded*: the V3.8 chain pull pins the joint locus distance at 2.69σ (chain-Gaussian, $\rho = -0.894$) and 2.40σ (posterior-mass; Λ CDM itself sits at 2.83σ in the same convention — DESI’s quoted 2.8σ ; marginal w_0 within 1.3σ). **What this does not establish:** a derivation of $V_{\text{min}} = 0$ (the Weinberg wall, [Open]), or the $L0$ -reservoir / entropy-dilution structure (an LMU interpretive layer, [Hypothesis]). Section 8 adds this session’s independent Run 2 (A_i fixed, m shot); §9–11 carry the V3.4 archive’s own PNGB cross-audit, honest status, and attribution.

1. Equations — the three-party balance

A canonical scalar minimally coupled to gravity, homogeneous, in a flat FRW background:

$$\text{field (Klein–Gordon): } \ddot{A} + 3H\dot{A} + \frac{dV}{dA} = 0, \quad V(A) = \frac{1}{2}m^2 A^2 \quad (1)$$

$$\text{energy bookkeeping: } \frac{d}{dt} \left(\underbrace{\frac{1}{2}\dot{A}^2}_K + V \right) = -3H\dot{A}^2 \quad (2)$$

$$\text{equation of state: } \rho_A = \frac{1}{2}\dot{A}^2 + V, \quad p_A = \frac{1}{2}\dot{A}^2 - V, \quad w_A = \frac{\frac{1}{2}\dot{A}^2 - V}{\frac{1}{2}\dot{A}^2 + V} \quad (3)$$

$$\text{Friedmann: } 3H^2 = \frac{\rho_{m0}}{a^3} + \rho_A, \quad \rho_{m0} = 3\Omega_{m0} = 0.93 \quad (4)$$

$$\text{DESI fitting form (CPL): } w(a) = w_0 + w_a(1 - a) \quad (5)$$

The three “parties” of the field equation are inertia $\ddot{A}$, Hubble friction $3H\dot{A}$ (the dissipative sink), and the restoring force dV/dA . All equations are standard [Fact] (Klein–Gordon–FRW; thawing class: Scherrer & Sen 2008; CPL: Chevallier–Polarski 2001, Linder 2003).

2. Result 1 — $w(a)$ versus DESI (the m -scan)

For each field mass m , the initial field value is shot so that $\Omega_{\text{DE}}(a=1) = 0.69$; then w_0 and $w_a = -dw/da|_{a=1}$ are read off. [Fact]

m/H_0	w_0	w_a	$w(a \rightarrow 0)$ via CPL	note
0.3	-0.987	-0.019	-1.006	near- Λ
0.5	-0.964	-0.054	-1.018	
0.8	-0.908	-0.144	-1.052	
1.0	-0.857	-0.233	-1.090	matches DESI w_0
1.5	-0.681	-0.588	-1.269	matches DESI w_a
2.0	-0.445	-1.176	-1.621	
DESI	-0.84	-0.62	-1.46	phantom past

Verdict. At $m \approx 1$ the model matches DESI's w_0 but predicts $w_a \approx -0.23$, not -0.62 ; matching w_a instead forces $w_0 \approx -0.68$. *One or the other, not both.* The structural reason: the DESI central implies $w_0 + w_a = -1.46$, i.e. $w < -1$ in the past (phantom crossing), whereas a canonical scalar obeys $w = (K - V)/(K + V) \geq -1$ identically (verified: $\min w = -1.0000$). **[Theorem]** (bound) Hence canonical thawing is *barred from the DESI central region by principle* — but lies on the same $w \geq -1$ side that Fork Y wants, and sits at a joint distance of 2.69σ from the DESI central on the released chain covariance ($\rho = -0.894$), 2.40σ in the posterior-mass convention — *closer to the data than Λ CDM itself* ($3.04\sigma/2.83\sigma$ in the same two metrics; marginal w_0 alone within 1.3σ) — the V3.8 chain pull replaces the V3.7 covariance grid here.




Figure 1: Left: $w(a)$ for thawing (blue) stays ≥ -1 ; the DESI CPL line crosses below -1 in the past (purple, forbidden for a canonical field). Right: the thawing locus in the (w_0, w_a) plane never reaches the DESI point, which sits in the past-phantom region $w_0 + w_a < -1$.

3. Result 2 — the $V \rightarrow 0$ endgame (Fork Y's actual goal)

Fork Y does not seek phantom crossing; it seeks $A \rightarrow \approx 0$ ($V \rightarrow 0$, Minkowski endgame). For $V = \frac{1}{2}m^2 A^2$ (minimum at $A = 0$), once $H < m$ the field is underdamped and **oscillates through** $A = 0$: A flips sign, V touches 0 at each crossing, and w swings between -1 and $+1$ with $\langle w \rangle \rightarrow 0$ — i.e. it behaves like *matter* (virial of a quadratic: $\langle K \rangle = \langle V \rangle$). **[Fact]** (Turner 1983)

Integrated to $a = 20.6$: the oscillation envelope decays $0.34 \rightarrow 0.004$; $\rho_{\text{DE}} : 2.07 \rightarrow 2.6 \times 10^{-4}$; $\rho_m \rightarrow 10^{-4}$; $H : 1.0 \rightarrow 0.011$. Power laws $\rho \propto a^{-3.00}$, $H \propto a^{-1.50}$, $\langle w \rangle = 0.003$ — exact matter-like dilution carrying $A \rightarrow \approx 0$, $H \rightarrow 0$. **[Fact]**

Decisive point. $H \rightarrow 0$ (not a constant) precisely because $V_{\min} = 0$. Were $V_{\min} > 0$, $\rho_{\text{DE}} \rightarrow V_{\min}$ and $H \rightarrow \text{const}$ = eternal de Sitter; $V_{\min} < 0$ gives recollapse (crunch). Thus $V_{\min} = 0$ is exactly the choice that yields the Minkowski endgame — and *why* $V_{\min} = 0$ is the Weinberg (1989) cosmological-constant wall, **[Open]** for the entire field.




Figure 2: Massive scalar rolling then oscillating: A alternates sign, V touches 0 at each $A = 0$ crossing, w swings $-1 \leftrightarrow +1$, and the running average $\langle w \rangle \rightarrow 0$ (matter-like).

4. Result 3 — transient acceleration (“the mountain”)

Integrating A into the future, the cosmic acceleration $\ddot{a}/a$ **rises, peaks, and falls** — a transient “mountain,” unlike Λ CDM’s plateau. Verified values of $\ddot{a}/a$: $a=0.3 : -4.90$; $0.7 : +0.24$; $1.0 : +0.39$; $2.0 : +0.16$; $4.0 : -0.09$. Acceleration peaks at $a = 0.99$ ($\approx$ today) and ends at $a \approx 2.91$, after which the universe decelerates toward $H \rightarrow 0$. **[Fact]**

Honest caveat. The hump is **broad and shallow** (a $\sim 5\%$ variation in $\ddot{a}/a$ across $a = 0.85$ – 1.2), not a sharp peak; and “we sit at the summit” is a consequence of choosing m to fit w_0 , so the why-now coincidence is relocated into m , not removed.

The reason acceleration ends is that w_{tot} rises through $-1/3$ (equivalently $\ddot{a} = 0$) as A thaws. The CPL form must *not* be extrapolated past $a = 1$: with our (w_0, w_a) it gives $w(a=5) = +0.08$, $w(a=10) = +1.24$ (unphysical), whereas the real field stays bounded in $[-1, +1]$.

This transient-acceleration behaviour is published: **de Souza, Rodrigues & Alcaniz (2025), Phys. Rev. D, arXiv:2504.16337**, who find thawing competitive with Λ CDM ($\Delta \ln B = -0.33$ vs CPL’s -1.87) and predict acceleration ending at $z \approx -0.15$. **[Fact]**




Figure 3: Endstate: $A \rightarrow \approx 0$, all densities $\rightarrow 0$ as a^{-3} , and $H \rightarrow 0$ as $a^{-3/2}$ — asymptotic Minkowski.




Figure 4: Acceleration $\ddot{a}/a$: A-thawing (blue) rises, peaks near today, then falls; Λ CDM (green dashed) rises to a plateau and never falls. Red dot = today.

5. Result 4 — cross-check across potentials (no dialing)

Test of robustness: pick *raw* shape parameters (not chosen to hit any w_0), fix only the overall scale to the *measured* $\Omega_{\text{DE}} = 0.69$, and see what comes out.

potential (raw shape, no dialing)	w_0 (output)	w_a	accel-ends a
Quadratic $\frac{1}{2}m^2 A^2$, $m=1$, $A_i=2$	-0.806	-0.328	2.09
Quadratic, $m=1$, $A_i=4$	-0.961	-0.059	—
Exponential $e^{-\lambda A}$, $\lambda=1$	-0.847	-0.191	—
Exponential, $\lambda=0.5$	-0.962	-0.053	—
Exponential, $\lambda=0.3$	-0.987	-0.019	—
PNGB $1 - \cos(A/f)$, $f=3$, $A_i=2$	-0.820	-0.309	2.18

What is invariant (output, not dialed): every potential lands in $w_0 \in [-0.81, -0.99]$ on the $w > -1$ side, with $w_a < 0$ (thawing) and $w \geq -1$ throughout. A clear relation emerges *by itself* — w_0 near $-1 \Rightarrow$ small $|w_a|$; $w_0 \approx -0.8 \Rightarrow$ larger $|w_a|$ — i.e. all potentials fall on one locus in the (w_0, w_a) plane. This is the signature of a canonical scalar; it appears without fitting. **[Fact]**

What is model-discriminating (not rounding): the precise (w_0, w_a) and the end-of-acceleration time differ with potential shape by factors of ~ 2 – 3 (a quadratic with a minimum oscillates to




Figure 5: Future: the real integrated $w_A(a)$ (blue) stays in $[-1, 1]$ and oscillates; the CPL linear extrapolation (red dashed) diverges and is unphysical for $a > 1$. Lower panel: acceleration ends, universe decelerates (“slowing”).

matter and ends acceleration; a runaway exponential keeps mild negative pressure longer). These differences are real physics, and they are what observations use to tell potentials apart.

6. Layer separation (anti-drift)

statement	status
$w = (K - V)/(K + V) \geq -1$; canonical scalar never phantom	[Theorem]/[Fact]
Thawing $w(a)$, transient acceleration, $A \rightarrow \approx 0$, $H \rightarrow 0$ (asymptotic Minkowski)	[Fact]
Oscillating quadratic scalar $\Rightarrow \langle w \rangle = 0$ (matter-like), $\rho \propto a^{-3}$	[Fact]
Global energy not conserved in expansion (FRW has no time-translation symmetry)	[Fact]
L_0 has an exact timelike Killing vector / exact conservation	only as $t \rightarrow \infty$; finite-time L_0 is <i>quasi-static</i> ([Fact] asymptotically)
“Island matter merges into the L_0 reservoir”	[Hypothesis] (LMU interpretation; not from the field equation)
“Entropy diluted in an infinite $L_0 \Rightarrow$ second-law-safe”	[Hypothesis] (rests on infinite- L_0 ; departs from Penrose)
“BH by-product is the source of A ”	[Dead-end] (fails energy budget / timing / homogeneity)




Figure 6: Left: $w(a)$ for several potentials (when matched in w_0) nearly coincide and bracket the de Souza EoS; all stay ≥ -1 . Right: all show the same transient-acceleration hump. The universal (qualitative) physics is potential-independent; the quantitative w_a /timing is shape-dependent.

Bottom line: the equations deliver $A \rightarrow \approx 0$, $H \rightarrow 0$ (asymptotic Minkowski). “Exact Minkowski + exact conservation + reservoir banking + entropy dilution” is an interpretive layer the framework adds, not a consequence of the field equation. The two must always be tagged separately. [\[Auditor\]](#)

Why this table is mandatory (V3.5). It is the anti-drift guard in practice: in the session that produced Operator B' it is what caught the auditor’s Λ -import error (Correction 1) — the interlude background had been computed on a constant- Λ de Sitter despite row 2’s verified $H \rightarrow 0$. [\[Auditor\]](#)

7. DESI status and the pre-registered falsifier

DR1→DR2 (6M→14M redshifts): the central (w_0, w_a) drifts slightly toward -1 while error bars shrink faster, so the preference for evolving dark energy *does not diminish*; it stays $\lesssim 3\sigma$, below the 5σ discovery threshold, with SN-calibration systematics unresolved, and some analyses dispute robustness. [\[Fact\]](#) The thawing class is supported only at “weak/inconclusive” Bayesian levels but is *not excluded*; non-phantom models are compatible at 2σ , and the phantom-crossing evidence is the *weaker* part of DESI. **V3.8 quantification (measured covariance):** on the released DR2+Pantheon+ chains $\rho(w_0, w_a) = -0.894$, and the thawing locus’s minimum joint distance is 2.69σ (chain-Gaussian) / 2.40σ (posterior-mass, the convention of DESI’s own 2.8σ Λ CDM exclusion, reproduced as 2.83σ); the V3.7 ρ -grid (2.00 – 2.92σ) is retired to the pin record; “compatible at 2σ ” was the optimistic edge, and the proper test remains a direct non-CPL fit (Dinda & Maartens 2025; de Souza et al. 2025). [\[Auditor\]](#)

Pre-registered falsifier (committed now). Fork Y (canonical A , $w \geq -1$, Minkowski endgame) is falsified *iff* DESI-Y5 (~ 2027) sustains $w(z) < -1$ at high z at $\gtrsim 3\sigma$ *with systematics resolved*. It is *not* falsified by $w \geq -1$ evolving data, nor by a sub- 3σ /systematics-limited phantom hint. The 3σ line is fixed and must not be moved if data later push against the model.

8. Run 2 — A_i fixed, m shot (this session’s cross-validation)

A second, independent shooting strategy was run for the equation document (fresh KG+Friedmann integration, Radau): instead of fixing m and shooting A_i (the m -scan of §2), **Run 2 fixes $A_i = 2.70 M_p$ and shoots m** so that $\Omega_{A,0} = 0.689$. Result: $m = 0.804 H_0 = 1.16 \times 10^{-33}$ eV; $w_0 = -0.91$, $w_a = -0.15$ (matching the thawing relation $w_a \simeq -1.5(1 + w_0) = -0.14$ to 4%); acceleration window $a \in (0.61, 5.5)$; first $A = 0$ crossing at $a = 9.4$ (μ -dependent, “few–20” across the verified band); $\langle w \rangle \rightarrow 0$ and $H \rightarrow 0$ confirmed (Fig. 7). Canonical $w \geq -1$ throughout. **[Fact]**

Cross-validation. Run 2 lands on the m -scan’s $m = 0.8$ row, $(w_0, w_a) = (-0.908, -0.144)$ — two independent shooting strategies (m fixed/ A_i shot vs. A_i fixed/ m shot), same point on the thawing locus. The V3.5 assembly recomputed both fresh (R6, Radau and DOP853): A_i shot at $m = 0.8$ gives $A_i = 2.715 M_p$, $(w_0, w_a) = (-0.9080, -0.1438)$; m shot at $A_i = 2.70 M_p$ gives $m = 0.8030 H_0$, $(-0.9071, -0.1456)$. **[Auditor]**




Figure 7: Run 2 — the A-field through one zone’s life: thaw, transient acceleration, oscillation to dust, $H \rightarrow 0$. Every curve is from the equation document’s own integration (R6). Owners: eqs. (1)–(3), Turner 1983.

9. The PNGB cross-audit against DESI DR2 (the V3.4 archive)

The following two sections are the V3.4 consolidated pass’s own dark-energy module (its Part II §§1–2), carried as the archive record: an independent integration on a different thawing potential (PNGB rather than quadratic), in different variables (e -folds), reaching the same structural verdict. Redundancy here is deliberate — it is the cross-check. [\[Auditor\]](#)

10. The field, and what the data ask of it

The field A is a single canonical scalar with a rolling potential (Ratra–Peebles 1988; Wetterich 1988; Caldwell–Dave–Steinhardt 1998), treated *per zone* rather than as a universal constant. Its job in the framework is twofold and the two are linked: it must (a) *accelerate the zone now* (the only observed fact), and (b) *weaken toward $A \rightarrow 0$* so the zone returns to the timeless reservoir and the loop can close. DESI DR2 sharpened (a) into a quantitative target: the data prefer an *evolving* equation of state,

$$w(a) = w_0 + w_a(1 - a), \quad w_0 = -0.838 \pm 0.055, \quad w_a = -0.62 \pm 0.2 \quad (\text{DESI} + \text{Pantheon+}), \quad (6)$$

i.e. dark energy was *stronger* in the past and is weakening — which is, qualitatively, exactly what requirement (b) predicts. The open joint O2 is whether a *canonical* field (the spine the framework chose) can match this quantitatively, and which way its potential’s minimum sits.

11. Computation — Klein–Gordon vs DESI (O2)

The field obeys the Klein–Gordon equation in an expanding background; in e -folds $N = \ln a$, with $8\pi G = 1$ and H solved self-consistently from the Friedmann constraint,

$$\phi'' + \left(3 + \frac{\dot{H}}{H^2}\right)\phi' + \frac{1}{H^2} \frac{dV}{d\phi} = 0, \quad H^2 = \frac{\rho_m + V(\phi)}{3 - \frac{1}{2}\phi'^2}, \quad w_\phi = \frac{\frac{1}{2}H^2\phi'^2 - V}{\frac{1}{2}H^2\phi'^2 + V}. \quad (7)$$

We integrate a thawing pseudo-Nambu–Goldstone potential $V(\phi) = A_0[1 + \cos(\phi/f)]$ (Frieman et al. 1995) — the canonical thawing case, whose minimum sits at $V = 0$ — from a frozen high-redshift start, tuned so $\Omega_\phi(\text{today}) = 0.69$ with $\Omega_m = 0.31$, and scan the initial displacement ϕ_0 (how far the field thaws by today).

Result. The integration reproduces, from first principles, the tension the main text states heuristically:

- The thawing field reaches $w_0 \approx -0.88$, $w_a \approx -0.17$ at its DESI-closest displacement — it *approaches* the data but misses the central $(w_0, w_a) = (-0.838, -0.62)$: w_a is too shallow, and a canonical field **cannot cross** $w = -1$ (Vikman 2005), which is where DESI’s central law goes. The miss is the $\sim 2\text{--}3\sigma$ gap the main text quotes. *This is a miss of the central value, not an exclusion*: a non-phantom field ($w \geq -1$) remains compatible with DESI at $\sim 2\sigma$, and the phantom-crossing signal is the weakest part of the data — so “cannot reach the central” must not be read as “ruled out as thawing.” The honest claim is *consistent at 2σ* , not best-fit. [\[Hypothesis\]](#)
- **The endgame is the payoff.** The same thawing field rolls to the minimum of its potential at $\phi = \pi f$, where $V = 0$ — a Minkowski, $A \rightarrow 0$ end-state, exactly the reservoir the loop needs. And the dependence runs the right way: the more the field thaws to fit DESI’s evolution, the sooner it reaches $V = 0$. So DESI’s preference for *weakening* dark energy and the framework’s need for the field to *switch off* are the same statement seen from two ends. [\[Hypothesis\]](#)




Figure 8: (a) $w(z)$ for the thawing field at several initial displacements: it thaws upward from $w = -1$, while the DESI central law (dashed) dives *below* -1 at low z — the phantom region a canonical field cannot enter. (b) The thawing locus in (w_0, w_a) : it runs from Λ CDM toward the DESI DR2 ellipse but stays above/right of the central preference — w_a too shallow, and w_0 overshoots before w_a deepens.

What it means for O2. The field that delivers the endgame the framework requires ($V \rightarrow 0$, Minkowski) is the same field that is mildly *disfavoured* by the present w_0 - w_a data (too shallow, no phantom crossing). The competing reading — the $\sim 94\%$ future-AdS preference in the DESI analyses — points instead to $V_{\min} < 0$, a collapsing zone, which is the rival ($f(R)$ /crunch) closure rather than the reservoir. **The sign of $V_{\min}$ is therefore the whole of O2**, and it is exactly what DESI DR3 (and Euclid/Roman) will pin down. The framework is self-consistent on the endgame it needs and is honest that today’s data lean slightly the other way. **[Open]**

External validation and the committed falsifier. The transient-acceleration behaviour of this field — $\ddot{a}/a$ rises, peaks near today, then falls, unlike Λ CDM’s plateau — is independently published: de Souza, Rodrigues & Alcaniz (2025, PRD, arXiv:2504.16337) find thawing competitive with Λ CDM ($\Delta \ln B = -0.33$ vs CPL’s -1.87) and acceleration ending near $z \approx -0.15$. Honest cost: the hump is broad and shallow, and “we sit at the summit” follows from tuning m to fit w_0 — so the why-now coincidence is relocated into m , not removed, leaving the field even with Λ CDM on that count, not ahead. The pre-registered falsifier (fixed now, not to be moved): Fork Y is falsified *iff* DESI-Y5 (~ 2027) sustains $w(z) < -1$ at high z at $\gtrsim 3\sigma$ with systematics resolved; it is *not* falsified by $w \geq -1$ evolving data or a sub- 3σ /systematics-limited phantom hint. The full no-dialing cross-check across six potentials (quadratic, exponential, PNGB), the min $w = -1$ bound, the $V \rightarrow 0 \Rightarrow H \rightarrow 0$ Minkowski endgame, and the figures are recorded in §§1–7 of this part (the verification record, integrated in V3.5 from the formerly separate companion document *Verification of the LMU Dark-Energy Field A*); this section is the archive summary. **[Fact]** / **[Auditor]**

Is the DESI “phantom past” real, or a CPL artefact?

This is the load-bearing question for O2, and it sharpens the falsifier above. Taking every value consistently from the DESI DR2 key paper (Abdul-Karim et al. 2025, arXiv:2503.14738, single source) to avoid cross-sample mixing:

DESI combination	w_0	w_a	Λ CDM exclusion
DR2 + Pantheon+	-0.838 ± 0.055	$-0.62^{+0.22}_{-0.19}$	2.8σ
DR2 + Union3	-0.667 ± 0.088	$-1.09^{+0.31}_{-0.27}$	3.8σ
DR2 + DESY5	-0.752 ± 0.057	$-0.86^{+0.23}_{-0.20}$	$4.2\sigma \rightarrow 3.2\sigma$ (DES reanalysis)
A-thawing ($m=1$)	-0.857	-0.233	—

Every SN combination’s CPL central crosses below -1 in the past (at $z = 0.5$, $w_{\text{CPL}} \approx -1.04$ for all three samples, versus A-thawing’s -0.934 ; at $z = 1$ all near -1.15 to -1.21), so canonical A — pinned at $w \geq -1$ — does not trace the CPL central for $z \gtrsim 0.5$, though it agrees at $z = 0$. **But this crossing is a property of the CPL straight-line ansatz, not a measurement that the field is phantom:** Dinda & Maartens (2025, MNRAS 542 L31; arXiv:2504.15190) and de Souza et al. (2025, PRD; arXiv:2504.16337) show that fitting physical non-phantom thawing models (rather than CPL) gives fits comparable to CPL — equally good on a curved background (Dinda–Maartens, whose stated conclusion is that the crossing is an artefact of a parametrization not based on a physical model) and competitive with ΛCDM ($\Delta \ln B = -0.33$, de Souza et al.) — so the data do not enforce phantom crossing. The dominant systematic is the SN-sample spread (w_0 from -0.84 to -0.67 ; 2.8σ for Pantheon+ up to 4.2σ for DESY5, the latter weakened to 3.2σ by the DES reanalysis), with Pantheon+ the least discrepant. So A-thawing is *disfavoured at the CPL central but not excluded*, and the Y5 falsifier reduces to one sharp question: does the phantom crossing survive a physical (non-CPL) re-analysis with systematics resolved? If yes, canonical A fails; if it is a parametrization artefact, A stays competitive. **[Open]** / **[Auditor]**

12. Honest status

- The quintessence integration is standard scalar-field cosmology; it does not discriminate the framework from ΛCDM or from other thawing-quintessence models. Its value is to confirm O2 from a real integration and to show the endgame ($V \rightarrow 0$) is realised by the DESI-approaching field.
- The PNGB choice is representative of canonical thawing; a freezing or hilltop potential would shift (w_0, w_a) and, for a hilltop, could overshoot to $V < 0$ (the crunch branch) — which is the O2 fork, not a detail.
- The z_{aeon} result is an impossibility argument, not a calculation; it rests on the conformal invariance of massless-radiation entropy, which is secure, and on 3.4’s scoping of the boundary to the massless channel.
- Everything tied to A (roll rate, the sign of V_{min}) remains hostage to DR3 and to the cosmological-constant fine-tuning problem (Weinberg 1989), which this module does not address. **[Open]**

Merged bottom line (record + archive). The equations deliver $A \rightarrow \approx 0$, $H \rightarrow 0$; the reservoir reading of that endgame is the framework’s interpretive layer; the field’s quantitative fate is left, honestly, to DESI (Y5 falsifier above). **[Auditor]**

13. Attribution

Element	Owner / source
Per-zone rolling field; “ $V \rightarrow 0$ closes the loop” framing	Pitarn ([Design])
Quintessence (rolling scalar dark energy)	Ratra & Peebles (1988); Wetterich (1988); Caldwell, Dave & Steinhardt (1998)
Thawing PNGB potential	Frieman, Hill, Stebbins & Waga (1995)
No phantom crossing for a canonical field	Vikman (2005)
DESI DR2 w_0 - w_a constraint (single-source values)	DESI Collaboration / Abdul-Karim et al. (2025, arXiv:2503.14738)
Transient (thawing) acceleration vs DESI; Bayesian comparison	de Souza, Rodrigues & Alcaniz (2025, PRD, arXiv:2504.16337)
Phantom crossing as a CPL artefact; non-CPL thawing fit	Dinda & Maartens (2025, MNRAS 542 L31; arXiv:2504.15190); de Souza et al. (2025; arXiv:2504.16337)
Conformal cyclic cosmology / massless-entropy invariance	Penrose (2010)
Black-hole entropy; radiation-to-BH entropy ratio (4/3)	Bekenstein (1973); Hawking (1975); Zurek (1982)
Klein–Gordon integration; conformal-boundary audit	editor pass ([Auditor])

Module A is the dark-energy half of the V3.4 math pass. It shows, from a real field integration, that the canonical thawing spine approaches but does not yet match DESI (O2, w_0 - w_a too shallow, no phantom crossing) while delivering the $V \rightarrow 0$ endgame the loop requires; and it shows, from the conformal invariance of massless entropy, that the boundary’s z_{aeon} is undeterminable in principle — which explains the old ambiguity and confirms the framework’s “dilute, don’t reset” entropy stance. The field’s fate is left, honestly, to DESI DR3.

Part III

Closure of the loop — the $L0$ ledger

What this module settles. The framework’s loop closes its matter, energy, and entropy budgets through a reservoir state $L0$ (a timeless, $A \approx 0$, *asymptotically* Minkowski background between aeons (at finite time: a quasi-static *dust era*, §2)). The verbal argument (main text §6) is here made quantitative: what enters $L0$ and at what rate, why the energy budget balances exactly (in=out), where “energy conserved” is and is not the same as “the loop can run,” and — the point of the module — a clean separation of the *four* closure worries that are answered by physics from the *one* that remains an explicit, undecidable assumption. The honest end-state: the loop is closed up to a single load-bearing premise (an actually-infinite reservoir), and that premise is the toll every cyclic and eternal-inflation cosmology pays, not a defect peculiar to this one. **[Design]** (the reservoir architecture and ledger) / **[Fact]**, **[Hypothesis]** (the physics terms) / **[Open]** (the one premise).

1. The reservoir and what enters it

Let a single zone (one causally bound island) end its aeon. Three channels route its mass–energy into the shared reservoir $L0$ — but at different stages: one evaporates *in* $L0$ (after the zone has relaxed toward Minkowski — the dust era, §2), one fires *inside* the still-accelerating zone with its radiation entering $L0$ only as the zone relaxes, and one simply drifts in. All three are standard physics, assembled by Pitarn into one ledger.

(1) The survivor evaporates (the slow source). The lone $L1$, having swept its reach and spun down (Page 1976, which reaches $a_* \rightarrow 0$ *before* the end), evaporates by Hawking radiation. For a hole of mass $M_\bullet$ the temperature, luminosity, and lifetime are

$$T_H = \frac{\hbar c^3}{8\pi G M_\bullet k_B}, \quad L = \frac{\hbar c^6}{15360 \pi G^2 M_\bullet^2}, \quad \tau_{\text{evap}} = \frac{5120 \pi G^2 M_\bullet^3}{\hbar c^4} \simeq 2.1 \times 10^{67} \left(\frac{M_\bullet}{M_\odot} \right)^3 \text{ yr.} \quad (1)$$

For a survivor at the canonical band top, $10^{11} M_\odot$: $\tau_{\text{evap}} = 2.1 \times 10^{100}$ yr and $S_{\text{BH}} = 1.05 \times 10^{99} k_B$ (a $10^{12} M_\odot$ *upper illustration* gives $T_H \sim 6 \times 10^{-20}$ K, $\tau_{\text{evap}} \sim 2 \times 10^{103}$ yr): the survivor is not a one-off blast but a **cold source that bleeds its full rest energy $M_\bullet c^2$ into $L0$ over an enormous time**. Because the interlude background is a *decelerating dust era* with no event horizon (§2), this slow bleed is not a loss channel: radiation not yet re-gathered dilutes only power-law ($a^3 \sim 10^{180}$ per interlude) while matter *clusters* on a far faster clock (the race, §2 and §4), and a not-yet-evaporated hole is itself a standing mass that anchors where the next structure can grow. **[Fact]** (Hawking 1975; lifetime as quoted).

(2) Ceiling-limited holes pop (the fast source) — a *within-zone* event. While a zone is still accelerating ($A > 0$, an effective de Sitter phase with Hubble rate H), a black hole cannot exceed the Nariai mass, set by the merging of the black-hole and cosmological horizons in Schwarzschild–de Sitter geometry. The degenerate-horizon condition $9G^2 M^2 \Lambda / c^4 = 1$ (with $\Lambda = 3H^2/c^2$) gives

$$M_{\text{Nariai}} = \frac{c^3}{3\sqrt{3}GH} \approx 1.7 \times 10^{22} M_\odot \quad (H \sim H_0), \quad (2)$$

not the naive “hole the size of the Hubble radius” value $c^3/(2GH)$, which overestimates the coefficient by $\sqrt{27}/2 \approx 2.6$. A hole that reaches this ceiling before it can evaporate decays by the Nariai instability into radiation (Route A). *This happens inside the zone, not in $L0$* : once the zone relaxes to $A \approx 0$ there is no cosmological horizon ($H \rightarrow 0 \Rightarrow M_{\text{Nariai}} \rightarrow \infty$), so there is no Nariai ceiling in the reservoir itself — the pop is a de Sitter-phase event whose radiation enters $L0$ only

as the zone winds down. **[Hypothesis]** (Nariai 1950; instability: Ginsparg & Perry 1983; Bousso & Hawking 1996 — the post-Nariai fate is less established.)

(3) Debris escapes as $A \rightarrow 0$ (the cold source). Mass not bound into the survivor — mid-aeon ejecta, un-captured outskirts (the two ejection routes of the ladder) — is left adrift as the zone relaxes to Minkowski, and drifts into $L0$ carrying its rest mass. This is the *material* that seeds later rounds (it has a genuine past; it is not a vacuum fluctuation). **[Hypothesis]**

Crucially, by unitarity the evaporation leaves *no* remnant: the entire $M_\bullet c^2$ is returned to the exterior as radiation. So a zone’s collapsed mass is **not** lost down a hole — it re-emerges in $L0$.

2. The interlude background — a dust era (Correction 1)

(*New in V3.5, from the Operator-B’ session; the constant- Λ interlude of earlier drafts is retired — Part V §4.*) After oscillation onset (Part II §3) the zone and its surroundings are a **dust universe**:

$$a \propto t^{2/3}, \quad H = \frac{2}{3t} \rightarrow 0, \quad \text{no event horizon} \quad (3)$$

(**[Fact]**, FRW dust). Over one interlude ($\tau = 2.1 \times 10^{100}$ yr at the $10^{11} M_\odot$ band top, eq. (1)) the comoving causal reach is $\sim 8 \times 10^{56} \text{ m} \approx 10^{26}$ island radii; mean densities dilute by $a^3 \sim 10^{180}$ (matter) while structure *grows*:

$$\delta \propto a^p, \quad p = \frac{-1 + \sqrt{1 + 24f}}{4} \quad (f = \text{clustering fraction}); \quad t_{\text{coll}} \simeq t_{\text{osc}} \left(\frac{1.686}{\delta_i} \right)^{3/2} \quad (4)$$

(**[Fact]**: linear growth in a mixed background; Gunn–Gott 1972). And the live-phase Nariai ceiling does not operate here: $M_{\text{Nariai}} = c^3/(3\sqrt{3}GH) \rightarrow \infty$ as $H \rightarrow 0$ — eq. (2) is a de Sitter-window statement only.

$\hookrightarrow$ **Couplings and results.** (i) Pair collapse beats evaporation by $\sim 10^{89}$ for $\delta_i \sim 1$ (Fig. 1; this assembly’s R6 run gives 10^{89-90} at $\delta_i \sim 1$ and still 10^{85} at $\delta_i = 10^{-3}$): survivor–survivor mergers and corpse aggregation are *generic*; the cross-island forced fork (same merger map, $S(2M) = 4S(M)$) is the main channel, not the exotic branch. (ii) The aggregation *rate* is now **computed** (item F, resolved 2026-06-10; Fig. 2): the two branches are *scale-separated* by the field’s Jeans length. Below the comoving $\lambda_J = 1.68 \times 10^{26} \text{ m} = 0.38 R_{\text{obs}}$ at oscillation onset (shrinking $\propto a^{-1/4}$; Hu–Barkana–Gruzinov 2000), the A-dust is pressure-supported: with the Part II late-time $\rho_A/\rho_m = 63$ the clustering fraction is $f = 0.016$, $p = 0.044$, growth $\times \sim 4 \times 10^2$ per interlude — nearest neighbours still merge. On island scales the A-dust is **super-Jeans from onset** ($R_{\text{isl}}/\lambda_J = 2.9 \times 10^3$ at the live $m = 0.803 H_0$, the margin growing as $a^{1/4}$; oscillating scalar as dust, Turner 1983): $p \rightarrow 1$ and the hierarchy runs open. The gate over the field mass closes only at $m \lesssim 0.13 H_0$ — by *horizon exile* (>100 e-folds of late slow-roll inflation, $N \approx A_i^2/4$, Linde 1983), not Jeans pressure — a regime already excluded by the model’s own DESI-thaw identification; island modes re-enter at $a/a_{\text{osc}} = 2.6 \times 10^7$ and any seed $\delta \gtrsim 10^{-52}$ collapses before τ , while the Poisson discreteness floor alone ($\delta \sim 6 \times 10^{-11}$; Afshordi–McDonald–Spergel 2003 analog) exceeds the requirement by $\sim 10^{42}$. **[Fact-theory]** (conditional on $V_{\text{min}} = 0$, item B; seed origin stays open in item D) (iii) New zones are born at aggregation sites (“where the material has gathered”), which restores the author’s gathering picture with a physical mechanism. **[Hypothesis]** riding fork 5.

3. Energy balances exactly — the in=out ledger

Why conservation holds here, and not inside a live zone. General relativity has *no* global energy-conservation law in a generic expanding spacetime — there is no timelike Killing vector, and a photon’s energy redshifts away with no local sink (this is why “energy is not conserved”




Figure 1: The interlude timing race on the corrected background. Everything in the shaded region merges before any flash. The dashed line marks the smooth-A-dust worst case — now scoped to sub-Jeans scales only; island scales cluster (item F, Fig. 2).

in an accelerating zone with $A > 0$). The reservoir is the exception *asymptotically*: $L0$ tends to $A \approx 0$, i.e. Minkowski as $t \rightarrow \infty$, which *does* admit a timelike Killing vector $\xi^\mu = (\partial_t)^\mu$, and hence a conserved total energy

$$E_{L0} = \int_{\Sigma} T_{\mu\nu} \xi^\mu n^\nu d\Sigma = \text{const.} \quad (5)$$

Per the layer-separation table (Part II §6), the exact Killing vector and exact conservation hold *only asymptotically*; the finite-time interlude is a quasi-static dust era (§2) with $H = 2/(3t) \rightarrow 0$, and the conservation statement is its $t \rightarrow \infty$ limit, used as the bookkeeping frame. [Auditor] This is the firm ground under Pitarn’s “in=out”: conservation is asserted *at the reservoir level*, where the symmetry that guarantees it actually exists, not across the conformal boundary and not inside an expanding zone.

The ledger. Summing the three channels, each zone deposits its rest energy $E_{\text{zone}} \simeq \sum_i M_i c^2$ into $L0$ as radiation; the reservoir in turn feeds the assembly of new zones. At the level of the whole reservoir, with $\dot{E}_{L0}^{\text{in}}$ the rate of deposit from dying zones and $\dot{E}_{L0}^{\text{out}}$ the rate drawn into forming ones,

$$\boxed{\dot{E}_{L0}^{\text{in}} = \dot{E}_{L0}^{\text{out}} \iff E_{L0} = \text{const (closed reservoir).}} \quad (6)$$

This is an *accounting identity given closure*, not a theorem proved from below: it holds *if* $L0$ is a closed system (nothing leaks beyond it). “ $L0$ is closed” is the same statement as “ $L0$ is all there is” — i.e. the infinite-reservoir premise of §6. So (6) is exact, but it names the assumption it rests on rather than removing it. [Auditor]

V3.7 boundary scope: (6) is the $L0$ debris channel only. Across the conformal boundary the energy ledger does not close: per child, the pre-rescale flash share is $\sim 3 \times 10^{60}$ J against a child matter content $\sim 2 \times 10^{80}$ J today, so every crossing injects a factor $\Omega \sim 10^{26}$ (pair threshold) to 10^{43-47} (GUT gluing) — permitted by GR (no global conservation off the Killing frame), supplied by nothing, identical to the irreducible z_{aeon} of Part IV §2. The photon-count/entropy closure across the boundary is an *identity* (bookkeeping), not a test. [Auditor]

Ledger principle (V3.9, author): *nothing is lost without a receipt* — every LMU ledger either closes exactly (entropy per comoving patch, $B1'$; local $\nabla_\mu T^{\mu\nu} = 0$), is reset by declaration (the $B3'$




Figure 2: Item F resolved. **Left:** comoving scales over the live interlude — the island bound $R_{\text{isl}} = 1107 R_{\text{obs}}$ sits a factor 2895 above λ_J at onset and the margin grows ($\lambda_J^{\text{com}} \propto a^{-1/4}$); island modes re-enter the horizon at $a/a_{\text{osc}} = 2.6 \times 10^7$ and collapse from the Poisson floor long before τ . **Right:** the gate over the field mass — $\delta_{\text{min}}(m)$ versus the Poisson seed floor; open for $m \gtrsim 0.14 H_0$, closed below $\approx 0.13 H_0$ by horizon exile (a regime outside the DESI-thaw identification). Reproduced by `lm_u_itemF_jeans_gate.py`.

re-init list: baryon number via the Sakharov conditions, DM, A_i , $P(k)$, Ω_k), or carries a named price (the $\times \Omega$ boundary injection; T_{reheat}). No silent channel exists; ruling (a) extends the closed entropy ledger to the past direction (finite aeon depth per tree).

4. Energy is not the same as usable capacity

The ledger (6) closes the *energy* budget. It does *not*, by itself, guarantee the loop can keep running, because heat death was never about energy disappearing — it is about *usable capacity* (the ability to build structure) degrading while energy is conserved. The relevant question is whether radiation deposited cold in $L0$ can be *re-gathered into mass* to seed a new round.

Re-formation needs a region driven back above the relevant production threshold; for e^+e^- this is

$$k_B T \gtrsim 2m_e c^2 \Rightarrow T \gtrsim 1.2 \times 10^{10} \text{ K}, \quad (7)$$

reached only where gravity re-concentrates the diluted radiation to high density (mass leads, radiation follows; main paper §12.4). The loop's *capacity* therefore closes only if the reclamation rate keeps pace with dilution,

$$\Gamma_{\text{reclaim}}(\text{gravity re-concentration}) \gtrsim \Gamma_{\text{dilute}}(\text{drift, redshift in } L0), \quad (8)$$

in the net over infinite time. **This inequality is now quantified** on the corrected dust background (§2): Gunn–Gott collapse, eq. (4), beats the survivor's evaporation clock by $\sim 10^{89}$ – 10^{90} at $\delta_i \sim 1$ and still by $\sim 10^{85}$ at $\delta_i = 10^{-3}$ — generic, not marginal — so (8) holds wherever *any* component clusters. The aggregation-rate gate is now **resolved** (item F): sub-Jeans scales keep the suppressed branch ($f = 0.016$, $p = 0.044$, $\times \sim 4 \times 10^2$ — nearest neighbours still collapse); island scales are super-Jeans from onset and run the hierarchy open ($p \rightarrow 1$) for every $m \gtrsim 0.14 H_0$, the closed branch ($m \lesssim 0.13$, horizon exile) lying outside the live model (§2, Fig. 2). Status: the inequality (8) is **[Hypothesis]** (*quantified*); the rate gate is closed in the live model, **[Fact-theory]** conditional on $V_{\text{min}} = 0$; the remaining role of the infinite-reservoir premise (next sections) is capacity-*forever*, not capacity-per-interlude. **[Open] (capacity only) V3.7 exchange rate:** with the measured $g = 4.0 \times 10^{14}$, colonization must supply $10^{14.6}$ fresh island volumes per generation; any *finite* $L0$ of $10^X V_{\text{isl}}$ is exhausted after $n_{\text{max}} = X/14.6$ generations ($10^3 \rightarrow 0.2$;

$10^{100} \rightarrow 6.8$; $10^{1000} \rightarrow 68.5$) — “large” buys $\log_{10}/14.6$ rounds; only actually-infinite runs forever. [Auditor]

5. Four worries answered by physics

Stated together so the contrast with the one remaining premise is clean. Each is closed by a mechanism with an owner, not by assumption.

- **Energy loss — closed.** Unitary Hawking evaporation returns the full $M_{\bullet}c^2$; no remnant, no hole swallows energy permanently; $L0$ ’s Minkowski Killing vector (5) conserves the total. Energy in=out. [Fact]
- **Boltzmann brains — closed.** The seeds of a new round are *real debris with a past* (channel 3), not vacuum fluctuations. A *material* floor does not over-produce freak observers the way a fluctuation floor would ($P \propto e^{-\Delta S}$ favours a lone brain over a universe only when the floor is fluctuational). The framework’s floor is material, so the reductio that kills fluctuation-based eternal cosmologies does not bite. [Hypothesis]
- **Classical heat death (uniformity) — closed.** Gravity is not gas-in-a-box: a self-gravitating system has negative heat capacity and *no* maximum-entropy equilibrium (Antonov 1962; Lynden-Bell & Wood 1968, the gravothermal catastrophe). Smoothness is maximally unstable (Jeans), so a near-still state is the most unstable starting point, not a dead end — the perturbation that restarts the engine is a theorem, not a hope. [Fact]
- **Running down to zero — closed.** Rounds are not one tank draining on a global clock; they are born at *different places* in an unbounded volume, each with its own cycle. There is no global time along which a per-round rate declines; “Goldilocks always available *somewhere*” replaces “the reservoir slowly empties.” [Hypothesis]

6. The one assumption that remains

What all four resolutions quietly use — and what (6) and (8) reduce to — is a single premise:

$L0$ is genuinely infinite (or growing), with low-density “Goldilocks” regions always remaining. (9)

If $L0$ were merely *large* but finite, the deposit sites would be finite, every region would eventually thermalise, and the running-down worry would return. The whole structure rests on *infinite*, not *large* — and whether space is actually infinite is unmeasurable in principle (the observable horizon is finite; everything beyond is assumption).

Why this escapes the standard no-go results. The Borde–Guth–Vilenkin theorem makes a spacetime of net positive average expansion past-incomplete; the dust-era endgame gives $H_{\text{avg}} \rightarrow 0$, so the BGV premise can *fail asymptotically*, and the Aguirre–Gratton construction is the documented route through it — past-eternity is now an *adopted* axiom-level choice (item H, decided 2026-06-10; the explicit per-patch construction remains open as item H1, entangled with item A); forward-eternity is free (branching ≥ 1). The Kinney–Stein bind — that zero average expansion leaves no expansion with which to dilute the accumulating entropy — is then escaped *not* by expansion but by the infinite reservoir itself: the entropy density

$$\rho_S = \frac{S_{\text{total}}}{V_{L0}} \rightarrow 0 \quad (V_{L0} \rightarrow \infty) \quad (10)$$

falls even though S_{total} rises every round (second-law-safe). *This is precisely why the infinite- $L0$ premise, not an entropy reset, is the load-bearer.* [Open] (BGV: Borde, Guth & Vilenkin 2003; the bind: Kinney & Stein 2021, arXiv:2110.15380.)

The rate of the bill is measured, not bounded (V3.5). The per-round entropy growth factor $g = (4/3) \Sigma S_{\text{BH}}/S_{\text{bath}}$ is *intensive* — an entropy-density ratio that island size cannot tune — and is now measured: floor 1.7×10^{13} (BHMF holes $\geq 10^9 M_\odot$ only), fiducial 4.1×10^{14} (Egan & Lineweaver 2010 budget), post-ladder $\sim 2 \times 10^{16}$ (assumption-dependent; V3.7 recompute: 7.4×10^{16} , Part VII §1 — OoM-stable). Behind the demographics: Soltan 1982; Marconi & Hunt 2004; Shankar, Weinberg & Miralda-Escudé 2009. **[Fact]** (the measurement; details Part IV §1)

Fork-5 condition (background-independent counting): the population must colonize fresh comoving extent at $\geq (1+g) \approx 10^{14.6}$ per generation, forever $\iff L0$ **unbounded**. Kill: any finite $L0$, or sub-exponential capacity.

The old bound $g \in (0, 10]$ and its $\times 2$ /round kill figure are **retired** (Part V §4); the premise above is unchanged — what changed is that its price per generation is now a number. **[Auditor]**

7. Entropy — diluted, not reset (and why it cannot be reset)

The premise above also fixes how entropy is handled, and a companion result (Part IV §2) shows the alternative is closed off: the survivor’s terminal radiation is massless, and the entropy of massless radiation is *conformally invariant* (photon number is preserved under $g \rightarrow \Omega^2 g$). The Penrose rescale that lights the next hot start therefore *cannot* reset entropy — it carries the radiation entropy $S_{\text{rad}} \simeq \frac{4}{3} S_{\text{BH}}$ (the 4/3 ratio is Zurek’s, 1982) across unchanged. So the second law’s debt must be paid by *dilution in the infinite reservoir*, not by erasure at the boundary. The boundary audit and the closure choice agree: **the rescale cannot reset entropy, so the reservoir must absorb it** — which is the framework’s stated stance and a deliberate departure from Penrose. **[Auditor]**

8. Honest status — theorem vs assumption

Settled by physics (theorem / measured)	Rests on the one premise (undecidable)
Energy returns ($M_\bullet c^2$ via unitary Hawking), no remnant	Energy/capacity in=out at the <i>reservoir total</i>
Minkowski $L0$ conserves total energy (Killing vector)	“Closed system” $\equiv$ “ $L0$ is all there is”
No max-entropy equilibrium for gravity (Antonov; LBW)	Goldilocks regions <i>always</i> remain (measure)
Material floor $\Rightarrow$ no Boltzmann-brain reductio	$L0$ genuinely infinite, not merely large
Rescale cannot reset entropy (massless conformal invariance)	Island-scale seed origin beyond the old horizon (item D/F3; the rate gate itself is closed — item F)
Pair collapse beats evaporation by $\gtrsim 10^{85}$ (Gunn–Gott on the dust era)	Colonization $\geq (1+g) \approx 10^{14.6}$ per generation, forever

The left column is closed. The right column is one premise wearing several hats. It is undecidable in principle (beyond the horizon), and it is the toll the whole genre pays — eternal inflation rests on the same infinite-measure assumption. The framework’s contribution is to *name* it as the single load-bearer and show that nothing else in the closure requires the impossible (a global low-entropy reset). **[Open]**

9. Attribution

Element	Owner / source
$L0$ reservoir; three-source ledger; in=out accounting; material-floor closure	Pitarn ([Design])
Hawking evaporation (T_H , L , τ_{evap}); unitary return	Hawking (1975); Page (1976, spin-down before flash)
Black-hole entropy S_{BH}	Bekenstein (1973)
Nariai ceiling (SdS degenerate horizon) and its instability (Route A)	Nariai (1950); Ginsparg & Perry (1983); Bousso & Hawking (1996)
Radiation-to-black-hole entropy ratio ($S_{\text{rad}} = \frac{4}{3}S_{\text{BH}}$)	Zurek (1982)
No max-entropy equilibrium for self-gravity	Antonov (1962); Lynden-Bell & Wood (1968)
Conformal rescale (scoped); massless-entropy invariance	Penrose (2010)
Cyclic-entropy problem (the thing being answered)	Tolman (1934)
Past-incompleteness; the zero-expansion bind	Borde, Guth & Vilenkin (2003); Kinney & Stein
Energy ledger, Killing-vector argument, proven/assumed audit	editor pass ([Auditor])

Module C is the closure half of the V3.4 math pass. It makes the $L0$ ledger quantitative — three sources (survivor evaporation, Nariai pops, escaped debris), a Minkowski Killing vector that conserves the reservoir total, and an in=out balance that is exact given closure — and it separates the four closure worries that physics settles (energy return, Boltzmann brains, classical heat death, running-down) from the single premise that remains (an actually-infinite reservoir with Goldilocks regions always available). Energy in=out is closed; usable-capacity in=out, and everything else, reduces to that one undecidable premise — correctly the load-bearer, honestly the genre’s shared toll.

Part IV

Operator B' — the aeon transition (export–partition form)

What this part formalizes. The transition between aeons is now an explicit operator with four stages: **B1'** export (what leaves a dying zone, and at what measured rate), **B2'** sorting and partition (what crosses the conformal boundary, what rides the reservoir, and how many children one corpse hosts), **B3'** per-round re-initialization (the unexplained initial data, named out loud), and **B4'** minting (how the next zone’s baryons and dark matter are made from the exported entropy). The old implicit $1 \rightarrow 1$ inheritance of earlier drafts is **superseded**: the material lineage is $1 \rightarrow \sim 10^{14}$, and when aggregation feeds a new zone there is *no unique parent* (§6). **[Design]** (the composition) / owners per stage below.

1. B1' — the export ledger and the measured rate g

$$S_{\text{out}}^{(N)} = S_{\text{leftover}}^{(N)} + \frac{4}{3} \sum_{\text{all } i} S_{\text{BH}}(M_i^{(N)}) = (1 + g) S_{\text{fix}}, \quad g = \frac{(4/3) \Sigma S_{\text{BH}}}{S_{\text{bath}}} \text{Zurek 1982; Tolman 1934} \quad (1)$$

g is *intensive and measured* — an entropy-density ratio that island size cannot tune:

g floor (only $n \geq 10^9 M_\odot$ holes, 10^{-5} Mpc^{-3})	1.7×10^{13}	BHMF only
g fiducial (observable-universe budget)	4.1×10^{14}	Egan–Lineweaver 2010
g post-ladder (mass density in $10^{10-11} M_\odot$ survivors)	2×10^{16} (V3.7: 7.4×10^{16})	$S \propto M^2$

behind it: Soltan 1982; Marconi–Hunt 2004; Shankar, Weinberg & Miralda-Escudé 2009. **[Fact]** (R6 recomputation for this assembly: floor 1.65×10^{13} ; fiducial 4.01×10^{14} , $\log_{10} = 14.60$, from the Egan–Lineweaver S_{BH} budget $3.1 \times 10^{104} k_B$ over the observed bath $1.03 \times 10^{90} k_B$ per observable volume; the post-ladder figure is assumption-dependent — it locks the Soltan-chain mass density into $10^{11} M_\odot$ holes — and reproduces to order of magnitude.)

2. Why entropy is the invariant channel — the conformal boundary and z_{aeon}

(Relocated from the archive’s dark-energy module in V3.5: *this is the justification of B1' — entropy is the cross-aeon bridge variable because the rescale cannot touch it.*) The survivor’s terminal flash is all-massless, and 3.4 keeps Penrose’s conformal rescale *scoped to that channel only* (massive debris persists in the reservoir L0; it never enters the boundary). The rescale is a Weyl map $\tilde{g}_{\mu\nu} = \Omega^2 g_{\mu\nu}$, and the open item z_{aeon} is the magnitude of that rescale — the “blueshift” that gilds the cold evaporated end-state into a hot dense start.

2.1 What the map fixes (the derivable part)

Under $\tilde{g} = \Omega^2 g$, the transformation rules are standard: lengths and proper times scale as Ω , energies and temperatures as Ω^{-1} , a mass ruler as Ω^{-1} (the V1.5 “ $M \sim \Omega^{-1}$ ” scaling, correct for this channel), while *massless* field equations are invariant. The earlier “ $\mathcal{M}_\Omega$ carry operator” that mapped mass, spin axis, and position across the boundary is therefore **superseded for massive**

content in 3.4 — such content rides the reservoir, not the rescale — and survives only as the conformal map on the massless radiation. [\[Auditor\]](#)

2.2 Why the entropy route is blocked (the auditor result)

The proposed first-principles derivation — z_{aeon} from a critical-to-initial entropy ratio $S_{\text{crit}}/S_{\text{initial}}$ — **cannot work**, for a reason internal to the construction. The boundary channel is, by design, the genuinely scale-free one; and the entropy of massless radiation is *conformally invariant* (photon number is preserved under $g \rightarrow \Omega^2 g$ — this is the very fact Penrose’s cyclic cosmology rests on). The survivor of mass $M_\bullet$ evaporates with a radiation entropy (4/3 ratio from Zurek 1982)

$$S_{\text{rad}} \simeq \frac{4}{3} S_{\text{BH}} \simeq \frac{4}{3} (1.05 \times 10^{77}) (M_\bullet/M_\odot)^2 k_B \approx 1.4 \times 10^{101} k_B \quad (M_\bullet \sim 10^{12} M_\odot), \quad (2)$$

matching the per-zone entropy scale (at the V3.5 canonical band top $10^{11} M_\odot$: $S_{\text{rad}} = 1.40 \times 10^{99} k_B$, the order of the per-island $S_{\text{fix}} = 2.8 \times 10^{99} k_B$ of Part V; the 10^{12} figure shown is the upper illustration). Because this entropy is carried across the boundary unchanged, $S_{\text{initial}} = S_{\text{rad}}$: **the rescale generates no entropy ratio to fix z_{aeon} with.** What is left to set the scale is the gluing condition itself — the temperature at which the cold evaporation (with $T_{\text{Hawking}} \sim 10^{-20}$ K for a $10^{12} M_\odot$ hole) is matched to a hot start (a “blueshift” $\sim T_{\text{hot}}/T_{\text{cold}} \sim 10^{47}$) — and that depends on the global structure beyond the causal horizon.

Two consequences.

- **It explains the ambiguity.** The factor- 10^{20} uncertainty previously attached to the $S_{\text{crit}}/S_{\text{initial}}$ heuristic was not a precision that better work would sharpen; the ratio it relied on does not exist, because the relevant entropy is preserved, not reset.
- **It validates 3.4’s entropy stance.** If the conformal rescale cannot reset entropy (it preserves the massless entropy), then the population’s ever-rising entropy must be handled some other way — which is exactly 3.4’s choice: *diluted in the infinite reservoir, not reset*. The conformal-boundary audit and the closure choice agree.

Verdict. z_{aeon} is not a *solvable* open item awaiting a derivation; it is an *irreducible* one, set by a beyond-horizon gluing, in the same epistemic class as the Weyl-smoothness initial condition (O1) and the infinite-reservoir assumption (O3). The honest math-pass outcome is not a number but a boundary on knowledge: here is precisely why no number is derivable. [\[Open\]](#)

3. B2’ — sorting and partition

Massless content crosses; every rest-mass lump goes one-way to $L0$ (re-entry only as the seed-flux channel). The corpse hosts $N_{\text{child}} = f_{\text{part}}(1 + g)$ children, each rescaling a comoving patch holding exactly S_{fix} ; the un-nucleated remainder dilutes in $L0$ forever. Crossing is conformal *per patch* (Penrose 2010; matching conditions Tod), a deliberate departure from CCC’s global rescale; the explicit x^μ rule for lumps and patch boundaries is the committed-but-unwritten item [\[Open\]](#) (F2, blast radius: operator level). [\[Design\]](#)/[\[Hypothesis\]](#)

4. B3’ — per-round re-initialization

$A \rightarrow A_i = 2.70 M_p$; η at baryogenesis (the three conditions: Sakharov 1967); $P(k)$ amplitude and shape; spin axes $\hat{n}$ per the carry rules. All mechanisms [\[Open\]](#) — this is where the framework’s unexplained initial data lives, by design and said out loud.

5. B4' — minting

$$N_\gamma = \frac{f_\gamma S_{\text{fix}}}{3.60 k_B}, \quad M_b = \eta m_p N_\gamma, \quad M_{\text{DM}} = \frac{\Omega_c}{\Omega_b} M_b = 5.36 M_b \eta, \quad \Omega: \text{Planck 2018; bookkeeping: Kolb-Turner} \quad (3)$$

$\hookrightarrow$ recomputed constants (R6, fresh for this assembly): $f_\gamma = 0.5116$, $s/n_\gamma = 3.6016 k_B$, $n_b = 0.2505 \text{ m}^{-3}$, coherence $\rho_b/\Omega_b \rho_c = 1.00$ (definition-level — η is measured as this ratio; recorded as internal consistency, not a test). DM is minted fresh each round as a collisionless thermal relic, symmetric with baryons (passes the Bullet Cluster). **The 5:1 ratio is matched, not predicted:** freeze-out gives $\Omega_{\text{DM}} \propto 1/\langle\sigma v\rangle$, and the canonical $\langle\sigma v\rangle \simeq 3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$ is the value *chosen* to land it (Steigman et al. 2012). Direct-detection nulls (LZ 2024: $\sigma_{\text{SI}} \lesssim \text{few} \times 10^{-48} \text{ cm}^2$ at tens of GeV; XENONnT) close the classic WIMP corridor, so the realization must live in still-open corridors (e.g. dark-sector freeze-out), with $N_{\text{eff}} = 2.99 \pm 0.17$ (Planck 2018) bounding light relics. The parent's mass reaches each child *only as radiation*, re-minted here: the material lineage is $1 \rightarrow \sim 10^{14}$, not $1 \rightarrow 1$. **[Fact-theory]** components, **[Hypothesis]** composition.

6. Endowment gathering — no unique parent (the surviving S7 addendum)

When a new zone forms in $L0$ it gathers from whatever dead-loop outputs lie in its region:

$$S_{\text{endow}} = \sum_k w_k S_{\text{out},k} \quad (\text{gathering weights } w_k \text{ over reachable dead loops}),$$

so the 1:1 parent $\rightarrow$ child lineage of earlier drafts was too strict, in two ways. **(i) No unique parent:** the lump channel is likewise generic, not exotic — new zones form from *mixed* $L0$ debris fields, so carry-through seeds need no special neighbour geometry (the explicit x^μ rule remains item F2, **[Open]**). **(ii) M_{L1} responds to endowment — slowly:** the survivor is the *maximum* of the merger-tree tail, so extreme-value scaling applies; with the exponential cluster tail folded through the *cluster-end owned* slopes $M_\bullet \propto M_{\text{halo}}^\beta$, $\beta \approx 1.05\text{--}1.35$ (Bogdán et al. 2018, observed; Bassini et al. 2019, simulations — item E resolved; the old $\beta \approx 0.7$ was an orphan matching the *stellar* BCG-halo slope), $d \log_{10} M_{L1} \approx 0.15\text{--}0.27$ dex per decade of endowment. Gathering $10\times$ more dead loops moves the survivor only $\times 1.4\text{--}1.9$; gathering half, -0.05 to -0.08 dex. The fixed point of earlier drafts is therefore a **fixed band with a multiplicative walk**: unbiased gathering keeps $\log M_{L1}$ within $\pm 0.7\text{--}1.2$ dex of 11 over 50 rounds. The drifted branch that stacked a gathering $\propto L0$ -growth link on top (and its round count to a ceiling) is **retired** — the link is unestablished (R2: cross-round extrapolation from zero measured rounds) and its ceiling premise died with the constant- Λ interlude (Part V, retired table). **[Hypothesis]** (the band and EV scaling) / **[Dead-end]** (the drift branch). **V3.7 stability audit:** the deterministic return map is *flat* — the next round's M_{L1} is set by gathering statistics, not the parent's mass; slope $\lambda = \beta_{\text{eff}} \cdot 2 \cdot (\text{survivor share of island export}) \approx 1.3 \times 10^{-15}$, so the band is *restoring*. The $\pm 0.7\text{--}1.2$ dex/50-round walk is the pessimistic random-walk-endowment envelope: Monte Carlo (2×10^4 walkers) finds the floor unreachable in 50 rounds in every model, with median first passage 902–1714 rounds under the document's own RW pessimism and never under i.i.d. gathering. **[Auditor]**

Part V

Results of the composed system

1. Per-island fixed band

S_{fix}	$2.80 \times 10^{99} k_B$ — exact fixed point under equal partition
V (today-equivalent)	$9.69 \times 10^{89} \text{ m}^3$
$M_b; M_{\text{DM}}$	$2.04 \times 10^{32}; 1.09 \times 10^{33} M_{\odot}$
M_{L1}	$10^{10}\text{--}10^{11} M_{\odot}$ band , extreme-value scaling ≈ 0.14 dex per decade of endowment

The band slope follows Gumbel/extreme-value statistics (Fisher–Tippett 1928; Gnedenko 1943) for an exponential cluster tail with $M_{\text{max}} \approx 5\text{--}7 M_*$, folded through $M_{\bullet} \propto M_{\text{halo}}^{\beta}$ with the **cluster-end owned slopes** $\beta \approx 1.05$ (Bogdán et al. 2018, observed, via the Lovisari et al. 2015 kT – M_{500} chain) to 1.35 ± 0.15 (Bassini et al. 2019, simulations), giving $\approx 0.15\text{--}0.27$ dex per decade of endowment (up to 0.33 under galaxy-end slopes: Bandara et al. 2009; Ferrarese 2002; Li et al. 2024 weak lensing). The previously quoted 0.14 used an unowned $\beta \approx 0.7$ — numerically coincident with the range’s lower edge, but 0.7 matches the *stellar* $M_{*,\text{BCG}}\text{--}M_{500}$ slope (Kravtsov et al. 2018; Bassini et al. 2019), not the black-hole one; the derivation is corrected and the orphan retired (item E, resolved 2026-06-10). Two knobs stay explicit: steeper Press–Schechter-type tails push the slope *down*; the owned β range pushes it *up* from the old headline. Aggregation statistics drive any walk of the band (Part IV §6; $\pm 0.7\text{--}1.2$ dex over 50 rounds); the old fork-5–linked drift is retired (§4). **[Hypothesis] V3.7 floor:** the band has a computed lower boundary — $R_{\text{isl}}(M) \propto M^{2/3}$ meets λ_J at $M_{\text{floor}} \approx 6.4 \times 10^5 M_{\odot}$, 4.69 dex below the band centre (below it island modes fall sub-Jeans and the open hierarchy stalls) — and no operative ceiling ($M_{\text{Nariai}} \rightarrow \infty$ on the dust interlude; the live ceiling sits +11.8 dex away). **[Auditor]**

This island-volume scale is also a **lower bound on island size**: the measured entropy density of the bath plus the S_{BH} of a band-top L1 survivor require $V \geq 1.36 \times 10^9$ observable volumes ($R \geq 1107 R_{\text{obs}}$, computed from the survivor’s radiation entropy $S_{\text{rad}} = \frac{4}{3} S_{\text{BH}}$ alone; this assembly’s R6 run, which uses the island’s full S_{fix} , gives $V/V_{\text{obs}} = 2.7 \times 10^9$, $R = 1395 R_{\text{obs}}$ — above the bound by exactly $S_{\text{fix}}/S_{\text{rad}} = 2.0$: a bound plus a consistency check, not the same statement twice). **[Fact]** (the arithmetic) / **[Design]** (the reading)

2. The two-sector verdict and the Tolman ledger

$$S_{\text{tot}}^{(n)} = (1 + g)^n S_{\text{tot}}^{(0)} \quad (\text{monotone; the second law as a rate equation}) \text{ Tolman 1934} \quad (1)$$

with g the *measured* rate of Part IV §1 (floor 1.7×10^{13} ; fiducial 4.1×10^{14} ; post-ladder $\sim 2 \times 10^{16}$, V3.7 recompute 7.4×10^{16} — reproduced here because this is the ledger that pays it).

Fork-5 condition (background-independent counting): the population must colonize fresh comoving extent at $\geq (1 + g) \approx 10^{14.6}$ per generation, forever $\iff$ **L0 unbounded**. Kill: any finite L0, or sub-exponential capacity.

Per island the loop closes in matter — densities, the ladder, the BHMF, and the M_{L1} band repeat — and ratchets in *extent* (Fig. 1). The entropy resolution is by dilution into expanding space (Steinhardt–Turok 2002) plus jettison outside the recycling patch (Baum–Frampton 2007), with a steady mortal-islands/eternal-population structure (Aguirre–Gratton 2002; population bookkeeping Garriga–Vilenkin; structural echo of the perfect cosmological principle, Bondi–Gold–Hoyle

1948, applied only at the unobservable L0 level). The dilution lane’s newest realization: Itzhaki, Peleg & Steinhardt 2026 (JCAP 05, 084; arXiv:2508.09745) — a complete cyclic bouncing cosmology from string-theory components, under perturbative control throughout; nearest-neighbour competitor to fork 5 alongside CCC, to be read against the Kinney 2026 re-kill of IS-cyclic carried in the borrowed-legs audit (V3.8) — whether that argument reaches the string construction is unchecked. [Auditor]Eckstein 2023 dissolves the Tolman growth at the definition level via entanglement entropy of a global pure state; LMU keeps the counting version and carries Eckstein as the literature alternative. L0 is *closed in state space per island*, an *irreversible branching tree globally* — “closed circuit” in the first sense only. [Hypothesis] riding fork 5.




Figure 1: Per-island quantities are flat (fixed band); total entropy and island count grow geometrically with the *measured* g . The whole Tolman load sits on fork 5.

3. Predictions and standing tensions

Carry-through predictions: relic spin-axis alignment, testable by NewAthena and LISA (late 2030s); “too-early” lump-seeded holes in young zones (the seed-flux channel). [Hypothesis]

Standing tension — mandatory flag (travels with every reuse of the ladder): the cluster spin-down expectation ($a_\star \sim 0.55$) is in tension with the measured M87 spin $a_\star \gtrsim 0.8$ (Drew et al. 2025) — status *leaning-against*, not falsified; the relic-side spin decoupling is unmeasurable in principle (Part VI §21). **V3.7 inversion:** the spin walk $a = a_0/\sqrt{1 + N}$ (Hughes & Blandford 2003) turns $a_\star \geq 0.8$ into $N_{\text{misaligned}} \leq 0.56\text{--}2.26$ (lower: pure walk from $a_0 = 0.998$; upper: walk allowing accretion re-spin, lift $\times 1.45$) against the modeled 5.9 — tension factor 2.6–10.6, i.e. an aligned-merger fraction $\geq 62\%$ where the model assumed isotropy. Watch for new measurements (ngEHT/BHEX, ~ 2030 s).

4. Retired numbers (V3.5) — nothing on this list may be cited as live

Each entry names the superseded figure, why it died, and the live replacement. Any later document citing one of these as current is in error.

Retired item	Why retired	Live replacement
$g \in (0, 10]$ and the $\times 2/\text{round}$ kill bound “ ≈ 267 rounds to Nariai” (band drift to a ceiling)	survivor-class truncation understated g by ~ 14 orders both premises dead: the $\times 2/\text{round}$ link is unestablished (R2) and the constant- Λ ceiling does not operate on the dust interlude	measured g : floor 1.7×10^{13} , fiducial 4.1×10^{14} (Part IV §1) none; band walk per Part IV §6 ($M_{\text{Nariai}} \rightarrow \infty$ as $H \rightarrow 0$)
Constant- Λ interlude package: e-folds 10^{90} ; de Sitter horizon $1.65 \times 10^{26} \text{ m}$; $s_{L0} = 2.5 \times 10^{-67} k_B \text{ m}^{-3}$; the $0.17 M_\odot$ <i>transport verdict</i>	Correction 1: the interlude is a dust era, not de Sitter (the A-field has $V_{\min} = 0$, $H \rightarrow 0$)	dust-era kinematics $a \propto t^{2/3}$, $H = 2/(3t)$; the timing race (Part III §2)
Nariai $1.8 \times 10^{22} M_\odot$ as a <i>universal</i> ceiling	the ceiling is a live-phase (acceleration-window) state-ment only	scoped ceiling, Part III §1 (eq. Nariai); $M_{\text{Nariai}} \rightarrow \infty$ on the interlude
“Encounter rate $\rightarrow 0$ ” / cross-island terms default-negligible	de Sitter artifact (no event horizon on the dust era)	cross-island forced fork = the main interlude channel (Part III §2); F5 closed by retirement — no successor will be derived (Part VII, G)
“Survivor isolated forever”	isolation holds only during the transient acceleration window, $a \approx 0.6 \rightarrow \text{few-20}$	scoped isolation clause (Part I; Part III §2); survival-based L1 definition unchanged
NGC 1277 mass $1.7 \times 10^{10} M_\odot$ (van den Bosch et al. 2012)	superseded by the adaptive-optics dynamical mass	Walsh et al. 2016: $(4.9 \pm 1.6) \times 10^9 M_\odot$; extreme-overmass flagship: TON 618

Scope notes for the grep-audit: the 10^{90} yr duration of the L1 endgame (Part VI §10) and the w_a value -0.17 (Part II) are different quantities that happen to share digits with retired items; they are live. **[Auditor]**

5. R6 recomputation record (this assembly)

All Part IV–V numbers were recomputed fresh for the V3.5 pass, none inherited, and carry into V3.6 unchanged (the V3.6 ledger pass adds the item-F gate and corrected-band numbers, each R6-recomputed in its session record): $S_{\text{BH}}(10^{11} M_\odot) = 1.049 \times 10^{99} k_B$ (coefficient 1.049×10^{77}); $\tau_{\text{evap}}(10^{11}) = 2.10 \times 10^{100} \text{ yr}$, $(10^{12}) = 2.10 \times 10^{103} \text{ yr}$; $M_{\text{Nariai}}(H_0=67.4) = 1.789 \times 10^{22} M_\odot$; $f_\gamma = 0.5116$; $s_\gamma/n_\gamma = 3.6016 k_B$; $n_b = 0.2504 \text{ m}^{-3}$; $g_{\text{floor}} = 1.65 \times 10^{13}$, $g_{\text{fid}} = 4.01 \times 10^{14}$ ($\log_{10} = 14.60$); $S_{\text{fix}} = s_{\text{bath}} V = 2.800 \times 10^{99} k_B$; $M_b = 2.040 \times 10^{32} M_\odot$; $M_{\text{DM}} = 1.094 \times 10^{33} M_\odot$; Gunn–Gott race $\tau_{\text{evap}}/t_{\text{coll}}(\delta_i=1) \sim 10^{89-90}$, $(\delta_i=10^{-3}) \sim 10^{85}$; $p(f=1/64) = 0.043$, growth $\times 4 \times 10^2$; KG $m=0.8$: $(w_0, w_a) = (-0.9080, -0.1438)$; NGC 384 $\Delta M_\sigma = +0.296$; relic six $+0.227 \pm 0.099$. **[Fact]** (this run)

Part VI

Verification and empirical confrontation

What this part adds. Parts I–III state the framework’s equations and results; this part records an independent re-derivation of them and then confronts the one genuinely distinctive, falsifiable prediction — the relic spin — with real measured data. The honest outcome, stated up front: every number reproduces, the ladder runs as a single forward map that sorts real galaxies by eating-position, and the spin decoupling is the mass-robust core that survives every stress test — yet that same decoupling is, for every object where it would discriminate, currently *unmeasurable*. The framework’s sharpest claim is therefore correctly **[Open]**, neither confirmed nor dead. **[Auditor]**

Update, 2026-06 (restored from the archive’s title page). A later audit pass sharpened the status of the one distinctive claim (the relic/cluster spin decoupling). Two findings, folded into §7 and the closing §21 of this part: (i) the relic side is *unmeasurable in principle* — a relic’s defining quiescence disables every spin probe (X-ray reflection, shadow illumination, merger-GW), so only a still-accreting relic is reachable; and (ii) the testable cluster side — M87 — now trends to high spin, which would not merely tension but *collapse* the decoupling into the coherence parameter f_{coh} . The claim’s status moves from “blocked-but-sharp” to **leaning-against**, decided on the M87 side within ~ 10 years (ngEHT/BHEX). **[Auditor]**

1. Independent verification of the analytic and ODE sectors

Every closed-form number in Parts I–III was recomputed from fundamental constants alone; all reproduce to rounding — Hawking $T_H, L, \tau_{\text{evap}}$; $M_{\text{Nariai}} = 1.79 \times 10^{22} M_\odot$ at $H_0 = 67.4$; $S_{\text{BH}} = 1.05 \times 10^{77} (M/M_\odot)^2 k_B$; $S_{\text{rad}} = \frac{4}{3} S_{\text{BH}} = 1.4 \times 10^{101} k_B$ at $10^{12} M_\odot$ (upper illustration; $1.40 \times 10^{99} k_B$ at the band top $10^{11} M_\odot$); the clustering threshold $\pi^2/4 = 2.47$; the Eddington seed-growth times. Both ODE sectors re-integrate to the same answers: the Klein–Gordon field reaches $w_0 \approx -0.89$ at its DESI-closest displacement and never crosses $w = -1$, and the matter-ladder spin decoupling reproduces (§3). The two analytic corrections folded into Part III — the Nariai coefficient $1/3\sqrt{3}$ and the Zurek $4/3$ entropy ratio — are confirmed. **[Fact]**

2. The ladder as a single forward map

Across the environment plane the five equations partition (σ, n_{gal}) into the four rungs (Fig. 1, left): shallow hosts $\rightarrow$ stuck-L4, deep-and-companion-rich $\rightarrow$ L2, deep-and-isolated-with-a-cocoon $\rightarrow$ L3-relic, the rest $\rightarrow$ L3. Four real galaxies, one per class, fall in their observed bins — Sgr A* (L3), M87 (L2), NGC 1277 (L3-relic), NGC 4395 (stuck-L4). **The decisive demonstration is M87 vs NGC 1277:** near-identical dispersion ($\sigma \approx 324$ vs 333) sorted into *different* rungs by companions and gas history, not mass — “ranking by eating-position, not size,” shown on two real objects. **This is a consistency classifier, not predictive validation:** the class is set by inputs (n_{gal} , gas) that are read in, the sample is one object per class, and (Part I) nothing here discriminates the framework from Λ CDM. **[Design]** (the ranking) / **[Auditor]** (the classifier and check).

3. Mass–spin separability — the decoupling is mass-robust

A natural worry: the framework keys off spin, spin co-evolves with mass, so if the black-hole *mass* is wrong the spin and rung are wrong too — the equations would carry incomplete variables. This




Figure 1: Left: run as one system across the environment plane, the five equations partition (σ, n_{gal}) into the four rungs; four real galaxies (one per class) land in their observed bins. **Right:** the spin decoupling $\Delta a_* = a_*^{\text{relic}} - a_*^{\text{cluster}}$ is positive in 100% of a 160-draw coefficient sweep (median 0.49).

does not hold, for a structural reason. The spin equation of Part I depends on $\dot{M}^{\text{acc}}/M$ and Γ_{mrg} , both invariant under a rescaling of the mass *scale* (accretion and merger rates are each $\propto M$). Re-integrating the relic and cluster archetypes with the seed scaled by $\times 0.1, \times 1, \times 10$ leaves the decoupling intact:

seed scale	relic a_*	cluster a_*	Δa_*
$\times 0.1$	0.998	0.549	+0.45
$\times 1$	0.998	0.522	+0.48
$\times 10$	0.998	0.493	+0.51

and a ± 0.2 -dex shift in M_σ moves the regulated spin by < 0.01 . The decoupling’s *sign and size* are set by the merger term’s presence, not the mass scale; the absolute spin values carry the mass calibration, the falsifiable signature does not. **Choosing spin (not mass) as the axis is the correct response to exactly this worry.** [Auditor]

4. The spin-direction model — magnitude robust, axis fragile

Promoting spin to a vector (magnitude + axis), grounded in chaotic-versus-coherent accretion (King & Pringle 2006), the spin-up budget (Bardeen 1970), and disk alignment (Bardeen–Petterson 1975): **(A)** the magnitude decoupling survives, now with its hidden assumption exposed — “relic spins high” holds provided the cocoon accretion is coherent ($f_{\text{coh}} \gtrsim 0.15$, a mild condition for a single gas reservoir; Fig. 2A). **(B)** an *inherited* spin axis is erased, tilted past 45° once the survivor accretes ~ 0.77 of its mass in the new aeon (Fig. 2B), so the carry-through axis prediction (C5) is a *selection-limited* falsifier, alive only for the least-fed survivors. The variable *refines, does not expand* the predictions; the robust content stays the spin-magnitude decoupling. Coefficients are illustrative (the qualitative behaviour is the literature’s; the exact numbers are not). [Hypothesis]

5. Confronting the distinctive prediction with data

The framework’s one distinctive, falsifiable claim: over-massive *isolated* relics spin *high* (accretion-built, bucking the general high-mass $\rightarrow$ low-spin trend), while merger-built cluster BCGs spin *low*. Confronted with real measurements:




Figure 2: (A) The magnitude decoupling survives once spin is a vector, but “relic spins high” is now explicitly conditional on coherent cocoon feeding ($f_{\text{coh}} \gtrsim 0.15$). (B) An inherited spin axis is tilted past 45° after the survivor accretes ~ 0.77 of its mass in the new aeon — the carry-through axis is a selection-limited falsifier.

The measured-spin trend (X-ray reflection sample, ~ 50 holes; Reynolds 2013/2021 and the 2025 compilation) does show high spin at low mass and lower spin at high mass, attributed to accretion versus mergers — *consistent* with the decoupling, but this is the *standard* interpretation (cosmological simulations predict the same), and the sample is too heterogeneous and selection-biased (X-ray reflection needs active accretion) to discriminate. **Consistent, non-discriminating.**

The relics themselves are unmeasurable. NGC 1277 and Mrk 1216 are quiescent — no accretion, hence no X-ray reflection, no $\text{Fe K}\alpha$, *no spin*. The distinctive test cannot be run. Their *masses* are well measured: for NGC 1277 the high-resolution adaptive-optics dynamical value is $M_\bullet = 4.9 \times 10^9 M_\odot$ ($\log M_\bullet = 9.69 \pm 0.14$; Walsh et al. 2016, confirmed by Krajnović et al. 2018), revising down the original low-resolution van den Bosch et al. (2012) value (now retired; Part V §4); intermediate optical values run $\log M_\bullet \approx 9.1$ (Graham et al. 2016) to 10.1 (Yıldırım et al. 2015). At the adopted $\sigma = 317 \text{ km s}^{-1}$ the Walsh mass sits **+0.24 dex above** the McConnell–Ma M – σ relation — consistent with the relic $+0.22 \pm 0.09$ dex offset (NGC 1277 is one of the $n = 6$ calibrators). *The framework’s calibration should use the Walsh 9.69, not the higher Yıldırım 10.1; the dispersion of literature values (~ 1.1 dex) is itself a caution.* **[Fact]** / **[Auditor]**

M87 is the one near-term test, and its spin is unconstrained. As the canonical L2 the framework predicts a moderate-to-low spin ($a_* \sim 0.5$). Published M87* spin estimates span $a_* \approx 0.1$ – 0.98 and are strongly model-dependent: EHT imaging does not pin it; jet-power / Blandford–Znajek and Doppler-beaming methods lean moderate-to-high (~ 0.5 – 0.9 , one 2025 method giving ~ 0.8 as a lower limit), while a jet-boundary-shape method gives low (~ 0.2 – 0.3). The framework’s ~ 0.5 sits in the middle of that range — *consistent with everything, discriminating nothing*; only a confirmed near-maximal spin ($a_* \gtrsim 0.9$) would tension the merger-spin-down picture. **[Open]**

Verdict. The distinctive spin signature is structurally robust — mass-invariant (§3), surviving the coefficient sweep and the coherence/axis refinement (§4) — yet *empirically blocked for every object where it would discriminate*: the relics are quiescent (no spin) and M87’s spin spans the whole allowed range (no constraint). The framework bets against the observed high-mass $\rightarrow$ low-spin trend for its key objects, with no data able to settle it. Correct status: **[Open]**, neither confirmed nor dead. Unlock routes: NewAthena (~ 2037 , any weakly-accreting relic), LISA (merger-remnant spins), ngEHT (M87 via polarimetry). **[Auditor]**

6. Attribution

Element	Owner / source
Eating-position ranking; relic-spin falsifier; spin-as-axis choice	Pitarn ([Design])
Forward-map classifier; real-galaxy check; mass-spin separability tests; vector-spin model	editor pass ([Auditor])
Chaotic/coherent accretion; spin budget; disk alignment	King & Pringle (2006); Bardeen (1970); Bardeen & Petterson (1975)
Measured SMBH spins	Reynolds (2013, 2021) and 2025 compilation
NGC 1277 black-hole mass	Walsh et al. (2016); Krajnović et al. (2018); cf. van den Bosch et al. (2012), Yıldırım et al. (2015), Graham et al. (2016)
M87* spin estimates	EHT Collaboration; Feng & Wu (2017); Nokhrina et al. (2019); and others (model-dependent)

Part IV is the verification-and-confrontation half of the math pass. The analytic and ODE sectors reproduce independently; the ladder sorts four real galaxies by eating-position (a consistency classifier, not a discriminator); the spin decoupling is shown mass-robust (the correct answer to “mass wrong $\Rightarrow$ all wrong”) and survives a vector-spin refinement that exposes its one condition (coherent feeding) and limits the carry-through axis to least-fed survivors. Confronted with data, the distinctive relic-spin claim is consistent-but-non-discriminating where spins exist, and unmeasurable for the relics themselves (quiescent) and unconstrained for M87 (a_* spans 0.1–0.98). Correctly [**Open**]— the sharpest empirical claim is blocked by the physics of the measurement, not by being wrong.

Part V — Out-of-sample confrontation & 2026 audit update

Part IV tested the spin claim and the ladder-as-classifier. Part V folds in the 2026 confrontation that was not yet included: a genuine *out-of-sample* relic ($M_\bullet$ measured directly, outside the calibration set), the mass-free merger diagnostic, the dwarf empty-centre axes, and the structural reason the distinctive content is either mainstream or measurement-blocked. New owners named; [**Auditor**] marks an audit step.

7. The relic over-massiveness against the 2025 dynamical-mass set

Six red-nugget relics now have robust *direct* dynamical $M_\bullet$ (Cohn et al. 2025, arXiv:2504.00172): NGC 1277, NGC 1271, Mrk 1216 (stellar-dynamical; Walsh et al. 2015/16/17) and UGC 2698, PGC 11179, NGC 384 (molecular-gas-dynamical, ALMA; Cohn et al. 2021/23/24). On $M-\sigma$ every relic lies *above the mean line but within the intrinsic scatter* (~ 0.4 dex), read by Cohn et al. as tentative *mild positive evolution* of $M-\sigma$, not a distinct population. On $M-M_{\text{bul}}$ the stellar-dynamical relics are over-massive but the gas-dynamical UGC 2698 and NGC 384 are *on*-relation; Cohn et al. (2024) flag a possible stellar- vs gas-dynamical method systematic. The interpretation is mainstream — BHs largely finish by $z \sim 2$, then dry mergers add outskirts stars without BH growth (van Dokkum et al. 2010; Ferré-Mateu et al. 2015) — and the Cohn et al. (2025) title, “evolutionary pathway-dependent black-hole scaling relations,” is the framework’s “rank by cause, not mass” as a mainstream result. **LMU is convergent with, not ahead of, this literature.** [**Fact**] / [**Auditor**]

8. NGC 384 — the out-of-sample test point

NGC 384 was *not* in the calibration $n = 6$ from which the $+0.22$ offset was derived, so it is a genuine out-of-sample test. Cohn et al. (2024): $\sigma_\star = 221 \pm 6 \text{ km s}^{-1}$ (circular aperture at R_e), $M_\bullet = 7.26 \times 10^8 M_\odot$ ($\log = 8.86$), ALMA gas-dynamical. On McConnell–Ma (2013), $\log M_\bullet^{\text{pred}} = 8.32 + 5.64 \log(221/200) = 8.565$, so

$$\Delta_{M\sigma}(\text{NGC 384}) = 8.861 - 8.565 = +0.30 \text{ dex (over the mean, within scatter)}. \quad (1)$$

NGC 384 is on-relation on M – M_{bul} but *over*-massive on M – σ — it *confirms* the $+0.22$ -above- M – σ pattern out of sample. Swapping the framework’s NGC 1281 (mass = the less-robust Yıldırım Table-3, M – σ -fixed value) for NGC 384 gives the literature’s robust six:

galaxy	f_{halo} (Zhu 2025)	$\Delta_{M\sigma}$ (dex)
NGC 1277	0.05	+0.24
NGC 1271	0.19	+0.21
Mrk 1216	0.21	+0.31
PGC 11179	0.05	+0.26
UGC 2698	0.38	+0.04 (lone low outlier)
NGC 384	0.14	+0.30 (out-of-sample)
mean $\Delta_{M\sigma} = +0.227 \pm 0.099$ — numerically the framework’s $+0.22 \pm 0.09$		

So the M – σ offset is real and now out-of-sample-supported, but it stays *within the scatter* (not a distinct population) and the M – M_{bul} offset is partly a method artifact. The merger-history *modulation* does not survive: with the hot-inner-halo fraction f_{halo} (Zhu et al. 2025) as a mass-free merger proxy, f_{halo} vs $\Delta_{M\sigma}$ gives Pearson $r = -0.72$ ($p = 0.10$), but leave-one-out shows it is *entirely* driven by UGC 2698 (drop it $\Rightarrow r = +0.24$); the frozen five are flat at $+0.26 \pm 0.04$ across a $4\times$ range in f_{halo} , itself confounded with $M_\star$ ($r = +0.64$). “Merger erases over-massiveness” rests on $n=1$ (R2). **[Auditor]** / **[Dead-end]**

9. The dwarf empty-centre axes (position/occupation, mass-free)

The dwarf bifurcation is, on inspection, a *position/occupation* axis, not a mass axis (dwarf $M_\bullet$ swing 2 – $36\times$; R1). (a) *Wandering*: $\sim 50\%$ of dwarf MBHs are off-centre $> 1 \text{ kpc}$ (Bellovary et al. 2019/2021; Reines et al. 2020), but the mechanism is *dynamical-friction failure (the BH never sinks)*, not GW recoil; the framework’s f_{eject} (ejection from the galaxy, recoil $> v_{\text{esc}} \approx 2\sigma$) is a different quantity from the observed bound-but-displaced population, and the same shallow-potential/long- t_{df} physics that yields a large f_{eject} also prevents the BH–BH merger recoil requires — so $f_{\text{eject}} \leftrightarrow \text{off-nuclear}$ is not a valid test. **[Dead-end]** (b) *Empty centre*: M33 has a central BH $\leq 1500 M_\odot$ (Gebhardt et al. 2001) / ≤ 3000 (Merritt et al. 2001), ~ 3 dex below scaling — a pure disc, no bulge, but an NSC $\sim 2 \times 10^6 M_\odot$. This is *not* “a relic without a BH” (the relic is the over-massive opposite); it is the occupation axis, and its tight upper limit is a cleaner mass-free anchor than off-nuclear-AGN fractions. **[Auditor]**

10. Galaxy spin as phase — complement to the BH-spin claim of Part IV

Part IV’s distinctive claim is high *black-hole* spin $a_\star$. Its galaxy-scale counterpart — “system spin tells forming vs dissipating” — is the fast/slow-rotator paradigm (λ_R ; Emsellem et al. 2011; Cappellari): slow rotators are merger-built/spun-down, fast rotators accretion-built, high- z quiescent

galaxies fast rotators that become slow via later merging. It is measurable but *(i)* galaxy $\lambda_R \neq \text{BH}$ spin a_* ; *(ii)* “stopped = dissipating” conflates quenched/passive (stable for a Hubble time) with terminal/evaporating (the L1 endgame, $\sim 10^{90}$ yr, unobservable); *(iii)* it is mainstream. Tested directly it cannot separate the relics — they are *all* fast rotators by selection. As in Part IV, the one distinctive spin diagnostic (relic / M87 a_*) is blocked by the measurement. [Open]

11. The audit verdict — a structural result

The 2026 confrontation closes the same way Part IV did, now with the out-of-sample point in hand: **wherever the framework’s content is testable today it is mainstream** (over-massive relics, BH-first, pathway-dependent scaling; the dwarf position and fast/slow-rotator axes), **and the one place it is genuinely distinctive — high BH spin a_* in relics and M87 — is unmeasurable until the late 2030s** (NewAthena + LISA + ngEHT). A distinct relic population *above* $M-\sigma$ would need the offset to exceed the intrinsic scatter; at +0.22 it does not. So LMU V3.4 is not at present distinguishable from standard two-phase / pathway-dependent galaxy+BH evolution by available data — “nothing new comes from the literature” is a *structural* fact, not a search failure. [Open]

Empirical sweep (V3.12) — looked at, not load-bearing. *(i)* The NANOGrav 15-yr stochastic GW background (Agazie et al. 2023, $3.5-4\sigma$, amplitude consistent with a SMBH-binary population) supports the ladder’s working assumption that SMBH mergers complete — the final-parsec problem is empirically eased, in the framework’s favour. *(ii)* Limits on stupendously large black holes (Carr–Kühnel–Visinelli 2020, $M \gtrsim 10^{11} M_\odot$ from accretion and lensing) are the observational face of the top-sparse mass-function claim. *(iii)* Dwarf BH occupation fractions (Greene–Strader–Ho 2020; Miller et al. 2015) give the dwarf sequence its statistical backdrop beyond the four named objects. *(iv)* The frozen A -field at BBN: $\rho_A/\rho_{\text{rad}} \sim 10^{-35}$ at $T \sim 1$ MeV — decades below early-dark-energy limits; A is uncoupled to the SM by construction [[Design]] (coupled variants would face Cassini/Eöt-Wash fifth-force bounds). *(v)* Spatial topology: Planck 2015 XVIII finds no multiply-connected signature and floors the fundamental domain at the last-scattering scale; the framework’s own island bound $R \geq 1107 R_{\text{obs}}$ sits orders of magnitude above that floor — a consistency pass, and the reason a 3-torus remains *permitted as topology* while dead as a Λ -source (dead-end log A2). *(vi)* Mission-date hygiene: Athena was redesigned as NewAthena (launch ~ 2037); the former “Athena ~ 2035 ” instances are updated inline (this rebuild). *(vii)* Gigaparsec tracer patterns: the Giant Arc (Lopez, Clowes & Williger 2022, MNRAS 516, 1557) and the Big Ring (Lopez, Clowes & Williger 2024, JCAP) — same sky region, same $z \approx 0.8$, possibly one connected structure, so effectively $n = 1$ (R2) — are claimed cosmological-principle violations; Sawala et al. 2025 find comparable structures in FLAMINGO-10K ΛCDM simulations and in random fields; Lopez & Clowes 2025 (arXiv:2504.14940) rebut. Status: contested — the Hawking-point epistemic shape (pattern detection vs. null-simulation marginalization). Scope note: at Gpc scales nothing assembles within a Hubble time, so a confirmed structure is initial-condition physics — it lands as an added target on the item-(D) generator import (excess large-scale coherence), not on the ladder or the FRW spine; a *cross-aeon* reading, if ever established, falls in the same differential-falsifier basket as Hawking points. [Open] (contested) [Auditor]

12. Action items for the main V3.4 paper

Three edits follow (the main paper is edited separately): **(1)** the main text §12.3 still carries the Nariai coefficient $c^3/2GH$ and $\sim 10^{23} M_\odot$; replace with $c^3/(3\sqrt{3}GH) \approx 1.79 \times 10^{22} M_\odot$ and re-credit the 4/3 entropy ratio to Zurek (1982) — both already correct in this pass. **(2)** Swap NGC 1281 $\rightarrow$ NGC 384 in the relic sample (robust ALMA mass, in the literature’s six; NGC 1281’s

mass is likely $M\text{--}\sigma$ -fixed). **(3)** Reword the distinctive claim from “+0.22 dex distinctively above $M\text{--}\sigma$ ” to “ $\sim +0.23$ dex above the $M\text{--}\sigma$ *mean*, within the intrinsic scatter, consistent with mild $M\text{--}\sigma$ evolution (Cohn et al. 2025); the $M\text{--}M_{\text{bul}}$ offset is partly method-dependent (Cohn et al. 2024).” **[Auditor]**

Part VI — Independent stress-test & open-joint audit (2026)

An independent re-implementation (the author’s) recomputed every checkable number from fundamental constants, and the three open joints flagged in Parts I–III were gated against current (2024–2026) literature. The rule is the project’s: numbers must reproduce, wording must not out-run the physics, every borrowed result is attributed. Two magnitude caveats and one new internal tension surfaced; all are recorded. **[Auditor]**

13. Stress-test — every checkable number reproduces independently

Re-derived from fundamental constants alone, no value inherited from the document:

quantity	re-derived	document	match
M_{Nariai}	$1.789 \times 10^{22} M_{\odot}$	1.79×10^{22}	✓
naive/Nariai ratio	$2.598 = \sqrt{27}/2$	$\sqrt{27}/2$	✓ exact
Hawking T_H ($10^{12} M_{\odot}$)	$6.17 \times 10^{-20} \text{ K}$	$\sim 6 \times 10^{-20}$	✓
τ_{evap}	$2.10 \times 10^{103} \text{ yr}$	$\sim 2 \times 10^{103}$	✓
S_{BH}	$1.05 \times 10^{101} k_B$	1.05×10^{101}	✓
$S_{\text{rad}} = \frac{4}{3} S_{\text{BH}}$	1.40×10^{101}	$\sim 1.4 \times 10^{101}$	✓
pair-production T	$1.186 \times 10^{10} \text{ K}$	1.2×10^{10}	✓
$\pi^2/4$	2.4674	2.47	✓

Both analytic corrections the pass claims (the Nariai $\sqrt{27}/2$ coefficient and the Zurek 4/3 ratio) are confirmed. The gluing blueshift $\sim 10^{47}$ implies $T_{\text{hot}} \sim 6 \times 10^{27} \text{ K}$ (sub-Planck, GUT-scale) — self-consistent, but an estimate, matching the document’s **[Open]** tag. **[Fact]**

14. Stress-test — the dark-energy field reproduces the no-crossing claim

Re-integrating the Klein–Gordon equation on the PNGB potential for three displacements:

ϕ_0/f	w_0	w_a	min w	crosses -1 ?
0.9	−0.958	−0.066	−1.000	no
1.2	−0.910	−0.141	−1.000	no
1.5	−0.810	−0.300	−1.000	no

The field never crosses $w = -1$ (min $w = -1.000$ throughout) — the Vikman theorem confirmed numerically; the thawing locus runs from ΛCDM toward DESI but w_a stays too shallow to reach the central (−0.84, −0.62), matching the document. One honest difference: the independent run finds a $\sim 1.7\sigma$ gap, the document 2–3 σ ; this is f -choice and dataset sensitivity (Pantheon+ vs DES), not an error — the same conclusion (cannot reach the central value; a phantom excursion would be required). **[Fact]** / **[Auditor]**

15. Stress-test — spin decoupling: sign and mass-invariance reproduce; magnitude does not

Re-implementing the ladder spin sector independently:

test	re-implemented	document
$\Delta a_* > 0$ fraction	100.0%	100% ✓
mass-invariance ($\times 0.1/1/10$)	$\Delta = +0.238/+0.247/+0.256$ (relic $\rightarrow 0.998$)	stable ✓
median Δa_*	0.246	0.45 (differs)
fiducial cluster a_*	0.751	~ 0.55 (differs)

The two load-bearing properties reproduce under independent re-implementation: the *sign* is positive in 100% of draws (topology-forced — the relic has no spin-down term, so it is driven to $a_{\text{eq}} = 0.998$, while the cluster always carries one and must sit lower), and the decoupling is *mass-invariant* (the relic reaches 0.998 regardless of seed; Δa_* barely moves), confirming the falsifiable signature does not ride on the BH-mass calibration. The *magnitude* does not reproduce — the independent run gives a median $\Delta a_* \approx 0.25$ (cluster spinning down only to 0.75), not 0.45 — because it simplifies Γ_{mrg} (dropping the Chandrasekhar and $(\sigma/\sigma_0 - 1)$ factors the document carries in full), lowering the cluster spin-down. **This makes the document’s own “size depends on coefficients = calibration” concrete:** the gap is uncertain at the $\sim 2\times$ level, so the falsifier must be stated as “*sign + a conserved gap*,” not “ $\Delta a_* = 0.45$ ” — a cluster measured at $a_* \sim 0.7$ in the late 2030s would not falsify the 0.25-gap version even as it appears to contradict the 0.45 one. [\[Auditor\]](#)

16. Stress-test — Part V empirical numbers reproduce exactly

check	re-derived	document
NGC 384 $\Delta_{M\sigma}$	+0.296	+0.30 ✓
relic-six mean	$+0.227 \pm 0.099$	$+0.227 \pm 0.099$ exact
within scatter?	$+0.227 < 0.4$: yes, not a distinct pop.	✓
f_{halo} corr. (all 6)	$r = -0.724$	-0.72 ✓
drop UGC 2698	$r = +0.242$ (sign flips)	+0.24, $n=1$ -driven ✓

Numbers and honest verdicts both reproduce: “+0.227 within scatter = not a distinct population” and “merger-erase = $n=1$ [\[Dead-end\]](#)” are confirmed. The data inputs (NGC 384 $\sigma_* = 221 \pm 6$; Cohn et al. 2025 title and content) are verified against the current literature. [\[Fact\]](#)

17. Open joint 1 — super-Eddington feedback: strengthened, but “evades” is the wrong word

The hypothesis that an isolated hole can over-grow (the sole route to a relic over-mass, and so the engine under the relic branch and the spin decoupling) is *stronger* than the document’s lone Inayoshi–Haiman–Ostriker (2016) citation: 2024–25 simulations (Massonneau 2023; Bennett 2024; Pacucci & Narayan 2024; Rennehan 2024, uncapped) and JWST broad-line AGN without X-rays reproduce at-or-above-Eddington growth, and LID-568 accretes at $> 4000\%$ Eddington at $z \sim 4$. [\[Hypothesis\]](#) **But the framing must change.** LID-568 drives a strong nuclear outflow (-500 to -600 km s^{-1}): super-Eddington does *not* “evade” the wind — it launches one. The actual mechanism is *radiative (photon) trapping* in the dense slim disc (Abramowicz et al. 1988): the radiative efficiency drops, so the hole can swallow fast *despite* feedback. The open question

— the live falsifier of §9 — is whether that wind shuts off the hole’s *own* accretion (M – σ self-regulation) or only the host’s star formation; LID-568 shows the latter, not the former. **[Open]** Two further points. (i) *A structural strength worth highlighting*: disc self-shadowing can inflate *virial single-epoch* masses by up to 1 dex (Lupi et al. 2024b; GRAVITY/Abuter et al. 2024 find 0.43–0.73 dex offsets), so high- z AGN over-massiveness is partly suspect — but the relics are anchored on *dynamical* masses (Cohn 2025), immune to this artifact; choosing the relic (dynamical) anchor over high- z AGN (virial) is what makes the relic claim robust. **[Auditor]** (ii) *The engine is not the lock*: super-Eddington supports fast *growth* at high z , but “frozen, permanently over-massive at $z \sim 2$ ” additionally needs the BH-first *timing* step (Cohn’s interpretation) — a second, still-assumed link the engine alone does not supply. **[Open]**

18. Open joint 2 — sign of $V_{\min}$: the framework sits on the less-favoured fork

The closure of Part III needs $V_{\min} = 0$ (Minkowski reservoir, no contraction); the rival is $V_{\min} < 0$ (AdS, crunch — the Steinhardt/CBE contracting-cyclic route; Andrei–Ijjas–Steinhardt 2022). Current evidence leans rival, harder than the document states:

- **Data.** Gialamas et al. (2025, arXiv:2506.21542) find a 93.8% preference for a future AdS transition — but it is *model-dependent* (a Higgs-like potential), the no-phantom case still fits at 2σ , and the robust signal is that dark energy is *decaying* (consistent with the framework’s $V \rightarrow 0$). So: leans $V < 0$, but weakly and model-dependently. **[Open]**
- **Theory.** A de Sitter ($V > 0$) minimum is disfavoured by the swampland program and an AdS ($V < 0$) minimum is string-natural, so $V = 0$ is a knife-edge between a forbidden and a natural option — but the swampland conjectures are themselves unproven, so this is a *motivation*, not a verdict. **[Hypothesis] Owners attached (V3.13):** the dS-disfavour leg is the de Sitter swampland conjecture — Obied, Ooguri, Spodyneiko & Vafa 2018 (arXiv:1806.08362); refined: Ooguri, Palti, Shiu & Vafa 2019 (Phys. Lett. B 788, 180); cosmological corollary (thawing quintessence toward $V \rightarrow 0$ — the A-field’s spec): Agrawal, Obied, Steinhardt & Vafa 2018 (Phys. Lett. B 784, 271); Minkowski endpoints are exempt from the conjecture. Independent quantum route, new here: de Sitter as a quantum state self-depletes on the quantum break-time $t_Q \sim M_p^2/(H^3 N_{\text{sp}})$, i.e. $N_Q \sim (M_p/H)^2$ e-folds (Dvali & Gómez 2014, arXiv:1412.8077; Dvali, Gómez & Zell 2017, JCAP 06, 028), on the Gibbons–Hawking 1977 (PRD 15, 2738) finite-entropy base. Mandatory counter-leg: string dS constructions are defended (Kachru–Kallosh–Linde–Trivedi 2003, PRD 68, 046005); all legs **[Hypothesis]**.

Two honest refinements over the document. The framework is *singularly* committed to the $V = 0$ knife-edge, whereas the rival needs no sign tuning (AdS is generic) — so on *naturalness* the framework is worse off *specifically here*, not merely sharing the (value) cosmological-constant problem. Against that, $V = 0$ and $V < 0$ are decided by the *same* observable, so the framework is *as falsifiable as* the rival — an honest, not an escape, position. The verdict waits on DESI DR3 / Euclid / Roman. **[Auditor]**

19. Open joint 3 — f_{coh} coherence: sound logic, but “mild” is optimistic and the cocoon bet is unsettled

The coherent/chaotic dichotomy is established (coherent, fixed angular-momentum axis $\rightarrow a > 0.9$: Dotti et al. 2010, Thorne 1974; chaotic, isotropic $\rightarrow a = 0.1$ – 0.3 : King & Pringle 2006). The document’s f_{coh} has a *sound* structural origin: in the “chaotic-II” regime (Fanidakis et al.

2011) a *massive* hole cannot align with each clump and averages toward zero-angular-momentum accretion $\rightarrow$ low spin, so a high-spin relic ($\sim 10^{10} M_{\odot}$) *forces* coherent feeding — f_{coh} is required, not invented. **[Hypothesis]** But two cracks. (i) “ $f_{\text{coh}} \gtrsim 0.15$, mild” is optimistic: retrograde accretion spins down more efficiently than prograde spins up (the retrograde ISCO is farther out), so in a 15/85 mix the efficient retrograde share pushes the true threshold *higher* unless the coherent fraction is sustained-prograde — “conditional,” not “mild.” (ii) Whether a relic’s dense early cocoon is truly a single coherent reservoir is a *bet against* the observed massive-BH $\rightarrow$ low-spin trend (the document admits “bucking the trend”), and simulations may overstate coherence (NewHorizon-type resolution effects), pushing real coherence *below* what is assumed. The cluster (low-spin) side is, by contrast, well supported (chaotic cold accretion $\rightarrow$ low-spin equilibrium with misaligned torques and Blandford–Znajek spin-down; BlackHoleWeather 2025). Settled directly only by NewAthena (~ 2037): if relic spins are low, coherence *and* the decoupling fail together. **[Open]**

20. Open joint 4 (new) — the engine and the spin condition pull against each other

Surfaced by reading joints 1 and 3 together: super-Eddington over-growth needs *low* radiative efficiency (photon trapping), while a high relic spin needs *high* efficiency. A relic must therefore occupy a “coherent-but-trapped” window satisfying both at once — a composite, untested condition, not two independent supports. This also bites the verified Pass 3: its scalar model bakes in the relic reaching $a_{\text{eq}} = 0.998$ (coherence assumed), so if real coherence is lower (joint 3), the relic’s a_{eq} is < 0.998 and the “100% positive sign” is itself *conditional on* f_{coh} . The sign stays topology-protected (relic above cluster) as long as the relic remains the *less*-spun-down of the pair; what is conditional is the magnitude, and with it the discriminating power. **[Open]**

21. Audit verdict — one distinctive claim on a three-link conditional chain

The framework’s single non-circular, distinctive expectation — the spin decoupling — hangs on a chain in which every link is live but unconfirmed, and two links are in tension:

$$\begin{aligned}
 &\text{super-Eddington over-growth} \quad \mathbf{[Hypothesis]} \\
 \rightarrow &\text{relic over-massive} \quad (\text{Cohn timing}) \quad \mathbf{[Open]} \\
 \rightarrow &\text{coherent feeding} \Rightarrow \text{high spin} \quad \mathbf{[Hypothesis]} \\
 \rightarrow &\Delta a_* > 0 \quad (\text{sign verified; magnitude}) \quad \mathbf{[Open]}
 \end{aligned}$$

with joints 1 and 3 pulling against each other (joint 4). The expectation is genuinely *non-circular and falsifiable* (NewAthena, late 2030s), and its *sign* is topology-protected and reproduces independently — but it is *conditional-deep*, not standalone-robust. The pattern across all four joints is uniform and, on honesty, to the framework’s credit: each open joint is flagged by the document itself, the physics is sound, and the only corrections needed are three over-confident phrases — “evades” (joint 1), “slightly the other way” (joint 2), “mild” (joint 3) — to be softened to “trapping despite outflow,” “data and (conjectural) theory lean to the rival,” and “conditional, against the trend.” No physics is wrong; the distinctive content is sharp enough to test and honest about waiting for the data. **[Auditor]**

References added in this audit

Abramowicz et al. (1988, slim discs); Andrei, Ijjas & Steinhardt (2022, slow contraction); Bennett et al. (2024); Dotti et al. (2010); Fanidakis et al. (2011, chaotic-II); Gialamas et al. (2025, arXiv:2506.21542); GRAVITY/Abuter et al. (2024, BLR offsets); Inayoshi & Ichikawa (2024, LRD density $\rightarrow$ high spin); King & Pringle (2006, chaotic accretion); Lupi et al. (2024b, disc self-shadowing); Massonneau et al. (2023); Pacucci & Narayan (2024); Rennehan et al. (2024, uncapped accretion); Reynolds (2024, coherent-accretion preference); Suh et al. (2025, LID-568); BlackHoleWeather (2025, chaotic-cold-accretion low spin). All used as owned results; the framework supplies only the readings. [\[Auditor\]](#)

Parts I–III write the worldview’s claims as equations on a spine of standard physics (every equation owned and tagged); Part IV verifies the analytics and tests the spin claim; Part V confronts the distinctive content with 2023–2026 direct-mass data (the out-of-sample relic NGC 384); Part VI re-derives every checkable number independently and gates the now-four open joints against current literature. The verdict is consistent across all six: the picture is internally second-law-consistent and its qualitative readings match the mainstream, the one new equation (the ladder) does not discriminate from Λ CDM, the closure rests on the $V_{\min} = 0$ knife-edge the data presently disfavour, and the single distinctive signature — the relic/cluster spin decoupling — is non-circular and falsifiable but conditional-deep, settled only by NewAthena/LISA (late 2030s) and DESI DR3. Worldview and provenance map; not a peer-reviewed theory.

Part VII

Stress audit with measured inputs, and Decision Record DR-Vmin

What this part is. Two 2026-06-11 sessions, carried as one record. §1: every stage of the composed loop $\Phi = \Lambda_{\text{live}} \circ B' \circ \Lambda_0$ driven by *measured* values only (Planck 2018; Fixsen 2009; DESI DR2+Pantheon+, Abdul-Karim et al. 2025; EHT 2019; GRAVITY 2022; Walsh 2016; Drew 2025; Egan–Lineweaver 2010; Soltan/Shankar), with the system’s stability quantified. §§2–6: the V_{min} Decision Record — the trichotomy with prices, the fourth runner Ω_k , the per-zone reading, and the cascade-to-zero mechanism. **Status of the DR: RULED-CONDITIONAL (2026-06-11, author):** paths 4+4b adopted — $V = 0$ read as the dynamically selected attractor, every lineage’s pathway to $L0$ set by its birth draw; spine numbers unchanged (all branches identical below ϵ_{DESI}). **Conditions carried:** the K1 legs (§5), re-roll (item L), the why-zero conjecture leg. **Scripts:** lmu_v36_stress_test.py, lmu_dr_vmin_scan.py, curvature_fourth_runner.py, l0_pull_numbers.py, l1_zoo_endgame.py, three_floors.py.

1. The stress audit — the spine reproduces; seven findings

Reproduction. From measured inputs alone, this pass independently recovers: $m = 0.805 H_0$ at $A_i = 2.70$ with the locus row $m = 0.80 \rightarrow (w_0, w_a) = (-0.9080, -0.1438)$ exact; the acceleration window $(0.60, 5.49)$; $a_{\text{osc}} = 9.37$; $\min w = -1.000000$; late slopes $(-3.001, -1.500)$; $f_\gamma = 0.5116$; g floor/fiducial/post = $1.65 \times 10^{13}/4.01 \times 10^{14}/7.4 \times 10^{16}$; $V_{\text{isl}} = 1.356 \times 10^9 V_{\text{obs}}$ ($R = 1107 R_{\text{obs}}$); minting $M_b = 2.05 \times 10^{32}$, $M_{\text{DM}} = 1.10 \times 10^{33} M_\odot$; $\lambda_J = 1.682 \times 10^{26} \text{ m}$ (margin 2896); $\tau_{\text{evap}}(10^{11}) = 2.097 \times 10^{100} \text{ yr}$; $M_{\text{Nariai}} = 1.789 \times 10^{22} M_\odot$; $\text{race} = 1.2 \times 10^{89}$. **[Fact]** (this run)

Findings (full report: LMU_V3_6_STRESS_TEST_realdata.md): **F1** DESI joint distance pinned on the released chains: $\rho = -0.894, 2.69\sigma/2.40\sigma$ Gaussian/posterior-mass (applied in Part II). **F2** the V_{min} tolerance tiers (applied at items B/J2). **F3** the M87 inversion (applied at the mandatory flag). **F4/F5** band restoring, $\lambda \approx 1.3 \times 10^{-15}$; Jeans floor $6.4 \times 10^5 M_\odot$, no ceiling (applied in Parts IV–V). **F6** the fork-5 exchange rate (applied in Part III). **F7** the boundary injects $\times \Omega \in [10^{26}, 10^{47}]$ per crossing; entropy/photon-count closure is an identity (applied in Part III §3). Insensitivities: the Hubble tension shifts everything $\leq 18\%$; the $\times 4500$ spread in g changes the bill, never a verdict; Ω_m $0.310 \rightarrow 0.3153$ moves (w_0, w_a) by ≤ 0.001 . **[Auditor]**

2. DR-Vmin — the trichotomy with prices

$\epsilon \equiv |V_{\text{min}}|/\rho_{\text{DE,obs}}$. The field equation contains no V_{min} (the derivative of a constant vanishes), so the thaw-and-oscillate history is branch-blind until the floor’s budget share grows: $|\Delta A| < 4.6 \times 10^{-3}$ for $a < 2$ at $\epsilon = 10^{-2}$ (Fig. 2). Today’s observables move as $|\Delta w_0| \approx 0.185 \epsilon$: $\epsilon_{\text{DESI}} = 3.0 \times 10^{-2}$ at the $0.1\sigma_{w_0}$ convention — below it the three branches are observationally identical and share the pre-registered DESI-Y5 falsifier ($w \geq -1$, canonical). **[Fact-theory]**

$V_{\text{min}} = -\epsilon$ (**the rival, now integrated**). Turnaround at $a = 5.8/13.9/30.4$ for $\epsilon = 10^{-1}/10^{-2}/10^{-3}$ ($t_{\text{turn}} = 2.2 \times 10^{18}/6.2 \times 10^{18}/1.9 \times 10^{19} \text{ s}$): the ladder completes in every case; the survivor never evaporates — B' is replaced wholesale by the bounce. The LQC effective equations (Ashtekar–Pawlowski–Singh 2006; $\rho_c = 0.41\rho_{\text{Pl}}$) were integrated through the bounce: $\rho/\rho_c = 1.0000$ at $a_{\text{min}} > 0$, H flips $-285.7 \rightarrow +285.7$ (demo units), constraint violation $\leq 8 \times 10^{-7}$; real contraction spans ~ 42 decades, $a_{\text{bounce}}/a_{\text{turn}} \sim 10^{-42}$. Item (B) dissolves (*any* $\epsilon < 0$ closes a cycle); the landmine is BKL anisotropy in contraction (needs an ekpyrotic-class smoother, Erickson et al. 2004); BH-through-bounce **[Open]** (Quintin & Brandenberger 2016). **[Fact-theory]** (mechanics)

/ **[Hypothesis]** (adoption)

$V_{\min} = +\epsilon$ (**survival ceilings; B16 correction**). The dS refreeze at $t_{\text{freeze}} = [6\pi G \epsilon \rho_{\text{DE}}]^{-1/2}$ truncates each stage. Computed ceilings: strict-B2' loop (children = conformal patches of one corpse) $\epsilon < 3 \times 10^{-2}$ (the DESI ceiling — evaporation and the flash complete for any relevant ϵ : $T_{\text{dS}} \ll T_H$, $M_{\text{Nariai}}(H_\infty) \gg \text{band}$); intra-island aggregation $\epsilon < 4.3 \times 10^{-3}$; **cross-island endowment gathering of k corpses** $\epsilon < 2.2 \times 10^{-26} (k/2)^{-3.5}$ (horizon re-entry + Poisson seeds); old-style unbounded colonization $\epsilon < 3.1 \times 10^{-181}$. *Correction of record (B16)*: a chat-level estimate had placed neighbour gathering at the intra-island tier (4×10^{-3}); the computed cross-island ceiling is 23 decades tighter. Fork-5 volume is paid by expansion itself: 11.2 dS e-folds per generation mint the $(1+g)$ volume in $10^{20}\text{--}10^{32}\text{ s} \ll \tau$ — the infinite-reservoir premise is *replaced* by eternal expansion (Steinhardt–Turok dilution; the Kinney–Stein bind is void), at the price that BGV 2003 reactivates and item (H) becomes load-bearing on Aguirre–Gratton 2002. Item (A) is *import-resolved* on this branch: spacelike $\mathcal{I}^+$ returns and the Penrose 2010 / Eckstein 2023 crossing works natively. **[Auditor]**

LMU DR-Vmin -- the V_min trichotomy with prices (R6 scan)




Figure 1: The trichotomy with prices: (a) observational invisibility below $\epsilon_{\text{DESI}} = 3 \times 10^{-2}$; (b) $+\epsilon$ survival ceilings; (c) $-\epsilon$ turnaround family; (d) the LQC bounce, integrated — the flip before zero.

3. The fourth runner — spatial curvature Ω_k

The endgame race had three runners (matter a^{-3} , floor a^0 , shear a^{-6}); curvature (a^{-2}) was missing. Closed: turnaround at $a = D/(3|\Omega_k|)$; open: linear growth freezes under curvature domination (Peebles 1980). Ceilings on the $V_{\min} = 0$ branch: first pairs free (< 1.24); island hierarchy $< 2.9 \times 10^{-18}$; eternity $< 5.1 \times 10^{-60}$ (≈ 68 e-folds of flattening per aeon, Guth 1981




Figure 2: Same wave, three floors: the field equation is $V_{\min}$ -blind; only the Friedmann floor differs. The wave reveals the floor only when the endgame has already arrived — the damped oscillation is evidence for all three branches equally.

scaling). At today's observational bound $|\Omega_k| \sim 10^{-3}$, a closed sign crunches the interlude at 7.7×10^{15} yr. Consequences: the $V_{\min} = 0$ branch carries *two* knife-edges; Ω_k joins the B3' re-init list (item I); the $+\epsilon$ branch is self-curing (dS e-folds re-flatten); $-\epsilon$ indifferent. Pattern note: this is again **Type-B** — a known GR variable whose pricing converges on a mainstream requirement. **[Auditor] Literature anchor (audit add):** Dinda & Maartens (2025, MNRAS 542 L31; arXiv:2504.15190) fit thawing quintessence on a *curved* background to DR2 and find it matches flat $w_0 w_a$ — the fourth runner is already a live, partially w -degenerate degree of freedom in the current data fits. **[Fact]**

4. Path 4 — the per-zone reading

$V_{\min}$ can be read as B3' re-initialization *data* rather than universal law: frozen vacuum draws per zone (landscape mechanism: Bousso–Polchinski 2000; Susskind 2003; the small-but-nonzero prediction: Weinberg 1987). $R_{\text{isl}} \geq 1107 R_{\text{obs}}$ hides per-zone variance three decades beyond any horizon — isotropy data do not constrain it. The trichotomy then ceases to be a fork to choose: the population walks all three according to its distribution, and selection acts (flash-lineages out-reproduce bounce-lineages by $\sim 10^{14.6}$ per generation). **[Hypothesis]**

5. Path 4b — cascade-to-zero (author wiring), and the absorbing state

Vacua do not average by size; at any contact the *lower* V eats the higher through domain-wall/vacuum-decay dynamics (Coleman 1977; Coleman–De Luccia 1980). Two computed anchors: (i) $L0$ at $V = 0$ is a **fortress** — nucleating AdS of depth ρ_{DE} inside Minkowski is gravitationally blocked unless the wall tension is below $\sigma_{\text{crit}} = 5.8 \times 10^{16} \text{ J m}^{-2}$ ($\approx (24 \text{ MeV})^3$); every wall at or above the electroweak scale exceeds this by 10^{10} – 10^{53} (a QCD-scale wall by only $\sim 10^2$). **[Fact-theory]** (ii) $+\epsilon$ **zones are genuinely pulled to 0**: $\text{dS} \rightarrow \text{Minkowski}$ decay is allowed with thin-wall exponent $\log_{10} B \approx 166$ (electroweak wall) — the wait is $e^{10^{166}}$, glacial but finite against the branch’s eternity, so $+\epsilon$ and 0 merge asymptotically. **The cascade guarantees the destination, not the interlude** ($e^{10^{166}} \gg$ every loop deadline $\sim 10^{107} \text{ s}$); the tier table of §2 still governs mid-flight. Why zero (and not a deeper floor) ends the cascade is the conjecture leg: non-SUSY AdS unstable (Ooguri–Vafa 2017), SUSY Minkowski uniquely stable (positive-energy theorem, Witten 1981). **[Hypothesis]**

Absorbing state and the completed triad. With a bounce-side re-draw of V_{min} (Wheeler reprocessing / Smolin CNS 1992; **[Hypothesis]**, item L — plain LQC conserves the potential), every zone reaches $V = 0$ through one of three doors: $\text{draw} \approx 0$ merges directly (clock $\tau \propto M^3$); $+\epsilon$ evaporates, flashes, and its leftover vacuum is pulled (the cascade never pulls an L1 — the survivor is $10^{(10^{166})}$ -fold gone by then; L1 is a *role*, not an object); $-\epsilon$ bounces and re-rolls until it exits to ≥ 0 (geometric waiting; per-zone Tolman accumulation capped as a side effect). Nothing draws exact zero — zero is an **attractor**, consistent with the standing nothing-is-exactly-0/1 principle. Heredity completes the Darwinian triad: *laws inherited perfectly* (children are patches of the corpse — same V_{min} , same constants), *state re-rolled* (fresh $P(k)$); fertility $N_{\text{child}} = 1 + g$ varies $\times 4500$ with the zone’s BH census — the variance selection needs. Generative non-stillness is *carried in by the corpse* (flash + debris + endowment), not grown from the void: Minkowski at $T = 0$ births nothing real (vacuum-jitter rebirth would import the Boltzmann-brain pathology, Dyson–Kleban–Susskind 2002; Albrecht–Sorbo 2004); the one owned exception is the $+\epsilon$ far-future thermal channel (recycling universe, Garriga–Vilenkin 1998). Keystone — **resolved-conditional (K1, 2026-06-11)**: the σ_{crit} fortress blocks only quantum nucleation; the classical account of *born- $-\epsilon$ zones* closes with three locks (`k1_wall_accounting.py`; record `LMU_K1_wall_accounting.md`). **(C) flat-space**: a wall grows only beyond $R^* = 2\sigma/(\epsilon\rho_{\text{DE}})$ (thin wall, Coleman 1977); island-scale $-\epsilon$ domains sit $R^*/R_{\text{isl}} \geq 1.6 \times 10^{10}$ *below* growth for an EW wall at the DESI ceiling — the tension crushes them inward (even a QCD wall clears by $\times 3.6$ at the worst point, and the bound collapses to sub-eV^3 as ϵ falls). **(G) gravity (decisive)**: any island-scale wall of tension $\geq \text{QCD}$ is in the Zel’dovich–Kobzarev–Okun 1974 catastrophic regime — $R_{\text{isl}}/R_{\text{grav}} = 1.4 \times 10^6$ (QCD) to 6×10^{15} (EW), wall mass 10^{32} – $10^{42} M_{\odot}$ deep inside its own Schwarzschild radius: planar walls repel (Vilenkin 1981; Ipser–Sikivie 1984), closed walls pinch off behind horizons (Blau–Guendelman–Guth 1987; Berezin–Kuzmin–Tkachev 1987), and exterior conversion is obstructed outright (Farhi–Guth 1987). Implosion, never invasion; the crunch ($2.2 \times 10^{11} \text{ yr}$ at ceiling ϵ) and island light-crossing ($5.1 \times 10^{13} \text{ yr}$) sit ~ 86 orders inside the 10^{100} -yr budget. **Two-sided ledger**: outside, the $-\epsilon$ zone deposits as an ordinary BH corpse ($\tau \propto M^3$, standard $L0$ furniture — the vacuum book is never touched); inside, crunch $\rightarrow$ LQC bounce $\rightarrow$ re-roll — **item (L) acquires its geometric home inside the collapse horizon, Smolin’s native CNS setting.** **[Fact-theory]** (the three locks) / **[Hypothesis]** (re-roll)

6. Status, rulings required, and the audit trail

Ruling (item K, 2026-06-11): paths 4+4b **adopted, conditional** — the population walks all three branches by birth draw; $V = 0$ is the selected attractor; K1 computed and closed (§5).




Figure 3: The L1 zoo — ten unlike zones, integrated: *whether* a zone enters $V = 0$ at $L0$ is decided by its $(V_{\min}, \Omega_k)$ draw alone; M_{L1} and A_i set *when* ($\tau \propto M^3$), never *whether*. Late slopes check: dust -1.500 , Milne -1.032 .

Spine numbers unchanged. What would reopen this: a sub-QCD-tension wall mechanism between vacuum draws, a failure of the bounce re-roll (L), or the why-zero conjecture leg falling (Ooguri–Vafa/Witten). If $-\epsilon$ is ever *measured* ($w < -1$ sustained at DESI-Y5 with systematics resolved), the B' -replacement build spec opens regardless of this ruling. **Failure log of these sessions** (R4, carried in the session records): two file-write truncations (mitigated by chunked writes); the B16 tier conflation (corrected, §2); ϵ_{DESI} is now computed on the chain covariance ($4.6 \times 10^{-2} \rho_{\text{DE}}$, joint 0.1σ ; the 0.1σ -on- w_0 convention retired in V3.8); gathering ceilings are estimate-grade (Poisson seeds, $\delta_c = 1.686$); one missed file-presentation step (checklist updated). Deliverable records: LMU_V3_6_STRESS_TEST_realdata.md, LMU_DR_Vmin_trichotomy.md + addenda (curvature, 4b, absorbing state, triad). [\[Auditor\]](#)

7. The (A)-window — the crossing restored on the ruled spine

Item (A) was the boundary problem of a zone at $V_{\min} = 0$ *exactly*: null $\mathcal{I}^+$, no Penrose/Eckstein crossing. Under the (K) ruling nothing draws exact zero (the attractor is approached, never sampled — and exact zero has measure zero besides), so every lineage carries $\pm\epsilon'$: the $-\epsilon'$ half bounces inside its collapse horizon (K1) and never needs $\mathcal{I}^+$; the $+\epsilon'$ half ends de Sitter for *any* $\epsilon' > 0$ — $H \rightarrow H_\infty = H_0 \sqrt{0.69 \epsilon'}$, conformal time $\eta = -e^{-H_\infty \tau} / H_\infty$, $ds^2 = (H_\infty \eta)^{-2} (-d\eta^2 + d\vec{x}^2)$, $\mathcal{I}^+$ spacelike — and the CCC machinery runs *native* (Penrose 2010 itself requires $\Lambda > 0$). [\[Fact-theory\]](#)

The clocks (computed; `lmu_A_window.py`). Vacuum domination at $a_{\text{dom}} = (1.45/\epsilon')^{1/3}$, $t_{\text{dom}} = t_0 a_{\text{dom}}^{3/2}$; the split $\boxed{\epsilon'_{\text{crit}} = 6.3 \times 10^{-181}}$ — flash-in-dS above, flash-in-dust-then-dS below, *both* admitting the slice — re-derives the full-program tier of §2 from the other direction (the same arithmetic, inverted). Evaporation is undisturbed everywhere on the grid ($T_H/T_{\text{dS}} \geq 10^{12}$; $M_{\text{Nariai}}(H_\infty) \geq 10^{23} M_\odot \gg \text{band}$). The window $[\max(t_{\text{dom}}, \tau_{\text{evap}}), t_{\text{decay}}] \sim [10^{108} \text{ s}, e^{10^{166}} \text{ s}]$ supplies $\sim e^{10^{166}}$ e-folds; the cascade rate $\Gamma \sim e^{-10^{166}}$ is hyper-subcritical, so the dS phase is globally future-eternal (eternal-inflation persistence) and un-decayed comoving patches are certain

on the slice. **[Fact-theory] Quantum-breaking caveat (V3.13).** The same lane that backs $V_{\min} = 0$ (J2, Part VI §18) cuts against *unconditional* future-eternity here: if de Sitter quantum-breaks (Dvali–Gómez–Zell 2017), the usable budget is $N_Q \sim (M_p/H_\infty)^2$ e-folds — R6-recomputed: 1.39×10^{122} at $\epsilon' = \epsilon_{\text{DESI}} = 3 \times 10^{-2}$ and 6.60×10^{300} at $\epsilon'_{\text{crit}} = 6.3 \times 10^{-181}$ (a species factor $N_{\text{sp}} \sim 10^2$ shifts the exponent by -2) — replacing the decay-rate budget $\sim e^{10^{166}}$. The crossing itself is indifferent: it needs only $N = O(10^2)$ ($e^{-2N} = 10^{-87}$ at $N = 100$), margin $\gtrsim 10^{120}$; but “globally future-eternal” must be read as conditional on the absence of quantum breaking. **[Hypothesis]** (the cut) / **[Fact-theory]** (the margin arithmetic).

The drafted construction (equations first, proof later — author doctrine). Matching slice Σ_N at N e-folds past $\max(t_{\text{dom}}, \tau_{\text{evap}})$: dust residual $\propto e^{-3N}$; Weyl-cleanliness met exponentially for any finite N , the rate set by the *slowest* channel — frozen super-horizon tensors at e^{-2N} (electric) / e^{-3N} (magnetic, the Tod-2007 jump channel $\partial_n B$); fluid-sourced Weyl e^{-4N} (V3.9 erratum, computed). Crossing: $\hat{g} = \Omega^2 g$ with $\Omega = -1/(H_\infty \eta)$, reciprocal continuation $\hat{\Omega} = -1/\Omega$ (Penrose 2010), Tod matching conditions on Σ_N ; per-patch scope unchanged from B2' (child i = comoving ball $B_i \subset \Sigma_N$ holding exactly S_{fix}); entropy invariant (B1' unchanged); $z_{\text{aeon}}(\epsilon') = T_{\text{reheat}}/T_{\text{dS}}(\epsilon')$ — now *parametrized* per zone, with T_{reheat} the same beyond-horizon irreducible as before. **[Design] Proof obligations (V3.9 status): P1** the Tod junction theorem for $(g, \hat{g})$ on Σ_N — **FRW core closed** (V3.9): a component of EoS w enters $1/a$ at $\eta^{3(1+w)}$, integer for every carried component; the oscillating A -residue makes the crossing exactly $C^{2,1}$ (C^2 , the Einstein/FPX requirement, holds with zero margin; C^3 log-periodically blocked); the child bang is radiation-exact; the FPX hypotheses hold at the root window. *Open remainder (reduced, V3.10): G1* — reciprocal gluing, now *perturbative openness off the Kopiński–Valiente Kroon 2022 conformally-flat matching point*; mismatch channels priced ([G1-h/B] automatic; [G1-E] e^{-2N} ; [G1-j] e^{-3N}), conditional travelling: [G1- Ω] (the $\hat{\Omega}$ prescription: Newman/Tod/Nurowski/Markwell–Stevens), now also absorbing the $C^{2,1}$ knife-edge — [G1-d] *closed* (V3.11): on the flat-conformal-class spine LT09 holds trivially and the kink constrains $\hat{\Omega}$ only; in the TT sector the LT09 ladder formalizes the horns (smooth horn passes, price π_{ij} ; class horn = impulsive-wave $C^{1,1}$ interface outside LT09 class (i), amplitude e^{-2N}). **G2** — contaminated-bang well-posedness, reduced to one named theorem (inhomogeneous radiation $+\Lambda+ w' = -2/3$ image + regular-branch scalar, at $C^{2,1}$; nearest anchors Anguige–Tod 1999, Tod 2007, Lee–Nungesser–Tod 2020), one computed condition (G2-c: the $t^{-1/2}$ stiff-branch amplitude vanishes on Σ), the named tool *executed* (V3.11, [G2-d]: the Gover–Kopiński–Waldron 2023 criterion — every LMU passenger lands in an allowed trace slot, the dust-image is geometrized as H^{ext} , and the stiff branch has no slot, so G2-c is doubly derived; cross-aeon stress matching inherits [G1- Ω]) [math-GR, **[Open]**; records LMU_G1_G2_pass.md, LMU_G1d_G2d_pass.md]. **P2** — **closed-conditional** (V3.9, horizon exile): Ω is global within a zone; every wall/ ∂B_i comoving scale exits the H_∞ horizon in finite N , so the crossing region never sees them (item (C) closes on the same budget; falls with (K)). **P3** $P(k)$ through Σ_N — *item (D), re-scoped by DR- $P(k)$ (ruled, V3.12)*: the spectrum is *per-aeon*, not inherited — the ruling formalizes the computation (all channels die at $e^{-2N}/e^{-3N}/e^{-4N}$; the slowest survivor is tensor, not adiabatic scalar — wrong type and spectrum for A_s). Each aeon must therefore contain its own fluctuation generator [**[Open]**, named import]; targets it must hit: $n_s = 0.9649 \pm 0.0042$, $A_s = 2.1 \times 10^{-9}$, $r < 0.036$ (BICEP/Keck 2021), $f_{\text{NL}}^{\text{loc}} = -0.9 \pm 5.1$, percent-level isocurvature (Planck 2018). Generator existence is *necessary for loop closure*: no $\delta \sim 10^{-5} \Rightarrow$ no galaxies $\Rightarrow$ no ladder $\Rightarrow$ no next flash. Discount specific to this framework: horizon and flatness are already paid by the crossover (causal contact from the parent’s infinite past; Penrose 2010), so the import is a *generator*, not full inflation — but a flash-as-seed shortcut is pre-blocked by the dead-end-log precedent (flash-as-DE died on 13–49 orders; same burden of amplitude + spectrum + Gaussianity applies). Wording rule, document-wide: the flash lights the *pre-generator* start; the observed hot start (BBN/CMB) is lit by the generator’s reheating. The transfer question survives only as the *cleanliness side*: what leaks across must stay below $r < 0.036$ and PTA stochastic-background limits — both passed automatically at large N . Forced stance: **LMU predicts the absence of CCC-style Hawking points and concentric circles** — horizon

exile puts every other island’s flash outside the horizon — a clean, falsifiable departure from Penrose’s CCC (claims: Gurzadyan–Penrose 2010; An–Meissner–Nurowski–Penrose 2020; mechanism: Meissner–Penrose 2025, arXiv:2503.24263 — the GWE crossover; critiques: Moss–Scott–Zibin 2011; Jow–Scott 2020). **Sharpened (V3.13).** Meissner–Penrose 2025 upgrades Hawking points from claimed signal to *structural requirement* of CCC’s crossover (conformally smooth *except* at Hawking points), making this falsifier symmetric: confirmed HPs falsify LMU’s horizon exile; sustained nulls leave CCC’s mechanism with an unfulfilled requirement. Detection status: contested — An–Meissner–Nurowski–Penrose 2020 (MNRAS 495, 3403) claim 99.98%; Jow & Scott 2020 (JCAP 03, 021) find 87% ($\sim 1\sigma$) after ring-size marginalization. **Stakes flag, mandatory:** a confirmed HP detection does not merely close DR-P(k) — it breaks horizon exile itself, re-opening item (C), the transfer side of (D), and P2; everything stamped “rides (K)/(A)” re-opens with it. The background map (the c_3/c_4 series) stands; perturbation-level cleanliness **[Open]**. (H1) gains its arena: the $+\epsilon'$ window is the future-side Aguirre–Gratton stage. No new debt — conditions inherited verbatim from (K); if (K) falls, this falls with it. **[Auditor]**

Part VIII

Open ledger and dead-end pointers

1. The open ledger (one list, no duplicates)

- (A) **Non-conformal rebirth — resolved-conditional / drafted** (2026-06-11, rides the (K) ruling): under “nothing draws exact zero,” every lineage is $\pm\epsilon'$ — the $-\epsilon'$ half bounces inside its horizon (K1, no $\mathcal{I}^+$ needed); the $+\epsilon'$ half ends de Sitter for *any* $\epsilon' > 0$, restoring the spacelike $\mathcal{I}^+$ that Penrose’s and Eckstein’s crossings require natively. Window computed: $\epsilon'_{\text{crit}} = 6.3 \times 10^{-181}$ (the full-program tier, re-derived inverted) splits flash-in-dS from flash-in-dust-then-dS; both admit the matching slice Σ_N ; budget $\sim e^{10^{166}}$ e-folds; evaporation undisturbed ($T_H/T_{\text{dS}} \geq 10^{12}$). Equation set E1–E4 drafted with proof obligations P1–P3 (Part VII §7); P2 = item (C) sharpened, P3 = item (D) sharpened. *Item (A) was the boundary condition of a zone that, under the ruling, never exists.* No new debt — conditions inherited verbatim from (K). **[Fact-theory]** (geometry) / **[Design]** (construction)
- (B) $V_{\text{min}} = 0$ **vs. Weinberg 1989**: a QFT-level wall; classical intuition cannot reach it. **[Open]** **Priced (V3.7)**: the interlude survives a residual $|V_{\text{min}}|/\rho_{\text{DE}}$ only below 4.3×10^{-3} (first pairs) / 5.5×10^{-56} (island hierarchy) / 3.1×10^{-181} (full program to τ) — ~ 59 decades beyond the classic 10^{-122} problem; in Planck units the floor must sit at $\sim 10^{-303}$. Branches and a candidate dissolving mechanism: Part VII-DR.
- (C) **F2**: the explicit x^μ /patch-boundary rule for lumps across a per-zone rescale. Sharpened by the (A)-window: the rule now lives on a concrete surface — Ω regularity at $\partial B_i \subset \Sigma_N$ (obligation P2, Part VII §7) — **closed-conditional** (V3.9, horizon exile: every finite wall/patch scale exits the H_∞ horizon in finite N ; rides the (K)/(A) budget, falls with (K)). **[Fact-theory]** (geometry, conditional)
- (D) **F3**: the $P(k)$ re-initialization mechanism — **re-scoped by DR-P(k) (ruled, V3.12)**: the spectrum is *per-aeon*, not inherited (every transfer channel dies at e^{-2N} or faster; the slowest survivor is tensor — wrong type and spectrum for A_s). The open content is now the *in-aeon fluctuation generator* [named import] with targets n_s , A_s , $r < 0.036$, f_{NL} (P3, Part VII §7); the cross- Σ side survives only as cleanliness bounds, passed at large N . **[Open]** (generator)
- (E) β **owner — resolved** (2026-06-10): $\beta \approx 0.7$ has no owner as an $M_\bullet$ – M_{halo} slope; all owned values are ≥ 1 (Bogdán et al. 2018 ≈ 1.05 , observed cluster end; Bassini et al. 2019 1.35 ± 0.15 , sims; Bandara et al. 2009; Ferrarese 2002; Li et al. 2024 lensing ≈ 1.7). The 0.7 matches the stellar $M_{*,\text{BCG}}$ – M_{500} slope (Kravtsov et al. 2018) — a relation mix-up. Band slope corrected to 0.15–0.27 dex/decade; the headline survives at the range’s lower edge, the derivation is replaced (Part V §1). **[Fact]**
- (F) **A-dust Jeans gate — resolved-conditional** (2026-06-10): island scales are super-Jeans from oscillation onset for all $m \gtrsim 0.14 H_0$ ($R_{\text{isl}}/\lambda_J = 2.9 \times 10^3$ at the live point, growing as $a^{1/4}$); the A-dust clusters with $p \rightarrow 1$, island modes re-enter at $a/a_{\text{osc}} \approx 2.6 \times 10^7$, and any seed $\delta \gtrsim 10^{-52}$ collapses before τ (the Poisson floor alone exceeds the requirement by $\sim 10^{42}$). The gate closes only for $m \lesssim 0.13 H_0$ via horizon exile (> 100 e-folds of late inflation), a regime outside the DESI-thaw identification. The suppressed branch ($p = 0.044$, $\times 4 \times 10^2$) is scoped to sub-Jeans scales ($< 0.38 R_{\text{obs}}$ comoving at onset). Conditional on $V_{\text{min}} = 0$ (item B); seed origin stays open in (D). Files: `lmu_itemF_jeans_gate.py/.png`. **[Fact-theory]**
- (G) **F5 — closed by retirement** (2026-06-10, author ruling): no constant- Λ medium exists at any stage (dark energy is the A-field on the $L1$ cycle; $L0$ has $A \approx 0$), so the F5 constant- Λ source terms get no successor and the check-before-quote obligation is discharged by renun-

ciation. $\rho^* \sim 10^{-116}$ and $0.17 M_\odot$ stay permanently quarantined in the retired table; fork-5 cross-island *delivery* (Part III §2, coupling iv) is unaffected. Any future source terms are a new item, derived on the dust background. [\[Design\]](#)

- (H) **Past-eternity — decided** (2026-06-10): adopted as an axiom-level choice — the population is eternal in both time directions, no global beginning; islands die and are born, locations shifting through matter/radiation exchange. Forward-eternity is free (branching ≥ 1); the past side is carried by the Aguirre–Gratton 2002 construction already cited in Part V, with Borde–Guth–Vilenkin 2003 standing as the theorem the axiom must answer. [\[Design\]](#)
- (H1) **The explicit construction — drafted-conditional** (V3.8): anchor ruled (a) per-lineage (author, 2026-06-11); equations H-E1–H-E4 — per-tree past eternal in proper time, *finite in aeon count*; AG carries the arrow, FPX25 the metric continuation, never conflated. Obligations: **P1** (Tod junction; FRW core closed V3.9 — the crossing is $C^{2,1}$, zero margin at C^2 ; the Tod-2007 obstruction channel is sourced at the computed e^{-3N} ; remainder G1+G2 — reduced V3.10, checks run V3.11: G1 = perturbation off the KVK22 solved sector, [G1- Ω] travelling ([G1-d] closed, merged into [G1- Ω]); G2 = one named theorem + condition G2-c (doubly derived) + [G2-d] executed) and **P4** (geodesic exhaustion; reduces to P1 $\wedge$ the KMS24–FPX25 dispute $\wedge$ LQC-effective, modulo wall-riding pathologies). Rides (K)/(A); falls with (K). [\[Open\]](#) (P1, P4)
- (I) $A_i = 2.70 M_p$ **mechanism** (with η , $P(k)$, $\hat{n}$, and Ω_k — **added V3.7, the fourth endgame runner**: the B3' re-initialization data). [\[Open\]](#)
- (K) **DR-Vmin — ruled-conditional** (2026-06-11, author): paths 4+4b adopted — $V_{\min}$ is per-zone birth data (variation), $V = 0$ the dynamically selected attractor (cascade + absorbing state), every pathway to $L0$ set by the $(V_{\min}, \Omega_k)$ draw. Spine numbers unchanged (branch-blind below ϵ_{DESI}). **K1 wall accounting resolved-conditional** (Part VII §5): $-\epsilon$ zones cannot convert $L0$ — quantum door blocked (σ_{crit}), classical door crushed ($R^*/R_{\text{isl}} \geq 1.6 \times 10^{10}$ EW), gravity door catastrophic-to-self (ZKO; $R_{\text{isl}}/R_{\text{grav}} \geq 1.4 \times 10^6$); outside = BH corpse, inside = bounce+re-roll. Conditions: walls $\geq$ QCD scale; thin-wall junctions; item (L); the why-zero conjecture leg. [\[Design\]](#) (the ruling) / [\[Fact-theory\]](#) (K1)
- (L) **Bounce re-roll of $V_{\min}$** (path-4b leg): plain LQC conserves the potential; the re-draw is Wheeler-reprocessing / Smolin-CNS class; K1 locates it geometrically inside the $-\epsilon$ collapse horizon — Smolin's native setting. Ownership of the inside-the-horizon-universe picture itself: Pathria 1972 (Nature 240, 298) first; Popławski 2010 (Phys. Lett. B 687, 110; 694, 181) the modern Einstein–Cartan bounce realization — the $-\epsilon'$ branch is their setting with a re-drawn $V_{\min}$ added. [\[Hypothesis\]](#)

The four empirical joints of the ladder audit live in Part VI §§17–20 and are listed here only for the single-ledger view:

- (J1) super-Eddington feedback — “trapping despite outflow,” not “evades” (Part VI §17). [\[Open\]](#)
- (J2) the sign of $V_{\min}$ — the framework sits on the less-favoured fork; same observable as (B)'s data side (Part VI §18). [\[Open\]](#) **Mapped (V3.7)**: all three branches are observationally identical today below $\epsilon_{\text{DESI}} = 3 \times 10^{-2}$ and share the DESI-Y5 falsifier; the discriminators are theoretical economy, not near-term data (Part VII-DR).
- (J3) f_{coh} coherence — “conditional, against the trend,” not “mild” (Part VI §19). [\[Open\]](#)
- (J4) the engine and the spin condition pull against each other — the composite “coherent-but-trapped” window (Part VI §20). [\[Open\]](#)

With the M87 *leaning-against* flag (Parts I, V, VI) attached wherever the ladder is reused.

2. Dead-end pointers (check before extending anything)

A1 — DM origin, killed $\times 5$ (Bullet Cluster + timing + 5:1); correct form: thermal relic re-minted each aeon. **[Dead-end]** **A2** — DE/ Λ origin: rotation/vorticity (Saadeh et al. 2016 kills), Hawking-tail-as-DE, flash-as-A-source (short by 13–49 orders) — all dead; in V3.4/3.5 dark energy is the A-field, no constant remains. **[Dead-end]** **B** — gravity-beats-expansion: **now scoped** — valid only during the acceleration window; void on the dust interlude (Correction 1). Mid-aeon baryogenesis stays dead (hot boundary only). **[Dead-end]/[Design]** **B-W** — Weyl-residue disposal/repurposing (both directions dead). *Disposal*: cancellation or absorption of the frozen $E/\partial_n B$ residues dies on (i) super-horizon causality, (ii) incoherent power addition (stochastic GW add in power, never subtract), (iii) category — the gap is mathematical (the Anguige–Tod class demands exact zero), so any “reduce the residue” mechanism is structurally redundant against the unlimited e-fold budget. *Repurposing* (residue as asymmetry seed): dead on the DR-P(k) ruling (slowest survivor is tensor — wrong type/spectrum for A_s), on scaling (a knob flowing to zero with N cannot set a constant; η sits at B3', Sakharov per round), and on the Weyl Curvature Hypothesis (Penrose 1979) running the *opposite* way — the arrow lives in the residue’s smallness, not in the residue. Residues have one job: pricing the G1 perturbation. **[Dead-end]** **CCBH** (Crocker–Weiner; Farrah et al. 2023): context, coupling scoped off on the spine; R8 applies to rebuttals.

Part IX

Attribution — the master table (R7, deduplicated)

Per-part attribution sections (Parts I, II, III, VI) are retained where they stand, as the archive record. This table is the deduplicated union: one row per borrowed component, its owner, and its role in the loop. Nothing here is claimed by the author; the author’s own contribution is the single closing row.

Component	Owner	LMU role
FRW background	Friedmann 1922	per-zone clock
Quintessence field equation	Ratra–Peebles 1988; Wetterich 1988; Caldwell, Dave & Steinhardt 1998	A-field slot
Thawing class, w_a relation	Caldwell–Linder 2005; Scherrer–Sen 2008	verified band
Thawing PNGB potential	Frieman, Hill, Stebbins & Waga 1995	archive cross-audit
CPL parametrization w_0, w_a	Chevallier–Polarski 2001; Linder 2003	DESI fitting form
No phantom crossing (canonical bound)	Vikman 2005	the $w \geq -1$ wall
Oscillation $\rightarrow$ dust, $\langle w \rangle = 0$	Turner 1983	interlude background
Transient thawing vs DESI; Bayesian comparison	de Souza, Rodrigues & Alcaniz 2025 (PRD; arXiv:2504.16337)	published engine check
Phantom crossing as a CPL artefact	Dinda & Maartens 2025 (MNRAS 542 L31; arXiv:2504.15190)	the Y5 question
DESI DR2 data	DESI Collaboration / Abdul-Karim et al. 2025 (arXiv:2503.14738)	A-field test
Cosmological-constant problem	Weinberg 1989	the $V_{\min} = 0$ wall (B)
Halo mass function	Press–Schechter 1974; Sheth et al. 2001	ladder input
Velocity-dispersion function	Sheth et al. 2003	BHMF composition
Seeds; Eddington/Salpeter e -fold; hyper-Eddington	Salpeter 1964; Inayoshi, Haiman & Ostriker 2016	low-mass end
Bondi accretion	Bondi 1952	gas-supply term
Coagulation	Smoluchowski 1916	merger kernel
$M_{\bullet}-\sigma$	McConnell–Ma 2013; Kormendy–Ho 2013	ceiling map
AGN feedback / M_{σ} self-regulation	Silk & Rees 1998; King 2003	the ceiling’s engine
Relaxation; dynamical friction; GW inspiral	Chandrasekhar 1943; White & Rees 1978; Peters 1964	rates
Spin-up limit; merger spin-down	Thorne 1974; Hughes & Blandford 2003; Berti & Volonteri 2008	spin decoupling
Chaotic vs coherent accretion spin	King & Pringle 2006; Bardeen 1970; Bardeen & Petterson 1975	spin-state inputs
Escape speed; recoil/superkick	NFW 1996; González et al. 2007; Campanelli et al. 2007	survivor protection
Starvation quenching	Bullock et al. 2000	relic freezing
Two-phase relic growth	van Dokkum et al. 2010; Ferré-Mateu et al. 2015/2017	relic story
Fast/slow rotators (λ_R)	Emsellem et al. 2011; Cappellari	galaxy-spin contrast
Relic dynamical masses	Walsh et al. 2016; Cohn et al. 2021–2024 (ALMA); Krajnović et al. 2018	the $n = 6$ anchors
Relic sample; out-of-sample relics	Yıldırım et al. 2017; Newman et al. 2025 (MRG-M0138)	over-massiveness
Measured BH spins (X-ray reflection)	Reynolds 2013/2021; compilations to 2025	spin-trend check

Component	Owner	LMU role
M87 spin measurement	Drew et al. 2025 (and EHT-era methods)	the leaning-against flag
UHZ1 status (contested)	Alvarez-Márquez et al. 2026; Fujimoto et al. 2024	dwarf-timeline caveat
Superradiance caveat	Arvanitaki & Dubovsky 2011; Brito et al. 2015	spin-extraction loophole
Isotropy bound (kills rotation-DE)	Saadeh et al. 2016	dead-end A2
Spherical collapse	Gunn–Gott 1972	interlude aggregation
Ultralight Jeans suppression, $k_J \propto a^{1/4}$	Hu–Barkana–Gruzinov 2000	$p(m)$ gate — (F) resolved
Scalar-field Jeans scale	Khlopov–Malomed–Zeldovich 1985	gate formula lineage
Mixed-component growth $p(f)$	linear theory (Peebles 1980)	sub-Jeans branch
Slow-roll e-folds $N \approx A_i^2/4$	Linde 1983	horizon-exile closure
Poisson seeds from discrete sources	Afshordi–McDonald–Spergel 2003	island seed floor
Evaporation; spin-down before flash	Hawking 1975; Page 1976	interlude clock
S_{BH} ; radiation 4/3	Bekenstein 1973; Zurek 1982	export ledger
Nariai ceiling; instability	Nariai 1950; Ginsparg & Perry 1983; Bousso & Hawking 1996	scoped live-phase ceiling
Gravothermal catastrophe	Antonov 1962; Lynden-Bell & Wood 1968	no-heat-death theorem
Entropy budget (measured g)	Egan–Lineweaver 2010; Soltan 1982; Marconi–Hunt 2004; Shankar et al. 2009	the ratchet rate
Entropy-growth problem	Tolman 1934	the bill
Zero-expansion entropy bind	Kinney & Stein	the bind fork 5 answers
Dilution resolution	Steinhardt–Turok 2002	fork-5 reading
Jettison / causal-patch cycling	Baum–Frampton 2007	partition reading
Steady eternal population	Aguirre–Gratton 2002; Garriga–Vilenkin	L0 structure; (H) decided axiom
Past-eternity theorem	Borde–Guth–Vilenkin 2003	the standing theorem behind (H); construction drafted (H1); proofs open (P1, P4)
Perfect cosmological principle	Bondi–Gold–Hoyle 1948	structural echo (L0 only)
Conformal crossing	Penrose 2010; Tod	per-patch, scoped
Definition-level Tolman dissolution	Eckstein 2023	literature alternative
Isotropic-singularity IVP shelf	Goode–Wainwright 1985; Newman 1993; Claudel–Newman 1998; Anguige–Tod 1999 I/II; Anguige 2000; Tod 2003, 2007; Lee–Nungesser–Tod 2020	G2 anchors
$\Lambda > 0$ smooth- $\mathcal{I}^+$ shelf	Friedrich 1986, 2017, 2023; Lübke–Valiente Kroon 2013; Joudioux–Thaller–Valiente Kroon 2021; Hadžić–Speck 2015	parent boundary closed
CCC two-sided matching (base point)	Kopiński–Valiente Kroon 2022	G1 solved sector
$\hat{\Omega}$ prescriptions (mutually inconsistent)	Newman 2014; Tod 2015; Nurowski 2021; Markwell–Stevens 2023	conditional [G1- Ω]
Generic-stress bang admissibility	Gover–Kopiński–Waldron 2023	[G2-d] executed (V3.11)
Conformal-gauge-singularity extension	Lübke–Tod 2008, 2009; Lübke 2009, 2013	[G1-d] closed (V3.11)
Conformal fundamental forms / matching	Blitz–Gover–Waldron; Gover–Kopiński 2023; Nurowski 2021	GKW §VI dictionary (inherits [G1- Ω])
Baryogenesis conditions; η , Ω 's	Sakharov 1967; Planck 2018	minting
CMB monopole	Fixsen 2009	anchors

Component	Owner	LMU role
Extreme-value scaling $M_{\bullet} \propto M_{\text{halo}}^{\beta}$ (cluster end)	Fisher–Tippett 1928; Gnedenko 1943 Bogdán et al. 2018; Bassini et al. 2019 (range: Bandara et al. 2009; Ferrarese 2002; Li et al. 2024)	M_{L1} band band fold, $\beta \approx 1.05\text{--}1.35$ — (E) resolved
CCBH (context only) Primordial-spectrum targets SMBH-binary GW background	Croker–Weiner; Farrah et al. 2023 Planck 2018; BICEP/Keck 2021 NANOGrav: Agazie et al. 2023	scoped off the spine DR-P(k) generator spec merger support; cleanli- ness bound
DM exclusion landscape	LZ 2024; XENONnT; Steigman et al. 2012	minting corridor
Hawking points: claim / critique	Gurzadyan–Penrose 2010; An et al. 2020 / Moss–Scott–Zibin 2011; Jow–Scott 2020	LMU predicts absence
SLABs; dwarf occupation; topol- ogy	Carr–Kühnel–Visinelli 2020; Greene–Strader–Ho 2020; Planck 2015 XVIII	sweep (V3.12)
Author’s own: the zone/L-ladder architecture, the forced-fork coupling choice, the survival-based L1 definition, the export–partition composition of B' , the eating-position ranking, the reservoir/ledger architecture of Part III, and the assignment of the equations above to the loop — a hybrid ; every mechanism is owned in the rows above. Integrations, sweeps, and the proven/assumed audits: editor pass ([Auditor]).		

End of the V3.13 consolidated pass. The components are physics; the wiring is the worldview; the bill is on fork 5 — now with its exchange rate — and the $V_{\min}$ floor carries a price tag and a candidate cascade.