



Assessing Ecological Risk of Invasive Alien Plants in South Korea

Author(s): Ji-Hyon Kil, Kew-Cheol Shim and Ho-Joon Lee

Source: Weed Technology, 2004, Vol. 18, Invasive Weed Symposium (2004), pp. 1490-1492

Published by: Cambridge University Press on behalf of the Weed Science Society of

America

Stable URL: https://www.jstor.org/stable/3989679

REFERENCES

Linked references are available on JSTOR for this article: https://www.jstor.org/stable/3989679?seq=1&cid=pdf-reference#references_tab_contents
You may need to log in to JSTOR to access the linked references.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at https://about.jstor.org/terms



Weed Science Society of America and Cambridge University Press are collaborating with JSTOR to digitize, preserve and extend access to Weed Technology

Symposium =

Assessing Ecological Risk of Invasive Alien Plants in South Korea¹

JI-HYON KIL, KEW-CHEOL SHIM, and HO-JOON LEE2

Abstract: Invasive alien plant species represent an ecological risk to indigenous plants and ecosystems. In this study, the dichotomous system is suggested to assess the ecological risk of invasive plant species. Case studies of noxious alien plants in South Korea were used to develop the system based on environmental risk factors. Assessing ecological risk of invasive alien plants was considered with the following factors: harmful effects to animals or humans, ecological characteristics, invasion time, present state after invasion (dispersion), and availability of resources. Invasive plant risk assessment was based on the methodology for predictive risk assessment used to manage and monitor invasive alien plants in South Korea. Our system enabled us to classify invasive plants in Korea into four grades: (1) noxious species (three categories), (2) monitoring species (five categories), (3) harmless species (innocuous species), and (4) beneficial species. Our assessment system can help land managers monitor invasive alien plant species.

Additional index words: Dichotomous system, ranking system.

INTRODUCTION

The threats to the natural environment by invasive species are a global issue (De Poorter 2002; Vitousek et al. 1996). Invasive species threaten to affect other plant species and communities as they expand their ranges and invade new areas. The invasion of exotic plants into ecosystems is detrimental on an ecological level because it can potentially alter the balance between the native species (Coblentz 1990; OTA 1993; Vitousek et al. 1996).

² Researcher, Division of Plant Ecology, National Institute of Environmental Research, Environmental Research Complex, Gyungseo-dong, Seogu, Incheon 404-170, Republic of Korea; Assistant Professor, Department of Biology Education, Kongju National University, Shinkwan-dong, Gongju, Chungnam 314-701, Republic of Korea; Professor, Department of Biological Sciences, Konkuk University, Hwayang-dong, Gwangjin-gu, Seoul 143-701, Republic of Korea. Corresponding author's E-mail: skcshim@kongju.ac.kr.




Figure 1. Priority factors to assess an ecological risk of invasive plants in South Korea.

Invasive alien plants also are harmful to humans and domestic livestock (Harper et al. 1996). Because invasive species are increasing, we recognize the need for a classification system to help management planning with invaded ecosystems (Heffernan et al. 2001; Koh et al. 2002).

Many countries attempting to solve the problems of invasive plants want to develop a system of invasive plant management in an ecological, environmentally

A1. harmful to human, animals, and crop plants	
	Maniana anastra t
B1. harmful to human, animals and crop plants in the country	Noxious species i
B2. harmful to human, animals and crop plants out of country,	
but never reported to be harmful to human, animals and crop plants	
C1. disturbed ecologically in the country	- Noxious species II
C2. disturbed ecologically out of the country,	
but never reported to disturbed ecologically in the country	
D1. less than 20 years since invasion	
E1. dispersed widely	
E2. dispersed locally	- Monitoring species III
D2. more than 20 years since invasion	0 1
E3. dispersed widely	- Monitoring species II
E4. dispersed locally	
A2. never reported to be harmful to human, animals, and crop plants in o	
B3. disturbed ecologically	· · · · · · · · · · · · · · · · · · ·
C3. disturbed ecologically in the country	Noxious species III
C4. disturbed ecologically out of the country, but never disturbed ec	
D3. less than 20 years since invasion in the country	orogramity in the country
E5. dispersed widely	- Monitoring species I
E6. dispersed locally	Monitoring species III
D4. more than 20 years since invasion in the country	Monitoring species in
E7. dispersed widely	- Monitoring species II
E8. dispersed locally	
B4. never disturbed ecologically in or out of the country	- Monitoring species iv
C5. never valuable in commerce, agriculture and economy	
D5. less than 20 years since invasion in the country, and dispersed	eri da ter
D6. more than 20 years since invasion in the country	
C6. valuable in commerce, agriculture and economy	

Figure 2. Dichotomous key to selected grades and categories of invasive alien plant species by ecological risk in South Korea.

¹ Received for publication February 6, 2004, and in revised form May 5, 2004

Table 1. Grades, categories, and characteristics of ecological risk, and examples of invasive alien plants in South Korea.

Grade and category	Characteristics	Plant species
Noxious species		
I	Found to be harmful to humans, animals (domes- tics), and crops in and out of the country Potential to induce damages directly by toxic chem-	Solanum carolinense L., Ambrosia artemisiifolia L., Ambrosia trifida L.
II	icals Found to be harmful to humans, animals (domestics), and crops out of the country but not in the country	Eupatorium rugosum Houtt.
Ш	Predictable to be harmful in the country Not found to be harmful to humans, animals (domestics), and crops in and out of the country Found to disturb ecologically on the ecosystem in and out of the country	Paspalum distichum L., Paspalum distichum L. var. indutum Shinners
Monitoring species		
I	Less than 20 y since invasion and distributed widely in the country Found to disturb ecologically on the ecosystem out of the country but not in the country Predictable to disturb ecologically in the country	Coronopus didymus Smith, Paspalum dilatatum Poir.
II	More than 20 y since invasion and distributed widely in the country Found to disturb ecologically on the ecosystem out of the country but not in the country	Senecio vulgaris L., Agropyron repens (L.) Beauv., Festuca arundinacea Schreb, Chenopodium al- bum L., Phytolacca americana L., Thlaspi ar- vense L., Medicago sativa L., Convolvulus ar- vensis L., Datura stramonium L.
Ш	Less than 20 y since invasion and distributed local- ly in the country Found to disturb ecologically on the ecosystem out of the country but not in the country	Cirsium arvense Scop., Hordeum murinum L., Sorghum halepense (L.) Pers., Raphanus raphanistrum L., Sinapis arvensis L., Conium maculatum L.
IV	More than 20 y since invasion and distributed lo- cally in the country Found to disturb ecologically on the ecosystem out of the country but not in the country	Lolium temulentum L.
V	Less than 20 y since invasion in the country Not found to disturb ecologically on the ecosystem in or out of the country	Hordeum jubatum L., Papaver dubium L.
Innocuous species	More than 20 y since invasion in the country Not found to disturb ecologically on the ecosystem in or out of the country	Erechtites hieracifolia Raf., Crassocephalum crepi- dioides S. Moore, Veronica persica Poiret
Beneficial species	Valuable in commerce, agriculture, and economy Not found to disturb ecologically on the ecosystem in or out of the country	Chenopodium ambrosioides L., Oenothera biennis L., Tagetes minuta L.

sound, and cost-effective way (Hobbs and Humphries 1995; Koh et al. 2001, 2002; Suter 1993). By understanding ecological interactions, we suggest that ecological risk of any species can be partially predicted by examining its life history and geographic distribution. Thus, we developed a ranking system based on case studies of invasive alien plants in South Korea and other countries to assess ecological risk.

METHODOLOGY

The approach used in this investigation was to assemble published data from South Korea, Japan, United States, Canada, United Kingdom, and Australia. A ranking system to assess an ecological risk of invasive alien plants was developed as follows: (1) listing of invasive plant species in South Korea, (2) surveying and assem-

bling the published data on available invasive plants in South Korea and other countries, (3) isolating and identifying ecological risk factors, (4) positioning a hierarchy of ecological risk factors, and (5) developing a ranking system to grade the ecological risk of invasive alien plants.

The factors of ecological risk were selected, and the hierarchical system was developed to assess ecological risk to manage and monitor invasive alien plants. The ranks of invasive alien plant species were classified by dichotomous method. The ranking system was verified by applying to alien plants in South Korea.

RESULTS AND DISCUSSION

To manage invasive alien plant species effectively, species need to be prioritized based on ecological risk




Figure 3. Examples of assessing an ecological risk of invasive alien plants.

(Hiebert 1997; Hiebert and Stubbendieck 1993; Koh et al. 2002). But many risk-ranking systems consider few factors and emphasize only management and control aspects of invasive alien plants (Heffernan 1998; Hiebert and Stubbendieck 1993; Holm et al. 1991, 1997; Koh et al. 2001). We suggest that it is important also to examine relative probabilities and magnitudes of harm to human health or animal health (Figure 1). Also, invasion time, present state after invasion (dispersion), and availability (valuable to use commercially, economically, and agriculturally) should be considered when assessing ecological risk. Considering these additional risk factors enabled us to rank invasive alien species by a dichotomous classification method (Figure 2). Four grades of ecological risk were suggested: noxious species, monitoring species, innocuous species, and beneficial species. The examples of assessing an ecological risk of invasive plants are shown in Figure 3.

The ranking system was verified by applying to invasive alien plants in South Korea, and they were classified to grades and categories (Table 1). The ranking system can be used to manage and monitor invasive alien

plants. In addition, the ecological risk potential of invaders can be examined by investigating their life-history characteristics as well as geographic and ecological distribution.

LITERATURE CITED

- Coblentz, B. E. 1990. Exotic organisms: a dilemma for conservation biology. Conserv. Biol. 4:261–265.
- De Poorter, M. 2002. Alien species that threaten ecosystems, habitats or species (Agenda item 22). The Sixth Meeting of Conference of the Parties to the Conservation on Biological Diversity. IUCN Information: Web page: http://www.iucn.org/themes/pbia/wl/docs/biodiversity/cop6/invasives.doc. Accessed: October 30, 2002.
- Harper, K. T., R. van Buren, and S. G. Kitchen. 1996. Invasion of alien annuals and ecological consequences in salt desert shrublands of western Utah. *In J. R. Barrow*, E. D. McArthur, R. E. Sosebee, and R. J. Tausch, eds. Shrubland Ecosystem Dynamics in a Changing Environment. Technical Rep. INT-GTR-338. Ogden, UT: Diane Publishing. Pp. 58–65.
- Heffernan, K. E. 1998. Managing Invasive Alien Plants in Natural Areas,
 Parks, and Small Woodlands. Natural Heritage Technical Rep. 98-25.
 Richmond, VA: Virginia Department of Conservation and Recreation,
 Division of Natural Heritage. 17 p.
- Heffernan, K. E., P. P. Coulling, J. F. Townsend, and C. J. Hutto. 2001. Ranking Invasive Exotic Plant Species in Virginia. Natural Heritage Technical Rep. 01-13. Richmond, VA: Virginia Department of Conservation and Recreation, Division of Natural Heritage. 27 p.
- Hiebert, R. D. 1997. Prioritizing invasive plants and planning for management. *In J. O. Luken and J. W. Thieret*, eds. Assessment and Management of Plant Invasions. New York: Springer-Verlag. Pp. 195–212.
- Hiebert, R. D. and J. Stubbendieck. 1993. Handbook for Ranking Exotic Plants for Management and Control. Natural Resources Rep. NPS/NRMWRO/ NRR-93/08. Denver, CO: United States Department of the Interior National Park Service Natural Resources Publication Office. 30 p.
- Hobbs, R. J. and S. E. Humphries. 1995. An integrated approach to management of plant invasions. Conserv. Biol. 9:761–770.
- Holm, L., J. Doll, E. Holm, J. Pancho, and J. Herberger. 1997. World Weeds: Natural Histories and Distribution. New York: J. Wiley. 1129 p.
- Holm, L. G., D. L. Plucknett, J. V. Pancho, and J. P. Herberger. 1991. The World's Worst Weeds: Distribution and Biology. Malabar, FL: Krieger. 609 p.
- Koh, K. S., M. H. Suh, J. H. Kil, Y. B. Ku, H. K. Oh, D. G. Rhee, S. H. Park, E. S. Cheon, and Y. H. Yang. 2002. Research on the Effects of Alien Plants on Ecosystem and Their Management (III). Report of National Institute of Environmental Research, Republic of Korea. 24:121–134.
- Koh, K. S., M. H. Suh, J. H. Kil, Y. B. Ku, H. K. Oh, S. U. Suh, S. H. Park, E. S. Cheon, and Y. H. Yang. 2001. Research on the Effects of Alien Plants on Ecosystem and Their Management (II). Report of National Institute of Environmental Research, Republic of Korea. 23:515–528.
- [OTA] Office of Technology Assessment. 1993. Harmful Non-indigenous Species in the United States. OTA-F-565. Washington, DC: U.S. Government Printing Office.
- Suter, G. 1993. Exotic organisms. *In G. Suter*, ed. Ecological Risk Assessment. Chelsea, MI: Lewis. Pp. 391–401.
- Vitousek, P. M., C. M. D'Antonio, L. L. Loope, and R. Westbrooks. 1996. Biological invasion as global environmental change. Am. Sci. 84:468–478.