

Rule 1: $\text{class } A \text{ extends } B \text{ if } A \text{ is a subclass of } B$
 $CA = CB \rightarrow$

6. releases

Final conclusion

$$[r_1(a, b)] \quad \{z = b\}$$
[illegible]
$$[R_{4/2} - 5] - [R_{6/2} - 5]$$
$$[P_1(C_1)] \quad (2=6)$$

3 - classes

$\{ \text{Rules} \} \Rightarrow ? \text{ or } (yes) \Rightarrow \vee \text{ Success}$

$$L((u,v) \rightarrow z) = \text{rel}((u,v), \text{rel}(z,u)) \wedge \text{rel}(z,v)$$

? closed, 12.5 $(x(1.1) - f_0)$

$$[R_{\alpha} \vee \beta] \rightarrow [R_{\alpha} \vee \beta] \quad (C_5(234)) \quad (Z = F)$$

L 2 - Guss, u

$$f \in R_{\mathbb{A}} \Rightarrow f = \sum_{i=0}^{\infty} a_i x^i$$
$$= (2 + i\sqrt{2}) \rightarrow 2 - i\sqrt{2} \quad (2 + i\sqrt{2}) \cdot (2 - i\sqrt{2}) = 2^2 - (i\sqrt{2})^2 = 4 - (-2) = 4 + 2 = 6$$

$E \{ \text{class} \} \rightarrow \{ \text{class} \}$

$\{rule4\} \rightarrow \{rule3\}$
 $\{rule2\} \rightarrow \{rule1\}$

2)

$$P(X, Y) = P(Y, Z) = \text{Result} = \text{Success}$$

$$P/2 = P/2, X=Y, Y=Z, X=Y=Z$$

$$\text{not put } X=3, Y=3$$

$$Q(X, X) = Q(1, 1) \text{ fail}$$

$$Q/2 = Q/2, X=1, X=2, X \text{ cannot equal 1 and 2}$$

So false

$$M(X, Y) = M(a, Y) \text{ Success}$$

$$M/2 = M/2, f(X) = f(a) \Rightarrow X=a, b=Y \Rightarrow Y=b$$
$$X=a, Y=b = \text{not put}$$

$$K(X, Y) = K(a) \text{ fail}$$

$K(X, Y)$ has arity 2, $K(a)$ has arity 1, function $K/2$ and $K/1$ are different

$$S+(1, 1, 1, \text{empty}, B) = S+(A, B) \text{ Success}$$

$$S/2 = S/2, 1=A, S/1=1, S+(a, \text{empty}) = B \Rightarrow B = S+(a, \text{empty})$$

not put: $A=1, B = S+(a, \text{empty})$