

# IMD0033 - Probabilidade

## Lesson 23 - The Mean

Ivanovitch Silva  
December, 2018



# Agenda

---

- The mean
- The mean as a balance point
- Defining the mean algebraically
- Estimating the population mean
- Estimates from low-sized samples

# Atualizar o repositório


---


```
git clone https://github.com/ivanovitchm/imd0033_2018_2.git
```

Ou ....

```
git pull
```

# PREVIOUSLY ON...





Depending on the particular characteristics of a distribution, we'll see that we can summarize it using the **mean**, the **weighted mean**, the **median**, or the **mode**.

We'll also learn to measure the **variability** in a distribution

$$A = [3, 3, 3, 3]$$


$$B = [30, 1, 15, 43]$$

We can clearly see that there's much more variability (diversity) in B. We'll learn to quantify variability using measures like **variance** and **standard deviation**.


# The Mean

[0, 1, 4, 7, 8, 10]

$$\frac{0 + 1 + 4 + 7 + 8 + 10}{6} = \frac{30}{6} = 5$$



# The Mean as a Balance Point



[0,2,3,3,3,4,13]

Values that are <b>below</b> the mean	Distance from the mean
0	4 units
2	2 units
3	1 unit
3	1 unit
3	1 unit
<hr/>	
Total distance: <b>9 units</b>	

Values that are <b>above</b> the mean	Distance from the mean
13	9 units
<hr/>	
Total distance: <b>9 units</b>	



# Defining the mean algebraically

$$\mu = \frac{x_1 + x_2 + \dots + x_N}{N}$$

$N = 7$   
 $\left[ \begin{array}{cccccc} 0 & 2 & 3 & 3 & 3 & 4 & 13 \end{array} \right]$   
 $\vdots$   
 $x_1 \quad x_2 \quad x_3 \quad x_4 \quad x_5 \quad x_6 \quad x_7$

$$\mu = \frac{0 + 2 + 3 + 3 + 3 + 4 + 13}{7} = \frac{28}{7} = 4$$

**Population Mean**

$$\bar{X} = \frac{x_1 + x_2 + \dots + x_n}{n}$$

$n = 3$   
 $\left[ \begin{array}{ccc} 2 & 3 & 4 \end{array} \right]$   
 $\vdots$   
 $x_1 \quad x_2 \quad x_3$

$$\bar{X} = \frac{2 + 3 + 4}{3} = \frac{9}{3} = 3$$

**Sample Mean**

	Population	Sample
Mean	$\mu$	$\bar{x}, \bar{x}_n, \bar{X}, M$
Number of values	$N$	$n$

# Defining the mean algebraically

---

$$\mu = \frac{x_1 + x_2 + \dots + x_N}{N} = \frac{\sum X}{N} = \frac{\sum_{i=1}^N x_i}{N}$$

**Population Mean**

$$\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{\sum X}{n} = \frac{\sum_{i=1}^n x_i}{n}$$

**Sample Mean**



## Ames, Iowa: Alternative to the Boston Housing Data as an End of Semester Regression Project

[Dean De Cock](#)

Truman State University

*Journal of Statistics Education* Volume 19, Number 3(2011),  
[www.amstat.org/publications/jse/v19n3/decock.pdf](http://www.amstat.org/publications/jse/v19n3/decock.pdf)

Copyright © 2011 by Dean De Cock all rights reserved. This text may be freely shared among individuals, but it may not be republished in any medium without express written consent from the author and advance notification of the editor.

---

**Key Words:** Multiple Regression; Linear Models; Assessed Value; Group Project.

### Abstract

This paper presents a data set describing the sale of individual residential property in Ames, Iowa from 2006 to 2010. The data set contains 2930 observations and a large number of explanatory variables (23 nominal, 23 ordinal, 14 discrete, and 20 continuous) involved in assessing home values. I will discuss my previous use of the Boston Housing Data Set and I will suggest methods for incorporating this new data set as a final project in an undergraduate regression course.



```


1 import pandas as pd
2 pd.set_option('display.max_columns', 500)
3
4 houses = pd.read_csv("AmesHousing_1.txt", sep='\t')
5 houses.shape

```

__	Order	PID	MS SubClass	MS Zoning	Lot Frontage	Lot Area	Street	Alley	Mo Sold	Yr Sold	Sale Type	Sale Condition	SalePrice
0	1	526301100	20	RL	141.0	131770	Pave	0	5	2010	WD	Normal	215000
1	2	526350040	20	RH	80.0	11622	Pave	0	6	2010	WD	Normal	105000
2	3	526351010	20	RL	81.0	14267	Pave	12500	6	2010	WD	Normal	172000
3	4	526353030	20	RL	93.0	11160	Pave	0	4	2010	WD	Normal	244000
4	5	527105010	60	RL	74.0	13830	Pave	0	3	2010	WD	Normal	189900



# Estimating the population mean







---



The general tendency for the sampling error is to decrease as the sample size increases.

# Estimates from low-sized samples





# Next Steps

---

In the next mission, we'll explore a few edge cases where it's either impossible to compute the mean, or it's possible but not theoretically sound.





```
index.js
import React, { useState } from 'react';
import './index.css';

function App() {
  const [contacts, setContacts] = useState([
    { name: 'John Doe', phone: '123-456-7890' },
    { name: 'Jane Smith', phone: '987-654-3210' }
  ]);

  const handleClick = () => {
    // TODO: Add new contact logic
  };

  return (
    <div>
      <h1>Contact List</h1>
      <ul>
        {contacts.map(contact => (
          <li>{contact.name} {contact.phone}</li>
        ))}
      </ul>
      <button onClick={handleClick}>Add New Contact</button>
    </div>
  );
}

export default App;
```

```
index.html
<!DOCTYPE html>
<html>
  <head>
    <script src="index.js"></script>
  </head>
  <body>
    <div></div>
  </body>
</html>
```