



WRFDA Overview

Zhiquan (Jake) Liu

NCAR/MMM

WRFDA is a Data Assimilation system built within the WRF software framework, used for application in both research and operational environments....

Outline

- Basic principal of data assimilation
 - Scalar case
 - Two state variables case
 - General n-dimensional case
- Introduction to WRF Data Assimilation

Outline

- Basic principal of data assimilation
 - Scalar case
 - Two state variables case
 - General n-dimensional case
- Introduction to WRF Data Assimilation

What is data assimilation?

- A **statistical** method to obtain the **best** estimate of **state variables**
- In the atmospheric sciences, DA involves combining **model forecast (prior)** and **observations**, along with their respective errors characterization, to produce an ***analysis (Posterior)*** that can initialize a numerical weather prediction model (e.g., WRF)

Scalar Case

- State variable to estimate “ x ”, e.g., consider today’s temperature of Boulder at 12 UTC.
- Now we have a “background” (or “prior”) information x_b of x , which is from a 6-h GFS or WRF forecast initiated from 06 UTC today.
- We also have an **observation** y of x at a surface station in Boulder
- What is the best estimate (**analysis**) x_a of x ?

Scalar Case

- We can simply average them: $x_a = \frac{1}{2}(x_b + y)$
 - This means we trust equally the background and observation.
- But if their accuracy is different and we have some estimation of their errors
 - e.g., for background, we have statistics (e.g., mean and variance) of $x_b - y$ from the past
 - For observation, we have instrument error information from manufacturer

Scalar Case

- Then we can do a weighted mean: $x_a = ax_b + by$ in a least square sense, i.e.,

- Minimize $J(x) = \frac{1}{2} \frac{(x-x_b)^2}{\sigma_b^2} + \frac{1}{2} \frac{(x-y)^2}{\sigma_o^2}$

- Requires $\frac{dJ(x)}{dx} = \frac{(x-x_b)}{\sigma_b^2} + \frac{(x-y)}{\sigma_o^2} = 0$

- Then we can easily get

$$x_a = \frac{\sigma_o^2}{\sigma_b^2 + \sigma_o^2} x_b + \frac{\sigma_b^2}{\sigma_b^2 + \sigma_o^2} y$$

- We can also write in the form of analysis increment


$$x_a - x_b = \frac{\sigma_b^2}{\sigma_b^2 + \sigma_o^2} (y - x_b)$$

Scalar Case

- Minimize $J(x) = \frac{1}{2} \frac{(x-x_b)^2}{\sigma_b^2} + \frac{1}{2} \frac{(x-y)^2}{\sigma_o^2}$
- Is actually equivalent to maximizing a Gaussian PDF

$$ce^{-J(x)}$$

Assume errors of X_b and y are unbiased



Two state variables case


- Consider two state variables to estimate: Boulder and Denver's temperatures x_1 and x_2 at 12 UTC today.
- Background from 6-h forecast: x_1^b and x_2^b
 - and their error covariance with correlation c

$$\mathbf{B} = \begin{bmatrix} \sigma_1^2 & c\sigma_1\sigma_2 \\ c\sigma_1\sigma_2 & \sigma_2^2 \end{bmatrix}$$

- We only have an observation y_1 at a Boulder station and its error variance σ_o^2

2D PDF

$$PDF(x, y) = \frac{1}{2\pi\sqrt{1-c^2}} \exp\left\{-\frac{1}{2(1-c^2)}(x^2 - 2cxy + y^2)\right\}$$




Analysis increment for two variables

$$x_1^a - x_1^b = \frac{\sigma_1^2}{\sigma_1^2 + \sigma_o^2} (y_1 - x_1^b)$$

$$x_2^a - x_2^b = \frac{c\sigma_1\sigma_2}{\sigma_1^2 + \sigma_o^2} (y_1 - x_1^b)$$

Unobserved variable x_2 gets updated through the error correlation c in the background error covariance.



This correlation can be correlation between two locations (spatial), two variables (multivariate), or two times (temporal).



General Case

Observations
 y^0 , $\sim 10^5\text{-}10^6$

Model state
 x , $\sim 10^7$



General Case: vector and matrix notation

state vector

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{bmatrix}$$

observation vector

$$y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}$$

background error covariance

$$\mathbf{B} = \begin{bmatrix} \sigma_1^2 & c_{12}\sigma_1\sigma_2 & \dots & \dots \\ c_{12}\sigma_1\sigma_2 & \sigma_2^2 & \dots & \dots \\ \dots & \dots & \ddots & \dots \\ \dots & \dots & \dots & \sigma_m^2 \end{bmatrix}$$

Observation error covariance

$$\mathbf{R} = \begin{bmatrix} \sigma_{o1}^2 & 0 & \dots & 0 \\ 0 & \sigma_{o2}^2 & \dots & 0 \\ \vdots & \dots & \ddots & \vdots \\ 0 & \dots & \dots & \sigma_{on}^2 \end{bmatrix}$$

General Case: cost function

$$J(x) = \frac{1}{2}(x - x^b)^T \mathbf{B}^{-1}(x - x^b) + \frac{1}{2}[\mathbf{H}x - y]^T \mathbf{R}^{-1}[\mathbf{H}x - y]$$

\mathbf{H} maps x to y space, e. g., interpolation.

Terminology in DA: **observation operator**

Minimize $J(x)$ is equivalent to maximizing a multi-dimensional Gaussian PDF

$$\text{Constant} * e^{-J(x)}$$

General Case: analytical solution

Again, minimizing J requires its gradient (a vector) with respect to x equal to zero:

$$\nabla J_x(x) = \mathbf{B}^{-1}(x - x_b) - \mathbf{H}^T \mathbf{R}^{-1}[\mathbf{y} - \mathbf{H}x] = 0$$

This leads to analytical solution for the analysis increment:

$$x^a - x^b = \mathbf{B} \mathbf{H}^T (\mathbf{H} \mathbf{B} \mathbf{H}^T + \mathbf{R})^{-1} [\mathbf{y} - \mathbf{H}x^b]$$

$\mathbf{H} \mathbf{B} \mathbf{H}^T$: projection of background error covariance
in observation space


$\mathbf{B} \mathbf{H}^T$: projection of background error covariance
in background-observation space

Iterative algorithm to find minimum of cost function

- **Descending algorithms**

- Descending direction: γ_n (N-dimensional vector)
- Descending step: μ_n

$$x_{n+1} = x_n + \mu_n \gamma_n$$



from Bouttier and Courtier 1999

Precision of Analysis with optimal B and R

$$\mathbf{A}^{-1} = \mathbf{B}^{-1} + \mathbf{H}^T \mathbf{R}^{-1} \mathbf{H}$$

Generalization of scalar case $\frac{1}{\sigma_a^2} = \frac{1}{\sigma_b^2} + \frac{1}{\sigma_o^2}$

Or in another form: $\mathbf{A} = (\mathbf{I} - \mathbf{K}\mathbf{H})\mathbf{B}$

With

$$\mathbf{K} = \mathbf{B}\mathbf{H}^T(\mathbf{H}\mathbf{B}\mathbf{H}^T + \mathbf{R})^{-1}$$

called Kalman gain matrix

Precision of analysis: more general formulation

$$\mathbf{A} = (\mathbf{I} - \mathbf{K}\mathbf{H})\mathbf{B}_t(\mathbf{I} - \mathbf{K}\mathbf{H})^T + \mathbf{K}\mathbf{R}_t\mathbf{K}^T$$

where \mathbf{B}_t and \mathbf{R}_t are “true” background and observation error covariances.

This formulation is valid for any given gain matrix \mathbf{K} , which could be suboptimal (e.g., due to incorrect estimation/specification of \mathbf{B} and \mathbf{R}).

Analysis increment with a single humidity observation


$$x^a - x^b = \mathbf{B}\mathbf{H}^T(\mathbf{H}\mathbf{B}\mathbf{H}^T + \mathbf{R})^{-1}[y - \mathbf{H}x^b]$$

$$x_l^a - x_l^b = \frac{c_{lk}\sigma_l\sigma_k}{\sigma_k^2 + \sigma_{ok}^2}(y_k - x_k^b)$$

It is generalization of previous two variables case:

$$x_1^a - x_1^b = \frac{\sigma_1^2}{\sigma_1^2 + \sigma_o^2}(y_1 - x_1^b)$$


$$x_2^a - x_2^b = \frac{c\sigma_1\sigma_2}{\sigma_1^2 + \sigma_o^2}(y_1 - x_1^b)$$



cv_options=6 in WRFDA

Other Remarks

- Observation operator can be non-linear and thus analysis error PDF is not necessarily Gaussian
- $J(x)$ can have multiple local minima. Final solution of least square depends on starting point of iteration, e.g., choose the background x_b as the first guess.




Other Remarks


- **B** matrix is of very large dimension, explicit inverse of **B** is impossible, substantial efforts in data assimilation were given to the estimation and modeling of **B**.
- **B** shall be spatially-varied and time-evolving according to weather regime.
- Analysis can be sub-optimal if using inaccurate estimate of **B** and **R**.
- Could use non-Gaussian PDF
 - Thus not a least square cost function
 - Difficult (usually slow) to solve; could transform into Gaussian problem via variable transform

Two helpful books

Albert Tarantola



Edwin Thompson Jaynes



<http://www.ipgp.fr/~tarantola/Files/Professional/Books/>

**Probability Theory :
The Logic of Science**


Freely available book!

NUMERICAL RECIPES™

The Art of
Scientific Computing

Third Edition

Click on any image below to display in the right column more information about the product or service.



The book.



The on-line electronic book.



The machine-readable source code (all versions).



NR readers forum.

Numerical Recipes Home Page

We are [nr.com](#), Numerical Recipes Software. We are one of the oldest continuously operating sites on the Web, dating back to 1993 when there were only about 25,000 domains in the Internet. (Today, that number is about 200,000,000.) In partnership with Cambridge University Press, we develop the *Numerical Recipes* series of books on scientific computing and related software products.



Latest News!


We're now on Facebook as [Numerical Recipes Software](#). Give us a "like" or make a post and you might get a free NR3 ebook lifetime subscription! (This is not an official contest -- we're just feeling Facebook-friendly.) Check regularly [at this link](#) to see if you're a winner.

Numerical Recipes in Java™! High-quality translations of our version 3.04 C++ code have been contributed by a Numerical Recipes user. They are

Outline

- Basic principal of data assimilation
 - Scalar case
 - Two state variables case
 - General n-dimensional case
- Introduction to WRF Data Assimilation

WRFDA in the WRF Modeling System



What WRFDA can do?

- Provide Initial conditions for the WRF model forecast
- Verification and validation via difference b.w. obs and model
- Observing system design, monitoring and assessment
- Reanalysis
- Better understanding:
 - Data assimilation methods
 - Model errors
 - Data errors
 - ...

DA algorithms currently available in WRFDA

- 3DVAR and FGAT
 - Different options for choice of control variables (e.g., Psi/Chi or U/V) and background error covariance modeling (e.g., vertical EOF or vertical recursive filter)
- 4DVAR
 - TL/Adjoint (i.e., WRFPlus code) of WRF up-to-date with WRF
 - Allow LBC control variable and Jc-DFI
- Hybrid-3DEnVar and **Hybrid-4DEnVar (since V3.9)**
 - Can run in dual-resolution mode
 - Can ingest ensemble from global or regional sources
- ETKF: for generating ensemble analysis

WRFDA Observations

- **In-Situ:**
 - SYNOP
 - METAR
 - SHIP
 - BUOY
 - TEMP
 - PIBAL
 - AIREP, AIREP humidity
 - TAMDAR
- **Bogus:**
 - TC bogus
 - Global bogus
- **Radiances:**

- HIRS	NOAA-16, NOAA-17, NOAA-18, NOAA-19, METOP-A
- AMSU-A	NOAA-15, NOAA-16, NOAA-18, NOAA-19, EOS-Aqua, METOP-A, METOP-B
- AMSU-B	NOAA-15, NOAA-16, NOAA-17
- MHS	NOAA-18, NOAA-19, METOP-A, METOP-B
- AIRS	EOS-Aqua
- SSMIS	DMSP-16, DMSP-17, DMSP-18
- IASI	METOP-A, METOP-B
- ATMS	Suomi-NPP
- MWTS	FY-3
- MWHS	FY-3
- SEVIRI	METEOSAT
- AMSR2	GCOM-W1 (new in V3.8)
- **Remotely sensed retrievals:**
 - Atmospheric Motion Vectors (geo/polar)
 - SATEM thickness
 - Ground-based GPS **TPW or ZTD**
 - SSM/I oceanic surface wind speed and TPW
 - Scatterometer oceanic surface winds
 - Wind Profiler
 - **Radar data (reflectivity/retrieved rainwater, and radial-wind)**
 - Satellite temperature/humidity/thickness profiles
 - GPS refractivity (e.g. COSMIC)
 - **Stage IV precipitation/rain rate data (4D-Var only)**




WRFDA is flexible to allow assimilation of different formats of observations:

- **Little_r (ascii), HDF, Binary**
- **NOAA MADIS (netcdf),**
- **NCEP PrepBufr,**
- **NCEP radiance bufr**


WRFDA

Radiance Assimilation


- Two RTM interfaces
 - RTTOV or CRTM
- Variational Bias Correction
- Modular code design to ease adding new satellite sensors
- Capability for cloudy radiance DA



New in V3.9: all-sky radiance DA: AMSR2




Channel	Frequency (GHz)	Polarization	Footprint (along scan* along track)
1,2	6.925	V,H	35*61 km
3,4	7.3	V,H	35*61 km
5,6	10.65	V,H	24*41 km
7,8	18.7	V,H	13*22 km
9,10	23.8	V,H	15*26 km
11,12	36.5	V,H	7*12 km
13,14	89.0	V,H	3*5 km




3DVAR (Barker et al. 2004)


$$J(x) = \frac{1}{2}(x - x_b)^T B^{-1}(x - x_b) + \frac{1}{2}[H(x) - y]^T R^{-1}[H(x) - y]$$







4DVAR (Huang et al. 2009)

$$J(\mathbf{x}_0) = \frac{1}{2}(\mathbf{x}_0 - \mathbf{x}_0^b)^T \mathbf{B}^{-1} (\mathbf{x}_0 - \mathbf{x}_0^b) + \frac{1}{2} \sum_{i=1}^N [\mathbf{H}_i(M_i(\mathbf{x}_0)) - \mathbf{y}_i]^T \mathbf{R}_i^{-1} [\mathbf{H}_i(M_i(\mathbf{x}_0)) - \mathbf{y}_i]$$









Radar DA for hydrological application

STEP Hydromet Real Time Exp. during spring time




- The goal is to improve local-scale QPF in coupled hydromet system
- < 1 h rapid update
- Radar radial velocity and reflectivity assimilation
- High resolution vs. ensemble
- Impact of terrain
- Improved results in capturing localized storms





Impact of Aircraft T VarBC on rainfall forecast

(a) TAMDAR coverage on January 15, 2016


| Time Window (hour): -3/+3 |





Mexico station precipitation data 2016.03.08-03.09



GOES imager radiance DA at convection-permitting scale (4km, hourly-cycling, hybrid-3DVAR)




Mexico station precipitation data 2016.01.04-01.05



Yang et al., 2017,
JGR.

24h accumulated rainfall field initialized at 2016071912



Himawari-8 AHI radiance DA impact

Other ongoing work

- Continue developing Multi-Resolution Incremental 4DVAR (MRI-4DVAR)
- Continue developing cloudy radiance/product DA
- High spatial- and temporal-resolution geostationary satellite DA
- Improving surface data assimilation
- Improving radar DA
- WRFPlus-Chem & WRFDA-Chem

Last Remarks

- We welcome contributions from external users/developers.
 - Contact wrfhelp@ucar.edu or directly email to me liuz@ucar.edu for contributing back your code
- We maintain a WRFDA-related publications list, please inform us your papers to be included
 - <http://www2.mmm.ucar.edu/wrf/users/wrfda/publications.html>