Neuron

CellPress
OPEN ACCESS

Correction

Dually innervated dendritic spines develop in the absence of excitatory activity and resist plasticity through tonic inhibitory crosstalk

Mason S. Kleinjan, William C. Buchta, Roberto Ogelman, In-Wook Hwang, Masaaki Kuwajima, Dusten D. Hubbard, Dean J. Kareemo, Olga Prikhodko, Samantha L. Olah, Luis E. Gomez Wulschner, Wickliffe C. Abraham, Santos J. Franco, Kristen M. Harris, Won Chan Oh,* and Matthew J. Kennedy*
*Correspondence: wonchan.oh@cuanschutz.edu (W.C.O.), matthew.kennedy@cuanschutz.edu (M.J.K.)
https://doi.org/10.1016/j.neuron.2023.04.003

(Neuron *111*, 362–371.e1–e6; February 1, 2023)

In the originally published version of the paper, the scale bar in Figure 1G was erroneously labeled as 500 nm when it was, in fact, 377 nm. The corrected figure with 500 nm scale bar is published here, and the paper has been updated online. This change does not affect any conclusions of the manuscript. The authors apologize for the error.




Figure 1G. DiSs are present in all layers of hippocampus (corrected)




Figure 1G. DiSs are present in all layers of hippocampus (original)

This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).