

# Physics 916: Homework #4

Due on April 13, 2020 at 5pm

*Professor Jean Marcel Ngoko*

Robert Tabb

## Problem 1

Consider a physical system whose three-dimensional state space is spanned by an orthonormal basis  $\{|u_1\rangle, |u_2\rangle, |u_3\rangle\}$ . In that state space, consider two operators  $L_z$  and  $S$  defined by:

$$\begin{aligned} L_z |u_1\rangle &= |u_1\rangle, L_z |u_2\rangle = |0\rangle, L_z |u_3\rangle = -|u_3\rangle \\ S |u_1\rangle &= |u_3\rangle, S |u_2\rangle = |u_2\rangle, S |u_3\rangle = |u_1\rangle \end{aligned}$$

(a) Write the matrices which represent, in the  $\{|u_1\rangle, |u_2\rangle, |u_3\rangle\}$  basis, the operators  $L_z, L_z^2, S$ , and  $S^2$ . Are these operators observables?

(b) Give the form of the most general matrix, which represents an operator which commutes with  $L_z$ . Same for  $L_z^2$  and  $S^2$ .

(c) Do  $L_z^2$  and  $S^2$  form a CSCO? Give a basis of common eigenvectors.

### Solution

#### Part a

The matrix representation of these two operators is found by applying to each  $|u_i\rangle$  and simply seeing what each row and column must be to bring about the given transformations. They are:

$$L_z = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad S = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

Then we square them to get the other two matrices:

$$L_z^2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad S^2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

These are all observables because they are Hermitian:

$$\begin{aligned} L_z^\dagger &= L_z, (L_z^2)^\dagger = L_z^2 \\ S^\dagger &= S, (S^2)^\dagger = S^2 \end{aligned}$$

#### Part b

To find the most general matrix,  $A$ , which commutes with  $L_z$ , we need to solve:

$$[L_z, A] = L_z A - A L_z = 0$$

Let's define  $A$  in a general way as:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

Then we have:

$$\begin{aligned}
 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} &= \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} \\
 \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ 0 & 0 & 0 \\ -a_{31} & -a_{32} & -a_{33} \end{pmatrix} &= \begin{pmatrix} a_{11} & 0 & -a_{13} \\ a_{21} & 0 & -a_{23} \\ a_{31} & 0 & -a_{33} \end{pmatrix} \\
 \Rightarrow a_{11} = a_{11}, a_{12} = 0, a_{13} = -a_{13} \\
 a_{21} = 0, a_{22} = a_{22}, a_{23} = 0 \\
 a_{31} = 0, a_{32} = 0, a_{33} = a_{33} \\
 \Rightarrow A = \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{pmatrix}
 \end{aligned}$$

Let's use the same general definition of A to find the general matrix which commutes with  $L_z^2$ :

$$\begin{aligned}
 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} &= \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
 \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ 0 & 0 & 0 \\ a_{31} & a_{32} & a_{33} \end{pmatrix} &= \begin{pmatrix} a_{11} & 0 & a_{13} \\ a_{21} & 0 & a_{23} \\ a_{31} & 0 & a_{33} \end{pmatrix} \\
 \Rightarrow a_{11} = a_{11}, a_{12} = 0, a_{13} = a_{13} \\
 a_{21} = 0, a_{22} = a_{22}, a_{23} = 0 \\
 a_{31} = a_{31}, a_{32} = 0, a_{33} = a_{33} \\
 \Rightarrow A = \begin{pmatrix} a_{11} & 0 & a_{13} \\ 0 & a_{22} & 0 \\ a_{31} & 0 & a_{33} \end{pmatrix}
 \end{aligned}$$

And now for  $S^2$ :

$$\begin{aligned}
 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} &= \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
 \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} &= \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \\
 A &= \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}
 \end{aligned}$$

The commuting matrix for  $L_z$  must be diagonal, for  $L_z^2$  it must have the structure shown above, and for  $S^2$  everything commutes because  $S^2 = I$ , the identity.

### Part c

The first thing to note is that both  $L_z^2$  and  $S^2$  are degenerate. This means that the eigenvalues of either alone cannot fully specify the state of a vector.

Next, I want to see if  $L_z^2$  commutes with  $S^2$ . If you refer back to the matrix representation of  $S^2$  from part a, you'll see that  $S^2 = I$ , where  $I$  is the identity matrix.  $\Rightarrow [S^2, L_z^2] = [I, L_z^2] = 0$ .

But do they form a CSCO? We can quickly find the eigenvectors for  $L_z^2$ . Since the matrix is diagonal, we can simply read off the eigenvalues,  $\lambda = 1, 0$ . Plugging these into the characteristic equation, we get:

For  $\lambda = 0$ :

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\Rightarrow c_1 = 0, c_3 = 0$$

$$|v_{\lambda=0}\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

For  $\lambda = 1$ :

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\Rightarrow c_2 = 0$$

$$|v_{\lambda=1}\rangle = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$$

But we can split  $|v_{\lambda=1}\rangle$  into two orthogonal vectors:

$$|v_{\lambda=1}\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

Renaming the vectors as  $v_1, v_2, v_3$ :

$$|v_1\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, |v_2\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, |v_3\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

Now we have three mutually orthogonal eigenvectors fully specifying the eigenspace. These are easy to check:

$$\begin{aligned} L_z^2 |v_1\rangle &= |v_1\rangle, S^2 |v_1\rangle = |v_1\rangle \\ L_z^2 |v_2\rangle &= |0\rangle, S^2 |v_2\rangle = |v_2\rangle \\ L_z^2 |v_3\rangle &= |v_3\rangle, S^2 |v_3\rangle = |v_3\rangle \end{aligned}$$

This has not gotten rid of our degeneracy. We still have  $\lambda = 1$  for  $v_1$  and  $v_3$ . Thus,  $L_z^2$  and  $S^2$  do not form a CSCO. Also note; the choice of basis vectors here is not unique since  $|v_1\rangle$  and  $|v_3\rangle$  were chosen arbitrarily. There are an infinite number of vectors in the plane spanned by these that could act as orthonormal basis vectors for the eigenspace.

## Problem 2

(a) Consider a linear operator  $A$  in a state space. Show that the trace of this operator is invariant under the change of basis. The trace of a matrix  $A$ , noted as  $Tr\{A\}$ , is defined as the sum of its diagonal elements.

(b) Assume that the eigenvalues,  $a_n$ , of the observable  $A$  are degenerate with a degree  $g_n$ . Calculate  $Tr\{A\}$  in terms of the eigenvalues and the degree of degeneracy.

### Solution

(a) We have a linear operator,  $A$  and a unitary transformation  $U$  which acts to change the basis such that  $|v\rangle = U|v'\rangle$ . Acting on the operator  $A$  with  $U$  is accomplished like this:  $U^\dagger A U$ . So for the trace to remain unchanged, we must have  $Tr\{A\} = Tr\{U^\dagger A U\}$ . Let's rewrite this in tensor notation for convenience using the Einstein summation convention:

$$\begin{aligned} Tr\{A\} &= A_{ii} \\ Tr\{U^\dagger A U\} &= U_{ij}^\dagger A_{jk} U_{ki} \\ &= U_{ki} U_{ij}^\dagger A_{jk} \\ &= (U U^\dagger)_{kj} A_{jk} \end{aligned}$$

Since  $U$  is unitary,  $U^\dagger U = I$ , which in Einstein notation, is the Kronecker delta:

$$\begin{aligned} (U^\dagger U)_{kj} A_{jk} &= \delta_{kj} A_{jk} \\ &= A_{kk} \\ &= Tr\{A\} \end{aligned}$$

(b) I have already shown that the trace is invariant under a change of basis. It is also well known that one can create a matrix through a change of basis such that the diagonals of the observable are its eigenvalues.

$$U^\dagger A U = \begin{pmatrix} a_1 & * & \cdots & * \\ 0 & a_2 & \cdots & * \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_n \end{pmatrix}$$

The trace of  $U^\dagger A U$  is then just the sum of the eigenvalues:

$$Tr\{U^\dagger A U\} = \sum_i^n a_i$$

And since I just showed in part a that the trace of  $A$  is unchanged under a change of basis, we have:

$$\begin{aligned} Tr\{A\} &= Tr\{U^\dagger A U\} \\ &= \sum_i^n a_i \end{aligned}$$

So the trace of a linear operator is the sum of its eigenvalues. But if there are degeneracies present, simply summing the eigenvalues is not enough since any degenerate eigenvalues would be present more than once. The degree of degeneracy,  $g_n$  is the number of times a degenerate eigenvalue is present. Therefore we can simply multiply the degree of degeneracy by the eigenvalues in the trace:

$$Tr\{A\} = \sum_i^n g_i a_i$$

### Problem 3

Let  $|\phi_n\rangle$  be the eigenstates of a Hermitian operator  $H$  ( $H$  is the Hamiltonian of an arbitrary physical system for this problem). Assume that the states,  $|\phi_n\rangle$  form a discrete orthonormal basis. The operator  $U(m, n)$  is defined by:

$$U(m, n) = |\phi_m\rangle \langle \phi_n|$$

- (a) Calculate  $U^\dagger(m, n)$  of  $U(m, n)$ .
- (b) Calculate the commutator,  $[H, U(m, n)]$
- (c) Prove the relation:  $U(m, n)U^\dagger(m, n) = \delta_{nq}(m, p)$
- (d) Calculate  $Tr\{U(m, n)\}$ , the trace of the operator  $U(m, n)$ .
- (e) Let  $A$  be an operator with matrix elements,  $A_{mn} = \langle \phi_m | A | \phi_n \rangle$ . Prove the relation:  $A = \sum_{m,n} A_{mn} U(m, n)$
- (f) Show that  $A_{pq} = Tr\{AU^\dagger(m, n)\}$ .

### Solution

(a)

$$\begin{aligned} U(m, n) &= |\phi_m\rangle \langle \phi_n| \\ U^\dagger(m, n) &= [|\phi_m\rangle \langle \phi_n|]^\dagger \\ &= \langle \phi_n |^\dagger |\phi_m\rangle^\dagger \\ &= |\phi_n\rangle \langle \phi_m| \end{aligned}$$

(b)

$$\begin{aligned} [H, U(m, n)] &= HU(m, n) - U(m, n)H \\ &= H |\phi_m\rangle \langle \phi_n| - |\phi_m\rangle \langle \phi_n| H \end{aligned}$$

Since  $|\phi_n\rangle$  is an eigenstate of  $H$ , we can say that  $H |\phi_n\rangle = E_n |\phi_n\rangle$  and also that the hermitian conjugate of this is also true:  $\langle \phi_n | H = E_n \langle \phi_n |$ . Note that since  $H$  is hermitian,  $H^\dagger = H$ , so in the previous equation I simply left it written as  $H$ . Using this, the commutation relation can be rewritten:

$$\begin{aligned} H |\phi_m\rangle \langle \phi_n| - |\phi_m\rangle \langle \phi_n| H &= E_m |\phi_m\rangle \langle \phi_n| - E_n |\phi_m\rangle \langle \phi_n| \\ &= (E_m - E_n) |\phi_m\rangle \langle \phi_n| \\ &= (E_m - E_n) U(m, n) \end{aligned}$$

(c) Using the definition of  $U^\dagger$  determined in part (a):

$$U(m, n)U^\dagger(m, n) = |\phi_m\rangle \langle \phi_n| |\phi_n\rangle \langle \phi_m|$$

## Problem 4