

# Selected Topics

William Ruth

Université de Montréal

# Topics

- Adaptive Pareto Smoothed Importance Sampling
- Multilevel Causal Mediation Analysis
- Modelling Tuberculosis in Foreign-Born Canadians

- **Adaptive Pareto Smoothed Importance Sampling**
- Multilevel Causal Mediation Analysis
- Modelling Tuberculosis in Foreign-Born Canadians

# Outline

- Importance sampling
- Measuring performance
- Improving performance
  - Modifications
  - Optimization

# Importance Sampling

- Need to compute an expected value
  - $\mathbb{E}_F \varphi(X)$
- Can't do the sum/integral
- Monte Carlo approximation
  - Simulating from  $F$  might be hard

# Importance Sampling

- Introduce “proposal distribution”,  $G$ :

$$\begin{aligned}\mathbb{E}_F \varphi(X) &= \mathbb{E}_G \left[ \varphi(X) \cdot \frac{f(X)}{g(X)} \right] \\ &= \mathbb{E}_G [\varphi(X) \cdot w(X)]\end{aligned}$$


- $G$  can be nearly anything\*
  - \*Some choices will be better than others

# Example: Mystery Target


- $f$  unknown, but can be evaluated
- $\varphi(X) = X^2$
- Try some proposals:
  - $G_1 \sim N(0, 2^2)$
  - $G_2 \sim N(0, 0.6^2)$
- Use  $M = 1000$  samples from proposal
  - $\hat{\mathbb{E}}_1 = 0.99, \hat{SD} = 1.97$
  - $\hat{\mathbb{E}}_2 = 1.10, \hat{SD} = 2.32$

# Example: Mystery Target

$$G_1 = N(0, 2^2)$$



$$G_2 = N(0, 0.5^5)$$






# Importance Sampling

- $G_1$  weights look fine
- $G_2$  weights dominated by one large value
- We can make this difference precise
- “Effective Sample Size”:

$$ESS = \frac{[\sum_i w(X_i)]^2}{\sum_i w(X_i)^2}$$


# Example: Mystery Target

$$G_1 = N(0, 2^2)$$



$$ESS_1 \approx 662$$

$$G_2 = N(0, 0.5^5)$$



$$ESS_2 \approx 54$$

# Importance Sampling

- Problem: Low ESS  $\rightarrow$  hard to estimate means
- But ESS is based on means
  - (Chatterjee and Diaconis, 2018)

# Improving IS


- Choose a good proposal
- Modify large weights
  - Truncated IS
  - Pareto Smoothed IS

# Improving IS


- Truncated Importance Sampling:
  - (Ionides, 2008)
- 1. Choose a threshold
- 2. Apply hard thresholding to any large weights
- Still consistent for the target

# Example: Mystery Target

$$G_1 = N(0, 2^2)$$




$$G_2 = N(0, 0.5^5)$$




# Example: Mystery Target

$$G_1 = N(0, 2^2)$$



$$ESS_1 \approx 662$$
$$ESS_1^{(\text{trunc})} \approx 662$$

$$G_2 = N(0, 0.5^5)$$



$$ESS_2 \approx 54$$
$$ESS_2^{(\text{trunc})} \approx 245$$


# Improving IS

- Pareto Smoothed Importance Sampling:
    - (Vehtari et al., 2024)
1. Choose a threshold
    - Weights above threshold represent tail of their dist.
  2. Approximate tail with Generalized Pareto Dist.
    - Fit GPD to weights above threshold
    - (Zhang and Stephens, 2009)
  3. Replace large weights with quantiles of fitted GPD




# Example: Mystery Target


$$G_1 = N(0, 2^2)$$




$$G_2 = N(0, 0.5^5)$$



# Example: Mystery Target




$$\hat{k}_1 \approx -1.81$$




$$\hat{k}_2 \approx 0.72$$

# Example: Mystery Target

$$G_1 = N(0, 2^2)$$




$$G_2 = N(0, 0.5^5)$$




# Example: Mystery Target


$$G_1 = N(0, 2^2)$$




$$G_2 = N(0, 0.5^5)$$



# Example: Mystery Target



$$\begin{aligned} ESS_1 &\approx 662 \\ ESS_1^{(\text{trunc})} &\approx 662 \\ ESS_1^{(\text{PS})} &\approx 662 \end{aligned}$$



$$\begin{aligned} ESS_2 &\approx 54 \\ ESS_2^{(\text{trunc})} &\approx 245 \\ ESS_2^{(\text{PS})} &\approx 160 \end{aligned}$$

# Adaptive IS

- Alternative approach: directly optimize ESS
  - Adaptive Importance Sampling:
    - (Akyildiz and Míguez, 2021)
1. Choose a (parametric) family of proposals
  2. Iteratively update the proposal to maximize ESS

# Stochastic Approximation

- Actually, we want to maximize a population-level analog:  $\rho \approx \frac{N}{ESS}$
- If we had  $\rho$ , we would do gradient descent
  - $\theta_{k+1} = \theta_k - \alpha \nabla \rho(\theta_k)$
- Instead, do gradient descent on  $\hat{\rho}$ 
  - $\hat{\theta}_{k+1} = \hat{\theta}_k - \alpha_k \nabla \hat{\rho}(\hat{\theta}_k)$

# Stochastic Approximation


- Originally developed for root finding with noise
  - (Robbins and Monro, 1951)
- Quickly adapted for optimization
  - Use noisy evaluations for finite difference
  - (Kiefer and Wolfowitz, 1952)




# Stochastic Approximation

- Very well developed theory
- Step size  $\rightarrow 0$
- Stochastic gradient descent
  - Resample a (very) large dataset


# Example: Mystery Target




# Example: Mystery Target



# Example: Mystery Target



# Example: Mystery Target



# Our Method

- Recall: Be careful using IS means to diagnose IS
- Vehtari et al. give an alternative
  - Shape parameter of fitted tail distribution,  $\hat{k}$

# Our Method

- Use diagnostic as objective function
- Apply stochastic approximation to minimize  $\hat{k}$ 
  - More precisely, its population analog:  $k(\theta)$

# Our Method

- Results: plot trajectories for the same values of  $\sigma$  used above for ESS



# Recap

- Importance sampling and extensions
  - Truncation
  - Pareto Smoothing
- Diagnostics for importance sampling
  - Effective sample size
  - Pareto tail index
- Adaptive importance sampling
  - Stochastic approximation

- Adaptive Pareto Smoothed Importance Sampling
- **Multilevel Causal Mediation Analysis**
- Modelling Tuberculosis in Foreign-Born Canadians

# Topics

- Add slides from my SSC talk

- Adaptive Pareto Smoothed Importance Sampling
- Multilevel Causal Mediation Analysis
- **Modelling Tuberculosis in Foreign-Born Canadians**

# Topics

- Give brief overview and mention directions for future research
- One of the profs in the department, Cristina Anton, does numerical SDEs. Mention potential collaboration

# Acknowledgements

## Collaborators:

- Payman (UBC), Richard (SFU)
- Bouchra (UdeM), Bruno (HEC), Rado (UdeM), Rowin (UdeM)
- Jeremy (SFU), Albert (Langara)

## Funding:

- CANSSI

# Thank You

# Some References

- Akyildiz, Ö. D. and Míguez, J. (2021). Convergence rates for optimized adaptive importance samplers. *Statistics and Computing*, 31(12).
- Chatterjee, S. and Diaconis, P. (2018). The sample size required in importance sampling. *The Annals of Applied Probability*, 28(2).
- Ionides, E. L. (2008). Truncated importance sampling. *Journal of Computational and Graphical Statistics*, 17(2).
- Kiefer, J. and Wolfowitz, J. (1952). Stochastic estimation of the maximum of a regression function. *The Annals of Mathematical Statistics*, 23(3).
- Robbins, H. and Monro, S. (1951). A stochastic approximation method. *The Annals of Mathematical Statistics*, 22(3).
- Vehtari, A., Simpson, D., Gelman, A., Yao, Y., and Gabry, J. (2022). Pareto smoothed importance sampling. *ArXiv*.
- Vehtari, A., Simpson, D., Gelman, A., Yao, Y., and Gabry, J. (2024). Pareto smoothed importance sampling. *Journal of Machine Learning Research*, 25(72).
- Zhang, J. and Stephens, M. A. (2009). A new and efficient estimation method for the generalized Pareto distribution. *Technometrics*, 51(3).