

Corrections for “Solution of the Skyrme–HF+BCS equation on a 3D mesh, II: A new version of the Ev8 code”

Wouter Ryssens*

*Institut d’Astronomie et d’Astrophysique, Université Libre de Bruxelles,
Campus de la Plaine CP 226, BE-1050 Brussels, Belgium*

Michael Bender†

*Université de Lyon, Université Claude Bernard Lyon 1,
CNRS, IP2I Lyon / IN2P3, UMR 5822, F-69622, Villeurbanne, France*

This document is an unfortunately growing set of corrections for the **original** published version of the paper on the Ev8 code, Ref. [1]. Please note that this does not concern any errors in the source code; when found, these errors will be corrected (and documented) in history of this git repository.

1. There is a missing factor of 1/2 in the expression for the direct Coulomb energy, Eq. (43). The correct formula is

$$E_{\text{Coul}}^{\text{d}} = \frac{1}{2} \int d^3r U(\mathbf{r}) \rho_p(\mathbf{r}). \quad (1)$$

2. The equations for the pairing cutoff, Eqs. (51) and (52), are affected by a misprint: neither should have a square root. Additionally, the exponent in the second factor of Eq. (52) has the wrong overall sign. Equations (51) and (52) should read respectively

$$f_k = [1 + e^{(+\epsilon_k - \lambda_q - \Delta\epsilon_q)/\mu_q}]^{-1}, \quad (2)$$

$$f_k = [1 + e^{(+\epsilon_k - \lambda_q - \Delta\epsilon_q)/\mu_q}]^{-1} [1 + e^{(-\epsilon_k + \lambda_q - \Delta\epsilon_q)/\mu_q}]^{-1}. \quad (3)$$

3. Eq.(68) and Eq.(70) of the original paper are affected by an error of notation: all squares should affect *expectation values* instead of *operators*. In addition, there is a missing factor of 2 in Eq.(68), in front of $\langle \hat{Q}_{22} \rangle^2$. Corrected, Eqs.(68-69-70) should read:

$$q = \sqrt{\frac{16\pi}{5} \left(\langle \hat{Q}_{20} \rangle^2 + 2\langle \hat{Q}_{22} \rangle^2 \right)} = \sqrt{\frac{2}{3} \left(\langle \hat{Q}_x \rangle^2 + \langle \hat{Q}_y \rangle^2 + \langle \hat{Q}_z \rangle^2 \right)}, \quad (4)$$

$$\gamma = 2 \arctan \left(\frac{\sqrt{2} \langle \hat{Q}_{22} \rangle}{\sqrt{\langle \hat{Q}_{20} \rangle^2 + 2\langle \hat{Q}_{22} \rangle^2 + \langle \hat{Q}_{20} \rangle}} \right) = 2 \arctan \left(\frac{\langle \hat{Q}_x \rangle - \langle \hat{Q}_y \rangle}{\sqrt{2 \left(\langle \hat{Q}_z \rangle^2 - \langle \hat{Q}_x \rangle \langle \hat{Q}_y \rangle \right) + \langle \hat{Q}_x \rangle - \langle \hat{Q}_y \rangle}} \right). \quad (5)$$

4. Eq. (96) in the original paper contains several errors. Two of these were already corrected in the corrigendum [2]: a global sign and a factor $\frac{\pi^2}{3}$ given in the second line that should not be squared. A third error was not corrected: the equation for the second derivative in Ref. [2] still contains a superfluous factor of two for $i \neq j$. The correct version of Eq. (96) reads:

$$\left. \frac{d^2 f_t(x)}{dx^2} \right|_{x=x_s} = \begin{cases} (-1)^{t-s+1} 2 \left(\frac{\pi}{Ndx} \right)^2 \frac{\cos[\pi(t-s)/N]}{\sin^2[\pi(t-s)/N]} & \text{for } t \neq s, \\ -\frac{\pi^2}{3dx^2} \left(1 - \frac{1}{N^2} \right) & \text{for } t = s. \end{cases} \quad (6)$$

-
- [1] W. Ryssens, V. Hellemans, M. Bender and P.-H. Heenen, *Solution of the Skyrme-HF+BCS equation on a 3D mesh, II: A new version of the Ev8 code*, Computer Physics Communications **187**, 175-194 (2015). DOI:10.1016/j.cpc.2014.10.001.
[2] W. Ryssens, V. Hellemans, M. Bender and P.-H. Heenen, *Corrigendum to “Solution of the Skyrme-HF+BCS equation on a 3D mesh, II: A new version of the Ev8 code”*, Computer Physics Communications **190**, 231 (2015). DOI:10.1016/j.cpc.2015.01.011.

* wryssens@ulb.ac.be

† bender@ip2i.in2p3.fr