


Contents

I	Mathematical Logic and Foundations	1
1	Category Theory	2
1.1	Motivation and Examples	2
1.2	Categories	2
1.3	Objects, Morphisms, and Composition	2
1.4	Identity Morphisms	2
1.5	Commutative Diagrams	2
1.6	Isomorphisms	2
1.7	Monomorphisms and Epimorphisms	3
1.8	Initial and Terminal Objects	3
1.9	Products and Coproducts	3
1.10	Equalizers and Coequalizers	3
1.11	Pullbacks and Pushouts	3
1.12	Functors	3
1.13	Natural Transformations	3
1.14	Equivalence of Categories	3
1.15	Universal Properties	3
1.16	Categories of Mathematical Structures	3
1.17	Category Theory as Language for Algebra, Topology, and Logic	4
1.18	Capstone: Universal Properties in Familiar Mathematics	4
2	Model Theory	9
2.1	Languages and Structures	10
2.1.1	First-Order Languages	10
2.1.1.1	L-Structures	11
2.1.1.2	Variable Assignments	12
2.1.1.3	Term Evaluation	12
2.1.1.4	Satisfaction [Stub]	12
2.1.1.5	Elementary Equivalence and Substructures [Stub]	13
2.2	Satisfaction	13
2.3	Theories and Models	13
2.4	Elementary Equivalence	13
2.5	Substructures and Extensions	13
2.6	Isomorphisms	13
2.7	Compactness	13
2.8	Löwenheim–Skolem	14
2.9	Definability	14
2.10	Categoricity and Nonstandard Models	14

2.11	Types	14
2.12	Saturated Models	14
3	Proof Theory	15
4	Type Theory	19
4.1	Motivation: Types as Classifiers	19
4.2	Typed Syntax	19
4.3	Simple Types	19
4.4	Contexts and Typing Judgments	19
4.5	Formation, Introduction, Elimination, and Computation Rules	19
4.6	Function Types	20
4.7	Product Types	20
4.8	Sum Types	20
4.9	Unit Type and Empty Type	20
4.10	Propositions as Types	20
4.11	Curry–Howard Correspondence	20
4.12	Simply Typed Lambda Calculus	20
4.13	Dependent Types	20
4.14	Dependent Function Types	20
4.15	Dependent Pair Types	20
4.16	Identity Types	20
4.17	Inductive Types	21
4.18	Natural Numbers as an Inductive Type	21
4.19	Type Theory and Proof Assistants	21
4.20	Capstone: Encoding Logical Proofs as Typed Terms	21
5	Lambda Calculus	22
5.1	Motivation: Functions as Computation	23
5.2	Lambda Terms	23
5.3	Variables, Abstraction, and Application	24
5.4	Free and Bound Variables	24
5.5	Substitution	24
5.6	Alpha-Equivalence	24
5.7	Beta-Reduction	25
5.8	Eta-Reduction	25
5.9	Reduction Strategies	25
5.10	Normal Forms	25
5.11	Church Booleans	25
5.12	Church Numerals	25
5.13	Pairs and Lists	25
5.14	Fixed Points and Recursion	25
5.15	The Y-Combinator	25
5.16	Lambda-Definable Functions	25
5.17	Untyped Lambda Calculus and Computability	25
5.18	Simply Typed Lambda Calculus	26
5.19	Strong Normalization	26
5.20	Curry–Howard Connection	26
5.21	Capstone: Encoding Arithmetic in Lambda Calculus	26

Volume VIII

Mathematical Logic and Foundations



Frontispiece placeholder.

David Hilbert, 1862–1943

Volume VIII

Mathematical Logic and Foundations

Self-Study Notes

William Sollers

June 11, 2026

Part I

Mathematical Logic and Foundations

Chapter 1

Category Theory

category-theory

Numerical PDE $\rightarrow$ **Category Theory** $\rightarrow$ Model Theory

Status: Planned

Active mathematical content has not yet been added.

1.1 Motivation and Examples

To be populated.

1.2 Categories

To be populated.

1.3 Objects, Morphisms, and Composition

To be populated.

1.4 Identity Morphisms

To be populated.

1.5 Commutative Diagrams

To be populated.

1.6 Isomorphisms

To be populated.

1.7 Monomorphisms and Epimorphisms

To be populated.

1.8 Initial and Terminal Objects

To be populated.

1.9 Products and Coproducts

To be populated.

1.10 Equalizers and Coequalizers

To be populated.

1.11 Pullbacks and Pushouts

To be populated.

1.12 Functors

To be populated.

1.13 Natural Transformations

To be populated.

1.14 Equivalence of Categories

To be populated.

1.15 Universal Properties

To be populated.

1.16 Categories of Mathematical Structures

To be populated.

1.17 Category Theory as Language for Algebra, Topology, and Logic

To be populated.

1.18 Capstone: Universal Properties in Familiar Mathematics

To be populated.

Category Theory Notes

This chapter is planned. Active mathematical content has not yet been added.

Proofs

Proof entries for this chapter are planned. No proof files have been regenerated.

Exercise Proofs

Capstone Problem

Problem. TODO: state one synthesizing problem for Category Theory, using only concepts at or before this chapter in the registry.

Remark (Dependency ceiling). The capstone may use only results routed at or before this chapter.

Exercises

Exercises for this chapter are planned. No exercise content has been restored here yet.

Chapter 2

Model Theory

model-theory

Category Theory $\rightarrow$ **Model Theory** $\rightarrow$ Proof Theory

Structural Roadmap

Each major topic is organised as:

Definitions $\longrightarrow$ Main Theorems $\longrightarrow$ Consequences and Structural Insight

The global progression is:

1. **First-order languages and $\mathcal{L}$ -structures.** Syntax of languages; domains, function interpretations, relation interpretations, constant interpretations.
2. **Variable assignments and term evaluation.** How terms acquire values in a structure.
3. **The satisfaction relation.** Inductive definition of $\mathcal{M} \models \varphi[s]$; truth and falsity in a structure.
4. **Elementary equivalence and substructures.** $\mathcal{M} \equiv \mathcal{N}$, $\mathcal{N} \prec \mathcal{M}$, the Tarski–Vaught criterion.
5. **Theories and models.** *[Planned]*
6. **Compactness and Löwenheim–Skolem.** *[Planned]*

Remark 2.0.1 (Primary sources). Marker, *Model Theory: An Introduction* (Springer GTM 217), Chapter 1. Enderton, *A Mathematical Introduction to Logic*, Chapter 2. Hodges, *A Shorter Model Theory*, Chapter 1.

2.1 Languages and Structures

L-Structures Toolkit Quick Reference

Concept	Symbol / Notation	Detail
First-order language	$\mathcal{L}$	
L-structure	$\mathcal{M} = (M, \mathcal{L}^{\mathcal{M}})$	
Domain / universe	$ \mathcal{M} $ or M	
Interpretation of f	$f^{\mathcal{M}} : M^n \rightarrow M$	
Interpretation of R	$R^{\mathcal{M}} \subseteq M^n$	
Interpretation of c	$c^{\mathcal{M}} \in M$	
Variable assignment	$s : \text{Var} \rightarrow M$	
Term evaluation	$s(t) \in M$	
Satisfaction	$\mathcal{M} \models \varphi[s]$	

2.1.1 First-Order Languages

Definition (First-Order Language $\mathcal{L}$)

A **first-order language** $\mathcal{L}$ consists of three (possibly empty) sets of non-logical symbols:

- a set $\mathcal{R}$ of **relation symbols** (also called *predicate symbols*), each assigned a fixed arity $n \geq 1$;
- a set $\mathcal{F}$ of **function symbols**, each assigned a fixed **arity** $n \geq 0$ (0-ary function symbols are *constant symbols*);
- a set $\mathcal{C}$ of **constant symbols** (equivalently, 0-ary function symbols listed separately for emphasis).

Together with the logical symbols common to all first-order languages (variables, connectives $\neg, \wedge, \vee, \rightarrow, \leftrightarrow$, quantifiers $\forall, \exists$, equality $=$, parentheses), the triple $(\mathcal{F}, \mathcal{R}, \mathcal{C})$ determines $\mathcal{L}$.

Remark (Fully quantified form). $\mathcal{L} = (\mathcal{F}, \mathcal{R}, \mathcal{C})$ where $\text{ar}(f) \in \mathbb{N}$ for each $f \in \mathcal{F}$ and $\text{ar}(R) \in \mathbb{N}_{\geq 1}$ for each $R \in \mathcal{R}$.

Remark (Intuition). A language is a *vocabulary*: it names the operations, relations, and distinguished elements that can appear in sentences, without yet saying what those names mean. The meaning is supplied by a structure.

Remark (Source note). Some sources (e.g. Marker §1.1, Hodges) absorb constant symbols into 0-ary function symbols. Others (e.g. Chang-Keisler) list them separately. The content is equivalent; only the bookkeeping differs.

Remark (Consequence). The set of $\mathcal{L}$ -terms and the set of $\mathcal{L}$ -formulas are both defined inductively from $\mathcal{L}$; see [term evaluation](#) below.

Example 2.1.1 (Standard examples of first-order languages). 1. **Language of ordered fields** $\mathcal{L}_{\text{of}}$: function symbols $\{+, \cdot, -\}$ of arities 2, 2, 1; relation

symbol $\{<\}$ of arity 2; constant symbols $\{0,1\}$. The real field $(\mathbb{R}, +, \cdot, -, <, 0, 1)$ is an $\mathcal{L}_{\text{of}}$ -structure.

2. **Language of graphs** $\mathcal{L}_{\text{gr}}$: no function symbols, no constants; one binary relation symbol $\{E\}$ (for "edge").
3. **Language of groups** $\mathcal{L}_{\text{gp}}$: function symbols $\{\cdot, ^{-1}\}$ of arities 2,1; constant symbol $\{e\}$.
4. **Pure equality language** $\mathcal{L}_{=}$: no non-logical symbols at all. Structures are arbitrary non-empty sets.

2.1.1.1 L-Structures

Definition ($\mathcal{L}$ -Structure)

Let $\mathcal{L} = (\mathcal{F}, \mathcal{R}, \mathcal{C})$ be a first-order language. An $\mathcal{L}$ -**structure** $\mathcal{M}$ consists of:

1. a non-empty set M called the **domain** (or **universe**) of $\mathcal{M}$, written $|\mathcal{M}|$;
2. for each n -ary relation symbol $R \in \mathcal{R}$, a relation $R^{\mathcal{M}} \subseteq M^n$;
3. for each n -ary function symbol $f \in \mathcal{F}$, a function $f^{\mathcal{M}} : M^n \rightarrow M$;
4. for each constant symbol $c \in \mathcal{C}$, a distinguished element $c^{\mathcal{M}} \in M$.

The superscript notation $f^{\mathcal{M}}, R^{\mathcal{M}}, c^{\mathcal{M}}$ denotes the **interpretation** of each symbol in $\mathcal{M}$.

Remark (Fully quantified form). $\mathcal{M}$ is an $\mathcal{L}$ -structure iff $M \neq \emptyset$, and $\forall f \in \mathcal{F}_{(n)}, f^{\mathcal{M}} : M^n \rightarrow M$, and $\forall R \in \mathcal{R}_{(n)}, R^{\mathcal{M}} \subseteq M^n$, and $\forall c \in \mathcal{C}, c^{\mathcal{M}} \in M$.

Remark (Intuition). A structure is an $\mathcal{L}$ -vocabulary *made concrete*: each symbol is assigned an actual mathematical object of the correct type. The domain provides the raw material; the interpretations give the symbols their meaning inside that domain.

Remark (Common error). The domain M must be *non-empty*. First-order logic with an empty domain is pathological: $\forall x \varphi$ would be vacuously true and $\exists x \varphi$ vacuously false for every φ , breaking the duality of the quantifiers.

Remark (Source note). Some sources write $\mathfrak{M}$ (Fraktur) for structures and reserve M for the domain; others use $\mathcal{M}$ throughout and write $|\mathcal{M}|$ for the domain. The latter convention is used here. Marker uses $\mathcal{M}$; Enderton uses $\mathfrak{A}$.

Example 2.1.2 (L-structures for familiar mathematical objects). 1. $(\mathbb{R}, +^{\mathbb{R}}, \cdot^{\mathbb{R}}, -^{\mathbb{R}}, <^{\mathbb{R}}, 0^{\mathbb{R}}, 1^{\mathbb{R}})$ is an $\mathcal{L}_{\text{of}}$ -structure with domain $M = \mathbb{R}$.

2. $(\mathbb{Q}, +^{\mathbb{Q}}, \cdot^{\mathbb{Q}}, -^{\mathbb{Q}}, <^{\mathbb{Q}}, 0^{\mathbb{Q}}, 1^{\mathbb{Q}})$ is another $\mathcal{L}_{\text{of}}$ -structure with domain $M = \mathbb{Q}$.

3. The complete graph K_3 on $\{1, 2, 3\}$ is an $\mathcal{L}_{\text{gr}}$ -structure with $E^{K_3} = \{(1, 2), (2, 1), (1, 3), (3, 1), (2, 3), (3, 2)\}$.

Two different structures can share the same language the language only fixes the *type* of the interpretations, not their specific content.

2.1.1.2 Variable Assignments

Definition (Variable Assignment)

Let $\mathcal{M}$ be an $\mathcal{L}$ -structure with domain M , and let $\text{Var} = \{v_1, v_2, v_3, \dots\}$ be a fixed countable set of first-order variables.

A **variable assignment** (or **environment**) for $\mathcal{M}$ is a function $s : \text{Var} \rightarrow M$.

For a variable assignment s , a variable v_i , and an element $a \in M$, write $s[v_i \mapsto a]$ for the assignment that agrees with s on all variables except v_i , where it returns a :

$$s[v_i \mapsto a](v_j) = \begin{cases} a & \text{if } j = i, \\ s(v_j) & \text{if } j \neq i. \end{cases}$$

Remark (Fully quantified form). $s : \text{Var} \rightarrow M$, and $s[v_i \mapsto a](v_j) = a$ if $j = i$, else $s(v_j)$.

Remark (Intuition). A variable assignment gives temporary values to free variables before evaluating a formula. The $s[v_i \mapsto a]$ notation is the standard tool for handling quantifier rules: to check $\exists v_i \varphi$, test whether $\mathcal{M} \models \varphi[s[v_i \mapsto a]]$ for some $a \in M$.

2.1.1.3 Term Evaluation

Definition (Term Evaluation)

Let $\mathcal{M}$ be an $\mathcal{L}$ -structure and $s : \text{Var} \rightarrow M$ a variable assignment. The **evaluation** $s(t) \in M$ of an $\mathcal{L}$ -term t under s is defined inductively:

1. If $t = v_i$ is a variable, then $s(t) = s(v_i)$.
2. If $t = c$ is a constant symbol, then $s(t) = c^{\mathcal{M}}$.
3. If $t = f(t_1, \dots, t_n)$ for an n -ary function symbol f , then $s(t) = f^{\mathcal{M}}(s(t_1), \dots, s(t_n))$.

Remark (Fully quantified form). $s : \text{Var} \rightarrow M$, and $s(t) \in M$ for every $\mathcal{L}$ -term t . Formally: $s(\cdot)$ is the unique function on terms satisfying clauses (1)–(3).

Remark (Intuition). Term evaluation is the "computation" step: given concrete values for all variables, every term reduces to an element of M . The inductive definition mirrors the inductive definition of terms themselves.

2.1.1.4 Satisfaction [Stub]

Definition (Satisfaction, $\mathcal{M} \models \varphi[s]$) Stub

Let $\mathcal{M}$ be an $\mathcal{L}$ -structure, s a variable assignment, and φ an $\mathcal{L}$ -formula. The **satisfaction relation** $\mathcal{M} \models \varphi[s]$ (read: " $\mathcal{M}$ satisfies φ under s ") is defined inductively on the structure of φ :

[To be filled in clauses for atomic formulas, negation, conjunction, disjunction, implication, existential quantifier, universal quantifier.]

Remark (Fully quantified form). *[Stub to be completed.]*

Remark (Intuition). Satisfaction is the semantic notion of truth: $\mathcal{M} \models \varphi[s]$ means φ is true in $\mathcal{M}$ when each free variable v_i takes the value $s(v_i)$. The definition proceeds by induction on formula complexity, grounding out on atomic formulas evaluated via term evaluation.

2.1.1.5 Elementary Equivalence and Substructures [Stub]

Status: Planned

The following topics are planned for this section:

1. Substructures ($\mathcal{N} \subseteq \mathcal{M}$).
2. Elementary substructures ($\mathcal{N} \prec \mathcal{M}$) and the Tarski–Vaught criterion.
3. Elementary equivalence ($\mathcal{M} \equiv \mathcal{N}$).
4. Isomorphisms of $\mathcal{L}$ -structures.

2.2 Satisfaction

To be populated.

2.3 Theories and Models

To be populated.

2.4 Elementary Equivalence

To be populated.

2.5 Substructures and Extensions

To be populated.

2.6 Isomorphisms

To be populated.

2.7 Compactness

To be populated.

2.8 Löwenheim-Skolem

To be populated.

2.9 Definability

To be populated.

2.10 Categoricity and Nonstandard Models

To be populated.

2.11 Types

To be populated.

2.12 Saturated Models

To be populated.

Chapter 3

Proof Theory

proof-theory

Model Theory $\rightarrow$ **Proof Theory** $\rightarrow$ Type Theory

Status: Planned

This chapter is planned. Active mathematical content has not yet been added.

Proof Theory Notes

This chapter is planned. Active mathematical content has not yet been added.

Proofs

Proof entries for this chapter are planned. No proof files have been regenerated.

Exercise Proofs

Capstone Problem

Problem. TODO: state one synthesizing problem for Proof Theory, using only concepts at or before this chapter in the registry.

Remark (Dependency ceiling). The capstone may use only results routed at or before this chapter.

Exercises

Exercises for this chapter are planned. No exercise content has been restored here yet.

Chapter 4

Type Theory

type-theory

Proof Theory $\rightarrow$ **Type Theory** $\rightarrow$ Lambda Calculus

Status: Planned

Coming Soon

Notes, proofs, and exercises will appear here in a future revision.

4.1 Motivation: Types as Classifiers

To be populated.

4.2 Typed Syntax

To be populated.

4.3 Simple Types

To be populated.

4.4 Contexts and Typing Judgments

To be populated.

4.5 Formation, Introduction, Elimination, and Computation Rules

To be populated.

4.6 Function Types

To be populated.

4.7 Product Types

To be populated.

4.8 Sum Types

To be populated.

4.9 Unit Type and Empty Type

To be populated.

4.10 Propositions as Types

To be populated.

4.11 Curry-Howard Correspondence

To be populated.

4.12 Simply Typed Lambda Calculus

To be populated.

4.13 Dependent Types

To be populated.

4.14 Dependent Function Types

To be populated.

4.15 Dependent Pair Types

To be populated.

4.16 Identity Types

To be populated.

4.17 Inductive Types

To be populated.

4.18 Natural Numbers as an Inductive Type

To be populated.

4.19 Type Theory and Proof Assistants

To be populated.

4.20 Capstone: Encoding Logical Proofs as Typed Terms

To be populated.

Proofs

To be populated.

Capstone

To be populated.

Chapter 5

Lambda Calculus

lambda-calculus

Type Theory $\rightarrow$ **Lambda Calculus** $\rightarrow$ Algebras of Sets

Chapter Overview

Lambda calculus (λ -calculus) is a formal system for expressing computation through function abstraction and application. Introduced by Alonzo Church (1936) to formalise computability, it is the theoretical foundation of functional programming and is equivalent in power to Turing machines.

Why it belongs here: Lambda calculus sits at the intersection of logic, set theory, and computation. The Curry–Howard correspondence connects it directly to the predicate logic and type theory in this volume: propositions are types, proofs are programs, and implication corresponds to function type.

Primary source: Barendregt & Barendsen, *Introduction to Lambda Calculus* (free PDF).
Secondary: Rojas, *A Tutorial Introduction to the Lambda Calculus*. Bridge to types: Pierce, *Types and Programming Languages* Ch. 5, 8, 9.

Planned Content

1. **Syntax** — terms, variables, abstraction $\lambda x.M$, application MN , free and bound variables
2. **Reduction** — α -conversion, β -reduction, η -reduction, normal forms
3. **Church Encodings** — booleans, natural numbers (Church numerals), pairs, lists
4. **Fixed Points** — the Y-combinator, recursive functions
5. **Computability** — λ -definability, equivalence with Turing machines, undecidability of β -equality
6. **Typed Lambda Calculus** — simple types, type inference, Curry–Howard correspondence

5.1 Motivation: Functions as Computation

To be populated.

5.2 Lambda Terms

Section Toolkit — Quick Reference

Concept	One-line summary
λ -term	Variable x , abstraction $\lambda x.M$, application MN
α -equivalence	$\lambda x.M =_\alpha \lambda y.M[y/x]$ when y fresh
β -reduction	$(\lambda x.M)N \rightarrow_\beta M[N/x]$
Normal form	Term with no β -redex; may not exist (non-termination)
Church boolean T	$\lambda x.\lambda y.x$
Church boolean F	$\lambda x.\lambda y.y$
Church numeral $\bar{n}$	$\lambda f.\lambda x.f^n(x)$ (f applied n times to x)
Type $\tau \rightarrow \sigma$	Function type: input of type τ , output of type σ
Typing judgment	$\Gamma \vdash M : \tau$ reads “ M has type τ in context Γ ”
Curry–Howard	Propositions are types; proofs are terms; $A \rightarrow B$ is implication

Untyped Lambda Calculus

Background References

- **First-order logic:** Enderton [1972], Barwise [1977a].
- **General topology:** Kelly [1955].
- **Set theory:** Halmos [1960], van Dalen et al. [1978].
- **Recursion theory:** Rogers [1967].
- **Category theory:** Arbib and Manes [1975], MacLane [1972].

Definition — Lambda Terms

Definition 5.2.1 (Lambda Terms). The set Λ of λ -terms is defined inductively as follows:

1. Every variable x is a λ -term.
2. If M is a λ -term and x is a variable, then

$$\lambda x.M$$

is a λ -term.

3. If M and N are λ -terms, then

$$MN$$

is a λ -term.

Remark (Interpretation). The λ -terms are the smallest set closed under three constructions: variables, abstraction (function formation), and application (function evaluation). This inductive definition plays the same role as recursive definitions in analysis or algebra: every λ -term is built from variables by finitely many applications of abstraction and application.

Remark (Structural View). Every λ -term has a tree structure:

- variables are leaves,
- abstractions $\lambda x.M$ are unary nodes,
- applications MN are binary nodes.

Thus Λ is a freely generated syntactic algebra.

5.3 Variables, Abstraction, and Application

To be populated.

5.4 Free and Bound Variables

To be populated.

5.5 Substitution

To be populated.

5.6 Alpha-Equivalence

To be populated.

5.7 Beta-Reduction

To be populated.

5.8 Eta-Reduction

To be populated.

5.9 Reduction Strategies

To be populated.

5.10 Normal Forms

To be populated.

5.11 Church Booleans

To be populated.

5.12 Church Numerals

To be populated.

5.13 Pairs and Lists

To be populated.

5.14 Fixed Points and Recursion

To be populated.

5.15 The Y-Combinator

To be populated.

5.16 Lambda-Definable Functions

To be populated.

5.17 Untyped Lambda Calculus and Computability

To be populated.

5.18 Simply Typed Lambda Calculus

To be populated.

5.19 Strong Normalization

To be populated.

5.20 Curry-Howard Connection

To be populated.

5.21 Capstone: Encoding Arithmetic in Lambda Calculus

To be populated.

Chapter 6

Algebras of Sets

algebras-of-sets

Lambda Calculus $\rightarrow$ **Algebras of Sets** $\rightarrow$ Torus Renders

Toolkit — Rings of Sets

Concept family: The ring axioms (Halmos–Givant §1) as the algebraic foundation for Boolean algebras of sets.

Atomic items in this section:

- Definition: Ring

Key predicate:

$\text{Ring}(R, +, \cdot, -, 0, 1) \equiv \text{AbelianGroup}(R, +, -, 0) \wedge \text{Monoid}(R, \cdot, 1) \wedge \text{Distributes}(\cdot, +, R).$

Axiom numbering follows Halmos–Givant; axiom (4) (commutativity of multiplication) is not a ring axiom but a theorem in the Boolean setting.

Definition — Ring

Definition 6.0.1 (Ring). A **ring** is a non-empty set R together with two binary operations $+$ and $\cdot$ on R , a unary operation $-$ on R , and two distinguished elements $0, 1 \in R$, satisfying the following axioms for all $p, q, r \in R$:

- (1) $p + (q + r) = (p + q) + r$ *(additive associativity)*
- (2) $p \cdot (q \cdot r) = (p \cdot q) \cdot r$ *(multiplicative associativity)*
- (3) $p + q = q + p$ *(additive commutativity)*
- (5) $p + 0 = p$ *(additive identity)*
- (6) $p \cdot 1 = p$ *(multiplicative identity)*
- (7) $p + (-p) = 0$ *(additive inverse)*
- (8) $p \cdot (q + r) = p \cdot q + p \cdot r$ *(left distributivity)*
- (9) $(q + r) \cdot p = q \cdot p + r \cdot p$ *(right distributivity)*

Axiom numbering follows Halmos-Givant ?; axiom (4) is commutativity of multiplication, which is not a ring axiom but a theorem in the Boolean setting. A ring possessing a distinguished element 1 satisfying axiom (6) is called a **ring with unit**. A ring satisfying $p \cdot q = q \cdot p$ for all $p, q \in R$ is called a **commutative ring**.

Remark (Standard quantified statement).

$$\begin{aligned}
 (R, +, \cdot, -, 0, 1) \text{ is a ring} &\iff \forall p, q, r \in R: \\
 &\quad p + (q + r) = (p + q) + r \\
 &\quad \wedge \quad p \cdot (q \cdot r) = (p \cdot q) \cdot r \\
 &\quad \wedge \quad p + q = q + p \\
 &\quad \wedge \quad p + 0 = p \\
 &\quad \wedge \quad p \cdot 1 = p \\
 &\quad \wedge \quad p + (-p) = 0 \\
 &\quad \wedge \quad p \cdot (q + r) = p \cdot q + p \cdot r \\
 &\quad \wedge \quad (q + r) \cdot p = q \cdot p + r \cdot p.
 \end{aligned}$$

Remark (Definition predicate reading).

$$\text{Ring}(R, +, \cdot, -, 0, 1) \iff \text{AbelianGroup}(R, +, -, 0) \wedge \text{Monoid}(R, \cdot, 1) \wedge \text{Distributes}(\cdot, +, R)$$

Remark (Negated quantified statement).

$$\begin{aligned}
 (R, +, \cdot, -, 0, 1) \text{ is not a ring} &\iff \exists p, q, r \in R: \\
 &\quad p + (q + r) \neq (p + q) + r \\
 \vee \quad &\quad p \cdot (q \cdot r) \neq (p \cdot q) \cdot r \\
 \vee \quad &\quad p + q \neq q + p \\
 \vee \quad &\quad p + 0 \neq p \\
 \vee \quad &\quad p \cdot 1 \neq p \\
 \vee \quad &\quad p + (-p) \neq 0 \\
 \vee \quad &\quad p \cdot (q + r) \neq p \cdot q + p \cdot r \\
 \vee \quad &\quad (q + r) \cdot p \neq q \cdot p + r \cdot p.
 \end{aligned}$$

Remark (Failure mode decomposition). The eight axioms partition into three structural groups:

$$\underbrace{(1), (3), (5), (7)}_{\text{AbelianGroup}(R, +, -, 0)} \quad \underbrace{(2), (6)}_{\text{Monoid}(R, \cdot, 1)} \quad \underbrace{(8), (9)}_{\text{Distributes}(\cdot, +, R)}$$

A structure violating any axiom in the first group lacks a coherent additive group. A structure violating any axiom in the second group lacks a multiplicative identity or associativity. A structure violating the third group possesses two individually well-behaved operations that do not interact coherently; distributivity is the sole structural bridge between $+$ and $\cdot$, and its failure is sufficient to destroy the ring property.

Remark (Interpretation). A ring abstracts the arithmetic of the integers. The additive structure is always a commutative group. The multiplicative structure is a monoid -- associative with unit -- but need not be commutative; axiom (4) of Halmos-Givant, commutativity of multiplication, is absent from the ring axioms. In the Boolean setting, multiplicative commutativity is not assumed but derived as a theorem from idempotence alone. Distributivity encodes the coherent interaction between the two operations: it is the axiom that makes $\cdot$ and $+$ form a single algebraic system rather than two independent ones.

Concept family: Boolean rings (Halmos–Givant §1) and their immediate consequences from the single idempotence axiom.

Atomic items in this section:

- Definition: Boolean Ring
- Definition: Idempotent Element
- Definition: Characteristic 2
- Lemma: Self Additive Inverse ($p + p = 0$)
- Lemma: Self Negation ($-p = p$)

Key predicate:

$$\text{BooleanRing}(R, +, \cdot, -, 0, 1) \equiv \text{Ring}(R, +, \cdot, -, 0, 1) \wedge \text{IdempotentRing}(R, \cdot).$$

Central theorem chain: Idempotence $\Rightarrow$ Characteristic 2 $\Rightarrow$ Self Negation ($-p = p$) $\Rightarrow$ Commutativity of $\cdot$.

Definition — Boolean Ring

Definition 6.0.2 (Boolean Ring). A **Boolean ring** is an idempotent ring with unit.

That is, a Boolean ring is a ring $(R, +, \cdot, -, 0, 1)$ satisfying the additional axiom:

$$p \cdot p = p \quad \text{for all } p \in R. \quad (11)$$

An element p satisfying $p \cdot p = p$ is called **idempotent**. A ring in which every element is idempotent is called an **idempotent ring**.

The official axiom set for a Boolean ring consists of axioms (1)–(3), (5), (6), (8)–(11), where axiom (7) (the additive inverse law) is replaced by axiom (10) (characteristic 2), which is derived as a theorem. Axiom numbering follows Halmos–Givant ?.

Go to proof that every Boolean ring has characteristic 2.

Remark (Standard quantified statement).

$$\text{BooleanRing}(R, +, \cdot, -, 0, 1) \iff \text{Ring}(R, +, \cdot, -, 0, 1) \wedge \forall p \in R: p \cdot p = p.$$

Remark (Definition predicate reading).

$$\text{BooleanRing}(R, +, \cdot, -, 0, 1) \iff \text{Ring}(R, +, \cdot, -, 0, 1) \wedge \text{IdempotentRing}(R, \cdot)$$

Remark (Negated quantified statement).

$$\neg \text{BooleanRing}(R, +, \cdot, -, 0, 1) \iff \neg \text{Ring}(R, +, \cdot, -, 0, 1) \vee \exists p \in R: p \cdot p \neq p.$$

Remark (Failure mode decomposition). A structure fails to be a Boolean ring

in exactly one of two ways:

$$\underbrace{\neg \text{Ring}(R, +, \cdot, -, 0, 1)}_{\text{fails a ring axiom}} \quad \vee \quad \underbrace{\exists p \in R: p \cdot p \neq p}_{\text{some element is not idempotent}}$$

The second failure mode is the structurally interesting one. A ring in which even one element fails $p \cdot p = p$ is not Boolean. The simplest example: the ring $\mathbb{Z}$ of integers fails idempotence at every element except 0 and 1.

Remark (Canonical examples).

- $\mathbf{2} = \{0, 1\}$ with addition mod 2 and ordinary multiplication is the simplest non-degenerate Boolean ring.
- $\mathbf{2}^n$, the set of all n -tuples of elements of $\mathbf{2}$, with coordinatewise addition and multiplication, is a Boolean ring with 2^n elements. The zero is $(0, \dots, 0)$ and the unit is $(1, \dots, 1)$.
- $(\mathcal{P}(X), \Delta, \cap)$, the power set of any set X with symmetric difference as addition and intersection as multiplication, is a Boolean ring. Under the correspondence $A \leftrightarrow \mathbf{1}_A$ (the characteristic function of A), this ring is isomorphic to $\mathbf{2}^X$.

Remark (Interpretation). A Boolean ring is the minimal algebraic structure that forces the full Boolean calculus to emerge from a single additional axiom -- idempotence of multiplication. The ring axioms supply the additive abelian group and the multiplicative monoid with distributivity. Idempotence then forces characteristic 2 (every element equals its own additive inverse), forces commutativity of multiplication, and identifies the ring with a field of sets via the representation theorem (Halmos-Givant, Chapter 22). The abstract definition thus captures exactly the algebra of set-theoretic operations: symmetric difference plays the role of addition and intersection plays the role of multiplication.

Remark (Dependencies). • **Definition: Ring**

The simplest non-degenerate Boolean ring is the two-element ring $\mathbf{2} = \{0, 1\}$. Its addition and multiplication are given by the following tables.

$+$	0	1	$\cdot$	0	1
0	0	1	0	0	0
1	1	0	1	0	1

Addition in $\mathbf{2}$ is addition modulo 2: $1 + 1 = 0$. Multiplication in $\mathbf{2}$ is ordinary multiplication of integers restricted to $\{0, 1\}$. Every element of $\mathbf{2}$ satisfies $p + p = 0$ and $p \cdot p = p$, making $\mathbf{2}$ the canonical example against which both axioms (10) and (11) are verified.

Definition — Idempotent Element

Definition 6.0.3 (Idempotent Element). An element p of a ring R is called **idempotent** if

$$p \cdot p = p.$$

A ring in which every element is idempotent is called an **idempotent ring**.

Remark (Standard quantified statement).

$$\text{IdempotentElement}(p, R, \cdot) \iff p \cdot p = p.$$

$$\text{IdempotentRing}(R, \cdot) \iff \forall p \in R: p \cdot p = p.$$

Remark (Definition predicate reading).

$$\text{IdempotentElement}(p, R, \cdot) \iff \text{IdempotentRing}(R, \cdot) \iff \forall p \in R: \text{IdempotentElement}(p, R, \cdot)$$

Remark (Negated quantified statement).

$$\neg \text{IdempotentElement}(p, R, \cdot) \iff p \cdot p \neq p.$$

Remark (Interpretation). Idempotence is the single additional axiom that forces Boolean structure. In the two-element ring **2**, verification is immediate: $0 \cdot 0 = 0$ and $1 \cdot 1 = 1$. In a power set algebra $\mathcal{P}(X)$, idempotence of intersection -- $A \cap A = A$ -- is the set-theoretic counterpart. The force of the axiom is not in these examples but in what it implies for an arbitrary ring: characteristic and commutativity of multiplication both follow from idempotence alone, as the theorems below establish.

Remark (Dependencies). • [Definition: Ring](#)

Definition — Characteristic 2

Definition 6.0.4 (Characteristic 2). A ring R is said to have characteristic 2 if

$$p + p = 0 \quad \text{for all } p \in R. \quad (10)$$

Remark (Standard quantified statement).

$$\text{Characteristic2}(R, +, 0) \iff \forall p \in R: p + p = 0.$$

Remark (Definition predicate reading).

$$\text{Characteristic2}(R, +, 0) \iff \forall p \in R: \text{IdempotentElement}(p, R, +)$$

Remark (Negated quantified statement).

$$\neg \text{Characteristic2}(R, +, 0) \iff \exists p \in R: p + p \neq 0.$$

Remark (Interpretation). Characteristic 2 means that every element is its own additive inverse: $p + p = 0$ says precisely $p = -p$. The operation of negation therefore becomes the identity map -- applying $-$ to any element returns that element unchanged. Halmos observes that in a ring of characteristic 2, the minus sign for additive inverses is never needed and is henceforth dropped. Axiom (7), $p + (-p) = 0$, is replaced in the official Boolean ring axiom set by the stronger equation (10), $p + p = 0$, which is a theorem derived from idempotence.

Remark (Dependencies). • [Definition: Boolean Ring](#)

• [Definition: Idempotent Element](#)

Lemma 6.0.5 (Self Additive Inverse). *In any Boolean ring R ,*

$$p + p = 0 \quad \text{for all } p \in R.$$

Go to proof.

Remark (Standard quantified statement).

$$\forall p \in R: p + p = 0.$$

Remark (Definition predicate reading).

$$\text{BooleanRing}(R, +, \cdot, -, 0, 1) \implies \text{Characteristic2}(R, +, 0).$$

Remark (Interpretation). This is Halmos consequence (a): idempotence of multiplication forces the additive structure to have characteristic 2. The proof expands $(p+q)^2$ by distributivity and idempotence, cancels p and q , then sets $p = q$ to obtain $p+p=0$. The Lean 4 formalisation is `AdditiveIdempotence` in `BooleanAlgebra`.

Remark (Dependencies). • [Definition: Boolean Ring](#)

• [Definition: Characteristic 2](#)

Lemma 6.0.6 (Self Negation). *In any Boolean ring R ,*

$$-p = p \quad \text{for all } p \in R.$$

Go to proof.

Remark (Standard quantified statement).

$$\forall p \in R: -p = p.$$

Remark (Interpretation). This follows immediately from [6.0.5](#). Since $p+p=0$ and $p+(-p)=0$ both hold, the uniqueness of additive inverses gives $-p=p$. The negation operation is therefore the identity map on any Boolean ring. Halmos uses this to drop the minus sign entirely from his notation for the rest of the book: since $-p=p$, writing $-p$ and writing p are the same thing.

Remark (Dependencies). • [Lemma: Self Additive Inverse](#)

Toolkit — Boolean Ring Examples

Concept family: Concrete Boolean ring examples and structural constraints.

Atomic items in this section:

- Remark: No Boolean ring of order 3 exists
- Example: The four-element Boolean ring $\mathbf{2}^2$
- Remark: Boolean rings have order 2^n

Key constraint: $(R, +)$ in any Boolean ring is an elementary abelian 2-group, so $|R| = 2^n$ for some $n \geq 0$.

Remark (No Boolean ring of order 3). A Boolean ring of order 3 does not exist. The additive group $(R, +)$ of any Boolean ring satisfies $p + p = 0$ for all p , by Lemma 6.0.5. Every element therefore has additive order dividing 2, so $(R, +)$ is an elementary abelian 2-group. Elementary abelian 2-groups have order 2^n for some $n \geq 0$. Since 3 is not a power of 2, no Boolean ring of order 3 can exist.

Example (The four-element Boolean ring 2^2). The smallest Boolean ring with more than two elements is $2^2 = \{(0,0), (0,1), (1,0), (1,1)\}$, the set of ordered pairs from $2 = \{0,1\}$. Addition and multiplication are defined coordinatewise:

$$(p_0, p_1) + (q_0, q_1) = (p_0 + q_0, p_1 + q_1), \quad (p_0, p_1) \cdot (q_0, q_1) = (p_0 \cdot q_0, p_1 \cdot q_1),$$

where each coordinate is computed in 2 . The zero element is $(0,0)$ and the unit is $(1,1)$.

Addition table.

+	(0,0)	(0,1)	(1,0)	(1,1)
(0,0)	(0,0)	(0,1)	(1,0)	(1,1)
(0,1)	(0,1)	(0,0)	(1,1)	(1,0)
(1,0)	(1,0)	(1,1)	(0,0)	(0,1)
(1,1)	(1,1)	(1,0)	(0,1)	(0,0)

Multiplication table.

.	(0,0)	(0,1)	(1,0)	(1,1)
(0,0)	(0,0)	(0,0)	(0,0)	(0,0)
(0,1)	(0,0)	(0,1)	(0,0)	(0,1)
(1,0)	(0,0)	(0,0)	(1,0)	(1,0)
(1,1)	(0,0)	(0,1)	(1,0)	(1,1)

Verification of idempotence. Every element squares to itself:

p	$p \cdot p$
(0,0)	(0,0)
(0,1)	(0,1)
(1,0)	(1,0)
(1,1)	(1,1)

Verification of characteristic 2. Every element is its own additive inverse:

p	$p + p$
(0,0)	(0,0)
(0,1)	(0,0)
(1,0)	(0,0)
(1,1)	(0,0)

Remark (Boolean rings have order 2^n). Every finite Boolean ring is isomorphic to 2^n for some $n \geq 0$, and therefore has exactly 2^n elements. This follows from the representation theorem (Halmos-Givant, Chapter 22): every Boolean algebra is isomorphic to a field of sets, and every finite field of sets is isomorphic to $\mathcal{P}(X)$ for some finite set X with $|X| = n$, giving $|\mathcal{P}(X)| = 2^n$. The permissible finite sizes are $1, 2, 4, 8, 16, \dots$ and no other cardinality is achievable.

Capstone Problem

Problem. TODO: state one synthesizing problem for Algebras Of Sets, using only concepts at or before this chapter in the registry.

Remark (Dependency ceiling). The capstone may use only results routed at or before this chapter.

Remark (Return). [Return to Definition: Ring](#)

Remark (Note on proof type). This file accompanies the Ring definition rather than a theorem. The "proof" here is a verification that the axiom set is internally consistent (no axiom is derivable from the others in a way that would create circular dependencies) and that the structure as given is non-empty by example. It is not a proof of a theorem but a justification that the definition is well-posed.

Proposition (The Ring Axioms Are Well-Defined). *The axiom set $\{(1), (2), (3), (5), (6), (7)\}$ of Halmos-Givant is non-redundant and consistent: each axiom fails in some structure satisfying all the others, and the integers $(\mathbb{Z}, +, \cdot, -, 0, 1)$ witness consistency.*

Proof. TODO

□

Remark (Return). [Return to Definition: Boolean Ring](#)

Remark (Relationship to existing proof). The proof of this theorem is contained in full in [Self Additive Inverse \(prf:boolean-ring-self-additive-inverse\)](#), which proves $p+p=0$ directly from idempotence. The label `prf:boolean-ring-characteristic-2` is the entry point linked from the Boolean Ring definition; the content is shared.

Theorem (Every Boolean Ring Has Characteristic 2). *In any Boolean ring R ,*

$$p + p = 0 \quad \text{for all } p \in R.$$

Equivalently, $\text{Characteristic2}(R, +, 0)$ holds in every Boolean ring.

Proof. Professional Standard Proof.

Apply idempotence to the element $p + p$:

$$(p + p) \cdot (p + p) = p + p. \tag{i}$$

Expand the left side using right distributivity (9), then left distributivity (8), then collapse each $p \cdot p = p$ by idempotence (11):

$$(p+p) \cdot (p+p) \stackrel{(9)}{=} p \cdot (p+p) + p \cdot (p+p) \stackrel{(8)}{=} (p^2 + p^2) + (p^2 + p^2) \stackrel{p^2=p}{=} (p+p) + (p+p). \tag{ii}$$

Combining (i) and (ii): $(p + p) + (p + p) = p + p$. Setting $s = p + p$, the equation $s + s = s$ gives $s = 0$ by adding $-s$ and applying associativity, the inverse law, and the identity law. Hence $p + p = 0$. $\square$

Remark (Dependencies).

- [Definition: Boolean Ring](#)
- [Definition: Idempotent Element](#)

Remark (Return). [Return to Lemma](#)

Lemma (Self Additive Inverse). *In any Boolean ring R ,*

$$p + p = 0 \quad \text{for all } p \in R.$$

Proof. **Professional Standard Proof.**

Apply idempotence to the element $p + p$:

$$(p + p) \cdot (p + p) = p + p. \quad (\text{i})$$

Expand the left side using right distributivity (9), then left distributivity (8), then collapse each $p \cdot p = p$ by idempotence (11):

$$(p + p) \cdot (p + p) \stackrel{(9)}{=} p \cdot (p + p) + p \cdot (p + p) \stackrel{(8)}{=} (p^2 + p^2) + (p^2 + p^2) \stackrel{p^2=p}{=} (p + p) + (p + p). \quad (\text{ii})$$

Combining (i) and (ii): $(p + p) + (p + p) = p + p$. Setting $s = p + p$, the equation $s + s = s$ gives $s = 0$ by adding $-s$ and applying associativity, the inverse law, and the identity law. Hence $p + p = 0$. $\square$

Proof. **Detailed Learning Proof.**

Step 1. Apply idempotence to the element $p + p$. The element $p + p$ is a member of the carrier of R . Axiom (11) (multiplicative idempotence) asserts every element of R is its own square. Apply this to the element $p + p$:

$$(p + p) \cdot (p + p) = p + p. \quad (\text{i})$$

This is the pivot of the proof. It says $p + p$ squares to itself. Nothing about $p + p = 0$ has been assumed; that is the conclusion to be derived.

Step 2. Expand $(p + p) \cdot (p + p)$ using right distributivity. Axiom (9) (right distributivity) states:

$$(q + r) \cdot t = q \cdot t + r \cdot t \quad \forall q, r, t \in R.$$

Set $q = p$, $r = p$, $t = p + p$:

$$(p + p) \cdot (p + p) = p \cdot (p + p) + p \cdot (p + p). \quad (\text{ii})$$

The product has split into two identical copies of $p \cdot (p + p)$. This step holds in any ring; no idempotence has been used.

Step 3. Expand each $p \cdot (p + p)$ using left distributivity. Axiom (8) (left distributivity) states:

$$t \cdot (q + r) = t \cdot q + t \cdot r \quad \forall t, q, r \in R.$$

Apply to $p \cdot (p + p)$ with $t = p$, $q = p$, $r = p$:

$$p \cdot (p + p) = p \cdot p + p \cdot p.$$

Apply to both copies in (ii):

$$p \cdot (p + p) + p \cdot (p + p) = (p \cdot p + p \cdot p) + (p \cdot p + p \cdot p). \quad (\text{iii})$$

The right side contains four occurrences of $p \cdot p$. This expansion holds in any ring; idempotence has still not been used.

Step 4. Collapse each $p \cdot p$ to p by idempotence. Axiom (11) gives $p \cdot p = p$. Substitute into all four occurrences in (iii):

$$(p \cdot p + p \cdot p) + (p \cdot p + p \cdot p) = (p + p) + (p + p). \quad (\text{iv})$$

This is the second and final use of idempotence.

Step 5. Chain (i) through (iv). Equations (i), (ii), (iii), (iv) combine as:

$$p + p = (p + p) \cdot (p + p) = (p + p) + (p + p).$$

Hence

$$(p + p) + (p + p) = p + p. \quad (\text{v})$$

Step 6. Conclude $p + p = 0$ by cancellation. Let $s = p + p$. Equation (v) states $s + s = s$. Add the additive inverse $-s$ to the right of both sides, then apply associativity, the inverse law, and the identity law:

$$s + s = s \implies (s + s) + (-s) = s + (-s) \implies s + \underbrace{(s + (-s))}_{=0} = 0 \implies s + 0 = 0 \implies s = 0.$$

Since $s = p + p$, the conclusion is $p + p = 0$. □

Remark (Proof structure). The proof is direct. The strategy applies idempotence twice: once to $p + p$ as a whole (Step 1) and once to p individually (Step 4), with distributivity (Steps 2-3) connecting the two applications. The conclusion follows by a standard additive cancellation (Step 6) using only the group axioms for $(R, +)$. No circularity arises: the proof never assumes $1 + 1 = 0$ or any consequence of characteristic 2. Both $p + p = 0$ and $1 + 1 = 0$ are consequences of idempotence; this proof derives the former directly without appealing to the latter. The Lean 4 formalisation is `AdditiveIdempotence` in `LRA/VolumeI/BooleanAl`.

Remark (Dependencies).

- [Definition of Ring](#) -- axioms (1) associativity, (5) identity, (7) inverse, (8) left distributivity, (9) right distributivity.
- [Definition of Boolean Ring](#) -- axiom (11) multiplicative idempotence, applied in Steps 1 and 4.
- [Definition of Idempotent Element](#) -- names the property $p \cdot p = p$.

Remark (Return). [Return to Lemma](#)

Lemma (Self Negation). *In any Boolean ring R ,*

$$-p = p \quad \text{for all } p \in R.$$

Proof. Professional Standard Proof.

By [Self Additive Inverse](#), $p + p = 0$. By the additive inverse axiom (7), $p + (-p) = 0$. Since both p and $-p$ satisfy the equation $p + x = 0$, and additive inverses are unique in any group (a standard group theory fact applied to the abelian group $(R, +)$), we conclude $-p = p$. $\square$

Proof. Detailed Learning Proof.

Step 1. Recall the two equations that both hold in R .

From Self Additive Inverse ([Lemma 6.0.5](#)):

$$p + p = 0. \tag{i}$$

From the additive inverse axiom (7) of the ring definition:

$$p + (-p) = 0. \tag{ii}$$

Step 2. Apply uniqueness of additive inverses.

The additive group $(R, +, 0)$ is a group (abelian, by axiom (3)). In any group, if $a + b = 0$ and $a + c = 0$, then:

$$b = 0 + b = (a + c) + b \stackrel{\text{assoc}}{=} a + (c + b) \stackrel{\text{comm}}{=} a + (b + c) \stackrel{\text{assoc}}{=} (a + b) + c = 0 + c = c.$$

Apply with $a = p$, $b = p$ (from (i)) and $c = -p$ (from (ii)): both p and $-p$ are additive inverses of p , so $p = -p$. $\square$

Remark (Proof structure). A one-step uniqueness argument. Self Additive Inverse shows p is its own additive inverse; the ring axiom asserts $-p$ is also an additive inverse of p ; uniqueness of inverses in the underlying group forces $p = -p$. The proof has no calculation content -- it is purely structural.

Remark (Dependencies).

- [Lemma: Self Additive Inverse](#)
- [Definition: Ring](#) (axiom (7), additive inverse law)

Remark (Return). [Return to Theorem \(Self Additive Inverse\)](#).

Theorem (Self Additive Inverse). *In any Boolean ring R ,*

$$p + p = 0 \quad \text{for all } p \in R.$$

Proof. **Professional Standard Proof.** TODO: professional standard proof for self-additive-inverse. □

Proof. **Detailed Learning Proof.** TODO: detailed learning proof for self-additive-inverse. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Bibliography