


FROM CANTOR TO ITO

# Origins of Numbers

## The Continuum



Karl Weierstrass

1815-1897

*Self-Study Notes*

William Sollers

July 1, 2026

*For every reader learning to make mathematics their own.*

# Contents

|          |                                                             |            |
|----------|-------------------------------------------------------------|------------|
| <b>1</b> | <b>Rational Numbers (<math>\mathbb{Q}</math>)</b>           | <b>1</b>   |
| 1.1      | Construction of $\mathbb{Q}$ . . . . .                      | 1          |
| 1.2      | Operations on $\mathbb{Q}$ -Classes . . . . .               | 6          |
| 1.3      | Field Structure of $\mathbb{Q}$ . . . . .                   | 18         |
| 1.4      | Order on $\mathbb{Q}$ . . . . .                             | 22         |
| 1.5      | Ordered Field Structure of $\mathbb{Q}$ . . . . .           | 32         |
| 1.6      | Arbitrary Closeness in $\mathbb{Q}$ . . . . .               | 34         |
| 1.7      | Rational Gaps . . . . .                                     | 37         |
| 1.8      | Embedding $\mathbb{Z}$ into $\mathbb{Q}$ . . . . .          | 39         |
| 1.9      | Transition to $\mathbb{R}$ . . . . .                        | 42         |
| 1.10     | Canonical Rational Number Statement Plan . . . . .          | 42         |
| <b>2</b> | <b>Real Numbers (<math>\mathbb{R}</math>)</b>               | <b>178</b> |
| 2.1      | Construction of $\mathbb{R}$ . . . . .                      | 178        |
| 2.2      | Order on $\mathbb{R}$ . . . . .                             | 180        |
| 2.3      | Completeness of $\mathbb{R}$ . . . . .                      | 183        |
| 2.4      | Addition on $\mathbb{R}$ . . . . .                          | 184        |
| 2.5      | Multiplication on $\mathbb{R}$ . . . . .                    | 187        |
| 2.6      | Field and Ordered-Field Structure of $\mathbb{R}$ . . . . . | 192        |
| 2.7      | Embedding $\mathbb{Q}$ into $\mathbb{R}$ . . . . .          | 194        |
| 2.8      | The Gap Filled . . . . .                                    | 196        |
| 2.9      | Characterization of $\mathbb{R}$ . . . . .                  | 197        |
| 2.10     | Transition and Forward . . . . .                            | 198        |
| <b>3</b> | <b>Complex Numbers</b>                                      | <b>249</b> |
| 3.1      | Construction of $\mathbb{C}$ . . . . .                      | 249        |
| 3.2      | Operations on $\mathbb{C}$ -Classes . . . . .               | 252        |
| 3.3      | Field Structure of $\mathbb{C}$ . . . . .                   | 256        |
| 3.4      | Embedding $\mathbb{R}$ into $\mathbb{C}$ . . . . .          | 258        |
| <b>4</b> | <b>Number Lines and Intervals</b>                           | <b>291</b> |
| 4.1      | Number Lines . . . . .                                      | 291        |
| 4.2      | Intervals on a Number Line . . . . .                        | 292        |

|          |                                                                                                          |            |
|----------|----------------------------------------------------------------------------------------------------------|------------|
| <b>5</b> | <b>Embedding the Number Systems</b>                                                                      | <b>294</b> |
| 5.1      | Embeddings as Structure Preserving Maps . . . . .                                                        | 294        |
| 5.2      | Embedding $\mathbb{N}$ Into $\mathbb{W}$ . . . . .                                                       | 297        |
| 5.3      | Embedding $\mathbb{W}$ Into $\mathbb{Z}$ . . . . .                                                       | 298        |
| 5.4      | Embedding $\mathbb{Z}$ Into $\mathbb{Q}$ . . . . .                                                       | 299        |
| 5.5      | Embedding $\mathbb{Q}$ Into $\mathbb{R}$ . . . . .                                                       | 300        |
| 5.6      | Compatibility of Arithmetic and Order . . . . .                                                          | 301        |
| 5.7      | One Chain Many Structures . . . . .                                                                      | 302        |
| <b>6</b> | <b>Summary: The Number Systems</b>                                                                       | <b>310</b> |
| 6.1      | Axiom Systems for $\mathbb{N}$ , $\mathbb{W}$ , $\mathbb{Z}$ , $\mathbb{Q}$ , and $\mathbb{R}$ . . . . . | 310        |
| 6.2      | Number Systems Summary . . . . .                                                                         | 311        |

# Chapter 1

## Rational Numbers ( $\mathbb{Q}$ )



### 1.1 Construction of $\mathbb{Q}$

#### Prefractions

##### Definition

**Definition 1.1.1** (Nonzero Integers). The set of nonzero integers is

$$\mathbb{Z}^* := \mathbb{Z} \setminus \{0_{\mathbb{Z}}\}.$$

*Remark* (Dependencies).

- [Integers](#)
- [Zero in  \$\mathbb{Z}\$](#)

##### Definition

**Definition 1.1.2** (Prefraction). A *prefraction* is an ordered pair  $(a, b) \in \mathbb{Z} \times \mathbb{Z}^*$ , read as the formal quotient  $a/b$ .

*Remark* (Dependencies).

- [Ordered Pair](#)
- [Integers](#)
- [Nonzero Integers](#)

### Definition

**Definition 1.1.3** (Prefraction Set). The prefraction set is

$$\text{Pre } \mathbb{Q} := \mathbb{Z} \times \mathbb{Z}^*.$$

*Remark* (Dependencies).

- [Cartesian Product](#)
- [Integers](#)
- [Nonzero Integers](#)

## Fraction Equivalence

### Definition

**Definition 1.1.4** (Fraction Equivalence). For  $(a, b), (c, d) \in \text{Pre } \mathbb{Q}$ , define

$$(a, b) \sim_{\mathbb{Q}} (c, d) \quad :\Longleftrightarrow \quad a \cdot d = b \cdot c.$$

*Remark* (Dependencies).

- [Relation](#)
- [Prefraction Set](#)
- [Multiplication on  \$\mathbb{Z}\$](#)

### Lemma

**Lemma 1.1.5** (Fraction Equivalence Is Reflexive). *For every  $(a, b) \in \text{Pre } \mathbb{Q}$ ,*

$$(a, b) \sim_{\mathbb{Q}} (a, b).$$

*Remark* (Dependencies).

- [Fraction Equivalence](#)
- [Multiplication on  \$\mathbb{Z}\$  Is Commutative](#)

### Lemma

**Lemma 1.1.6** (Fraction Equivalence Is Symmetric). *For all prefractions  $(a, b), (c, d)$ ,*

$$(a, b) \sim_{\mathbb{Q}} (c, d) \implies (c, d) \sim_{\mathbb{Q}} (a, b).$$

*Remark* (Dependencies).

- [Fraction Equivalence](#)

- [Multiplication on  \$\mathbb{Z}\$  Is Commutative](#)

#### Lemma

**Lemma 1.1.7** (Fraction Equivalence Is Transitive). *For all prefractions  $(a, b), (c, d), (e, f)$ ,*

$$(a, b) \sim_{\mathbb{Q}} (c, d) \wedge (c, d) \sim_{\mathbb{Q}} (e, f) \implies (a, b) \sim_{\mathbb{Q}} (e, f).$$

*Remark* (Dependencies).

- [Multiplicative Cancellation on  \$\mathbb{Z}\$](#)

#### Theorem

**Theorem 1.1.8** (Fraction Equivalence Is an Equivalence Relation). *The relation  $\sim_{\mathbb{Q}}$  is an equivalence relation on  $\text{Pre } \mathbb{Q}$ .*

*Remark* (Dependencies).

- [Fraction Equivalence](#)
- [Fraction Equivalence Is Reflexive](#)
- [Fraction Equivalence Is Symmetric](#)
- [Fraction Equivalence Is Transitive](#)

## Definition of $\mathbb{Q}$

#### Definition

**Definition 1.1.9** (Rational Numbers). The rational numbers are the quotient

$$\mathbb{Q} := \text{Pre } \mathbb{Q} / \sim_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [Quotient Set](#)
- [Prefraction Set](#)
- [Fraction Equivalence](#)

#### Definition

**Definition 1.1.10** (Rational Class Notation). The notation  $[a, b]$  denotes the  $\sim_{\mathbb{Q}}$ -equivalence class of the prefraction  $(a, b)$ , where  $b \neq 0_{\mathbb{Z}}$ .

*Remark* (Dependencies).



- [Rational Numbers](#)
- [Fraction Equivalence](#)

#### Proposition

**Proposition 1.1.11** (Equality Criterion for Rational Classes). *For prefractions  $(a, b), (c, d) \in \text{Pre } \mathbb{Q}$ ,*

$$[a, b] = [c, d] \iff a \cdot d = b \cdot c.$$

*Remark* (Dependencies).

- [Rational Class Notation](#)
- [Fraction Equivalence](#)
- [Quotient Set](#)
- [Equivalence Class](#)

## Zero and One in $\mathbb{Q}$

#### Definition

**Definition 1.1.12** (Zero in  $\mathbb{Q}$ ). Define

$$0_{\mathbb{Q}} := [0_{\mathbb{Z}}, 1_{\mathbb{Z}}].$$

*Remark* (Dependencies).

- [Rational Class Notation](#)
- [Zero in  \$\mathbb{Z}\$](#)
- [One in  \$\mathbb{Z}\$](#)

#### Definition

**Definition 1.1.13** (One in  $\mathbb{Q}$ ). Define

$$1_{\mathbb{Q}} := [1_{\mathbb{Z}}, 1_{\mathbb{Z}}].$$

*Remark* (Dependencies).

- [Rational Class Notation](#)
- [One in  \$\mathbb{Z}\$](#)

### Lemma

**Lemma 1.1.14** (Vanishing Criterion in  $\mathbb{Q}$ ). *For every prefraction  $(a, b) \in \text{Pre } \mathbb{Q}$ ,*

$$[a, b] = 0_{\mathbb{Q}} \iff a = 0_{\mathbb{Z}}.$$

*Remark* (Dependencies).

- [Rational Class Notation](#)
- [Fraction Equivalence](#)
- [Zero in  \$\mathbb{Q}\$](#)
- [Zero in  \$\mathbb{Z}\$](#)

### Theorem

**Theorem 1.1.15** (Zero Is a Rational). *The class  $0_{\mathbb{Q}}$  is an element of  $\mathbb{Q}$ .*

*Remark* (Dependencies).

- [Rational Numbers](#)
- [Zero in  \$\mathbb{Q}\$](#)
- [Rational Class Notation](#)

### Theorem

**Theorem 1.1.16** (One Is a Rational). *The class  $1_{\mathbb{Q}}$  is an element of  $\mathbb{Q}$ .*

*Remark* (Dependencies).

- [Rational Numbers](#)
- [One in  \$\mathbb{Q}\$](#)
- [Rational Class Notation](#)

### Theorem

**Theorem 1.1.17** (Zero Is Not One in  $\mathbb{Q}$ ). *In  $\mathbb{Q}$ ,*

$$0_{\mathbb{Q}} \neq 1_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [Zero Is Not One in  \$\mathbb{Z}\$](#)

## 1.2 Operations on $\mathbb{Q}$ -Classes

### Addition on $\mathbb{Q}$ -Classes

#### Definition

**Definition 1.2.1** (Addition on  $\mathbb{Q}$ -Classes). Define addition on representatives by

$$[a, b] + [c, d] := [a \cdot d + b \cdot c, b \cdot d].$$

*Remark* (Dependencies).

- [Rational Class Notation](#)
- [Addition on  \$\mathbb{Z}\$](#)
- [Multiplication on  \$\mathbb{Z}\$](#)

#### Lemma

**Lemma 1.2.2** (Addition Lands in  $\text{Pre } \mathbb{Q}$ ). If  $b \neq 0_{\mathbb{Z}}$  and  $d \neq 0_{\mathbb{Z}}$ , then

$$b \cdot d \neq 0_{\mathbb{Z}}.$$

*Remark* (Dependencies).

- [\$\mathbb{Z}\$  Has No Zero Divisors](#)

#### Lemma

**Lemma 1.2.3** (Addition Respects Fraction Equivalence on the Left). If  $[a, b] = [a', b']$ , then

$$[a, b] + [c, d] = [a', b'] + [c, d].$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{Q}\$](#)
- [Fraction Equivalence](#)
- [Equality Criterion for Rational Classes](#)
- [Multiplicative Cancellation on  \$\mathbb{Z}\$](#)

#### Lemma

**Lemma 1.2.4** (Addition Respects Fraction Equivalence on the Right). If  $[c, d] = [c', d']$ , then

$$[a, b] + [c, d] = [a, b] + [c', d'].$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{Q}\$](#)
- [Fraction Equivalence](#)
- [Equality Criterion for Rational Classes](#)
- [Multiplicative Cancellation on  \$\mathbb{Z}\$](#)

#### Lemma

**Lemma 1.2.5** (Addition Respects Fraction Equivalence). *If  $[a, b] = [a', b']$  and  $[c, d] = [c', d']$ , then*

$$[a, b] + [c, d] = [a', b'] + [c', d'].$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{Q}\$](#)
- [Addition Respects Fraction Equivalence on the Left](#)
- [Addition Respects Fraction Equivalence on the Right](#)

#### Theorem

**Theorem 1.2.6** (Addition on  $\mathbb{Q}$  Is Well-Defined). *Addition determines a well-defined binary operation  $\mathbb{Q} \times \mathbb{Q} \rightarrow \mathbb{Q}$ .*

*Remark* (Dependencies).

- [Addition on  \$\mathbb{Q}\$](#)
- [Addition Respects Fraction Equivalence](#)

#### Theorem

**Theorem 1.2.7** (Addition on  $\mathbb{Q}$  Is Associative). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$(x + y) + z = x + (y + z).$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Addition on  \$\mathbb{Z}\$  Is Associative](#)
- [Multiplication on  \$\mathbb{Z}\$  Is Associative](#)
- [Multiplication Distributes over Addition on  \$\mathbb{Z}\$](#)

### Theorem

**Theorem 1.2.8** (Addition on  $\mathbb{Q}$  Is Commutative). *For all  $x, y \in \mathbb{Q}$ ,*

$$x + y = y + x.$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Addition on  \$\mathbb{Z}\$  Is Commutative](#)

### Theorem

**Theorem 1.2.9** (Zero Is a Left Additive Identity on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} + x = x.$$

*Remark* (Dependencies).

- [Zero in  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Zero Is an Additive Identity on  \$\mathbb{Z}\$](#)
- [One Is a Multiplicative Identity on  \$\mathbb{Z}\$](#)

### Corollary

**Corollary 1.2.10** (Zero Is a Right Additive Identity on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$x + 0_{\mathbb{Q}} = x.$$

*Remark* (Dependencies).

- [Zero Is a Left Additive Identity on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$  Is Commutative](#)

### Theorem

**Theorem 1.2.11** (Zero Is an Additive Identity on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} + x = x \quad \text{and} \quad x + 0_{\mathbb{Q}} = x.$$

*Remark* (Dependencies).

- [Zero Is a Left Additive Identity on  \$\mathbb{Q}\$](#)
- [Zero Is a Right Additive Identity on  \$\mathbb{Q}\$](#)

## Negation on $\mathbb{Q}$ -Classes

### Definition

**Definition 1.2.12** (Negation on  $\mathbb{Q}$ -Classes). Define negation on representatives by

$$-[a, b] := [-a, b].$$

*Remark* (Dependencies).

- [Rational Class Notation](#)
- [Negation on  \$\mathbb{Z}\$](#)

### Lemma

**Lemma 1.2.13** (Negation Respects Fraction Equivalence). *If  $[a, b] = [a', b']$ , then*

$$-[a, b] = -[a', b'].$$

*Remark* (Dependencies).

- [Negation on  \$\mathbb{Q}\$](#)
- [Fraction Equivalence](#)
- [Sign Rule on  \$\mathbb{Z}\$](#)

### Theorem

**Theorem 1.2.14** (Negation on  $\mathbb{Q}$  Is Well-Defined). *Negation determines a well-defined unary operation  $\mathbb{Q} \rightarrow \mathbb{Q}$ .*

*Remark* (Dependencies).

- [Negation on  \$\mathbb{Q}\$](#)
- [Negation Respects Fraction Equivalence](#)

### Theorem

**Theorem 1.2.15** (Negation Gives a Left Additive Inverse on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$(-x) + x = 0_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [Negation on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$](#)

- [Negation on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Addition on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Negation Gives a Left Additive Inverse on  \$\mathbb{Z}\$](#)

#### Corollary

**Corollary 1.2.16** (Negation Gives a Right Additive Inverse on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$x + (-x) = 0_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [Negation Gives a Left Additive Inverse on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$  Is Commutative](#)

#### Theorem

**Theorem 1.2.17** (Every Rational Has an Additive Inverse). *For every  $x \in \mathbb{Q}$  there exists  $y \in \mathbb{Q}$  such that*

$$x + y = 0_{\mathbb{Q}} \quad \text{and} \quad y + x = 0_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [Negation on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Negation Gives a Left Additive Inverse on  \$\mathbb{Q}\$](#)
- [Negation Gives a Right Additive Inverse on  \$\mathbb{Q}\$](#)

## Subtraction on $\mathbb{Q}$

#### Definition

**Definition 1.2.18** (Subtraction on  $\mathbb{Q}$ ). For  $x, y \in \mathbb{Q}$ , define

$$x - y := x + (-y).$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{Q}\$](#)
- [Negation on  \$\mathbb{Q}\$](#)

#### Theorem

**Theorem 1.2.19** (Subtraction on  $\mathbb{Q}$  Is Well-Defined). *Subtraction determines a well-defined binary operation  $\mathbb{Q} \times \mathbb{Q} \rightarrow \mathbb{Q}$ .*

*Remark* (Dependencies).

- [Subtraction on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Negation on  \$\mathbb{Q}\$  Is Well-Defined](#)

## Multiplication on $\mathbb{Q}$ -Classes

### Definition

**Definition 1.2.20** (Multiplication on  $\mathbb{Q}$ -Classes). Define multiplication on representatives by

$$[a, b] \cdot [c, d] := [a \cdot c, b \cdot d].$$

*Remark* (Dependencies).

- [Rational Class Notation](#)
- [Multiplication on  \$\mathbb{Z}\$](#)

### Lemma

**Lemma 1.2.21** (Multiplication Lands in  $\text{Pre } \mathbb{Q}$ ). If  $b \neq 0_{\mathbb{Z}}$  and  $d \neq 0_{\mathbb{Z}}$ , then

$$b \cdot d \neq 0_{\mathbb{Z}}.$$

*Remark* (Dependencies).

- [\$\mathbb{Z}\$  Has No Zero Divisors](#)

### Lemma

**Lemma 1.2.22** (Multiplication Respects Fraction Equivalence on the Left). If  $[a, b] = [a', b']$ , then

$$[a, b] \cdot [c, d] = [a', b'] \cdot [c, d].$$

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{Q}\$](#)
- [Fraction Equivalence](#)
- [Equality Criterion for Rational Classes](#)
- [Multiplicative Cancellation on  \$\mathbb{Z}\$](#)

### Lemma

**Lemma 1.2.23** (Multiplication Respects Fraction Equivalence on the Right). If  $[c, d] = [c', d']$ , then

$$[a, b] \cdot [c, d] = [a, b] \cdot [c', d'].$$

*Remark* (Dependencies).



- [Multiplication on  \$\mathbb{Q}\$](#)
- [Fraction Equivalence](#)
- [Equality Criterion for Rational Classes](#)
- [Multiplicative Cancellation on  \$\mathbb{Z}\$](#)

#### Lemma

**Lemma 1.2.24** (Multiplication Respects Fraction Equivalence). *If  $[a, b] = [a', b']$  and  $[c, d] = [c', d']$ , then*

$$[a, b] \cdot [c, d] = [a', b'] \cdot [c', d'].$$

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{Q}\$](#)
- [Multiplication Respects Fraction Equivalence on the Left](#)
- [Multiplication Respects Fraction Equivalence on the Right](#)

#### Theorem

**Theorem 1.2.25** (Multiplication on  $\mathbb{Q}$  Is Well-Defined). *Multiplication determines a well-defined binary operation  $\mathbb{Q} \times \mathbb{Q} \rightarrow \mathbb{Q}$ .*

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{Q}\$](#)
- [Multiplication Respects Fraction Equivalence](#)

#### Theorem

**Theorem 1.2.26** (Multiplication on  $\mathbb{Q}$  Is Associative). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$(x \cdot y) \cdot z = x \cdot (y \cdot z).$$

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Multiplication on  \$\mathbb{Z}\$  Is Associative](#)

#### Theorem

**Theorem 1.2.27** (Multiplication on  $\mathbb{Q}$  Is Commutative). *For all  $x, y \in \mathbb{Q}$ ,*

$$x \cdot y = y \cdot x.$$

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Multiplication on  \$\mathbb{Z}\$  Is Commutative](#)

#### Theorem

**Theorem 1.2.28** (One Is a Left Multiplicative Identity on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$1_{\mathbb{Q}} \cdot x = x.$$

*Remark* (Dependencies).

- [One in  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [One Is a Multiplicative Identity on  \$\mathbb{Z}\$](#)

#### Corollary

**Corollary 1.2.29** (One Is a Right Multiplicative Identity on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$x \cdot 1_{\mathbb{Q}} = x.$$

*Remark* (Dependencies).

- [One Is a Left Multiplicative Identity on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Commutative](#)

#### Theorem

**Theorem 1.2.30** (One Is a Multiplicative Identity on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$1_{\mathbb{Q}} \cdot x = x \quad \text{and} \quad x \cdot 1_{\mathbb{Q}} = x.$$

*Remark* (Dependencies).

- [One Is a Left Multiplicative Identity on  \$\mathbb{Q}\$](#)
- [One Is a Right Multiplicative Identity on  \$\mathbb{Q}\$](#)

## Reciprocal on Nonzero $\mathbb{Q}$ -Classes

### Lemma

**Lemma 1.2.31** (Nonzero Predicate Is Well-Defined on  $\mathbb{Q}$ ). *For every prefraction  $(a, b) \in \text{Pre } \mathbb{Q}$ ,*

$$[a, b] \neq 0_{\mathbb{Q}} \iff a \neq 0_{\mathbb{Z}}.$$

*Thus nonzeroness is independent of the representative.*

*Remark* (Dependencies).

- [Zero in  \$\mathbb{Q}\$](#)
- [Vanishing Criterion in  \$\mathbb{Q}\$](#)

### Definition

**Definition 1.2.32** (Reciprocal on Nonzero  $\mathbb{Q}$ -Classes). For  $[a, b] \neq 0_{\mathbb{Q}}$ , define

$$[a, b]^{-1} := [b, a].$$

*Remark* (Dependencies).

- [Rational Class Notation](#)
- [Vanishing Criterion in  \$\mathbb{Q}\$](#)

### Lemma

**Lemma 1.2.33** (Reciprocal Lands in  $\text{Pre } \mathbb{Q}$ ). *If  $[a, b] \neq 0_{\mathbb{Q}}$ , then  $a \neq 0_{\mathbb{Z}}$ , so  $(b, a)$  is a prefraction.*

*Remark* (Dependencies).

- [Prefraction](#)
- [Prefraction Set](#)
- [Vanishing Criterion in  \$\mathbb{Q}\$](#)
- [Zero in  \$\mathbb{Z}\$](#)

### Lemma

**Lemma 1.2.34** (Reciprocal Respects Fraction Equivalence). *If  $[a, b] = [a', b']$  and  $[a, b] \neq 0_{\mathbb{Q}}$ , then*

$$[a, b]^{-1} = [a', b']^{-1}.$$

*Remark* (Dependencies).

- [Reciprocal on  \$\mathbb{Q}\$](#)

- [Fraction Equivalence](#)
- [Reciprocal Lands in Pre  \$\mathbb{Q}\$](#)

#### Theorem

**Theorem 1.2.35** (Reciprocal on  $\mathbb{Q}$  Is Well-Defined). *Reciprocal determines a well-defined map*

$$\mathbb{Q} \setminus \{0_{\mathbb{Q}}\} \rightarrow \mathbb{Q} \setminus \{0_{\mathbb{Q}}\}.$$

*Remark* (Dependencies).

- [Reciprocal on  \$\mathbb{Q}\$](#)
- [Reciprocal Lands in Pre  \$\mathbb{Q}\$](#)
- [Reciprocal Respects Fraction Equivalence](#)

#### Theorem

**Theorem 1.2.36** (Reciprocal Gives a Left Multiplicative Inverse on  $\mathbb{Q}$ ). *For every nonzero  $x \in \mathbb{Q}$ ,*

$$x^{-1} \cdot x = 1_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [Reciprocal on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$](#)
- [One in  \$\mathbb{Q}\$](#)
- [Reciprocal on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Well-Defined](#)

#### Corollary

**Corollary 1.2.37** (Reciprocal Gives a Right Multiplicative Inverse on  $\mathbb{Q}$ ). *For every nonzero  $x \in \mathbb{Q}$ ,*

$$x \cdot x^{-1} = 1_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [Reciprocal Gives a Left Multiplicative Inverse on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Commutative](#)

### Theorem

**Theorem 1.2.38** (Every Nonzero Rational Has a Multiplicative Inverse). *For every  $x \in \mathbb{Q}$  with  $x \neq 0_{\mathbb{Q}}$ , there exists  $y \in \mathbb{Q}$  such that*

$$x \cdot y = 1_{\mathbb{Q}} \quad \text{and} \quad y \cdot x = 1_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [Reciprocal Gives a Left Multiplicative Inverse on  \$\mathbb{Q}\$](#)
- [Reciprocal Gives a Right Multiplicative Inverse on  \$\mathbb{Q}\$](#)

## Division on $\mathbb{Q}$

### Definition

**Definition 1.2.39** (Division on  $\mathbb{Q}$ ). For  $x, y \in \mathbb{Q}$  with  $y \neq 0_{\mathbb{Q}}$ , define

$$x/y := x \cdot y^{-1}.$$

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{Q}\$](#)
- [Reciprocal on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.2.40** (Division on  $\mathbb{Q}$  Is Well-Defined). *Division determines a well-defined map*

$$\mathbb{Q} \times (\mathbb{Q} \setminus \{0_{\mathbb{Q}}\}) \rightarrow \mathbb{Q}.$$

*Remark* (Dependencies).

- [Division on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Reciprocal on  \$\mathbb{Q}\$  Is Well-Defined](#)

## Distributivity on $\mathbb{Q}$ -Classes

### Theorem

**Theorem 1.2.41** (Left Distributivity on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x \cdot (y + z) = x \cdot y + x \cdot z.$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Multiplication Distributes over Addition on  \$\mathbb{Z}\$](#)

#### Corollary

**Corollary 1.2.42** (Right Distributivity on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$(y + z) \cdot x = y \cdot x + z \cdot x.$$

*Remark* (Dependencies).

- [Left Distributivity on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Commutative](#)

#### Theorem

**Theorem 1.2.43** (Multiplication Distributes over Addition on  $\mathbb{Q}$ ). *Multiplication distributes over addition on both sides in  $\mathbb{Q}$ .*

*Remark* (Dependencies).

- [Left Distributivity on  \$\mathbb{Q}\$](#)
- [Right Distributivity on  \$\mathbb{Q}\$](#)

## Class-Level Field Summary

#### Theorem

**Theorem 1.2.44** ( $\mathbb{Q}$  Is an Additive Abelian Group). *The structure  $(\mathbb{Q}, +, 0_{\mathbb{Q}})$  is an abelian group.*

*Remark* (Dependencies).

- [Addition on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$  Is Associative](#)
- [Addition on  \$\mathbb{Q}\$  Is Commutative](#)
- [Zero Is an Additive Identity on  \$\mathbb{Q}\$](#)
- [Every Rational Has an Additive Inverse](#)

### Theorem

**Theorem 1.2.45** (Nonzero  $\mathbb{Q}$  Is a Multiplicative Abelian Group). *The structure  $(\mathbb{Q} \setminus \{0_{\mathbb{Q}}\}, \cdot, 1_{\mathbb{Q}})$  is an abelian group.*

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{Q}\$](#)
- [Nonzero Predicate Is Well-Defined on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Associative](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Commutative](#)
- [One Is a Multiplicative Identity on  \$\mathbb{Q}\$](#)
- [Every Nonzero Rational Has a Multiplicative Inverse](#)

### Theorem

**Theorem 1.2.46** ( $\mathbb{Q}$  Is a Field). *The structure  $(\mathbb{Q}, +, \cdot, 0_{\mathbb{Q}}, 1_{\mathbb{Q}})$  is a field.*

*Remark* (Dependencies).

- [\$\mathbb{Q}\$  Is an Additive Abelian Group](#)
- [Nonzero  \$\mathbb{Q}\$  Is a Multiplicative Abelian Group](#)
- [Multiplication Distributes over Addition on  \$\mathbb{Q}\$](#)
- [Zero Is Not One in  \$\mathbb{Q}\$](#)

## 1.3 Field Structure of $\mathbb{Q}$

### Derived Field Laws on $\mathbb{Q}$

#### Corollary

**Corollary 1.3.1** (Zero Product on  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$x \cdot y = 0_{\mathbb{Q}} \iff x = 0_{\mathbb{Q}} \text{ or } y = 0_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [\$\mathbb{Q}\$  Is a Field](#)

#### Corollary

**Corollary 1.3.2** (Double Negation on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$-(-x) = x.$$

*Remark* (Dependencies).

- [ℚ Is a Field](#)

#### Corollary

**Corollary 1.3.3** (Negation of a Product on  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$(-x) \cdot y = -(x \cdot y) \quad \text{and} \quad x \cdot (-y) = -(x \cdot y).$$

*Remark* (Dependencies).

- [ℚ Is a Field](#)

#### Corollary

**Corollary 1.3.4** (Product of Negatives on  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$(-x) \cdot (-y) = x \cdot y.$$

*Remark* (Dependencies).

- [Negation of a Product on  \$\mathbb{Q}\$](#)
- [Double Negation on  \$\mathbb{Q}\$](#)

#### Corollary

**Corollary 1.3.5** (Additive Cancellation on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x + z = y + z \implies x = y.$$

*Remark* (Dependencies).

- [ℚ Is an Additive Abelian Group](#)

#### Corollary

**Corollary 1.3.6** (Multiplicative Cancellation on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$z \neq 0_{\mathbb{Q}} \wedge x \cdot z = y \cdot z \implies x = y.$$

*Remark* (Dependencies).



- [ℚ Is a Field](#)

#### Corollary

**Corollary 1.3.7** (Uniqueness of Additive Inverses on  $\mathbb{Q}$ ). *Every additive inverse in  $\mathbb{Q}$  is unique.*

*Remark* (Dependencies).

- [ℚ Is an Additive Abelian Group](#)

#### Corollary

**Corollary 1.3.8** (Uniqueness of Multiplicative Inverses on  $\mathbb{Q}$ ). *Every multiplicative inverse of a nonzero rational number is unique.*

*Remark* (Dependencies).

- [Nonzero ℚ Is a Multiplicative Abelian Group](#)

#### Corollary

**Corollary 1.3.9** (The Inverse of One Is One on  $\mathbb{Q}$ ). *In  $\mathbb{Q}$ ,*

$$1_{\mathbb{Q}}^{-1} = 1_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [One in ℚ](#)
- [Reciprocal Gives a Left Multiplicative Inverse on ℚ](#)
- [One Is a Multiplicative Identity on ℚ](#)

#### Corollary

**Corollary 1.3.10** (Inverse of a Product on  $\mathbb{Q}$ ). *For all nonzero  $x, y \in \mathbb{Q}$ ,*

$$(x \cdot y)^{-1} = x^{-1} \cdot y^{-1}.$$

*Remark* (Dependencies).

- [ℚ Is a Field](#)
- [Reciprocal on ℚ Is Well-Defined](#)
- [Uniqueness of Multiplicative Inverses on ℚ](#)

#### Corollary

**Corollary 1.3.11** (Inverse of an Inverse on  $\mathbb{Q}$ ). *For every nonzero  $x \in \mathbb{Q}$ ,*

$$(x^{-1})^{-1} = x.$$

*Remark* (Dependencies).

## Integer Powers on $\mathbb{Q}$

### Definition

**Definition 1.3.12** (Integer Power on  $\mathbb{Q}$ ). Let  $x \in \mathbb{Q}$ . For  $n \in \mathbb{W}$ , define  $x^n$  by the usual recursive power operation. If  $x \neq 0_{\mathbb{Q}}$  and  $n \in \mathbb{Z}$  is negative, define

$$x^n := (x^{-1})^{-n}.$$

*Remark* (Dependencies).

- [ℚ Is a Field](#)
- [Every Nonzero Rational Has a Multiplicative Inverse](#)

### Corollary

**Corollary 1.3.13** (Exponent Addition Law on  $\mathbb{Q}$ ). For every nonzero  $x \in \mathbb{Q}$  and all  $m, n \in \mathbb{Z}$ ,

$$x^{m+n} = x^m \cdot x^n.$$

*Remark* (Dependencies).

- [Integer Power on  \$\mathbb{Q}\$](#)
- [Integer Power on  \$\mathbb{Q}\$](#)

### Corollary

**Corollary 1.3.14** (Power of a Power on  $\mathbb{Q}$ ). For every nonzero  $x \in \mathbb{Q}$  and all  $m, n \in \mathbb{Z}$ ,

$$(x^m)^n = x^{m \cdot n}.$$

*Remark* (Dependencies).

- [Integer Power on  \$\mathbb{Q}\$](#)
- [Integer Power on  \$\mathbb{Q}\$](#)

### Corollary

**Corollary 1.3.15** (Power of a Product on  $\mathbb{Q}$ ). For all nonzero  $x, y \in \mathbb{Q}$  and every  $n \in \mathbb{Z}$ ,

$$(x \cdot y)^n = x^n \cdot y^n.$$

*Remark* (Dependencies).

- [Integer Power on  \$\mathbb{Q}\$](#)
- [Integer Power on  \$\mathbb{Q}\$](#)
- [Inverse of a Product on  \$\mathbb{Q}\$](#)

## 1.4 Order on $\mathbb{Q}$

### Positivity on $\mathbb{Q}$

#### Definition

**Definition 1.4.1** (Positive Rational). A rational number  $[a, b] \in \mathbb{Q}$  is *positive* if

$$a \cdot b > 0_{\mathbb{Z}}.$$

*Remark* (Dependencies).

- [Rational Class Notation](#)
- [Multiplication on  \$\mathbb{Z}\$](#)
- [Order on  \$\mathbb{Z}\$](#)

#### Lemma

**Lemma 1.4.2** (Positivity Respects Fraction Equivalence on  $\mathbb{Q}$ ). If  $[a, b] = [c, d]$ , then

$$a \cdot b > 0_{\mathbb{Z}} \iff c \cdot d > 0_{\mathbb{Z}}.$$

*Remark* (Dependencies).

- [\$\mathbb{Z}\$  Is a Totally Ordered Set](#)
- [\$\mathbb{Z}\$  Has No Zero Divisors](#)

#### Theorem

**Theorem 1.4.3** (Positivity on  $\mathbb{Q}$  Is Well-Defined). The predicate “ $x$  is positive” is well-defined on equivalence classes in  $\mathbb{Q}$ .

*Remark* (Dependencies).

- [Positive Rational](#)
- [Positivity Respects Fraction Equivalence on  \$\mathbb{Q}\$](#)

#### Definition

**Definition 1.4.4** (Strict Order on  $\mathbb{Q}$ ). For  $x, y \in \mathbb{Q}$ , define

$$x < y \quad :\iff \quad y - x \text{ is positive.}$$

*Remark* (Dependencies).

- [Positive Rational](#)
- [Subtraction on  \$\mathbb{Q}\$](#)

### Definition

**Definition 1.4.5** (Order on  $\mathbb{Q}$ ). For  $x, y \in \mathbb{Q}$ , define

$$x \leq y \quad :\Longleftrightarrow \quad x < y \text{ or } x = y.$$

*Remark* (Dependencies).

- [Strict Order on  \$\mathbb{Q}\$](#)

### Proposition

**Proposition 1.4.6** (Product of Positives Is Positive on  $\mathbb{Q}$ ). For all  $x, y \in \mathbb{Q}$ ,

$$0_{\mathbb{Q}} < x \wedge 0_{\mathbb{Q}} < y \implies 0_{\mathbb{Q}} < x \cdot y.$$

*Remark* (Dependencies).

- [Positive Rational](#)
- [Multiplication on  \$\mathbb{Q}\$](#)
- [\$\mathbb{Z}\$  Has No Zero Divisors](#)

## Trichotomy on $\mathbb{Q}$

### Theorem

**Theorem 1.4.7** (Sign Trichotomy on  $\mathbb{Q}$ ). For every  $x \in \mathbb{Q}$ , exactly one of

$$x < 0_{\mathbb{Q}}, \quad x = 0_{\mathbb{Q}}, \quad 0_{\mathbb{Q}} < x$$

holds.

*Remark* (Dependencies).

- [Positive Rational](#)
- [Strict Order on  \$\mathbb{Q}\$](#)
- [Zero in  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.8** (Trichotomy on  $\mathbb{Q}$ ). For all  $x, y \in \mathbb{Q}$ , exactly one of

$$x < y, \quad x = y, \quad y < x$$

holds.

*Remark* (Dependencies).

- [Strict Order on  \$\mathbb{Q}\$](#)
- [Subtraction on  \$\mathbb{Q}\$](#)
- [Sign Trichotomy on  \$\mathbb{Q}\$](#)

## Total Order on $\mathbb{Q}$

### Theorem

**Theorem 1.4.9** (Strict Order on  $\mathbb{Q}$  Is Irreflexive). *For every  $x \in \mathbb{Q}$ , not  $x < x$ .*

*Remark* (Dependencies).

- [Strict Order on  \$\mathbb{Q}\$](#)
- [Trichotomy on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.10** (Strict Order on  $\mathbb{Q}$  Is Asymmetric). *For all  $x, y \in \mathbb{Q}$ ,*

$$x < y \implies \neg(y < x).$$

*Remark* (Dependencies).

- [Strict Order on  \$\mathbb{Q}\$](#)
- [Trichotomy on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.11** (Strict Order on  $\mathbb{Q}$  Is Transitive). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x < y \wedge y < z \implies x < z.$$

*Remark* (Dependencies).

- [Strict Order on  \$\mathbb{Q}\$](#)
- [Addition Preserves Strict Order on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.12** (Order on  $\mathbb{Q}$  Is Reflexive). *For every  $x \in \mathbb{Q}$ ,  $x \leq x$ .*

*Remark* (Dependencies).

- [Order on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.13** (Order on  $\mathbb{Q}$  Is Antisymmetric). *For all  $x, y \in \mathbb{Q}$ ,*

$$x \leq y \wedge y \leq x \implies x = y.$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{Q}\$](#)
- [Strict Order on  \$\mathbb{Q}\$  Is Asymmetric](#)

### Theorem

**Theorem 1.4.14** (Order on  $\mathbb{Q}$  Is Transitive). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x \leq y \wedge y \leq z \implies x \leq z.$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{Q}\$](#)
- [Strict Order on  \$\mathbb{Q}\$  Is Transitive](#)

### Theorem

**Theorem 1.4.15** (Order on  $\mathbb{Q}$  Is Total). *For all  $x, y \in \mathbb{Q}$ ,*

$$x \leq y \text{ or } y \leq x.$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{Q}\$](#)
- [Trichotomy on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.16** ( $\mathbb{Q}$  Is a Totally Ordered Set). *The ordered pair  $(\mathbb{Q}, \leq)$  is a totally ordered set.*

*Remark* (Dependencies).

- [Ordered Set](#)
- [Total Order](#)
- [Order on  \$\mathbb{Q}\$](#)
- [Order on  \$\mathbb{Q}\$  Is Reflexive](#)
- [Order on  \$\mathbb{Q}\$  Is Antisymmetric](#)
- [Order on  \$\mathbb{Q}\$  Is Transitive](#)
- [Order on  \$\mathbb{Q}\$  Is Total](#)

## Addition Order Compatibility on $\mathbb{Q}$

### Theorem

**Theorem 1.4.17** (Left Addition Preserves Strict Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x < y \implies z + x < z + y.$$

*Remark* (Dependencies).

- [Strict Order on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$](#)
- [Positive Rational](#)

### Corollary

**Corollary 1.4.18** (Right Addition Preserves Strict Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x < y \implies x + z < y + z.$$

*Remark* (Dependencies).

- [Left Addition Preserves Strict Order on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$  Is Commutative](#)

### Theorem

**Theorem 1.4.19** (Addition Preserves Strict Order on  $\mathbb{Q}$ ). *Addition preserves strict order on both sides in  $\mathbb{Q}$ .*

*Remark* (Dependencies).

- [Left Addition Preserves Strict Order on  \$\mathbb{Q}\$](#)
- [Right Addition Preserves Strict Order on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.20** (Left Addition Preserves Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x \leq y \implies z + x \leq z + y.$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{Q}\$](#)
- [Left Addition Preserves Strict Order on  \$\mathbb{Q}\$](#)

### Corollary

**Corollary 1.4.21** (Right Addition Preserves Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x \leq y \implies x + z \leq y + z.$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{Q}\$](#)
- [Right Addition Preserves Strict Order on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.22** (Addition Preserves Order on  $\mathbb{Q}$ ). *Addition preserves non-strict order on both sides in  $\mathbb{Q}$ .*

*Remark* (Dependencies).

- [Left Addition Preserves Order on  \$\mathbb{Q}\$](#)
- [Right Addition Preserves Order on  \$\mathbb{Q}\$](#)

### Corollary

**Corollary 1.4.23** (Addition Reflects Order on  $\mathbb{Q}$ ). *Addition by a fixed rational reflects both strict and non-strict order.*

*Remark* (Dependencies).

- [Addition Preserves Strict Order on  \$\mathbb{Q}\$](#)
- [Addition Preserves Order on  \$\mathbb{Q}\$](#)
- [Negation Gives a Left Additive Inverse on  \$\mathbb{Q}\$](#)

## Multiplication Order Compatibility on $\mathbb{Q}$

### Theorem

**Theorem 1.4.24** (Left Multiplication by Positives Preserves Strict Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < z \wedge x < y \implies z \cdot x < z \cdot y.$$

*Remark* (Dependencies).

- [Strict Order on  \$\mathbb{Q}\$](#)
- [Product of Positives Is Positive on  \$\mathbb{Q}\$](#)



- [Multiplication on  \$\mathbb{Q}\$](#)

#### Corollary

**Corollary 1.4.25** (Right Multiplication by Positives Preserves Strict Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < z \wedge x < y \implies x \cdot z < y \cdot z.$$

*Remark* (Dependencies).

- [Left Multiplication by Positives Preserves Strict Order on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Commutative](#)

#### Theorem

**Theorem 1.4.26** (Multiplication by Positives Preserves Strict Order on  $\mathbb{Q}$ ). *Multiplication by a positive rational preserves strict order on both sides.*

*Remark* (Dependencies).

- [Left Multiplication by Positives Preserves Strict Order on  \$\mathbb{Q}\$](#)
- [Right Multiplication by Positives Preserves Strict Order on  \$\mathbb{Q}\$](#)

#### Theorem

**Theorem 1.4.27** (Left Multiplication by Positives Preserves Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < z \wedge x \leq y \implies z \cdot x \leq z \cdot y.$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{Q}\$](#)
- [Left Multiplication by Positives Preserves Strict Order on  \$\mathbb{Q}\$](#)

#### Corollary

**Corollary 1.4.28** (Right Multiplication by Positives Preserves Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < z \wedge x \leq y \implies x \cdot z \leq y \cdot z.$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{Q}\$](#)
- [Right Multiplication by Positives Preserves Strict Order on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.29** (Multiplication by Positives Preserves Order on  $\mathbb{Q}$ ). *Multiplication by a positive rational preserves non-strict order on both sides.*

*Remark* (Dependencies).

- [Left Multiplication by Positives Preserves Order on  \$\mathbb{Q}\$](#)
- [Right Multiplication by Positives Preserves Order on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.30** (Multiplication by Negatives Reverses Strict Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$z < 0_{\mathbb{Q}} \wedge x < y \implies z \cdot y < z \cdot x.$$

*Remark* (Dependencies).

- [Strict Order on  \$\mathbb{Q}\$](#)
- [Negation of a Product on  \$\mathbb{Q}\$](#)
- [Positive Multiplication Preserves Strict Order on  \$\mathbb{Q}\$](#)

### Corollary

**Corollary 1.4.31** (Multiplication by Negatives Reverses Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$z < 0_{\mathbb{Q}} \wedge x \leq y \implies z \cdot y \leq z \cdot x.$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{Q}\$](#)
- [Multiplication by Negatives Reverses Strict Order on  \$\mathbb{Q}\$](#)

## Reciprocal Order Laws on $\mathbb{Q}$

### Theorem

**Theorem 1.4.32** (Sign of a Reciprocal on  $\mathbb{Q}$ ). *For every nonzero  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < x \implies 0_{\mathbb{Q}} < x^{-1}, \quad x < 0_{\mathbb{Q}} \implies x^{-1} < 0_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [Reciprocal on  \$\mathbb{Q}\$](#)
- [Order on  \$\mathbb{Q}\$](#)

- [Product of Positives Is Positive on  \$\mathbb{Q}\$](#)
- [Reciprocal Is a Left Multiplicative Inverse on  \$\mathbb{Q}\$](#)

#### Theorem

**Theorem 1.4.33** (Reciprocal Reverses Order on Positives in  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < x < y \implies 0_{\mathbb{Q}} < y^{-1} < x^{-1}.$$

*Remark* (Dependencies).

- [Reciprocal on  \$\mathbb{Q}\$](#)
- [Order on  \$\mathbb{Q}\$](#)
- [Sign of a Reciprocal on  \$\mathbb{Q}\$](#)
- [Positive Multiplication Preserves Strict Order on  \$\mathbb{Q}\$](#)

## Absolute Value on $\mathbb{Q}$

#### Definition

**Definition 1.4.34** (Absolute Value on  $\mathbb{Q}$ ). For  $x \in \mathbb{Q}$ , define

$$|x| := \begin{cases} x, & 0_{\mathbb{Q}} \leq x, \\ -x, & x < 0_{\mathbb{Q}}. \end{cases}$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{Q}\$](#)
- [Zero in  \$\mathbb{Q}\$](#)
- [Negation on  \$\mathbb{Q}\$](#)

#### Theorem

**Theorem 1.4.35** (Absolute Value Is Nonnegative on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} \leq |x|.$$

*Remark* (Dependencies).

- [Absolute Value on  \$\mathbb{Q}\$](#)
- [Order on  \$\mathbb{Q}\$](#)
- [Trichotomy on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.36** (Absolute Value Vanishes Only at Zero on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$|x| = 0_{\mathbb{Q}} \iff x = 0_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [Absolute Value on  \$\mathbb{Q}\$](#)
- [Absolute Value Is Nonnegative on  \$\mathbb{Q}\$](#)
- [Trichotomy on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.37** (Absolute Value Is Multiplicative on  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$|x \cdot y| = |x| \cdot |y|.$$

*Remark* (Dependencies).

- [Absolute Value on  \$\mathbb{Q}\$](#)
- [Positive Multiplication Preserves Order on  \$\mathbb{Q}\$](#)
- [Product of Negatives on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.38** (Triangle Inequality on  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$|x + y| \leq |x| + |y|.$$

*Remark* (Dependencies).

- [Absolute Value on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$](#)
- [Order on  \$\mathbb{Q}\$](#)
- [\$\mathbb{Q}\$  Is a Totally Ordered Set](#)
- [Addition Preserves Order on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.4.39** (Absolute Value of a Reciprocal on  $\mathbb{Q}$ ). *For every nonzero  $x \in \mathbb{Q}$ ,*

$$|x^{-1}| = |x|^{-1}.$$

*Remark* (Dependencies).

- [Absolute Value on  \$\mathbb{Q}\$](#)
- [Reciprocal on  \$\mathbb{Q}\$  Is Well-Defined](#)
- [Sign of a Reciprocal on  \$\mathbb{Q}\$](#)
- [Absolute Value Is Multiplicative on  \$\mathbb{Q}\$](#)

## 1.5 Ordered Field Structure of $\mathbb{Q}$

### Ordered Field Instantiation

#### Theorem

**Theorem 1.5.1** ( $\mathbb{Q}$  Is an Ordered Field). *The structure*

$$(\mathbb{Q}, +, \cdot, 0_{\mathbb{Q}}, 1_{\mathbb{Q}}, \leq)$$

*is an ordered field.*

*Remark* (Dependencies).

- [\$\mathbb{Q}\$  Is a Field](#)
- [\$\mathbb{Q}\$  Is a Totally Ordered Set](#)
- [Addition Preserves Order on  \$\mathbb{Q}\$](#)
- [Multiplication by Positives Preserves Order on  \$\mathbb{Q}\$](#)

#### Corollary

**Corollary 1.5.2** (Ordered Field Laws Hold on  $\mathbb{Q}$ ). *Every theorem that depends only on the ordered-field axioms applies to  $\mathbb{Q}$  with its constructed operations and order.*

*Remark* (Dependencies).

- [\$\mathbb{Q}\$  Is an Ordered Field](#)

### Standard Ordered-Field Consequences

#### Corollary

**Corollary 1.5.3** (Squares Are Nonnegative on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} \leq x^2.$$

*Remark* (Dependencies).

- [\$\mathbb{Q}\$  Is an Ordered Field](#)

- [Product of Positives Is Positive on  \$\mathbb{Q}\$](#)

#### Corollary

**Corollary 1.5.4** (One Is Positive in  $\mathbb{Q}$ ). *In  $\mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < 1_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [\$\mathbb{Q}\$  Is an Ordered Field](#)
- [One in  \$\mathbb{Q}\$](#)

#### Corollary

**Corollary 1.5.5** (Natural Numbers Are Positive in  $\mathbb{Q}$ ). *Every natural number embedded in  $\mathbb{Q}$  is positive.*

*Remark* (Dependencies).

- [One Is Positive in  \$\mathbb{Q}\$](#)

#### Corollary

**Corollary 1.5.6** (Positive Rationals Have Positive Inverses). *For every  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < x \implies 0_{\mathbb{Q}} < x^{-1}.$$

*Remark* (Dependencies).

- [\$\mathbb{Q}\$  Is an Ordered Field](#)
- [Reciprocal Reverses Order on Positives in  \$\mathbb{Q}\$](#)

#### Corollary

**Corollary 1.5.7** (Fraction Comparison in  $\mathbb{Q}$ ). *For rational classes with positive denominator product,*

$$[a, b] < [c, d] \iff a \cdot d < c \cdot b.$$

*Remark* (Dependencies).

- [\$\mathbb{Q}\$  Is an Ordered Field](#)
- [Product of Positives Is Positive on  \$\mathbb{Q}\$](#)
- [One in  \$\mathbb{Q}\$](#)
- [Reciprocal Reverses Order on Positives in  \$\mathbb{Q}\$](#)
- [Positive Multiplication Preserves Order on  \$\mathbb{Q}\$](#)
- [Reciprocal Reverses Order on Positives in  \$\mathbb{Q}\$](#)

## 1.6 Arbitrary Closeness in $\mathbb{Q}$

### Density and the Archimedean Property

#### Definition

**Definition 1.6.1** (Two in  $\mathbb{Q}$ ). Define

$$2_{\mathbb{Q}} := 1_{\mathbb{Q}} + 1_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [One in  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Q}\$](#)

#### Lemma

**Lemma 1.6.2** (Two Is Nonzero in  $\mathbb{Q}$ ). *In  $\mathbb{Q}$ ,*

$$2_{\mathbb{Q}} \neq 0_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [One Is Positive in  \$\mathbb{Q}\$](#)
- [Trichotomy on  \$\mathbb{Q}\$](#)

#### Lemma

**Lemma 1.6.3** (Midpoint Lies Strictly Between Rationals). *For all  $x, y \in \mathbb{Q}$ ,*

$$x < y \implies x < (x + y) \cdot 2_{\mathbb{Q}}^{-1} < y.$$

*Remark* (Dependencies).

- [\$\mathbb{Q}\$  Is an Ordered Field](#)
- [Two Is Nonzero in  \$\mathbb{Q}\$](#)

#### Theorem

**Theorem 1.6.4** (Density of  $\mathbb{Q}$  in Itself). *For all  $x, y \in \mathbb{Q}$ ,*

$$x < y \implies \exists r \in \mathbb{Q} (x < r < y).$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{Q}\$](#)
- [Midpoint Lies Strictly Between Rationals](#)

### Theorem

**Theorem 1.6.5** (Archimedean Property of  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < x \implies \exists n \in \mathbb{N} (y < n \cdot x).$$

*Remark* (Dependencies).

- [Well-Ordering Principle](#)
- [Division Algorithm on  \$\mathbb{Z}\$](#)
- [\$\mathbb{Z}\$  Is a Totally Ordered Set](#)

## From Discreteness to Arbitrary Closeness

*Remark 1.6.6* (Discrete Systems Before  $\mathbb{Q}$ ). The systems  $\mathbb{N}$ ,  $\mathbb{W}$ , and  $\mathbb{Z}$  are discrete in the order sense: consecutive integer points have no same-system point between them. The rational numbers are the first number system in this construction chain where intervals can be subdivided internally without leaving the system.

### Definition

**Definition 1.6.7** (Arbitrarily Close Distinct Points). A number system has arbitrarily close distinct points if for every point  $x$  and every positive tolerance  $\varepsilon$ , there is a point  $y \neq x$  such that

$$|x - y| < \varepsilon.$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{Q}\$](#)
- [Absolute Value on  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 1.6.8** ( $\mathbb{Q}$  Has Arbitrarily Close Distinct Points). *For every  $x \in \mathbb{Q}$  and every  $\varepsilon \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < \varepsilon \implies \exists y \in \mathbb{Q} (y \neq x \text{ and } |x - y| < \varepsilon).$$

*Remark* (Dependencies).

- [Reciprocals Become Arbitrarily Small in  \$\mathbb{Q}\$](#)
- [Addition Preserves Strict Order on  \$\mathbb{Q}\$](#)

### Corollary

**Corollary 1.6.9** ( $\mathbb{Q}$  Has No Adjacent Points). *There do not exist rational numbers  $x < y$  with no rational number strictly between them.*

*Remark* (Dependencies).



- [Density of  \$\mathbb{Q}\$  in Itself](#)

#### Proposition

**Proposition 1.6.10** ( $\mathbb{Q}$  Has No Least Positive Rational). *For every  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < x \implies \exists y \in \mathbb{Q} (0_{\mathbb{Q}} < y < x).$$

*Remark* (Dependencies).

- [Density of  \$\mathbb{Q}\$  in Itself](#)

#### Corollary

**Corollary 1.6.11** (Reciprocals Become Arbitrarily Small in  $\mathbb{Q}$ ). *For every  $\varepsilon \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < \varepsilon \implies \exists n \in \mathbb{N} (0_{\mathbb{Q}} < n^{-1} < \varepsilon).$$

*Remark* (Dependencies).

- [Archimedean Property of  \$\mathbb{Q}\$](#)
- [Sign of a Reciprocal on  \$\mathbb{Q}\$](#)

## Limit-Like Behavior Before Limits

#### Proposition

**Proposition 1.6.12** (One-Sided Rational Approximation). *For every  $x \in \mathbb{Q}$  and every  $\varepsilon \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < \varepsilon \implies \exists y, z \in \mathbb{Q} (x - \varepsilon < y < x < z < x + \varepsilon).$$

*Remark* (Dependencies).

- [\$\mathbb{Q}\$  Has Arbitrarily Close Distinct Points](#)
- [Density of  \$\mathbb{Q}\$  in Itself](#)

*Remark 1.6.13* (Refinement and Bounds). The Archimedean property supplies arbitrarily small rational increments, so a rational point can be approached from one side at any prescribed tolerance. Mediants supply a complementary order-refinement mechanism: once two rational bounds are known, their mediant is a new rational bound strictly between them. The following approximation topics make these two mechanisms explicit through dyadic grids, mediant refinement, Stern–Brocot paths, and rational Cauchy sequences.

## 1.7 Rational Gaps

### No Rational Square Root of Two

#### Theorem

**Theorem 1.7.1** (No Rational Square Root of Two). *There is no rational number  $q \in \mathbb{Q}$  such that*

$$q \cdot q = 2_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [ℚ Is a Field](#)
- [ℤ Is an Integral Domain](#)
- TODO: formalize the parity or infinite-descent facts in  $\mathbb{Z}$ .

### The Square-Two Gap

#### Definition

**Definition 1.7.2** (The Gap Set in  $\mathbb{Q}$ ). Define

$$S := \{q \in \mathbb{Q} : 0_{\mathbb{Q}} < q \text{ and } q \cdot q < 2_{\mathbb{Q}}\}.$$

*Remark* (Dependencies).

- [Rationals](#)
- [Order on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$](#)
- [Zero in  \$\mathbb{Q}\$](#)

#### Proposition

**Proposition 1.7.3** (The Gap Set Is Bounded Above in  $\mathbb{Q}$ ). *The set  $S$  is nonempty and bounded above in  $\mathbb{Q}$ .*

*Remark* (Dependencies).

- [ℚ Is an Ordered Field](#)
- [Natural Numbers Are Positive in  \$\mathbb{Q}\$](#)

#### Theorem

**Theorem 1.7.4** (The Square-Two Gap Has No Rational Supremum). *The set  $S$  has no least upper bound in  $\mathbb{Q}$ .*

*Remark* (Dependencies).

- [No Rational Square Root of Two](#)
- [Density of  \$\mathbb{Q}\$  in Itself](#)
- [\$\mathbb{Q}\$  Is an Ordered Field](#)

#### Corollary

**Corollary 1.7.5** ( $\mathbb{Q}$  Is Order-Incomplete). *The ordered field  $\mathbb{Q}$  is not order-complete: there exists a nonempty bounded-above subset of  $\mathbb{Q}$  with no supremum in  $\mathbb{Q}$ .*

*Remark* (Dependencies).

- [The Gap Set Is Bounded Above in  \$\mathbb{Q}\$](#)
- [The Square-Two Gap Has No Rational Supremum](#)

## Cauchy Sequences in $\mathbb{Q}$

#### Definition

**Definition 1.7.6** (Cauchy Sequence in  $\mathbb{Q}$ ). A sequence  $(x_n)$  in  $\mathbb{Q}$  is Cauchy if

$$\forall \varepsilon \in \mathbb{Q} (0_{\mathbb{Q}} < \varepsilon \implies \exists N \in \mathbb{N} \forall m, n \geq N |x_m - x_n| < \varepsilon).$$

*Remark* (Dependencies).

- [Rationals](#)
- [Order on  \$\mathbb{Q}\$](#)
- [Absolute Value on  \$\mathbb{Q}\$](#)
- [Subtraction on  \$\mathbb{Q}\$](#)

#### Definition

**Definition 1.7.7** (Null Sequence in  $\mathbb{Q}$ ). A sequence  $(x_n)$  in  $\mathbb{Q}$  is null if

$$\forall \varepsilon \in \mathbb{Q} (0_{\mathbb{Q}} < \varepsilon \implies \exists N \in \mathbb{N} \forall n \geq N |x_n| < \varepsilon).$$

*Remark* (Dependencies).

- [Rationals](#)
- [Order on  \$\mathbb{Q}\$](#)
- [Absolute Value on  \$\mathbb{Q}\$](#)

### Lemma

**Lemma 1.7.8** (Rational Limits Are Unique). *A convergent sequence in  $\mathbb{Q}$  has at most one rational limit.*

*Remark* (Dependencies).

- [Triangle Inequality on  \$\mathbb{Q}\$](#)
- [\$\mathbb{Q}\$  Is an Ordered Field](#)

## 1.8 Embedding $\mathbb{Z}$ into $\mathbb{Q}$

### The Integer Embedding

#### Definition

**Definition 1.8.1** (Embedding of  $\mathbb{Z}$  into  $\mathbb{Q}$ ). Define

$$\iota : \mathbb{Z} \rightarrow \mathbb{Q}, \quad \iota(n) := [n, 1_{\mathbb{Z}}].$$

*Remark* (Dependencies).

- [Rational Numbers](#)
- [One in  \$\mathbb{Z}\$](#)
- [Zero Is Not One in  \$\mathbb{Z}\$](#)

#### Theorem

**Theorem 1.8.2** (Embedding Preserves Zero). *The embedding sends integer zero to rational zero:*

$$\iota(0_{\mathbb{Z}}) = 0_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [Embedding of  \$\mathbb{Z}\$  into  \$\mathbb{Q}\$](#)
- [Zero in  \$\mathbb{Z}\$](#)
- [Zero in  \$\mathbb{Q}\$](#)

#### Theorem

**Theorem 1.8.3** (Embedding Preserves One). *The embedding sends integer one to rational one:*

$$\iota(1_{\mathbb{Z}}) = 1_{\mathbb{Q}}.$$

*Remark* (Dependencies).

- [Embedding of  \$\mathbb{Z}\$  into  \$\mathbb{Q}\$](#)
- [One in  \$\mathbb{Z}\$](#)
- [One in  \$\mathbb{Q}\$](#)

#### Theorem

**Theorem 1.8.4** (Embedding  $\mathbb{Z}$  into  $\mathbb{Q}$  Is Injective). *For all  $m, n \in \mathbb{Z}$ ,*

$$\iota(m) = \iota(n) \implies m = n.$$

*Remark* (Dependencies).

- [Equality Criterion for Rational Classes](#)
- [One Is a Multiplicative Identity on  \$\mathbb{Z}\$](#)

## Preservation of Algebra and Order

#### Theorem

**Theorem 1.8.5** (Embedding Preserves Addition). *For all  $m, n \in \mathbb{Z}$ ,*

$$\iota(m + n) = \iota(m) + \iota(n).$$

*Remark* (Dependencies).

- [Embedding  \$\mathbb{Z}\$  into  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Z}\$](#)
- [Addition on  \$\mathbb{Q}\$](#)
- [Addition on  \$\mathbb{Z}\$  Is Well-Defined](#)
- [Addition on  \$\mathbb{Q}\$  Is Well-Defined](#)

#### Theorem

**Theorem 1.8.6** (Embedding Preserves Multiplication). *For all  $m, n \in \mathbb{Z}$ ,*

$$\iota(m \cdot n) = \iota(m) \cdot \iota(n).$$

*Remark* (Dependencies).

- [Embedding  \$\mathbb{Z}\$  into  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Z}\$](#)
- [Multiplication on  \$\mathbb{Q}\$](#)

- [Multiplication on  \$\mathbb{Z}\$  Is Well-Defined](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Well-Defined](#)

#### Theorem

**Theorem 1.8.7** (Embedding Preserves Order). *For all  $m, n \in \mathbb{Z}$ ,*

$$m \leq n \iff \iota(m) \leq \iota(n).$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{Q}\$ -Classes](#)
- [Multiplication on  \$\mathbb{Q}\$ -Classes](#)
- [Order on  \$\mathbb{Q}\$](#)
- [\$\mathbb{Z}\$  Is a Totally Ordered Set](#)

#### Theorem

**Theorem 1.8.8** ( $\mathbb{Z}$  Embeds into  $\mathbb{Q}$ ). *The map  $\iota : \mathbb{Z} \rightarrow \mathbb{Q}$  is an injective order-preserving ring homomorphism.*

*Remark* (Dependencies).

- [Embedding Preserves Zero](#)
- [Embedding Preserves One](#)
- [Embedding  \$\mathbb{Z}\$  into  \$\mathbb{Q}\$  Is Injective](#)
- [Embedding Preserves Addition](#)
- [Embedding Preserves Multiplication](#)
- [Embedding Preserves Order](#)

## Field of Fractions

#### Theorem

**Theorem 1.8.9** ( $\mathbb{Q}$  Is the Field of Fractions of  $\mathbb{Z}$ ). *For every  $x \in \mathbb{Q}$ , there exist  $m, n \in \mathbb{Z}$  with  $n \neq 0_{\mathbb{Z}}$  such that*

$$x = \iota(m) \cdot \iota(n)^{-1}.$$

*Remark* (Dependencies).

- [Rational Class Notation](#)

- [Every Nonzero Rational Has a Multiplicative Inverse](#)
- [Vanishing Criterion](#)

#### Corollary

**Corollary 1.8.10** (The Embedding of  $\mathbb{Z}$  into  $\mathbb{Q}$  Is Not Surjective). *The image of  $\iota : \mathbb{Z} \rightarrow \mathbb{Q}$  is a proper subset of  $\mathbb{Q}$ .*

*Remark* (Dependencies).

- [Density of  \$\mathbb{Q}\$  in Itself](#)
- [Embedding Preserves Order](#)

## 1.9 Transition to $\mathbb{R}$

### Why the Real Numbers Are Introduced

*Remark 1.9.1* (Why the Real Numbers Are Introduced). The rational numbers form an ordered field, are dense in themselves, and satisfy the Archimedean property. These facts make  $\mathbb{Q}$  a rich arithmetic and order structure, but they do not make it complete.

The square-two gap shows the defect precisely: there is a nonempty bounded-above subset of  $\mathbb{Q}$  with no supremum in  $\mathbb{Q}$ . The real numbers are introduced as a completion of  $\mathbb{Q}$ , so that bounded-above sets acquire suprema and Cauchy sequences acquire limits.

*Remark* (Dependencies).

- [\$\mathbb{Q}\$  Is an Ordered Field](#)
- [Density of  \$\mathbb{Q}\$  in Itself](#)
- [Archimedean Property of  \$\mathbb{Q}\$](#)
- [\$\mathbb{Q}\$  Is Order-Incomplete](#)
- [Cauchy Sequence in  \$\mathbb{Q}\$](#)

## 1.10 Canonical Rational Number Statement Plan

### Architecture Rule for the Rational Numbers

The rational-number chapter is organized around a local-versus-generic separation. Local results open the quotient construction box: prefractions, fraction equivalence, well-defined operations, field axioms, order, density, gaps, and the embedding of  $\mathbb{Z}$  into  $\mathbb{Q}$ . Generic field and ring consequences are not re-proved here. They are cited as corollaries from the general algebraic infrastructure developed earlier.

*Remark* (Partiality of reciprocal). The new structural issue relative to  $\mathbb{Z}$  is that reciprocal is a partial operation. It is defined only on nonzero rational classes, so the nonzero predicate must be proved well-defined before reciprocal is introduced.

## Construction of $\mathbb{Q}$

- Def def:nonzero-integers:  $\mathbb{Z}^* := \mathbb{Z} \setminus \{0\}$ .
- Def def:prefraction: a prefraction is  $(a, b) \in \mathbb{Z} \times \mathbb{Z}^*$ , read as  $a/b$ .
- Def def:prefraction-set:  $\text{PreQ} := \mathbb{Z} \times \mathbb{Z}^*$ .
- Def def:fraction-equivalence:  $(a, b) \sim_Q (c, d)$  iff  $ad = bc$ .
- Lem lem:fraction-equivalence-reflexive:  $(a, b) \sim_Q (a, b)$ .
- Lem lem:fraction-equivalence-symmetric:  $(a, b) \sim_Q (c, d) \Rightarrow (c, d) \sim_Q (a, b)$ .
- Lem lem:fraction-equivalence-transitive: fraction equivalence is transitive; depends on nonzero multiplicative cancellation in  $\mathbb{Z}$ .
- Thm thm:fraction-equivalence-relation:  $\sim_Q$  is an equivalence relation on  $\text{PreQ}$ .
- Def def:rational:  $\mathbb{Q} := \text{PreQ} / \sim_Q$ .
- Def def:rational-class-notation:  $[a, b]$  denotes the  $\sim_Q$ -class of  $(a, b)$  with  $b \neq 0$ .
- Def def:zero-in-q:  $0_Q := [0, 1]$ .
- Def def:one-in-q:  $1_Q := [1, 1]$ .
- Lem lem:vanishing-criterion-q:  $[a, b] = 0_Q$  iff  $a = 0$ .
- Thm thm:zero-is-a-rational:  $0_Q \in \mathbb{Q}$ .
- Thm thm:one-is-a-rational:  $1_Q \in \mathbb{Q}$ .
- Thm thm:zero-not-one-in-q:  $0_Q \neq 1_Q$ ; depends on  $0_Z \neq 1_Z$ .

## Operations on Rational Classes

- Def def:addition-on-q-classes:  $[a, b] + [c, d] := [ad + bc, bd]$ .
- Lem lem:addition-denominator-nonzero-q:  $b \neq 0$  and  $d \neq 0$  imply  $bd \neq 0$ .
- Lem lem:addition-on-q-respects-left-equivalence: addition respects fraction equivalence in the left argument.
- Lem lem:addition-on-q-respects-right-equivalence: addition respects fraction



equivalence in the right argument.

- Lem lem:addition-on-q-respects-equivalence: addition respects fraction equivalence in both arguments.
- Thm thm:addition-on-q-well-defined: addition is a well-defined operation  $\mathbb{Q} \times \mathbb{Q} \rightarrow \mathbb{Q}$ .
- Thm thm:addition-on-q-associative:  $(x + y) + z = x + (y + z)$ .
- Thm thm:addition-on-q-commutative:  $x + y = y + x$ .
- Thm thm:zero-left-additive-identity-on-q:  $0_Q + x = x$ .
- Cor cor:zero-right-additive-identity-on-q:  $x + 0_Q = x$ .
- Def def:negation-on-q-classes:  $-[a, b] := [-a, b]$ .
- Lem lem:negation-on-q-respects-equivalence: negation respects fraction equivalence.
- Thm thm:negation-on-q-well-defined: negation is well-defined on  $\mathbb{Q}$ .
- Thm thm:negation-left-additive-inverse-on-q:  $(-x) + x = 0_Q$ .
- Cor cor:negation-right-additive-inverse-on-q:  $x + (-x) = 0_Q$ .
- Thm thm:every-rational-has-additive-inverse: every rational has a two-sided additive inverse.
- Def def:subtraction-on-q:  $x - y := x + (-y)$ .
- Thm thm:subtraction-on-q-well-defined: subtraction is well-defined.
- Def def:multiplication-on-q-classes:  $[a, b] \cdot [c, d] := [ac, bd]$ .
- Lem lem:multiplication-denominator-nonzero-q:  $b \neq 0$  and  $d \neq 0$  imply  $bd \neq 0$ .
- Lem lem:multiplication-on-q-respects-left-equivalence: multiplication respects equivalence in the left argument.
- Lem lem:multiplication-on-q-respects-right-equivalence: multiplication respects equivalence in the right argument.
- Lem lem:multiplication-on-q-respects-equivalence: multiplication respects equivalence in both arguments.
- Thm thm:multiplication-on-q-well-defined: multiplication is well-defined on  $\mathbb{Q}$ .
- Thm thm:multiplication-on-q-associative:  $(xy)z = x(yz)$ .

- Thm `thm:multiplication-on-q-commutative`:  $xy = yx$ .
- Thm `thm:one-left-multiplicative-identity-on-q`:  $1_Q x = x$ .
- Cor `cor:one-right-multiplicative-identity-on-q`:  $x 1_Q = x$ .
- Lem `lem:nonzero-predicate-well-defined-q`:  $[a, b] \neq 0_Q$  iff  $a \neq 0$ .
- Def `def:reciprocal-on-q-classes`: for  $a \neq 0$ ,  $[a, b]^{-1} := [b, a]$ .
- Lem `lem:reciprocal-denominator-nonzero-q`:  $a \neq 0$  puts  $(b, a)$  in  $\text{PreQ}$ .
- Lem `lem:reciprocal-on-q-respects-equivalence`: reciprocal respects equivalence on nonzero classes.
- Thm `thm:reciprocal-on-q-well-defined`: reciprocal is a well-defined map  $\mathbb{Q} \setminus \{0_Q\} \rightarrow \mathbb{Q} \setminus \{0_Q\}$ .
- Thm `thm:reciprocal-left-multiplicative-inverse-on-q`:  $x \neq 0_Q \Rightarrow x^{-1}x = 1_Q$ .
- Cor `cor:reciprocal-right-multiplicative-inverse-on-q`:  $x \neq 0_Q \Rightarrow xx^{-1} = 1_Q$ .
- Thm `thm:every-nonzero-rational-has-multiplicative-inverse`: every nonzero rational has a two-sided multiplicative inverse.
- Def `def:division-on-q`: for  $y \neq 0_Q$ ,  $x/y := x \cdot y^{-1}$ .
- Thm `thm:division-on-q-well-defined`: division is well-defined on  $\mathbb{Q} \times (\mathbb{Q} \setminus \{0_Q\})$ .
- Thm `thm:left-distributivity-on-q`:  $x(y + z) = xy + xz$ .
- Cor `cor:right-distributivity-on-q`:  $(y + z)x = yx + zx$ .

## Field Structure and Generic Field Laws

- Thm `thm:q-additive-abelian-group`:  $(\mathbb{Q}, +, 0_Q)$  is an abelian group.
- Thm `thm:q-nonzero-multiplicative-abelian-group`:  $(\mathbb{Q} \setminus \{0_Q\}, \cdot, 1_Q)$  is an abelian group.
- Thm `thm:q-is-a-field`:  $\mathbb{Q}$  is a field.
- Cor `cor:zero-product-on-q`:  $xy = 0_Q$  iff  $x = 0_Q$  or  $y = 0_Q$ ; cites the generic field zero-product theorem.
- Cor `cor:double-negation-on-q`:  $-(-x) = x$ ; cites the generic ring result.

- Cor `cor:negation-of-product-on-q`:  $(-x)y = -(xy) = x(-y)$ ; cites the generic ring result.
- Cor `cor:product-of-negatives-on-q`:  $(-x)(-y) = xy$ ; cites the generic ring result.
- Cor `cor:additive-cancellation-on-q`:  $x + z = y + z \Rightarrow x = y$ ; cites generic group cancellation.
- Cor `cor:multiplicative-cancellation-on-q`:  $z \neq 0_Q$  and  $xz = yz$  imply  $x = y$ ; cites generic field cancellation.
- Cor `cor:uniqueness-additive-inverse-on-q`: additive inverses are unique.
- Cor `cor:uniqueness-multiplicative-inverse-on-q`: multiplicative inverses of nonzero elements are unique.
- Cor `cor:inverse-of-one-is-one-on-q`:  $1_Q^{-1} = 1_Q$ .
- Cor `cor:inverse-of-product-on-q`: if  $x, y \neq 0$ , then  $(xy)^{-1} = x^{-1}y^{-1}$ .
- Cor `cor:inverse-of-inverse-on-q`: if  $x \neq 0$ , then  $(x^{-1})^{-1} = x$ .
- Def `def:integer-power-on-q`:  $x^n$  for  $n \in \mathbb{Z}$ , with  $x \neq 0_Q$  when  $n < 0$ .
- Cor `cor:exponent-addition-on-q`:  $x^{m+n} = x^m x^n$ .
- Cor `cor:power-of-power-on-q`:  $(x^m)^n = x^{mn}$ .
- Cor `cor:power-of-product-on-q`:  $(xy)^n = x^n y^n$ .

## Order on $\mathbb{Q}$

- Def `def:positive-rational`:  $[a, b]$  is positive iff  $ab > 0$  in  $\mathbb{Z}$ .
- Lem `lem:positivity-respects-equivalence-q`: positivity respects fraction equivalence.
- Thm `thm:positivity-on-q-well-defined`: positivity is well-defined on  $\mathbb{Q}$ .
- Def `def:strict-order-on-q`:  $x < y$  iff  $y - x$  is positive.
- Def `def:order-on-q`:  $x \leq y$  iff  $x < y$  or  $x = y$ .
- Prop `prop:product-of-positives-is-positive-q`: positive rationals have positive product.
- Thm `thm:sign-trichotomy-on-q`: exactly one of  $x < 0_Q$ ,  $x = 0_Q$ ,  $0_Q < x$  holds.
- Thm `thm:trichotomy-on-q`: exactly one of  $x < y$ ,  $x = y$ ,  $y < x$  holds.

- Thm thm:strict-order-on-q-irreflexive:  $\text{not } x < x$ .
- Thm thm:strict-order-on-q-asymmetric:  $x < y \Rightarrow \neg(y < x)$ .
- Thm thm:strict-order-on-q-transitive:  $x < y$  and  $y < z$  imply  $x < z$ .
- Thm thm:order-on-q-reflexive:  $x \leq x$ .
- Thm thm:order-on-q-antisymmetric:  $x \leq y$  and  $y \leq x$  imply  $x = y$ .
- Thm thm:order-on-q-transitive:  $x \leq y$  and  $y \leq z$  imply  $x \leq z$ .
- Thm thm:order-on-q-total:  $x \leq y$  or  $y \leq x$ .
- Thm thm:q-totally-ordered-set:  $(\mathbb{Q}, \leq)$  is a total order.
- Thm thm:left-addition-preserves-strict-order-on-q:  $x < y \Rightarrow z + x < z + y$ .
- Cor cor:right-addition-preserves-strict-order-on-q:  $x < y \Rightarrow x + z < y + z$ .
- Thm thm:left-addition-preserves-order-on-q:  $x \leq y \Rightarrow z + x \leq z + y$ .
- Cor cor:right-addition-preserves-order-on-q:  $x \leq y \Rightarrow x + z \leq y + z$ .
- Cor cor:addition-reflects-order-on-q: addition by a fixed rational reflects strict and non-strict order.
- Thm thm:left-positive-multiplication-preserves-strict-order-on-q: positive left multiplication preserves strict order.
- Cor cor:right-positive-multiplication-preserves-strict-order-on-q: positive right multiplication preserves strict order.
- Thm thm:positive-multiplication-preserves-order-on-q: positive multiplication preserves non-strict order.
- Thm thm:negative-multiplication-reverses-strict-order-on-q: negative multiplication reverses strict order.
- Thm thm:reciprocal-reverses-order-on-positives-q:  $0_Q < x < y \Rightarrow 0_Q < y^{-1} < x^{-1}$ .
- Thm thm:sign-of-reciprocal-q: reciprocals preserve positive and negative sign.
- Def def:absolute-value-on-q:  $|x| := x$  if  $0_Q \leq x$ , and  $|x| := -x$  otherwise.
- Thm thm:absolute-value-nonnegative-on-q:  $0_Q \leq |x|$ .
- Thm thm:absolute-value-zero-iff-on-q:  $|x| = 0_Q$  iff  $x = 0_Q$ .

- Thm thm:absolute-value-multiplicative-on-q:  $|xy| = |x||y|$ .
- Thm thm:triangle-inequality-on-q:  $|x + y| \leq |x| + |y|$ .
- Thm thm:absolute-value-of-reciprocal-q:  $x \neq 0_Q \Rightarrow |x^{-1}| = |x|^{-1}$ .
- Thm thm:q-is-an-ordered-field:  $\mathbb{Q}$  is an ordered field.

## Density, Gaps, and Embedding

- Lem lem:midpoint-between-q:  $x < y \Rightarrow x < (x + y)2_Q^{-1} < y$ .
- Thm thm:density-of-q: between any two rationals lies a rational.
- Prop prop:no-least-positive-rational: every positive rational has a smaller positive rational.
- Thm thm:archimedean-property-of-q: for  $0_Q < x$ , some  $n \in \mathbb{N}$  has  $y < nx$ .
- Cor cor:reciprocals-arbitrarily-small-q: reciprocals of naturals become arbitrarily small.
- Thm thm:no-rational-square-root-of-two: no rational square is  $2_Q$ .
- Def def:gap-set-q:  $S := \{q \in \mathbb{Q} : 0_Q < q \wedge q^2 < 2_Q\}$ .
- Prop prop:gap-set-bounded-above-q: the gap set is nonempty and bounded above.
- Thm thm:gap-set-no-rational-supremum-q: the gap set has no rational supremum.
- Cor cor:q-order-incomplete:  $\mathbb{Q}$  is not order-complete.
- Def def:cauchy-sequence-in-q: rational Cauchy sequence.
- Def def:null-sequence-in-q: rational null sequence.
- Lem lem:rational-limit-unique: a convergent rational sequence has at most one rational limit.
- Def def:embedding-z-into-q:  $\iota : \mathbb{Z} \rightarrow \mathbb{Q}, \iota(n) := [n, 1]$ .
- Thm thm:embedding-z-into-q-preserves-zero:  $\iota(0_Z) = 0_Q$ .
- Thm thm:embedding-z-into-q-preserves-one:  $\iota(1_Z) = 1_Q$ .
- Thm thm:embedding-z-into-q-injective:  $\iota(m) = \iota(n) \Rightarrow m = n$ .
- Thm thm:embedding-z-into-q-preserves-addition:  $\iota(m + n) = \iota(m) + \iota(n)$ .
- Thm thm:embedding-z-into-q-preserves-multiplication:  $\iota(mn) = \iota(m)\iota(n)$ .

- Thm `thm:embedding-z-into-q-preserves-order`:  $m \leq n$  iff  $\iota(m) \leq \iota(n)$ .
- Thm `thm:z-embeds-into-q`:  $\iota$  is an injective order-preserving ring homomorphism.
- Thm `thm:q-is-field-of-fractions-of-z`: every rational is a quotient of integer images.
- Cor `cor:embedding-z-into-q-not-surjective`:  $\iota(\mathbb{Z}) \subsetneq \mathbb{Q}$ .
- Rem `rem:why-real-numbers-are-introduced`:  $\mathbb{Q}$  is an ordered field but order-incomplete, motivating  $\mathbb{R}$ .

## Proofs

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Fraction Equivalence Is Reflexive). *For every  $(a, b) \in \text{Pre } \mathbb{Q}$ ,*

$$(a, b) \sim_{\mathbb{Q}} (a, b).$$

*Professional Standard Proof.*

TODO: professional standard proof for fraction-equivalence-reflexive. □

*Detailed Learning Proof.*

TODO: detailed learning proof for fraction-equivalence-reflexive. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Fraction Equivalence Is Symmetric). *For all prefractions  $(a, b), (c, d)$ ,*

$$(a, b) \sim_{\mathbb{Q}} (c, d) \implies (c, d) \sim_{\mathbb{Q}} (a, b).$$

*Professional Standard Proof.*

TODO: professional standard proof for fraction-equivalence-symmetric. □

*Detailed Learning Proof.*

TODO: detailed learning proof for fraction-equivalence-symmetric. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Lemma** (Fraction Equivalence Is Transitive). *For all prefractions  $(a, b), (c, d), (e, f)$ ,*

$$(a, b) \sim_{\mathbb{Q}} (c, d) \wedge (c, d) \sim_{\mathbb{Q}} (e, f) \implies (a, b) \sim_{\mathbb{Q}} (e, f).$$

*Professional Standard Proof.*

TODO: professional standard proof for fraction-equivalence-transitive. □

*Detailed Learning Proof.*

TODO: detailed learning proof for fraction-equivalence-transitive. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Multiplicative Cancellation on  \$\mathbb{Z}\$](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Fraction Equivalence Is an Equivalence Relation). *The relation  $\sim_{\mathbb{Q}}$  is an equivalence relation on  $\text{Pre } \mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for fraction-equivalence-relation. □

*Detailed Learning Proof.*

TODO: detailed learning proof for fraction-equivalence-relation. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Proposition** (Equality Criterion for Rational Classes). *For prefractions  $(a, b), (c, d) \in \text{Pre } \mathbb{Q}$ ,*

$$[a, b] = [c, d] \iff a \cdot d = b \cdot c.$$

*Professional Standard Proof.*

TODO: professional standard proof for equality-criterion-for-rational-classes. □

*Detailed Learning Proof.*

TODO: detailed learning proof for equality-criterion-for-rational-classes. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Vanishing Criterion in  $Q$ ). *For every prefraction  $(a, b) \in \text{Pre } \mathbb{Q}$ ,*

$$[a, b] = 0_{\mathbb{Q}} \iff a = 0_{\mathbb{Z}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for vanishing-criterion-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for vanishing-criterion-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Zero Is a Rational). *The class  $0_{\mathbb{Q}}$  is an element of  $\mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for zero-is-a-rational. □

*Detailed Learning Proof.*

TODO: detailed learning proof for zero-is-a-rational. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (One Is a Rational). *The class  $1_{\mathbb{Q}}$  is an element of  $\mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for one-is-a-rational. □

*Detailed Learning Proof.*

TODO: detailed learning proof for one-is-a-rational. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Zero Is Not One in  $\mathbb{Q}$ ). *In  $\mathbb{Q}$ ,*

$$0_{\mathbb{Q}} \neq 1_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for zero-not-one-in-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for zero-not-one-in-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Zero Is Not One in  \$\mathbb{Z}\$](#)

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Addition Lands in  $PreQ$ ). If  $b \neq 0_{\mathbb{Z}}$  and  $d \neq 0_{\mathbb{Z}}$ , then

$$b \cdot d \neq 0_{\mathbb{Z}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for addition-denominator-nonzero-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-denominator-nonzero-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℤ Has No Zero Divisors](#)



*Remark* (Return). [Return to Theorem](#)

**Lemma** (Addition Respects Fraction Equivalence on the Left). *If  $[a, b] = [a', b']$ , then*

$$[a, b] + [c, d] = [a', b'] + [c, d].$$

*Professional Standard Proof.*

TODO: professional standard proof for addition-on-q-respects-left-equivalence. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-on-q-respects-left-equivalence. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Addition Respects Fraction Equivalence on the Right). *If  $[c, d] = [c', d']$ , then*

$$[a, b] + [c, d] = [a, b] + [c', d'].$$

*Professional Standard Proof.*

TODO: professional standard proof for addition-on-q-respects-right-equivalence. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-on-q-respects-right-equivalence. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Addition Respects Fraction Equivalence). *If  $[a, b] = [a', b']$  and  $[c, d] = [c', d']$ , then*

$$[a, b] + [c, d] = [a', b'] + [c', d'].$$

*Professional Standard Proof.*

TODO: professional standard proof for addition-on-q-respects-equivalence. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-on-q-respects-equivalence. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Addition on  $\mathbb{Q}$  Is Well-Defined). *Addition determines a well-defined binary operation  $\mathbb{Q} \times \mathbb{Q} \rightarrow \mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for addition-on-q-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-on-q-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Addition on  $\mathbb{Q}$  Is Associative). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$(x + y) + z = x + (y + z).$$

*Professional Standard Proof.*

TODO: professional standard proof for addition-on-q-associative. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-on-q-associative. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Addition on  $\mathbb{Q}$  Is Commutative). *For all  $x, y \in \mathbb{Q}$ ,*

$$x + y = y + x.$$

*Professional Standard Proof.*

TODO: professional standard proof for addition-on-q-commutative. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-on-q-commutative. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Zero Is a Left Additive Identity on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} + x = x.$$

*Professional Standard Proof.*

TODO: professional standard proof for zero-left-additive-identity-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for zero-left-additive-identity-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Zero Is a Right Additive Identity on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$x + 0_{\mathbb{Q}} = x.$$

*Professional Standard Proof.*

TODO: professional standard proof for zero-right-additive-identity-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for zero-right-additive-identity-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Theorem** (Zero Is an Additive Identity on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} + x = x \quad \text{and} \quad x + 0_{\mathbb{Q}} = x.$$

*Professional Standard Proof.*

TODO: professional standard proof for zero-additive-identity-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for zero-additive-identity-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Negation Respects Fraction Equivalence). *If  $[a, b] = [a', b']$ , then*

$$-[a, b] = -[a', b'].$$

*Professional Standard Proof.*

TODO: professional standard proof for negation-on-q-respects-equivalence. □

*Detailed Learning Proof.*

TODO: detailed learning proof for negation-on-q-respects-equivalence. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Negation on  $Q$  Is Well-Defined). *Negation determines a well-defined unary operation  $\mathbb{Q} \rightarrow \mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for negation-on-q-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for negation-on-q-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Negation Gives a Left Additive Inverse on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$(-x) + x = 0_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for negation-left-additive-inverse-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for negation-left-additive-inverse-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Negation Gives a Right Additive Inverse on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$x + (-x) = 0_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for negation-right-additive-inverse-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for negation-right-additive-inverse-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Every Rational Has an Additive Inverse). *For every  $x \in \mathbb{Q}$  there exists  $y \in \mathbb{Q}$  such that*

$$x + y = 0_{\mathbb{Q}} \quad \text{and} \quad y + x = 0_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for every-rational-has-additive-inverse. □

*Detailed Learning Proof.*

TODO: detailed learning proof for every-rational-has-additive-inverse. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Subtraction on  $\mathbb{Q}$  Is Well-Defined). *Subtraction determines a well-defined binary operation  $\mathbb{Q} \times \mathbb{Q} \rightarrow \mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for subtraction-on-q-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for subtraction-on-q-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Multiplication Lands in  $PreQ$ ). *If  $b \neq 0_{\mathbb{Z}}$  and  $d \neq 0_{\mathbb{Z}}$ , then*

$$b \cdot d \neq 0_{\mathbb{Z}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-denominator-nonzero-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-denominator-nonzero-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℤ Has No Zero Divisors](#)



*Remark* (Return). [Return to Theorem](#)

**Lemma** (Multiplication Respects Fraction Equivalence on the Left). *If  $[a, b] = [a', b']$ , then*

$$[a, b] \cdot [c, d] = [a', b'] \cdot [c, d].$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-on-q-respects-left-equivalence. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-on-q-respects-left-equivalence. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Multiplication Respects Fraction Equivalence on the Right). *If  $[c, d] = [c', d']$ , then*

$$[a, b] \cdot [c, d] = [a, b] \cdot [c', d'].$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-on-q-respects-right-equivalence. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-on-q-respects-right-equivalence. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Multiplication Respects Fraction Equivalence). *If  $[a, b] = [a', b']$  and  $[c, d] = [c', d']$ , then*

$$[a, b] \cdot [c, d] = [a', b'] \cdot [c', d'].$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-on-q-respects-equivalence. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-on-q-respects-equivalence. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication on  $\mathbb{Q}$  Is Well-Defined). *Multiplication determines a well-defined binary operation  $\mathbb{Q} \times \mathbb{Q} \rightarrow \mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-on-q-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-on-q-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication on  $\mathbb{Q}$  Is Associative). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$(x \cdot y) \cdot z = x \cdot (y \cdot z).$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-on-q-associative. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-on-q-associative. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication on  $\mathbb{Q}$  Is Commutative). *For all  $x, y \in \mathbb{Q}$ ,*

$$x \cdot y = y \cdot x.$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-on-q-commutative. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-on-q-commutative. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (One Is a Left Multiplicative Identity on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$1_{\mathbb{Q}} \cdot x = x.$$

*Professional Standard Proof.*

TODO: professional standard proof for one-left-multiplicative-identity-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for one-left-multiplicative-identity-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (One Is a Right Multiplicative Identity on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$x \cdot 1_{\mathbb{Q}} = x.$$

*Professional Standard Proof.*

TODO: professional standard proof for one-right-multiplicative-identity-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for one-right-multiplicative-identity-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Theorem** (One Is a Multiplicative Identity on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$1_{\mathbb{Q}} \cdot x = x \quad \text{and} \quad x \cdot 1_{\mathbb{Q}} = x.$$

*Professional Standard Proof.*

TODO: professional standard proof for one-multiplicative-identity-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for one-multiplicative-identity-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Nonzero Predicate Is Well-Defined on  $Q$ ). *For every prefraction  $(a, b) \in \text{Pre } \mathbb{Q}$ ,*

$$[a, b] \neq 0_{\mathbb{Q}} \iff a \neq 0_{\mathbb{Z}}.$$

*Thus nonzeroness is independent of the representative.*

*Professional Standard Proof.*

TODO: professional standard proof for nonzero-predicate-well-defined-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for nonzero-predicate-well-defined-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Reciprocal Lands in  $PreQ$ ). *If  $[a, b] \neq 0_{\mathbb{Q}}$ , then  $a \neq 0_{\mathbb{Z}}$ , so  $(b, a)$  is a prefraction.*

*Professional Standard Proof.*

TODO: professional standard proof for reciprocal-denominator-nonzero-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for reciprocal-denominator-nonzero-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Reciprocal Respects Fraction Equivalence). *If  $[a, b] = [a', b']$  and  $[a, b] \neq 0_{\mathbb{Q}}$ , then*

$$[a, b]^{-1} = [a', b']^{-1}.$$

*Professional Standard Proof.*

TODO: professional standard proof for reciprocal-on-q-respects-equivalence. □

*Detailed Learning Proof.*

TODO: detailed learning proof for reciprocal-on-q-respects-equivalence. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Reciprocal on  $\mathbb{Q}$  Is Well-Defined). *Reciprocal determines a well-defined map*

$$\mathbb{Q} \setminus \{0_{\mathbb{Q}}\} \rightarrow \mathbb{Q} \setminus \{0_{\mathbb{Q}}\}.$$

*Professional Standard Proof.*

TODO: professional standard proof for reciprocal-on-q-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for reciprocal-on-q-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Reciprocal Gives a Left Multiplicative Inverse on  $\mathbb{Q}$ ). *For every nonzero  $x \in \mathbb{Q}$ ,*

$$x^{-1} \cdot x = 1_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for reciprocal-left-multiplicative-inverse-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for reciprocal-left-multiplicative-inverse-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Reciprocal Gives a Right Multiplicative Inverse on  $\mathbb{Q}$ ). *For every nonzero  $x \in \mathbb{Q}$ ,*

$$x \cdot x^{-1} = 1_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for reciprocal-right-multiplicative-inverse-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for reciprocal-right-multiplicative-inverse-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Every Nonzero Rational Has a Multiplicative Inverse). *For every  $x \in \mathbb{Q}$  with  $x \neq 0_{\mathbb{Q}}$ , there exists  $y \in \mathbb{Q}$  such that*

$$x \cdot y = 1_{\mathbb{Q}} \quad \text{and} \quad y \cdot x = 1_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for every-nonzero-rational-has-multiplicative-inverse.  $\square$

*Detailed Learning Proof.*

TODO: detailed learning proof for every-nonzero-rational-has-multiplicative-inverse.  $\square$

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Theorem** (Division on  $\mathbb{Q}$  Is Well-Defined). *Division determines a well-defined map*

$$\mathbb{Q} \times (\mathbb{Q} \setminus \{0_{\mathbb{Q}}\}) \rightarrow \mathbb{Q}.$$

*Professional Standard Proof.*

TODO: professional standard proof for division-on-q-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for division-on-q-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Left Distributivity on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x \cdot (y + z) = x \cdot y + x \cdot z.$$

*Professional Standard Proof.*

TODO: professional standard proof for left-distributivity-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for left-distributivity-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Right Distributivity on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$(y + z) \cdot x = y \cdot x + z \cdot x.$$

*Professional Standard Proof.*

TODO: professional standard proof for right-distributivity-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for right-distributivity-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication Distributes over Addition on  $\mathbb{Q}$ ). *Multiplication distributes over addition on both sides in  $\mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-distributes-over-addition-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-distributes-over-addition-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** ( $\mathbb{Q}$  Is an Additive Abelian Group). *The structure  $(\mathbb{Q}, +, 0_{\mathbb{Q}})$  is an abelian group.*

*Professional Standard Proof.*

TODO: professional standard proof for q-additive-abelian-group. □

*Detailed Learning Proof.*

TODO: detailed learning proof for q-additive-abelian-group. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Nonzero  $\mathbb{Q}$  Is a Multiplicative Abelian Group). *The structure  $(\mathbb{Q} \setminus \{0_{\mathbb{Q}}\}, \cdot, 1_{\mathbb{Q}})$  is an abelian group.*

*Professional Standard Proof.*

TODO: professional standard proof for q-nonzero-multiplicative-abelian-group. □

*Detailed Learning Proof.*

TODO: detailed learning proof for q-nonzero-multiplicative-abelian-group. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** ( $\mathbb{Q}$  Is a Field). *The structure  $(\mathbb{Q}, +, \cdot, 0_{\mathbb{Q}}, 1_{\mathbb{Q}})$  is a field.*

*Professional Standard Proof.*

TODO: professional standard proof for  $\mathbb{Q}$ -is-a-field. □

*Detailed Learning Proof.*

TODO: detailed learning proof for  $\mathbb{Q}$ -is-a-field. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Zero Is Not One in  \$\mathbb{Q}\$](#)

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Zero Product on  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$x \cdot y = 0_{\mathbb{Q}} \iff x = 0_{\mathbb{Q}} \text{ or } y = 0_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for zero-product-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for zero-product-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℚ Is a Field](#)



*Remark* (Return). [Return to Theorem](#)

**Corollary** (Double Negation on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$-(-x) = x.$$

*Professional Standard Proof.*

TODO: professional standard proof for double-negation-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for double-negation-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Negation of a Product on  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$(-x) \cdot y = -(x \cdot y) \quad \text{and} \quad x \cdot (-y) = -(x \cdot y).$$

*Professional Standard Proof.*

TODO: professional standard proof for negation-of-product-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for negation-of-product-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Product of Negatives on  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$(-x) \cdot (-y) = x \cdot y.$$

*Professional Standard Proof.*

TODO: professional standard proof for product-of-negatives-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for product-of-negatives-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Additive Cancellation on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x + z = y + z \implies x = y.$$

*Professional Standard Proof.*

TODO: professional standard proof for additive-cancellation-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for additive-cancellation-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Multiplicative Cancellation on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$z \neq 0_{\mathbb{Q}} \wedge x \cdot z = y \cdot z \implies x = y.$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplicative-cancellation-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplicative-cancellation-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Uniqueness of Additive Inverses on  $\mathbb{Q}$ ). *Every additive inverse in  $\mathbb{Q}$  is unique.*

*Professional Standard Proof.*

TODO: professional standard proof for uniqueness-additive-inverse-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for uniqueness-additive-inverse-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Uniqueness of Multiplicative Inverses on  $\mathbb{Q}$ ). *Every multiplicative inverse of a nonzero rational number is unique.*

*Professional Standard Proof.*

TODO: professional standard proof for uniqueness-multiplicative-inverse-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for uniqueness-multiplicative-inverse-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (The Inverse of One Is One on  $\mathbb{Q}$ ). *In  $\mathbb{Q}$ ,*

$$1_{\mathbb{Q}}^{-1} = 1_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for inverse-of-one-is-one-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for inverse-of-one-is-one-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Corollary** (Inverse of a Product on  $Q$ ). *For all nonzero  $x, y \in \mathbb{Q}$ ,*

$$(x \cdot y)^{-1} = x^{-1} \cdot y^{-1}.$$

*Professional Standard Proof.*

TODO: professional standard proof for inverse-of-product-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for inverse-of-product-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Inverse of an Inverse on  $Q$ ). *For every nonzero  $x \in \mathbb{Q}$ ,*

$$(x^{-1})^{-1} = x.$$

*Professional Standard Proof.*

TODO: professional standard proof for inverse-of-inverse-on-q.

□

*Detailed Learning Proof.*

TODO: detailed learning proof for inverse-of-inverse-on-q.

□

*Remark* (Proof structure).

*Remark* (Dependencies).

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Exponent Addition Law on  $\mathbb{Q}$ ). *For every nonzero  $x \in \mathbb{Q}$  and all  $m, n \in \mathbb{Z}$ ,*

$$x^{m+n} = x^m \cdot x^n.$$

*Professional Standard Proof.*

TODO: professional standard proof for exponent-addition-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for exponent-addition-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Power of a Power on  $\mathbb{Q}$ ). *For every nonzero  $x \in \mathbb{Q}$  and all  $m, n \in \mathbb{Z}$ ,*

$$(x^m)^n = x^{m \cdot n}.$$

*Professional Standard Proof.*

TODO: professional standard proof for power-of-power-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for power-of-power-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Power of a Product on  $\mathbb{Q}$ ). *For all nonzero  $x, y \in \mathbb{Q}$  and every  $n \in \mathbb{Z}$ ,*

$$(x \cdot y)^n = x^n \cdot y^n.$$

*Professional Standard Proof.*

TODO: professional standard proof for power-of-product-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for power-of-product-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Integer Power on  \$\mathbb{Q}\$](#)
- [Integer Power on  \$\mathbb{Q}\$](#)
- [Inverse of a Product on  \$\mathbb{Q}\$](#)

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Positivity Respects Fraction Equivalence on  $Q$ ). *If  $[a, b] = [c, d]$ , then*

$$a \cdot b > 0_{\mathbb{Z}} \iff c \cdot d > 0_{\mathbb{Z}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for positivity-respects-equivalence-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for positivity-respects-equivalence-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℤ Is a Totally Ordered Set](#)
- [ℤ Has No Zero Divisors](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Positivity on  $Q$  Is Well-Defined). *The predicate “ $x$  is positive” is well-defined on equivalence classes in  $\mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for positivity-on-q-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for positivity-on-q-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Proposition** (Product of Positives Is Positive on  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < x \wedge 0_{\mathbb{Q}} < y \implies 0_{\mathbb{Q}} < x \cdot y.$$

*Professional Standard Proof.*

TODO: professional standard proof for product-of-positives-is-positive-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for product-of-positives-is-positive-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Theorem** (Sign Trichotomy on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ , exactly one of*

$$x < 0_{\mathbb{Q}}, \quad x = 0_{\mathbb{Q}}, \quad 0_{\mathbb{Q}} < x$$

*holds.*

*Professional Standard Proof.*

TODO: professional standard proof for sign-trichotomy-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for sign-trichotomy-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Trichotomy on  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ , exactly one of*

$$x < y, \quad x = y, \quad y < x$$

*holds.*

*Professional Standard Proof.*

TODO: professional standard proof for trichotomy-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for trichotomy-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Strict Order on  $\mathbb{Q}$  Is Irreflexive). *For every  $x \in \mathbb{Q}$ , not  $x < x$ .*

*Professional Standard Proof.*

TODO: professional standard proof for strict-order-on-q-irreflexive. □

*Detailed Learning Proof.*

TODO: detailed learning proof for strict-order-on-q-irreflexive. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Strict Order on  $\mathbb{Q}$  Is Asymmetric). *For all  $x, y \in \mathbb{Q}$ ,*

$$x < y \implies \neg(y < x).$$

*Professional Standard Proof.*

TODO: professional standard proof for strict-order-on-q-asymmetric. □

*Detailed Learning Proof.*

TODO: detailed learning proof for strict-order-on-q-asymmetric. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Strict Order on  $\mathbb{Q}$  Is Transitive). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x < y \wedge y < z \implies x < z.$$

*Professional Standard Proof.*

TODO: professional standard proof for strict-order-on-q-transitive. □

*Detailed Learning Proof.*

TODO: detailed learning proof for strict-order-on-q-transitive. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Order on  $Q$  Is Reflexive). *For every  $x \in \mathbb{Q}$ ,  $x \leq x$ .*

*Professional Standard Proof.*

TODO: professional standard proof for order-on-q-reflexive. □

*Detailed Learning Proof.*

TODO: detailed learning proof for order-on-q-reflexive. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Order on  $\mathbb{Q}$  Is Antisymmetric). *For all  $x, y \in \mathbb{Q}$ ,*

$$x \leq y \wedge y \leq x \implies x = y.$$

*Professional Standard Proof.*

TODO: professional standard proof for order-on-q-antisymmetric. □

*Detailed Learning Proof.*

TODO: detailed learning proof for order-on-q-antisymmetric. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Order on  $\mathbb{Q}$  Is Transitive). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x \leq y \wedge y \leq z \implies x \leq z.$$

*Professional Standard Proof.*

TODO: professional standard proof for order-on-q-transitive. □

*Detailed Learning Proof.*

TODO: detailed learning proof for order-on-q-transitive. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Theorem** (Order on  $\mathbb{Q}$  Is Total). *For all  $x, y \in \mathbb{Q}$ ,*

$$x \leq y \text{ or } y \leq x.$$

*Professional Standard Proof.*

TODO: professional standard proof for order-on-q-total. □

*Detailed Learning Proof.*

TODO: detailed learning proof for order-on-q-total. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** ( $\mathbb{Q}$  Is a Totally Ordered Set). *The ordered pair  $(\mathbb{Q}, \leq)$  is a totally ordered set.*

*Professional Standard Proof.*

TODO: professional standard proof for q-totally-ordered-set. □

*Detailed Learning Proof.*

TODO: detailed learning proof for q-totally-ordered-set. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Left Addition Preserves Strict Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x < y \implies z + x < z + y.$$

*Professional Standard Proof.*

TODO: professional standard proof for left-addition-preserves-strict-order-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for left-addition-preserves-strict-order-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Right Addition Preserves Strict Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x < y \implies x + z < y + z.$$

*Professional Standard Proof.*

TODO: professional standard proof for right-addition-preserves-strict-order-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for right-addition-preserves-strict-order-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Addition Preserves Strict Order on  $\mathbb{Q}$ ). *Addition preserves strict order on both sides in  $\mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for addition-preserves-strict-order-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-preserves-strict-order-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Left Addition Preserves Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x \leq y \implies z + x \leq z + y.$$

*Professional Standard Proof.*

TODO: professional standard proof for left-addition-preserves-order-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for left-addition-preserves-order-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Right Addition Preserves Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$x \leq y \implies x + z \leq y + z.$$

*Professional Standard Proof.*

TODO: professional standard proof for right-addition-preserves-order-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for right-addition-preserves-order-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Addition Preserves Order on  $\mathbb{Q}$ ). *Addition preserves non-strict order on both sides in  $\mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for addition-preserves-order-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-preserves-order-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Corollary** (Addition Reflects Order on  $\mathbb{Q}$ ). *Addition by a fixed rational reflects both strict and non-strict order.*

*Professional Standard Proof.*

TODO: professional standard proof for addition-reflects-order-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-reflects-order-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Left Multiplication by Positives Preserves Strict Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < z \wedge x < y \implies z \cdot x < z \cdot y.$$

*Professional Standard Proof.*

TODO: professional standard proof for left-positive-multiplication-preserves-strict-order-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for left-positive-multiplication-preserves-strict-order-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Right Multiplication by Positives Preserves Strict Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < z \wedge x < y \implies x \cdot z < y \cdot z.$$

*Professional Standard Proof.*

TODO: professional standard proof for right-positive-multiplication-preserves-strict-order-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for right-positive-multiplication-preserves-strict-order-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication by Positives Preserves Strict Order on  $\mathbb{Q}$ ). *Multiplication by a positive rational preserves strict order on both sides.*

*Professional Standard Proof.*

TODO: professional standard proof for positive-multiplication-preserves-strict-order-on-q.  $\square$

*Detailed Learning Proof.*

TODO: detailed learning proof for positive-multiplication-preserves-strict-order-on-q.  $\square$

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Left Multiplication by Positives Preserves Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < z \wedge x \leq y \implies z \cdot x \leq z \cdot y.$$

*Professional Standard Proof.*

TODO: professional standard proof for left-positive-multiplication-preserves-order-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for left-positive-multiplication-preserves-order-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Right Multiplication by Positives Preserves Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < z \wedge x \leq y \implies x \cdot z \leq y \cdot z.$$

*Professional Standard Proof.*

TODO: professional standard proof for right-positive-multiplication-preserves-order-on-q.  $\square$

*Detailed Learning Proof.*

TODO: detailed learning proof for right-positive-multiplication-preserves-order-on-q.  $\square$

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication by Positives Preserves Order on  $\mathbb{Q}$ ). *Multiplication by a positive rational preserves non-strict order on both sides.*

*Professional Standard Proof.*

TODO: professional standard proof for positive-multiplication-preserves-order-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for positive-multiplication-preserves-order-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication by Negatives Reverses Strict Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$z < 0_{\mathbb{Q}} \wedge x < y \implies z \cdot y < z \cdot x.$$

*Professional Standard Proof.*

TODO: professional standard proof for negative-multiplication-reverses-strict-order-on-q.  $\square$

*Detailed Learning Proof.*

TODO: detailed learning proof for negative-multiplication-reverses-strict-order-on-q.  $\square$

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Corollary** (Multiplication by Negatives Reverses Order on  $\mathbb{Q}$ ). *For all  $x, y, z \in \mathbb{Q}$ ,*

$$z < 0_{\mathbb{Q}} \wedge x \leq y \implies z \cdot y \leq z \cdot x.$$

*Professional Standard Proof.*

TODO: professional standard proof for negative-multiplication-reverses-order-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for negative-multiplication-reverses-order-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Sign of a Reciprocal on  $\mathbb{Q}$ ). *For every nonzero  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < x \implies 0_{\mathbb{Q}} < x^{-1}, \quad x < 0_{\mathbb{Q}} \implies x^{-1} < 0_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for sign-of-reciprocal-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for sign-of-reciprocal-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Reciprocal Reverses Order on Positives in  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < x < y \implies 0_{\mathbb{Q}} < y^{-1} < x^{-1}.$$

*Professional Standard Proof.*

TODO: professional standard proof for reciprocal-reverses-order-on-positives-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for reciprocal-reverses-order-on-positives-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Absolute Value Is Nonnegative on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} \leq |x|.$$

*Professional Standard Proof.*

TODO: professional standard proof for absolute-value-nonnegative-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for absolute-value-nonnegative-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Absolute Value Vanishes Only at Zero on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$|x| = 0_{\mathbb{Q}} \iff x = 0_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for absolute-value-zero-iff-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for absolute-value-zero-iff-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Absolute Value Is Multiplicative on  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$|x \cdot y| = |x| \cdot |y|.$$

*Professional Standard Proof.*

TODO: professional standard proof for absolute-value-multiplicative-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for absolute-value-multiplicative-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Triangle Inequality on  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$|x + y| \leq |x| + |y|.$$

*Professional Standard Proof.*

TODO: professional standard proof for triangle-inequality-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for triangle-inequality-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Absolute Value of a Reciprocal on  $\mathbb{Q}$ ). *For every nonzero  $x \in \mathbb{Q}$ ,*

$$|x^{-1}| = |x|^{-1}.$$

*Professional Standard Proof.*

TODO: professional standard proof for absolute-value-of-reciprocal-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for absolute-value-of-reciprocal-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Theorem** ( $\mathbb{Q}$  Is an Ordered Field). *The structure*

$$(\mathbb{Q}, +, \cdot, 0_{\mathbb{Q}}, 1_{\mathbb{Q}}, \leq)$$

*is an ordered field.*

*Professional Standard Proof.*

TODO: professional standard proof for  $\mathbb{Q}$ -is-an-ordered-field. □

*Detailed Learning Proof.*

TODO: detailed learning proof for  $\mathbb{Q}$ -is-an-ordered-field. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℚ Is a Field](#)
- [ℚ Is a Totally Ordered Set](#)
- [Addition Preserves Order on ℚ](#)
- [Multiplication by Positives Preserves Order on ℚ](#)

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Ordered Field Laws Hold on  $\mathbb{Q}$ ). *Every theorem that depends only on the ordered-field axioms applies to  $\mathbb{Q}$  with its constructed operations and order.*

*Professional Standard Proof.*

TODO: professional standard proof for ordered-field-laws-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for ordered-field-laws-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℚ Is an Ordered Field](#)

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Squares Are Nonnegative on  $\mathbb{Q}$ ). *For every  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} \leq x^2.$$

*Professional Standard Proof.*

TODO: professional standard proof for squares-nonnegative-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for squares-nonnegative-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (One Is Positive in  $\mathbb{Q}$ ). *In  $\mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < 1_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for one-positive-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for one-positive-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Natural Numbers Are Positive in  $\mathbb{Q}$ ). *Every natural number embedded in  $\mathbb{Q}$  is positive.*

*Professional Standard Proof.*

TODO: professional standard proof for natural-numbers-positive-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for natural-numbers-positive-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Positive Rationals Have Positive Inverses). *For every  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < x \implies 0_{\mathbb{Q}} < x^{-1}.$$

*Professional Standard Proof.*

TODO: professional standard proof for positive-rationals-have-positive-inverses. □

*Detailed Learning Proof.*

TODO: detailed learning proof for positive-rationals-have-positive-inverses. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Fraction Comparison in  $\mathbb{Q}$ ). *For rational classes with positive denominator product,*

$$[a, b] < [c, d] \iff a \cdot d < c \cdot b.$$

*Professional Standard Proof.*

TODO: professional standard proof for fraction-comparison-on-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for fraction-comparison-on-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℚ Is an Ordered Field](#)
- [Product of Positives Is Positive on ℚ](#)
- [One in ℚ](#)
- [Reciprocal Reverses Order on Positives in ℚ](#)
- [Positive Multiplication Preserves Order on ℚ](#)
- [Reciprocal Reverses Order on Positives in ℚ](#)

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Two Is Nonzero in  $\mathbb{Q}$ ). *In  $\mathbb{Q}$ ,*

$$2_{\mathbb{Q}} \neq 0_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for two-nonzero-in-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for two-nonzero-in-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [One Is Positive in  \$\mathbb{Q}\$](#)
- [Trichotomy on  \$\mathbb{Q}\$](#)



*Remark* (Return). [Return to Theorem](#)

**Lemma** (Midpoint Lies Strictly Between Rationals). *For all  $x, y \in \mathbb{Q}$ ,*

$$x < y \implies x < (x + y) \cdot 2_{\mathbb{Q}}^{-1} < y.$$

*Professional Standard Proof.*

TODO: professional standard proof for midpoint-between-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for midpoint-between-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℚ Is an Ordered Field](#)
- [Two Is Nonzero in ℚ](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Density of  $\mathbb{Q}$  in Itself). *For all  $x, y \in \mathbb{Q}$ ,*

$$x < y \implies \exists r \in \mathbb{Q} (x < r < y).$$

*Professional Standard Proof.*

TODO: professional standard proof for density-of-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for density-of-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Archimedean Property of  $\mathbb{Q}$ ). *For all  $x, y \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < x \implies \exists n \in \mathbb{N} (y < n \cdot x).$$

*Professional Standard Proof.*

TODO: professional standard proof for archimedean-property-of-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for archimedean-property-of-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Well-Ordering Principle](#)
- [Division Algorithm on  \$\mathbb{Z}\$](#)
- [\$\mathbb{Z}\$  Is a Totally Ordered Set](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** ( $\mathbb{Q}$  Has Arbitrarily Close Distinct Points). *For every  $x \in \mathbb{Q}$  and every  $\varepsilon \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < \varepsilon \implies \exists y \in \mathbb{Q} (y \neq x \text{ and } |x - y| < \varepsilon).$$

*Professional Standard Proof.*

TODO: professional standard proof for q-has-arbitrarily-close-distinct-points. □

*Detailed Learning Proof.*

TODO: detailed learning proof for q-has-arbitrarily-close-distinct-points. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Reciprocals Become Arbitrarily Small in  \$\mathbb{Q}\$](#)
- [Addition Preserves Strict Order on  \$\mathbb{Q}\$](#)

*Remark* (Return). [Return to Theorem](#)

**Corollary** ( $\mathbb{Q}$  Has No Adjacent Points). *There do not exist rational numbers  $x < y$  with no rational number strictly between them.*

*Professional Standard Proof.*

TODO: professional standard proof for q-has-no-adjacent-points. □

*Detailed Learning Proof.*

TODO: detailed learning proof for q-has-no-adjacent-points. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Density of  \$\mathbb{Q}\$  in Itself](#)

*Remark* (Return). [Return to Theorem](#)

**Proposition** ( $\mathbb{Q}$  Has No Least Positive Rational). *For every  $x \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < x \implies \exists y \in \mathbb{Q} (0_{\mathbb{Q}} < y < x).$$

*Professional Standard Proof.*

TODO: professional standard proof for no-least-positive-rational. □

*Detailed Learning Proof.*

TODO: detailed learning proof for no-least-positive-rational. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Density of  \$\mathbb{Q}\$  in Itself](#)

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Reciprocals Become Arbitrarily Small in  $\mathbb{Q}$ ). *For every  $\varepsilon \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < \varepsilon \implies \exists n \in \mathbb{N} \ (0_{\mathbb{Q}} < n^{-1} < \varepsilon) .$$

*Professional Standard Proof.*

TODO: professional standard proof for reciprocals-arbitrarily-small-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for reciprocals-arbitrarily-small-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Archimedean Property of  \$\mathbb{Q}\$](#)
- [Sign of a Reciprocal on  \$\mathbb{Q}\$](#)

*Remark* (Return). [Return to Theorem](#)

**Proposition** (One-Sided Rational Approximation). *For every  $x \in \mathbb{Q}$  and every  $\varepsilon \in \mathbb{Q}$ ,*

$$0_{\mathbb{Q}} < \varepsilon \implies \exists y, z \in \mathbb{Q} (x - \varepsilon < y < x < z < x + \varepsilon).$$

*Professional Standard Proof.*

TODO: professional standard proof for one-sided-rational-approximation. □

*Detailed Learning Proof.*

TODO: detailed learning proof for one-sided-rational-approximation. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℚ Has Arbitrarily Close Distinct Points](#)
- [Density of ℚ in Itself](#)



*Remark* (Return). [Return to Theorem](#)

**Theorem** (No Rational Square Root of Two). *There is no rational number  $q \in \mathbb{Q}$  such that*

$$q \cdot q = 2_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for no-rational-square-root-of-two. □

*Detailed Learning Proof.*

TODO: detailed learning proof for no-rational-square-root-of-two. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℚ Is a Field](#)
- [ℤ Is an Integral Domain](#)
- parity or infinite descent facts in  $\mathbb{Z}$

*Remark* (Return). [Return to Theorem](#)

**Proposition** (The Gap Set Is Bounded Above in  $\mathbb{Q}$ ). *The set  $S$  is nonempty and bounded above in  $\mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for gap-set-bounded-above-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for gap-set-bounded-above-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℚ Is an Ordered Field](#)
- [Natural Numbers Are Positive in ℚ](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (The Square-Two Gap Has No Rational Supremum). *The set  $S$  has no least upper bound in  $\mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for gap-set-no-rational-supremum-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for gap-set-no-rational-supremum-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [No Rational Square Root of Two](#)
- [Density of  \$\mathbb{Q}\$  in Itself](#)
- [\$\mathbb{Q}\$  Is an Ordered Field](#)

*Remark* (Return). [Return to Theorem](#)

**Corollary** ( $\mathbb{Q}$  Is Order-Incomplete). *The ordered field  $\mathbb{Q}$  is not order-complete: there exists a nonempty bounded-above subset of  $\mathbb{Q}$  with no supremum in  $\mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for  $\mathbb{Q}$ -order-incomplete. □

*Detailed Learning Proof.*

TODO: detailed learning proof for  $\mathbb{Q}$ -order-incomplete. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [The Gap Set Is Bounded Above in  \$\mathbb{Q}\$](#)
- [The Square-Two Gap Has No Rational Supremum](#)

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Rational Limits Are Unique). *A convergent sequence in  $\mathbb{Q}$  has at most one rational limit.*

*Professional Standard Proof.*

TODO: professional standard proof for rational-limit-unique. □

*Detailed Learning Proof.*

TODO: detailed learning proof for rational-limit-unique. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Triangle Inequality on  \$\mathbb{Q}\$](#)
- [ℚ Is an Ordered Field](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Embedding Preserves Zero). *The embedding sends integer zero to rational zero:*

$$\iota(0_{\mathbb{Z}}) = 0_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for embedding-z-into-q-preserves-zero. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-z-into-q-preserves-zero. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Embedding Preserves One). *The embedding sends integer one to rational one:*

$$\iota(1_{\mathbb{Z}}) = 1_{\mathbb{Q}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for embedding-z-into-q-preserves-one. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-z-into-q-preserves-one. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Embedding  $Z$  into  $Q$  Is Injective). *For all  $m, n \in \mathbb{Z}$ ,*

$$\iota(m) = \iota(n) \implies m = n.$$

*Professional Standard Proof.*

TODO: professional standard proof for embedding-z-into-q-injective. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-z-into-q-injective. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Equality Criterion for Rational Classes](#)
- [One Is a Multiplicative Identity on  \$\mathbb{Z}\$](#)



*Remark* (Return). [Return to Theorem](#)

**Theorem** (Embedding Preserves Addition). *For all  $m, n \in \mathbb{Z}$ ,*

$$\iota(m + n) = \iota(m) + \iota(n).$$

*Professional Standard Proof.*

TODO: professional standard proof for embedding-z-into-q-preserves-addition. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-z-into-q-preserves-addition. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Embedding Preserves Multiplication). *For all  $m, n \in \mathbb{Z}$ ,*

$$\iota(m \cdot n) = \iota(m) \cdot \iota(n).$$

*Professional Standard Proof.*

TODO: professional standard proof for embedding-z-into-q-preserves-multiplication. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-z-into-q-preserves-multiplication. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Embedding Preserves Order). *For all  $m, n \in \mathbb{Z}$ ,*

$$m \leq n \iff \iota(m) \leq \iota(n).$$

*Professional Standard Proof.*

TODO: professional standard proof for embedding-z-into-q-preserves-order. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-z-into-q-preserves-order. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Addition on  \$\mathbb{Q}\$ -Classes](#)
- [Multiplication on  \$\mathbb{Q}\$ -Classes](#)
- [Order on  \$\mathbb{Q}\$](#)
- [\$\mathbb{Z}\$  Is a Totally Ordered Set](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** ( $\mathbb{Z}$  Embeds into  $\mathbb{Q}$ ). *The map  $\iota : \mathbb{Z} \rightarrow \mathbb{Q}$  is an injective order-preserving ring homomorphism.*

*Professional Standard Proof.*

TODO: professional standard proof for z-embeds-into-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for z-embeds-into-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Embedding Preserves Zero](#)
- [Embedding Preserves One](#)
- [Embedding  \$\mathbb{Z}\$  into  \$\mathbb{Q}\$  Is Injective](#)
- [Embedding Preserves Addition](#)
- [Embedding Preserves Multiplication](#)
- [Embedding Preserves Order](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (*Q Is the Field of Fractions of Z*). *For every  $x \in \mathbb{Q}$ , there exist  $m, n \in \mathbb{Z}$  with  $n \neq 0_{\mathbb{Z}}$  such that*

$$x = \iota(m) \cdot \iota(n)^{-1}.$$

*Professional Standard Proof.*

TODO: professional standard proof for q-is-field-of-fractions-of-z. □

*Detailed Learning Proof.*

TODO: detailed learning proof for q-is-field-of-fractions-of-z. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Rational Class Notation](#)
- [Every Nonzero Rational Has a Multiplicative Inverse](#)
- [Vanishing Criterion](#)

*Remark* (Return). [Return to Theorem](#)

**Corollary** (The Embedding of  $\mathbb{Z}$  into  $\mathbb{Q}$  Is Not Surjective). *The image of  $\iota : \mathbb{Z} \rightarrow \mathbb{Q}$  is a proper subset of  $\mathbb{Q}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for embedding-z-into-q-not-surjective. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-z-into-q-not-surjective. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Density of  \$\mathbb{Q}\$  in Itself](#)
- [Embedding Preserves Order](#)

# Chapter 2

## Real Numbers ( $\mathbb{R}$ )



### 2.1 Construction of $\mathbb{R}$

#### Dedekind Cuts

##### Definition

**Definition 2.1.1** (Dedekind Cut). A *Dedekind cut* is a set  $\alpha \subseteq \mathbb{Q}$  satisfying:

$$\alpha \neq \emptyset, \quad \alpha \neq \mathbb{Q},$$

$$p \in \alpha \wedge q < p \implies q \in \alpha,$$

and

$$p \in \alpha \implies \exists r \in \alpha (p < r).$$

*Remark* (Dependencies).

- [Density of  \$\mathbb{Q}\$](#)

##### Definition

**Definition 2.1.2** (The Real Numbers). The real numbers are the set

$$\mathbb{R} := \{\alpha : \alpha \text{ is a Dedekind cut in } \mathbb{Q}\}.$$

*Remark* (Dependencies).

- [Dedekind Cut](#)

## Rational Cuts

### Definition

**Definition 2.1.3** (Rational Cut). For  $q \in \mathbb{Q}$ , define the rational cut

$$q^* := \{r \in \mathbb{Q} : r < q\}.$$

*Remark* (Dependencies).

- [Rational Numbers](#)
- [Order on  \$\mathbb{Q}\$](#)

### Lemma

**Lemma 2.1.4** (Rational Cuts Are Cuts). *For every  $q \in \mathbb{Q}$ , the set  $q^*$  is a Dedekind cut.*

*Remark* (Dependencies).

- [Density of  \$\mathbb{Q}\$](#)
- [No Least Positive Rational](#)

## Zero and One in $\mathbb{R}$

### Definition

**Definition 2.1.5** (Zero in  $\mathbb{R}$ ). Define

$$0_{\mathbb{R}} := (0_{\mathbb{Q}})^* = \{r \in \mathbb{Q} : r < 0_{\mathbb{Q}}\}.$$

*Remark* (Dependencies).

- [Rational Cut](#)
- [Zero in  \$\mathbb{Q}\$](#)

### Definition

**Definition 2.1.6** (One in  $\mathbb{R}$ ). Define

$$1_{\mathbb{R}} := (1_{\mathbb{Q}})^* = \{r \in \mathbb{Q} : r < 1_{\mathbb{Q}}\}.$$

*Remark* (Dependencies).

- [Rational Cut](#)
- [One in  \$\mathbb{Q}\$](#)



### Theorem

**Theorem 2.1.7** (Zero and One Are Reals). *The cuts  $0_{\mathbb{R}}$  and  $1_{\mathbb{R}}$  are elements of  $\mathbb{R}$ .*

*Remark* (Dependencies).

- [The Real Numbers](#)
- [Zero in  \$\mathbb{R}\$](#)
- [One in  \$\mathbb{R}\$](#)
- [Rational Cuts Are Cuts](#)

### Theorem

**Theorem 2.1.8** (Zero Is Not One in  $\mathbb{R}$ ). *In  $\mathbb{R}$ ,*

$$0_{\mathbb{R}} \neq 1_{\mathbb{R}}.$$

*Remark* (Dependencies).

- [Zero in  \$\mathbb{R}\$](#)
- [One in  \$\mathbb{R}\$](#)
- [One Is Positive in  \$\mathbb{Q}\$](#)

## 2.2 Order on $\mathbb{R}$

### Order by Inclusion

#### Definition

**Definition 2.2.1** (Order on  $\mathbb{R}$ ). For cuts  $\alpha, \beta \in \mathbb{R}$ , define

$$\alpha \leq \beta \quad :\Longleftrightarrow \quad \alpha \subseteq \beta.$$

*Remark* (Dependencies).

- [The Real Numbers](#)
- [Subset](#)

#### Definition

**Definition 2.2.2** (Strict Order on  $\mathbb{R}$ ). For cuts  $\alpha, \beta \in \mathbb{R}$ , define

$$\alpha < \beta \quad :\Longleftrightarrow \quad \alpha \subsetneq \beta.$$

*Remark* (Dependencies).

- [The Real Numbers](#)
- [Order on  \$\mathbb{R}\$](#)
- [Proper Subset](#)

#### Theorem

**Theorem 2.2.3** (Order on  $\mathbb{R}$  Is Reflexive). *For every  $\alpha \in \mathbb{R}$ ,  $\alpha \leq \alpha$ .*

*Remark* (Dependencies).

- [Order on  \$\mathbb{R}\$](#)
- [Subset](#)

#### Theorem

**Theorem 2.2.4** (Order on  $\mathbb{R}$  Is Antisymmetric). *For all  $\alpha, \beta \in \mathbb{R}$ ,*

$$\alpha \leq \beta \wedge \beta \leq \alpha \implies \alpha = \beta.$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{R}\$](#)
- [Set Equality](#)
- [Subset](#)

#### Theorem

**Theorem 2.2.5** (Order on  $\mathbb{R}$  Is Transitive). *For all  $\alpha, \beta, \gamma \in \mathbb{R}$ ,*

$$\alpha \leq \beta \wedge \beta \leq \gamma \implies \alpha \leq \gamma.$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{R}\$](#)
- [Subset](#)

#### Theorem

**Theorem 2.2.6** (Comparability of Cuts). *For all cuts  $\alpha, \beta \in \mathbb{R}$ ,*

$$\alpha \subseteq \beta \vee \beta \subseteq \alpha.$$

*Remark* (Dependencies).

- [Dedekind Cut](#)

- [ℚ Is a Totally Ordered Set](#)

#### Theorem

**Theorem 2.2.7** (Order on  $\mathbb{R}$  Is Total). *For all  $\alpha, \beta \in \mathbb{R}$ ,*

$$\alpha \leq \beta \vee \beta \leq \alpha.$$

*Remark* (Dependencies).

- [Order on  \$\mathbb{R}\$](#)
- [Comparability of Cuts](#)

#### Theorem

**Theorem 2.2.8** (Trichotomy on  $\mathbb{R}$ ). *For all  $\alpha, \beta \in \mathbb{R}$ , exactly one of*

$$\alpha < \beta, \quad \alpha = \beta, \quad \beta < \alpha$$

*holds.*

*Remark* (Dependencies).

- [Strict Order on  \$\mathbb{R}\$](#)
- [Order on  \$\mathbb{R}\$  Is Antisymmetric](#)
- [Order on  \$\mathbb{R}\$  Is Total](#)

#### Theorem

**Theorem 2.2.9** ( $\mathbb{R}$  Is a Totally Ordered Set). *The ordered pair  $(\mathbb{R}, \leq)$  is a total order.*

*Remark* (Dependencies).

- [Ordered Set](#)
- [Total Order](#)
- [Order on  \$\mathbb{R}\$](#)
- [Order on  \$\mathbb{R}\$  Is Reflexive](#)
- [Order on  \$\mathbb{R}\$  Is Antisymmetric](#)
- [Order on  \$\mathbb{R}\$  Is Transitive](#)
- [Order on  \$\mathbb{R}\$  Is Total](#)

## 2.3 Completeness of $\mathbb{R}$

### Definition

**Definition 2.3.1** (Upper Bound in  $\mathbb{R}$ ). Let  $S \subseteq \mathbb{R}$ . A cut  $\gamma \in \mathbb{R}$  is an upper bound of  $S$  if

$$\forall \alpha \in S (\alpha \leq \gamma).$$

The set  $S$  is bounded above if it has an upper bound in  $\mathbb{R}$ .

*Remark* (Dependencies).

- [The Real Numbers](#)
- [Order on  \$\mathbb{R}\$](#)
- [Subset](#)

### Definition

**Definition 2.3.2** (Supremum in  $\mathbb{R}$ ). For  $S \subseteq \mathbb{R}$ , a cut  $\sigma$  is the supremum of  $S$  if  $\sigma$  is an upper bound of  $S$  and every upper bound  $\gamma$  of  $S$  satisfies  $\sigma \leq \gamma$ .

*Remark* (Dependencies).

- [Upper Bound in  \$\mathbb{R}\$](#)
- [Order on  \$\mathbb{R}\$](#)

### Definition

**Definition 2.3.3** (Least-Upper-Bound Property on  $\mathbb{R}$ ). The ordered set  $\mathbb{R}$  has the *least-upper-bound property* if every nonempty subset of  $\mathbb{R}$  that is bounded above has a supremum in  $\mathbb{R}$ .

*Remark* (Dependencies).

- [The Real Numbers](#)
- [Order on  \$\mathbb{R}\$](#)
- [Upper Bound in  \$\mathbb{R}\$](#)
- [Supremum in  \$\mathbb{R}\$](#)

### Lemma

**Lemma 2.3.4** (Union of a Bounded Nonempty Family of Cuts Is a Cut). *If  $S \subseteq \mathbb{R}$  is nonempty and bounded above, then*

$$\bigcup_{\alpha \in S} \alpha$$

*is a Dedekind cut.*

*Remark* (Dependencies).

- [Dedekind Cut](#)
- [Upper Bound in  \$\mathbb{R}\$](#)
- [Union](#)

### Theorem

**Theorem 2.3.5** (Least-Upper-Bound Property). *Every nonempty subset  $S \subseteq \mathbb{R}$  that is bounded above has a least upper bound, namely*

$$\sup S = \bigcup_{\alpha \in S} \alpha.$$

*Remark* (Dependencies).

- [Least-Upper-Bound Property on  \$\mathbb{R}\$](#)
- [Union of a Bounded Nonempty Family of Cuts Is a Cut](#)
- [Supremum in  \$\mathbb{R}\$](#)

### Corollary

**Corollary 2.3.6** (Greatest-Lower-Bound Property). *Every nonempty subset  $S \subseteq \mathbb{R}$  that is bounded below has a greatest lower bound.*

*Remark* (Dependencies).

- [Least-Upper-Bound Property](#)

## 2.4 Addition on $\mathbb{R}$

### Definition

**Definition 2.4.1** (Addition of Cuts). For cuts  $\alpha, \beta \in \mathbb{R}$ , define

$$\alpha + \beta := \{p + q : p \in \alpha, q \in \beta\}.$$

*Remark* (Dependencies).

- [The Real Numbers](#)
- [Addition on  \$\mathbb{Q}\$](#)

### Lemma

**Lemma 2.4.2** (Sum of Cuts Is a Cut). *If  $\alpha, \beta \in \mathbb{R}$ , then  $\alpha + \beta$  is a Dedekind cut.*

*Remark* (Dependencies).

- [Addition on  \$\mathbb{R}\$](#)
- [Dedekind Cut](#)
- [Q Is an Ordered Field](#)

#### Theorem

**Theorem 2.4.3** (Addition on  $\mathbb{R}$  Is Associative). *For all  $\alpha, \beta, \gamma \in \mathbb{R}$ ,*

$$(\alpha + \beta) + \gamma = \alpha + (\beta + \gamma).$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{R}\$](#)
- [Sum of Cuts Is a Cut](#)
- [Addition on  \$\mathbb{Q}\$  Is Associative](#)

#### Theorem

**Theorem 2.4.4** (Addition on  $\mathbb{R}$  Is Commutative). *For all  $\alpha, \beta \in \mathbb{R}$ ,*

$$\alpha + \beta = \beta + \alpha.$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{R}\$](#)
- [Sum of Cuts Is a Cut](#)
- [Addition on  \$\mathbb{Q}\$  Is Commutative](#)

#### Theorem

**Theorem 2.4.5** (Zero Is an Additive Identity on  $\mathbb{R}$ ). *For every  $\alpha \in \mathbb{R}$ ,*

$$\alpha + 0_{\mathbb{R}} = \alpha.$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{R}\$](#)
- [Zero in  \$\mathbb{R}\$](#)
- [Zero Is an Additive Identity on  \$\mathbb{Q}\$](#)

### Definition

**Definition 2.4.6** (Negation of a Cut). For  $\alpha \in \mathbb{R}$ , define

$$-\alpha := \{p \in \mathbb{Q} : \exists r \in \mathbb{Q} (r > 0 \wedge -p - r \notin \alpha)\}.$$

*Remark* (Dependencies).

- [The Real Numbers](#)
- [Rational Numbers](#)
- [Positive Rational](#)
- [Order on  \$\mathbb{Q}\$](#)

### Lemma

**Lemma 2.4.7** (Negation of a Cut Is a Cut). *If  $\alpha \in \mathbb{R}$ , then  $-\alpha$  is a Dedekind cut.*

*Remark* (Dependencies).

- [Negation on  \$\mathbb{R}\$](#)
- [Dedekind Cut](#)
- [\$\mathbb{Q}\$  Is an Ordered Field](#)

### Theorem

**Theorem 2.4.8** (Negation Is an Additive Inverse on  $\mathbb{R}$ ). *For every  $\alpha \in \mathbb{R}$ ,*

$$\alpha + (-\alpha) = 0_{\mathbb{R}}.$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{R}\$](#)
- [Negation on  \$\mathbb{R}\$](#)
- [Negation of a Cut Is a Cut](#)
- [Zero in  \$\mathbb{R}\$](#)

### Theorem

**Theorem 2.4.9** ( $\mathbb{R}$  Is an Additive Abelian Group). *The structure  $(\mathbb{R}, +, 0_{\mathbb{R}})$  is an abelian group.*

*Remark* (Dependencies).

- [Addition on  \$\mathbb{R}\$](#)

- [Addition on  \$\mathbb{R}\$  Is Associative](#)
- [Addition on  \$\mathbb{R}\$  Is Commutative](#)
- [Zero Is an Additive Identity on  \$\mathbb{R}\$](#)
- [Negation Is an Additive Inverse on  \$\mathbb{R}\$](#)

#### Definition

**Definition 2.4.10** (Subtraction on  $\mathbb{R}$ ). For cuts  $\alpha, \beta \in \mathbb{R}$ , define

$$\alpha - \beta := \alpha + (-\beta).$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{R}\$](#)
- [Negation on  \$\mathbb{R}\$](#)

## 2.5 Multiplication on $\mathbb{R}$

#### Definition

**Definition 2.5.1** (Nonnegative Cut). A cut  $\alpha \in \mathbb{R}$  is nonnegative if  $0_{\mathbb{R}} \leq \alpha$  and positive if  $0_{\mathbb{R}} < \alpha$ .

*Remark* (Dependencies).

- [The Real Numbers](#)
- [Zero in  \$\mathbb{R}\$](#)
- [Order on  \$\mathbb{R}\$](#)

#### Definition

**Definition 2.5.2** (Product of Nonnegative Cuts). For nonnegative cuts  $\alpha, \beta \in \mathbb{R}$ , define

$$\alpha \cdot \beta := \{r \in \mathbb{Q} : r < 0_{\mathbb{Q}}\} \cup \{p \cdot q : p \in \alpha, q \in \beta, p \geq 0_{\mathbb{Q}}, q \geq 0_{\mathbb{Q}}\}.$$

*Remark* (Dependencies).

- [Nonnegative Cut](#)
- [Multiplication on  \$\mathbb{Q}\$](#)
- [Order on  \$\mathbb{Q}\$](#)



### Lemma

**Lemma 2.5.3** (Product of Nonnegative Cuts Is a Cut). *If  $\alpha, \beta \geq 0_{\mathbb{R}}$ , then  $\alpha \cdot \beta$  is a Dedekind cut.*

*Remark* (Dependencies).

- [Product of Nonnegative Cuts](#)
- [Dedekind Cut](#)
- [ℚ Is an Ordered Field](#)

### Definition

**Definition 2.5.4** (Absolute Value on ℝ). For  $\alpha \in \mathbb{R}$ , define

$$|\alpha| := \begin{cases} \alpha, & 0_{\mathbb{R}} \leq \alpha, \\ -\alpha, & \alpha < 0_{\mathbb{R}}. \end{cases}$$

*Remark* (Dependencies).

- [Order on ℝ](#)
- [Strict Order on ℝ](#)
- [Zero in ℝ](#)
- [Negation on ℝ](#)

### Theorem

**Theorem 2.5.5** (Triangle Inequality). *For all  $\alpha, \beta \in \mathbb{R}$ ,*

$$|\alpha + \beta| \leq |\alpha| + |\beta|.$$

*Go to proof.*

*Remark* (Dependencies).

- [Absolute Value on ℝ](#)
- [ℝ Is an Ordered Field](#)

### Definition

**Definition 2.5.6** (Multiplication of Cuts). For cuts  $\alpha, \beta \in \mathbb{R}$ , define  $\alpha \cdot \beta$  by sign cases: use  $|\alpha| \cdot |\beta|$  when  $\alpha$  and  $\beta$  have the same sign, use  $-(|\alpha| \cdot |\beta|)$  when they have opposite signs, and use  $0_{\mathbb{R}}$  when either factor is  $0_{\mathbb{R}}$ .

*Remark* (Dependencies).

- [The Real Numbers](#)
- [Product of Nonnegative Cuts](#)
- [Absolute Value on  \$\mathbb{R}\$](#)
- [Negation on  \$\mathbb{R}\$](#)
- [Zero in  \$\mathbb{R}\$](#)

#### Lemma

**Lemma 2.5.7** (Product of Cuts Is a Cut). *If  $\alpha, \beta \in \mathbb{R}$ , then  $\alpha \cdot \beta \in \mathbb{R}$ .*

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{R}\$](#)
- [Product of Nonnegative Cuts](#)
- [Product of Nonnegative Cuts Is a Cut](#)
- [Negation of a Cut Is a Cut](#)

#### Theorem

**Theorem 2.5.8** (Multiplication on  $\mathbb{R}$  Is Associative). *For all  $\alpha, \beta, \gamma \in \mathbb{R}$ ,*

$$(\alpha \cdot \beta) \cdot \gamma = \alpha \cdot (\beta \cdot \gamma).$$

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{R}\$](#)
- [Product of Cuts Is a Cut](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Associative](#)

#### Theorem

**Theorem 2.5.9** (Multiplication on  $\mathbb{R}$  Is Commutative). *For all  $\alpha, \beta \in \mathbb{R}$ ,*

$$\alpha \cdot \beta = \beta \cdot \alpha.$$

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{R}\$](#)
- [Product of Cuts Is a Cut](#)
- [Multiplication on  \$\mathbb{Q}\$  Is Commutative](#)

### Theorem

**Theorem 2.5.10** (One Is a Multiplicative Identity on  $\mathbb{R}$ ). *For every  $\alpha \in \mathbb{R}$ ,*

$$\alpha \cdot 1_{\mathbb{R}} = \alpha.$$

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{R}\$](#)
- [One in  \$\mathbb{R}\$](#)
- [One Is a Multiplicative Identity on  \$\mathbb{Q}\$](#)

### Definition

**Definition 2.5.11** (Reciprocal of a Positive Cut). For  $\alpha > 0_{\mathbb{R}}$ , define

$$\alpha^{-1} := \{r \in \mathbb{Q} : r \leq 0_{\mathbb{Q}}\} \cup \{r \in \mathbb{Q} : r > 0_{\mathbb{Q}} \wedge \exists s \in \mathbb{Q} (s > r \wedge s^{-1} \notin \alpha)\}.$$

*Remark* (Dependencies).

- [The Real Numbers](#)
- [Order on  \$\mathbb{R}\$](#)
- [Order on  \$\mathbb{Q}\$](#)
- [Reciprocal on  \$\mathbb{Q}\$](#)

### Definition

**Definition 2.5.12** (Reciprocal of a Nonzero Cut). For  $\alpha \neq 0_{\mathbb{R}}$ , define  $\alpha^{-1}$  to be  $|\alpha|^{-1}$  if  $\alpha > 0_{\mathbb{R}}$  and  $-(|\alpha|^{-1})$  if  $\alpha < 0_{\mathbb{R}}$ .

*Remark* (Dependencies).

- [Reciprocal of a Positive Cut](#)
- [Absolute Value on  \$\mathbb{R}\$](#)
- [Negation on  \$\mathbb{R}\$](#)
- [Zero in  \$\mathbb{R}\$](#)
- [Order on  \$\mathbb{R}\$](#)

### Lemma

**Lemma 2.5.13** (Reciprocal of a Nonzero Cut Is a Cut). *If  $\alpha \neq 0_{\mathbb{R}}$ , then  $\alpha^{-1} \in \mathbb{R}$ .*

*Remark* (Dependencies).

- [Reciprocal on  \$\mathbb{R}\$](#)
- [Reciprocal of a Positive Cut](#)
- [Dedekind Cut](#)
- [Q Is an Ordered Field](#)

#### Theorem

**Theorem 2.5.14** (Reciprocal Is a Multiplicative Inverse on  $\mathbb{R}$ ). *If  $\alpha \neq 0_{\mathbb{R}}$ , then*

$$\alpha \cdot \alpha^{-1} = 1_{\mathbb{R}}.$$

*Remark* (Dependencies).

- [Reciprocal on  \$\mathbb{R}\$](#)
- [Multiplication on  \$\mathbb{R}\$](#)
- [Reciprocal of a Nonzero Cut Is a Cut](#)
- [One in  \$\mathbb{R}\$](#)

#### Definition

**Definition 2.5.15** (Division on  $\mathbb{R}$ ). For  $\beta \neq 0_{\mathbb{R}}$ , define

$$\alpha/\beta := \alpha \cdot \beta^{-1}.$$

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{R}\$](#)
- [Reciprocal on  \$\mathbb{R}\$](#)
- [Zero in  \$\mathbb{R}\$](#)

#### Theorem

**Theorem 2.5.16** (Distributivity on  $\mathbb{R}$ ). *For all  $\alpha, \beta, \gamma \in \mathbb{R}$ ,*

$$\alpha \cdot (\beta + \gamma) = \alpha \cdot \beta + \alpha \cdot \gamma.$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{R}\$](#)
- [Multiplication on  \$\mathbb{R}\$](#)
- [Multiplication Distributes over Addition on Q](#)

## 2.6 Field and Ordered-Field Structure of $\mathbb{R}$

### Theorem

**Theorem 2.6.1** (Nonzero  $\mathbb{R}$  Is a Multiplicative Abelian Group). *The structure  $(\mathbb{R} \setminus \{0_{\mathbb{R}}\}, \cdot, 1_{\mathbb{R}})$  is an abelian group.*

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{R}\$](#)
- [Multiplication on  \$\mathbb{R}\$  Is Associative](#)
- [Multiplication on  \$\mathbb{R}\$  Is Commutative](#)
- [One Is a Multiplicative Identity on  \$\mathbb{R}\$](#)
- [Reciprocal Is a Multiplicative Inverse on  \$\mathbb{R}\$](#)

### Theorem

**Theorem 2.6.2** ( $\mathbb{R}$  Is a Field). *The structure  $(\mathbb{R}, +, \cdot, 0_{\mathbb{R}}, 1_{\mathbb{R}})$  is a field.*

*Remark* (Dependencies).

- [\$\mathbb{R}\$  Is an Additive Abelian Group](#)
- [Nonzero  \$\mathbb{R}\$  Is a Multiplicative Abelian Group](#)
- [Distributivity on  \$\mathbb{R}\$](#)
- [Zero Is Not One in  \$\mathbb{R}\$](#)

### Corollary

**Corollary 2.6.3** (Field Laws Hold on  $\mathbb{R}$ ). *Every theorem proved generically for fields holds in  $\mathbb{R}$ , including zero product, double negation, sign rules, cancellation, uniqueness of inverses, inverse laws, and exponent laws.*

*Remark* (Dependencies).

## Order Compatibility

### Theorem

**Theorem 2.6.4** (Addition Preserves Order on  $\mathbb{R}$ ). *For all  $\alpha, \beta, \gamma \in \mathbb{R}$ ,*

$$\alpha < \beta \implies \alpha + \gamma < \beta + \gamma,$$

*and the corresponding non-strict statement also holds.*

*Remark* (Dependencies).

- [Addition on  \$\mathbb{R}\$](#)
- [Order on  \$\mathbb{R}\$](#)
- [Strict Order on  \$\mathbb{R}\$](#)
- [Addition Preserves Order on  \$\mathbb{Q}\$](#)

#### Theorem

**Theorem 2.6.5** (Multiplication by Positives Preserves Order on  $\mathbb{R}$ ). *For all  $\alpha, \beta, \gamma \in \mathbb{R}$ ,*

$$0_{\mathbb{R}} < \gamma \wedge \alpha < \beta \implies \gamma \cdot \alpha < \gamma \cdot \beta.$$

*Remark* (Dependencies).

- [Multiplication on  \$\mathbb{R}\$](#)
- [Order on  \$\mathbb{R}\$](#)
- [Strict Order on  \$\mathbb{R}\$](#)
- [Positive Multiplication Preserves Order on  \$\mathbb{Q}\$](#)

#### Theorem

**Theorem 2.6.6** ( $\mathbb{R}$  Is an Ordered Field). *The structure  $(\mathbb{R}, +, \cdot, 0_{\mathbb{R}}, 1_{\mathbb{R}}, \leq)$  is an ordered field.*

*Remark* (Dependencies).

- [\$\mathbb{R}\$  Is a Field](#)
- [\$\mathbb{R}\$  Is a Totally Ordered Set](#)
- [Addition Preserves Order on  \$\mathbb{R}\$](#)
- [Multiplication by Positives Preserves Order on  \$\mathbb{R}\$](#)

#### Corollary

**Corollary 2.6.7** (Ordered-Field Laws Hold on  $\mathbb{R}$ ). *Every theorem proved generically for ordered fields holds in  $\mathbb{R}$ .*

*Remark* (Dependencies).

#### Theorem

**Theorem 2.6.8** ( $\mathbb{R}$  Is a Complete Ordered Field). *The ordered field  $\mathbb{R}$  satisfies the least-upper-bound property.*

*Remark* (Dependencies).

- [Complete Ordered Field](#)
- [ℝ Is an Ordered Field](#)
- [Least-Upper-Bound Property](#)

## 2.7 Embedding ℚ into ℝ

### Definition

**Definition 2.7.1** (Embedding of ℚ into ℝ). Define  $\iota : \mathbb{Q} \rightarrow \mathbb{R}$  by

$$\iota(q) := q^*.$$

*Remark* (Dependencies).

- [Rational Numbers](#)
- [The Real Numbers](#)
- [Rational Cut](#)

### Theorem

**Theorem 2.7.2** (Embedding Preserves Zero and One). *The embedding satisfies*

$$\iota(0_{\mathbb{Q}}) = 0_{\mathbb{R}} \quad \text{and} \quad \iota(1_{\mathbb{Q}}) = 1_{\mathbb{R}}.$$

*Remark* (Dependencies).

- [Embedding of ℚ into ℝ](#)
- [Zero in ℚ](#)
- [One in ℚ](#)
- [Zero in ℝ](#)
- [One in ℝ](#)

### Theorem

**Theorem 2.7.3** (Embedding ℚ into ℝ Is Injective). *For all  $p, q \in \mathbb{Q}$ ,*

$$\iota(p) = \iota(q) \implies p = q.$$

*Remark* (Dependencies).

- [Embedding of ℚ into ℝ](#)
- [Rational Cut](#)

- [ℚ Is a Totally Ordered Set](#)

#### Theorem

**Theorem 2.7.4** (Embedding Preserves Addition and Multiplication). *For all  $p, q \in \mathbb{Q}$ ,*

$$\iota(p + q) = \iota(p) + \iota(q) \quad \text{and} \quad \iota(p \cdot q) = \iota(p) \cdot \iota(q).$$

*Remark* (Dependencies).

- [Embedding of ℚ into ℝ](#)
- [Addition on ℝ](#)
- [Multiplication on ℝ](#)
- [Addition on ℚ](#)
- [Multiplication on ℚ](#)

#### Theorem

**Theorem 2.7.5** (Embedding Preserves Order). *For all  $p, q \in \mathbb{Q}$ ,*

$$p \leq q \iff \iota(p) \leq \iota(q).$$

*Remark* (Dependencies).

- [Embedding of ℚ into ℝ](#)
- [Order on ℚ](#)
- [Order on ℝ](#)
- [Rational Cut](#)

#### Theorem

**Theorem 2.7.6** (ℚ Embeds into ℝ as an Ordered Field). *The map  $\iota$  is an injective order-preserving field homomorphism.*

*Remark* (Dependencies).

- [Embedding of ℚ into ℝ](#)
- [Embedding Preserves Zero and One](#)
- [Embedding ℚ into ℝ Is Injective](#)
- [Embedding Preserves Addition and Multiplication](#)
- [Embedding Preserves Order](#)



### Theorem

**Theorem 2.7.7** (Density of  $\mathbb{Q}$  in  $\mathbb{R}$ ). *For all  $\alpha, \beta \in \mathbb{R}$ ,*

$$\alpha < \beta \implies \exists q \in \mathbb{Q} (\alpha < \iota(q) < \beta).$$

*Remark* (Dependencies).

- [Embedding of  \$\mathbb{Q}\$  into  \$\mathbb{R}\$](#)
- [Strict Order on  \$\mathbb{R}\$](#)
- [Density of  \$\mathbb{Q}\$  in Itself](#)
- [Dedekind Cut](#)

### Theorem

**Theorem 2.7.8** (Archimedean Property of  $\mathbb{R}$ ). *For  $\alpha, \beta \in \mathbb{R}$ ,*

$$0_{\mathbb{R}} < \alpha \implies \exists n \in \mathbb{N} (\beta < \iota(n) \cdot \alpha).$$

*Remark* (Dependencies).

- [Least-Upper-Bound Property](#)

## 2.8 The Gap Filled

### Definition

**Definition 2.8.1** (The Square-Root-of-Two Cut). Define

$$\sqrt{2} := \{r \in \mathbb{Q} : r < 0_{\mathbb{Q}} \vee r \cdot r < 2_{\mathbb{Q}}\}.$$

*Remark* (Dependencies).

- [Rational Numbers](#)
- [Order on  \$\mathbb{Q}\$](#)
- [Multiplication on  \$\mathbb{Q}\$](#)
- [Two in  \$\mathbb{Q}\$](#)

### Lemma

**Lemma 2.8.2** (The Square-Root-of-Two Cut Is a Cut). *The set  $\sqrt{2}$  is a Dedekind cut.*

*Remark* (Dependencies).

- [Density of  \$\mathbb{Q}\$](#)

### Theorem

**Theorem 2.8.3** (Its Square Is Two). *In  $\mathbb{R}$ ,*

$$\sqrt{2} \cdot \sqrt{2} = 2_{\mathbb{R}},$$

*where  $2_{\mathbb{R}} := \iota(2_{\mathbb{Q}})$ .*

*Remark* (Dependencies).

- [The Square-Root-of-Two Cut](#)
- [The Square-Root-of-Two Cut Is a Cut](#)
- [Multiplication on  \$\mathbb{R}\$](#)
- [Embedding of  \$\mathbb{Q}\$  into  \$\mathbb{R}\$](#)
- [Two in  \$\mathbb{Q}\$](#)

### Corollary

**Corollary 2.8.4** ( $\mathbb{R}$  Contains the Supremum  $\mathbb{Q}$  Was Missing). *Let*

$$S = \{q \in \mathbb{Q} : 0_{\mathbb{Q}} < q \wedge q \cdot q < 2_{\mathbb{Q}}\}.$$

*Then*

$$\sqrt{2} = \sup \iota(S).$$

*Thus the bounded-above rational set with no rational supremum has a supremum in  $\mathbb{R}$ .*

*Remark* (Dependencies).

- [The Gap Set in  \$\mathbb{Q}\$](#)
- [The Gap Set Has No Rational Supremum](#)

## 2.9 Characterization of $\mathbb{R}$

### Definition

**Definition 2.9.1** (Complete Ordered Field). A *complete ordered field* is an ordered field satisfying the least-upper-bound property.

*Remark* (Dependencies).

- [Least-Upper-Bound Property on  \$\mathbb{R}\$](#)

### Theorem

**Theorem 2.9.2** (Uniqueness of the Complete Ordered Field). *Any two complete ordered fields are isomorphic by a unique order-isomorphism.*

*Remark* (Dependencies).

- [Complete Ordered Field](#)
- [ℝ Is a Complete Ordered Field](#)

### Corollary

**Corollary 2.9.3** (ℝ Is the Real Numbers). *The field ℝ is determined up to unique order-isomorphism by being a complete ordered field.*

*Remark* (Dependencies).

- [Complete Ordered Field](#)
- [Uniqueness of the Complete Ordered Field](#)
- [ℝ Is a Complete Ordered Field](#)

## 2.10 Transition and Forward

*Remark 2.10.1* (What Completeness Buys). The least-upper-bound property promotes the arbitrary closeness available in  $\mathbb{Q}$  to guaranteed limits in  $\mathbb{R}$ : bounded monotone sequences converge, Cauchy sequences converge, and suprema exist. The general theory of these phenomena belongs to analysis; this chapter supplies the complete ordered field in which they live.

*Remark 2.10.2* (Parallel Constructions). Cauchy sequences, nested intervals, and dyadic expansions give alternative constructions of complete ordered fields. By [Uniqueness of the Complete Ordered Field](#), each yields this same  $\mathbb{R}$  up to unique isomorphism.

## Proofs

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Rational Cuts Are Cuts). *For every  $q \in \mathbb{Q}$ , the set  $q^*$  is a Dedekind cut.*

*Professional Standard Proof.*

TODO: professional standard proof for rational-cut-is-a-cut. □

*Detailed Learning Proof.*

TODO: detailed learning proof for rational-cut-is-a-cut. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Density of  \$\mathbb{Q}\$](#)
- [No Least Positive Rational](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Zero and One Are Reals). *The cuts  $0_{\mathbb{R}}$  and  $1_{\mathbb{R}}$  are elements of  $\mathbb{R}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for zero-one-are-reals. □

*Detailed Learning Proof.*

TODO: detailed learning proof for zero-one-are-reals. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Zero Is Not One in  $R$ ). *In  $\mathbb{R}$ ,*

$$0_{\mathbb{R}} \neq 1_{\mathbb{R}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for zero-not-one-in-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for zero-not-one-in-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Zero in  \$\mathbb{R}\$](#)
- [One in  \$\mathbb{R}\$](#)
- [One Is Positive in  \$\mathbb{Q}\$](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Order on  $R$  Is Reflexive). *For every  $\alpha \in \mathbb{R}$ ,  $\alpha \leq \alpha$ .*

*Professional Standard Proof.*

TODO: professional standard proof for order-on-r-reflexive. □

*Detailed Learning Proof.*

TODO: detailed learning proof for order-on-r-reflexive. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Order on  $\mathbb{R}$  Is Antisymmetric). *For all  $\alpha, \beta \in \mathbb{R}$ ,*

$$\alpha \leq \beta \wedge \beta \leq \alpha \implies \alpha = \beta.$$

*Professional Standard Proof.*

TODO: professional standard proof for order-on-r-antisymmetric. □

*Detailed Learning Proof.*

TODO: detailed learning proof for order-on-r-antisymmetric. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Theorem** (Order on  $\mathbb{R}$  Is Transitive). *For all  $\alpha, \beta, \gamma \in \mathbb{R}$ ,*

$$\alpha \leq \beta \wedge \beta \leq \gamma \implies \alpha \leq \gamma.$$

*Professional Standard Proof.*

TODO: professional standard proof for order-on-r-transitive. □

*Detailed Learning Proof.*

TODO: detailed learning proof for order-on-r-transitive. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Comparability of Cuts). *For all cuts  $\alpha, \beta \in \mathbb{R}$ ,*

$$\alpha \subseteq \beta \vee \beta \subseteq \alpha.$$

*Professional Standard Proof.*

TODO: professional standard proof for comparability-of-cuts.

□

*Detailed Learning Proof.*

TODO: detailed learning proof for comparability-of-cuts.

□

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Order on  $R$  Is Total). *For all  $\alpha, \beta \in \mathbb{R}$ ,*

$$\alpha \leq \beta \vee \beta \leq \alpha.$$

*Professional Standard Proof.*

TODO: professional standard proof for order-on-r-total. □

*Detailed Learning Proof.*

TODO: detailed learning proof for order-on-r-total. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Trichotomy on  $\mathbb{R}$ ). *For all  $\alpha, \beta \in \mathbb{R}$ , exactly one of*

$$\alpha < \beta, \quad \alpha = \beta, \quad \beta < \alpha$$

*holds.*

*Professional Standard Proof.*

TODO: professional standard proof for trichotomy-on-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for trichotomy-on-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (*R Is a Totally Ordered Set*). *The ordered pair  $(\mathbb{R}, \leq)$  is a total order.*

*Professional Standard Proof.*

TODO: professional standard proof for r-totally-ordered-set. □

*Detailed Learning Proof.*

TODO: detailed learning proof for r-totally-ordered-set. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Union of a Bounded Nonempty Family of Cuts Is a Cut). *If  $S \subseteq \mathbb{R}$  is nonempty and bounded above, then*

$$\bigcup_{\alpha \in S} \alpha$$

*is a Dedekind cut.*

*Professional Standard Proof.*

TODO: professional standard proof for union-of-cuts-is-a-cut. □

*Detailed Learning Proof.*

TODO: detailed learning proof for union-of-cuts-is-a-cut. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Least-Upper-Bound Property). *Every nonempty subset  $S \subseteq \mathbb{R}$  that is bounded above has a least upper bound, namely*

$$\sup S = \bigcup_{\alpha \in S} \alpha.$$

*Professional Standard Proof.*

TODO: professional standard proof for lub-property-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for lub-property-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Greatest-Lower-Bound Property). *Every nonempty subset  $S \subseteq \mathbb{R}$  that is bounded below has a greatest lower bound.*

*Professional Standard Proof.*

TODO: professional standard proof for glb-property-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for glb-property-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Least-Upper-Bound Property](#)



*Remark* (Return). [Return to Theorem](#)

**Lemma** (Sum of Cuts Is a Cut). *If  $\alpha, \beta \in \mathbb{R}$ , then  $\alpha + \beta$  is a Dedekind cut.*

*Professional Standard Proof.*

TODO: professional standard proof for sum-of-cuts-is-a-cut. □

*Detailed Learning Proof.*

TODO: detailed learning proof for sum-of-cuts-is-a-cut. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Addition on  $\mathbb{R}$  Is Associative). *For all  $\alpha, \beta, \gamma \in \mathbb{R}$ ,*

$$(\alpha + \beta) + \gamma = \alpha + (\beta + \gamma).$$

*Professional Standard Proof.*

TODO: professional standard proof for addition-on-r-associative. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-on-r-associative. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Addition on  $\mathbb{R}$  Is Commutative). *For all  $\alpha, \beta \in \mathbb{R}$ ,*

$$\alpha + \beta = \beta + \alpha.$$

*Professional Standard Proof.*

TODO: professional standard proof for addition-on-r-commutative. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-on-r-commutative. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Zero Is an Additive Identity on  $\mathbb{R}$ ). *For every  $\alpha \in \mathbb{R}$ ,*

$$\alpha + 0_{\mathbb{R}} = \alpha.$$

*Professional Standard Proof.*

TODO: professional standard proof for zero-additive-identity-on-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for zero-additive-identity-on-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Negation of a Cut Is a Cut). *If  $\alpha \in \mathbb{R}$ , then  $-\alpha$  is a Dedekind cut.*

*Professional Standard Proof.*

TODO: professional standard proof for negation-of-cut-is-a-cut. □

*Detailed Learning Proof.*

TODO: detailed learning proof for negation-of-cut-is-a-cut. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Negation Is an Additive Inverse on  $\mathbb{R}$ ). *For every  $\alpha \in \mathbb{R}$ ,*

$$\alpha + (-\alpha) = 0_{\mathbb{R}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for negation-additive-inverse-on-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for negation-additive-inverse-on-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** ( $\mathbb{R}$  Is an Additive Abelian Group). *The structure  $(\mathbb{R}, +, 0_{\mathbb{R}})$  is an abelian group.*

*Professional Standard Proof.*

TODO: professional standard proof for r-additive-abelian-group. □

*Detailed Learning Proof.*

TODO: detailed learning proof for r-additive-abelian-group. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Product of Nonnegative Cuts Is a Cut). *If  $\alpha, \beta \geq 0_{\mathbb{R}}$ , then  $\alpha \cdot \beta$  is a Dedekind cut.*

*Professional Standard Proof.*

TODO: professional standard proof for product-nonnegative-is-a-cut. □

*Detailed Learning Proof.*

TODO: detailed learning proof for product-nonnegative-is-a-cut. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Lemma** (Product of Cuts Is a Cut). *If  $\alpha, \beta \in \mathbb{R}$ , then  $\alpha \cdot \beta \in \mathbb{R}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for product-of-cuts-is-a-cut. □

*Detailed Learning Proof.*

TODO: detailed learning proof for product-of-cuts-is-a-cut. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Triangle Inequality). *For all  $\alpha, \beta \in \mathbb{R}$ ,*

$$|\alpha + \beta| \leq |\alpha| + |\beta|.$$

*Professional Standard Proof.*

TODO: professional standard proof for triangle-inequality. □

*Detailed Learning Proof.*

TODO: detailed learning proof for triangle-inequality. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Absolute Value on  \$\mathbb{R}\$](#)
- [ℝ Is an Ordered Field](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication on  $\mathbb{R}$  Is Associative). *For all  $\alpha, \beta, \gamma \in \mathbb{R}$ ,*

$$(\alpha \cdot \beta) \cdot \gamma = \alpha \cdot (\beta \cdot \gamma).$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-on-r-associative. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-on-r-associative. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication on  $R$  Is Commutative). *For all  $\alpha, \beta \in \mathbb{R}$ ,*

$$\alpha \cdot \beta = \beta \cdot \alpha.$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-on-r-commutative. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-on-r-commutative. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (One Is a Multiplicative Identity on  $R$ ). *For every  $\alpha \in \mathbb{R}$ ,*

$$\alpha \cdot 1_{\mathbb{R}} = \alpha.$$

*Professional Standard Proof.*

TODO: professional standard proof for one-multiplicative-identity-on-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for one-multiplicative-identity-on-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Reciprocal of a Nonzero Cut Is a Cut). *If  $\alpha \neq 0_{\mathbb{R}}$ , then  $\alpha^{-1} \in \mathbb{R}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for reciprocal-is-a-cut. □

*Detailed Learning Proof.*

TODO: detailed learning proof for reciprocal-is-a-cut. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Reciprocal Is a Multiplicative Inverse on  $R$ ). *If  $\alpha \neq 0_{\mathbb{R}}$ , then*

$$\alpha \cdot \alpha^{-1} = 1_{\mathbb{R}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for reciprocal-multiplicative-inverse-on-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for reciprocal-multiplicative-inverse-on-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Distributivity on  $\mathbb{R}$ ). *For all  $\alpha, \beta, \gamma \in \mathbb{R}$ ,*

$$\alpha \cdot (\beta + \gamma) = \alpha \cdot \beta + \alpha \cdot \gamma.$$

*Professional Standard Proof.*

TODO: professional standard proof for distributivity-on-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for distributivity-on-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Theorem** (Nonzero  $\mathbb{R}$  Is a Multiplicative Abelian Group). *The structure  $(\mathbb{R} \setminus \{0_{\mathbb{R}}\}, \cdot, 1_{\mathbb{R}})$  is an abelian group.*

*Professional Standard Proof.*

TODO: professional standard proof for r-nonzero-multiplicative-abelian-group. □

*Detailed Learning Proof.*

TODO: detailed learning proof for r-nonzero-multiplicative-abelian-group. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (*R Is a Field*). *The structure  $(\mathbb{R}, +, \cdot, 0_{\mathbb{R}}, 1_{\mathbb{R}})$  is a field.*

*Professional Standard Proof.*

TODO: professional standard proof for r-is-a-field. □

*Detailed Learning Proof.*

TODO: detailed learning proof for r-is-a-field. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Zero Is Not One in  \$\mathbb{R}\$](#)

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Field Laws Hold on  $R$ ). *Every theorem proved generically for fields holds in  $\mathbb{R}$ , including zero product, double negation, sign rules, cancellation, uniqueness of inverses, inverse laws, and exponent laws.*

*Professional Standard Proof.*

TODO: professional standard proof for field-laws-on-r.

□

*Detailed Learning Proof.*

TODO: detailed learning proof for field-laws-on-r.

□

*Remark* (Proof structure).

*Remark* (Dependencies).

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Addition Preserves Order on  $\mathbb{R}$ ). *For all  $\alpha, \beta, \gamma \in \mathbb{R}$ ,*

$$\alpha < \beta \implies \alpha + \gamma < \beta + \gamma,$$

*and the corresponding non-strict statement also holds.*

*Professional Standard Proof.*

TODO: professional standard proof for addition-preserves-order-on-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-preserves-order-on-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication by Positives Preserves Order on  $\mathbb{R}$ ). *For all  $\alpha, \beta, \gamma \in \mathbb{R}$ ,*

$$0_{\mathbb{R}} < \gamma \wedge \alpha < \beta \implies \gamma \cdot \alpha < \gamma \cdot \beta.$$

*Professional Standard Proof.*

TODO: professional standard proof for positive-multiplication-preserves-order-on-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for positive-multiplication-preserves-order-on-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (*R Is an Ordered Field*). *The structure  $(\mathbb{R}, +, \cdot, 0_{\mathbb{R}}, 1_{\mathbb{R}}, \leq)$  is an ordered field.*

*Professional Standard Proof.*

TODO: professional standard proof for r-is-an-ordered-field.

□

*Detailed Learning Proof.*

TODO: detailed learning proof for r-is-an-ordered-field.

□

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (Ordered-Field Laws Hold on  $R$ ). *Every theorem proved generically for ordered fields holds in  $\mathbb{R}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for ordered-field-laws-on-r.

□

*Detailed Learning Proof.*

TODO: detailed learning proof for ordered-field-laws-on-r.

□

*Remark* (Proof structure).

*Remark* (Dependencies).

*Remark* (Return). [Return to Theorem](#)

**Theorem** (*R Is a Complete Ordered Field*). *The ordered field  $\mathbb{R}$  satisfies the least-upper-bound property.*

*Professional Standard Proof.*

TODO: professional standard proof for r-is-complete-ordered-field.

□

*Detailed Learning Proof.*

TODO: detailed learning proof for r-is-complete-ordered-field.

□

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℝ Is an Ordered Field](#)
- [Least-Upper-Bound Property](#)



*Remark* (Return). [Return to Theorem](#)

**Theorem** (Embedding Preserves Zero and One). *The embedding satisfies*

$$\iota(0_{\mathbb{Q}}) = 0_{\mathbb{R}} \quad \text{and} \quad \iota(1_{\mathbb{Q}}) = 1_{\mathbb{R}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for embedding-q-into-r-preserves-zero-one. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-q-into-r-preserves-zero-one. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Embedding  $Q$  into  $R$  Is Injective). *For all  $p, q \in \mathbb{Q}$ ,*

$$\iota(p) = \iota(q) \implies p = q.$$

*Professional Standard Proof.*

TODO: professional standard proof for embedding-q-into-r-injective. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-q-into-r-injective. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Embedding Preserves Addition and Multiplication). *For all  $p, q \in \mathbb{Q}$ ,*

$$\iota(p + q) = \iota(p) + \iota(q) \quad \text{and} \quad \iota(p \cdot q) = \iota(p) \cdot \iota(q).$$

*Professional Standard Proof.*

TODO: professional standard proof for embedding-q-into-r-preserves-arithmetic. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-q-into-r-preserves-arithmetic. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Embedding Preserves Order). *For all  $p, q \in \mathbb{Q}$ ,*

$$p \leq q \iff \iota(p) \leq \iota(q).$$

*Professional Standard Proof.*

TODO: professional standard proof for embedding-q-into-r-preserves-order. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-q-into-r-preserves-order. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** ( $Q$  Embeds into  $R$  as an Ordered Field). *The map  $\iota$  is an injective order-preserving field homomorphism.*

*Professional Standard Proof.*

TODO: professional standard proof for q-embeds-into-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for q-embeds-into-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Density of  $\mathbb{Q}$  in  $\mathbb{R}$ ). *For all  $\alpha, \beta \in \mathbb{R}$ ,*

$$\alpha < \beta \implies \exists q \in \mathbb{Q} (\alpha < q < \beta).$$

*Professional Standard Proof.*

TODO: professional standard proof for density-of-q-in-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for density-of-q-in-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Archimedean Property of  $\mathbb{R}$ ). For  $\alpha, \beta \in \mathbb{R}$ ,

$$0_{\mathbb{R}} < \alpha \implies \exists n \in \mathbb{N} (\beta < \iota(n) \cdot \alpha).$$

*Professional Standard Proof.*

TODO: professional standard proof for archimedean-property-of-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for archimedean-property-of-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Least-Upper-Bound Property](#)

*Remark* (Return). [Return to Theorem](#)

**Lemma** (The Square-Root-of-Two Cut Is a Cut). *The set  $\sqrt{2}$  is a Dedekind cut.*

*Professional Standard Proof.*

TODO: professional standard proof for sqrt-two-is-a-cut. □

*Detailed Learning Proof.*

TODO: detailed learning proof for sqrt-two-is-a-cut. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Density of  \$\mathbb{Q}\$](#)



*Remark* (Return). [Return to Theorem](#)

**Theorem** (Its Square Is Two). *In  $\mathbb{R}$ ,*

$$\sqrt{2} \cdot \sqrt{2} = 2_{\mathbb{R}},$$

*where  $2_{\mathbb{R}} := \iota(2_{\mathbb{Q}})$ .*

*Professional Standard Proof.*

TODO: professional standard proof for sqrt-two-squared-is-two. □

*Detailed Learning Proof.*

TODO: detailed learning proof for sqrt-two-squared-is-two. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (*R Contains the Supremum Q Was Missing*). *Let*

$$S = \{q \in \mathbb{Q} : 0_{\mathbb{Q}} < q \wedge q \cdot q < 2_{\mathbb{Q}}\}.$$

*Then*

$$\sqrt{2} = \sup \iota(S).$$

*Thus the bounded-above rational set with no rational supremum has a supremum in  $\mathbb{R}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for r-fills-the-gap. □

*Detailed Learning Proof.*

TODO: detailed learning proof for r-fills-the-gap. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [The Gap Set in  \$\mathbb{Q}\$](#)
- [The Gap Set Has No Rational Supremum](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Uniqueness of the Complete Ordered Field). *Any two complete ordered fields are isomorphic by a unique order-isomorphism.*

*Professional Standard Proof.*

TODO: professional standard proof for uniqueness-of-cof. □

*Detailed Learning Proof.*

TODO: detailed learning proof for uniqueness-of-cof. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Corollary** (*R Is the Real Numbers*). *The field  $\mathbb{R}$  is determined up to unique order-isomorphism by being a complete ordered field.*

*Professional Standard Proof.*

TODO: professional standard proof for r-is-the-reals.

□

*Detailed Learning Proof.*

TODO: detailed learning proof for r-is-the-reals.

□

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

# Capstone

# Chapter 3

## Complex Numbers



### 3.1 Construction of $\mathbb{C}$

#### Complex Prepairs

##### The Last Number System Extension

The complex numbers extend the real numbers by adding a formal square root of  $-1$ . Unlike the construction of  $\mathbb{Z}$  from  $\mathbb{W}$  or  $\mathbb{Q}$  from  $\mathbb{Z}$ , no nontrivial identification is needed: ordered pairs of real numbers already have the correct equality. We still use the same construction pattern: representatives, an equivalence relation, equivalence classes, then operations whose well-definedness is checked before they are used.

#### Definition

**Definition 3.1.1** (Complex Prepair). A *complex prepair* is an ordered pair  $(a, b) \in \mathbb{R} \times \mathbb{R}$ , read informally as the expression  $a + bi$ .

*Remark* (Dependencies).

- [Ordered Pair](#)
- [The Real Numbers](#)

#### Definition

**Definition 3.1.2** (Complex Prepair Set). The complex prepair set is

$$\text{Pre } \mathbb{C} := \mathbb{R} \times \mathbb{R}.$$

*Remark* (Dependencies).

- [Cartesian Product](#)
- [The Real Numbers](#)

## Coordinate Equivalence

### Definition

**Definition 3.1.3** (Complex Coordinate Equivalence). For  $(a, b), (c, d) \in \text{Pre } \mathbb{C}$ , define

$$(a, b) \sim_{\mathbb{C}} (c, d) \quad :\Longleftrightarrow \quad a = c \text{ and } b = d.$$

*Remark* (Predicate reading).

$$(a, b) \sim_{\mathbb{C}} (c, d) \Longleftrightarrow (a = c) \wedge (b = d).$$

Thus the complex-number quotient is formally present, but the equivalence classes are singleton coordinate-equality classes.

*Remark* (Dependencies).

- [Relation](#)
- [Complex Prepair Set](#)

### Lemma

**Lemma 3.1.4** (Complex Coordinate Equivalence Is Reflexive). For every  $(a, b) \in \text{Pre } \mathbb{C}$ ,

$$(a, b) \sim_{\mathbb{C}} (a, b).$$

### Lemma

**Lemma 3.1.5** (Complex Coordinate Equivalence Is Symmetric). For all prepairs  $(a, b), (c, d)$ ,

$$(a, b) \sim_{\mathbb{C}} (c, d) \Longrightarrow (c, d) \sim_{\mathbb{C}} (a, b).$$

### Lemma

**Lemma 3.1.6** (Complex Coordinate Equivalence Is Transitive). For all prepairs  $(a, b), (c, d), (e, f)$ ,

$$(a, b) \sim_{\mathbb{C}} (c, d) \wedge (c, d) \sim_{\mathbb{C}} (e, f) \Longrightarrow (a, b) \sim_{\mathbb{C}} (e, f).$$

### Theorem

**Theorem 3.1.7** (Complex Coordinate Equivalence Is an Equivalence Relation). The relation  $\sim_{\mathbb{C}}$  is an equivalence relation on  $\text{Pre } \mathbb{C}$ .

*Remark* (Proof sketch). Reflexivity, symmetry, and transitivity are checked coordinate by coordinate using equality in  $\mathbb{R}$ . Thus this quotient construction has the same formal shape as the constructions of  $\mathbb{Z}$  and  $\mathbb{Q}$ , even though the equivalence classes are singleton classes.

*Remark* (Dependencies).

- [Complex Coordinate Equivalence](#)
- [Complex Coordinate Equivalence Is Reflexive](#)
- [Complex Coordinate Equivalence Is Symmetric](#)
- [Complex Coordinate Equivalence Is Transitive](#)

## Definition of $\mathbb{C}$

### Definition

**Definition 3.1.8** (Complex Numbers). The complex numbers are the quotient

$$\mathbb{C} := \text{Pre } \mathbb{C} / \sim_{\mathbb{C}}.$$

*Remark* (Dependencies).

- [Quotient Set](#)
- [Complex Prepair Set](#)
- [Complex Coordinate Equivalence](#)

### Definition

**Definition 3.1.9** (Complex Class Notation). The notation  $[a, b]_{\mathbb{C}}$  denotes the  $\sim_{\mathbb{C}}$ -equivalence class of the prepair  $(a, b)$ .

*Remark* (Standard form). Since coordinate equivalence is equality of coordinates, every complex number has a unique representative

$$[a, b]_{\mathbb{C}}.$$

We later write this same element as  $a + bi$ .

### Definition

**Definition 3.1.10** (Zero, One, and the Imaginary Unit in  $\mathbb{C}$ ). Define

$$0_{\mathbb{C}} := [0, 0]_{\mathbb{C}}, \quad 1_{\mathbb{C}} := [1, 0]_{\mathbb{C}}, \quad i := [0, 1]_{\mathbb{C}}.$$

*Remark* (Dependencies).

- [Equivalence Class](#)
- [Complex Class Notation](#)
- [The Real Numbers](#)



## 3.2 Operations on $\mathbb{C}$ -Classes

### Addition on $\mathbb{C}$ -Classes

#### Definition

**Definition 3.2.1** (Addition on  $\mathbb{C}$ -Classes). Define addition on representatives by

$$[a, b]_{\mathbb{C}} + [c, d]_{\mathbb{C}} := [a + c, b + d]_{\mathbb{C}}.$$

*Remark* (Dependencies).

- [Complex Class Notation](#)
- [Addition on  \$\mathbb{R}\$](#)

#### Lemma

**Lemma 3.2.2** (Addition Respects Complex Coordinate Equivalence). *If  $[a, b]_{\mathbb{C}} = [a', b']_{\mathbb{C}}$  and  $[c, d]_{\mathbb{C}} = [c', d']_{\mathbb{C}}$ , then*

$$[a, b]_{\mathbb{C}} + [c, d]_{\mathbb{C}} = [a', b']_{\mathbb{C}} + [c', d']_{\mathbb{C}}.$$

*Remark* (Dependencies).

- [Addition on  \$\mathbb{C}\$](#)
- [Complex Coordinate Equivalence](#)

#### Theorem

**Theorem 3.2.3** (Addition on  $\mathbb{C}$  Is Well-Defined). *Addition determines a well-defined binary operation*

$$\mathbb{C} \times \mathbb{C} \rightarrow \mathbb{C}.$$

*Remark* (Proof sketch). Suppose the input classes are represented in two ways. Coordinate equivalence gives  $a = a'$ ,  $b = b'$ ,  $c = c'$ , and  $d = d'$ . Addition in  $\mathbb{R}$  gives  $a + c = a' + c'$  and  $b + d = b' + d'$ , so the resulting complex classes agree.

*Remark* (Dependencies).

- [Function](#)
- [Addition on  \$\mathbb{C}\$](#)
- [Addition Respects Complex Coordinate Equivalence](#)

## Negation and Subtraction

### Definition

**Definition 3.2.4** (Negation on  $\mathbb{C}$ -Classes). Define negation by

$$-[a, b]_{\mathbb{C}} := [-a, -b]_{\mathbb{C}}.$$

### Lemma

**Lemma 3.2.5** (Negation Respects Complex Coordinate Equivalence). If  $[a, b]_{\mathbb{C}} = [a', b']_{\mathbb{C}}$ , then

$$-[a, b]_{\mathbb{C}} = -[a', b']_{\mathbb{C}}.$$

### Theorem

**Theorem 3.2.6** (Negation on  $\mathbb{C}$  Is Well-Defined). Negation determines a well-defined unary operation  $\mathbb{C} \rightarrow \mathbb{C}$ .

*Remark* (Dependencies).

- [Function](#)
- [Negation on  \$\mathbb{C}\$](#)
- [Negation Respects Complex Coordinate Equivalence](#)

*Remark* (Proof sketch). If  $[a, b]_{\mathbb{C}} = [a', b']_{\mathbb{C}}$ , then  $a = a'$  and  $b = b'$ . Hence  $-a = -a'$  and  $-b = -b'$ , so the negated representatives determine the same complex class.

### Definition

**Definition 3.2.7** (Subtraction on  $\mathbb{C}$ ). For  $z, w \in \mathbb{C}$ , define

$$z - w := z + (-w).$$

### Theorem

**Theorem 3.2.8** (Subtraction on  $\mathbb{C}$  Is Well-Defined). Subtraction determines a well-defined binary operation  $\mathbb{C} \times \mathbb{C} \rightarrow \mathbb{C}$ .

*Remark* (Dependencies).

- [Function](#)
- [Subtraction on  \$\mathbb{C}\$](#)
- [Addition on  \$\mathbb{C}\$  Is Well-Defined](#)
- [Negation on  \$\mathbb{C}\$  Is Well-Defined](#)

## Multiplication on $\mathbb{C}$ -Classes

### Definition

**Definition 3.2.9** (Multiplication on  $\mathbb{C}$ -Classes). Define multiplication on representatives by

$$[a, b]_{\mathbb{C}} \cdot [c, d]_{\mathbb{C}} := [ac - bd, ad + bc]_{\mathbb{C}}.$$

*Remark* (Why this formula). The rule is forced by the intended identity  $i^2 = -1$ :

$$(a + bi)(c + di) = (ac - bd) + (ad + bc)i.$$

*Remark* (Dependencies).

- [Complex Class Notation](#)
- [Multiplication on  \$\mathbb{R}\$](#)
- [Addition on  \$\mathbb{R}\$](#)

### Lemma

**Lemma 3.2.10** (Multiplication Respects Complex Coordinate Equivalence). If  $[a, b]_{\mathbb{C}} = [a', b']_{\mathbb{C}}$  and  $[c, d]_{\mathbb{C}} = [c', d']_{\mathbb{C}}$ , then

$$[a, b]_{\mathbb{C}} \cdot [c, d]_{\mathbb{C}} = [a', b']_{\mathbb{C}} \cdot [c', d']_{\mathbb{C}}.$$

### Theorem

**Theorem 3.2.11** (Multiplication on  $\mathbb{C}$  Is Well-Defined). *Multiplication determines a well-defined binary operation*

$$\mathbb{C} \times \mathbb{C} \rightarrow \mathbb{C}.$$

*Remark* (Proof sketch). Changing representatives preserves each coordinate. Therefore the real expressions  $ac - bd$  and  $ad + bc$  are unchanged after replacing  $(a, b)$  by an equivalent prepair and  $(c, d)$  by an equivalent prepair. The product class is independent of the chosen representatives.

*Remark* (Dependencies).

- [Function](#)
- [Multiplication on  \$\mathbb{C}\$](#)
- [Multiplication Respects Complex Coordinate Equivalence](#)

## Conjugation and Inverses

### Definition

**Definition 3.2.12** (Complex Conjugation). For  $z = [a, b]_{\mathbb{C}}$ , define its *complex conjugate* by

$$\bar{z} := [a, -b]_{\mathbb{C}}.$$

### Theorem

**Theorem 3.2.13** (Complex Conjugation Is Well-Defined). *Complex conjugation determines a well-defined unary operation  $\mathbb{C} \rightarrow \mathbb{C}$ .*

*Remark* (Dependencies).

- [Function](#)
- [Complex Conjugation](#)

*Remark* (Proof sketch). Equivalent prepairs have equal first coordinates and equal second coordinates. Keeping the first coordinate and negating the second preserves that equality, so conjugation does not depend on the representative.

### Lemma

**Lemma 3.2.14** (Product with the Conjugate Is Real and Nonnegative). *For  $z = [a, b]_{\mathbb{C}}$ ,*

$$z\bar{z} = [a^2 + b^2, 0]_{\mathbb{C}}.$$

*Moreover, if  $z \neq 0_{\mathbb{C}}$ , then  $a^2 + b^2 \neq 0$ .*

### Definition

**Definition 3.2.15** (Multiplicative Inverse in  $\mathbb{C}$ ). For  $z = [a, b]_{\mathbb{C}} \neq 0_{\mathbb{C}}$ , define

$$z^{-1} := \left[ \frac{a}{a^2 + b^2}, \frac{-b}{a^2 + b^2} \right]_{\mathbb{C}}.$$

### Theorem

**Theorem 3.2.16** (Inversion on  $\mathbb{C}^{\times}$  Is Well-Defined). *Inversion determines a well-defined operation on the nonzero complex numbers.*

*Remark* (Dependencies).

- [Function](#)
- [Multiplicative Inverse in  \$\mathbb{C}\$](#)
- [Product with the Conjugate Is Real and Nonnegative](#)

*Remark* (Proof sketch). If  $z = [a, b]_{\mathbb{C}} \neq 0_{\mathbb{C}}$ , then  $a$  and  $b$  are not both zero, so  $a^2 + b^2 \neq 0$  in  $\mathbb{R}$ . Equivalent representatives have the same coordinates, hence the same denominator and the same inverse coordinates.

## 3.3 Field Structure of $\mathbb{C}$

### Additive Structure

#### Theorem

**Theorem 3.3.1** (Addition on  $\mathbb{C}$  Is Associative). *For all  $z, w, u \in \mathbb{C}$ ,*

$$(z + w) + u = z + (w + u).$$

#### Theorem

**Theorem 3.3.2** (Addition on  $\mathbb{C}$  Is Commutative). *For all  $z, w \in \mathbb{C}$ ,*

$$z + w = w + z.$$

#### Theorem

**Theorem 3.3.3** (Zero Is an Additive Identity on  $\mathbb{C}$ ). *For every  $z \in \mathbb{C}$ ,*

$$0_{\mathbb{C}} + z = z \quad \text{and} \quad z + 0_{\mathbb{C}} = z.$$

#### Theorem

**Theorem 3.3.4** (Every Complex Number Has an Additive Inverse). *For every  $z \in \mathbb{C}$ ,*

$$z + (-z) = 0_{\mathbb{C}} \quad \text{and} \quad (-z) + z = 0_{\mathbb{C}}.$$

### Multiplicative Structure

#### Theorem

**Theorem 3.3.5** (Multiplication on  $\mathbb{C}$  Is Associative). *For all  $z, w, u \in \mathbb{C}$ ,*

$$(zw)u = z(wu).$$

#### Theorem

**Theorem 3.3.6** (Multiplication on  $\mathbb{C}$  Is Commutative). *For all  $z, w \in \mathbb{C}$ ,*

$$zw = wz.$$

### Theorem

**Theorem 3.3.7** (One Is a Multiplicative Identity on  $\mathbb{C}$ ). *For every  $z \in \mathbb{C}$ ,*

$$1_{\mathbb{C}}z = z \quad \text{and} \quad z1_{\mathbb{C}} = z.$$

### Theorem

**Theorem 3.3.8** (Every Nonzero Complex Number Has a Multiplicative Inverse). *For every  $z \in \mathbb{C}$  with  $z \neq 0_{\mathbb{C}}$ ,*

$$zz^{-1} = 1_{\mathbb{C}} \quad \text{and} \quad z^{-1}z = 1_{\mathbb{C}}.$$

## Distributivity and the Field Theorem

### Theorem

**Theorem 3.3.9** (Multiplication Distributes over Addition on  $\mathbb{C}$ ). *For all  $z, w, u \in \mathbb{C}$ ,*

$$z(w + u) = zw + zu \quad \text{and} \quad (w + u)z = wz + uz.$$

### Theorem

**Theorem 3.3.10** ( $\mathbb{C}$  Is a Field). *With the operations defined above,*

$$(\mathbb{C}, +, \cdot, 0_{\mathbb{C}}, 1_{\mathbb{C}})$$

*is a field.*

*Remark* (Dependencies).

- [Addition on  \$\mathbb{C}\$  Is Well-Defined](#)
- [Multiplication on  \$\mathbb{C}\$  Is Well-Defined](#)
- [Every Nonzero Complex Number Has a Multiplicative Inverse](#)

## The Imaginary Unit

### Theorem

**Theorem 3.3.11** (The Imaginary Unit Squares to Negative One). *In  $\mathbb{C}$ ,*

$$i^2 = -1_{\mathbb{C}}.$$

*Remark* (Interpretation). The equation  $i^2 = -1$  is the structural reason for introducing  $\mathbb{C}$ . It makes the polynomial  $x^2 + 1$  have a root, while the field operations remain compatible with the real arithmetic already constructed.

### Theorem

**Theorem 3.3.12** ( $\mathbb{C}$  Is Not an Ordered Field). *There is no order relation on  $\mathbb{C}$  making  $\mathbb{C}$  an ordered field and extending the usual order on  $\mathbb{R}$ .*

*Remark* (Reason). In any ordered field, every square is nonnegative. If  $\mathbb{C}$  were an ordered field extending the real order,  $i^2$  would be nonnegative. But  $i^2 = -1$ , and  $-1 < 0$  in the real subfield.

## 3.4 Embedding $\mathbb{R}$ into $\mathbb{C}$

### The Real Axis

#### Definition

**Definition 3.4.1** (Canonical Embedding of  $\mathbb{R}$  into  $\mathbb{C}$ ). Define

$$\iota : \mathbb{R} \rightarrow \mathbb{C}, \quad \iota(a) := [a, 0]_{\mathbb{C}}.$$

*Remark* (Dependencies).

- [The Real Numbers](#)
- [Complex Numbers](#)
- [Complex Class Notation](#)

### Theorem

**Theorem 3.4.2** (The Real Embedding into  $\mathbb{C}$  Is Injective). *If  $\iota(a) = \iota(b)$ , then  $a = b$ .*

### Theorem

**Theorem 3.4.3** (The Real Embedding Preserves Addition and Multiplication). *For all  $a, b \in \mathbb{R}$ ,*

$$\iota(a + b) = \iota(a) + \iota(b), \quad \iota(ab) = \iota(a)\iota(b).$$

### Theorem

**Theorem 3.4.4** (The Real Embedding Preserves Zero and One).

$$\iota(0) = 0_{\mathbb{C}}, \quad \iota(1) = 1_{\mathbb{C}}.$$

*Remark* (Identification). After these preservation results, it is standard to identify  $a \in \mathbb{R}$  with  $[a, 0]_{\mathbb{C}} \in \mathbb{C}$ . Under this convention,

$$[a, b]_{\mathbb{C}} = [a, 0]_{\mathbb{C}} + [0, b]_{\mathbb{C}} = a + bi.$$

## Real and Imaginary Parts

### Definition

**Definition 3.4.5** (Real and Imaginary Parts). For  $z = [a, b]_{\mathbb{C}}$ , define

$$\operatorname{Re}(z) := a, \quad \operatorname{Im}(z) := b.$$

### Theorem

**Theorem 3.4.6** (Real and Imaginary Parts Are Well-Defined). *The functions*

$$\operatorname{Re}, \operatorname{Im} : \mathbb{C} \rightarrow \mathbb{R}$$

*are well-defined.*

*Remark* (The completed number-system ladder). The construction of  $\mathbb{C}$  completes the standard ladder of number systems in this volume:

$$\mathbb{N} \subseteq \mathbb{W} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}.$$

The final step sacrifices order compatibility in exchange for algebraic power:  $-1$  now has a square root.

## Proofs



*Remark* (Return). [Return to Theorem](#)

**Lemma** (Complex Coordinate Equivalence Is Reflexive). *For every  $(a, b) \in \text{Pre } \mathbb{C}$ ,*

$$(a, b) \sim_{\mathbb{C}} (a, b).$$

*Professional Standard Proof.*

TODO: professional standard proof for complex-coordinate-equivalence-reflexive. □

*Detailed Learning Proof.*

TODO: detailed learning proof for complex-coordinate-equivalence-reflexive. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Complex Coordinate Equivalence Is Symmetric). *For all prepairs  $(a, b), (c, d)$ ,*

$$(a, b) \sim_{\mathbb{C}} (c, d) \implies (c, d) \sim_{\mathbb{C}} (a, b).$$

*Professional Standard Proof.*

TODO: professional standard proof for complex-coordinate-equivalence-symmetric. □

*Detailed Learning Proof.*

TODO: detailed learning proof for complex-coordinate-equivalence-symmetric. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Complex Coordinate Equivalence Is Transitive). *For all prepairs  $(a, b), (c, d), (e, f)$ ,*

$$(a, b) \sim_{\mathbb{C}} (c, d) \wedge (c, d) \sim_{\mathbb{C}} (e, f) \implies (a, b) \sim_{\mathbb{C}} (e, f).$$

*Professional Standard Proof.*

TODO: professional standard proof for complex-coordinate-equivalence-transitive. □

*Detailed Learning Proof.*

TODO: detailed learning proof for complex-coordinate-equivalence-transitive. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Complex Coordinate Equivalence Is an Equivalence Relation). *The relation  $\sim_{\mathbb{C}}$  is an equivalence relation on  $\text{Pre } \mathbb{C}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for complex-coordinate-equivalence-relation. □

*Detailed Learning Proof.*

TODO: detailed learning proof for complex-coordinate-equivalence-relation. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Complex Coordinate Equivalence](#)
- [Complex Coordinate Equivalence Is Reflexive](#)
- [Complex Coordinate Equivalence Is Symmetric](#)
- [Complex Coordinate Equivalence Is Transitive](#)

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Addition Respects Complex Coordinate Equivalence). *If  $[a, b]_{\mathbb{C}} = [a', b']_{\mathbb{C}}$  and  $[c, d]_{\mathbb{C}} = [c', d']_{\mathbb{C}}$ , then*

$$[a, b]_{\mathbb{C}} + [c, d]_{\mathbb{C}} = [a', b']_{\mathbb{C}} + [c', d']_{\mathbb{C}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for addition-on-c-respects-equivalence. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-on-c-respects-equivalence. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Addition on  \$\mathbb{C}\$](#)
- [Complex Coordinate Equivalence](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Addition on  $\mathbb{C}$  Is Well-Defined). *Addition determines a well-defined binary operation*

$$\mathbb{C} \times \mathbb{C} \rightarrow \mathbb{C}.$$

*Professional Standard Proof.*

TODO: professional standard proof for addition-on-c-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-on-c-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Function](#)
- [Addition on  \$\mathbb{C}\$](#)
- [Addition Respects Complex Coordinate Equivalence](#)

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Negation Respects Complex Coordinate Equivalence). *If  $[a, b]_{\mathbb{C}} = [a', b']_{\mathbb{C}}$ , then*

$$-[a, b]_{\mathbb{C}} = -[a', b']_{\mathbb{C}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for negation-on-c-respects-equivalence. □

*Detailed Learning Proof.*

TODO: detailed learning proof for negation-on-c-respects-equivalence. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Negation on  $\mathbb{C}$  Is Well-Defined). *Negation determines a well-defined unary operation  $\mathbb{C} \rightarrow \mathbb{C}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for negation-on-c-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for negation-on-c-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Function](#)
- [Negation on  \$\mathbb{C}\$](#)
- [Negation Respects Complex Coordinate Equivalence](#)



*Remark* (Return). [Return to Theorem](#)

**Theorem** (Subtraction on  $\mathbb{C}$  Is Well-Defined). *Subtraction determines a well-defined binary operation  $\mathbb{C} \times \mathbb{C} \rightarrow \mathbb{C}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for subtraction-on-c-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for subtraction-on-c-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Function](#)
- [Subtraction on  \$\mathbb{C}\$](#)
- [Addition on  \$\mathbb{C}\$  Is Well-Defined](#)
- [Negation on  \$\mathbb{C}\$  Is Well-Defined](#)

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Multiplication Respects Complex Coordinate Equivalence). *If  $[a, b]_{\mathbb{C}} = [a', b']_{\mathbb{C}}$  and  $[c, d]_{\mathbb{C}} = [c', d']_{\mathbb{C}}$ , then*

$$[a, b]_{\mathbb{C}} \cdot [c, d]_{\mathbb{C}} = [a', b']_{\mathbb{C}} \cdot [c', d']_{\mathbb{C}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-on-c-respects-equivalence. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-on-c-respects-equivalence. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication on  $\mathbb{C}$  Is Well-Defined). *Multiplication determines a well-defined binary operation*

$$\mathbb{C} \times \mathbb{C} \rightarrow \mathbb{C}.$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-on-c-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-on-c-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Function](#)
- [Multiplication on  \$\mathbb{C}\$](#)
- [Multiplication Respects Complex Coordinate Equivalence](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Complex Conjugation Is Well-Defined). *Complex conjugation determines a well-defined unary operation  $\mathbb{C} \rightarrow \mathbb{C}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for complex-conjugation-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for complex-conjugation-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Function](#)
- [Complex Conjugation](#)

*Remark* (Return). [Return to Theorem](#)

**Lemma** (Product with the Conjugate Is Real and Nonnegative). *For  $z = [a, b]_{\mathbb{C}}$ ,*

$$z\bar{z} = [a^2 + b^2, 0]_{\mathbb{C}}.$$

*Moreover, if  $z \neq 0_{\mathbb{C}}$ , then  $a^2 + b^2 \neq 0$ .*

*Professional Standard Proof.*

TODO: professional standard proof for complex-product-with-conjugate. □

*Detailed Learning Proof.*

TODO: detailed learning proof for complex-product-with-conjugate. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Inversion on  $\mathbb{C}^\times$  Is Well-Defined). *Inversion determines a well-defined operation on the nonzero complex numbers.*

*Professional Standard Proof.*

TODO: professional standard proof for inversion-on-c-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for inversion-on-c-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Function](#)
- [Multiplicative Inverse in  \$\mathbb{C}\$](#)
- [Product with the Conjugate Is Real and Nonnegative](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Addition on  $\mathbb{C}$  Is Associative). *For all  $z, w, u \in \mathbb{C}$ ,*

$$(z + w) + u = z + (w + u).$$

*Professional Standard Proof.*

TODO: professional standard proof for addition-on-c-associative. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-on-c-associative. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Addition on  $\mathbb{C}$  Is Commutative). *For all  $z, w \in \mathbb{C}$ ,*

$$z + w = w + z.$$

*Professional Standard Proof.*

TODO: professional standard proof for addition-on-c-commutative. □

*Detailed Learning Proof.*

TODO: detailed learning proof for addition-on-c-commutative. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.



*Remark* (Return). [Return to Theorem](#)

**Theorem** (Zero Is an Additive Identity on  $\mathbb{C}$ ). *For every  $z \in \mathbb{C}$ ,*

$$0_{\mathbb{C}} + z = z \quad \text{and} \quad z + 0_{\mathbb{C}} = z.$$

*Professional Standard Proof.*

TODO: professional standard proof for zero-additive-identity-on-c. □

*Detailed Learning Proof.*

TODO: detailed learning proof for zero-additive-identity-on-c. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Every Complex Number Has an Additive Inverse). *For every  $z \in \mathbb{C}$ ,*

$$z + (-z) = 0_{\mathbb{C}} \quad \text{and} \quad (-z) + z = 0_{\mathbb{C}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for every-complex-has-additive-inverse. □

*Detailed Learning Proof.*

TODO: detailed learning proof for every-complex-has-additive-inverse. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication on  $\mathbb{C}$  Is Associative). *For all  $z, w, u \in \mathbb{C}$ ,*

$$(zw)u = z(wu).$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-on-c-associative. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-on-c-associative. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication on  $\mathbb{C}$  Is Commutative). *For all  $z, w \in \mathbb{C}$ ,*

$$zw = wz.$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-on-c-commutative. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-on-c-commutative. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (One Is a Multiplicative Identity on  $\mathbb{C}$ ). *For every  $z \in \mathbb{C}$ ,*

$$1_{\mathbb{C}}z = z \quad \text{and} \quad z1_{\mathbb{C}} = z.$$

*Professional Standard Proof.*

TODO: professional standard proof for one-multiplicative-identity-on-c. □

*Detailed Learning Proof.*

TODO: detailed learning proof for one-multiplicative-identity-on-c. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Every Nonzero Complex Number Has a Multiplicative Inverse). *For every  $z \in \mathbb{C}$  with  $z \neq 0_{\mathbb{C}}$ ,*

$$zz^{-1} = 1_{\mathbb{C}} \quad \text{and} \quad z^{-1}z = 1_{\mathbb{C}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for every-nonzero-complex-has-inverse. □

*Detailed Learning Proof.*

TODO: detailed learning proof for every-nonzero-complex-has-inverse. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Multiplication Distributes over Addition on  $\mathbb{C}$ ). *For all  $z, w, u \in \mathbb{C}$ ,*

$$z(w + u) = zw + zu \quad \text{and} \quad (w + u)z = wz + uz.$$

*Professional Standard Proof.*

TODO: professional standard proof for multiplication-distributes-over-addition-on-c. □

*Detailed Learning Proof.*

TODO: detailed learning proof for multiplication-distributes-over-addition-on-c. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** ( $\mathbb{C}$  Is a Field). *With the operations defined above,*

$$(\mathbb{C}, +, \cdot, 0_{\mathbb{C}}, 1_{\mathbb{C}})$$

*is a field.*

*Professional Standard Proof.*

TODO: professional standard proof for  $\mathbb{C}$ -is-a-field. □

*Detailed Learning Proof.*

TODO: detailed learning proof for  $\mathbb{C}$ -is-a-field. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Addition on  \$\mathbb{C}\$  Is Well-Defined](#)
- [Multiplication on  \$\mathbb{C}\$  Is Well-Defined](#)
- [Every Nonzero Complex Number Has a Multiplicative Inverse](#)



*Remark* (Return). [Return to Theorem](#)

**Theorem** (The Imaginary Unit Squares to Negative One). *In  $\mathbb{C}$ ,*

$$i^2 = -1_{\mathbb{C}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for i-squared-negative-one. □

*Detailed Learning Proof.*

TODO: detailed learning proof for i-squared-negative-one. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (*C Is Not an Ordered Field*). *There is no order relation on  $\mathbb{C}$  making  $\mathbb{C}$  an ordered field and extending the usual order on  $\mathbb{R}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for c-is-not-an-ordered-field.

□

*Detailed Learning Proof.*

TODO: detailed learning proof for c-is-not-an-ordered-field.

□

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (The Real Embedding into  $\mathbb{C}$  Is Injective). *If  $\iota(a) = \iota(b)$ , then  $a = b$ .*

*Professional Standard Proof.*

TODO: professional standard proof for embedding-r-into-c-injective. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-r-into-c-injective. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (The Real Embedding Preserves Addition and Multiplication). *For all  $a, b \in \mathbb{R}$ ,*

$$\iota(a + b) = \iota(a) + \iota(b), \quad \iota(ab) = \iota(a)\iota(b).$$

*Professional Standard Proof.*

TODO: professional standard proof for embedding-r-into-c-preserves-operations. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-r-into-c-preserves-operations. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (The Real Embedding Preserves Zero and One).

$$\iota(0) = 0_{\mathbb{C}}, \quad \iota(1) = 1_{\mathbb{C}}.$$

*Professional Standard Proof.*

TODO: professional standard proof for embedding-r-into-c-preserves-zero-one. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-r-into-c-preserves-zero-one. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Real and Imaginary Parts Are Well-Defined). *The functions*

$$\operatorname{Re}, \operatorname{Im} : \mathbb{C} \rightarrow \mathbb{R}$$

*are well-defined.*

*Professional Standard Proof.*

TODO: professional standard proof for real-imaginary-parts-well-defined. □

*Detailed Learning Proof.*

TODO: detailed learning proof for real-imaginary-parts-well-defined. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: list mathematical dependencies.

# Capstone

# Chapter 4

## Number Lines and Intervals



### 4.1 Number Lines

#### The Geometric Model of an Ordered System

A number line is the geometric realization of a linearly ordered system: points on a line arranged so that “left of” matches the order. The construction so far is algebraic; the number line is the picture that accompanies it, and it becomes exact for  $\mathbb{R}$ .

#### Definition (Number Line)

**Definition 4.1.1** (Number Line). A *number line* for a linearly ordered system  $(A, <_A)$  is a linearly ordered set of points  $L$  together with an order isomorphism  $\lambda : A \rightarrow L$ , so that for all  $a, a' \in A$ ,

$$a <_A a' \iff \lambda(a) \text{ lies to the left of } \lambda(a').$$

*Remark* (Standard quantified statement).

$$\lambda : A \rightarrow L \text{ is a bijection with } \forall a, a' \in A (a <_A a' \iff \lambda(a) \prec \lambda(a')).$$

*Remark* (Interpretation). The number line records only order, not arithmetic: it is the visual home of betweenness and length. For  $\mathbb{R}$  the order isomorphism is onto a full continuum, which is why the real line has no gaps; for  $\mathbb{Q}$  the same picture has holes at every irrational location.

*Remark* (Dependencies).

- [Order Embedding](#)



## 4.2 Intervals on a Number Line

### The Basic Pieces of the Line

Intervals are the natural named pieces of a number line. This section gives each basic interval shape its notation and spells out which endpoints are included.

#### Definition (Open Interval)

**Definition 4.2.1** (Open Interval). Let  $a, b \in \mathbb{R}$  with  $a \leq b$ . The *open interval* with endpoints  $a$  and  $b$  is

$$(a, b) = \{x \in \mathbb{R} : a < x < b\}.$$

*Remark* (Standard quantified statement).

$$(a, b) = \{x \in \mathbb{R} : a < x < b\}.$$

*Remark* (Interpretation). The open interval excludes both endpoints. It is one of the four bounded interval shapes.

*Remark* (Dependencies).

#### Definition (Closed Interval)

**Definition 4.2.2** (Closed Interval). Let  $a, b \in \mathbb{R}$  with  $a \leq b$ . The *closed interval* with endpoints  $a$  and  $b$  is

$$[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}.$$

*Remark* (Standard quantified statement).

$$[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}.$$

*Remark* (Interpretation). The closed interval includes both endpoints. It is one of the four bounded interval shapes.

*Remark* (Dependencies).

#### Definition (Left-Half-Open Interval)

**Definition 4.2.3** (Left-Half-Open Interval). Let  $a, b \in \mathbb{R}$  with  $a \leq b$ . The *left-half-open interval* with endpoints  $a$  and  $b$  is

$$(a, b] = \{x \in \mathbb{R} : a < x \leq b\}.$$

*Remark* (Standard quantified statement).

$$(a, b] = \{x \in \mathbb{R} : a < x \leq b\}.$$

*Remark* (Interpretation). The left-half-open interval excludes the left endpoint, includes the right. It is one of the four bounded interval shapes.

*Remark* (Dependencies).

#### Definition (Right-Half-Open Interval)

**Definition 4.2.4** (Right-Half-Open Interval). Let  $a, b \in \mathbb{R}$  with  $a \leq b$ . The *right-half-open interval* with endpoints  $a$  and  $b$  is

$$[a, b) = \{x \in \mathbb{R} : a \leq x < b\}.$$

*Remark* (Standard quantified statement).

$$[a, b) = \{x \in \mathbb{R} : a \leq x < b\}.$$

*Remark* (Interpretation). The right-half-open interval includes the left endpoint, excludes the right. It is one of the four bounded interval shapes.

*Remark* (Dependencies).

#### Definition (Ray)

**Definition 4.2.5** (Ray). Let  $a \in \mathbb{R}$ . The four *rays* determined by  $a$  are

$$(a, \infty) = \{x : x > a\}, [a, \infty) = \{x : x \geq a\}, (-\infty, a) = \{x : x < a\}, (-\infty, a] = \{x : x \leq a\}. \blacksquare$$

*Remark* (Standard quantified statement).

$$\begin{aligned} (a, \infty) &= \{x \in \mathbb{R} : a < x\}, & [a, \infty) &= \{x \in \mathbb{R} : a \leq x\}, \\ (-\infty, a) &= \{x \in \mathbb{R} : x < a\}, & (-\infty, a] &= \{x \in \mathbb{R} : x \leq a\}. \end{aligned}$$

*Remark* (Interpretation). Rays are the unbounded one-sided intervals. Together with the bounded intervals and the whole line  $(-\infty, \infty) = \mathbb{R}$ , they exhaust the interval shapes.

*Remark* (Dependencies).

## Proofs

## Capstone

# Chapter 5

## Embedding the Number Systems



### 5.1 Embeddings as Structure Preserving Maps

#### From Construction to Identification

The preceding chapters built  $\mathbb{N}$ ,  $\mathbb{W}$ ,  $\mathbb{Z}$ ,  $\mathbb{Q}$ , and  $\mathbb{R}$  one at a time. In that construction-first development the symbol 2 denotes a different object in each system: a Peano element, a whole number, an integer difference object, a rational fraction object, and a real number are literally distinct sets. Nothing in the constructions has made them identical.

An *embedding* is the device that closes this gap without pretending the objects were the same to begin with. It is an injective map that preserves the relevant structure present in the source system — the distinguished constants, the operations, and the order being considered. Once such a map is fixed, the source system is identified with its image inside the target, and only then is the familiar chain  $\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R}$  written down. The inclusion symbol records an identification along an embedding, not a literal subset relation.

#### Definition (Structure-Preserving Map)

**Definition 5.1.1** (Structure-Preserving Map). Let  $A$  and  $B$  be ordered arithmetic systems with a specified common source structure: selected constants, selected operations, and a strict order. A map  $\varphi : A \rightarrow B$  is *structure-preserving* exactly when it preserves each selected constant and, for every selected binary operation  $\star$  and all  $a, a' \in A$ ,

$$\varphi(a \star_A a') = \varphi(a) \star_B \varphi(a'),$$

and  $a <_A a' \implies \varphi(a) <_B \varphi(a')$ .

*Remark* (Standard quantified statement).

$$\begin{aligned} \varphi : A \rightarrow B \text{ is structure-preserving} \\ \iff \left( \begin{aligned} &\text{every selected constant is preserved} \\ &\wedge \text{ every selected operation is preserved} \\ &\wedge \forall a, a' \in A (a <_A a' \Rightarrow \varphi(a) <_B \varphi(a')) \end{aligned} \right). \end{aligned}$$

*Remark* (Interpretation). A structure-preserving map carries the chosen arithmetic and order from  $A$  into  $B$  without distortion. The choice of structure matters: the positive part of  $\mathbb{W}$  carries the Peano successor from  $\mathbb{N}$ , while the later ring and field embeddings focus on 0, 1, addition, multiplication, and order. A structure-preserving map need not be injective; collapsing distinct elements is still allowed at this stage. Injectivity is added separately to obtain an embedding.

*Remark* (Dependencies).

- [ℕ Embeds into ℚ as the Positive Part](#)

#### Definition (Order Embedding)

**Definition** (Order Embedding). Let  $A$  and  $B$  be ordered arithmetic systems and  $\varphi : A \rightarrow B$  a map. Then  $\varphi$  is an *order embedding* exactly when, for all  $a, a' \in A$ ,

$$a <_A a' \iff \varphi(a) <_B \varphi(a').$$

#### Definition (Embedding of Number Systems)

**Definition 5.1.2** (Embedding of Number Systems). Let  $A$  and  $B$  be ordered arithmetic systems. A map  $\varphi : A \rightarrow B$  is an *embedding* exactly when  $\varphi$  is injective and structure-preserving and is an order embedding.

*Remark* (Standard quantified statement).

$$\begin{aligned} \varphi \text{ is an embedding} \\ \iff \left( \begin{aligned} &\varphi \text{ is injective} \wedge \varphi \text{ is structure-preserving} \\ &\wedge \forall a, a' \in A (a <_A a' \Leftrightarrow \varphi(a) <_B \varphi(a')) \end{aligned} \right). \end{aligned}$$

*Remark* (Negated quantified statement).

$$\begin{aligned} \varphi \text{ is not an embedding} \\ \iff \left( \begin{aligned} &\exists a, a' \in A (a \neq a' \wedge \varphi(a) = \varphi(a')) \\ &\vee \varphi \text{ fails to preserve a constant, sum, or product} \\ &\vee \exists a, a' \in A ((a <_A a' \wedge \neg(\varphi(a) <_B \varphi(a'))) \vee (\varphi(a) <_B \varphi(a') \wedge \neg(a <_A a'))) \end{aligned} \right). \end{aligned}$$

*Remark* (Failure modes). An embedding can fail in three structurally distinct ways: it may collapse two distinct elements to the same image (injectivity fails), fail to transport one of the selected constants or operations (algebraic preservation fails), or fail to preserve or reflect a strict-order comparison (order embedding fails).

*Remark* (Interpretation). An embedding is the precise meaning of “ $A$  sits inside  $B$ .” It is injective, so distinct elements stay distinct; it preserves the arithmetic and reflects the order, so the copy of  $A$  inside  $B$  behaves exactly as  $A$  did. This is what licenses treating  $2 \in \mathbb{N}$  and  $2 \in \mathbb{R}$  as “the same number.”

*Remark* (Interpretation). Most embeddings after  $\mathbb{W}$  transport the *ordered ring/field* structure: constants, the two operations, and order. The first bridge  $\mathbb{N} \hookrightarrow \mathbb{W}$  is slightly different because  $\mathbb{N}$  is one-based and carries Peano successor. That local theorem is proved in the whole-number chapter. In every case, induction is not transported to the larger target: the phrase “the naturals sit inside the integers” must not be read as carrying induction across all of  $\mathbb{Z}$ .

*Remark* (Dependencies).

- [Structure-Preserving Map](#)
- [Order Embedding](#)

#### Definition (Image of an Embedding)

**Definition 5.1.3** (Image of an Embedding). Let  $\varphi : A \rightarrow B$  be an embedding. The *image* of  $\varphi$  is the set  $\varphi(A) = \{ \varphi(a) : a \in A \} \subseteq B$ , equipped with the selected constants, operations, and order of  $B$  restricted to  $\varphi(A)$ .

*Remark* (Standard quantified statement).

$$\varphi(A) = \{ b \in B : \exists a \in A (b = \varphi(a)) \}.$$

*Remark* (Interpretation). The image is the literal subset of  $B$  that plays the role of  $A$ . The identification theorem below states that  $\varphi$ , viewed as a map onto its image, is an isomorphism, so  $A$  and  $\varphi(A)$  are the same ordered arithmetic system under two names.

*Remark* (Dependencies).

- [Embedding of Number Systems](#)

#### Theorem (Embeddings Identify a System with Its Image)

**Theorem 5.1.4** (Embeddings Identify a System with Its Image). *Let  $\varphi : A \rightarrow B$  be an embedding of ordered arithmetic systems relative to a selected source structure. Then the corestriction  $\varphi : A \rightarrow \varphi(A)$  is an isomorphism onto its image: it is a bijection that preserves and reflects the selected constants, operations, and order. Consequently  $A$  may be identified with the corresponding substructure  $\varphi(A) \subseteq B$ . [Go to proof.](#)*

*Remark* (Standard quantified statement).

$\varphi : A \rightarrow B$  an embedding  
 $\implies \varphi : A \rightarrow \varphi(A)$  is a bijective isomorphism for the selected structure.

*Remark* (Interpretation). This is the theorem that makes the inclusion notation legitimate. After identification,  $A \subseteq B$  abbreviates “ $A$  embeds via  $\varphi$  as the substructure  $\varphi(A)$ .” Every later use of  $\mathbb{N} \subseteq \mathbb{R}$  is shorthand for this identification.

*Remark* (Dependencies).

- [Embedding of Number Systems](#)
- [Image of an Embedding](#)

## 5.2 Embedding $\mathbb{N}$ Into $\mathbb{W}$

### Embedding $\mathbb{N}$ into $\mathbb{W}$

This section records, from the global embedding viewpoint, the local bridge proved in the whole-number chapter:  $\mathbb{N}$  sits inside  $\mathbb{W}$  as the positive part. The map is the bridge that lets elements of  $\mathbb{N}$  be regarded as nonzero elements of  $\mathbb{W}$  after identification.

#### Definition (Embedding $\mathbb{N}$ into $\mathbb{W}$ )

**Definition 5.2.1** (Embedding  $\mathbb{N}$  into  $\mathbb{W}$ ). The *canonical embedding*  $\iota : \mathbb{N} \rightarrow \mathbb{W}$  is defined by

$\iota(n) = n$  (the copy of each natural number inside the whole numbers).

*Remark* (Standard quantified statement).

$\iota : \mathbb{N} \rightarrow \mathbb{W}, \quad \iota(n) = n$  (the copy of each natural number inside the whole numbers).

*Remark* (Interpretation). The map sends each element of  $\mathbb{N}$  to its natural representative in  $\mathbb{W}$ . The next result certifies that this representative choice preserves all arithmetic and order, so  $\mathbb{N}$  may be identified with its image in  $\mathbb{W}$ .

*Remark* (Dependencies).

- [Construction of  \$\mathbb{W}\$  from  \$\mathbb{N}\$  by adjoining 0](#)
- [N Embeds into  \$\mathbb{W}\$  as the Positive Part](#)

#### Theorem ( $\mathbb{N}$ Embeds in $\mathbb{W}$ )

**Theorem 5.2.2** ( $\mathbb{N}$  Embeds in  $\mathbb{W}$ ). The canonical map  $\iota : \mathbb{N} \rightarrow \mathbb{W}$  identifies  $\mathbb{N}$  with the positive part  $\mathbb{W} \setminus \{0\}$  and preserves successor, 1, addition, multiplication, and order. [Go to proof.](#)

*Remark* (Standard quantified statement).

$\iota : \mathbb{N} \rightarrow \mathbb{W} \setminus \{0\}$  is bijective and preserves successor, arithmetic, and order.

*Remark* (Interpretation). Once this theorem is in place,  $\mathbb{N}$  may be treated as the positive substructure  $\iota(\mathbb{N}) = \mathbb{W} \setminus \{0\}$  inside  $\mathbb{W}$ . The new element of  $\mathbb{W}$  is exactly the adjoined zero.

*Remark* (Dependencies).

- [Construction of  \$\mathbb{W}\$  from  \$\mathbb{N}\$  by adjoining 0](#)
- [Embedding  \$\mathbb{N}\$  into  \$\mathbb{W}\$](#)
- [N Embeds into  \$\mathbb{W}\$  as the Positive Part](#)

## 5.3 Embedding $\mathbb{W}$ Into $\mathbb{Z}$

### Embedding $\mathbb{W}$ into $\mathbb{Z}$

This section fixes the canonical embedding of  $\mathbb{W}$  into  $\mathbb{Z}$  and records that it is injective, structure-preserving, and order-reflecting. The map is the bridge that lets elements of  $\mathbb{W}$  be regarded as elements of  $\mathbb{Z}$  after identification.

#### Definition (Embedding $\mathbb{W}$ into $\mathbb{Z}$ )

**Definition 5.3.1** (Embedding  $\mathbb{W}$  into  $\mathbb{Z}$ ). The *canonical embedding*  $\iota : \mathbb{W} \rightarrow \mathbb{Z}$  is defined by

$$\iota(w) = [(w, 0)] \quad (\text{the integer represented by } w \text{ and } 0).$$

*Remark* (Standard quantified statement).

$$\iota : \mathbb{W} \rightarrow \mathbb{Z}, \quad \iota(w) = [(w, 0)] \quad (\text{the integer represented by } w \text{ and } 0).$$

*Remark* (Interpretation). The map sends each element of  $\mathbb{W}$  to its natural representative in  $\mathbb{Z}$ . The next result certifies that this representative choice preserves all arithmetic and order, so  $\mathbb{W}$  may be identified with its image in  $\mathbb{Z}$ .

*Remark* (Dependencies).

- [Formal-Difference Equivalence Is an Equivalence Relation](#)
- [Embedding of Number Systems](#)

#### Theorem ( $\mathbb{W}$ Embeds in $\mathbb{Z}$ )

**Theorem 5.3.2** ( $\mathbb{W}$  Embeds in  $\mathbb{Z}$ ). *The canonical map  $\iota : \mathbb{W} \rightarrow \mathbb{Z}$  is an embedding of ordered arithmetic systems: it is injective, preserves 0, 1, addition, and multiplication, and reflects the order. [Go to proof.](#)*

*Remark* (Standard quantified statement).

$\iota : \mathbb{W} \rightarrow \mathbb{Z}$  is injective, structure-preserving, and order-reflecting.

*Remark* (Interpretation). Once this theorem is proved, the identification theorem of §13.1 applies, and  $\mathbb{W}$  may be treated as the substructure  $\iota(\mathbb{W}) \subseteq \mathbb{Z}$ .

*Remark* (Dependencies).

- [Formal-Difference Equivalence Is an Equivalence Relation](#)
- [Embedding  \$\mathbb{W}\$  into  \$\mathbb{Z}\$](#)
- [Embeddings Identify a System with Its Image](#)

## 5.4 Embedding $\mathbb{Z}$ Into $\mathbb{Q}$

### Embedding $\mathbb{Z}$ into $\mathbb{Q}$

This section fixes the canonical embedding of  $\mathbb{Z}$  into  $\mathbb{Q}$  and records that it is injective, structure-preserving, and order-reflecting. The map is the bridge that lets elements of  $\mathbb{Z}$  be regarded as elements of  $\mathbb{Q}$  after identification.

#### Definition (Embedding $\mathbb{Z}$ into $\mathbb{Q}$ )

**Definition 5.4.1** (Embedding  $\mathbb{Z}$  into  $\mathbb{Q}$ ). The *canonical embedding*  $\iota : \mathbb{Z} \rightarrow \mathbb{Q}$  is defined by

$$\iota(z) = [(z, 1)] \quad (\text{the rational represented by } z/1).$$

*Remark* (Standard quantified statement).

$$\iota : \mathbb{Z} \rightarrow \mathbb{Q}, \quad \iota(z) = [(z, 1)] \quad (\text{the rational represented by } z/1).$$

*Remark* (Interpretation). The map sends each element of  $\mathbb{Z}$  to its natural representative in  $\mathbb{Q}$ . The next result certifies that this representative choice preserves all arithmetic and order, so  $\mathbb{Z}$  may be identified with its image in  $\mathbb{Q}$ .

*Remark* (Dependencies).

- [Construction of  \$\mathbb{Q}\$  as fraction classes of integers](#)
- [Embedding of Number Systems](#)

#### Theorem ( $\mathbb{Z}$ Embeds in $\mathbb{Q}$ )

**Theorem 5.4.2** ( $\mathbb{Z}$  Embeds in  $\mathbb{Q}$ ). *The canonical map  $\iota : \mathbb{Z} \rightarrow \mathbb{Q}$  is an embedding of ordered arithmetic systems: it is injective, preserves 0, 1, addition, and multiplication, and reflects the order. [Go to proof.](#)*



*Remark* (Standard quantified statement).

$\iota : \mathbb{Z} \rightarrow \mathbb{Q}$  is injective, structure-preserving, and order-reflecting.

*Remark* (Interpretation). Once this theorem is proved, the identification theorem of §13.1 applies, and  $\mathbb{Z}$  may be treated as the substructure  $\iota(\mathbb{Z}) \subseteq \mathbb{Q}$ .

*Remark* (Dependencies).

- [Construction of  \$\mathbb{Q}\$  as fraction classes of integers](#)
- [Embedding  \$\mathbb{Z}\$  into  \$\mathbb{Q}\$](#)
- [Embeddings Identify a System with Its Image](#)

## 5.5 Embedding $\mathbb{Q}$ Into $\mathbb{R}$

### Embedding $\mathbb{Q}$ into $\mathbb{R}$

This section fixes the canonical embedding of  $\mathbb{Q}$  into  $\mathbb{R}$  and records that it is injective, structure-preserving, and order-reflecting. The map is the bridge that lets elements of  $\mathbb{Q}$  be regarded as elements of  $\mathbb{R}$  after identification.

#### Definition (Embedding $\mathbb{Q}$ into $\mathbb{R}$ )

**Definition 5.5.1** (Embedding  $\mathbb{Q}$  into  $\mathbb{R}$ ). The *canonical embedding*  $\iota : \mathbb{Q} \rightarrow \mathbb{R}$  is defined by

$$\iota(q) = q^* \quad (\text{the real number determined by } q).$$

*Remark* (Standard quantified statement).

$$\iota : \mathbb{Q} \rightarrow \mathbb{R}, \quad \iota(q) = q^* \quad (\text{the real number determined by } q).$$

*Remark* (Interpretation). The map sends each element of  $\mathbb{Q}$  to its natural representative in  $\mathbb{R}$ . The next result certifies that this representative choice preserves all arithmetic and order, so  $\mathbb{Q}$  may be identified with its image in  $\mathbb{R}$ .

*Remark* (Dependencies).

- [Construction of  \$\mathbb{R}\$  by Dedekind cuts](#)
- [Embedding of Number Systems](#)

#### Theorem ( $\mathbb{Q}$ Embeds in $\mathbb{R}$ )

**Theorem 5.5.2** ( $\mathbb{Q}$  Embeds in  $\mathbb{R}$ ). *The canonical map  $\iota : \mathbb{Q} \rightarrow \mathbb{R}$  is an embedding of ordered arithmetic systems: it is injective, preserves 0, 1, addition, and multiplication, and reflects the order. [Go to proof.](#)*

*Remark* (Standard quantified statement).

$\iota : \mathbb{Q} \rightarrow \mathbb{R}$  is injective, structure-preserving, and order-reflecting.

*Remark* (Interpretation). Once this theorem is proved, the identification theorem of §13.1 applies, and  $\mathbb{Q}$  may be treated as the substructure  $\iota(\mathbb{Q}) \subseteq \mathbb{R}$ .

*Remark* (Dependencies).

- [Construction of  \$\mathbb{R}\$  by Dedekind cuts](#)
- [Embedding  \$\mathbb{Q}\$  into  \$\mathbb{R}\$](#)
- [Embeddings Identify a System with Its Image](#)

## 5.6 Compatibility of Arithmetic and Order

### One Arithmetic Through the Chain

Each embedding preserves arithmetic and order individually. What remains is to confirm that the embeddings are mutually compatible: composing  $\mathbb{N} \hookrightarrow \mathbb{W} \hookrightarrow \mathbb{Z} \hookrightarrow \mathbb{Q} \hookrightarrow \mathbb{R}$  agrees with the direct embedding of  $\mathbb{N}$  into  $\mathbb{R}$ , so that “2” has one unambiguous meaning after all identifications.

#### Theorem (Compatibility of the Embedding Chain)

**Theorem 5.6.1** (Compatibility of the Embedding Chain). *Let  $\iota_1 : \mathbb{N} \rightarrow \mathbb{W}$ ,  $\iota_2 : \mathbb{W} \rightarrow \mathbb{Z}$ ,  $\iota_3 : \mathbb{Z} \rightarrow \mathbb{Q}$ , and  $\iota_4 : \mathbb{Q} \rightarrow \mathbb{R}$  be the canonical embeddings. For every  $n \in \mathbb{N}$  the composite  $(\iota_4 \circ \iota_3 \circ \iota_2 \circ \iota_1)(n)$  is the real number identified with  $n$ , and the composite preserves the arithmetic and order carried by the source system. Hence arithmetic computed in any system agrees with arithmetic on its image in  $\mathbb{R}$ . [Go to proof.](#)*

*Remark* (Standard quantified statement).

$\forall n, m \in \mathbb{N} \left( \Phi(n+m) = \Phi(n) + \Phi(m) \wedge \Phi(n \cdot m) = \Phi(n) \cdot \Phi(m) \wedge (n < m \Leftrightarrow \Phi(n) < \Phi(m)) \right), \quad \Phi = \iota_4 \circ \iota_3 \circ \iota_2 \circ \iota_1$

*Remark* (Interpretation). Compatibility is what makes the single symbol 2 well-defined across the whole chain: the answer to  $2 + 2$  is the same whether computed in  $\mathbb{N}$  or after embedding into  $\mathbb{R}$ . Without this coherence the inclusion notation would silently carry two different meanings for the same numeral.

*Remark* (Dependencies).

- [N Embeds into W as the Positive Part](#)
- [W Embeds in Z](#)
- [Z Embeds in Q](#)
- [Q Embeds in R](#)

## 5.7 One Chain Many Structures

### One Chain, Many Structures

The embeddings assemble into a single chain  $\mathbb{N} \hookrightarrow \mathbb{W} \hookrightarrow \mathbb{Z} \hookrightarrow \mathbb{Q} \hookrightarrow \mathbb{R}$ . Each arrow is injective, preserves the relevant arithmetic, and reflects order; each step also *adds* structure that the previous system lacked.

From  $\mathbb{N}$  to  $\mathbb{W}$  an additive identity  $0$  appears. From  $\mathbb{W}$  to  $\mathbb{Z}$  additive inverses appear, completing a commutative ring. From  $\mathbb{Z}$  to  $\mathbb{Q}$  multiplicative inverses of nonzero elements appear, completing an ordered field. From  $\mathbb{Q}$  to  $\mathbb{R}$  order-completeness appears: the gaps in the rational line are filled. The numerals carry through unchanged by compatibility, but the ambient structure grows at every step. This chain is the bridge from number construction to the analysis of the real line.

*Remark* (Exposition). Read as a ladder of structures: counting ( $\mathbb{N}$ )  $\rightarrow$  counting with zero ( $\mathbb{W}$ )  $\rightarrow$  signed counting / ring ( $\mathbb{Z}$ )  $\rightarrow$  ordered field ( $\mathbb{Q}$ )  $\rightarrow$  complete ordered field ( $\mathbb{R}$ ). Only the last rung is order-complete, and that completeness is the property the structure-of-the-real-line chapter turns into metric topology.

## Proofs

*Remark* (Return). [Return to Theorem](#)

**Theorem** (Embeddings Identify a System with Its Image). *Let  $\varphi : A \rightarrow B$  be an embedding of ordered arithmetic systems. Then the corestriction  $\varphi : A \rightarrow \varphi(A)$  is an isomorphism of ordered arithmetic systems: it is a bijection that preserves and reflects the constants, both operations, and the order. Consequently  $A$  may be identified with the substructure  $\varphi(A) \subseteq B$ .*

*Professional Standard Proof.*

TODO: professional standard proof for embedding-identifies-with-image. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-identifies-with-image. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Embedding of Number Systems](#)
- [Image of an Embedding](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (*N Embeds in W*). *The canonical map  $\iota : \mathbb{N} \rightarrow \mathbb{W}$  identifies  $\mathbb{N}$  with the positive part  $\mathbb{W} \setminus \{0\}$  and preserves successor, 1, addition, multiplication, and order.*

*Professional Standard Proof.*

TODO: professional standard proof for n-embeds-in-w. □

*Detailed Learning Proof.*

TODO: detailed learning proof for n-embeds-in-w. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [Construction of  \$\mathbb{W}\$  from  \$\mathbb{N}\$  by adjoining 0](#)
- [Embedding  \$\mathbb{N}\$  into  \$\mathbb{W}\$](#)
- [N Embeds into  \$\mathbb{W}\$  as the Positive Part](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (*W Embeds in Z*). *The canonical map  $\iota : \mathbb{W} \rightarrow \mathbb{Z}$  is an embedding of ordered arithmetic systems: it is injective, preserves 0, 1, addition, and multiplication, and reflects the order.*

*Professional Standard Proof.*

TODO: professional standard proof for w-embeds-in-z. □

*Detailed Learning Proof.*

TODO: detailed learning proof for w-embeds-in-z. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: link to the construction of  $\mathbb{Z}$  as difference classes of whole numbers.
- [Embedding  \$\mathbb{W}\$  into  \$\mathbb{Z}\$](#)
- [Embeddings Identify a System with Its Image](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (*Z Embeds in Q*). *The canonical map  $\iota : \mathbb{Z} \rightarrow \mathbb{Q}$  is an embedding of ordered arithmetic systems: it is injective, preserves 0, 1, addition, and multiplication, and reflects the order.*

*Professional Standard Proof.*

TODO: professional standard proof for z-embeds-in-q. □

*Detailed Learning Proof.*

TODO: detailed learning proof for z-embeds-in-q. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: link to the construction of  $\mathbb{Q}$  as fraction classes of integers.
- [Embedding  \$\mathbb{Z}\$  into  \$\mathbb{Q}\$](#)
- [Embeddings Identify a System with Its Image](#)

*Remark* (Return). [Return to Theorem](#)

**Theorem** (*Q Embeds in R*). *The canonical map  $\iota : \mathbb{Q} \rightarrow \mathbb{R}$  is an embedding of ordered arithmetic systems: it is injective, preserves 0, 1, addition, and multiplication, and reflects the order.*

*Professional Standard Proof.*

TODO: professional standard proof for q-embeds-in-r. □

*Detailed Learning Proof.*

TODO: detailed learning proof for q-embeds-in-r. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- TODO: link to the construction of  $\mathbb{R}$  (Dedekind cuts or Cauchy sequences).
- [Embedding  \$\mathbb{Q}\$  into  \$\mathbb{R}\$](#)
- [Embeddings Identify a System with Its Image](#)



*Remark* (Return). [Return to Theorem](#)

**Theorem** (Compatibility of the Embedding Chain). *Let  $\iota_1 : \mathbb{N} \rightarrow \mathbb{W}$ ,  $\iota_2 : \mathbb{W} \rightarrow \mathbb{Z}$ ,  $\iota_3 : \mathbb{Z} \rightarrow \mathbb{Q}$ , and  $\iota_4 : \mathbb{Q} \rightarrow \mathbb{R}$  be the canonical embeddings. For every  $n \in \mathbb{N}$  the composite  $(\iota_4 \circ \iota_3 \circ \iota_2 \circ \iota_1)(n)$  is the real number identified with  $n$ , and the composite preserves the arithmetic and order carried by the source system. Hence arithmetic computed in any system agrees with arithmetic on its image in  $\mathbb{R}$ .*

*Professional Standard Proof.*

TODO: professional standard proof for embedding-chain-compatibility. □

*Detailed Learning Proof.*

TODO: detailed learning proof for embedding-chain-compatibility. □

*Remark* (Proof structure).

*Remark* (Dependencies).

- [ℕ Embeds into ℚ as the Positive Part](#)
- [ℚ Embeds in ℤ](#)
- [ℤ Embeds in ℚ](#)
- [ℚ Embeds in ℝ](#)

# Capstone

# Chapter 6

## Summary: The Number Systems



### 6.1 Axiom Systems for $\mathbb{N}$ , $\mathbb{W}$ , $\mathbb{Z}$ , $\mathbb{Q}$ , and $\mathbb{R}$

#### Derived Properties — Quick Reference

| Name                | Statement                             | Proof sketch                                                                       |
|---------------------|---------------------------------------|------------------------------------------------------------------------------------|
| Zero multiplication | $a \cdot 0 = 0$                       | $a \cdot 0 = a \cdot (0 + 0) = a \cdot 0 + a \cdot 0$ ; cancel $a \cdot 0$ via A4. |
| Negation rule       | $(-a) \cdot b = -(ab)$                | $ab + (-a)b = (a + (-a))b = 0 \cdot b = 0$ .                                       |
| $(-1)$ rule         | $(-1) \cdot a = -a$                   | Special case: $b = a$ , $-a = (-1)a$ .                                             |
| Double negation     | $(-a)(-b) = ab$                       | $(-a)(-b) = -(a(-b)) = -(-(ab)) = ab$ .                                            |
| No zero divisors    | $ab = 0 \Rightarrow a = 0$ or $b = 0$ | $\mathbb{Z}$ is an integral domain; proved via well-ordering + sign analysis.      |

## 6.2 Number Systems Summary

### The Natural Numbers $\mathbb{N}$

#### One-Based Peano Axioms for $\mathbb{N}$

Let  $\mathbb{N}$  be a set with distinguished element 1 and successor operation  $n \mapsto n++$ .

**P1**  $1 \in \mathbb{N}$ .

**P2**  $n \in \mathbb{N} \Rightarrow n++ \in \mathbb{N}$ . *(successor closure)*

**P3**  $n++ \neq 1$  for all  $n \in \mathbb{N}$ . *(the first element has no predecessor)*

**P4**  $n++ = m++ \Rightarrow n = m$ . *(successor is injective)*

**P5** If  $P(0)$  holds and  $P(n) \Rightarrow P(n++)$ , then  $P(n)$  holds for all  $n \in \mathbb{N}$ .  
*(induction)*

*Remark.* P1–P4 rule out wrap-around, ceiling, and rogue elements. P5 eliminates anything unreachable from 1 by finite iteration of the successor. Addition, multiplication, and order are not axioms – they are *defined* recursively from the successor and then shown to satisfy familiar laws by induction.

*Remark* (Well-ordering and induction are equivalent over  $\mathbb{N}$ ). The following are logically equivalent:

- Principle of Mathematical Induction (P5).
- Well-Ordering Principle: every nonempty  $A \subseteq \mathbb{N}$  has a least element.
- Strong Induction:  $(\forall k < n, P(k)) \Rightarrow P(n)$  for all  $n$  implies  $P$  holds everywhere.

One must assume at least one; the others follow. Well-ordering is the engine behind the Division Algorithm, Bézout’s Identity, and unique prime factorization.

### The Whole Numbers $\mathbb{W}$

#### $\mathbb{W}$ as $\mathbb{N}$ with an Adjoined Zero

The whole numbers are

$$\mathbb{W} = \mathbb{N} \cup \{0\}, \quad 0 \notin \mathbb{N}.$$

The successor, addition, multiplication, and order on  $\mathbb{W}$  extend the corresponding structures on  $\mathbb{N}$ :

- $S_{\mathbb{W}}(0) = 1$ , and  $S_{\mathbb{W}}$  agrees with  $S_{\mathbb{N}}$  on  $\mathbb{N}$ ;
- 0 is the additive identity;
- 0 is absorbing for multiplication;
- 0 is the least element of  $\mathbb{W}$ .

*Remark* (What  $\mathbb{W}$  gains over  $\mathbb{N}$ , and what it lacks). **Gain:** an additive identity. This makes  $(\mathbb{W}, +, 0)$  a commutative monoid and prepares the ordered-pair construction of the integers.

**Lack:** additive inverses for nonzero elements. Subtraction is still not an internal operation in general, so  $\mathbb{W}$  is not a ring.

## The Integers $\mathbb{Z}$

### Axioms for $\mathbb{Z}$ — Ordered Commutative Ring

#### Addition

$$\mathbf{A1} \quad a + b = b + a \quad (\text{commutativity})$$

$$\mathbf{A2} \quad (a + b) + c = a + (b + c) \quad (\text{associativity})$$

$$\mathbf{A3} \quad \exists 0 \in \mathbb{Z} \text{ such that } a + 0 = a \quad (\text{additive identity})$$

$$\mathbf{A4} \quad \forall a, \exists -a \in \mathbb{Z} \text{ with } a + (-a) = 0 \quad (\text{additive inverses})$$

#### Multiplication

$$\mathbf{M1} \quad a \cdot b = b \cdot a \quad (\text{commutativity})$$

$$\mathbf{M2} \quad (a \cdot b) \cdot c = a \cdot (b \cdot c) \quad (\text{associativity})$$

$$\mathbf{M3} \quad \exists 1 \in \mathbb{Z}, 1 \neq 0, \text{ such that } a \cdot 1 = a \quad (\text{multiplicative identity})$$

#### Distributivity

$$\mathbf{D} \quad a \cdot (b + c) = a \cdot b + a \cdot c \quad (\text{distributivity})$$

#### Order

$$\mathbf{O1} \quad a \leq b \text{ or } b \leq a \quad (\text{totality})$$

$$\mathbf{O2} \quad a \leq b \wedge b \leq a \Rightarrow a = b \quad (\text{antisymmetry})$$

$$\mathbf{O3} \quad a \leq b \wedge b \leq c \Rightarrow a \leq c \quad (\text{transitivity})$$

$$\mathbf{O4} \quad b \leq c \Rightarrow a + b \leq a + c \quad (\text{order} + \text{addition})$$

$$\mathbf{O5} \quad a \geq 0 \wedge b \geq 0 \Rightarrow ab \geq 0 \quad (\text{order} + \text{multiplication})$$

*Remark* (What  $\mathbb{Z}$  gains over  $\mathbb{W}$ , and what it lacks). **Gain:** Axiom A4 – additive inverses. Subtraction is always defined in  $\mathbb{Z}$ ; it is not in  $\mathbb{W}$ .

**Lack:** Multiplicative inverses. There is no  $x \in \mathbb{Z}$  with  $2x = 1$ , so M4 (every nonzero element is invertible) fails. In the language of abstract algebra,  $\mathbb{Z}$  is a *commutative ordered ring* but not a field. The passage to  $\mathbb{Q}$  restores multiplicative inverses by formally adjoining fractions.

## Derived Properties of $\mathbb{Z}$ (from the Ring Axioms)

*Remark* (Zero multiplication — the step most often looked up). The identity  $a \cdot 0 = 0$  is not an axiom; it is derived. The proof uses only A3, A4, and D:

$$a \cdot 0 = a \cdot (0 + 0) \stackrel{\mathbf{D}}{=} a \cdot 0 + a \cdot 0.$$

Adding  $-(a \cdot 0)$  to both sides (A4) gives  $0 = a \cdot 0$ , i.e.  $a \cdot 0 = 0$ . No order axioms, no well-ordering, no properties of  $\mathbb{N}$  are needed — this holds in any ring.

## Structural Comparison: $\mathbb{N}$ , $\mathbb{W}$ , $\mathbb{Z}$ , $\mathbb{Q}$ , $\mathbb{R}$

| Axiom Comparison Table     |                          |                            |                       |                   |                      |
|----------------------------|--------------------------|----------------------------|-----------------------|-------------------|----------------------|
| Group                      | Axiom                    | $\mathbb{N}$               | $\mathbb{W}$          | $\mathbb{Z}$      | $\mathbb{Q}$         |
| Addition                   | A1 commutativity         | ✓                          | ✓                     | ✓                 | ✓                    |
|                            | A2 associativity         | ✓                          | ✓                     | ✓                 | ✓                    |
|                            | A3 identity (0)          | ×                          | ✓                     | ✓                 | ✓                    |
|                            | A4 inverses ( $-a$ )     | ×                          | ×                     | ✓                 | ✓                    |
| Multiplication             | M1 commutativity         | ✓                          | ✓                     | ✓                 | ✓                    |
|                            | M2 associativity         | ✓                          | ✓                     | ✓                 | ✓                    |
|                            | M3 identity (1)          | ✓                          | ✓                     | ✓                 | ✓                    |
|                            | M4 inverses ( $1/a$ )    | ×                          | ×                     | ×                 | ✓                    |
|                            | D distributivity         | ✓                          | ✓                     | ✓                 | ✓                    |
| Order                      | O1–O3 total order        | ✓                          | ✓                     | ✓                 | ✓                    |
|                            | O4–O5 order + arithmetic | ✓                          | ✓                     | ✓                 | ✓                    |
|                            | Well-ordering            | ✓                          | ✓                     | ×                 | ×                    |
| Completeness               | Least upper bound        | ×                          | ×                     | ×                 | ×                    |
| <b>Structure</b>           |                          | <i>positive arithmetic</i> | <i>comm. semiring</i> | <i>comm. ring</i> | <i>ordered field</i> |
| ✓ = satisfied    × = fails |                          |                            |                       |                   |                      |

*Remark* (The extension hierarchy). Each number system repairs exactly one deficiency of its predecessor:

$$\mathbb{N} \xrightarrow{+0} \mathbb{W} \xrightarrow{+A4} \mathbb{Z} \xrightarrow{+M4} \mathbb{Q} \xrightarrow{+LUB} \mathbb{R}.$$

$\mathbb{N} \rightarrow \mathbb{W}$ : adjoin 0 so addition has an identity.  $\mathbb{W} \rightarrow \mathbb{Z}$ : adjoin additive inverses so subtraction is always defined.  $\mathbb{Z} \rightarrow \mathbb{Q}$ : adjoin multiplicative inverses so division by any nonzero element is always defined.  $\mathbb{Q} \rightarrow \mathbb{R}$ : adjoin least upper bounds so every bounded increasing sequence converges.

Note that  $\mathbb{N}$  and  $\mathbb{W}$  are the well-ordered systems in the chain.  $\mathbb{Z}$  loses well-ordering the moment negative integers are introduced (the set of all negative integers has no least element).

## From Extension to Embedding

The hierarchy above describes what each new system adds. The next chapter looks at a separate question: how the earlier systems are identified inside the later ones. That identification is not merely a matter of writing subset symbols. It is carried by structure-preserving embeddings that justify treating a number from an earlier construction as the corresponding number in a larger system.

## Proofs

## Capstone

# Bibliography