


FROM CANTOR TO ITO

Classical Analysis

Bounds, Sequences, and Series



Bernard Bolzano

1781-1848

Self-Study Notes

William Sollers

July 1, 2026

For every reader learning to make mathematics their own.

Contents

1	Real Analysis	1
1.1	Foundational Analysis Properties	2
1.1.1	Foundational Analysis Properties	2
1.1.2	Manipulating Inequalities and the Logic of Bounding	9
1.1.3	Completeness as a Constructive Engine	14
1.1.4	Walking a Sequence with a Predicate	17
1.1.5	Interval Bisection and Nested Intervals	22
1.1.6	Subsequence Partition via Residue Classes	26
1.2	Asymptotic Notation	30
1.3	Inequality	33
1.4	Modulus	39
2	Structure of the Real Line	85
2.1	Real Line One Dimensional	85
2.2	Order Distance Length	85
2.3	Absolute Value as Distance	87
2.4	Intervals as Subsets	87
2.5	Neighborhoods and Balls	90
2.6	Open Sets	91
2.7	Closed Sets	92
2.8	Interior Exterior Boundary	93
2.9	Limit and Isolated Points	95
2.10	Unions and Intersections	96
2.11	Covers and Subcovers	96
2.12	Compactness on $\mathbb{R}$	97
2.13	Heine Borel	99
3	Bounds	110
3.1	Completeness Axiom	110
3.1.1	Nested Intervals	112
3.1.2	Archimedean Property	114
3.1.3	Density	115
3.1.4	Completeness Summary	122
3.2	Bounds and Extremal Values	123
3.2.1	Upper and Lower Bounds	123

3.2.2	Suprema and Infima	126
3.2.3	Maxima and Minima	135
3.2.4	Epsilon Characterization	139
3.2.5	Extremal Bounds Summary	142
3.3	Algebra of Bounds	142
3.3.1	Set Operations and Bounds	142
3.3.2	Lattice Operations and Bounds	148
3.3.3	Images, Inverse Images, and Extrema	152
3.3.4	Dilation and Displacement	160
3.3.5	Algebra of Suprema and Infima	166
4	Functions	277
4.1	Algebra of Functions	277
4.2	Real-Valued Functions	281
4.3	Subsets of $\mathbb{R}$	293
5	Discrete Calculus	312
5.1	Motivation	312
5.2	Foundations	315
5.2.1	Foundations	315
5.2.2	Summation Notation	315
5.2.3	The Binomial Theorem	317
5.3	Algebra of Summations	319
5.4	Difference Operators	322
5.4.1	Difference Operators	322
5.4.2	Higher Differences	323
5.4.3	Properties of the Difference Operator	324
5.4.4	Difference Tables and Polynomial Detection	325
5.5	Telescoping Sums	327
5.5.1	Telescoping Sums	327
5.6	Discrete Antiderivatives	329
5.6.1	Discrete Antiderivatives	329
5.7	Polynomial Basis	332
5.7.1	Polynomial Basis: Falling Factorials	332
5.8	Calculus Rules	335
5.8.1	Calculus Rules	335
5.9	Expansions	337
5.9.1	Newton's Forward Difference Expansion	337
5.10	Special Functions	339
5.10.1	Special Functions in Discrete Calculus	339
5.10.2	The Function 2^n in Discrete Calculus	340

6	Sequences	355
6.1	Sequence Definitions	355
6.1.1	Definitions	355
6.1.2	Null and Constant Sequences	357
6.2	Examples and Counterexamples	369
6.3	Convergence	371
6.3.1	Limits	374
6.3.2	Algebra of Limits	379
6.4	Tails	391
6.5	Monotonicity	395
6.6	Subsequences	406
6.7	Divergence	416
6.8	Liminf and Limsup	422
6.9	Order-Theoretic Limit Constructions	430
6.10	Cluster Values of Sequences	433
6.11	Cauchy	436
6.12	Applications of Sequences	445
7	Series	556
7.1	Finite Series	556
7.2	Infinite Series	565
8	Function Sequences	585

Chapter 1

Real Analysis



1.1 Foundational Analysis Properties

1.1.1 Foundational Analysis Properties

Foundational Analysis Properties — Quick Reference

Concept	Goal	Proof mechanism / logical form
Well-definedness	Consistency	$x \sim y \Rightarrow F(x) = F(y)$
Existence	Feasibility	$\exists x P(x)$
Uniqueness	Singularity	$P(x_1) \wedge P(x_2) \Rightarrow x_1 = x_2$
Linearity	Decomposition	$L(f + g) = L(f) + L(g)$ and $L(cf) = cL(f)$
Uniformity	One choice works globally	$\forall \varepsilon > 0, \exists \delta > 0, \forall x, y, Q(\varepsilon, \delta, x, y)$
Equivalence	Interchangeable descriptions	$A \Rightarrow B$ and $B \Rightarrow A$
Characterisation	Usable test	$P(x) \iff C(x)$
Estimate	Control	$A \leq B, A < \varepsilon$, or $d(A, B) < \varepsilon$
Boundedness	Two-sided control	$\exists l, u \in S \forall x \in A (l \leq x \leq u)$
Upper/lower control	One-sided domination	$\forall x \in A (x \leq u)$ or $\forall x \in A (l \leq x)$
Extremality	Realized optimality	$m \in A \wedge \text{Bound}(m, A)$
Least/greatest character	Optimal bound	$\text{Bound}(b, A) \wedge \forall c (\text{Bound}(c, A) \Rightarrow b \preceq c)$
Monotonicity	Order preservation	Start with $x \leq y$ and prove $f(x) \leq f(y)$, or the reverse
Closure	Staying inside a class	$P(x) \wedge P(y) \Rightarrow P(x \star y)$
Approximation	Arbitrary precision	$\forall \varepsilon > 0 \exists x d(x, a) < \varepsilon$

Many proofs in analysis are not isolated inventions. They prove recurring structural properties: a definition is independent of choices, an object exists, there is only one possible object, an operation respects addition and scalars, a single parameter works everywhere, or two definitions describe the same phenomenon. Naming the property first tells us what the proof must do.

Well-Definedness

Well-definedness appears whenever an object is defined using choices: representatives of equivalence classes, selected sequences, selected partitions, or selected approximations. The goal is to prove that different allowed choices give the same output.

$$x \sim x' \Rightarrow F(x) = F(x').$$

Remark 1.1.1 (Proof sketch). 1. Identify the choice built into the definition.

2. Take two allowed choices, usually x and x' with $x \sim x'$.

3. Compute or compare the two proposed outputs.

4. Show the outputs agree in the codomain.

Remark 1.1.2 (Crucial logic). The output must be independent of the chosen representative. If changing the representative changes the output, the proposed definition is not a function on the quotient or chosen domain.

Existence

Existence proves that at least one object satisfying a property is available:

$$\exists x P(x).$$

In analysis the object might be a limit, a supremum, a point in an interval, a subsequence, a bound, an index N , a radius δ , or a partition.

Remark 1.1.3 (Proof sketch). 1. Determine every requirement in the property P .

2. Produce a candidate explicitly, or invoke a theorem that guarantees one.

3. Verify the candidate belongs to the required domain.

4. Verify the candidate satisfies every part of P .

Remark 1.1.4 (Constructive and nonconstructive existence). A constructive proof names the object directly. A nonconstructive proof uses a theorem such as completeness, the intermediate value theorem, or Bolzano–Weierstrass. In both cases, the final burden is the same: the object must satisfy the stated property.

Uniqueness

Uniqueness proves that at most one object can satisfy a property:

$$P(x_1) \wedge P(x_2) \Rightarrow x_1 = x_2.$$

It does not by itself prove existence. It only proves that two successful candidates cannot be distinct.

Remark 1.1.5 (Proof sketch). 1. Assume x_1 and x_2 both satisfy P .

2. Use the defining property of x_1 and the defining property of x_2 .

3. Derive two-sided comparison, zero distance, or identical defining data.

4. Conclude $x_1 = x_2$.

Remark 1.1.6 (Common analysis forms). For suprema, show $s \leq t$ and $t \leq s$. For limits, show $|L - M| < \varepsilon$ for every $\varepsilon > 0$, hence $L = M$. In metric spaces, show $d(x_1, x_2) = 0$.

Linearity

Linearity proves that an operation decomposes over addition and scalar multiplication:

$$L(f + g) = L(f) + L(g), \quad L(cf) = cL(f).$$

The goal is not merely algebraic neatness. Linearity says that a complicated input can be analysed by splitting it into simpler pieces.

Remark 1.1.7 (Proof sketch). 1. Take arbitrary admissible inputs f, g and scalar c .

2. Expand $L(f + g)$ using the definition of L and the definition of addition.

3. Rearrange the expression until $L(f) + L(g)$ appears.

4. Repeat with $L(cf)$ and show it equals $cL(f)$.

Remark 1.1.8 (Typical analysis examples). Limits, derivatives, integrals, and many summation operators have linearity theorems. Each proof follows the same shape: expand the left side, split the expression, and identify the two pieces.

Uniformity

Uniformity means that one choice works for all relevant points at once. The prototype is not just uniform continuity, but any statement where a witness is chosen before the point is allowed to vary:

$$\forall \varepsilon > 0, \exists \delta > 0, \forall x, y, Q(\varepsilon, \delta, x, y).$$

The key phrase is “one choice.” The same N , δ , bound, or partition must work globally over the specified domain.

Remark 1.1.9 (Proof sketch). 1. Fix the tolerance or target requirement.

2. Choose a single witness that depends only on the allowed global data.

3. Let the point, pair of points, function, or index vary arbitrarily.

4. Verify that the same witness works in every permitted case.

Remark 1.1.10 (Contrast with pointwise statements). Pointwise statements allow the witness to depend on the point. Uniform statements do not. The order of quantifiers is the whole issue:

$$\forall x \exists \delta \text{ is pointwise, while } \exists \delta \forall x \text{ is uniform.}$$

Equivalence

Equivalence proves that two statements are logically interchangeable:

$$A \iff B.$$

Such theorems are common because analysis often has several descriptions of the same phenomenon: epsilon-delta, sequential, order-theoretic, topological, or integral.

Remark 1.1.11 (Proof sketch). 1. Prove $A \Rightarrow B$.

2. Prove $B \Rightarrow A$.

3. Keep the assumptions for the two directions separate.

4. After both directions are proved, either description may be used.

Remark 1.1.12 (Typical analysis examples). Sequential and epsilon-delta definitions of continuity are equivalent. Riemann and Darboux integrability are equivalent under the appropriate hypotheses. A set is closed exactly when it contains the limits of all convergent sequences from the set.

Characterisation

A characterisation is an equivalence whose purpose is to turn a definition into a usable test:

$$P(x) \iff C(x).$$

The theorem does not merely say two statements agree. It supplies a criterion that can be used in later proofs.

Remark 1.1.13 (Proof sketch). 1. Identify the original definition.

2. Identify the proposed criterion.

3. Prove definition $\Rightarrow$ criterion.

4. Prove criterion $\Rightarrow$ definition.

Remark 1.1.14 (Typical analysis examples). The epsilon characterisation of the supremum turns least-upper-bound language into an approximation rule. Cauchy criteria turn convergence questions into internal distance estimates. Sequential criteria turn topological conditions into statements about sequences.

Estimates and Bounds

An estimate proves control:

$$A \leq B, \quad |A| < \varepsilon, \quad d(A, B) < \varepsilon.$$

The aim is not to compute the exact value. The aim is to put the target quantity inside a range that is strong enough for the theorem.

Remark 1.1.15 (Proof sketch). 1. Identify the quantity that must be controlled.

2. Replace it by a larger or simpler quantity using valid inequalities.

3. Use hypotheses to replace variable terms by constants or small errors.

4. End with the required bound.

Remark 1.1.16 (Typical analysis examples). Showing a sequence is bounded, proving an error term tends to zero, bounding a Riemann-sum error, or proving $|f(x) - L| < \varepsilon$ are all estimate problems. The proof is a chain of control.

Boundedness

Boundedness proves two-sided control. In an ordered setting, the usual form is that a set or sequence is bounded above and bounded below:

$$\exists l, u \in S \forall x \in A (l \leq x \leq u).$$

Remark 1.1.17 (Proof sketch). 1. Prove bounded above by finding a candidate upper bound.

2. Prove bounded below by finding a candidate lower bound.

3. Verify both candidates work for every element under consideration.

4. Conclude that the object is bounded.

Upper and Lower Control

Upper and lower control prove one-sided domination. The proof focuses on a candidate bound and checks that every element lies on the required side of it:

$$\forall x \in A (x \leq u), \quad \forall x \in A (l \leq x).$$

Remark 1.1.18 (Proof sketch). 1. Name the candidate upper or lower bound.

2. Let an arbitrary element of the set be given.

3. Use the hypotheses or definition of the set to compare that element with the candidate.

4. Since the element was arbitrary, conclude the one-sided bound.

Extremality

Extremality proves that an optimal value is actually realized by the object being studied. A maximum is not merely an upper bound; it is an upper bound that belongs to the set. A minimum is the corresponding realized lower bound.

$$\text{Maximum}(m, A) \iff (m \in A) \wedge \text{UpperBound}(m, A),$$

$$\text{Minimum}(m, A) \iff (m \in A) \wedge \text{LowerBound}(m, A).$$

Remark 1.1.19 (Proof sketch). 1. Prove the candidate belongs to the set.

2. Prove the candidate is the appropriate one-sided bound.

3. Combine membership with the bound condition.

4. Conclude maximum or minimum.

Least and Greatest Character

Least and greatest character proves optimality among competitors. For a supremum, the candidate must first be an upper bound, and then it must be no larger than every other upper bound. Infimum proofs reverse the direction.

$$\text{Supremum}(s, A) \iff \text{UpperBound}(s, A) \wedge \forall u \in S (\text{UpperBound}(u, A) \Rightarrow s \leq u),$$

$$\text{Infimum}(i, A) \iff \text{LowerBound}(i, A) \wedge \forall l \in S (\text{LowerBound}(l, A) \Rightarrow l \leq i).$$

Remark 1.1.20 (Proof sketch). 1. Prove the candidate is a bound of the required kind.

2. Let an arbitrary competing bound be given.
3. Compare the candidate with that competing bound.
4. Conclude the candidate is the least upper bound or greatest lower bound.

Monotonicity

Monotonicity proves order preservation or order reversal:

$$x \leq y \Rightarrow f(x) \leq f(y), \quad x \leq y \Rightarrow f(y) \leq f(x).$$

The proof begins with an arbitrary ordered pair and carries that order through the definitions.

Remark 1.1.21 (Proof sketch). 1. Take arbitrary points x and y with $x \leq y$.

2. Expand the definitions of the function or operation.
3. Use order rules and hypotheses to compare the outputs.
4. Conclude that the order is preserved or reversed.

Closure

Closure proves that a class is stable under an operation. One starts with objects already known to belong to the class, combines them, and then verifies that the result still satisfies the defining property.

$$P(x) \wedge P(y) \Rightarrow P(x \star y).$$

For closure under scalar operations, the same pattern becomes

$$P(x) \Rightarrow P(c \star x)$$

for every admissible scalar or parameter c .

Remark 1.1.22 (Proof sketch). 1. Take arbitrary objects from the class.

2. Form the object produced by the operation.
3. Check every condition in the definition of the class.
4. Conclude the result remains inside the class.

Approximation

Approximation proves arbitrary precision. The proof usually begins with a tolerance and constructs an object close enough to a target:

$$\forall \varepsilon > 0 \exists x d(x, a) < \varepsilon.$$

Remark 1.1.23 (Proof sketch). 1. Fix an arbitrary tolerance $\varepsilon > 0$.

2. Choose or construct an approximating object.
3. Estimate the error between the approximation and the target.
4. Show the error is below ε .

A Classification Pass

Before proving a theorem, classify its job:

- Does the statement depend on choices? Check well-definedness.
- Does it ask for an object? Prove existence.
- Does it say only one object can work? Prove uniqueness.
- Does it say an operation splits across sums and scalars? Prove linearity.
- Does one witness have to work for every point? Prove uniformity.
- Does it say two descriptions agree? Prove equivalence.
- Does it give a test for a definition? Prove a characterisation.
- Does it ask for smallness or control? Prove an estimate.
- Does it ask for two-sided order control? Prove boundedness.
- Does it ask for one-sided order control? Prove an upper or lower bound.
- Does it ask for a realized best element? Prove extremality.
- Does it ask for the best bound among all bounds? Prove least or greatest character.
- Does it preserve or reverse order? Prove monotonicity.
- Does it stay inside a class after an operation? Prove closure.
- Does it ask for arbitrary precision? Prove approximation.

Remark 1.1.24 (The point). The classification does not solve the proof, but it tells us what a completed proof must contain. For example, an existence proof must exhibit or obtain an object, while a uniqueness proof must compare two candidates. An equivalence proof must have two directions, while a uniformity proof must choose one witness before the arbitrary point is fixed.

1.1.2 Manipulating Inequalities and the Logic of Bounding

Inequality Toolkit — Quick Reference

Tool	Statement	Typical use
Triangle inequality	$ a + b \leq a + b $	Split a distance at an intermediate point
Distance form	$ a - b \leq a - c + c - b $	Introduce pivot c to separate two estimates
Reverse triangle ineq.	$ a - b \leq a - b $	Bound magnitudes using closeness
Absolute value interval	$ x \leq y \iff -y \leq x \leq y$	Convert $ \cdot $ bound to two-sided inequality
Boundedness replacement	$ b_n \leq M$ (constant): replace $ b_n $ by M	Eliminate variable factors in products
ε -budget split	$ A < \frac{\varepsilon}{2}$ and $ B < \frac{\varepsilon}{2} \Rightarrow A + B < \varepsilon$	Combine two independent estimates
+1 trick	Replace $ L $ by $ L + 1$ in denominators	Avoid division by zero when L may vanish

The Fundamental Asymmetry Between Equality and Inequality

Equality is bilateral: $a = b$ licenses substitution in either direction and carries no information about magnitude. An inequality is directional: $a \leq b$ locates a relative to b on the line, says nothing about the gap, and does not support backwards substitution.

This asymmetry is the source of a strategic shift. In an equality argument the goal is to show that two expressions are the same object. In a bounding argument the goal is to show that one expression is *controlled* by another. Control is inherently one-sided: a ceiling above $|f(x)|$ is useful regardless of how far below it $|f(x)|$ actually falls.

Remark 1.1.25 (Consequence for proof strategy). The question in analysis is rarely “what is $|a_n - L|$?” It is “is $|a_n - L|$ smaller than ε ?” A valid crude bound that answers the second question is more useful than an exact answer to the first. Deliberate over-estimation — absent from algebra — is a required habit.

Remark 1.1.26 (Common error). Treating an inequality as if it were an equality by reversing direction. Given $a \leq b$, negation gives $-a \geq -b$, not $-a \leq -b$. Multiplication by a negative constant reverses direction. Failing to track sign changes when multiplying or negating is the most frequent algebraic error in ε - δ computations.

The Basic Rules of Inequality Arithmetic

Proposition 1.1.27 (Order arithmetic). *Let $a, b, c, d \in \mathbb{R}$.*

- (i) *If $a \leq b$ and $c \leq d$, then $a + c \leq b + d$.*
- (ii) *If $a \leq b$ and $c > 0$, then $ac \leq bc$.*
- (iii) *If $a \leq b$ and $c < 0$, then $ac \geq bc$.*
- (iv) *$|x| \leq y$ (with $y \geq 0$) if and only if $-y \leq x \leq y$.*

Go to proof.

Remark.

Remark 1.1.28 (Fully quantified form). (i) $\forall a, b, c, d \in \mathbb{R}, (a \leq b) \wedge (c \leq d) \Rightarrow a + c \leq b + d$.
(iv) $\forall x \in \mathbb{R}, \forall y \geq 0, |x| \leq y \Leftrightarrow -y \leq x \leq y$.

Remark 1.1.29 (Inequalities add but do not subtract). Rule (i) licenses adding inequalities in the same direction. There is no subtraction rule: $a \leq b$ and $c \leq d$ yields no conclusion about $a - c$ versus $b - d$ in general. The direction of $a - c$ versus $b - d$ depends on the relative sizes of $(b - a)$ and $(d - c)$, which are not controlled.

Remark 1.1.30 (Consequence for absolute values). Rule (iv) is the bridge between $|\cdot|$ bounds and two-sided linear inequalities. It is invoked silently every time an ε - N argument converts $|a_n - L| < \varepsilon$ into positional information about a_n .




Figure 1.1: The absolute value characterisation as a symmetric interval. The shaded region is $[-y, y]$; the point x lies inside with $|x| \leq y$. The equivalence $|x| \leq y \Leftrightarrow -y \leq x \leq y$ (Proposition 1.1.27(iv)) converts an absolute value bound into a two-sided linear constraint.

The Triangle Inequality and the Pivot Move

Theorem (Triangle Inequality)

For all $a, b \in \mathbb{R}$, $|a + b| \leq |a| + |b|$. Equivalently, for all $a, b, c \in \mathbb{R}$, $|a - b| \leq |a - c| + |c - b|$.

Remark 1.1.31 (Fully quantified form). $\forall a, b \in \mathbb{R}, |a + b| \leq |a| + |b|$.

Remark 1.1.32 (The pivot move). The distance form $|a - b| \leq |a - c| + |c - b|$ is the operative version in analysis. The element c is a *pivot*: an intermediate point that decomposes a hard distance into two pieces, each controllable by hypothesis. In the proof that $a_n + b_n \rightarrow L + M$, the pivot $c = L + b_n$ gives

$$|(a_n + b_n) - (L + M)| = |(a_n - L) + (b_n - M)| \leq |a_n - L| + |b_n - M|,$$

and the two pieces are controlled by the individual convergences.

Remark 1.1.33 (Reverse triangle inequality). $||a| - |b|| \leq |a - b|$. Proof: apply the standard inequality to $a = (a - b) + b$ to get $|a| - |b| \leq |a - b|$; symmetry gives the other side. Consequence: $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}$ is Lipschitz with constant 1, so closeness of inputs implies closeness of magnitudes.

Remark 1.1.34 (Common error). Writing $|a| + |b| \leq |a + b|$. The triangle inequality always produces an *upper* bound on a sum, never a lower bound. The reverse ($|1| + |-1| = 2 > |1 + (-1)| = 0$) is a one-line counterexample.




Figure 1.2: The triangle inequality in distance form. The direct distance $|a - b|$ (red) is bounded above by the sum of indirect distances $|a - c|$ and $|c - b|$ (blue) through pivot c . In analysis, c is chosen to split $|a - b|$ into two pieces each controlled by a separate hypothesis.

Bounding Is Not Solving

In algebra, a computation terminates when the exact value is found. In analysis, a bounding argument terminates when the expression is shown to fall within an acceptable range. These are structurally different goals.

Remark 1.1.35 (Deliberate crudeness is valid). To bound $|x^2 - 9x + 1|$ on $[-1, 5]$:

$$|x^2 - 9x + 1| \leq |x|^2 + 9|x| + 1 \leq 25 + 45 + 1 = 71.$$

The actual supremum is far smaller than 71. That is irrelevant. The bound is valid, explicit, and was obtained cheaply. The test for a bound is not “is this sharp?” but “is this true and sufficient for the argument’s purpose?”

Remark 1.1.36 (Consequence for proof reading). A reader who checks that each inequality is valid can trust the conclusion without caring whether the bounds are optimal. Optimality is irrelevant to logical correctness.

The ε -Budget Split

The prototype of all two-piece bounding arguments:

$$|A + B| \leq |A| + |B| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

The total tolerance ε is a budget allocated among the summands. With k summands, allocate ε/k each; with summands of unequal difficulty, split unevenly provided the shares sum to at most ε .

Remark 1.1.37 (The four-step procedure). Standard structure of an ε - N or ε - δ proof:

1. Apply the triangle inequality: express $|f - L|$ as a sum $|A_1| + \cdots + |A_k|$.
2. Assign budgets $\varepsilon_i > 0$ with $\sum_i \varepsilon_i = \varepsilon$.

3. For each piece, find N_i (or δ_i) so that $|A_i| < \varepsilon_i$ for all $n \geq N_i$ (or $|x - a| < \delta_i$).
4. Set $N = \max_i N_i$ (or $\delta = \min_i \delta_i$) to enforce all constraints simultaneously.

Step 4 reflects the logical structure of the conclusion: all conditions must hold at once, so the threshold is the most restrictive one.

Remark 1.1.38 (Common error). Taking $N = N_1$ and ignoring N_2 . For $n \geq N_1$ the first piece is controlled; the second is not. Both pieces must be within budget simultaneously, which requires $n \geq \max\{N_1, N_2\}$.




Figure 1.3: The ε -budget split. The total tolerance ε is distributed as shares $\varepsilon/2$ between two pieces. Each piece is controlled independently for $n \geq N_i$. Taking $N = \max\{N_1, N_2\}$ enforces both bounds simultaneously, giving $|A + B| < \varepsilon$ for all $n \geq N$. With k pieces, assign ε/k each and take $N = \max_i N_i$.

The Boundedness Replacement

Proposition 1.1.39 (Convergent sequences are bounded). *If $(b_n) \rightarrow b$, then there exists $M > 0$ such that $|b_n| \leq M$ for all $n \in \mathbb{N}$. [Go to proof.](#)*

Remark.

Remark 1.1.40 (Fully quantified form). $\forall (b_n) \subset \mathbb{R}, \lim_{n \rightarrow \infty} b_n = b \Rightarrow \exists M > 0, \forall n \in \mathbb{N}, |b_n| \leq M$.

Remark 1.1.41 (The replacement move). Given $|b_n| \leq M$:

$$|b_n| \cdot |a_n - L| \leq M \cdot |a_n - L|.$$

The variable factor is gone. Choose N so that $|a_n - L| < \varepsilon/M$ for $n \geq N$; the whole product is then less than ε . The constant M need not be tight.

Remark 1.1.42 (Common error). Choosing $|a_n - L| < \varepsilon/|b_n|$. This is circular: N would depend on n , since $|b_n|$ changes with n . The threshold N must be a fixed number, independent of n . Replacing $|b_n|$ by the constant M breaks the circularity.

The Work-Backwards Principle

ε - N proofs are *stated* forwards but *discovered* backwards. The proof asserts: “Let $\varepsilon > 0$. Choose $N = \lceil C/\varepsilon \rceil$.” The value N was found by scratchwork: “the expression simplifies to C/n ; this is less than ε when $n > C/\varepsilon$; so set $N = \lceil C/\varepsilon \rceil$.”

The scratchwork is a private computation that produces the *witness*. The proof states the witness and verifies it, working forwards.

Scratchwork (private): Work backwards from $|a_n - L| < \varepsilon$ to find what N must satisfy.

Proof (public): State N explicitly. Verify $n \geq N \Rightarrow |a_n - L| < \varepsilon$.

Remark 1.1.43 (Common error). Allowing scratchwork to appear in the proof: “we need $n > C/\varepsilon$, so choose n this large.” The proof must assert a fixed N and verify the conclusion holds. A threshold stated as “ n large enough” is an incomplete scratchwork note, not a proof.

Remark 1.1.44 (Connection to the geometric phase). The work-backwards process is the analytic counterpart of the geometric phase in limit proofs. The geometric phase identifies structure and chooses a pivot; the backwards computation determines the explicit threshold. The proof states both and verifies the chain forwards.

Chaining Inequalities

A standard proof move builds a chain $|f(x)| \leq A(x) \leq B \leq \varepsilon$, where each step uses a different tool: the triangle inequality, a domain bound, and the choice of δ .

Remark 1.1.45 (Replacing by a larger quantity). A quantity on the right-hand side of $\leq$ may be freely replaced by anything larger, since this only weakens the inequality. If $|x| \leq |x-1|+1$ and $|x-1| < \delta$, then $|x| < \delta+1$. The replacement of $|x-1|$ by δ is valid because $|x-1|$ appears on the right. The same replacement on the *left* would be invalid.

Remark 1.1.46 (Common error). Including a step where the inequality points the wrong direction, or where a step holds only generically rather than pointwise. A chain is only as strong as its weakest link; every $\leq$ must be unconditionally valid.

The +1 Trick for Denominators

When $\varepsilon/|L|$ appears in an estimate and L may be zero, replace $|L|$ by $|L| + 1 \geq 1$:

$$\frac{\varepsilon}{|L| + 1} \leq \varepsilon.$$

Remark 1.1.47 (Fully quantified form). $\forall L \in \mathbb{R}$, $|L| + 1 \geq 1 > 0$. Hence $\varepsilon/(|L| + 1)$ is well-defined and positive for all $\varepsilon > 0$ and all $L \in \mathbb{R}$.

Remark 1.1.48 (Generalisation). The +1 is an instance of a wider principle: any constant intended for a denominator must be guaranteed strictly positive. If the constant is a given quantity C that might be zero, replace it by $C + 1$. The choice of 1 is conventional; any fixed positive number serves the same purpose.

Remark 1.1.49 (Common error). Using $|L|$ as a denominator without first verifying $L \neq 0$. The product limit proof $a_n b_n \rightarrow LM$ is the canonical location for this error: the bound $\varepsilon/(2|L|)$ fails when $L = 0$, and the $L = 0$ case requires separate treatment or the use of $|L| + 1$.

Summary: The Bounding Mindset

Algebra	Analysis
Find the exact value of x	Show $ x - L < \varepsilon$
Equality: both sides are the same object	Inequality: one side controls the other
Work backwards and forwards symmetrically	Discover witness backwards; verify forwards
Exact answer is the goal	Any valid bound is sufficient
Fractions must be simplified	Crude over-estimates are acceptable
Multiply both sides freely	Track sign changes; direction is load-bearing

Remark 1.1.50 (The structure of a bounding proof). A bounding proof is a *chain of control*: inequalities connecting the target expression to ε , where each link uses one of the toolkit tools above. The proof is complete when the chain reaches ε . Every link must be unconditionally valid; every variable factor must be replaced by a constant before the chain can close.

1.1.3 Completeness as a Constructive Engine

Completeness Construction Toolkit — Quick Reference

Tool	What it supplies
ε-characterisation of sup	$\forall \varepsilon > 0, \exists x \in S, x > \sup S - \varepsilon$
Inductive selection	Monotone $(x_n) \subset S$ with $x_n \rightarrow \sup S$
Monotone Approximation	Such a sequence exists for any nonempty bounded S
LUB $\Leftrightarrow$ MCT	The two completeness statements are equivalent

The ε -Characterisation of Suprema and Infima

Proposition 1.1.51 (ε -characterisation of sup). *Let $S \subset \mathbb{R}$ be nonempty and bounded above, and let $s = \sup S$. Then for every $\varepsilon > 0$ there exists $x \in S$ with $x > s - \varepsilon$.* [Go to proof.](#)

Remark.

Remark 1.1.52 (Fully quantified form). $\forall \varepsilon > 0, \exists x \in S, s - \varepsilon < x \leq s$. Equivalently: any $M < s$ is not an upper bound for S .

Remark 1.1.53 (Proof strategy). If no such x existed for some $\varepsilon_0 > 0$, then $s - \varepsilon_0$ would be an upper bound for S smaller than s , contradicting $s = \sup S$.

Remark 1.1.54 (Operative role). The ε -characterisation is the *existential engine* of constructive completeness arguments. At each inductive step it is invoked with a specific ε value, discharging the universal quantifier and yielding a concrete existence claim. Existential instantiation then names a witness — call it x_{n+1} — and the rest of the argument uses only the single property $x_{n+1} > s - \varepsilon$. The witness is underspecified; the argument makes no commitment about which element of S is chosen, and no further properties of x_{n+1} are used.

Remark 1.1.55 (Symmetric form for inf). For $t = \inf S$: for every $\varepsilon > 0$ there exists $y \in S$ with $y < t + \varepsilon$. The proof is symmetric: any $M > t$ is not a lower bound.

Inductive Selection at Bounds

Theorem (Inductive Selection at sup)

Let $S \subset \mathbb{R}$ be nonempty and bounded above, and let $s = \sup S$. Then there exists a strictly increasing sequence $(x_n)_{n=1}^\infty \subset S$ with $x_n \rightarrow s$.

Remark 1.1.56 (Fully quantified form). $\exists(x_n) \subset S: \forall n, x_n < x_{n+1}; \forall n, s - \frac{1}{n} < x_n \leq s; \lim_{n \rightarrow \infty} x_n = s$.

Remark 1.1.57 (Why no algebraic formula is possible in general). The set S may have large gaps; any fixed formula might land outside S entirely. The construction must be inductive, selecting from S at each stage using the ε -characterisation rather than evaluating a closed expression.

Remark 1.1.58 (Proof strategy — the max device). The inductive step must simultaneously satisfy two lower bounds: $x_{n+1} > x_n$ (monotonicity) and $x_{n+1} > s - \frac{1}{n+1}$ (proximity). A single invocation of the existential can impose only one threshold. Setting

$$M_n := \max\left(x_n, s - \frac{1}{n+1}\right)$$

collapses both requirements into one: $x_{n+1} > M_n$ delivers both $x_{n+1} > x_n$ and $x_{n+1} > s - \frac{1}{n+1}$ simultaneously. Since $M_n < s$ (both entries are strictly below s whenever $s \notin S$, or the constant sequence $x_n = s$ handles the case $s \in S$), the ε -characterisation with $\varepsilon = s - M_n > 0$ supplies the required witness.

Base case. Set $\varepsilon = 1$. The ε -characterisation gives $x_1 \in S$ with $x_1 > s - 1$; since $x_1 \in S$, also $x_1 \leq s$.

Inductive step. Given x_n with $s - \frac{1}{n} < x_n \leq s$, form M_n as above and instantiate the ε -characterisation with $\varepsilon = s - M_n > 0$ to obtain $x_{n+1} \in S$ with $x_{n+1} > M_n$. Then:

$$\begin{aligned} x_{n+1} > M_n \geq x_n &\Rightarrow x_{n+1} > x_n, \\ x_{n+1} > M_n \geq s - \frac{1}{n+1} &\Rightarrow s - \frac{1}{n+1} < x_{n+1} \leq s. \end{aligned}$$

Convergence. $0 \leq s - x_n < \frac{1}{n}$; the squeeze theorem and the Archimedean Property give $x_n \rightarrow s$.

Remark 1.1.59 (What the construction requires). Three ingredients: (i) completeness of $\mathbb{R}$ ($\sup S$ exists); (ii) the ε -characterisation ([Proposition 1.1.51](#)), supplying the witness at each step; (iii) the Archimedean Property ([Theorem 3.1.4](#)), forcing $|s - x_n| \leq \frac{1}{n} \rightarrow 0$. The internal structure of S — its cardinality, openness, density — plays no role.

Remark 1.1.60 (Common error). Instantiating the existential with only the proximity threshold $s - \frac{1}{n+1}$ and ignoring x_n . The resulting witness satisfies proximity but not monotonicity: one might obtain $x_1 = 0.9, x_2 = 0.4, x_3 = 0.85$, each close to s but not increasing. The max is not a convenience; it is the minimal device that encodes both requirements into one threshold.

Remark 1.1.61 (Infimum case). Replace max by min, $s - \frac{1}{n}$ by $t + \frac{1}{n}$, and reverse all strict inequalities. The resulting sequence $(y_n) \subset S$ is strictly decreasing with $y_n \rightarrow t = \inf S$.

Monotone Approximation of Bounds

Proposition 1.1.62 (Monotone Approximation of Bounds). *Let $S \subset \mathbb{R}$ be nonempty and bounded. Then there exist monotone sequences $(x_n), (y_n) \subset S$ with $x_n \rightarrow \sup S$ and $y_n \rightarrow \inf S$. [Go to proof.](#)*

Remark.

Remark 1.1.63 (Fully quantified form). $\forall S \neq \emptyset$ bounded: $\exists (x_n) \subset S, x_n \uparrow \sup S$; $\exists (y_n) \subset S, y_n \downarrow \inf S$.

Remark 1.1.64 (Proof strategy). The strictly increasing sequence from [Section 1.1.3](#) is already monotone and converges to $\sup S$. Taking $X_n = \max\{x_1, \dots, x_n\}$ makes it explicitly non-decreasing (useful when the inductive step does not guarantee strict increase). The infimum case is symmetric.

Remark 1.1.65 (Consequence). Every supremum is the limit of a sequence from the set. The sequential characterisation of $\sup$ and $\inf$ — that they are realised as limits of internal sequences — follows from this construction, making [Proposition 1.1.62](#) the canonical proof of that characterisation.

Remark 1.1.66 (Forward connections). The inductive selection pattern reappears in:

- **Monotone Convergence Theorem** — every increasing bounded sequence converges to the supremum of its range; this construction exhibits that the supremum is always realised as such a limit.
- **Nested Interval Property** — the left and right endpoints of nested intervals form monotone sequences converging to a common point by exactly this mechanism.
- **Existence of square roots** — the standard proof sets $s = \sup\{x \in \mathbb{R} : x^2 < 2\}$ and extracts an approximating sequence from that set by inductive selection.

LUB $\iff$ MCT: The Completeness Equivalence

Definition (Least Upper Bound Property)

$\mathbb{R}$ has the **least upper bound property** if every nonempty set $A \subset \mathbb{R}$ that is bounded above has a supremum in $\mathbb{R}$.

Theorem (LUB $\iff$ MCT)

The following are equivalent over the Archimedean ordered field axioms:

- $\mathbb{R}$ has the least upper bound property (LUB).
- Every bounded monotone sequence of real numbers converges (MCT).

Remark 1.1.67 (Fully quantified form). (i) $\forall A \subset \mathbb{R}, (A \neq \emptyset \wedge \exists M, \forall a \in A, a \leq M) \Rightarrow \exists \sup A \in \mathbb{R}$.

(ii) $\forall (x_n) \subset \mathbb{R}, (\forall n, x_n \leq x_{n+1}) \wedge (\exists M, \forall n, x_n \leq M) \Rightarrow \exists L \in \mathbb{R}, \lim_{n \rightarrow \infty} x_n = L$.

Remark 1.1.68 (Proof strategy — LUB $\Rightarrow$ MCT). Let (x_n) be increasing and bounded. The image set $A = \{x_n : n \in \mathbb{N}\}$ is bounded above, so LUB gives $\alpha = \sup A$. The ε -characterisation produces N with $x_N > \alpha - \varepsilon$. Monotonicity promotes this to a tail statement: $\alpha - \varepsilon < x_n \leq \alpha$ for all $n \geq N$. Hence $x_n \rightarrow \alpha$.

The quantifier structure is:

$$\alpha = \sup A \Rightarrow (\forall \varepsilon > 0) \exists N (x_N > \alpha - \varepsilon) \xrightarrow{\text{monotone}} (\forall \varepsilon > 0) \exists N (\forall n \geq N) (|x_n - \alpha| < \varepsilon).$$

Remark 1.1.69 (Proof strategy — MCT $\Rightarrow$ LUB). Given $A \neq \emptyset$ and bounded above by u_1 , pick $l_1 \in A$. Build a bisection bracket: at each stage, let $m_n = (l_n + u_n)/2$; if m_n is an upper bound for A set $u_{n+1} = m_n$, otherwise set $l_{n+1} = m_n$ (and pick a witnessing $a \in A$ with $a > l_{n+1}$). The sequences (l_n) and (u_n) are monotone and bounded, so MCT gives limits $l_n \rightarrow s$ and $u_n \rightarrow s$ (since $u_n - l_n = (u_1 - l_1)/2^n \rightarrow 0$). The invariants — “ u_n is an upper bound for A ” and “ l_n is not” — force $s = \sup A$.

The construction maintains:

- a *universal* invariant: $(\forall a \in A)(a \leq u_n)$,
- a *failure* invariant: $(\exists a \in A)(a > l_n)$.

These two invariants, together with $u_n - l_n \rightarrow 0$, pin s as the least upper bound.

Remark 1.1.70 (Why this matters structurally). The chain LUB $\Leftrightarrow$ MCT $\Leftrightarrow$ NIP $\Leftrightarrow$ BW $\Leftrightarrow$ CC (Cauchy Criterion) is the backbone of the equivalence cluster at the heart of real analysis. All five statements assert that $\mathbb{R}$ has no gaps, each in its own language. The inductive selection technique — building sequences from sets using the ε -characterisation — is the constructive face of all five equivalences.

Remark 1.1.71 (Downstream reuse). The quantifier pattern

$$(\forall \varepsilon > 0)(\exists \text{witness}) \Rightarrow (\exists \text{sequence with controlled tail})$$

reappears in Bolzano–Weierstrass (extracting convergent subsequences), $\limsup/\liminf$ (tail suprema as a decreasing sequence), compactness (finite subcovers as finite witnesses), and approximation theorems throughout analysis.

1.1.4 Walking a Sequence with a Predicate

Predicate-Walking Toolkit — Quick Reference

Theorem	Predicate $P(n)$	Tool
Monotone Subseq.	n is a peak of (x_n)	Dichotomy on peak count
Bolzano–Weierstrass	interval has ∞ many terms	Bisection + pigeonhole
Arzelà–Ascoli	converges at $d_1, \dots, d_m$	Diagonal extraction

The Infinitely-Often / Eventually Dichotomy

Definition (Infinitely Often / Eventually)

Let (x_n) be a sequence and P a predicate on $\mathbb{N}$. P holds **infinitely often** if $\{n \in \mathbb{N} : P(n)\}$ is infinite; P holds **eventually** if there exists $N \in \mathbb{N}$ such that $P(n)$ holds for all $n \geq N$.

Remark 1.1.72 (Fully quantified form). P infinitely often: $\forall N \in \mathbb{N}, \exists n \geq N, P(n)$.

P eventually: $\exists N \in \mathbb{N}, \forall n \geq N, P(n)$.

Proposition 1.1.73 (The dichotomy). *For any predicate P on $\mathbb{N}$, exactly one of the following holds:*

(i) P holds infinitely often.

(ii) $\neg P$ holds eventually.

Go to proof.

Remark.

Remark 1.1.74 (Fully quantified form). $(\forall N, \exists n \geq N, P(n)) \vee (\exists N, \forall n \geq N, \neg P(n))$.

The two cases are exhaustive because $\mathbb{N}$ is infinite: if P fails for all but finitely many n , those indices have a maximum N , and $\neg P(n)$ holds for all $n > N$.

Remark 1.1.75 (Relation to Infinite Pigeonhole). The index set $\mathbb{N}$ is partitioned into $\{n : P(n)\}$ and $\{n : \neg P(n)\}$; an infinite set cannot have both parts finite. The “infinitely often” case provides an inexhaustible reservoir of indices from which to select subsequence terms inductively.

The Abstract Extraction Template

Remark 1.1.76 (Four-step pattern). Every predicate-walking argument follows the same skeleton:

1. *Define the predicate.* Identify the property $P(n)$ to propagate: peak, infinite-occupancy, pointwise convergence, etc.
2. *Apply the dichotomy.* Either P holds infinitely often, or $\neg P$ holds eventually. Handle each case.
3. *Extract the subsequence.* In the “infinitely often” case, enumerate indices satisfying P in increasing order: $n_1 < n_2 < \dots$. In the “eventually” case, work in the $\neg P$ -tail and build inductively, using $\neg P$ as the fuel.
4. *Verify the property.* Show (x_{n_k}) has the desired property, using the predicate and any additional hypotheses (boundedness, MCT, NIP, equicontinuity).

The critical structural requirement: at each inductive stage, one must find $n_{k+1} > n_k$ satisfying the relevant condition. The “infinitely often” guarantee ensures the reservoir is never exhausted.

The Peak Argument: Monotone Subsequence Theorem

Definition (Peak Index)

An index $m \in \mathbb{N}$ is a **peak** of (x_n) if $x_m \geq x_n$ for all $n \geq m$.

Remark 1.1.77 (Fully quantified form). m is a peak: $\forall n \in \mathbb{N}, n \geq m \Rightarrow x_m \geq x_n$.

Theorem (Monotone Subsequence)

Every sequence (x_n) in $\mathbb{R}$ has a monotone subsequence.

Remark 1.1.78 (Fully quantified form). $\forall (x_n) \subset \mathbb{R}, \exists (n_k)$ strictly increasing, (x_{n_k}) is monotone

Remark 1.1.79 (Proof strategy — peak dichotomy). Apply the dichotomy to $P(m) = “m$ is a peak.”

Case 1: infinitely many peaks. Let $n_1 < n_2 < \dots$ enumerate the peaks. Since n_k is a peak and $n_{k+1} > n_k$: $x_{n_k} \geq x_{n_{k+1}}$. The subsequence is decreasing.

Case 2: finitely many peaks. Let N be an index beyond all peaks. No $n \geq N$ is a peak, so for each such n there exists $m > n$ with $x_m > x_n$. Set $n_1 = N$. Since n_1 is not a peak, $\exists n_2 > n_1$ with $x_{n_2} > x_{n_1}$. Inductively, this produces a strictly increasing subsequence.

The no-peak condition $\neg P(n_k)$ is the fuel for each inductive step: it is precisely the statement “there exists a later term exceeding x_{n_k} ,” which supplies n_{k+1} .

Remark 1.1.80 (Consequence: Bolzano–Weierstrass). Every bounded sequence has a bounded monotone subsequence. By MCT, that subsequence converges. Bolzano–Weierstrass follows immediately.

Remark 1.1.81 (Common error). Assuming Case 2 produces an increasing subsequence “by default.” It does, but only because $\neg P(n)$ supplies a strictly larger term. If the predicate were defined differently, $\neg P$ might provide no useful structure. The conclusion depends on the specific content of $\neg P$, not the abstract case structure.




Figure 1.4: The two cases of the Monotone Subsequence Theorem (Section 1.1.4). *Left:* Infinitely many peaks (blue) yield a decreasing subsequence (dashed blue). *Right:* After the last peak, every $n \geq N$ is a non-peak, so a later term always exceeds the current one; this fuels the inductive construction of an increasing subsequence (dashed green).

The Infinite-Occupancy Predicate: Bolzano–Weierstrass via Bisection

Proposition 1.1.82 (Bolzano–Weierstrass via bisection). *Every bounded sequence in $\mathbb{R}$ has a convergent subsequence.* [Go to proof.](#)

Remark.

Remark 1.1.83 (Proof strategy — infinite-occupancy predicate). The predicate at each stage: $P(I)$ = “interval I contains infinitely many terms of (x_n) .” Since $(x_n) \subseteq [a_0, b_0]$ for some bounded interval, $P([a_0, b_0])$ holds.

Inductive step. Given $I_k = [a_k, b_k]$ with $P(I_k)$, bisect at $m_k = (a_k + b_k)/2$. The two halves cover I_k ; at least one contains infinitely many terms. Set I_{k+1} to be that half. Pick $x_{n_{k+1}} \in I_{k+1}$ with $n_{k+1} > n_k$. The predicate $P(I_{k+1})$ holds by construction.

Convergence. The intervals have $|I_k| = (b_0 - a_0)/2^k \rightarrow 0$. By NIP there exists a unique $L \in \bigcap I_k$. Since $a_k \leq x_{n_k} \leq b_k$ and both endpoints converge to L , the squeeze theorem gives $x_{n_k} \rightarrow L$.

Remark 1.1.84 (Comparison with the peak proof). The peak proof is combinatorial: it operates on the values of the sequence directly and produces a monotone subsequence without requiring boundedness. Convergence follows from MCT.

The bisection proof is geometric: it operates on intervals containing the sequence’s range and produces a subsequence squeezed to a limit directly. Boundedness is required from the outset to produce the initial interval.

Both proofs use the same dichotomy, but in the peak proof it is applied to an index predicate; in the bisection proof, to an interval predicate.

Diagonal Extraction: Arzelà–Ascoli

Remark 1.1.85 (Diagonal extraction pattern). The diagonal argument generalises predicate-walking to sequences of functions. Let (f_n) be bounded and equicontinuous in $C^0([a, b], \mathbb{R})$, and let $D = \{d_1, d_2, \dots\} = \mathbb{Q} \cap [a, b]$.

Stage 1. $(f_n(d_1))$ is bounded; extract by BW a subsequence $(f_{1,k})$ converging at d_1 .

Stage 2. $(f_{1,k}(d_2))$ is bounded; extract a sub-subsequence $(f_{2,k})$ converging at d_2 . As a subsequence of $(f_{1,k})$, it still converges at d_1 .

Stage m . Extract $(f_{m,k})$ as a subsequence of $(f_{m-1,k})$ converging at d_m . By induction, $(f_{m,k})$ converges at $d_1, \dots, d_m$.

Diagonal step. Define $g_m = f_{m,m}$. For any fixed j and $m \geq j$, the term $g_m = f_{m,m}$ is a term of $(f_{j,k})$ with index $k = m \geq j$. Hence $g_m(d_j) \rightarrow y_j$. The sequence (g_m) converges pointwise on all of D .

Equicontinuity then propagates pointwise convergence on D to uniform convergence on $[a, b]$ via an $\varepsilon/3$ argument.

Remark 1.1.86 (Why the diagonal works). At stage m , the predicate “converges at $d_1, \dots, d_m$ ” holds for all terms of $(f_{m,k})$. The diagonal term $g_m = f_{m,m}$ is a tail term of every earlier row $f_{j,k}$ for $j \leq m$, so it satisfies the predicate for all $j \leq m$ simultaneously. Sending $m \rightarrow \infty$ gives convergence at every d_j .

	d_1	d_2	d_3	d_4	$\dots$
Stage 1: $(f_{1,k})$	conv.	?	?	?	
Stage 2: $(f_{2,k})$	conv.	conv.	?	?	
Stage 3: $(f_{3,k})$	conv.	conv.	conv.	?	
Stage 4: $(f_{4,k})$	conv.	conv.	conv.	conv.	

(g_m) converges pointwise on all of $D = \{d_1, d_2, \dots\}$

Figure 1.5: Diagonal extraction in the Arzelà–Ascoli argument. Each row $(f_{m,k})$ converges at $d_1, \dots, d_m$ (green) and is a subsequence of the row above. The diagonal $g_m = f_{m,m}$ (red) is a tail of every row, so it converges at every d_j . Equicontinuity then extends pointwise convergence on the dense set D to uniform convergence on $[a, b]$.

Limitations of the Technique

Remark 1.1.87 (The predicate must be decidable at each stage). At each inductive step, one must determine which branch of the dichotomy applies. If the predicate requires knowledge of the limit — a quantity not yet known — the construction cannot proceed. Predicate-walking requires a finitary, combinatorial, or topological condition on the terms already seen.

Remark 1.1.88 (Multiple simultaneous predicates may exhaust the reservoir). The conjunction $P \wedge Q$ can hold only finitely often even when P and Q each hold infinitely often. The diagonal argument avoids this by handling one predicate at a time, composing via the diagonal.

Remark 1.1.89 (Subsequences, not sequences). Predicate-walking yields existence of a subsequence with the desired property, not a conclusion about the full sequence. The Monotone Subsequence Theorem does not assert (x_n) is monotone, or that the extracted subsequence is unique.

Remark 1.1.90 (Convergence requires a separate argument). The extracted subsequence has structural properties but not yet a limit. Converting structure into convergence requires a separate theorem: monotone subsequence $\rightarrow$ MCT (requires boundedness); bisection subsequence $\rightarrow$ NIP and squeeze; pointwise convergence on $D \rightarrow$ equicontinuity and $\varepsilon/3$.

Remark 1.1.91 (Quantitative control is absent). Predicate-walking proofs are pure existence arguments. They specify neither the rate of convergence nor which indices are selected. Explicit rates or constructive witnesses require a different approach.

Remark 1.1.92 (Countability is essential for the diagonal). The Arzelà–Ascoli diagonal requires a countable dense D so that points can be enumerated and handled one at a time. In non-separable metric spaces (those without a countable dense subset) the argument breaks down.

The Unifying Principle

Remark 1.1.93 (Infinite Pigeonhole as the common core). Every predicate-walking argument rests on the fact that an infinite set cannot be partitioned into two finite parts. Applied to $\mathbb{N}$:

if P holds only finitely often, the P -indices have a maximum, and $\neg P$ holds from that point on. The method reduces a complex existence question to a sequence of binary choices, each justified by the Infinite Pigeonhole Principle. The construction is inductive, automatic, and never runs dry, provided the predicate is well-chosen and the reservoir is genuinely infinite at each stage.

Remark 1.1.94 (Dependency summary). 1. Monotone Subsequence Theorem — peak dichotomy alone; no completeness.

2. Bolzano–Weierstrass via MCT — Monotone Subseq. + MCT (requires completeness via AoC).

3. Bolzano–Weierstrass via bisection — NIP + squeeze (requires completeness via NIP $\equiv$ AoC).

4. Arzelà–Ascoli — BW at each diagonal stage + equicontinuity.

The Monotone Subsequence Theorem is the only result in the cluster requiring no completeness. All others depend on completeness via MCT or NIP.

1.1.5 Interval Bisection and Nested Intervals

Bisection and NIP Toolkit — Quick Reference

Label	Statement	Proof method
NIP	$\bigcap_{n=1}^{\infty} I_n \neq \emptyset$ for nested closed bounded intervals	Supremum of left endpoints
IVT	f continuous, $f(a) < L < f(b) \Rightarrow \exists c \in (a, b), f(c) = L$	Bisection + NIP
UNC	$\mathbb{R}$ is uncountable	Nested interval diagonalisation

The Nested Interval Property

Theorem (Nested Interval Property)

For each $n \in \mathbb{N}$, let $I_n = [a_n, b_n]$ be a closed bounded interval with $I_1 \supseteq I_2 \supseteq I_3 \supseteq \cdots$.
Then $\bigcap_{n=1}^{\infty} I_n \neq \emptyset$.

Remark 1.1.95 (Fully quantified form). $\forall n \in \mathbb{N}, I_n = [a_n, b_n] \neq \emptyset, [a_{n+1}, b_{n+1}] \subseteq [a_n, b_n] \Rightarrow \exists x \in \mathbb{R}, \forall n, x \in I_n$.

Remark 1.1.96 (Proof strategy). Let $A = \{a_n : n \in \mathbb{N}\}$. Nestedness implies every b_n is an upper bound for A . By AoC, $x = \sup A$ exists. Then $a_n \leq x \leq b_n$ for all n , so $x \in I_n$ for all n .

Remark 1.1.97 (Closedness is essential). The open intervals $(0, 1/n)$ are nested but $\bigcap (0, 1/n) = \emptyset$. Closedness ensures the endpoints can serve as witnesses in the AoC argument.

Remark 1.1.98 (Boundedness is essential). The unbounded closed intervals $[n, \infty)$ are nested but their intersection is empty. Both hypotheses are necessary.

Remark 1.1.99 (Singleton intersection). If additionally $b_n - a_n \rightarrow 0$, the intersection is a singleton $\{x\}$, since $a_n \leq x \leq b_n$ and the squeeze theorem forces $a_n \rightarrow x$ and $b_n \rightarrow x$. This strengthened form is used in most applications.




Figure 1.6: The Nested Interval Property (Section 1.1.5). Left endpoints a_n increase; right endpoints b_n decrease; each b_n is an upper bound for $\{a_n\}$. By the Axiom of Completeness, $x = \sup\{a_n\}$ exists and satisfies $a_n \leq x \leq b_n$ for all n , so $x \in \bigcap I_n$.

Remark 1.1.100 (Equivalence with completeness). NIP is equivalent to AoC over the Archimedean ordered field axioms. In particular, NIP fails over $\mathbb{Q}$: the intervals $[1, \sqrt{2}] \cap \mathbb{Q}$ shrink to a gap that $\mathbb{Q}$ cannot fill.

The Bisection Technique

Definition (Bisection Sequence)

Let $I_0 = [a_0, b_0]$ carry a property P . The **bisection sequence** (I_n) is defined inductively: given $I_n = [a_n, b_n]$, let $m_n = (a_n + b_n)/2$ and set

$$I_{n+1} = \begin{cases} [a_n, m_n] & \text{if } [a_n, m_n] \text{ inherits } P, \\ [m_n, b_n] & \text{otherwise.} \end{cases}$$

Remark 1.1.101 (Fully quantified form). $I_0 = [a_0, b_0]$; $\forall n, I_{n+1} \subseteq I_n$; $|I_n| = (b_0 - a_0)/2^n$.

Remark 1.1.102 (Length control). Bisection gives automatic length control: $|I_n| = (b_0 - a_0)/2^n \rightarrow 0$. No separate argument for the length condition of NIP is required.

Remark 1.1.103 (Abstract template). Every bisection proof has the same five-step shape:

1. *Initialize*. Identify P and starting interval I_0 .
2. *Bisect*. Check which half inherits P .
3. *Extract*. Apply NIP to obtain $x \in \bigcap I_n$; note $a_n \rightarrow x$ and $b_n \rightarrow x$.
4. *Conclude*. Pass P through the limit, using continuity, MCT, or squeeze as appropriate.

2. *NIP path*. Bisection as above.

The two paths are equivalent over the Archimedean ordered field axioms.

Remark 1.1.108 (Applied interpretation). The bisection construction is directly algorithmic: it locates a root to any desired precision in finitely many steps. After n bisections the root is known to within $(b - a)/2^n$. This is the bisection method of numerical analysis.

Uncountability of $\mathbb{R}$ via Nested Intervals

Proposition 1.1.109. $\mathbb{R}$ is uncountable. [Go to proof](#).

Remark.

Remark 1.1.110 (Proof strategy — nested interval diagonalisation). Suppose for contradiction $\mathbb{R} = \{x_1, x_2, \dots\}$. Build nested intervals (I_n) inductively with $I_{n+1} \subseteq I_n$ and $x_n \notin I_{n+1}$: at each step I_n contains two disjoint closed subintervals; x_{n+1} lies in at most one, so choose the other.

By NIP, $\bigcap I_n \neq \emptyset$: pick $y \in \bigcap I_n$. Then $y \neq x_n$ for every n (since $y \in I_n$ but $x_n \notin I_n$), contradicting the assumption that $\{x_n\}$ enumerates $\mathbb{R}$.

Remark 1.1.111 (Relation to Cantor’s diagonal argument). This is the geometric form of Cantor’s diagonalisation: instead of altering a decimal digit, the n th real is excluded from the n th nested interval. The witness y is the interval analogue of the “diagonal” real.

Summary: When to Use Each Technique

	Feature	Bisection
<i>Remark 1.1.112</i> (Bisection vs. general nested intervals).	How built	Halving at midpoints
	Length control	Automatic: $(b_0 - a_0)/2^n$
	Typical use	Property propagation via sign
	Canonical examples	IVT, BW
	Guarantor	NIP $\equiv$ AoC

Remark 1.1.113 (Completeness is indispensable). Neither technique works over $\mathbb{Q}$. The bisection procedure for $f(x) = x^2 - 2$ on $[1, 2]$ runs mechanically over $\mathbb{Q}$, but the limiting point $\sqrt{2}$ is not in $\mathbb{Q}$. Every bisection proof is, at bottom, an appeal to the completeness of $\mathbb{R}$.

Remark 1.1.114 (Cross-section of the chapter). The four sections of this chapter are not independent: each builds on the last. Inequality bounding (§1.1.2) requires only the ordered field axioms. Completeness as a constructive engine (§1.1.3) requires AoC and the Archimedean Property. Predicate-walking (§1.1.4) requires completeness via MCT or NIP, depending on the variant. Bisection and NIP (§1.1.5) is the geometric synthesis: it encapsulates completeness in a form that drives the IVT, BW, and uncountability proofs simultaneously, connecting back to both the inductive-selection constructions of §1.1.3 and the predicate-walking patterns of §1.1.4.

1.1.6 Subsequence Partition via Residue Classes

Residue Partition Toolkit — Quick Reference

Tool	Statement	Use
k -Periodicity Criterion	$(a_n) \rightarrow L \Leftrightarrow \forall r, (a_{kn+r}) \rightarrow L$	Prove convergence by analysing k c
Forward direction	Subseqs. of convergent seqs. converge to L	Reduce to subsequence analysis
Divergence corollary	Two classes $\rightarrow$ different limits $\Rightarrow (a_n)$ diverges	Detect divergence by comparing res
Index translation	$m = kn + r \Rightarrow n \geq N_r$ once $m \geq k \max_r N_r$	Core of the backward direction

The Residue-Class Partition

For any sequence (a_n) and any integer $k \geq 2$, the division algorithm partitions $\mathbb{N}$ into k disjoint residue classes:

$$\mathbb{N} = \bigsqcup_{r=0}^{k-1} \{kn + r : n \geq 0\}.$$

Each class defines a subsequence: $(a_{kn+r})_{n \geq 0}$ selects every k th term of (a_n) beginning at position r . These k subsequences partition (a_n) exactly — every term appears in precisely one class, order within each class is preserved, and their union recovers (a_n) .

Remark 1.1.115 (Fully quantified form of the partition). $\forall n \in \mathbb{N}, \exists! r \in \{0, \dots, k-1\}, \exists! q \in \mathbb{N}, n = kq + r$. Hence $\{(a_{kn+r})_{n \geq 0} : r = 0, \dots, k-1\}$ is a partition of (a_n) into k subsequences.




Figure 1.8: The residue-class partition of (a_n) with modulus $k = 3$. Each term is coloured by residue class: blue ($r = 0$), red ($r = 1$), green ($r = 2$). The partition is exact; the k -Periodicity Criterion ([Section 1.1.6](#)) states that $(a_n) \rightarrow L$ if and only if each coloured subsequence converges to the same L .

The k -Periodicity Convergence Criterion

Theorem (k -Periodicity Convergence Criterion)

Let (a_n) be a sequence, $k \geq 2$ an integer, and $L \in \mathbb{R}$. The following are equivalent:

- (i) $(a_n) \rightarrow L$.
- (ii) For every $r \in \{0, 1, \dots, k-1\}$, the subsequence $(a_{kn+r}) \rightarrow L$.

Remark 1.1.116 (Fully quantified form). $(i) \Leftrightarrow (ii)$, where:

(i): $\forall \varepsilon > 0, \exists N \in \mathbb{N}, \forall m \geq N, |a_m - L| < \varepsilon$.

(ii): $\forall r \in \{0, \dots, k-1\}, \forall \varepsilon > 0, \exists N_r \in \mathbb{N}, \forall n \geq N_r, |a_{kn+r} - L| < \varepsilon$.

Remark 1.1.117 (Proof strategy). **Forward direction** $(i) \Rightarrow (ii)$. Each (a_{kn+r}) is a subsequence of (a_n) . Subsequences of a convergent sequence converge to the same limit ([Proposition 1.1.39](#)).

Backward direction $(ii) \Rightarrow (i)$. The operative direction. Given $\varepsilon > 0$, for each residue r let N_r satisfy $|a_{kn+r} - L| < \varepsilon$ for all $n \geq N_r$. Set $N^* = k \cdot \max_r N_r$. For any $m \geq N^*$, write $m = kn + r$ via the division algorithm. Then:

$$n = \frac{m - r}{k} \geq \frac{N^* - (k - 1)}{k} \geq \frac{k \max_r N_r - (k - 1)}{k} \geq N_r,$$

so $|a_m - L| = |a_{kn+r} - L| < \varepsilon$.

Remark 1.1.118 (The index translation). The key step converts a global index m to a local index n within its residue class via $m = kn + r$. The factor k in the definition of N^* absorbs the offset $r \leq k - 1$ from every class simultaneously, ensuring that $n \geq N_r$ holds for whichever class m lands in.

The Divergence Corollary

Proposition 1.1.119 (Residue divergence criterion). *If there exist residues $r, s \in \{0, \dots, k-1\}$ such that $(a_{kn+r}) \rightarrow L$ and $(a_{kn+s}) \rightarrow M$ with $L \neq M$, then (a_n) diverges.* [Go to proof.](#)

Remark.

Remark 1.1.120 (Fully quantified form). $(a_{kn+r}) \rightarrow L \wedge (a_{kn+s}) \rightarrow M \wedge L \neq M \Rightarrow (a_n)$ diverges.

Remark 1.1.121 (Proof strategy). Contrapositively: if $(a_n) \rightarrow L$, then every subsequence converges to L (forward direction of [Section 1.1.6](#)). If two residue-class subsequences converge to different limits, (a_n) cannot converge.

Remark 1.1.122 (Canonical example). The sequence $((-1)^n)$ has $a_{2n} = 1 \rightarrow 1$ and $a_{2n+1} = -1 \rightarrow -1$. Since $1 \neq -1$, the sequence diverges — established immediately without any ε - N computation.

The Alternating Series Test via Even/Odd Partition

The $k = 2$ case of [Section 1.1.6](#) drives the cleanest proof of the Alternating Series Test.

Theorem (Alternating Series Test)

Let (a_n) be a positive decreasing sequence with $a_n \rightarrow 0$. Then the alternating series $\sum_{n=1}^{\infty} (-1)^{n+1} a_n$ converges.

Remark 1.1.123 (Fully quantified form). $(\forall n, a_n > 0) \wedge (\forall n, a_{n+1} \leq a_n) \wedge (a_n \rightarrow 0) \Rightarrow \exists L, \sum_{n=1}^{\infty} (-1)^{n+1} a_n = L$.

Remark 1.1.124 (Proof strategy — even/odd partition). Let s_n denote the n th partial sum. Apply [Section 1.1.6](#) with $k = 2$: it suffices to show $(s_{2k}) \rightarrow L$ and $(s_{2k+1}) \rightarrow L$ for the same L .

Even subsequence (s_{2k}) : decreasing and bounded. Group consecutive pairs:

$$s_{2k} = \sum_{j=1}^k (-a_{2j-1} + a_{2j}).$$

Since (a_n) is decreasing, each summand $-a_{2j-1} + a_{2j} \leq 0$, so (s_{2k}) is decreasing. Also:

$$s_{2k} = -a_1 + \sum_{j=1}^{k-1} (a_{2j} - a_{2j+1}) + a_{2k} \geq -a_1,$$

since each term in the sum is non-negative. Thus (s_{2k}) is decreasing and bounded below by $-a_1$. By MCT, $s_{2k} \rightarrow L$ for some $L \geq -a_1$.

Odd subsequence (s_{2k+1}) : linked to the even. Observe $s_{2k+1} = s_{2k} + a_{2k+1}$. Since $s_{2k} \rightarrow L$ and $a_{2k+1} \rightarrow 0$, the Algebraic Limit Theorem gives $s_{2k+1} \rightarrow L$.

Applying the criterion. Both residue-class subsequences converge to the same L . By [Section 1.1.6](#) with $k = 2$, $(s_n) \rightarrow L$.

Remark 1.1.125 (Why the partition is the right tool). The even and odd partial sums have structurally different characters: the even sums are monotone (enabling MCT), the odd sums are then controlled via the algebraic relationship $s_{2k+1} = s_{2k} + a_{2k+1}$. No single uniform argument handles both. The partition separates the two cases into independent analyses, then the criterion reassembles the conclusion.




Figure 1.9: Even and odd partial sums of an alternating series $\sum (-1)^{n+1} a_n$ with (a_n) positive, decreasing, and tending to zero. Even sums (s_{2k}) (blue) decrease and are bounded below; by MCT they converge to L . Odd sums (s_{2k+1}) (red) satisfy $s_{2k+1} = s_{2k} + a_{2k+1} \rightarrow L$. Since both residue classes (mod $k = 2$) converge to L , the k -Periodicity Criterion ([Section 1.1.6](#)) gives $(s_n) \rightarrow L$.

Remark 1.1.126 (Alternative proofs). The Alternating Series Test admits three proofs from different completeness avatars:

1. *Cauchy criterion* — show (s_n) is eventually ε -steady directly.

2. *NIP* — the nested intervals $[s_{2k}, s_{2k-1}]$ shrink to a point by the Nested Interval Property.
3. *MCT + residue partition* — as above.

The partition proof is the most transparent structurally: the monotone behaviour and the limit identity are isolated into the two residue classes rather than handled simultaneously.

Higher k and Choosing the Right Period

For sequences with period $k > 2$, the criterion applies directly. The sequence $a_n = \cos(2\pi n/3)$ takes values $1, -\frac{1}{2}, -\frac{1}{2}, 1, -\frac{1}{2}, -\frac{1}{2}, \dots$, cycling with period 3. Partitioning modulo 3:

$$\begin{aligned} a_{3n} &= 1 \rightarrow 1, \\ a_{3n+1} &= -\frac{1}{2} \rightarrow -\frac{1}{2}, \\ a_{3n+2} &= -\frac{1}{2} \rightarrow -\frac{1}{2}. \end{aligned}$$

The three limits differ ($1 \neq -\frac{1}{2}$), so the sequence diverges by [Proposition 1.1.119](#).

Remark 1.1.127 (Choosing k to match the period). The right choice of k is the natural period of the sequence’s structure. Using $k = 2$ for a period-3 sequence produces residue classes that are themselves period-3 subsequences; convergence of each class then requires a further partition with $k = 3$. Choosing k to match the period compresses the argument to a single application of the criterion.

Remark 1.1.128 (Applying the criterion to prove convergence). To prove $(a_n) \rightarrow L$ via the criterion, one must:

1. Select k matching the period of the sequence’s behaviour.
2. Prove $(a_{kn+r}) \rightarrow L$ for each $r \in \{0, \dots, k-1\}$, using whatever tool fits that residue class.
3. Invoke [Section 1.1.6](#) to conclude $(a_n) \rightarrow L$.

The method does not produce L — it confirms a candidate. Identifying L requires separate work (direct computation, MCT applied to a class, etc.) before the criterion is invoked.

Contrast with Predicate-Walking

Remark 1.1.129 (Dual logical directions). The residue partition technique and predicate-walking (§1.1.4) operate in opposite logical directions.

Feature	Predicate-walking	Residue partition
Flow	Whole $\rightarrow$ one part	All parts $\rightarrow$ whole
Goal	Property of extracted subseq.	Convergence of full (a_n)
Coverage	One “good” subsequence	All k residue classes
Divergence use	Two classes, different limits	Same: two classes, different limits
Completeness role	Via MCT or NIP	Via MCT or algebraic limit theorem

Predicate-walking extracts what is needed; the residue partition reassembles from all pieces. The divergence criterion is available in both frameworks: two subsequences with different limits immediately imply divergence.

Limitations

Remark 1.1.130 (The same-limit condition is non-negotiable). The backward direction of [Section 1.1.6](#) requires all k residue-class subsequences to converge to the *same* L . There is no weakening: a convergent sequence has all its subsequences converging to the identical limit, so any two classes converging to different limits already certify divergence. The technique cannot produce convergence from mixed limits.

Remark 1.1.131 (The limit must be known in advance). The criterion confirms a candidate limit L ; it does not produce L from scratch. In practice, L is identified by direct computation on one residue class (e.g., the even partial sums converge to L by MCT), then the criterion assembles global convergence.

Remark 1.1.132 (Exact periodicity is required). The technique applies when the sequence's qualitative behaviour repeats with exact period k in the index. For sequences whose oscillation decays without exact periodicity, the residue classes are not individually tractable, and a direct ε - N argument or a squeeze is more appropriate.

Remark 1.1.133 (Common error). Applying the criterion with k smaller than the true period, then concluding convergence after verifying only the even residue class. If the odd class is not verified, the conclusion does not follow. Every residue class must be checked.

Remark 1.1.134 (Forward connections). The residue partition technique reappears in:

- **Cesàro means** — the Cesàro average of a sequence (a_n) can be analysed by partitioning the partial sums of the average into residue classes.
- **Fourier series** — the convergence of partial sums of a Fourier series is often established by analysing even and odd partial sums (or partial sums along specific subsequences of the approximation order).
- **Recursively defined sequences** — when a recurrence has period k , the residue-class subsequences each satisfy a simpler recurrence, enabling separate analysis.

1.2 Asymptotic Notation

Asymptotic Notation — Quick Reference

Concept/Statement	Meaning/Role
Little-o at a point	One function is negligible relative to another near a point.
Little-o at infinity	Negligibility along an unbounded input range.
Increment little-o	The local error notation $r(h) = o(h)$ as $h \rightarrow 0$.
Quotient criterion	Little- o is detected by a quotient tending to 0.
Sum rule	Negligible errors add.
Bounded factor rule	Bounded multipliers preserve negligible errors.

Little-o Notation

Asymptotic notation separates the main term of an expression from the part that becomes negligible in a limiting process. The notation $f = o(g)$ means that f is smaller than every fixed multiple of g near the limiting point. It does not say that f is zero. It says that f is negligible relative to the chosen scale g .

Definition 1.2.1 (Little-o at a Point). Let $a \in \mathbb{R}$, let f and g be real-valued functions defined on a punctured neighbourhood of a , and suppose $g(x) \neq 0$ for all x sufficiently close to a with $x \neq a$. We write

$$f(x) = o(g(x)) \quad (x \rightarrow a)$$

if and only if for every $\varepsilon > 0$ there exists $\delta > 0$ such that

$$0 < |x - a| < \delta \implies |f(x)| \leq \varepsilon |g(x)|.$$

Remark (Standard quantified statement).

$$f(x) = o(g(x)) \ (x \rightarrow a) \iff \forall \varepsilon > 0 \ \exists \delta > 0 \ \forall x \\ (0 < |x - a| < \delta \implies |f(x)| \leq \varepsilon |g(x)|).$$

Remark (Interpretation). The scale g is the unit of comparison. The assertion $f = o(g)$ says that no matter how small a tolerance multiple εg is chosen, f eventually fits inside that multiple.

Definition 1.2.2 (Little-o at Infinity). Let f and g be real-valued functions defined on some interval (M, ∞) , and suppose $g(x) \neq 0$ for all sufficiently large x . We write

$$f(x) = o(g(x)) \quad (x \rightarrow \infty)$$

if and only if for every $\varepsilon > 0$ there exists $R > 0$ such that

$$x > R \implies |f(x)| \leq \varepsilon |g(x)|.$$

Remark (Standard quantified statement).

$$f(x) = o(g(x)) \ (x \rightarrow \infty) \iff \forall \varepsilon > 0 \ \exists R > 0 \ \forall x \\ (x > R \implies |f(x)| \leq \varepsilon |g(x)|).$$

Remark (Interpretation). The point version and the infinity version have the same logical shape. Only the neighbourhood changes: small punctured intervals around a are replaced by tails (R, ∞) .

Definition 1.2.3 (Increment Little-o). Let r be a real-valued function defined for all sufficiently small nonzero h . We write

$$r(h) = o(h) \quad (h \rightarrow 0)$$

if and only if for every $\varepsilon > 0$ there exists $\delta > 0$ such that

$$0 < |h| < \delta \implies |r(h)| \leq \varepsilon |h|.$$

Remark (Standard quantified statement).

$$r(h) = o(h) \ (h \rightarrow 0) \iff \forall \varepsilon > 0 \ \exists \delta > 0 \ \forall h \ (0 < |h| < \delta \Rightarrow |r(h)| \leq \varepsilon |h|).$$

Remark (Interpretation). This is the form used in first-order approximation. A remainder $r(h)$ is smaller than the increment h itself, so the quotient $r(h)/h$ tends to zero.

Remark (Dependencies).

- [Little-o at a Point](#)

Proposition 1.2.4 (Little-o Quotient Characterisation). *Let $a \in \mathbb{R}$, and let f and g be real-valued functions defined on a punctured neighbourhood of a , with $g(x) \neq 0$ for x sufficiently close to a and $x \neq a$. Then*

$$f(x) = o(g(x)) \ (x \rightarrow a) \iff \frac{f(x)}{g(x)} \rightarrow 0 \ (x \rightarrow a).$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$[\forall \varepsilon > 0 \ \exists \delta > 0 \ 0 < |x - a| < \delta \Rightarrow |f(x)| \leq \varepsilon |g(x)|] \iff \frac{f(x)}{g(x)} \rightarrow 0.$$

Remark (Interpretation). The quotient form is often the fastest test for little-o. The epsilon form is often the safest proof form because it avoids manipulating a quotient until the nonvanishing of the scale has been checked.

Remark (Dependencies).

- [Little-o at a Point](#)

Proposition 1.2.5 (Sum Rule for Little-o). *If $f_1(x) = o(g(x))$ and $f_2(x) = o(g(x))$ as $x \rightarrow a$, then*

$$f_1(x) + f_2(x) = o(g(x)) \quad (x \rightarrow a).$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$f_1 = o(g) \wedge f_2 = o(g) \implies f_1 + f_2 = o(g).$$

Remark (Interpretation). Negligible errors can be added. The proof is the usual epsilon-budget split: make each error smaller than half the final tolerance.

Remark (Dependencies).

- [Little-o at a Point](#)

Proposition 1.2.6 (Bounded Factor Rule for Little-o). *If $f(x) = o(g(x))$ as $x \rightarrow a$ and m is bounded near a , then*

$$m(x)f(x) = o(g(x)) \quad (x \rightarrow a).$$

Go to proof.

Remark (Standard quantified statement).

$$f = o(g) \wedge \exists M > 0 \exists \eta > 0 \forall x (0 < |x - a| < \eta \Rightarrow |m(x)| \leq M) \implies mf = o(g).$$

Remark (Interpretation). A bounded multiplier cannot turn a negligible error into a main-order term. It only changes the constant in the epsilon estimate.

Remark (Dependencies).

- [Little-o at a Point](#)

1.3 Inequality

Inequality — Quick Reference

Concept/Statement	Meaning/Role
Add both sides	Adding the same term preserves strict inequalities.
Nonstrict add both sides	Addition also preserves nonstrict inequalities.
Add inequalities	Pairwise strict inequalities add.
Mixed addition	Strict and nonstrict hypotheses combine predictably.
Multiply by positive	Positive factors preserve order.
Multiply by negative	Negative factors reverse order.
Squeeze theorem	Double bounding by one value forces equality.
Reciprocal order	Reciprocals reverse positive inequalities.
Square monotonicity	Squaring preserves order on nonnegative inputs.
AM–GM	Arithmetic mean dominates geometric mean.
Bernoulli inequality	Powers of $1 + x$ dominate the tangent line.

Inequality

Theorem 1.3.1 (Addition Preserves Strict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a < b \Rightarrow a + c < b + c.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b, c \in \mathbb{R}, \quad a < b \Rightarrow a + c < b + c.$$

Remark (Dependencies).

[Ordered field](#), [ordered field axioms](#), [additive cancellation](#).

Theorem 1.3.2 (Addition Preserves Nonstrict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a \leq b \Rightarrow a + c \leq b + c.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b, c \in \mathbb{R}, \quad a \leq b \Rightarrow a + c \leq b + c.$$

Remark (Dependencies).

[Ordered field](#), [addition preserves strict inequality](#).

Theorem 1.3.3 (Addition of Strict Inequalities). *For all $a, b, c, d \in \mathbb{R}$,*

$$a < b \wedge c < d \Rightarrow a + c < b + d.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b, c, d \in \mathbb{R}, \quad a < b \wedge c < d \Rightarrow a + c < b + d.$$

Remark (Dependencies).

[addition preserves strict inequality](#), [transitivity of strict inequality](#).

Theorem 1.3.4 (Addition of Nonstrict Inequalities). *For all $a, b, c, d \in \mathbb{R}$,*

$$a \leq b \wedge c \leq d \Rightarrow a + c \leq b + d.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b, c, d \in \mathbb{R}, \quad a \leq b \wedge c \leq d \Rightarrow a + c \leq b + d.$$

Remark (Dependencies).

[addition preserves nonstrict inequality](#).

Theorem 1.3.5 (Mixed Addition of Inequalities). *For all $a, b, c, d \in \mathbb{R}$,*

$$a \leq b \wedge c < d \Rightarrow a + c < b + d.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b, c, d \in \mathbb{R}, \quad a \leq b \wedge c < d \Rightarrow a + c < b + d.$$

Remark (Dependencies).

[addition preserves nonstrict inequality](#), [addition preserves strict inequality](#), [mixed transitivity](#).

Theorem 1.3.6 (Multiplication by a Positive Scalar Preserves Strict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a < b \wedge 0 < c \Rightarrow ac < bc.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b, c \in \mathbb{R}, \quad a < b \wedge 0 < c \Rightarrow ac < bc.$$

Remark (Dependencies).

[Ordered field](#), multiply positive, [positive cone](#).

Theorem 1.3.7 (Multiplication by a Negative Scalar Reverses Strict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a < b \wedge c < 0 \Rightarrow ac > bc.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b, c \in \mathbb{R}, \quad a < b \wedge c < 0 \Rightarrow ac > bc.$$

Remark (Dependencies).

[Ordered field](#), multiply negative, [positive cone](#).

Theorem 1.3.8 (Multiplication by a Positive Scalar Preserves Nonstrict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a \leq b \wedge 0 < c \Rightarrow ac \leq bc.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b, c \in \mathbb{R}, \quad a \leq b \wedge 0 < c \Rightarrow ac \leq bc.$$

Remark (Dependencies).

[multiplication by positive preserves strict inequality](#).

Theorem 1.3.9 (Multiplication by a Nonneg Scalar Preserves Nonstrict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a \leq b \wedge 0 \leq c \Rightarrow ac \leq bc.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b, c \in \mathbb{R}, \quad a \leq b \wedge 0 \leq c \Rightarrow ac \leq bc.$$

Remark (Interpretation). This extends [the positive-scalar result](#) to include $c = 0$, where both sides become 0 and the inequality holds trivially.

Remark (Dependencies).

[multiplication by positive preserves nonstrict inequality](#), [ordered field](#).

Theorem (Squeeze)

Theorem 1.3.10 (Squeeze). *For all $a, b, c \in \mathbb{R}$,*

$$a \leq b \leq c \wedge a = c \Rightarrow b = a.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a, b, c \in \mathbb{R}, \quad a \leq b \leq c \wedge a = c \Rightarrow b = a.$$

Remark (Interpretation). The squeeze principle forces equality when a value is simultaneously bounded above and below by equal quantities. The continuous analogue (if $f(x) \leq g(x) \leq h(x)$ and $f(x) \rightarrow L$ and $h(x) \rightarrow L$ then $g(x) \rightarrow L$) is the central tool for computing limits. The algebraic version stated here is its foundation.

Remark (Dependencies).

[Ordered field](#), [additive cancellation](#).

Theorem 1.3.11 (Transitivity of Strict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a < b \wedge b < c \Rightarrow a < c.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a, b, c \in \mathbb{R}, \quad a < b \wedge b < c \Rightarrow a < c.$$

Remark (Dependencies).

[Ordered field](#), [positive cone](#).

Theorem 1.3.12 (Mixed Transitivity). *For all $a, b, c \in \mathbb{R}$,*

$$a \leq b \wedge b < c \Rightarrow a < c.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a, b, c \in \mathbb{R}, \quad a \leq b \wedge b < c \Rightarrow a < c.$$

Remark (Dependencies).

[Ordered field](#), [transitivity of strict inequality](#).

Theorem 1.3.13 (Reciprocal Reverses Strict Inequality for Positives). *For all $a, b \in \mathbb{R}$,*

$$0 < a < b \Rightarrow 0 < \frac{1}{b} < \frac{1}{a}.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b \in \mathbb{R}, \quad 0 < a < b \Rightarrow 0 < \frac{1}{b} < \frac{1}{a}.$$

Remark (Dependencies).

[Ordered field](#), [inverse positive](#), [reciprocal order for positive elements](#), [multiplication by positive preserves strict inequality](#).

Theorem 1.3.14 (Reciprocal Equivalence for Positives). *For all $a, b \in \mathbb{R}$ with $a > 0$ and $b > 0$,*

$$a < b \iff \frac{1}{b} < \frac{1}{a}.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b \in \mathbb{R}, \quad a > 0 \wedge b > 0 \Rightarrow (a < b \iff \frac{1}{b} < \frac{1}{a}).$$

Remark (Dependencies).

[reciprocal reverses strict inequality for positives](#), [reciprocal order for positive elements](#).

Theorem 1.3.15 (Square Is Strictly Monotone on Nonneg Reals). *For all $a, b \in \mathbb{R}$,*

$$0 \leq a < b \Rightarrow a^2 < b^2.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b \in \mathbb{R}, \quad 0 \leq a < b \Rightarrow a^2 < b^2.$$

Remark (Dependencies).

[Ordered field](#), [addition of strict inequalities](#), [multiplication by positive preserves strict inequality](#), [squares are nonnegative](#).

Theorem 1.3.16 (Square Root Is Monotone on Nonneg Reals). *For all $a, b \in \mathbb{R}$,*

$$0 \leq a \leq b \Rightarrow \sqrt{a} \leq \sqrt{b}.$$

Remark (Proof). [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b \in \mathbb{R}, \quad 0 \leq a \leq b \Rightarrow \sqrt{a} \leq \sqrt{b}.$$

Remark (Audit note). This theorem depends on `def:square-root`, which does not yet appear in the repository label inventory. That definition must be created and registered before this result can be fully formalised.

Remark (Dependencies).

`def:square-root` (not yet registered — see audit note above), [square is strictly monotone on nonneg reals](#).

Theorem 1.3.17 (AM-GM Inequality for Two Terms). *For all $a, b \geq 0$,*

$$\sqrt{ab} \leq \frac{a+b}{2}.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a, b \in \mathbb{R}, a \geq 0 \wedge b \geq 0 \Rightarrow \sqrt{ab} \leq \frac{a+b}{2}.$$

Remark (Interpretation). The arithmetic–geometric mean inequality for two terms is the simplest case of the AM-GM family. The standard proof proceeds by squaring both sides (using [square monotonicity](#)) and reducing to $(\sqrt{a} - \sqrt{b})^2 \geq 0$, which follows from [squares being nonnegative](#).

Remark (Dependencies).

[squares are nonnegative](#), [square is strictly monotone on nonneg reals](#), `def:square-root` (not yet registered — see audit note in [square root monotonicity](#)).

Theorem (Bernoulli's Inequality)

Theorem 1.3.18 (Bernoulli's Inequality). *For all $x \in \mathbb{R}$ with $x \geq -1$ and all $n \in \mathbb{N}$,*

$$(1+x)^n \geq 1+nx.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall x \in \mathbb{R}, \forall n \in \mathbb{N}, x \geq -1 \Rightarrow (1+x)^n \geq 1+nx.$$

Remark (Interpretation). Bernoulli's inequality is proved by induction and is a key estimate in analysis: it provides a linear lower bound for exponential growth and is used to establish the divergence of the harmonic series, bounds on compound interest, and properties of the exponential function.

Remark (Dependencies).

[Ordered field](#), [multiplication by nonneg preserves nonstrict inequality](#), [addition preserves nonstrict inequality](#).

1.4 Modulus

Modulus — Quick Reference

Concept/Statement	Meaning/Role
Absolute value	Magnitude stripped of sign.
Nonnegativity	Modulus is always nonnegative.
Zero characterisation	Only 0 has modulus 0.
Self or negation	$ a $ is either a or $-a$.
Symmetry	Negation does not change magnitude.
Product rule	Modulus is multiplicative.
Quotient rule	Division respects modulus when the denominator is nonzero.
Bounds by modulus	a lies between $- a $ and $ a $.
Nonstrict modulus inequality	$ a \leq r$ is equivalent to a two-sided bound.
Strict modulus inequality	$ a < r$ is equivalent to a strict two-sided bound.
Reverse triangle inequality	Magnitudes vary no faster than distance.
Finite triangle inequality	The modulus of a finite sum is bounded by summed moduli.

Modulus

Definition (Absolute Value)

Definition 1.4.1 (Absolute Value). Let $a \in \mathbb{R}$. The *absolute value* (or *modulus*) of a is

$$|a| := \begin{cases} a & \text{if } a \geq 0, \\ -a & \text{if } a < 0. \end{cases}$$

Equivalently, $|a| := \max(a, -a)$.

Remark (Standard quantified statement).

$$|a| = \max(a, -a) := \begin{cases} a & \text{if } a \geq 0, \\ -a & \text{if } a < 0. \end{cases}$$

Remark (Interpretation). The absolute value measures magnitude independent of sign. On the real line it equals the distance from a to 0: $|a| = d(a, 0)$ under the standard metric. The two-branch definition partitions $\mathbb{R}$ at 0 and reflects negative values to positive ones.

Remark (Dependencies).

[Ordered field](#), $\mathbb{R}$ is an [ordered field](#).

Theorem 1.4.2 (Absolute Value Is Nonnegative). For all $a \in \mathbb{R}$,

$$|a| \geq 0.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a \in \mathbb{R}, \quad |a| \geq 0.$$

Remark (Dependencies).

[Absolute value](#), [squares are nonnegative](#), [ordered field](#).

Theorem 1.4.3 (Absolute Value Zero Iff Zero). *For all $a \in \mathbb{R}$,*

$$|a| = 0 \iff a = 0.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a \in \mathbb{R}, \quad |a| = 0 \iff a = 0.$$

Remark (Dependencies).

[Absolute value](#), [absolute value is nonnegative](#), [ordered field](#).

Theorem 1.4.4 (Absolute Value Equals Self or Negation). *For all $a \in \mathbb{R}$, either $|a| = a$ or $|a| = -a$.*

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a \in \mathbb{R}, \quad |a| = a \vee |a| = -a.$$

Remark (Dependencies).

[Absolute value](#).

Theorem 1.4.5 (Absolute Value Is Symmetric). *For all $a \in \mathbb{R}$,*

$$|-a| = |a|.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a \in \mathbb{R}, \quad |-a| = |a|.$$

Remark (Dependencies).

[Absolute value](#), [double negation](#).

Theorem 1.4.6 (Absolute Value of a Product). *For all $a, b \in \mathbb{R}$,*

$$|ab| = |a| |b|.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a, b \in \mathbb{R}, \quad |ab| = |a| |b|.$$

Remark (Dependencies).

[Absolute value](#), negation of product, product of negatives, [squares are nonnegative](#).

Theorem 1.4.7 (Absolute Value of a Quotient). *For all $a \in \mathbb{R}$ and $b \in \mathbb{R}$ with $b \neq 0$,*

$$\left| \frac{a}{b} \right| = \frac{|a|}{|b|}.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a \in \mathbb{R}, \forall b \in \mathbb{R} \setminus \{0\}, \quad \left| \frac{a}{b} \right| = \frac{|a|}{|b|}.$$

Remark (Dependencies).

[Absolute value](#), [absolute value of a product](#), [absolute value zero iff zero](#), [uniqueness of multiplicative inverse](#).

Theorem 1.4.8 (Bound by Modulus). *For all $a \in \mathbb{R}$,*

$$-|a| \leq a \leq |a|.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a \in \mathbb{R}, \quad -|a| \leq a \leq |a|.$$

Remark (Dependencies).

[Absolute value](#), [absolute value is nonnegative](#), [ordered field](#).

Theorem (Nonstrict Modulus Inequality)

Theorem 1.4.9 (Nonstrict Modulus Inequality). *For all $a \in \mathbb{R}$ and $r \geq 0$,*

$$|a| \leq r \iff -r \leq a \leq r.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a \in \mathbb{R}, \forall r \in \mathbb{R}, r \geq 0 \Rightarrow (|a| \leq r \iff -r \leq a \leq r).$$

Remark (Interpretation). This biconditional is the primary tool for replacing a modulus inequality with a pair of simultaneous linear inequalities. It underlies the ε - δ manipulations throughout analysis: the condition $|x - a| \leq \varepsilon$ is equivalent to $a - \varepsilon \leq x \leq a + \varepsilon$.

Remark (Dependencies).

[Absolute value](#), [bound by modulus](#), [ordered field](#).

Theorem 1.4.10 (Strict Modulus Inequality). *For all $a \in \mathbb{R}$ and $r > 0$,*

$$|a| < r \iff -r < a < r.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a \in \mathbb{R}, \forall r \in \mathbb{R}, r > 0 \Rightarrow (|a| < r \iff -r < a < r).$$

Remark (Dependencies).

[Absolute value](#), [nonstrict modulus inequality](#), [ordered field](#).

Theorem (Triangle Inequality)

Theorem (Triangle Inequality). *For all $a, b \in \mathbb{R}$,*

$$|a + b| \leq |a| + |b|.$$

Remark (Standard quantified statement).

$$\forall a, b \in \mathbb{R}, \quad |a + b| \leq |a| + |b|.$$

Remark (Interpretation). The triangle inequality is the foundational estimate of analysis. Every convergence argument, continuity proof, and metric-space result that requires bounding a sum reduces, at some step, to this inequality. The name reflects its geometric meaning: the length of one side of a triangle does not exceed the sum of the other two.

Theorem 1.4.11 (Reverse Triangle Inequality). *For all $a, b \in \mathbb{R}$,*

$$||a| - |b|| \leq |a - b|.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall a, b \in \mathbb{R}, \quad ||a| - |b|| \leq |a - b|.$$

Remark (Interpretation). The reverse triangle inequality provides a lower bound on $|a - b|$ in terms of the difference of moduli. It is used to bound the modulus of a difference from below, particularly when proving that a function is not uniformly continuous or that a sequence does not converge.

Remark (Dependencies).

[Absolute value](#), [triangle inequality](#), [absolute value is symmetric](#).

Theorem 1.4.12 (Generalised Triangle Inequality). *For all $n \in \mathbb{N}$ and $a_1, \dots, a_n \in \mathbb{R}$,*

$$\left| \sum_{k=1}^n a_k \right| \leq \sum_{k=1}^n |a_k|.$$

Remark (Proof). [Go to proof](#).

Remark (Standard quantified statement).

$$\forall n \in \mathbb{N}, \forall a_1, \dots, a_n \in \mathbb{R}, \quad \left| \sum_{k=1}^n a_k \right| \leq \sum_{k=1}^n |a_k|.$$

Remark (Dependencies).

[Absolute value](#), [triangle inequality](#).

Proofs

Remark (Return). [Return to Theorem](#)

Proposition (Order arithmetic). *Let $a, b, c, d \in \mathbb{R}$.*

- (i) If $a \leq b$ and $c \leq d$, then $a + c \leq b + d$.*
- (ii) If $a \leq b$ and $c > 0$, then $ac \leq bc$.*
- (iii) If $a \leq b$ and $c < 0$, then $ac \geq bc$.*
- (iv) $|x| \leq y$ (with $y \geq 0$) if and only if $-y \leq x \leq y$.*

Professional Standard Proof.

TODO: professional standard proof for order-arithmetic. □

Detailed Learning Proof.

TODO: detailed learning proof for order-arithmetic. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Convergent sequences are bounded). *If $(b_n) \rightarrow b$, then there exists $M > 0$ such that $|b_n| \leq M$ for all $n \in \mathbb{N}$.*

Professional Standard Proof.

TODO: professional standard proof for convergent-bounded. □

Detailed Learning Proof.

TODO: detailed learning proof for convergent-bounded. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (-characterisation of). *Let $S \subset \mathbb{R}$ be nonempty and bounded above, and let $s = \sup S$. Then for every $\varepsilon > 0$ there exists $x \in S$ with $x > s - \varepsilon$.*

Professional Standard Proof.

TODO: professional standard proof for eps-char-sup. □

Detailed Learning Proof.

TODO: detailed learning proof for eps-char-sup. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Monotone Approximation of Bounds). *Let $S \subset \mathbb{R}$ be nonempty and bounded. Then there exist monotone sequences $(x_n), (y_n) \subset S$ with $x_n \rightarrow \sup S$ and $y_n \rightarrow \inf S$.*

Professional Standard Proof.

TODO: professional standard proof for monotone-approx-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for monotone-approx-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (The dichotomy). *For any predicate P on $\mathbb{N}$, exactly one of the following holds:*

(i) *P holds infinitely often.*

(ii) *$\neg P$ holds eventually.*

Professional Standard Proof.

TODO: professional standard proof for io-ev-dichotomy. □

Detailed Learning Proof.

TODO: detailed learning proof for io-ev-dichotomy. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Bolzano–Weierstrass via bisection). *Every bounded sequence in $\mathbb{R}$ has a convergent subsequence.*

Professional Standard Proof.

TODO: professional standard proof for bw-bisection. □

Detailed Learning Proof.

TODO: detailed learning proof for bw-bisection. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Intermediate Value Theorem). *Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. If $L \in \mathbb{R}$ satisfies $f(a) < L < f(b)$ or $f(a) > L > f(b)$, then there exists $c \in (a, b)$ with $f(c) = L$.*

Professional Standard Proof.

TODO: professional standard proof for ivt-bisection. □

Detailed Learning Proof.

TODO: detailed learning proof for ivt-bisection. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (r-uncountable). $\mathbb{R}$ is uncountable.

Professional Standard Proof.

TODO: professional standard proof for r-uncountable. □

Detailed Learning Proof.

TODO: detailed learning proof for r-uncountable. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Residue divergence criterion). *If there exist residues $r, s \in \{0, \dots, k-1\}$ such that $(a_{kn+r}) \rightarrow L$ and $(a_{kn+s}) \rightarrow M$ with $L \neq M$, then (a_n) diverges.*

Professional Standard Proof.

TODO: professional standard proof for residue-divergence. □

Detailed Learning Proof.

TODO: detailed learning proof for residue-divergence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Little-o Quotient Characterisation). *Let $a \in \mathbb{R}$, and let f and g be real-valued functions defined on a punctured neighbourhood of a , with $g(x) \neq 0$ for x sufficiently close to a and $x \neq a$. Then*

$$f(x) = o(g(x)) \ (x \rightarrow a) \iff \frac{f(x)}{g(x)} \rightarrow 0 \ (x \rightarrow a).$$

Professional Standard Proof.

By definition, $f = o(g)$ as $x \rightarrow a$ means that for every $\varepsilon > 0$, eventually $|f(x)| \leq \varepsilon|g(x)|$. Since $g(x) \neq 0$ eventually, this is equivalent to

$$\left| \frac{f(x)}{g(x)} \right| \leq \varepsilon$$

eventually. That is precisely the assertion that $f(x)/g(x) \rightarrow 0$ as $x \rightarrow a$. $\square$

Detailed Learning Proof.

Assume first that $f = o(g)$. Given $\varepsilon > 0$, choose $\delta > 0$ so that $0 < |x - a| < \delta$ implies $|f(x)| \leq \varepsilon|g(x)|$. On a smaller punctured neighbourhood, if necessary, $g(x) \neq 0$. Dividing by $|g(x)|$ gives $|f(x)/g(x)| \leq \varepsilon$, so $f(x)/g(x) \rightarrow 0$.

Conversely, assume $f(x)/g(x) \rightarrow 0$. Given $\varepsilon > 0$, choose $\delta > 0$ so that $0 < |x - a| < \delta$ implies $|f(x)/g(x)| \leq \varepsilon$, again within a punctured neighbourhood where g is nonzero. Multiplying by $|g(x)|$ gives $|f(x)| \leq \varepsilon|g(x)|$, hence $f = o(g)$. $\square$

Remark (Dependencies).

- [Little-o at a Point](#)

Remark (Return). [Return to Theorem](#)

Proposition (Sum Rule for Little-o). *If $f_1(x) = o(g(x))$ and $f_2(x) = o(g(x))$ as $x \rightarrow a$, then*

$$f_1(x) + f_2(x) = o(g(x)) \quad (x \rightarrow a).$$

Professional Standard Proof.

Let $\varepsilon > 0$. Since $f_1 = o(g)$ and $f_2 = o(g)$, each term is eventually bounded by $(\varepsilon/2)|g(x)|$ in absolute value. On the intersection of the two punctured neighbourhoods,

$$|f_1(x) + f_2(x)| \leq |f_1(x)| + |f_2(x)| \leq \varepsilon|g(x)|.$$

Therefore $f_1 + f_2 = o(g)$. □

Detailed Learning Proof.

Choose a target tolerance $\varepsilon > 0$. The first little-o hypothesis gives a punctured neighbourhood on which $|f_1(x)| \leq (\varepsilon/2)|g(x)|$. The second gives a punctured neighbourhood on which $|f_2(x)| \leq (\varepsilon/2)|g(x)|$. Taking the smaller radius makes both estimates hold simultaneously. The triangle inequality then gives the desired bound for the sum. □

Remark (Dependencies).

- [Little-o at a Point](#)

Remark (Return). [Return to Theorem](#)

Proposition (Bounded Factor Rule for Little-o). *If $f(x) = o(g(x))$ as $x \rightarrow a$ and m is bounded near a , then*

$$m(x)f(x) = o(g(x)) \quad (x \rightarrow a).$$

Professional Standard Proof.

Let $\varepsilon > 0$. Since m is bounded near a , choose $M > 0$ and a punctured neighbourhood on which $|m(x)| \leq M$. Since $f = o(g)$, choose a smaller punctured neighbourhood on which $|f(x)| \leq (\varepsilon/M)|g(x)|$. Then

$$|m(x)f(x)| \leq M|f(x)| \leq \varepsilon|g(x)|.$$

Thus $mf = o(g)$. □

Detailed Learning Proof.

The bounded factor can cost only a fixed multiplicative constant. If $|m(x)| \leq M$ near a , ask the little-o estimate for f to be ε/M instead of ε . Multiplying the two estimates gives the final tolerance ε . □

Remark (Dependencies).

- [Little-o at a Point](#)

Remark (Return). [Return to Theorem](#)

Theorem (Addition Preserves Strict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a < b \Rightarrow a + c < b + c.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-add-both-sides. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-add-both-sides. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Addition Preserves Nonstrict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a \leq b \Rightarrow a + c \leq b + c.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-nonstrict-add-both-sides. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-nonstrict-add-both-sides. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Addition of Strict Inequalities). *For all $a, b, c, d \in \mathbb{R}$,*

$$a < b \wedge c < d \Rightarrow a + c < b + d.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-add-inequalities. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-add-inequalities. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Addition of Nonstrict Inequalities). *For all $a, b, c, d \in \mathbb{R}$,*

$$a \leq b \wedge c \leq d \Rightarrow a + c \leq b + d.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-nonstrict-add-inequalities. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-nonstrict-add-inequalities. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Mixed Addition of Inequalities). *For all $a, b, c, d \in \mathbb{R}$,*

$$a \leq b \wedge c < d \Rightarrow a + c < b + d.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-mixed-add. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-mixed-add. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Multiplication by a Positive Scalar Preserves Strict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a < b \wedge 0 < c \Rightarrow ac < bc.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-multiply-positive. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-multiply-positive. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Multiplication by a Negative Scalar Reverses Strict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a < b \wedge c < 0 \Rightarrow ac > bc.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-multiply-negative. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-multiply-negative. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Multiplication by a Positive Scalar Preserves Nonstrict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a \leq b \wedge 0 < c \Rightarrow ac \leq bc.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-nonstrict-multiply-positive. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-nonstrict-multiply-positive. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Multiplication by a Nonneg Scalar Preserves Nonstrict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a \leq b \wedge 0 \leq c \Rightarrow ac \leq bc.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-nonstrict-multiply-nonneg. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-nonstrict-multiply-nonneg. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Squeeze). *For all $a, b, c \in \mathbb{R}$,*

$$a \leq b \leq c \wedge a = c \Rightarrow b = a.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-squeeze. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-squeeze. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Transitivity of Strict Inequality). *For all $a, b, c \in \mathbb{R}$,*

$$a < b \wedge b < c \Rightarrow a < c.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-transitivity-strict. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-transitivity-strict. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Mixed Transitivity). *For all $a, b, c \in \mathbb{R}$,*

$$a \leq b \wedge b < c \Rightarrow a < c.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-transitivity-mixed. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-transitivity-mixed. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Reciprocal Reverses Strict Inequality for Positives). *For all $a, b \in \mathbb{R}$,*

$$0 < a < b \Rightarrow 0 < \frac{1}{b} < \frac{1}{a}.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-reciprocal-positive. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-reciprocal-positive. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Reciprocal Equivalence for Positives). *For all $a, b \in \mathbb{R}$ with $a > 0$ and $b > 0$,*

$$a < b \iff \frac{1}{b} < \frac{1}{a}.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-reciprocal-flip. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-reciprocal-flip. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Square Is Strictly Monotone on Nonneg Reals). *For all $a, b \in \mathbb{R}$,*

$$0 \leq a < b \Rightarrow a^2 < b^2.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-square-monotone. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-square-monotone. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Square Root Is Monotone on Nonneg Reals). *For all $a, b \in \mathbb{R}$,*

$$0 \leq a \leq b \Rightarrow \sqrt{a} \leq \sqrt{b}.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-square-root-monotone. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-square-root-monotone. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (AM-GM Inequality for Two Terms). *For all $a, b \geq 0$,*

$$\sqrt{ab} \leq \frac{a+b}{2}.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-am-gm-two. □

Detailed Learning Proof.

TODO: detailed learning proof for ineq-am-gm-two. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Bernoulli's Inequality). *For all $x \in \mathbb{R}$ with $x \geq -1$ and all $n \in \mathbb{N}$,*

$$(1 + x)^n \geq 1 + nx.$$

Professional Standard Proof.

TODO: professional standard proof for ineq-bernoulli.

□

Detailed Learning Proof.

TODO: detailed learning proof for ineq-bernoulli.

□

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Absolute Value Is Nonnegative). *For all $a \in \mathbb{R}$,*

$$|a| \geq 0.$$

Professional Standard Proof.

TODO: professional standard proof for absolute-value-nonneg. □

Detailed Learning Proof.

TODO: detailed learning proof for absolute-value-nonneg. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Absolute Value Zero Iff Zero). *For all $a \in \mathbb{R}$,*

$$|a| = 0 \iff a = 0.$$

Professional Standard Proof.

TODO: professional standard proof for absolute-value-zero-iff-zero. □

Detailed Learning Proof.

TODO: detailed learning proof for absolute-value-zero-iff-zero. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Absolute Value Equals Self or Negation). *For all $a \in \mathbb{R}$, either $|a| = a$ or $|a| = -a$.*

Professional Standard Proof.

TODO: professional standard proof for absolute-value-self-or-neg. □

Detailed Learning Proof.

TODO: detailed learning proof for absolute-value-self-or-neg. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Absolute Value Is Symmetric). *For all $a \in \mathbb{R}$,*

$$|-a| = |a|.$$

Professional Standard Proof.

TODO: professional standard proof for absolute-value-symmetric. □

Detailed Learning Proof.

TODO: detailed learning proof for absolute-value-symmetric. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Absolute Value of a Product). *For all $a, b \in \mathbb{R}$,*

$$|ab| = |a| |b|.$$

Professional Standard Proof.

TODO: professional standard proof for absolute-value-product. □

Detailed Learning Proof.

TODO: detailed learning proof for absolute-value-product. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Absolute Value of a Quotient). *For all $a \in \mathbb{R}$ and $b \in \mathbb{R}$ with $b \neq 0$,*

$$\left| \frac{a}{b} \right| = \frac{|a|}{|b|}.$$

Professional Standard Proof.

TODO: professional standard proof for absolute-value-quotient.

□

Detailed Learning Proof.

TODO: detailed learning proof for absolute-value-quotient.

□

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Bound by Modulus). *For all $a \in \mathbb{R}$,*

$$-|a| \leq a \leq |a|.$$

Professional Standard Proof.

TODO: professional standard proof for absolute-value-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for absolute-value-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Nonstrict Modulus Inequality). *For all $a \in \mathbb{R}$ and $r \geq 0$,*

$$|a| \leq r \iff -r \leq a \leq r.$$

Professional Standard Proof.

TODO: professional standard proof for absolute-value-le-iff. □

Detailed Learning Proof.

TODO: detailed learning proof for absolute-value-le-iff. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Strict Modulus Inequality). *For all $a \in \mathbb{R}$ and $r > 0$,*

$$|a| < r \iff -r < a < r.$$

Professional Standard Proof.

TODO: professional standard proof for absolute-value-lt-iff. □

Detailed Learning Proof.

TODO: detailed learning proof for absolute-value-lt-iff. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Reverse Triangle Inequality). *For all $a, b \in \mathbb{R}$,*

$$||a| - |b|| \leq |a - b|.$$

Professional Standard Proof.

TODO: professional standard proof for reverse-triangle-inequality. □

Detailed Learning Proof.

TODO: detailed learning proof for reverse-triangle-inequality. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Generalised Triangle Inequality). *For all $n \in \mathbb{N}$ and $a_1, \dots, a_n \in \mathbb{R}$,*

$$\left| \sum_{k=1}^n a_k \right| \leq \sum_{k=1}^n |a_k|.$$

Professional Standard Proof.

TODO: professional standard proof for absolute-value-sum-bound. □

Detailed Learning Proof.

TODO: detailed learning proof for absolute-value-sum-bound. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Chapter 2

Structure of the Real Line



2.1 Real Line One Dimensional

The Line as Order Plus Distance

The real numbers carry two compatible structures at once: a linear order $<$ and a notion of distance $d(x, y) = |x - y|$. Together these make $\mathbb{R}$ a one-dimensional space — a line. This chapter develops only the metric topology of $\mathbb{R}$, not topology in general spaces. Every open set is built from intervals, and compactness is introduced only on $\mathbb{R}$, with Heine–Borel as the capstone.

The intuition is geometric: points are positions on a line, order is “left of,” and distance is the length of the segment between two points. The caution to keep throughout is that $\mathbb{R}$ is special. Order and metric agree here in a way that fails on a general space, so definitions are stated in the form that survives to metric spaces and manifolds, with $\mathbb{R}$ as the first instance rather than the only one.

2.2 Order Distance Length

Distance on the Line

The absolute value of a difference turns $\mathbb{R}$ into a metric space. Length of an interval is the distance between its endpoints. These are the quantitative tools underneath every later definition of nearness.

Definition (Distance on the Real Line)

Definition 2.2.1 (Distance on the Real Line). The *distance* on $\mathbb{R}$ is the map $d : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ given by

$$d(x, y) = |x - y|.$$

Remark (Standard quantified statement).

$$\forall x, y \in \mathbb{R} (d(x, y) = |x - y|).$$

Remark (Interpretation). The distance between two points is the length of the segment joining them. It is the metric that makes “close” precise and is the basis for balls, neighborhoods, and open sets.

Remark (Dependencies).

- [Triangle Inequality](#)

Definition (Length of a Bounded Interval)

Definition 2.2.2 (Length of a Bounded Interval). Let $a, b \in \mathbb{R}$ with $a \leq b$. The *length* of any bounded interval with endpoints a and b is

$$\ell = b - a.$$

Remark (Standard quantified statement).

$$\forall a, b \in \mathbb{R} (a \leq b \implies \ell([a, b]) = b - a = d(a, b)).$$

Remark (Interpretation). Length depends only on the endpoints, not on whether they are included. It is the distance $d(a, b)$ and measures the size of the basic pieces of the line.

Remark (Dependencies).

- [Distance on the Real Line](#)

Theorem (The Distance Function Is a Metric)

Theorem 2.2.3 (The Distance Function Is a Metric). The map $d(x, y) = |x - y|$ satisfies, for all $x, y, z \in \mathbb{R}$: (i) $d(x, y) \geq 0$ with $d(x, y) = 0$ iff $x = y$; (ii) $d(x, y) = d(y, x)$; and (iii) $d(x, z) \leq d(x, y) + d(y, z)$. Hence $(\mathbb{R}, d)$ is a metric space. [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall x, y, z \in \mathbb{R} (d(x, y) \geq 0 \wedge (d(x, y) = 0 \iff x = y) \wedge d(x, y) = d(y, x) \wedge d(x, z) \leq d(x, y) + d(y, z)). \blacksquare$$

Remark (Interpretation). The three metric axioms are exactly the properties of $|\cdot|$: nonnegativity with the zero-distance characterization, symmetry, and the triangle inequality. This is the precise sense in which $\mathbb{R}$ is a one-dimensional metric space.

Remark (Dependencies).

- [Distance on the Real Line](#)
- [Triangle Inequality](#)

2.3 Absolute Value as Distance

Absolute Value Is Distance to Zero

The metric d specializes to absolute value when one point is the origin. This is the geometric reading of $|a|$: how far a sits from 0 on the line.

Proposition (Absolute Value Is Distance to the Origin)

Proposition 2.3.1 (Absolute Value Is Distance to the Origin). *For every $a \in \mathbb{R}$, $|a| = d(a, 0)$. [Go to proof.](#)*

Remark (Standard quantified statement).

$$\forall a \in \mathbb{R} (|a| = d(a, 0)).$$

Remark (Interpretation). Absolute value is not a separate idea from distance; it is the distance to the origin. The two-sided bound $|x| \leq r \Leftrightarrow -r \leq x \leq r$ is then exactly the statement that x lies within distance r of 0, i.e. in the interval $[-r, r]$.

Remark (Dependencies).

- [Distance on the Real Line](#)

2.4 Intervals as Subsets

Intervals as the Basic Pieces

The interval shapes were introduced in §12.2. Here they are recast as subsets of the metric line and supplemented with boundedness, which is the size condition compactness will need.

Definition (Bounded Subset of the Real Line)

Definition 2.4.1 (Bounded Subset of the Real Line). A set $E \subseteq \mathbb{R}$ is *bounded* exactly when there exists $M \geq 0$ such that $|x| \leq M$ for all $x \in E$; equivalently, E is contained in some closed interval $[-M, M]$.

Remark (Standard quantified statement).

$$E \text{ is bounded} \iff \exists M \geq 0 \forall x \in E (|x| \leq M).$$

Remark (Negated quantified statement).

$$E \text{ is unbounded} \iff \forall M \geq 0 \exists x \in E (|x| > M).$$

Remark (Interpretation). Boundedness says E fits inside a single closed interval — it does not run off to infinity. Together with closedness it is one half of the Heine–Borel characterization of compact subsets of $\mathbb{R}$.

Remark (Dependencies).

- [Closed Interval](#)
- [Triangle Inequality](#)

Set Operations on Intervals

Intervals as Sets

Once intervals are regarded as subsets of $\mathbb{R}$, the usual set operations apply to them. This is the first place where intervals behave like the building blocks of topology rather than merely as pieces of notation on the number line. Intersections often remain intervals, while unions, complements, and differences may split into several interval pieces.

Definition (Union of Intervals)

Definition 2.4.2 (Union of Intervals). Let $I, J \subseteq \mathbb{R}$ be intervals. Their *union* is the set

$$I \cup J = \{x \in \mathbb{R} : x \in I \text{ or } x \in J\}.$$

Remark (Dependencies).

- [Open Interval](#)
- [Closed Interval](#)
- [Union](#)

Definition (Intersection of Intervals)

Definition 2.4.3 (Intersection of Intervals). Let $I, J \subseteq \mathbb{R}$ be intervals. Their *intersection* is the set

$$I \cap J = \{x \in \mathbb{R} : x \in I \text{ and } x \in J\}.$$

Remark (Dependencies).

- [Open Interval](#)
- [Closed Interval](#)
- [Intersection](#)

Definition (Complement of an Interval)

Definition 2.4.4 (Complement of an Interval). Let $I \subseteq \mathbb{R}$ be an interval. Its *complement* in $\mathbb{R}$ is the set

$$I^c = \mathbb{R} \setminus I = \{x \in \mathbb{R} : x \notin I\}.$$

Remark (Dependencies).

- [Open Interval](#)
- [Closed Interval](#)
- [Complement](#)

Definition (Difference of Intervals)

Definition 2.4.5 (Difference of Intervals). Let $I, J \subseteq \mathbb{R}$ be intervals. The *difference* of I by J is the set

$$I \setminus J = \{x \in \mathbb{R} : x \in I \text{ and } x \notin J\} = I \cap J^c.$$

Remark (Predicate readings). For any $x \in \mathbb{R}$,

$$x \in I \cup J \iff (x \in I) \vee (x \in J),$$

$$x \in I \cap J \iff (x \in I) \wedge (x \in J),$$

$$x \in I^c \iff x \notin I, \quad x \in I \setminus J \iff (x \in I) \wedge (x \notin J).$$

These are set-theoretic operations; no arithmetic operation is being performed on the endpoints.

Remark (Dependencies).

- [Open Interval](#)
- [Closed Interval](#)
- [Left-Half-Open Interval](#)
- [Right-Half-Open Interval](#)
- [Union of Intervals](#)
- [Intersection of Intervals](#)
- [Complement of an Interval](#)

Remark (Intersections of intervals). The intersection of two intervals is either empty or another interval. For example, if $a \leq b$ and $c \leq d$, then

$$[a, b] \cap [c, d] = \begin{cases} [\max\{a, c\}, \min\{b, d\}], & \max\{a, c\} \leq \min\{b, d\}, \\ \emptyset, & \max\{a, c\} > \min\{b, d\}. \end{cases}$$

Endpoint inclusion follows the endpoint rules of the original intervals.

Remark (Unions of intervals). The union of two intervals need not be an interval:

$$(0, 1) \cup (2, 3)$$

has a gap. If two intervals overlap or touch without a missing endpoint, their union is again an interval, for instance

$$(0, 2] \cup [2, 5) = (0, 5).$$

The missing or included endpoints determine whether the joined set is truly one interval.

Remark (Complements of intervals). The complement of a bounded interval usually splits into two rays:

$$(a, b)^c = (-\infty, a] \cup [b, \infty), \quad [a, b]^c = (-\infty, a) \cup (b, \infty).$$

Complements reverse endpoint inclusion: an endpoint excluded from the interval is included in the complement, and an endpoint included in the interval is excluded from the complement.

Remark (Differences of intervals). A difference is an intersection with a complement, so it can also split:

$$(0, 5) \setminus [2, 3] = (0, 2) \cup (3, 5).$$

If the removed interval only cuts off one side, the result may remain a single interval:

$$(0, 5) \setminus (3, \infty) = (0, 3].$$

The bracket at 3 appears because $3 \notin (3, \infty)$, so it is not removed.

Remark (Topological motivation). Open sets in $\mathbb{R}$ are built from unions of open intervals. Closed sets are understood through complements of open sets. Boundaries arise when a point is not cleanly inside a set or its complement. Thus interval set operations are the grammar behind the metric topology of the real line.

2.5 Neighborhoods and Balls

Balls Are Open Intervals

On the line, the ball of radius r about a point is an open interval centered at that point. Neighborhoods are sets that contain such a ball. These are the local objects from which open sets are assembled.

Definition (Open Ball on the Real Line)

Definition 2.5.1 (Open Ball on the Real Line). Let $x \in \mathbb{R}$ and $r > 0$. The *open ball* of radius r about x is

$$B_r(x) = \{y \in \mathbb{R} : |y - x| < r\} = (x - r, x + r).$$

Remark (Standard quantified statement).

$$B_r(x) = \{y \in \mathbb{R} : d(y, x) < r\} = (x - r, x + r).$$

Remark (Interpretation). A ball on the line is just a symmetric open interval: all points strictly within distance r of x . Writing balls as intervals keeps the chapter inside the metric topology of $\mathbb{R}$.

Remark (Dependencies).

- [Distance on the Real Line](#)

- [Open Interval](#)

Definition (Neighborhood of a Point)

Definition 2.5.2 (Neighborhood of a Point). A set $N \subseteq \mathbb{R}$ is a *neighborhood* of $x \in \mathbb{R}$ exactly when there exists $r > 0$ with $B_r(x) \subseteq N$.

Remark (Standard quantified statement).

$$N \text{ is a neighborhood of } x \iff \exists r > 0 (B_r(x) \subseteq N).$$

Remark (Interpretation). A neighborhood of x is any set with room to spare around x — it contains a whole ball, not just x itself. Neighborhoods are the language of “near x ” used to define interior points and limits.

Remark (Dependencies).

- [Open Ball on the Real Line](#)

2.6 Open Sets

Sets Built from Balls

An open set is one that contains a ball around each of its points. On the line these are exactly the unions of open intervals. The closure properties recorded here are the metric-topology core.

Definition (Open Set in $\mathbb{R}$)

Definition 2.6.1 (Open Set in $\mathbb{R}$). A set $U \subseteq \mathbb{R}$ is *open* exactly when

$$\forall x \in U \exists r > 0 (B_r(x) \subseteq U).$$

Remark (Standard quantified statement).

$$U \text{ is open} \iff \forall x \in U \exists r > 0 (B_r(x) \subseteq U).$$

Remark (Negated quantified statement).

$$U \text{ is not open} \iff \exists x \in U \forall r > 0 (B_r(x) \not\subseteq U).$$

Remark (Failure modes). Openness fails exactly when some point of U has no ball contained in U — a boundary point trapped inside U , every ball around which pokes outside.

Remark (Interpretation). An open set has interior room around each of its points: a small step in either direction remains inside the set. The empty set and $\mathbb{R}$ are open, and open intervals are the model examples.

Remark (Dependencies).

- [Open Ball on the Real Line](#)

Theorem (Open Intervals Are Open)

Theorem 2.6.2 (Open Intervals Are Open). *Every open interval (a, b) , every open ray (a, ∞) and $(-\infty, b)$, and $\mathbb{R}$ itself are open sets. [Go to proof.](#)*

Remark (Standard quantified statement).

$$\begin{aligned} &\forall a, b \in \mathbb{R} (a < b \implies (a, b) \text{ is open}), \\ &\forall a \in \mathbb{R} ((a, \infty) \text{ is open} \wedge (-\infty, a) \text{ is open}), \quad \mathbb{R} \text{ is open.} \end{aligned}$$

Remark (Interpretation). The name “open interval” is justified: each such interval satisfies the open-set condition, with the radius at x taken as the distance to the nearer endpoint.

Remark (Dependencies).

- [Open Set in \$\mathbb{R}\$](#)
- [Open Interval](#)

Theorem (Unions and Finite Intersections of Open Sets)

Theorem 2.6.3 (Unions and Finite Intersections of Open Sets). *An arbitrary union of open subsets of $\mathbb{R}$ is open, and a finite intersection of open subsets of $\mathbb{R}$ is open. [Go to proof.](#)*

Remark (Standard quantified statement).

$$\left(\forall i \in I \ U_i \text{ open} \right) \implies \left(\bigcup_{i \in I} U_i \right) \text{ open}; \quad \left(\bigwedge_{k=1}^n U_k \text{ open} \right) \implies \left(\bigcap_{k=1}^n U_k \right) \text{ open}.$$

Remark (Interpretation). Openness is preserved by arbitrary unions but only finite intersections. The infinite intersection $\bigcap_n (-1/n, 1/n) = \{0\}$ shows why “finite” cannot be dropped: a shrinking family of open intervals can close down onto a single point.

Remark (Dependencies).

- [Open Set in \$\mathbb{R}\$](#)

2.7 Closed Sets

Complements of Open Sets

Closed sets are the complements of open sets. They can equivalently be described as the sets that contain all of their limit points, the description used when reasoning about convergence.

Definition (Closed Set in $\mathbb{R}$)

Definition 2.7.1 (Closed Set in $\mathbb{R}$). A set $F \subseteq \mathbb{R}$ is *closed* exactly when its complement $\mathbb{R} \setminus F$ is open.

Remark (Standard quantified statement).

$$F \text{ is closed} \iff (\mathbb{R} \setminus F) \text{ is open.}$$

Remark (Interpretation). Closed and open are complementary, not opposite: a set can be both (e.g. $\emptyset, \mathbb{R}$) or neither (e.g. $[0, 1)$). Closed intervals $[a, b]$ are closed, as are finite sets and $\mathbb{R}$ itself.

Remark (Dependencies).

- [Open Set in \$\mathbb{R}\$](#)

Theorem (Closed Sets Contain Their Limit Points)

Theorem 2.7.2 (Closed Sets Contain Their Limit Points). A set $F \subseteq \mathbb{R}$ is closed if and only if it contains every one of its limit points. [Go to proof](#).

Remark (Standard quantified statement).

$$F \text{ closed} \iff \forall x \in \mathbb{R} (x \text{ is a limit point of } F \Rightarrow x \in F).$$

Remark (Interpretation). This is the convergence-flavored description of closedness: a closed set cannot be approached from outside without already containing the approached point. It is the form used when showing limits of sequences in F stay in F .

Remark (Dependencies).

- [Closed Set in \$\mathbb{R}\$](#)
- [Limit Point](#)

2.8 Interior Exterior Boundary

Inside, Outside, and Edge

Relative to a set E , each point of the line is interior to E , exterior to E , or on its boundary. These three regions partition $\mathbb{R}$ and refine the open/closed distinction.

Definition (Interior Point)

Definition 2.8.1 (Interior Point). Let $E \subseteq \mathbb{R}$ and $x \in \mathbb{R}$. Then x is an interior point of E exactly when

$$\exists r > 0 (B_r(x) \subseteq E).$$

Remark (Standard quantified statement).

$$x \text{ is an interior point of } E \iff \exists r > 0 (B_r(x) \subseteq E).$$

Remark (Interpretation). An interior point of E sits inside E with room to spare: a whole ball around it lies in E .

Remark (Dependencies).

- [Open Ball on the Real Line](#)

Definition (Exterior Point)

Definition 2.8.2 (Exterior Point). Let $E \subseteq \mathbb{R}$ and $x \in \mathbb{R}$. Then x is an exterior point of E exactly when

$$\exists r > 0 (B_r(x) \subseteq \mathbb{R} \setminus E).$$

Remark (Standard quantified statement).

$$x \text{ is an exterior point of } E \iff \exists r > 0 (B_r(x) \subseteq \mathbb{R} \setminus E).$$

Remark (Interpretation). An exterior point of E is interior to the complement: a whole ball around it misses E .

Remark (Dependencies).

- [Open Ball on the Real Line](#)

Definition (Boundary Point)

Definition 2.8.3 (Boundary Point). Let $E \subseteq \mathbb{R}$ and $x \in \mathbb{R}$. Then x is a boundary point of E exactly when

$$\forall r > 0 (B_r(x) \cap E \neq \emptyset \wedge B_r(x) \cap (\mathbb{R} \setminus E) \neq \emptyset).$$

Remark (Standard quantified statement).

$$x \text{ is a boundary point of } E \iff \forall r > 0 (B_r(x) \cap E \neq \emptyset \wedge B_r(x) \cap (\mathbb{R} \setminus E) \neq \emptyset).$$

Remark (Interpretation). A boundary point of E is approached from both sides: every ball around it meets both E and its complement.

Remark (Dependencies).

- [Open Ball on the Real Line](#)

Definition (Interior of a Set)

Definition 2.8.4 (Interior of a Set). The *interior* of $E \subseteq \mathbb{R}$, written $\text{int}(E)$, is the set of all interior points of E .

Remark (Standard quantified statement).

$$\text{int}(E) = \{ x \in \mathbb{R} : \exists r > 0 (B_r(x) \subseteq E) \}.$$

Remark (Interpretation). The interior is the largest open set inside E ; E is open exactly when $E = \text{int}(E)$. The boundary is what E and its complement share, and a set is closed exactly when it contains its boundary.

Remark (Dependencies).

- [Interior Point](#)

2.9 Limit and Isolated Points

Approachable and Detached Points

A limit point of E is one that E crowds arbitrarily closely, even with the point itself removed. An isolated point of E belongs to E but has a ball meeting E only at itself. These notions drive the convergence description of closed sets.

Definition (Limit Point)

Definition 2.9.1 (Limit Point). Let $E \subseteq \mathbb{R}$ and $x \in \mathbb{R}$. Then x is a *limit point* of E exactly when

$$\forall r > 0 \exists y \in E (0 < |y - x| < r).$$

Remark (Standard quantified statement).

$$x \text{ is a limit point of } E \iff \forall r > 0 \exists y \in E (0 < |y - x| < r).$$

Remark (Negated quantified statement).

$$x \text{ is not a limit point of } E \iff \exists r > 0 \forall y \in E (y = x \vee |y - x| \geq r).$$

Remark (Interpretation). Every punctured ball around a limit point meets E : the set accumulates at x . The strict inequality $0 < |y - x|$ excludes x itself, so x need not belong to E to be a limit point. This is the exact condition under which a sequence in E can converge to x .

Remark (Dependencies).

- [Open Ball on the Real Line](#)

Definition (Isolated Point)

Definition 2.9.2 (Isolated Point). Let $E \subseteq \mathbb{R}$ and $x \in E$. Then x is an *isolated point* of E exactly when

$$\exists r > 0 \forall y \in E (|y - x| < r \Rightarrow y = x).$$

Remark (Standard quantified statement).

$$x \text{ is an isolated point of } E \iff x \in E \wedge \exists r > 0 \forall y \in E (|y - x| < r \Rightarrow y = x).$$

Remark (Interpretation). An isolated point sits in E but stands apart: some ball around it contains no other point of E . A point of E is either isolated or a limit point of E , never both.

Remark (Dependencies).

- [Open Ball on the Real Line](#)

2.10 Unions and Intersections

Closure Laws for Open and Closed Sets

The open closure laws of §2.6 dualize to closed sets by complementation. Collecting them gives the bookkeeping rules used throughout the rest of the development.

Theorem (Closure Laws for Closed Sets)

Theorem 2.10.1 (Closure Laws for Closed Sets). *An arbitrary intersection of closed subsets of $\mathbb{R}$ is closed, and a finite union of closed subsets of $\mathbb{R}$ is closed. [Go to proof.](#)*

Remark (Standard quantified statement).

$$\left(\forall i \in I \ F_i \text{ closed} \right) \implies \left(\bigcap_{i \in I} F_i \right) \text{ closed}; \quad \left(\bigwedge_{k=1}^n F_k \text{ closed} \right) \implies \left(\bigcup_{k=1}^n F_k \right) \text{ closed}.$$

Remark (Interpretation). The roles of union and intersection swap relative to open sets: closed sets are stable under arbitrary intersection but only finite union. The asymmetry mirrors the open case under De Morgan's laws.

Remark (Dependencies).

- [Closed Set in \$\mathbb{R}\$](#)
- [Unions and Finite Intersections of Open Sets](#)

2.11 Covers and Subcovers

Covering a Set with Open Pieces

A cover of E is a family of sets whose union contains E ; an open cover uses open sets. A subcover keeps enough of the family to still cover, and a finite subcover keeps only finitely many. These are the exact ingredients of compactness, defined so as to survive beyond the line.

Definition (Open Cover)

Definition 2.11.1 (Open Cover). Let $E \subseteq \mathbb{R}$. An *open cover* of E is a family $\{U_i\}_{i \in I}$ of open subsets of $\mathbb{R}$ with

$$E \subseteq \bigcup_{i \in I} U_i.$$

Remark (Standard quantified statement).

$$\{U_i\}_{i \in I} \text{ is an open cover of } E \iff (\forall i \in I \ U_i \text{ open}) \wedge E \subseteq \bigcup_{i \in I} U_i.$$

Remark (Interpretation). An open cover is a way of surrounding every point of E with an open piece of the family. The index set I may be infinite; the question compactness asks is whether finitely many pieces already suffice.

Remark (Dependencies).

- [Open Set in \$\mathbb{R}\$](#)

Definition (Subcover and Finite Subcover)

Definition 2.11.2 (Subcover and Finite Subcover). Let $\{U_i\}_{i \in I}$ be an open cover of E . A *subcover* is a subfamily $\{U_i\}_{i \in J}$ with $J \subseteq I$ that still covers E . It is a *finite subcover* when J is finite.

Remark (Standard quantified statement).

$$\{U_i\}_{i \in J} \text{ is a finite subcover of } E \iff (J \subseteq I \wedge J \text{ is finite} \wedge E \subseteq \bigcup_{i \in J} U_i).$$

Remark (Interpretation). A subcover discards some pieces while still covering E ; a finite subcover discards all but finitely many. The existence of a finite subcover for *every* open cover is the defining property of compactness.

Remark (Dependencies).

- [Open Cover](#)

2.12 Compactness on $\mathbb{R}$

The Open-Cover Definition

Compactness is defined by the open-cover property, deliberately and not as “closed and bounded.” The open-cover formulation is the one that survives to general metric spaces and to manifolds such as the torus, where “bounded” has no intrinsic meaning. On $\mathbb{R}$ the two descriptions coincide, and that coincidence is the Heine–Borel theorem of the next section, proved rather than assumed.

Definition (Compact Set in $\mathbb{R}$)

Definition 2.12.1 (Compact Set in $\mathbb{R}$). A set $K \subseteq \mathbb{R}$ is *compact* exactly when every open cover of K admits a finite subcover:

$$\forall \{U_i\}_{i \in I} \text{ open cover of } K \exists \text{finite } J \subseteq I \left(K \subseteq \bigcup_{i \in J} U_i \right).$$

Remark (Standard quantified statement).

$$K \text{ compact} \iff \forall \{U_i\}_{i \in I} \left((\forall i \in I U_i \text{ open}) \wedge K \subseteq \bigcup_{i \in I} U_i \Rightarrow \exists \text{finite } J \subseteq I K \subseteq \bigcup_{i \in J} U_i \right).$$

Remark (Negated quantified statement).

$$K \text{ not compact} \iff \exists \{U_i\}_{i \in I} \left((\forall i \in I U_i \text{ open}) \wedge K \subseteq \bigcup_{i \in I} U_i \wedge \forall \text{finite } J \subseteq I K \not\subseteq \bigcup_{i \in J} U_i \right).$$

Remark (Failure modes). Compactness fails exactly when some open cover has no finite subcover. Two standard witnesses: $(0, 1]$ fails via the cover $\{(1/n, 2)\}_n$ (not closed), and $[0, \infty)$ fails via $\{(-1, n)\}_n$ (not bounded).

Remark (Interpretation). Compactness is a finiteness property phrased entirely in open sets: no matter how a set is covered by open pieces, finitely many already do the job. This is what later guarantees, on a compact domain, that continuous functions attain extrema and are uniformly continuous — the facts Analysis I relies on. Crucially, this definition does not mention order or boundedness, so it transfers unchanged to the torus.

Remark (Dependencies).

- [Open Cover](#)
- [Subcover and Finite Subcover](#)

Theorem (Compact Subsets of $\mathbb{R}$ Are Closed and Bounded)

Theorem 2.12.2 (Compact Subsets of $\mathbb{R}$ Are Closed and Bounded). *If $K \subseteq \mathbb{R}$ is compact, then K is closed and bounded. [Go to proof.](#)*

Remark (Standard quantified statement).

$$K \text{ compact} \implies (K \text{ closed} \wedge K \text{ bounded}).$$

Remark (Interpretation). This is the easy half of Heine–Borel and holds the same way in every metric space. Boundedness comes from covering K by growing balls about a fixed point; closedness comes from separating an outside point from K by disjoint balls. The converse — closed and bounded implies compact — is special to $\mathbb{R}^n$ and is the next theorem.

Remark (Dependencies).

- [Compact Set in \$\mathbb{R}\$](#)
- [Closed Set in \$\mathbb{R}\$](#)
- [Bounded Subset of the Real Line](#)

2.13 Heine Borel

The Capstone of the Real Line

Heine–Borel closes the chapter by proving the converse direction: on $\mathbb{R}$, closed and bounded is enough for compactness. The keystone is that every closed bounded interval $[a, b]$ is compact; the general statement follows because a closed subset of a compact set is compact.

Theorem (Closed Bounded Intervals Are Compact)

Theorem 2.13.1 (Closed Bounded Intervals Are Compact). *Every closed bounded interval $[a, b] \subseteq \mathbb{R}$ is compact. [Go to proof.](#)*

Remark (Standard quantified statement).

$$\forall a, b \in \mathbb{R} \ (a \leq b \implies [a, b] \text{ is compact}).$$

Remark (Interpretation). This is the hard half. The standard proof uses completeness directly: let S be the set of $x \in [a, b]$ such that $[a, x]$ has a finite subcover, and show $\sup S = b$ is attained. Bisection (the nested-interval method) gives the same result. Either route shows completeness is what makes the line compact in the large.

Remark (Dependencies).

- [Compact Set in \$\mathbb{R}\$](#)
- [Closed Interval](#)
- [Least-Upper-Bound Property](#)

Theorem (Heine–Borel Theorem for $\mathbb{R}$)

Theorem 2.13.2 (Heine–Borel Theorem for $\mathbb{R}$). *A set $K \subseteq \mathbb{R}$ is compact if and only if K is closed and bounded. [Go to proof.](#)*

Remark (Standard quantified statement).

$$K \text{ compact} \iff (K \text{ closed} \wedge K \text{ bounded}).$$

Remark (Interpretation). Heine–Borel is the bridge into Analysis I: it certifies that the closed bounded sets — the ones on which the Extreme Value and uniform-continuity theorems live — are exactly the compact ones. It is stated here as the equivalence *on $\mathbb{R}$* ; the open-cover side is the definition that later carries to the torus, while “closed and bounded” does not.

Remark (Exposition). With Heine–Borel in hand the chain is complete: number construction $\rightarrow$ embedded number systems $\rightarrow$ the real line as an ordered metric space $\rightarrow$ Analysis I. Completeness, introduced as the last rung of the embedding ladder, is exactly the property that makes the line compact in the large, and compactness is the hypothesis Analysis I leans on.

Remark (Dependencies).

- [Compact Subsets of \$\mathbb{R}\$ Are Closed and Bounded](#)
- [Closed Bounded Intervals Are Compact](#)

Proofs

Remark (Return). [Return to Theorem](#)

Theorem (The Distance Function Is a Metric). *The map $d(x, y) = |x - y|$ satisfies, for all $x, y, z \in \mathbb{R}$: (i) $d(x, y) \geq 0$ with $d(x, y) = 0$ iff $x = y$; (ii) $d(x, y) = d(y, x)$; and (iii) $d(x, z) \leq d(x, y) + d(y, z)$. Hence $(\mathbb{R}, d)$ is a metric space.*

Professional Standard Proof.

TODO: professional standard proof for distance-is-a-metric. □

Detailed Learning Proof.

TODO: detailed learning proof for distance-is-a-metric. □

Remark (Proof structure).

Remark (Dependencies).

- [Distance on the Real Line](#)
- TODO: link to the triangle inequality for absolute value.

Remark (Return). [Return to Theorem](#)

Proposition (Absolute Value Is Distance to the Origin). *For every $a \in \mathbb{R}$, $|a| = d(a, 0)$.*

Professional Standard Proof.

TODO: professional standard proof for abs-value-is-distance-to-zero. □

Detailed Learning Proof.

TODO: detailed learning proof for abs-value-is-distance-to-zero. □

Remark (Proof structure).

Remark (Dependencies).

- [Distance on the Real Line](#)

Remark (Return). [Return to Theorem](#)

Theorem (Open Intervals Are Open). *Every open interval (a, b) , every open ray (a, ∞) and $(-\infty, b)$, and $\mathbb{R}$ itself are open sets.*

Professional Standard Proof.

TODO: professional standard proof for open-interval-is-open. □

Detailed Learning Proof.

TODO: detailed learning proof for open-interval-is-open. □

Remark (Proof structure).

Remark (Dependencies).

- [Open Set in \$\mathbb{R}\$](#)
- [Open Interval](#)

Remark (Return). [Return to Theorem](#)

Theorem (Unions and Finite Intersections of Open Sets). *An arbitrary union of open subsets of $\mathbb{R}$ is open, and a finite intersection of open subsets of $\mathbb{R}$ is open.*

Professional Standard Proof.

TODO: professional standard proof for open-set-closure-operations. □

Detailed Learning Proof.

TODO: detailed learning proof for open-set-closure-operations. □

Remark (Proof structure).

Remark (Dependencies).

- [Open Set in \$\mathbb{R}\$](#)

Remark (Return). [Return to Theorem](#)

Theorem (Closed Sets Contain Their Limit Points). *A set $F \subseteq \mathbb{R}$ is closed if and only if it contains every one of its limit points.*

Professional Standard Proof.

TODO: professional standard proof for closed-iff-contains-limit-points. □

Detailed Learning Proof.

TODO: detailed learning proof for closed-iff-contains-limit-points. □

Remark (Proof structure).

Remark (Dependencies).

- [Closed Set in \$\mathbb{R}\$](#)
- [Limit Point](#)

Remark (Return). [Return to Theorem](#)

Theorem (Closure Laws for Closed Sets). *An arbitrary intersection of closed subsets of $\mathbb{R}$ is closed, and a finite union of closed subsets of $\mathbb{R}$ is closed.*

Professional Standard Proof.

TODO: professional standard proof for closed-set-closure-operations. □

Detailed Learning Proof.

TODO: detailed learning proof for closed-set-closure-operations. □

Remark (Proof structure).

Remark (Dependencies).

- [Closed Set in \$\mathbb{R}\$](#)
- [Unions and Finite Intersections of Open Sets](#)

Remark (Return). [Return to Theorem](#)

Theorem (Compact Subsets of $\mathbb{R}$ Are Closed and Bounded). *If $K \subseteq \mathbb{R}$ is compact, then K is closed and bounded.*

Professional Standard Proof.

TODO: professional standard proof for compact-implies-closed-bounded. □

Detailed Learning Proof.

TODO: detailed learning proof for compact-implies-closed-bounded. □

Remark (Proof structure).

Remark (Dependencies).

- [Compact Set in \$\mathbb{R}\$](#)
- [Closed Set in \$\mathbb{R}\$](#)
- [Bounded Subset of the Real Line](#)

Remark (Return). [Return to Theorem](#)

Theorem (Closed Bounded Intervals Are Compact). *Every closed bounded interval $[a, b] \subseteq \mathbb{R}$ is compact.*

Professional Standard Proof.

TODO: professional standard proof for closed-bounded-interval-compact. □

Detailed Learning Proof.

TODO: detailed learning proof for closed-bounded-interval-compact. □

Remark (Proof structure).

Remark (Dependencies).

- [Compact Set in \$\mathbb{R}\$](#)
- [Closed Interval](#)
- TODO: link to the Least Upper Bound Property / Nested Interval Property.

Remark (Return). [Return to Theorem](#)

Theorem (Heine–Borel Theorem for $\mathbb{R}$). *A set $K \subseteq \mathbb{R}$ is compact if and only if K is closed and bounded.*

Professional Standard Proof.

TODO: professional standard proof for heine-borel. □

Detailed Learning Proof.

TODO: detailed learning proof for heine-borel. □

Remark (Proof structure).

Remark (Dependencies).

- [Compact Subsets of \$\mathbb{R}\$ Are Closed and Bounded](#)
- [Closed Bounded Intervals Are Compact](#)

Chapter 3

Bounds



3.1 Completeness Axiom

Axiom of Completeness

Remark (Why completeness is needed). The ordered field axioms alone do not prevent “holes” in the number line. The rationals $\mathbb{Q}$ satisfy all field and order axioms, yet fail completeness. Consider the set

$$S = \{x \in \mathbb{Q} : x^2 < 2\}.$$

This set is nonempty (e.g. $1 \in S$) and bounded above in $\mathbb{Q}$ (e.g. 2 is an upper bound). Yet $\sup S$ does not exist as a rational number: the candidate $\sqrt{2}$ is irrational. The set S has no least upper bound in $\mathbb{Q}$ — a hole sits exactly where $\sqrt{2}$ should be.

The Completeness Axiom asserts that $\mathbb{R}$ has no such holes: every nonempty bounded-above set has a supremum *inside* $\mathbb{R}$. This single axiom is what distinguishes $\mathbb{R}$ from $\mathbb{Q}$, and it underlies every major theorem in analysis.

Definition (Least Upper Bound Property)

Definition 3.1.1 (Least Upper Bound Property). Let $(S, \leq)$ be an ordered set. The set S has the *Least Upper Bound Property* if and only if every nonempty subset of S that is bounded above in S has a supremum in S .

Remark (Standard quantified statement).

Let $(S, \leq)$ be an ordered set.

S has the Least Upper Bound Property

$$\iff \forall A \subseteq S \left[(A \neq \emptyset \wedge \exists u \in S \forall x \in A (x \leq u)) \rightarrow \exists s \in S \text{ Supremum}(s, A) \right].$$

Remark (Definition predicate reading).

$$\text{LeastUpperBoundProperty}(S) := \forall A \subseteq S \left[(A \neq \emptyset \wedge \exists u \in S \forall x \in A (x \leq u)) \rightarrow \exists s \in S \text{ Supremum}(s, A) \right]$$

Remark (Negated quantified statement).

Let $(S, \leq)$ be an ordered set.

S does not have the Least Upper Bound Property

$$\iff \exists A \subseteq S \left[A \neq \emptyset \wedge \exists u \in S \forall x \in A (x \leq u) \wedge \forall s \in S \neg \text{Supremum}(s, A) \right].$$

Remark (Negation predicate reading).

$$\neg \text{LeastUpperBoundProperty}(S) \iff \exists A \subseteq S \left[A \neq \emptyset \wedge \exists u \in S \forall x \in A (x \leq u) \wedge \forall s \in S \neg \text{Supremum}(s, A) \right]$$

Remark (Failure modes). The property fails when there is a nonempty subset of the ambient ordered set that has at least one upper bound but has no least upper bound inside that same ambient set.

Remark (Failure mode decomposition).

$$\neg \text{LeastUpperBoundProperty}(S) = \underbrace{\exists A \subseteq S (A \neq \emptyset)}_{\text{nonempty witness set}} \wedge \underbrace{\exists u \in S \forall x \in A (x \leq u)}_{\text{bounded above in } S} \wedge \underbrace{\forall s \in S \neg \text{Supremum}(s, A)}_{\text{no supremum in } S}$$

Remark (Interpretation). The Least Upper Bound Property says that upper-bounded nonempty subsets do not point toward missing ceiling elements. Whenever the order supplies at least one upper bound, it also supplies a smallest upper bound in the same ambient set. Failure therefore indicates a hole in the ordered structure rather than a defect in the subset alone.

Remark (Dependencies).

- [Ordered Set](#)
- [Subset](#)
- [Upper Bound](#)
- [Supremum](#)

Axiom (Completeness of $\mathbb{R}$)

Axiom 3.1.2 (Completeness of $\mathbb{R}$). The ordered field $\mathbb{R}$ has the Least Upper Bound Property.

Remark (Standard quantified statement).

$$\text{LeastUpperBoundProperty}(\mathbb{R}).$$

Equivalently,

$$\forall A \subseteq \mathbb{R} \left[(A \neq \emptyset \wedge \exists u \in \mathbb{R} \forall x \in A (x \leq u)) \rightarrow \exists s \in \mathbb{R} \text{ Supremum}(s, A) \right].$$

Remark (Axiom predicate reading).

$$\text{LeastUpperBoundProperty}(\mathbb{R}).$$

Remark (Negated quantified statement).

$$\neg \text{LeastUpperBoundProperty}(\mathbb{R}) \\ \iff \exists A \subseteq \mathbb{R} \left[A \neq \emptyset \wedge \exists u \in \mathbb{R} \forall x \in A (x \leq u) \wedge \forall s \in \mathbb{R} \neg \text{Supremum}(s, A) \right].$$

Remark (Negation predicate reading).

$$\neg \text{LeastUpperBoundProperty}(\mathbb{R}) \iff \exists A \subseteq \mathbb{R} \left[A \neq \emptyset \wedge \exists u \in \mathbb{R} \forall x \in A (x \leq u) \wedge \forall s \in \mathbb{R} \neg \text{Supremum}(s, A) \right].$$

Remark (Failure modes). Completeness would fail if some nonempty subset of $\mathbb{R}$ were bounded above in $\mathbb{R}$ but had no supremum in $\mathbb{R}$.

Remark (Failure mode decomposition).

$$\neg \text{LeastUpperBoundProperty}(\mathbb{R}) = \underbrace{\exists A \subseteq \mathbb{R} (A \neq \emptyset)}_{\text{nonempty witness set}} \wedge \underbrace{\exists u \in \mathbb{R} \forall x \in A (x \leq u)}_{\text{bounded above in } \mathbb{R}} \wedge \underbrace{\forall s \in \mathbb{R} \neg \text{Supremum}(s, A)}_{\text{missing real supremum}}$$

Remark (Interpretation). The completeness axiom asserts that the real line has no order gaps of the least-upper-bound kind. Nonempty sets of real numbers that are bounded above always determine a real number sitting exactly at the least possible ceiling. In Volume II, $\mathbb{R}$ is constructed as the complete ordered field arising from Dedekind cuts; in this volume, completeness is used as the standing least-upper-bound principle for real analysis. This is the structural property that distinguishes $\mathbb{R}$ from ordered fields such as $\mathbb{Q}$.

Remark (Dependencies).

- [The Real Numbers](#)
- [Ordered Field](#)
- [Least Upper Bound Property](#)

3.1.1 Nested Intervals

Nested Interval Property

Theorem (Nested Interval Property)

Theorem 3.1.3 (Nested Interval Property). *Let (I_n) be a sequence of nonempty closed intervals in $\mathbb{R}$. If $I_{n+1} \subseteq I_n$ for every $n \in \mathbb{N}$, then*

$$\bigcap_{n=1}^{\infty} I_n \neq \emptyset.$$

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned} & \forall (a_n) \subseteq \mathbb{R} \, \forall (b_n) \subseteq \mathbb{R} \\ & \left[(\forall n \in \mathbb{N} (a_n \leq b_n)) \wedge (\forall n \in \mathbb{N} ([a_{n+1}, b_{n+1}] \subseteq [a_n, b_n])) \right. \\ & \quad \left. \rightarrow \exists x \in \mathbb{R} \, \forall n \in \mathbb{N} (x \in [a_n, b_n]) \right]. \end{aligned}$$

Remark (Predicate reading).

$$\text{NestedClosedIntervals}(I_n) \implies \text{NonemptyIntersection} \left(\bigcap_{n=1}^{\infty} I_n \right).$$

Remark (Negated quantified statement).

$$\begin{aligned} & \exists (a_n) \subseteq \mathbb{R} \, \exists (b_n) \subseteq \mathbb{R} \\ & \left[(\forall n \in \mathbb{N} (a_n \leq b_n)) \wedge (\forall n \in \mathbb{N} ([a_{n+1}, b_{n+1}] \subseteq [a_n, b_n])) \right. \\ & \quad \left. \wedge \forall x \in \mathbb{R} \, \exists n \in \mathbb{N} (x \notin [a_n, b_n]) \right]. \end{aligned}$$

Remark (Negation predicate reading).

$$\text{NestedClosedIntervals}(I_n) \wedge \neg \text{NonemptyIntersection} \left(\bigcap_{n=1}^{\infty} I_n \right).$$

Remark (Failure modes). The theorem fails when a nested sequence of nonempty closed real intervals has no common real point.

Remark (Failure mode decomposition).

$$\underbrace{\text{NestedClosedIntervals}(I_n)}_{\text{closed intervals remain nested}} \wedge \underbrace{\forall x \in \mathbb{R} \, \exists n \in \mathbb{N} (x \notin I_n)}_{\text{no real point lies in every interval}}.$$

Remark (Interpretation). Nested closed intervals cannot keep shrinking or relocating in a way that loses every real point. Nesting keeps the intervals aligned, nonemptiness prevents trivial collapse at each stage, and completeness supplies a real point that survives all stages at once.

Remark (Dependencies).

[Axiom 3.1.2.](#)




Figure 3.1: Nested intervals $[a_1, b_1] \supseteq [a_2, b_2] \supseteq \cdots$ pinch toward a common point. The sequence of left endpoints is increasing and bounded above; its supremum x lies in every interval.

3.1.2 Archimedean Property

Archimedean Property

Remark (Why this requires proof). The Archimedean Property states that the natural numbers are unbounded in $\mathbb{R}$. That conclusion is not a formal consequence of the ordered-field axioms alone. There exist ordered fields satisfying all ordered-field axioms in which the natural numbers are bounded above, so the statement must be justified from stronger structure.

The needed structure is completeness. If $\mathbb{N}$ were bounded above in $\mathbb{R}$, then the Least Upper Bound Property would produce a supremum $\alpha \in \mathbb{R}$ for $\mathbb{N}$. But then $\alpha - 1$ cannot be an upper bound, so there exists $n \in \mathbb{N}$ with $n > \alpha - 1$. Hence $n + 1 > \alpha$, contradicting the fact that α is an upper bound of $\mathbb{N}$.

Go to proof.

Theorem (Archimedean Property)

Theorem 3.1.4 (Archimedean Property). *For every real number x , there exists a natural number n such that $n > x$.*

Go to proof.

Remark (Standard quantified statement).

$$\forall x \in \mathbb{R} \exists n \in \mathbb{N} (n > x).$$

Remark (Predicate reading).

$$\text{ArchimedeanProperty}(\mathbb{R}) \iff \forall x \in \mathbb{R} \exists n \in \mathbb{N} (n > x).$$

Remark (Negated quantified statement).

$$\exists x \in \mathbb{R} \forall n \in \mathbb{N} (n \leq x).$$

Remark (Negation predicate reading).

$$\neg \text{ArchimedeanProperty}(\mathbb{R}) \iff \exists x \in \mathbb{R} \forall n \in \mathbb{N} (n \leq x).$$

Remark (Failure modes). The theorem fails exactly when the natural numbers are bounded above by some real number.

Remark (Failure mode decomposition).

$$\neg \text{ArchimedeanProperty}(\mathbb{R}) = \underbrace{\exists x \in \mathbb{R}}_{\text{real upper bound}} \quad \underbrace{\forall n \in \mathbb{N} (n \leq x)}_{\text{all natural numbers remain below it}}.$$

Remark (Interpretation). The Archimedean Property says that the natural numbers do not stop below any real ceiling. No real number is so large that every natural number remains below it. In this chapter, the result is a consequence of completeness: a bounded copy of $\mathbb{N}$ would have a supremum, and that supremum is contradicted by shifting one natural number upward.

Remark (Dependencies).

Axiom 3.1.2, Definition 3.2.7.

Corollary (Archimedean Reciprocal Corollary)

Corollary 3.1.5 (Archimedean Reciprocal Corollary). *For every $x > 0$ and every $y \in \mathbb{R}$, there exists $n \in \mathbb{N}$ such that $nx > y$.*

Go to proof.

Remark (Standard quantified statement).

$$\forall x, y \in \mathbb{R} (x > 0 \rightarrow \exists n \in \mathbb{N} (nx > y)).$$

Remark (Predicate reading).

$$\text{ArchimedeanReciprocal}(\mathbb{R}) \iff \forall x, y \in \mathbb{R} (x > 0 \rightarrow \exists n \in \mathbb{N} (nx > y)).$$

Remark (Negated quantified statement).

$$\exists x, y \in \mathbb{R} (x > 0 \wedge \forall n \in \mathbb{N} (nx \leq y)).$$

Remark (Negation predicate reading).

$$\neg \text{ArchimedeanReciprocal}(\mathbb{R}) \iff \exists x, y \in \mathbb{R} (x > 0 \wedge \forall n \in \mathbb{N} (nx \leq y)).$$

Remark (Failure modes). The corollary fails when a positive real step size has all of its natural multiples bounded above by some real threshold.

Remark (Failure mode decomposition).

$$\neg \text{ArchimedeanReciprocal}(\mathbb{R}) = \underbrace{\exists x \in \mathbb{R} (x > 0)}_{\text{positive step size}} \wedge \underbrace{\exists y \in \mathbb{R} \forall n \in \mathbb{N} (nx \leq y)}_{\text{all natural multiples stay bounded}}.$$

Remark (Interpretation). Any fixed positive real step eventually passes any real threshold when repeated enough times. The statement is the Archimedean Property applied after rescaling by the positive number x : growth by units becomes growth by steps of size x .

Remark (Dependencies).

Theorem 3.1.4.

3.1.3 Density

Density

Definition (Dense Subset)

Definition 3.1.6 (Dense Subset). Let $D \subseteq S \subseteq \mathbb{R}$. The set D is *dense in* S if and only if for every $x \in S$ and every $\varepsilon > 0$ there exists $d \in D$ such that $|x - d| < \varepsilon$.

Remark (Standard quantified statement).

Let $D \subseteq S \subseteq \mathbb{R}$.

D is dense in $S \iff \forall x \in S \forall \varepsilon > 0 \exists d \in D (|x - d| < \varepsilon)$.

Remark (Definition predicate reading).

$\text{DenseSubset}(D, S) := \forall x \in S \forall \varepsilon > 0 \exists d \in D (|x - d| < \varepsilon)$.

Remark (Negated quantified statement).

Let $D \subseteq S \subseteq \mathbb{R}$.

D is not dense in $S \iff \exists x \in S \exists \varepsilon_0 > 0 \forall d \in D (|x - d| \geq \varepsilon_0)$.

Remark (Negation predicate reading).

$\neg \text{DenseSubset}(D, S) \iff \exists x \in S \exists \varepsilon_0 > 0 \forall d \in D (|x - d| \geq \varepsilon_0)$.

Remark (Failure modes). Density fails when some point of S has a positive-radius neighborhood that contains no point of D .

Remark (Failure mode decomposition).

$$\neg \text{DenseSubset}(D, S) = \underbrace{\exists x \in S}_{\text{point not approximated}} \quad \underbrace{\exists \varepsilon_0 > 0}_{\text{positive exclusion radius}} \quad \underbrace{\forall d \in D (|x - d| \geq \varepsilon_0)}_{\text{no point of } D \text{ enters the neighborhood}}.$$

Remark (Interpretation). Density is an approximation property inside S . It does not say that every point of S already belongs to D ; it says that points of D occur arbitrarily close to every point of S . Thus D can be a proper subset of S while still leaving no positive-width gap inside S .




Figure. Illustration of a
Dense Subset $D \subseteq S$.

Given any $x \in S$ (black dot) and any $\varepsilon > 0$
(red interval width), there exists $d \in D$
(red-filled blue dot) such that $|x - d| < \varepsilon$.

Definition (Order Dense Subset)

Definition 3.1.7 (Order Dense Subset). Let X be a linearly ordered set and let $A \subseteq X$. The subset A is *order dense in* X if and only if for every $a, b \in X$ with $a < b$ there exists $c \in A$ such that $a < c < b$.

Remark (Standard quantified statement).

Let X be a linearly ordered set and $A \subseteq X$.

A is order dense in $X \iff \forall a, b \in X (a < b \rightarrow \exists c \in A (a < c < b))$.

Remark (Definition predicate reading).

$\text{OrderDenseSubset}(A, X) := \forall a, b \in X (a < b \rightarrow \exists c \in A (a < c < b))$.

Remark (Negated quantified statement).

Let X be a linearly ordered set and $A \subseteq X$.

A is not order dense in $X \iff \exists a, b \in X (a < b \wedge \forall c \in A (c \leq a \vee b \leq c))$.

Remark (Negation predicate reading).

$\neg \text{OrderDenseSubset}(A, X) \iff \exists a, b \in X (a < b \wedge \forall c \in A (c \leq a \vee b \leq c))$.

Remark (Failure modes). Order density fails when there is a nonempty open interval of X that contains no point of A .

Remark (Failure mode decomposition).

$$\neg \text{OrderDenseSubset}(A, X) = \underbrace{\exists a, b \in X (a < b)}_{\text{nonempty interval}} \wedge \underbrace{\forall c \in A (c \leq a \vee b \leq c)}_{\text{no point of } A \text{ lies between } a \text{ and } b} .$$

Remark (Interpretation). Order density is an interval-penetration property. It ignores metric distance and asks only whether every ordered gap contains a point of the subset. A subset can therefore be order dense in X precisely when no nonempty open interval of X is missed entirely by that subset.

Remark (Dependencies).

- [Total Order](#)
- [Subset](#)

Theorem (Density of the Rational Numbers in $\mathbb{R}$)

Theorem 3.1.8 (Density of the Rational Numbers in $\mathbb{R}$). *For every $a, b \in \mathbb{R}$ with $a < b$, there exists $q \in \mathbb{Q}$ such that $a < q < b$. [Go to proof.](#)*

Remark (Standard quantified statement).

$\forall a, b \in \mathbb{R} (a < b \rightarrow \exists q \in \mathbb{Q} (a < q < b))$.

Remark (Predicate reading).

$\text{OrderDenseSubset}(\mathbb{Q}, \mathbb{R}) \iff \forall a, b \in \mathbb{R} (a < b \rightarrow \exists q \in \mathbb{Q} (a < q < b))$.

Remark (Negated quantified statement).

$$\exists a, b \in \mathbb{R} (a < b \wedge \forall q \in \mathbb{Q} (q \leq a \vee b \leq q)).$$

Remark (Negation predicate reading).

$$\neg \text{OrderDenseSubset}(\mathbb{Q}, \mathbb{R}) \iff \exists a, b \in \mathbb{R} (a < b \wedge \forall q \in \mathbb{Q} (q \leq a \vee b \leq q)).$$

Remark (Failure modes). The theorem would fail precisely if some nonempty real interval contained no rational number.

Remark (Failure mode decomposition).

$$\neg \text{OrderDenseSubset}(\mathbb{Q}, \mathbb{R}) = \underbrace{\exists a, b \in \mathbb{R} (a < b)}_{\text{nonempty real interval}} \wedge \underbrace{\forall q \in \mathbb{Q} (q \leq a \vee b \leq q)}_{\text{no rational lies inside}}.$$

Remark (Interpretation). Rational numbers penetrate every real interval. The theorem does not say that every real number is rational; it says that rational numbers are never separated from a real interval by a positive order gap. This is the interval form of rational density in the real line.

Remark (Dependencies).

[Definition 3.1.7](#), [Theorem 3.1.4](#).

Theorem (Density of the Irrational Numbers in $\mathbb{R}$)

Theorem 3.1.9 (Density of the Irrational Numbers in $\mathbb{R}$). *For every $a, b \in \mathbb{R}$ with $a < b$, there exists $s \in \mathbb{R} \setminus \mathbb{Q}$ such that $a < s < b$. [Go to proof](#).*

Remark (Standard quantified statement).

$$\forall a, b \in \mathbb{R} (a < b \rightarrow \exists s \in \mathbb{R} \setminus \mathbb{Q} (a < s < b)).$$

Remark (Predicate reading).

$$\text{OrderDenseSubset}(\mathbb{R} \setminus \mathbb{Q}, \mathbb{R}) \iff \forall a, b \in \mathbb{R} (a < b \rightarrow \exists s \in \mathbb{R} \setminus \mathbb{Q} (a < s < b)).$$

Remark (Negated quantified statement).

$$\exists a, b \in \mathbb{R} (a < b \wedge \forall s \in \mathbb{R} \setminus \mathbb{Q} (s \leq a \vee b \leq s)).$$

Remark (Negation predicate reading).

$$\neg \text{OrderDenseSubset}(\mathbb{R} \setminus \mathbb{Q}, \mathbb{R}) \iff \exists a, b \in \mathbb{R} (a < b \wedge \forall s \in \mathbb{R} \setminus \mathbb{Q} (s \leq a \vee b \leq s)).$$

Remark (Failure modes). The theorem would fail precisely if some nonempty real interval contained no irrational number.

Remark (Failure mode decomposition).

$$\neg \text{OrderDenseSubset}(\mathbb{R} \setminus \mathbb{Q}, \mathbb{R}) = \underbrace{\exists a, b \in \mathbb{R} (a < b)}_{\text{nonempty real interval}} \wedge \underbrace{\forall s \in \mathbb{R} \setminus \mathbb{Q} (s \leq a \vee b \leq s)}_{\text{no irrational lies inside}}.$$

Remark (Interpretation). Irrational numbers also penetrate every real interval. One way to see the result is to insert a rational point into a translated and rescaled interval, then shift it by an irrational amount. The outcome is that the irrational numbers, like the rational numbers, leave no interval gap in $\mathbb{R}$.

Remark (Dependencies).

[Definition 3.1.7](#), [Theorem 3.1.8](#).

Corollary (Rational and Irrational Points in Every Open Interval)

Corollary 3.1.10 (Rational and Irrational Points in Every Open Interval). *Every open interval in $\mathbb{R}$ contains both a rational number and an irrational number.* [Go to proof.](#)

Remark (Standard quantified statement).

$$\forall a, b \in \mathbb{R} \left[a < b \rightarrow \left(\exists q \in \mathbb{Q} (a < q < b) \wedge \exists s \in \mathbb{R} \setminus \mathbb{Q} (a < s < b) \right) \right].$$

Remark (Predicate reading).

$$\text{OrderDenseSubset}(\mathbb{Q}, \mathbb{R}) \wedge \text{OrderDenseSubset}(\mathbb{R} \setminus \mathbb{Q}, \mathbb{R}).$$

Remark (Negated quantified statement).

$$\exists a, b \in \mathbb{R} \left[a < b \wedge \left(\forall q \in \mathbb{Q} (q \leq a \vee b \leq q) \vee \forall s \in \mathbb{R} \setminus \mathbb{Q} (s \leq a \vee b \leq s) \right) \right].$$

Remark (Negation predicate reading).

$$\neg \left[\text{OrderDenseSubset}(\mathbb{Q}, \mathbb{R}) \wedge \text{OrderDenseSubset}(\mathbb{R} \setminus \mathbb{Q}, \mathbb{R}) \right].$$

Remark (Failure modes). The corollary fails if a nonempty real interval misses all rational points or misses all irrational points.

Remark (Failure mode decomposition).

$$\exists a, b \in \mathbb{R} (a < b) \wedge \left[\underbrace{\forall q \in \mathbb{Q} (q \leq a \vee b \leq q)}_{\text{no rational point}} \vee \underbrace{\forall s \in \mathbb{R} \setminus \mathbb{Q} (s \leq a \vee b \leq s)}_{\text{no irrational point}} \right].$$

Remark (Interpretation). The corollary combines the two density theorems. Every real interval, no matter how small, contains representatives of both arithmetic types. Thus neither the rational numbers nor the irrational numbers occupy isolated regions of the real line.

Remark (Dependencies).

[Theorem 3.1.8](#), [Theorem 3.1.9](#).




Figure 3.2: Visualization of the Density of $\mathbb{Q}$ and $\mathbb{R} \setminus \mathbb{Q}$ in $\mathbb{R}$. Zooming into any interval reveals that both sets are always present.

Remark (Zoom invariance and structural consequences).

Density in $\mathbb{R}$ has a scale-invariant consequence: no matter how small a nonempty open interval becomes, it still contains both rational and irrational points. The interleaving of $\mathbb{Q}$ and $\mathbb{R} \setminus \mathbb{Q}$ is therefore not a large-scale phenomenon. It persists at every interval scale.

This has three important consequences. First, the continuum does not become locally discrete under magnification. For every nonempty open interval (a, b) , both

$$\mathbb{Q} \cap (a, b) \quad \text{and} \quad (\mathbb{R} \setminus \mathbb{Q}) \cap (a, b)$$

remain nonempty, and in fact infinite. Density is therefore stable under repeated restriction to smaller intervals.

Second, this behavior sharply distinguishes the real line from discrete numerical models. Floating-point systems form a discrete set of representable values, so repeated magnification eventually reaches a scale at which adjacent representable numbers leave no further point strictly between them. At that stage, interval penetration fails in the discrete model even though it continues to hold in $\mathbb{R}$.

Third, the Archimedean Property supplies the scale mechanism behind this behavior. Given any positive scale ε , there exists $n \in \mathbb{N}$ with

$$\frac{1}{n} < \varepsilon.$$

This makes it possible to construct rational approximations at arbitrarily small interval scales, which is one of the basic engines behind density arguments.

The resulting contrast is structural. Smooth mathematical models are built in a setting where interval penetration persists at every scale, whereas finite computational models eventually encounter a smallest representable separation. The transition from continuum behavior to discrete behavior occurs exactly where nonempty intervals in the numerical model cease to contain further representable points.

Density Dependency Map




Figure 3.3: Dependency map for the density subchapter. The Floor Lemma and the Archimedean Corollary support the proof of the density of the rational numbers in $\mathbb{R}$. The irrational density theorem is then derived from the rational density result, and the final corollary combines both density theorems into a single interval statement. The definition records the local vocabulary of the subchapter.

3.1.4 Completeness Summary




Figure 3.4: Dependency map for the completeness subchapter. The Completeness Axiom yields the Nested Interval Property and the existence of $\sqrt{a}$; the Nested Interval Property then supports the Archimedean Property, the Floor Lemma, and the density results for $\mathbb{Q}$ and for the irrational numbers.

Remark (Connections forward). Completeness appears in every major limit theorem in the notes. The Monotone Convergence Theorem (convergence chapter) constructs the limit of a bounded monotone sequence as $\sup\{a_n : n \in \mathbb{N}\}$; existence of this supremum requires completeness. The Bolzano–Weierstrass Theorem applies the Nested Interval Property directly to extract a convergent subsequence from any bounded sequence. The Cauchy Criterion identifies convergent sequences purely in terms of their own spacing, without naming the limit in advance; its proof constructs the limit as a supremum and again requires completeness. In metric spaces (Distance chapter), completeness in the sense of Cauchy sequences is the natural generalization of the same idea to abstract settings. Every ε - δ proof that locates a limit or extremum is drawing on this chapter’s infrastructure.

3.2 Bounds and Extremal Values

3.2.1 Upper and Lower Bounds

Bounds are the first level of extremal information. They compare one candidate element with every element of a nonempty set, but they do not require the candidate to be sharp or to belong to the set.




Figure 3.5: Upper and lower bound conditions on a nonempty set.

Bound

Definition 3.2.1 (Bound). Let $(S, \leq)$ be an ordered set, let $A \subseteq S$ be nonempty, and let $b \in S$. A *bound* of A means a one-sided comparison between b and every element of A : either $x \leq b$ for all $x \in A$, or $b \leq x$ for all $x \in A$.

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad [(\forall x \in A)(x \leq b) \text{ or } (\forall x \in A)(b \leq x)].$$

Remark (Definition predicate reading).

$$\text{Bound}_{S, \leq}(b, A) := \text{UB}_{S, \leq}(b, A) \vee \text{LB}_{S, \leq}(b, A).$$

Remark (Negated quantified statement).

$$A = \emptyset \quad \text{or} \quad [(\exists x \in A)(b < x) \text{ and } (\exists y \in A)(y < b)].$$

Remark (Negation predicate reading).

$$\neg \text{Bound}_{S, \leq}(b, A) \iff A = \emptyset \vee (\neg \text{UB}_{S, \leq}(b, A) \wedge \neg \text{LB}_{S, \leq}(b, A)).$$

Remark (Failure modes). The statement fails if the set is empty, or if b lies strictly between two elements of A .

Remark (Failure mode decomposition). Failure of the upper-bound side means some element of A is strictly larger than b . Failure of the lower-bound side means some element of A is strictly smaller than b . Failure to be any bound requires both witnesses.

Upper Bound

Definition 3.2.2 (Upper Bound). Let $(S, \leq)$ be an ordered set, let $A \subseteq S$ be nonempty, and let $u \in S$. The element u is an *upper bound* of A if every element of A is less than or equal to u .

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad (\forall x \in A)(x \leq u).$$

Remark (Definition predicate reading).

$$\text{UB}_{S,\leq}(u, A) := A \neq \emptyset \wedge (\forall x \in A)(x \leq u).$$

Remark (Negated quantified statement).

$$A = \emptyset \quad \text{or} \quad (\exists x \in A)(u < x).$$

Remark (Negation predicate reading).

$$\neg \text{UB}_{S,\leq}(u, A) \iff A = \emptyset \vee (\exists x \in A)(u < x).$$

Remark (Failure modes). The set may be empty, or a counterexample element of A may sit strictly above the proposed bound.

Remark (Failure mode decomposition). To disprove that u is an upper bound, produce either the structural failure $A = \emptyset$ or a witness $x \in A$ with $u < x$.

Lower Bound

Definition 3.2.3 (Lower Bound). Let $(S, \leq)$ be an ordered set, let $A \subseteq S$ be nonempty, and let $\ell \in S$. The element ℓ is a *lower bound* of A if every element of A is greater than or equal to ℓ .

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad (\forall x \in A)(\ell \leq x).$$

Remark (Definition predicate reading).

$$\text{LB}_{S,\leq}(\ell, A) := A \neq \emptyset \wedge (\forall x \in A)(\ell \leq x).$$

Remark (Negated quantified statement).

$$A = \emptyset \quad \text{or} \quad (\exists x \in A)(x < \ell).$$

Remark (Negation predicate reading).

$$\neg \text{LB}_{S,\leq}(\ell, A) \iff A = \emptyset \vee (\exists x \in A)(x < \ell).$$

Remark (Failure modes). The set may be empty, or a counterexample element of A may sit strictly below the proposed bound.

Remark (Failure mode decomposition). To disprove that ℓ is a lower bound, produce either the structural failure $A = \emptyset$ or a witness $x \in A$ with $x < \ell$.

Bounded Above

Definition 3.2.4 (Bounded Above). Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. The set A is *bounded above* if it has at least one upper bound in S .

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad (\exists u \in S)(\forall x \in A)(x \leq u).$$

Remark (Definition predicate reading).

$$\text{BddAbove}_{S,\leq}(A) := (\exists u \in S) \text{UB}_{S,\leq}(u, A).$$

Remark (Negated quantified statement).

$$A = \emptyset \quad \text{or} \quad (\forall u \in S)(\exists x \in A)(u < x).$$

Remark (Negation predicate reading).

$$\neg \text{BddAbove}_{S,\leq}(A) \iff A = \emptyset \vee (\forall u \in S) \neg \text{UB}_{S,\leq}(u, A).$$

Remark (Failure modes). The set may be empty, or every proposed upper bound may be defeated by some larger element of A .

Remark (Failure mode decomposition). Unboundedness above is not one failed candidate; it is a uniform procedure that defeats every candidate $u \in S$.

Bounded Below

Definition 3.2.5 (Bounded Below). Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. The set A is *bounded below* if it has at least one lower bound in S .

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad (\exists \ell \in S)(\forall x \in A)(\ell \leq x).$$

Remark (Definition predicate reading).

$$\text{BddBelow}_{S,\leq}(A) := (\exists \ell \in S) \text{LB}_{S,\leq}(\ell, A).$$

Remark (Negated quantified statement).

$$A = \emptyset \quad \text{or} \quad (\forall \ell \in S)(\exists x \in A)(x < \ell).$$

Remark (Negation predicate reading).

$$\neg \text{BddBelow}_{S,\leq}(A) \iff A = \emptyset \vee (\forall \ell \in S) \neg \text{LB}_{S,\leq}(\ell, A).$$

Remark (Failure modes). The set may be empty, or every proposed lower bound may be defeated by some smaller element of A .

Remark (Failure mode decomposition). Unboundedness below is a uniform defeat of all lower-bound candidates.

Bounded Set

Definition 3.2.6 (Bounded Set). Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. The set A is *bounded* if it is bounded above and bounded below.

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad (\exists \ell, u \in S)(\forall x \in A)(\ell \leq x \leq u).$$

Remark (Definition predicate reading).

$$\text{Bdd}_{S,\leq}(A) := \text{BddBelow}_{S,\leq}(A) \wedge \text{BddAbove}_{S,\leq}(A).$$

Remark (Negated quantified statement).

$$A = \emptyset \quad \text{or} \quad (\forall \ell, u \in S)(\exists x \in A)(x < \ell \text{ or } u < x).$$

Remark (Negation predicate reading).

$$\neg \text{Bdd}_{S,\leq}(A) \iff A = \emptyset \vee \neg \text{BddBelow}_{S,\leq}(A) \vee \neg \text{BddAbove}_{S,\leq}(A).$$

Remark (Failure modes). The set may be empty, unbounded below, unbounded above, or both.

Remark (Failure mode decomposition). Boundedness has two independent one-sided requirements. A failure on either side is enough to destroy boundedness.

3.2.2 Suprema and Infima

Suprema and infima are sharp bounds. A supremum is the least upper bound; an infimum is the greatest lower bound. The set remains required to be nonempty throughout this section.




Figure 3.6: Suprema and infima are sharp bounds, whether or not they are elements of the set.

Supremum

Definition 3.2.7 (Supremum). Let $(S, \leq)$ be an ordered set, let $A \subseteq S$ be nonempty, and let $s \in S$. The element s is the *supremum* of A if s is an upper bound of A and no smaller element of S is an upper bound of A .

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad (\forall x \in A)(x \leq s) \quad \text{and} \quad (\forall u \in S)[(\forall x \in A)(x \leq u) \Rightarrow s \leq u].$$

Remark (Definition predicate reading).

$$\text{Sup}_{S,\leq}(s, A) := \text{UB}_{S,\leq}(s, A) \wedge (\forall u \in S)(\text{UB}_{S,\leq}(u, A) \Rightarrow s \leq u).$$

Remark (Negated quantified statement).

$$A = \emptyset \quad \text{or} \quad (\exists x \in A)(s < x) \quad \text{or} \quad (\exists u \in S)[(\forall x \in A)(x \leq u) \text{ and } u < s].$$

Remark (Negation predicate reading).

$$\neg \text{Sup}_{S,\leq}(s, A) \iff A = \emptyset \vee \neg \text{UB}_{S,\leq}(s, A) \vee (\exists u \in S)(\text{UB}_{S,\leq}(u, A) \wedge u < s).$$

Remark (Failure modes). The set may be empty; s may fail to be an upper bound; or s may be an upper bound that is not least among upper bounds.

Remark (Failure mode decomposition). The first clause is a domain failure. The second is a bound failure. The third is a sharpness failure witnessed by a strictly smaller upper bound.

Infimum

Definition 3.2.8 (Infimum). Let $(S, \leq)$ be an ordered set, let $A \subseteq S$ be nonempty, and let $i \in S$. The element i is the *infimum* of A if i is a lower bound of A and no larger element of S is a lower bound of A .

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad (\forall x \in A)(i \leq x) \quad \text{and} \quad (\forall \ell \in S)[(\forall x \in A)(\ell \leq x) \Rightarrow \ell \leq i].$$

Remark (Definition predicate reading).

$$\text{Inf}_{S,\leq}(i, A) := \text{LB}_{S,\leq}(i, A) \wedge (\forall \ell \in S)(\text{LB}_{S,\leq}(\ell, A) \Rightarrow \ell \leq i).$$

Remark (Negated quantified statement).

$$A = \emptyset \quad \text{or} \quad (\exists x \in A)(x < i) \quad \text{or} \quad (\exists \ell \in S)[(\forall x \in A)(\ell \leq x) \text{ and } i < \ell].$$

Remark (Negation predicate reading).

$$\neg \text{Inf}_{S,\leq}(i, A) \iff A = \emptyset \vee \neg \text{LB}_{S,\leq}(i, A) \vee (\exists \ell \in S)(\text{LB}_{S,\leq}(\ell, A) \wedge i < \ell).$$

Remark (Failure modes). The set may be empty; i may fail to be a lower bound; or i may be a lower bound that is not greatest among lower bounds.

Remark (Failure mode decomposition). The first clause is a domain failure. The second is a bound failure. The third is a sharpness failure witnessed by a strictly larger lower bound.

Uniqueness of the Supremum

Proposition 3.2.9 (Uniqueness of the Supremum). *Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. If $s, t \in S$ are both suprema of A , then $s = t$.*

Go to proof.

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad \text{Sup}_{S, \leq}(s, A) \quad \text{and} \quad \text{Sup}_{S, \leq}(t, A) \quad \implies \quad s = t.$$

Remark (Predicate reading).

$$\text{Sup}_{S, \leq}(s, A) \wedge \text{Sup}_{S, \leq}(t, A) \implies s = t.$$

Remark (Negated quantified statement).

$$A \neq \emptyset \quad \text{and} \quad \text{Sup}_{S, \leq}(s, A) \quad \text{and} \quad \text{Sup}_{S, \leq}(t, A) \quad \text{and} \quad s \neq t.$$

Remark (Negation predicate reading).

$$\text{Sup}_{S, \leq}(s, A) \wedge \text{Sup}_{S, \leq}(t, A) \wedge s \neq t.$$

Remark (Failure modes). Failure would require two distinct least upper bounds for the same nonempty set.

Remark (Failure mode decomposition). Each candidate is an upper bound and each lies below every upper bound; applying these minimality clauses to the other candidate gives both $s \leq t$ and $t \leq s$.

Uniqueness of the Infimum

Proposition 3.2.10 (Uniqueness of the Infimum). *Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. If $i, j \in S$ are both infima of A , then $i = j$.*

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad \text{Inf}_{S, \leq}(i, A) \quad \text{and} \quad \text{Inf}_{S, \leq}(j, A) \quad \implies \quad i = j.$$

Remark (Predicate reading).

$$\text{Inf}_{S, \leq}(i, A) \wedge \text{Inf}_{S, \leq}(j, A) \implies i = j.$$

Remark (Negated quantified statement).

$$A \neq \emptyset \quad \text{and} \quad \text{Inf}_{S, \leq}(i, A) \quad \text{and} \quad \text{Inf}_{S, \leq}(j, A) \quad \text{and} \quad i \neq j.$$

Remark (Negation predicate reading).

$$\text{Inf}_{S, \leq}(i, A) \wedge \text{Inf}_{S, \leq}(j, A) \wedge i \neq j.$$

Remark (Failure modes). Failure would require two distinct greatest lower bounds for the same nonempty set.

Remark (Failure mode decomposition). Each candidate is a lower bound and each lies above every lower bound; applying these maximality clauses to the other candidate gives both $i \leq j$ and $j \leq i$.

Upper Bound Property of the Supremum

Proposition 3.2.11 (Upper Bound Property of the Supremum). *Let $(S, \leq)$ be an ordered set, let $A \subseteq S$ be nonempty, and suppose that $s = \sup A$ exists in S . Then every element of A lies below s :*

$$(\forall x \in A)(x \leq s).$$

Go to proof.

Remark (Predicate reading).

$$\text{Sup}_{S, \leq}(s, A) \implies \text{UB}_{S, \leq}(s, A).$$

Remark (Negated quantified statement).

$$\text{Sup}_{S, \leq}(s, A) \quad \text{and} \quad (\exists x \in A)(s < x).$$

Remark (Failure modes). Failure would mean that a supremum is not an upper bound.

Remark (Failure mode decomposition). This is exactly the first clause in the definition of supremum.

Lower Bound Property of the Infimum

Proposition 3.2.12 (Lower Bound Property of the Infimum). *Let $(S, \leq)$ be an ordered set, let $A \subseteq S$ be nonempty, and suppose that $i = \inf A$ exists in S . Then every element of A lies above i :*

$$(\forall x \in A)(i \leq x).$$

Remark (Predicate reading).

$$\text{Inf}_{S, \leq}(i, A) \implies \text{LB}_{S, \leq}(i, A).$$

Remark (Negated quantified statement).

$$\text{Inf}_{S, \leq}(i, A) \quad \text{and} \quad (\exists x \in A)(x < i).$$

Remark (Failure modes). Failure would mean that an infimum is not a lower bound.

Remark (Failure mode decomposition). This is exactly the first clause in the definition of infimum.

Subset Inclusion Preserves Upper Bounds

Lemma 3.2.13 (Subset Inclusion Preserves Upper Bounds). *Let $A \subseteq B \subseteq S$ be nonempty subsets of an ordered set $(S, \leq)$. If u is an upper bound of B , then u is an upper bound of A .*

Go to proof.

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad A \subseteq B, \quad (\forall y \in B)(y \leq u) \implies (\forall x \in A)(x \leq u).$$

Remark (Predicate reading).

$$A \subseteq B \wedge \text{UB}_{S,\leq}(u, B) \implies \text{UB}_{S,\leq}(u, A).$$

Remark (Negated quantified statement).

$$A \subseteq B, \quad \text{UB}_{S,\leq}(u, B), \quad \text{and} \quad (\exists x \in A)(u < x).$$

Remark (Failure modes). Failure would require a counterexample in the smaller set even though the larger set is already bounded by u .

Remark (Failure mode decomposition). Any counterexample $x \in A$ is also an element of B , contradicting the upper-bound hypothesis for B .

Subset Inclusion Preserves Lower Bounds

Lemma 3.2.14 (Subset Inclusion Preserves Lower Bounds). *Let $A \subseteq B \subseteq S$ be nonempty subsets of an ordered set $(S, \leq)$. If ℓ is a lower bound of B , then ℓ is a lower bound of A .*

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad A \subseteq B, \quad (\forall y \in B)(\ell \leq y) \implies (\forall x \in A)(\ell \leq x).$$

Remark (Predicate reading).

$$A \subseteq B \wedge \text{LB}_{S,\leq}(\ell, B) \implies \text{LB}_{S,\leq}(\ell, A).$$

Remark (Negated quantified statement).

$$A \subseteq B, \quad \text{LB}_{S,\leq}(\ell, B), \quad \text{and} \quad (\exists x \in A)(x < \ell).$$

Remark (Failure modes). Failure would require a smaller-set counterexample to a lower bound that already works on the larger set.

Remark (Failure mode decomposition). Any counterexample $x \in A$ lies in B , contradicting the lower-bound hypothesis for B .

Supremum Is Monotone Under Inclusion

Theorem 3.2.15 (Supremum Is Monotone Under Inclusion). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded above. If $A \subseteq B$, then $\sup A \leq \sup B$.*

Go to proof.

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad B \neq \emptyset, \quad A \subseteq B, \quad \text{Sup}_{\mathbb{R},\leq}(s_A, A), \quad \text{Sup}_{\mathbb{R},\leq}(s_B, B) \implies s_A \leq s_B.$$

Remark (Predicate reading).

$$A \subseteq B \wedge \text{Sup}(s_A, A) \wedge \text{Sup}(s_B, B) \implies s_A \leq s_B.$$

Remark (Negated quantified statement).

$$A \subseteq B, \quad \text{Sup}(s_A, A), \quad \text{Sup}(s_B, B), \quad s_B < s_A.$$

Remark (Failure modes). Failure would mean enlarging a nonempty set lowers its least upper bound.

Remark (Failure mode decomposition). The value $\sup B$ bounds B , hence bounds A ; since $\sup A$ is least among upper bounds of A , $\sup A \leq \sup B$.

Infimum Is Monotone Under Inclusion

Theorem 3.2.16 (Infimum Is Monotone Under Inclusion). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded below. If $A \subseteq B$, then $\inf B \leq \inf A$.*

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad B \neq \emptyset, \quad A \subseteq B, \quad \text{Inf}_{\mathbb{R}, \leq}(i_A, A), \quad \text{Inf}_{\mathbb{R}, \leq}(i_B, B) \implies i_B \leq i_A.$$

Remark (Predicate reading).

$$A \subseteq B \wedge \text{Inf}(i_A, A) \wedge \text{Inf}(i_B, B) \implies i_B \leq i_A.$$

Remark (Negated quantified statement).

$$A \subseteq B, \quad \text{Inf}(i_A, A), \quad \text{Inf}(i_B, B), \quad i_A < i_B.$$

Remark (Failure modes). Failure would mean enlarging a nonempty set raises its greatest lower bound.

Remark (Failure mode decomposition). The value $\inf B$ bounds B below, hence bounds A below; since $\inf A$ is greatest among lower bounds of A , $\inf B \leq \inf A$.

Upper Bound Comparison with the Supremum

Proposition 3.2.17 (Upper Bound Comparison with the Supremum). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded above, and let $s = \sup A$. For $u \in \mathbb{R}$, the number u is an upper bound of A if and only if $s \leq u$.*

Go to proof.

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad \text{Sup}(s, A) \implies (\forall u \in \mathbb{R}) [\text{UB}(u, A) \iff s \leq u].$$

Remark (Predicate reading).

$$\text{Sup}(s, A) \implies \forall u (\text{UB}(u, A) \leftrightarrow s \leq u).$$

Remark (Negated quantified statement).

$$\text{Sup}(s, A) \quad \text{and} \quad (\exists u)[(\text{UB}(u, A) \wedge u < s) \vee (s \leq u \wedge \neg \text{UB}(u, A))].$$

Remark (Failure modes). Failure would mean an upper bound below the supremum, or a number above the supremum that fails to bound the set.

Remark (Failure mode decomposition). Leastness gives the forward implication. The fact that s bounds A gives the reverse implication by transitivity.

Lower Bound Comparison with the Infimum

Proposition 3.2.18 (Lower Bound Comparison with the Infimum). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded below, and let $i = \inf A$. For $\ell \in \mathbb{R}$, the number ℓ is a lower bound of A if and only if $\ell \leq i$.*

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad \text{Inf}(i, A) \implies (\forall \ell \in \mathbb{R})[\text{LB}(\ell, A) \iff \ell \leq i].$$

Remark (Predicate reading).

$$\text{Inf}(i, A) \implies \forall \ell (\text{LB}(\ell, A) \leftrightarrow \ell \leq i).$$

Remark (Negated quantified statement).

$$\text{Inf}(i, A) \quad \text{and} \quad (\exists \ell)[(\text{LB}(\ell, A) \wedge i < \ell) \vee (\ell \leq i \wedge \neg \text{LB}(\ell, A))].$$

Remark (Failure modes). Failure would mean a lower bound above the infimum, or a number below the infimum that fails to bound the set below.

Remark (Failure mode decomposition). Greatestness gives the forward implication. The fact that i bounds A below gives the reverse implication by transitivity.

Every Element Lies Below the Supremum

Proposition 3.2.19 (Every Element Lies Below the Supremum). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded above, and let $s = \sup A$. Then $x \leq s$ for every $x \in A$.
[Go to proof.](#)*

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad \text{Sup}(s, A) \implies (\forall x \in A)(x \leq s).$$

Remark (Predicate reading).

$$\text{Sup}(s, A) \implies \text{UB}(s, A).$$

Remark (Negated quantified statement).

$$\text{Sup}(s, A) \quad \text{and} \quad (\exists x \in A)(s < x).$$

Remark (Failure modes). Failure would mean the supremum is not an upper bound.

Remark (Failure mode decomposition). This is the upper-bound clause in the definition of supremum.

Every Element Lies Above the Infimum

Proposition 3.2.20 (Every Element Lies Above the Infimum). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded below, and let $i = \inf A$. Then $i \leq x$ for every $x \in A$.*

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad \text{Inf}(i, A) \implies (\forall x \in A)(i \leq x).$$

Remark (Predicate reading).

$$\text{Inf}(i, A) \implies \text{LB}(i, A).$$

Remark (Negated quantified statement).

$$\text{Inf}(i, A) \quad \text{and} \quad (\exists x \in A)(x < i).$$

Remark (Failure modes). Failure would mean the infimum is not a lower bound.

Remark (Failure mode decomposition). This is the lower-bound clause in the definition of infimum.

Infimum Is Less Than or Equal to the Supremum

Theorem 3.2.21 (Infimum Is Less Than or Equal to the Supremum). *Let $A \subseteq \mathbb{R}$ be nonempty, bounded below, and bounded above. Then $\inf A \leq \sup A$.*

Go to proof.

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad \text{Inf}(i, A), \quad \text{Sup}(s, A) \implies i \leq s.$$

Remark (Predicate reading).

$$\text{Inf}(i, A) \wedge \text{Sup}(s, A) \implies i \leq s.$$

Remark (Negated quantified statement).

$$\text{Inf}(i, A), \quad \text{Sup}(s, A), \quad \text{and} \quad s < i.$$

Remark (Failure modes). Failure would put the greatest lower bound strictly above the least upper bound of the same nonempty set.

Remark (Failure mode decomposition). Choose $a \in A$. Then $i \leq a$ because i is a lower bound, and $a \leq s$ because s is an upper bound.

Order Comparison of Suprema

Proposition 3.2.22 (Order Comparison of Suprema). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded above. If every $x \in A$ is below some $y \in B$, then $\sup A \leq \sup B$.*

Go to proof.

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad B \neq \emptyset, \quad (\forall x \in A)(\exists y \in B)(x \leq y) \implies \sup A \leq \sup B.$$

Remark (Predicate reading).

$$[\forall x \in A \exists y \in B (x \leq y)] \wedge \text{Sup}(s_A, A) \wedge \text{Sup}(s_B, B) \implies s_A \leq s_B.$$

Remark (Negated quantified statement).

$$[\forall x \in A \exists y \in B (x \leq y)] \quad \text{and} \quad \sup B < \sup A.$$

Remark (Failure modes). Failure would mean A reaches higher in supremum than B even though each element of A is dominated by an element of B .

Remark (Failure mode decomposition). Every element of B lies below $\sup B$, so every element of A also lies below $\sup B$. Thus $\sup B$ is an upper bound for A .

Least Upper Bound Property Implies Existence of Suprema

Theorem 3.2.23 (Least Upper Bound Property Implies Existence of Suprema). *Let $S \subseteq \mathbb{R}$ be nonempty and bounded above. By the least upper bound property of the real numbers, there exists a real number s such that $s = \sup S$.*

Remark (Standard quantified statement).

$$S \neq \emptyset \quad \text{and} \quad (\exists u \in \mathbb{R})(\forall x \in S)(x \leq u) \implies (\exists s \in \mathbb{R}) \text{Sup}_{\mathbb{R}, \leq}(s, S).$$

Remark (Predicate reading).

$$S \neq \emptyset \wedge \text{BddAbove}_{\mathbb{R}, \leq}(S) \implies (\exists s \in \mathbb{R}) \text{Sup}_{\mathbb{R}, \leq}(s, S).$$

Remark (Negated quantified statement).

$$S \neq \emptyset, \quad \text{BddAbove}_{\mathbb{R}, \leq}(S), \quad \text{and} \quad (\forall s \in \mathbb{R}) \neg \text{Sup}_{\mathbb{R}, \leq}(s, S).$$

Remark (Failure modes). In $\mathbb{R}$ the theorem cannot fail once the set is nonempty and bounded above; outside complete ordered settings, the existence clause can fail.

Remark (Failure mode decomposition). The hypotheses separate domain nonemptiness from one-sided boundedness. The conclusion asserts existence of the sharp upper bound.

Greatest Lower Bound Property Implies Existence of Infima

Theorem 3.2.24 (Greatest Lower Bound Property Implies Existence of Infima). *Let $S \subseteq \mathbb{R}$ be nonempty and bounded below. Then there exists a real number i such that $i = \inf S$.*

Remark (Standard quantified statement).

$$S \neq \emptyset \quad \text{and} \quad (\exists \ell \in \mathbb{R})(\forall x \in S)(\ell \leq x) \implies (\exists i \in \mathbb{R}) \text{Inf}_{\mathbb{R}, \leq}(i, S).$$

Remark (Predicate reading).

$$S \neq \emptyset \wedge \text{BddBelow}_{\mathbb{R}, \leq}(S) \implies (\exists i \in \mathbb{R}) \text{Inf}_{\mathbb{R}, \leq}(i, S).$$

Remark (Negated quantified statement).

$$S \neq \emptyset, \quad \text{BddBelow}_{\mathbb{R}, \leq}(S), \quad \text{and} \quad (\forall i \in \mathbb{R}) \neg \text{Inf}_{\mathbb{R}, \leq}(i, S).$$

Remark (Failure modes). In $\mathbb{R}$ the theorem cannot fail once the set is nonempty and bounded below.

Remark (Failure mode decomposition). Apply the least upper bound property to $-S$ or argue by the order-dual form of completeness.

3.2.3 Maxima and Minima

Maxima and minima are attained sharp bounds. They are stronger than suprema and infima because the extremal element must belong to the nonempty set itself.




Figure 3.7: Maxima and minima are suprema and infima that are attained by the set.

Maximum

Definition 3.2.25 (Maximum). Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. An element $m \in S$ is the *maximum* of A if $m \in A$ and every element of A is less than or equal to m .

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad m \in A \quad \text{and} \quad (\forall x \in A)(x \leq m).$$

Remark (Definition predicate reading).

$$\text{Max}_{S, \leq}(m, A) := m \in A \wedge \text{UB}_{S, \leq}(m, A).$$

Remark (Negated quantified statement).

$$A = \emptyset \quad \text{or} \quad m \notin A \quad \text{or} \quad (\exists x \in A)(m < x).$$

Remark (Negation predicate reading).

$$\neg \text{Max}_{S,\leq}(m, A) \iff A = \emptyset \vee m \notin A \vee \neg \text{UB}_{S,\leq}(m, A).$$

Remark (Failure modes). The set may be empty; the candidate may not be a member of the set; or a larger element of the set may exist.

Remark (Failure mode decomposition). Maximum combines membership and an upper-bound condition. Failure of either component prevents attainment.

Minimum

Definition 3.2.26 (Minimum). Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. An element $m \in S$ is the *minimum* of A if $m \in A$ and every element of A is greater than or equal to m .

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad m \in A \quad \text{and} \quad (\forall x \in A)(m \leq x).$$

Remark (Definition predicate reading).

$$\text{Min}_{S,\leq}(m, A) := m \in A \wedge \text{LB}_{S,\leq}(m, A).$$

Remark (Negated quantified statement).

$$A = \emptyset \quad \text{or} \quad m \notin A \quad \text{or} \quad (\exists x \in A)(x < m).$$

Remark (Negation predicate reading).

$$\neg \text{Min}_{S,\leq}(m, A) \iff A = \emptyset \vee m \notin A \vee \neg \text{LB}_{S,\leq}(m, A).$$

Remark (Failure modes). The set may be empty; the candidate may not be a member of the set; or a smaller element of the set may exist.

Remark (Failure mode decomposition). Minimum combines membership and a lower-bound condition. Failure of either component prevents attainment.

Uniqueness of the Maximum

Proposition 3.2.27 (Uniqueness of the Maximum). Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. If $m, n \in S$ are both maxima of A , then $m = n$.

Go to proof.

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad \text{Max}_{S,\leq}(m, A) \quad \text{and} \quad \text{Max}_{S,\leq}(n, A) \implies m = n.$$

Remark (Predicate reading).

$$\text{Max}_{S,\leq}(m, A) \wedge \text{Max}_{S,\leq}(n, A) \implies m = n.$$

Remark (Negated quantified statement).

$$A \neq \emptyset \quad \text{and} \quad \text{Max}_{S,\leq}(m, A) \quad \text{and} \quad \text{Max}_{S,\leq}(n, A) \quad \text{and} \quad m \neq n.$$

Remark (Negation predicate reading).

$$\text{Max}_{S,\leq}(m, A) \wedge \text{Max}_{S,\leq}(n, A) \wedge m \neq n.$$

Remark (Failure modes). Failure would require two distinct elements of A each to dominate the entire set.

Remark (Failure mode decomposition). Since m is maximal over all elements and $n \in A$, we get $n \leq m$. Since n is maximal over all elements and $m \in A$, we get $m \leq n$.

Uniqueness of the Minimum

Proposition 3.2.28 (Uniqueness of the Minimum). *Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. If $m, n \in S$ are both minima of A , then $m = n$.*

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad \text{Min}_{S,\leq}(m, A) \quad \text{and} \quad \text{Min}_{S,\leq}(n, A) \quad \implies \quad m = n.$$

Remark (Predicate reading).

$$\text{Min}_{S,\leq}(m, A) \wedge \text{Min}_{S,\leq}(n, A) \implies m = n.$$

Remark (Negated quantified statement).

$$A \neq \emptyset \quad \text{and} \quad \text{Min}_{S,\leq}(m, A) \quad \text{and} \quad \text{Min}_{S,\leq}(n, A) \quad \text{and} \quad m \neq n.$$

Remark (Negation predicate reading).

$$\text{Min}_{S,\leq}(m, A) \wedge \text{Min}_{S,\leq}(n, A) \wedge m \neq n.$$

Remark (Failure modes). Failure would require two distinct elements of A each to lie below the entire set.

Remark (Failure mode decomposition). Since m is minimal and $n \in A$, we get $m \leq n$. Since n is minimal and $m \in A$, we get $n \leq m$.

Maximum Implies Supremum

Proposition 3.2.29 (Maximum Implies Supremum). *Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. If $m = \max A$, then $m = \sup A$.*

Go to proof.

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad \text{Max}_{S,\leq}(m, A) \quad \implies \quad \text{Sup}_{S,\leq}(m, A).$$

Remark (Predicate reading).

$$\text{Max}_{S,\leq}(m, A) \implies \text{Sup}_{S,\leq}(m, A).$$

Remark (Negated quantified statement).

$$\text{Max}_{S,\leq}(m, A) \quad \text{and} \quad \neg \text{Sup}_{S,\leq}(m, A).$$

Remark (Failure modes). Failure would require an attained upper bound that is not least among upper bounds.

Remark (Failure mode decomposition). Membership gives $m \in A$. Any upper bound of A must lie above every element of A , hence above m itself.

Minimum Implies Infimum

Proposition 3.2.30 (Minimum Implies Infimum). *Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. If $m = \min A$, then $m = \inf A$.*

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad \text{Min}_{S,\leq}(m, A) \implies \text{Inf}_{S,\leq}(m, A).$$

Remark (Predicate reading).

$$\text{Min}_{S,\leq}(m, A) \implies \text{Inf}_{S,\leq}(m, A).$$

Remark (Negated quantified statement).

$$\text{Min}_{S,\leq}(m, A) \quad \text{and} \quad \neg \text{Inf}_{S,\leq}(m, A).$$

Remark (Failure modes). Failure would require an attained lower bound that is not greatest among lower bounds.

Remark (Failure mode decomposition). Membership gives $m \in A$. Any lower bound of A must lie below every element of A , hence below m itself.

Supremum in the Set is the Maximum

Proposition 3.2.31 (Supremum in the Set is the Maximum). *Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. If $s = \sup A$ and $s \in A$, then $s = \max A$.*

Go to proof.

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad \text{Sup}_{S,\leq}(s, A) \quad \text{and} \quad s \in A \implies \text{Max}_{S,\leq}(s, A).$$

Remark (Predicate reading).

$$\text{Sup}_{S,\leq}(s, A) \wedge s \in A \implies \text{Max}_{S,\leq}(s, A).$$

Remark (Negated quantified statement).

$$\text{Sup}_{S,\leq}(s, A) \quad \text{and} \quad s \in A \quad \text{and} \quad \neg \text{Max}_{S,\leq}(s, A).$$

Remark (Failure modes). Failure would require a supremum that belongs to the set but is not its maximum.

Remark (Failure mode decomposition). The supremum condition supplies the upper-bound clause. Membership supplies attainment.

Infimum in the Set is the Minimum

Proposition 3.2.32 (Infimum in the Set is the Minimum). *Let $(S, \leq)$ be an ordered set and let $A \subseteq S$ be nonempty. If $i = \inf A$ and $i \in A$, then $i = \min A$.*

Remark (Standard quantified statement).

$$A \neq \emptyset \quad \text{and} \quad \text{Inf}_{S,\leq}(i, A) \quad \text{and} \quad i \in A \quad \implies \quad \text{Min}_{S,\leq}(i, A).$$

Remark (Predicate reading).

$$\text{Inf}_{S,\leq}(i, A) \wedge i \in A \implies \text{Min}_{S,\leq}(i, A).$$

Remark (Negated quantified statement).

$$\text{Inf}_{S,\leq}(i, A) \quad \text{and} \quad i \in A \quad \text{and} \quad \neg \text{Min}_{S,\leq}(i, A).$$

Remark (Failure modes). Failure would require an infimum that belongs to the set but is not its minimum.

Remark (Failure mode decomposition). The infimum condition supplies the lower-bound clause. Membership supplies attainment.

3.2.4 Epsilon Characterization

ε -Characterization of Supremum

Definition (Epsilon Characterization of Supremum)

Definition 3.2.33 (Epsilon Characterization of Supremum). Let $A \subseteq \mathbb{R}$ and let $s \in \mathbb{R}$. The element s satisfies the *epsilon characterization of the supremum* of A if and only if s is an upper bound of A and for every $\varepsilon > 0$ there exists $a \in A$ such that $s - \varepsilon < a$.

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } A \subseteq \mathbb{R} \text{ and } s \in \mathbb{R}. \\ &s \text{ satisfies the epsilon characterization of } \sup A \\ &\iff \left[\forall x \in A (x \leq s) \right] \wedge \left[\forall \varepsilon > 0 \exists a \in A (s - \varepsilon < a) \right]. \end{aligned}$$

Remark (Definition predicate reading).

$$\text{EpsilonSupremum}(s, A) := \text{UpperBound}(s, A) \wedge \forall \varepsilon > 0 \exists a \in A (s - \varepsilon < a).$$

Remark (Negated quantified statement).

Let $A \subseteq \mathbb{R}$ and $s \in \mathbb{R}$.

s does not satisfy the epsilon characterization of $\sup A$

$$\iff \left[\exists x \in A (s < x) \right] \vee \left[\exists \varepsilon_0 > 0 \forall a \in A (a \leq s - \varepsilon_0) \right].$$

Remark (Negation predicate reading).

$$\neg \text{EpsilonSupremum}(s, A) \iff \left[\exists x \in A (s < x) \right] \vee \left[\exists \varepsilon_0 > 0 \forall a \in A (a \leq s - \varepsilon_0) \right].$$

Remark (Failure modes). The characterization fails either because s is not an upper bound of A , or because A remains at least some positive distance below s .

Remark (Failure mode decomposition).

$$\neg \text{EpsilonSupremum}(s, A) = \underbrace{\exists x \in A (s < x)}_{\text{upper-bound failure}} \vee \underbrace{\exists \varepsilon_0 > 0 \forall a \in A (a \leq s - \varepsilon_0)}_{\text{positive gap below } s}.$$

Remark (Interpretation). This condition says that s caps the set from above while points of A approach s from below at every positive scale. The upper-bound clause prevents A from crossing above s , and the epsilon clause prevents s from floating strictly above the set. Geometrically, s is a ceiling that the set presses against arbitrarily closely.

Remark (Dependencies).

[Definition 3.2.2.](#)




Figure 3.8: Refined visualization of the ε -characterization. Labels occupy staggered vertical lanes with leader lines to eliminate all horizontal collision, while the dashed spine preserves the geometric correspondence between the real line and the probe below it.

Definition (Epsilon Characterization of Infimum)

Definition 3.2.34 (Epsilon Characterization of Infimum). Let $A \subseteq \mathbb{R}$ and let $t \in \mathbb{R}$. The element t satisfies the *epsilon characterization of the infimum* of A if and only if t is a lower bound of A and for every $\varepsilon > 0$ there exists $a \in A$ such that $a < t + \varepsilon$.

Remark (Standard quantified statement).

$$\begin{aligned} & \text{Let } A \subseteq \mathbb{R} \text{ and } t \in \mathbb{R}. \\ & t \text{ satisfies the epsilon characterization of } \inf A \\ & \iff \left[\forall x \in A (t \leq x) \right] \wedge \left[\forall \varepsilon > 0 \exists a \in A (a < t + \varepsilon) \right]. \end{aligned}$$

Remark (Definition predicate reading).

$$\text{EpsilonInfimum}(t, A) := \text{LowerBound}(t, A) \wedge \forall \varepsilon > 0 \exists a \in A (a < t + \varepsilon).$$

Remark (Negated quantified statement).

$$\begin{aligned} & \text{Let } A \subseteq \mathbb{R} \text{ and } t \in \mathbb{R}. \\ & t \text{ does not satisfy the epsilon characterization of } \inf A \\ & \iff \left[\exists x \in A (x < t) \right] \vee \left[\exists \varepsilon_0 > 0 \forall a \in A (t + \varepsilon_0 \leq a) \right]. \end{aligned}$$

Remark (Negation predicate reading).

$$\neg \text{EpsilonInfimum}(t, A) \iff \left[\exists x \in A (x < t) \right] \vee \left[\exists \varepsilon_0 > 0 \forall a \in A (t + \varepsilon_0 \leq a) \right].$$

Remark (Failure modes). The characterization fails either because t is not a lower bound of A , or because A remains at least some positive distance above t .

Remark (Failure mode decomposition).


$$\neg \text{EpsilonInfimum}(t, A) = \underbrace{\exists x \in A (x < t)}_{\text{lower-bound failure}} \vee \underbrace{\exists \varepsilon_0 > 0 \forall a \in A (t + \varepsilon_0 \leq a)}_{\text{positive gap above } t}.$$

Remark (Interpretation). This condition says that t sits below the set while points of A approach t from above at every positive scale. The lower-bound clause prevents A from crossing below t , and the epsilon clause prevents t from sitting strictly below the set with a positive gap. Geometrically, t is a floor that the set presses against arbitrarily closely from above.

Remark (Dependencies).

[Definition 3.2.3.](#)

3.2.5 Extremal Bounds Summary



Remark (Equivalent formulations).

Over the ordered-field axioms, completeness is equivalent to each of the following:

- Every nonempty set bounded below has an infimum.
- Every Cauchy sequence in $\mathbb{R}$ converges.
- (*Nested Interval Property*) Every nested sequence of nonempty closed bounded intervals has nonempty intersection.

3.3 Algebra of Bounds

3.3.1 Set Operations and Bounds

Remark (Orientation). Set operations change the collection of possible counterexamples to a proposed bound. Each result below isolates one operation and one direction of bound transport.

Lemma (Union Preserves Upper Bounds)

Lemma 3.3.1 (Union Preserves Upper Bounds). *Let $A, B \subseteq \mathbb{R}$. If u_A is an upper bound of A and u_B is an upper bound of B , then $\max\{u_A, u_B\}$ is an upper bound of $A \cup B$. [Go to proof.](#)*

Remark (Standard quantified statement).

$$(\forall a \in A)(a \leq u_A) \wedge (\forall b \in B)(b \leq u_B) \implies (\forall x \in A \cup B)(x \leq \max\{u_A, u_B\}).$$

Remark (Predicate reading).

$$\text{UB}(u_A, A) \wedge \text{UB}(u_B, B) \implies \text{UB}(\max\{u_A, u_B\}, A \cup B).$$

Remark (Negated quantified statement).

$$(\forall a \in A)(a \leq u_A), \quad (\forall b \in B)(b \leq u_B), \quad (\exists x \in A \cup B)(\max\{u_A, u_B\} < x).$$

Remark (Negation predicate reading).

$$\text{UB}(u_A, A) \wedge \text{UB}(u_B, B) \wedge \neg \text{UB}(\max\{u_A, u_B\}, A \cup B).$$

Remark (Failure modes). A failure would require a union element larger than both proposed upper bounds.

Remark (Failure mode decomposition). Every $x \in A \cup B$ lies in A or in B . In the first case $x \leq u_A \leq \max\{u_A, u_B\}$; in the second case $x \leq u_B \leq \max\{u_A, u_B\}$.

Remark (Dependencies).

[Upper Bound](#).

Lemma (Union Preserves Lower Bounds)

Lemma 3.3.2 (Union Preserves Lower Bounds). *Let $A, B \subseteq \mathbb{R}$. If ℓ_A is a lower bound of A and ℓ_B is a lower bound of B , then $\min\{\ell_A, \ell_B\}$ is a lower bound of $A \cup B$. [Go to proof](#).*

Remark (Standard quantified statement).

$$(\forall a \in A)(\ell_A \leq a) \wedge (\forall b \in B)(\ell_B \leq b) \implies (\forall x \in A \cup B)(\min\{\ell_A, \ell_B\} \leq x).$$

Remark (Predicate reading).

$$\text{LB}(\ell_A, A) \wedge \text{LB}(\ell_B, B) \implies \text{LB}(\min\{\ell_A, \ell_B\}, A \cup B).$$

Remark (Negated quantified statement).

$$(\forall a \in A)(\ell_A \leq a), \quad (\forall b \in B)(\ell_B \leq b), \quad (\exists x \in A \cup B)(x < \min\{\ell_A, \ell_B\}).$$

Remark (Negation predicate reading).

$$\text{LB}(\ell_A, A) \wedge \text{LB}(\ell_B, B) \wedge \neg \text{LB}(\min\{\ell_A, \ell_B\}, A \cup B).$$

Remark (Failure modes). A failure would require a union element smaller than both proposed lower bounds.

Remark (Failure mode decomposition). Every $x \in A \cup B$ lies in A or in B . In either case the appropriate lower bound lies below x , and $\min\{\ell_A, \ell_B\}$ lies below that lower bound.

Remark (Dependencies).

Lower Bound.

Proposition (Union Is Bounded Above Exactly When Both Pieces Are)

Proposition 3.3.3 (Union Is Bounded Above Exactly When Both Pieces Are). *Let $A, B \subseteq \mathbb{R}$ be nonempty. Then $A \cup B$ is bounded above if and only if A is bounded above and B is bounded above. [Go to proof.](#)*

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad B \neq \emptyset \quad \implies \quad \text{BddAbove}(A \cup B) \iff \text{BddAbove}(A) \wedge \text{BddAbove}(B).$$

Remark (Predicate reading).

$$\text{BddAbove}(A \cup B) \iff \text{BddAbove}(A) \wedge \text{BddAbove}(B).$$

Remark (Negated quantified statement). The equivalence fails precisely if one side holds and the other does not.

Remark (Negation predicate reading).

$$\text{BddAbove}(A \cup B) \oplus (\text{BddAbove}(A) \wedge \text{BddAbove}(B)).$$

Remark (Failure modes). The forward direction could fail only if a bound for the union did not bound a piece. The reverse direction could fail only if bounds for both pieces did not combine into a bound for the union.

Remark (Failure mode decomposition). Use subset inheritance for the forward direction and [Union Preserves Upper Bounds](#) for the reverse direction.

Remark (Dependencies).

Bounded Above; Union Preserves Upper Bounds; Subsets Preserve Upper Bounds.

Proposition (Union Is Bounded Below Exactly When Both Pieces Are)

Proposition 3.3.4 (Union Is Bounded Below Exactly When Both Pieces Are). *Let $A, B \subseteq \mathbb{R}$ be nonempty. Then $A \cup B$ is bounded below if and only if A is bounded below and B is bounded below. [Go to proof.](#)*

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad B \neq \emptyset \quad \implies \quad \text{BddBelow}(A \cup B) \iff \text{BddBelow}(A) \wedge \text{BddBelow}(B).$$

Remark (Predicate reading).

$$\text{BddBelow}(A \cup B) \iff \text{BddBelow}(A) \wedge \text{BddBelow}(B).$$

Remark (Negated quantified statement). The equivalence fails precisely if one side holds and the other does not.

Remark (Negation predicate reading).

$$\text{BddBelow}(A \cup B) \oplus (\text{BddBelow}(A) \wedge \text{BddBelow}(B)).$$

Remark (Failure modes). The forward direction could fail only if a lower bound for the union did not bound a piece. The reverse direction could fail only if lower bounds for both pieces did not combine into a lower bound for the union.

Remark (Failure mode decomposition). Use subset inheritance for the forward direction and [Union Preserves Lower Bounds](#) for the reverse direction.

Remark (Dependencies).

[Bounded Below](#); [Union Preserves Lower Bounds](#); [Subsets Preserve Lower Bounds](#).

Corollary (Union Is Bounded Exactly When Both Pieces Are)

Corollary 3.3.5 (Union Is Bounded Exactly When Both Pieces Are). *Let $A, B \subseteq \mathbb{R}$ be nonempty. Then $A \cup B$ is bounded if and only if A is bounded and B is bounded. [Go to proof](#).*

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad B \neq \emptyset \quad \implies \quad \text{Bdd}(A \cup B) \iff \text{Bdd}(A) \wedge \text{Bdd}(B).$$

Remark (Predicate reading).

$$\text{Bdd}(A \cup B) \iff \text{Bdd}(A) \wedge \text{Bdd}(B).$$

Remark (Negated quantified statement). The two-sided boundedness equivalence fails if either the upper or lower one-sided equivalence fails.

Remark (Negation predicate reading).

$$\text{Bdd}(A \cup B) \oplus (\text{Bdd}(A) \wedge \text{Bdd}(B)).$$

Remark (Failure modes). Two-sided boundedness fails exactly through an upper-bound failure or a lower-bound failure.

Remark (Failure mode decomposition). Apply the bounded-above and bounded-below union equivalences separately.

Remark (Dependencies).

[Bounded Set](#); [Union Is Bounded Above Exactly When Both Pieces Are](#); [Union Is Bounded Below Exactly When Both Pieces Are](#).

Lemma (Subsets Preserve Upper Bounds)

Lemma 3.3.6 (Subsets Preserve Upper Bounds). *Let $C \subseteq A \subseteq \mathbb{R}$. If u is an upper bound of A , then u is an upper bound of C . [Go to proof](#).*

Remark (Standard quantified statement).

$$C \subseteq A \wedge (\forall a \in A)(a \leq u) \implies (\forall c \in C)(c \leq u).$$

Remark (Predicate reading).

$$C \subseteq A \wedge \text{UB}(u, A) \implies \text{UB}(u, C).$$

Remark (Negated quantified statement).

$$C \subseteq A, \quad (\forall a \in A)(a \leq u), \quad (\exists c \in C)(u < c).$$

Remark (Negation predicate reading).

$$C \subseteq A \wedge \text{UB}(u, A) \wedge \neg \text{UB}(u, C).$$

Remark (Failure modes). A failure would require a counterexample in C that is not already a counterexample in A .

Remark (Failure mode decomposition). Since $C \subseteq A$, every $c \in C$ is an element of A . The upper-bound condition on A therefore applies directly to c .

Remark (Dependencies).

Upper Bound.

Lemma (Subsets Preserve Lower Bounds)

Lemma 3.3.7 (Subsets Preserve Lower Bounds). *Let $C \subseteq A \subseteq \mathbb{R}$. If ℓ is a lower bound of A , then ℓ is a lower bound of C . [Go to proof.](#)*

Remark (Standard quantified statement).

$$C \subseteq A \wedge (\forall a \in A)(\ell \leq a) \implies (\forall c \in C)(\ell \leq c).$$

Remark (Predicate reading).

$$C \subseteq A \wedge \text{LB}(\ell, A) \implies \text{LB}(\ell, C).$$

Remark (Negated quantified statement).

$$C \subseteq A, \quad (\forall a \in A)(\ell \leq a), \quad (\exists c \in C)(c < \ell).$$

Remark (Negation predicate reading).

$$C \subseteq A \wedge \text{LB}(\ell, A) \wedge \neg \text{LB}(\ell, C).$$

Remark (Failure modes). A failure would require a point in C below ℓ , hence a point in A below ℓ .

Remark (Failure mode decomposition). Subset inheritance removes possible counterexamples; it cannot create a new one.

Remark (Dependencies).

Lower Bound.

Corollary (Intersections Inherit Bounds)

Corollary 3.3.8 (Intersections Inherit Bounds). *Let $A, B \subseteq \mathbb{R}$. Every upper bound of A is an upper bound of $A \cap B$, and every lower bound of A is a lower bound of $A \cap B$. [Go to proof.](#)*

Remark (Standard quantified statement).

$$A \cap B \subseteq A.$$

Remark (Predicate reading).

$$\text{UB}(u, A) \implies \text{UB}(u, A \cap B),$$

and similarly for lower bounds.

Remark (Negated quantified statement). A failure would be a failure of subset inheritance for $A \cap B \subseteq A$.

Remark (Negation predicate reading).

$$\text{UB}(u, A) \wedge \neg \text{UB}(u, A \cap B)$$

is contradictory, and the same holds for lower bounds.

Remark (Failure modes). Intersection can lose elements, but it cannot introduce an element outside the parent set A .

Remark (Failure mode decomposition). Apply subset inheritance to $C = A \cap B$.

Remark (Dependencies).

Subsets Preserve Upper Bounds; Subsets Preserve Lower Bounds.

Corollary (Differences Inherit Bounds)

Corollary 3.3.9 (Differences Inherit Bounds). *Let $A, B \subseteq \mathbb{R}$. Every upper bound of A is an upper bound of $A \setminus B$, and every lower bound of A is a lower bound of $A \setminus B$. [Go to proof.](#)*

Remark (Standard quantified statement).

$$A \setminus B \subseteq A.$$

Remark (Predicate reading).

$$\text{UB}(u, A) \implies \text{UB}(u, A \setminus B),$$

and similarly for lower bounds.

Remark (Negated quantified statement). A failure would be a failure of subset inheritance for $A \setminus B \subseteq A$.

Remark (Negation predicate reading).

$$\text{UB}(u, A) \wedge \neg \text{UB}(u, A \setminus B)$$

is contradictory, and the same holds for lower bounds.

Remark (Failure modes). Difference can remove elements from A , but it cannot create a new element outside A .

Remark (Failure mode decomposition). Apply subset inheritance to $C = A \setminus B$.

Remark (Dependencies).

[Subsets Preserve Upper Bounds](#); [Subsets Preserve Lower Bounds](#).

Corollary (Complements Inherit Ambient Bounds)

Corollary 3.3.10 (Complements Inherit Ambient Bounds). *Let $A \subseteq E \subseteq \mathbb{R}$, and take complements relative to E . Every upper bound of E is an upper bound of $E \setminus A$, and every lower bound of E is a lower bound of $E \setminus A$. [Go to proof](#).*

Remark (Standard quantified statement).

$$E \setminus A \subseteq E.$$

Remark (Predicate reading).

$$\text{UB}(u, E) \implies \text{UB}(u, E \setminus A), \quad \text{LB}(\ell, E) \implies \text{LB}(\ell, E \setminus A).$$

Remark (Negated quantified statement).

$$\text{UB}(u, E) \wedge \neg \text{UB}(u, E \setminus A)$$

or

$$\text{LB}(\ell, E) \wedge \neg \text{LB}(\ell, E \setminus A).$$

Remark (Negation predicate reading). Both negated configurations contradict subset inheritance because $E \setminus A \subseteq E$.

Remark (Failure modes). The complement does not inherit bounds from A ; it inherits them from the ambient set E .

Remark (Failure mode decomposition). Every point of $E \setminus A$ is a point of E , so every ambient bound of E applies to the complement.

Remark (Dependencies).

[Subsets Preserve Upper Bounds](#); [Subsets Preserve Lower Bounds](#).

3.3.2 Lattice Operations and Bounds

Remark (Orientation). In this section, *lattice* refers to the max–min structure already present in the usual order on $\mathbb{R}$. For two real numbers, their join is their maximum and their meet is their minimum:

$$x \vee y = \max\{x, y\}, \quad x \wedge y = \min\{x, y\}.$$

Thus lattice operations are not a new order relation; they are the operations created by the order. The set-level constructions below apply those pairwise operations to all pairs of points from two nonempty sets.

Definition (Pointwise Maximum and Minimum Sets)

Definition 3.3.11 (Pointwise Maximum and Minimum Sets). Let $A, B \subseteq \mathbb{R}$ be nonempty. Define

$$\max(A, B) := \{\max\{a, b\} : a \in A, b \in B\},$$

and

$$\min(A, B) := \{\min\{a, b\} : a \in A, b \in B\}.$$

Remark (Standard quantified statement).

$$x \in \max(A, B) \iff (\exists a \in A)(\exists b \in B)(x = \max\{a, b\}),$$

and similarly for $\min(A, B)$ with $\min\{a, b\}$.

Remark (Definition predicate reading).

$$\text{PointwiseMaxSet}(A, B) = \max(A, B), \quad \text{PointwiseMinSet}(A, B) = \min(A, B).$$

Remark (Negated quantified statement).

$$x \notin \max(A, B) \iff (\forall a \in A)(\forall b \in B)(x \neq \max\{a, b\}),$$

and similarly for $\min(A, B)$.

Remark (Negation predicate reading).

$$\neg \text{PointwiseMaxSet}(A, B) = C \quad \text{means} \quad (\exists x)[x \in C \triangle \max(A, B)],$$

and similarly for PointwiseMinSet .

Remark (Failure modes). The definition fails only if the image set omits a pairwise maximum or minimum, or if it includes a value not produced by one of the pairwise rules.

Remark (Failure mode decomposition). Every membership claim decomposes into two input witnesses and one order-based operation: either the larger input or the smaller input is selected.

Remark (Dependencies).

[Maximum](#); [Minimum](#).

Theorem (Supremum of the Pointwise Maximum Set)

Theorem 3.3.12 (Supremum of the Pointwise Maximum Set). Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded above. Then

$$\sup \max(A, B) = \max\{\sup A, \sup B\}.$$

[Go to proof](#).

Remark (Standard quantified statement).

$$A, B \neq \emptyset, \quad A, B \text{ bounded above} \implies \sup\{\max\{a, b\} : a \in A, b \in B\} = \max\{\sup A, \sup B\}. \blacksquare$$

Remark (Predicate reading).

$$\text{Supremum}(s_A, A) \wedge \text{Supremum}(s_B, B) \implies \text{Supremum}(\max\{s_A, s_B\}, \max(A, B)).$$

Remark (Negated quantified statement).

$$\sup \max(A, B) \neq \max\{\sup A, \sup B\}$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Supremum}(s_A, A) \wedge \text{Supremum}(s_B, B) \wedge \neg \text{Supremum}(\max\{s_A, s_B\}, \max(A, B)).$$

Remark (Failure modes). The formula could fail only if the maximum of the two suprema failed to bound all pairwise maxima, or if it were not the least such bound.

Remark (Failure mode decomposition). Every pairwise maximum is bounded by $\max\{\sup A, \sup B\}$ ■ Sharpness comes from holding one input fixed in the nonempty other set and letting the relevant set approach its own supremum.

Remark (Dependencies).

[Supremum](#); [Pointwise Maximum and Minimum Sets](#).

Theorem (Infimum of the Pointwise Maximum Set)

Theorem 3.3.13 (Infimum of the Pointwise Maximum Set). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded below. Then*

$$\inf \max(A, B) = \max\{\inf A, \inf B\}.$$

[Go to proof](#).

Remark (Standard quantified statement).

$$A, B \neq \emptyset, \quad A, B \text{ bounded below} \implies \inf\{\max\{a, b\} : a \in A, b \in B\} = \max\{\inf A, \inf B\}. \blacksquare$$

Remark (Predicate reading).

$$\text{Infimum}(t_A, A) \wedge \text{Infimum}(t_B, B) \implies \text{Infimum}(\max\{t_A, t_B\}, \max(A, B)).$$

Remark (Negated quantified statement).

$$\inf \max(A, B) \neq \max\{\inf A, \inf B\}$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Infimum}(t_A, A) \wedge \text{Infimum}(t_B, B) \wedge \neg \text{Infimum}(\max\{t_A, t_B\}, \max(A, B)).$$

Remark (Failure modes). The formula could fail only if the larger of the two lower thresholds were not a lower bound for all pairwise maxima, or if pairwise maxima could not approach that threshold from above.

Remark (Failure mode decomposition). Every pairwise maximum is at least both lower thresholds, hence at least their maximum. Approximation of the two infima gives pairwise maxima arbitrarily close to the larger lower threshold.

Remark (Dependencies).

[Infimum; Pointwise Maximum and Minimum Sets.](#)

Theorem (Supremum of the Pointwise Minimum Set)

Theorem 3.3.14 (Supremum of the Pointwise Minimum Set). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded above. Then*

$$\sup \min(A, B) = \min\{\sup A, \sup B\}.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$A, B \neq \emptyset, \quad A, B \text{ bounded above} \implies \sup\{\min\{a, b\} : a \in A, b \in B\} = \min\{\sup A, \sup B\}. \blacksquare$$

Remark (Predicate reading).

$$\text{Supremum}(s_A, A) \wedge \text{Supremum}(s_B, B) \implies \text{Supremum}(\min\{s_A, s_B\}, \min(A, B)).$$

Remark (Negated quantified statement).

$$\sup \min(A, B) \neq \min\{\sup A, \sup B\}$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Supremum}(s_A, A) \wedge \text{Supremum}(s_B, B) \wedge \neg \text{Supremum}(\min\{s_A, s_B\}, \min(A, B)).$$

Remark (Failure modes). The formula could fail only if the smaller supremum failed to bound all pairwise minima, or if that upper bound were not sharp.

Remark (Failure mode decomposition). Every pairwise minimum is below both suprema, hence below their minimum. Sharpness comes from taking both inputs near their own suprema.

Remark (Dependencies).

[Supremum; Pointwise Maximum and Minimum Sets.](#)

Theorem (Infimum of the Pointwise Minimum Set)

Theorem 3.3.15 (Infimum of the Pointwise Minimum Set). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded below. Then*

$$\inf \min(A, B) = \min\{\inf A, \inf B\}.$$

Go to proof.

Remark (Standard quantified statement).

$$A, B \neq \emptyset, \quad A, B \text{ bounded below} \implies \inf\{\min\{a, b\} : a \in A, b \in B\} = \min\{\inf A, \inf B\}. \blacksquare$$

Remark (Predicate reading).

$$\text{Infimum}(t_A, A) \wedge \text{Infimum}(t_B, B) \implies \text{Infimum}(\min\{t_A, t_B\}, \min(A, B)).$$

Remark (Negated quantified statement).

$$\inf \min(A, B) \neq \min\{\inf A, \inf B\}$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Infimum}(t_A, A) \wedge \text{Infimum}(t_B, B) \wedge \neg \text{Infimum}(\min\{t_A, t_B\}, \min(A, B)).$$

Remark (Failure modes). The formula could fail only if the smaller infimum failed to bound all pairwise minima from below, or if that lower bound were not sharp.

Remark (Failure mode decomposition). Every pairwise minimum is at least the smaller of the two infima. Sharpness comes from holding one set nonempty and approaching the smaller lower threshold.

Remark (Dependencies).

[Infimum](#); [Pointwise Maximum and Minimum Sets](#).

3.3.3 Images, Inverse Images, and Extrema

Remark (Orientation). One cannot usually move a supremum or infimum through an arbitrary function:

$$\sup f(A) \neq f(\sup A)$$


in general. Equality is an order-and-continuity statement. Monotone functions preserve or reverse the relevant inequalities, and continuity at the endpoint turns the endpoint bound into a sharp bound for the image. The inverse function statements are the same principle applied to f^{-1} .

Remark (Geometric meaning of the hypotheses). Think of a nonempty bounded set $A \subseteq \mathbb{R}$ as a cloud of points on the real line. The number $\sup A$ is the right-hand wall of that cloud: it may or may not belong to A , but points of A press arbitrarily close to it from the left. Applying a function f moves every point of the cloud and produces the image cloud

$$f(A) = \{f(a) : a \in A\}.$$


The question $\sup f(A) = f(\sup A)$ asks whether the function carries the right-hand wall of the original cloud to the right-hand wall of the image cloud.

Example (Increasing images keep the right edge on the right).




When f is increasing, order is preserved. A point that starts farther to the right cannot land to the left of a point that starts farther to the left. Thus the right edge of the image cloud is controlled by the right edge of the original cloud, provided continuity prevents a jump at the endpoint.

Example (Decreasing images swap the endpoint roles).



When f is decreasing, order is reversed. The left edge of the original cloud becomes the right edge of the image cloud, and the right edge of the original cloud becomes the left edge of the image cloud. This is why decreasing image theorems exchange suprema and infima.

Example (Nonmonotone images can cross).



If f is not monotone on the relevant domain, the arrows can cross. Then the right edge of $f(A)$ need not come from $\sup A$ or from $\inf A$ alone in a uniform way. The image edge may be produced by an interior point, by the left branch of a piecewise monotone function, or by the right branch. In that situation the set must be split into pieces where f is monotone, and the image pieces must be recombined using the algebra of suprema over unions.

Example (Why $x \mapsto x^2$ does not fit either global theorem). The function $f(x) = x^2$ is not increasing on all of $\mathbb{R}$. Indeed, $-2 \leq 1$, but

$$(-2)^2 = 4 > 1 = 1^2.$$

It is also not decreasing on all of $\mathbb{R}$. Indeed, $1 \leq 2$, but

$$1^2 = 1 < 4 = 2^2.$$

Thus neither the increasing image theorem nor the decreasing image theorem can be applied globally to $x \mapsto x^2$ on $\mathbb{R}$.

Let

$$A = (-1, 2).$$

Then $\inf A = -1$ and $\sup A = 2$. The image under the square map is

$$f(A) = \{x^2 : -1 < x < 2\} = [0, 4),$$

so

$$\sup f(A) = 4.$$

The increasing-theorem expression gives

$$f(\sup A) = f(2) = 4,$$

which happens to agree in this example. But the agreement is not justified by global increasingness, because x^2 is not globally increasing.

The decreasing-theorem expression gives

$$f(\inf A) = f(-1) = 1,$$

and this does not agree:

$$\sup f(A) = 4 \neq 1 = f(\inf A).$$

The failure occurs because the square map folds the real line at 0. It is decreasing on $(-\infty, 0]$ and increasing on $[0, \infty)$, so a set that straddles 0 must be analyzed branch by branch. For a nonempty bounded set $A \subseteq \mathbb{R}$ with finite endpoints $\alpha = \inf A$ and $\beta = \sup A$, the largest square is controlled by whichever endpoint is farther from 0:

$$\sup\{x^2 : x \in A\} = \max\{\alpha^2, \beta^2\}.$$

Theorem (Increasing Image Preserves Suprema)

Theorem 3.3.16 (Increasing Image Preserves Suprema). *Let $I \subseteq \mathbb{R}$, let $A \subseteq I$ be nonempty and bounded above, and let $f : I \rightarrow \mathbb{R}$ be increasing. Suppose $s = \sup A$ belongs to I and f is continuous at s . Then*

$$\sup f(A) = f(\sup A), \quad f(A) = \{f(a) : a \in A\}.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad s = \sup A \in I, \quad f \text{ increasing and continuous at } s \implies \sup\{f(a) : a \in A\} = f(s).$$

Remark (Predicate reading).

$$\text{Supremum}(s, A) \wedge \text{Increasing}(f, I) \wedge \text{ContinuousAt}(f, s) \implies \text{Supremum}(f(s), f(A)).$$

Remark (Negated quantified statement).

$$\sup f(A) \neq f(\sup A)$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Supremum}(s, A) \wedge \text{Increasing}(f, I) \wedge \text{ContinuousAt}(f, s) \wedge \neg \text{Supremum}(f(s), f(A)).$$

Remark (Failure modes). The equality can fail if f is not increasing, if $\sup A$ is not in the domain, or if the endpoint value of f is not approached by values $f(a)$ with $a \in A$.

Remark (Failure mode decomposition). Increasingness gives $f(a) \leq f(s)$ for every $a \in A$. The defining approximation property of the supremum gives points of A arbitrarily close to s from below, and continuity at s makes their images approach $f(s)$.

Remark (Dependencies).

[Supremum](#); [Increasing Function](#); [Continuity at a Point](#).

Theorem (Increasing Image Preserves Infima)

Theorem 3.3.17 (Increasing Image Preserves Infima). *Let $I \subseteq \mathbb{R}$, let $A \subseteq I$ be nonempty and bounded below, and let $f : I \rightarrow \mathbb{R}$ be increasing. Suppose $t = \inf A$ belongs to I and f is continuous at t . Then*

$$\inf f(A) = f(\inf A).$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad t = \inf A \in I, \quad f \text{ increasing and continuous at } t \implies \inf\{f(a) : a \in A\} = f(t).$$

Remark (Predicate reading).

$$\text{Infimum}(t, A) \wedge \text{Increasing}(f, I) \wedge \text{ContinuousAt}(f, t) \implies \text{Infimum}(f(t), f(A)).$$

Remark (Negated quantified statement).

$$\inf f(A) \neq f(\inf A)$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Infimum}(t, A) \wedge \text{Increasing}(f, I) \wedge \text{ContinuousAt}(f, t) \wedge \neg \text{Infimum}(f(t), f(A)).$$

Remark (Failure modes). The equality can fail if f is not increasing, if $\inf A$ is not in the domain, or if the endpoint value is not reached as a limit of image values.

Remark (Failure mode decomposition). Increasingness gives $f(t) \leq f(a)$ for every $a \in A$. Approximation of the infimum from above and continuity at t make this lower bound sharp.

Remark (Dependencies).

[Infimum](#); [Increasing Function](#); [Continuity at a Point](#).

Theorem (Decreasing Image Sends Infimum to Supremum)

Theorem 3.3.18 (Decreasing Image Sends Infimum to Supremum). *Let $I \subseteq \mathbb{R}$, let $A \subseteq I$ be nonempty and bounded below, and let $f : I \rightarrow \mathbb{R}$ be decreasing. Suppose $t = \inf A$ belongs to I and f is continuous at t . Then*

$$\sup f(A) = f(\inf A).$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad t = \inf A \in I, \quad f \text{ decreasing and continuous at } t \implies \sup\{f(a) : a \in A\} = f(t).$$

Remark (Predicate reading).

$$\text{Infimum}(t, A) \wedge \text{Decreasing}(f, I) \wedge \text{ContinuousAt}(f, t) \implies \text{Supremum}(f(t), f(A)).$$

Remark (Negated quantified statement).

$$\sup f(A) \neq f(\inf A)$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Infimum}(t, A) \wedge \text{Decreasing}(f, I) \wedge \text{ContinuousAt}(f, t) \wedge \neg \text{Supremum}(f(t), f(A)).$$

Remark (Failure modes). The equality can fail if f is not decreasing, if $\inf A$ is outside the domain, or if f is not continuous at the lower endpoint.

Remark (Failure mode decomposition). Decreasingness gives $f(a) \leq f(t)$ for every $a \in A$. Approximation of t from above and continuity at t make this upper bound sharp.

Remark (Dependencies).

[Infimum](#); [Supremum](#); [Decreasing Function](#); [Continuity at a Point](#).

Theorem (Decreasing Image Sends Supremum to Infimum)

Theorem 3.3.19 (Decreasing Image Sends Supremum to Infimum). *Let $I \subseteq \mathbb{R}$, let $A \subseteq I$ be nonempty and bounded above, and let $f : I \rightarrow \mathbb{R}$ be decreasing. Suppose $s = \sup A$ belongs to I and f is continuous at s . Then*

$$\inf f(A) = f(\sup A).$$

[Go to proof](#).

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad s = \sup A \in I, \quad f \text{ decreasing and continuous at } s \implies \inf\{f(a) : a \in A\} = f(s).$$

Remark (Predicate reading).

$$\text{Supremum}(s, A) \wedge \text{Decreasing}(f, I) \wedge \text{ContinuousAt}(f, s) \implies \text{Infimum}(f(s), f(A)).$$

Remark (Negated quantified statement).

$$\inf f(A) \neq f(\sup A)$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Supremum}(s, A) \wedge \text{Decreasing}(f, I) \wedge \text{ContinuousAt}(f, s) \wedge \neg \text{Infimum}(f(s), f(A)).$$

Remark (Failure modes). The equality can fail if f is not decreasing, if $\sup A$ is outside the domain, or if f is not continuous at the upper endpoint.

Remark (Failure mode decomposition). Decreasingness gives $f(s) \leq f(a)$ for every $a \in A$. Approximation of s from below and continuity at s make this lower bound sharp.

Remark (Dependencies).

[Supremum](#); [Infimum](#); [Decreasing Function](#); [Continuity at a Point](#).

Theorem (Increasing Inverse Preserves Suprema)

Theorem 3.3.20 (Increasing Inverse Preserves Suprema). *Let $I, J \subseteq \mathbb{R}$, and let $f : I \rightarrow J$ be bijective. Suppose $f^{-1} : J \rightarrow I$ is increasing and continuous. If $B \subseteq J$ is nonempty and bounded above, and if $\sup B \in J$, then*

$$\sup f^{-1}(B) = f^{-1}(\sup B).$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$B \neq \emptyset, \quad \sup B \in J, \quad f^{-1} \text{ increasing and continuous at } \sup B \implies \sup f^{-1}(B) = f^{-1}(\sup B). \blacksquare$$

Remark (Predicate reading).

$$\text{Supremum}(u, B) \wedge \text{Increasing}(f^{-1}, J) \wedge \text{ContinuousAt}(f^{-1}, u) \implies \text{Supremum}(f^{-1}(u), f^{-1}(B)). \blacksquare$$

Remark (Negated quantified statement).

$$\sup f^{-1}(B) \neq f^{-1}(\sup B)$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Supremum}(u, B) \wedge \text{Increasing}(f^{-1}, J) \wedge \text{ContinuousAt}(f^{-1}, u) \wedge \neg \text{Supremum}(f^{-1}(u), f^{-1}(B)).$$

Remark (Failure modes). The formula can fail if the inverse is not increasing, if $\sup B$ is outside the codomain set J , or if the inverse is not continuous at $\sup B$.

Remark (Failure mode decomposition). Apply the increasing image supremum theorem to $g = f^{-1}$ and the set B .

Remark (Dependencies).

[Increasing Image Preserves Suprema.](#)

Theorem (Increasing Inverse Preserves Infima)

Theorem 3.3.21 (Increasing Inverse Preserves Infima). *Let $I, J \subseteq \mathbb{R}$, and let $f : I \rightarrow J$ be bijective. Suppose $f^{-1} : J \rightarrow I$ is increasing and continuous. If $B \subseteq J$ is nonempty and bounded below, and if $\inf B \in J$, then*

$$\inf f^{-1}(B) = f^{-1}(\inf B).$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$B \neq \emptyset, \quad \inf B \in J, \quad f^{-1} \text{ increasing and continuous at } \inf B \implies \inf f^{-1}(B) = f^{-1}(\inf B).$$

Remark (Predicate reading).

$$\text{Infimum}(v, B) \wedge \text{Increasing}(f^{-1}, J) \wedge \text{ContinuousAt}(f^{-1}, v) \implies \text{Infimum}(f^{-1}(v), f^{-1}(B)).$$

Remark (Negated quantified statement).

$$\inf f^{-1}(B) \neq f^{-1}(\inf B)$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Infimum}(v, B) \wedge \text{Increasing}(f^{-1}, J) \wedge \text{ContinuousAt}(f^{-1}, v) \wedge \neg \text{Infimum}(f^{-1}(v), f^{-1}(B)).$$

Remark (Failure modes). The formula can fail if the inverse is not increasing, if $\inf B$ is outside the codomain set J , or if the inverse is not continuous at $\inf B$.

Remark (Failure mode decomposition). Apply the increasing image infimum theorem to $g = f^{-1}$ and the set B .

Remark (Dependencies).

[Increasing Image Preserves Infima.](#)

Theorem (Decreasing Inverse Sends Infimum to Supremum)

Theorem 3.3.22 (Decreasing Inverse Sends Infimum to Supremum). *Let $I, J \subseteq \mathbb{R}$, and let $f : I \rightarrow J$ be bijective. Suppose $f^{-1} : J \rightarrow I$ is decreasing and continuous. If $B \subseteq J$ is nonempty and bounded below, and if $\inf B \in J$, then*

$$\sup f^{-1}(B) = f^{-1}(\inf B).$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$B \neq \emptyset, \quad \inf B \in J, \quad f^{-1} \text{ decreasing and continuous at } \inf B \implies \sup f^{-1}(B) = f^{-1}(\inf B). \blacksquare$$

Remark (Predicate reading).

$$\text{Infimum}(v, B) \wedge \text{Decreasing}(f^{-1}, J) \wedge \text{ContinuousAt}(f^{-1}, v) \implies \text{Supremum}(f^{-1}(v), f^{-1}(B)).$$

Remark (Negated quantified statement).

$$\sup f^{-1}(B) \neq f^{-1}(\inf B)$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Infimum}(v, B) \wedge \text{Decreasing}(f^{-1}, J) \wedge \text{ContinuousAt}(f^{-1}, v) \wedge \neg \text{Supremum}(f^{-1}(v), f^{-1}(B)).$$

Remark (Failure modes). The formula can fail if the inverse is not decreasing, if $\inf B$ is outside J , or if the inverse is not continuous at $\inf B$.

Remark (Failure mode decomposition). Apply the decreasing image infimum-to-supremum theorem to $g = f^{-1}$ and B .

Remark (Dependencies).

[Decreasing Image Sends Infimum to Supremum.](#)

Theorem (Decreasing Inverse Sends Supremum to Infimum)

Theorem 3.3.23 (Decreasing Inverse Sends Supremum to Infimum). *Let $I, J \subseteq \mathbb{R}$, and let $f : I \rightarrow J$ be bijective. Suppose $f^{-1} : J \rightarrow I$ is decreasing and continuous. If $B \subseteq J$ is nonempty and bounded above, and if $\sup B \in J$, then*

$$\inf f^{-1}(B) = f^{-1}(\sup B).$$

[Go to proof.](#)

Remark (Standard quantified statement).

$B \neq \emptyset, \quad \sup B \in J, \quad f^{-1} \text{ decreasing and continuous at } \sup B \implies \inf f^{-1}(B) = f^{-1}(\sup B).$ ■

Remark (Predicate reading).

$\text{Supremum}(u, B) \wedge \text{Decreasing}(f^{-1}, J) \wedge \text{ContinuousAt}(f^{-1}, u) \implies \text{Infimum}(f^{-1}(u), f^{-1}(B)).$

Remark (Negated quantified statement).

$$\inf f^{-1}(B) \neq f^{-1}(\sup B)$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$\text{Supremum}(u, B) \wedge \text{Decreasing}(f^{-1}, J) \wedge \text{ContinuousAt}(f^{-1}, u) \wedge \neg \text{Infimum}(f^{-1}(u), f^{-1}(B)).$

Remark (Failure modes). The formula can fail if the inverse is not decreasing, if $\sup B$ is outside J , or if the inverse is not continuous at $\sup B$.

Remark (Failure mode decomposition). Apply the decreasing image supremum-to-infimum theorem to $g = f^{-1}$ and B .

Remark (Dependencies).

[Decreasing Image Sends Supremum to Infimum.](#)

3.3.4 Dilation and Displacement

Remark (Orientation). Dilation and displacement are affine operations on sets. Translation preserves the order of all comparisons, positive dilation preserves order, and negative dilation reverses order.

Definition (Displacement, Dilation, and Reflection)

Definition 3.3.24 (Displacement, Dilation, and Reflection). Let $A \subseteq \mathbb{R}$ and let $c, \lambda \in \mathbb{R}$. Define

$$A + c := \{a + c : a \in A\}, \quad A - c := \{a - c : a \in A\}, \quad \lambda A := \{\lambda a : a \in A\}.$$

The set $-A := (-1)A$ is the *reflection* of A through the origin.

Remark (Standard quantified statement).

$$x \in A + c \iff (\exists a \in A)(x = a + c), \quad x \in \lambda A \iff (\exists a \in A)(x = \lambda a).$$

Remark (Definition predicate reading).

$$\text{Translate}(A, c) = A + c, \quad \text{Dilate}(A, \lambda) = \lambda A, \quad \text{Reflect}(A) = -A.$$

Remark (Negated quantified statement).

$$x \notin A + c \iff (\forall a \in A)(x \neq a + c), \quad x \notin \lambda A \iff (\forall a \in A)(x \neq \lambda a).$$

Remark (Negation predicate reading).

$$\neg \text{Translate}(A, c) = B \quad \text{means} \quad (\exists x)[x \in B \Delta (A + c)],$$

and similarly for dilation and reflection.

Remark (Failure modes). The definition fails only if the image set omits an actual transformed point or contains a point that is not produced by transforming an element of A .

Remark (Failure mode decomposition). Every membership claim in a transformed set decomposes into an original element and the transformation rule that produced the new point.

Remark (Interpretation). Displacement slides the set, dilation stretches or collapses it, and reflection is the special dilation that reverses the real line.

Remark (Dependencies).

[Subset](#); [Multiplication on \$\mathbb{R}\$](#) ; [Addition on \$\mathbb{R}\$](#) .

Proposition (Translation Preserves Upper Bounds)

Proposition 3.3.25 (Translation Preserves Upper Bounds). Let $A \subseteq \mathbb{R}$ be nonempty. If u is an upper bound of A , then $u + c$ is an upper bound of $A + c$. [Go to proof.](#)

Remark (Standard quantified statement).

$$(\forall a \in A)(a \leq u) \implies (\forall y \in A + c)(y \leq u + c).$$

Remark (Predicate reading).

$$\text{UB}(u, A) \implies \text{UB}(u + c, A + c).$$

Remark (Negated quantified statement).

$$(\forall a \in A)(a \leq u), \quad (\exists y \in A + c)(u + c < y).$$

Remark (Negation predicate reading).

$$\text{UB}(u, A) \wedge \neg \text{UB}(u + c, A + c).$$

Remark (Failure modes). A failure would require a translated point above the translated upper bound.

Remark (Failure mode decomposition). Write $y = a + c$ with $a \in A$. Then $a \leq u$ gives $a + c \leq u + c$.

Remark (Dependencies).

[Upper Bound](#); [Displacement, Dilation, and Reflection](#).

Proposition (Translation Preserves Lower Bounds)

Proposition 3.3.26 (Translation Preserves Lower Bounds). *Let $A \subseteq \mathbb{R}$ be nonempty. If ℓ is a lower bound of A , then $\ell + c$ is a lower bound of $A + c$. [Go to proof](#).*

Remark (Standard quantified statement).

$$(\forall a \in A)(\ell \leq a) \implies (\forall y \in A + c)(\ell + c \leq y).$$

Remark (Predicate reading).

$$\text{LB}(\ell, A) \implies \text{LB}(\ell + c, A + c).$$

Remark (Negated quantified statement).

$$(\forall a \in A)(\ell \leq a), \quad (\exists y \in A + c)(y < \ell + c).$$

Remark (Negation predicate reading).

$$\text{LB}(\ell, A) \wedge \neg \text{LB}(\ell + c, A + c).$$

Remark (Failure modes). A failure would require a translated point below the translated lower bound.

Remark (Failure mode decomposition). Write $y = a + c$ with $a \in A$. Then $\ell \leq a$ gives $\ell + c \leq a + c$.

Remark (Dependencies).

[Lower Bound](#); [Displacement, Dilation, and Reflection](#).

Proposition (Positive Dilation Preserves Upper Bounds)

Proposition 3.3.27 (Positive Dilation Preserves Upper Bounds). *Let $A \subseteq \mathbb{R}$ be nonempty and let $\lambda > 0$. If u is an upper bound of A , then λu is an upper bound of λA . [Go to proof](#).*

Remark (Standard quantified statement).

$$\lambda > 0 \wedge (\forall a \in A)(a \leq u) \implies (\forall y \in \lambda A)(y \leq \lambda u).$$

Remark (Predicate reading).

$$\lambda > 0 \wedge \text{UB}(u, A) \implies \text{UB}(\lambda u, \lambda A).$$

Remark (Negated quantified statement).

$$\lambda > 0, \quad (\forall a \in A)(a \leq u), \quad (\exists y \in \lambda A)(\lambda u < y).$$

Remark (Negation predicate reading).

$$\lambda > 0 \wedge \text{UB}(u, A) \wedge \neg \text{UB}(\lambda u, \lambda A).$$

Remark (Failure modes). A failure would require positive multiplication to reverse an inequality.

Remark (Failure mode decomposition). Write $y = \lambda a$ with $a \in A$. Since $a \leq u$ and $\lambda > 0$, $\lambda a \leq \lambda u$.

Remark (Dependencies).

Upper Bound; Displacement, Dilation, and Reflection.

Proposition (Positive Dilation Preserves Lower Bounds)

Proposition 3.3.28 (Positive Dilation Preserves Lower Bounds). *Let $A \subseteq \mathbb{R}$ be nonempty and let $\lambda > 0$. If ℓ is a lower bound of A , then $\lambda\ell$ is a lower bound of λA . [Go to proof.](#)*

Remark (Standard quantified statement).

$$\lambda > 0 \wedge (\forall a \in A)(\ell \leq a) \implies (\forall y \in \lambda A)(\lambda\ell \leq y).$$

Remark (Predicate reading).

$$\lambda > 0 \wedge \text{LB}(\ell, A) \implies \text{LB}(\lambda\ell, \lambda A).$$

Remark (Negated quantified statement).

$$\lambda > 0, \quad (\forall a \in A)(\ell \leq a), \quad (\exists y \in \lambda A)(y < \lambda\ell).$$

Remark (Negation predicate reading).

$$\lambda > 0 \wedge \text{LB}(\ell, A) \wedge \neg \text{LB}(\lambda\ell, \lambda A).$$

Remark (Failure modes). A failure would require a positively dilated point below the positively dilated lower bound.

Remark (Failure mode decomposition). Write $y = \lambda a$ with $a \in A$. Since $\ell \leq a$ and $\lambda > 0$, $\lambda\ell \leq \lambda a$.

Remark (Dependencies).

Lower Bound; Displacement, Dilation, and Reflection.

Proposition (Negative Dilation Sends Lower Bounds to Upper Bounds)

Proposition 3.3.29 (Negative Dilation Sends Lower Bounds to Upper Bounds). *Let $A \subseteq \mathbb{R}$ be nonempty and let $\lambda < 0$. If ℓ is a lower bound of A , then $\lambda\ell$ is an upper bound of λA . Go to proof.*

Remark (Standard quantified statement).

$$\lambda < 0 \wedge (\forall a \in A)(\ell \leq a) \implies (\forall y \in \lambda A)(y \leq \lambda\ell).$$

Remark (Predicate reading).

$$\lambda < 0 \wedge \text{LB}(\ell, A) \implies \text{UB}(\lambda\ell, \lambda A).$$

Remark (Negated quantified statement).

$$\lambda < 0, \quad (\forall a \in A)(\ell \leq a), \quad (\exists y \in \lambda A)(\lambda\ell < y).$$

Remark (Negation predicate reading).

$$\lambda < 0 \wedge \text{LB}(\ell, A) \wedge \neg \text{UB}(\lambda\ell, \lambda A).$$

Remark (Failure modes). A failure would require negative multiplication not to reverse the lower-bound inequality.

Remark (Failure mode decomposition). Write $y = \lambda a$ with $a \in A$. Since $\ell \leq a$ and $\lambda < 0$, $\lambda a \leq \lambda\ell$.

Remark (Dependencies).

Lower Bound; Upper Bound; Displacement, Dilation, and Reflection.

Proposition (Negative Dilation Sends Upper Bounds to Lower Bounds)

Proposition 3.3.30 (Negative Dilation Sends Upper Bounds to Lower Bounds). *Let $A \subseteq \mathbb{R}$ be nonempty and let $\lambda < 0$. If u is an upper bound of A , then λu is a lower bound of λA . Go to proof.*

Remark (Standard quantified statement).

$$\lambda < 0 \wedge (\forall a \in A)(a \leq u) \implies (\forall y \in \lambda A)(\lambda u \leq y).$$

Remark (Predicate reading).

$$\lambda < 0 \wedge \text{UB}(u, A) \implies \text{LB}(\lambda u, \lambda A).$$

Remark (Negated quantified statement).

$$\lambda < 0, \quad (\forall a \in A)(a \leq u), \quad (\exists y \in \lambda A)(y < \lambda u).$$

Remark (Negation predicate reading).

$$\lambda < 0 \wedge \text{UB}(u, A) \wedge \neg \text{LB}(\lambda u, \lambda A).$$

Remark (Failure modes). A failure would require negative multiplication not to reverse the upper-bound inequality.

Remark (Failure mode decomposition). Write $y = \lambda a$ with $a \in A$. Since $a \leq u$ and $\lambda < 0$, $\lambda u \leq \lambda a$.

Remark (Dependencies).

Upper Bound; Lower Bound; Displacement, Dilation, and Reflection.

Corollary (Reflection Swaps Upper and Lower Bounds)

Corollary 3.3.31 (Reflection Swaps Upper and Lower Bounds). *Let $A \subseteq \mathbb{R}$ be nonempty. If ℓ is a lower bound of A , then $-\ell$ is an upper bound of $-A$. If u is an upper bound of A , then $-u$ is a lower bound of $-A$. [Go to proof.](#)*

Remark (Standard quantified statement).

$$(\forall a \in A)(\ell \leq a) \implies (\forall y \in -A)(y \leq -\ell),$$

and

$$(\forall a \in A)(a \leq u) \implies (\forall y \in -A)(-u \leq y).$$

Remark (Predicate reading).

$$\text{LB}(\ell, A) \implies \text{UB}(-\ell, -A), \quad \text{UB}(u, A) \implies \text{LB}(-u, -A).$$

Remark (Negated quantified statement). A failure is a failure of negative dilation at the scalar $\lambda = -1$.

Remark (Negation predicate reading).

$$\text{LB}(\ell, A) \wedge \neg \text{UB}(-\ell, -A)$$

or

$$\text{UB}(u, A) \wedge \neg \text{LB}(-u, -A).$$

Remark (Failure modes). Reflection reverses the real line, so lower information becomes upper information and upper information becomes lower information.

Remark (Failure mode decomposition). Apply negative dilation with $\lambda = -1$.

Remark (Dependencies).

Negative Dilation Sends Lower Bounds to Upper Bounds; Negative Dilation Sends Upper Bounds to Lower Bounds.

3.3.5 Algebra of Suprema and Infima

Current Algebraic Properties of Supremum and Infimum

Remark (Section boundary). Basic bound, supremum, infimum, maximum, and minimum facts now live in [Upper and Lower Bounds](#), [Suprema and Infima](#), and [Maxima and Minima](#). This section begins with algebraic transformations of those extremal values.

Definition (Set Arithmetic Images)

Definition 3.3.32 (Set Arithmetic Images). Let $A, B \subseteq \mathbb{R}$. Define

$$\begin{aligned} A + B &:= \{a + b : a \in A, b \in B\}, & A - B &:= \{a - b : a \in A, b \in B\}, \\ AB &:= \{ab : a \in A, b \in B\}, & A/B &:= \{a/b : a \in A, b \in B, b \neq 0\}. \end{aligned}$$

Remark (Standard quantified statement).

$$x \in A + B \iff (\exists a \in A)(\exists b \in B)(x = a + b),$$

and similarly for difference, product, and quotient with the quotient condition $b \neq 0$.

Remark (Definition predicate reading).

$$\text{SetSum}(A, B) = A + B, \quad \text{SetDifference}(A, B) = A - B, \quad \text{SetProduct}(A, B) = AB, \quad \text{SetQuotient}(A, B) = A/B$$

Remark (Negated quantified statement).

$$x \notin A + B \iff (\forall a \in A)(\forall b \in B)(x \neq a + b),$$

with the analogous readings for the other arithmetic images.

Remark (Negation predicate reading).

$$\neg \text{SetProduct}(A, B) = C \quad \text{means} \quad (\exists x)[x \in C \triangle AB],$$

and similarly for sum, difference, and quotient.

Remark (Failure modes). An arithmetic-image definition fails only if it omits a value produced by the rule or includes a value not produced by the rule.

Remark (Failure mode decomposition). Every membership claim decomposes into witnesses from the input sets and the arithmetic operation that produced the image point.

Remark (Dependencies).

[Subset](#); [Addition on \$\mathbb{R}\$](#) ; [Multiplication on \$\mathbb{R}\$](#) .

Definition (Reciprocal Image)

Definition 3.3.33 (Reciprocal Image). Let $A \subseteq \mathbb{R}$ and suppose $0 \notin A$. The reciprocal image of A is

$$A^{-1} := \{1/a : a \in A\}.$$

Remark (Standard quantified statement).

$$x \in A^{-1} \iff (\exists a \in A)(x = 1/a).$$

Remark (Definition predicate reading).

$$\text{ReciprocalImage}(A) = A^{-1}.$$

Remark (Negated quantified statement).

$$x \notin A^{-1} \iff (\forall a \in A)(x \neq 1/a).$$

Remark (Negation predicate reading).

$$\neg \text{ReciprocalImage}(A) = B \quad \text{means} \quad (\exists x)[x \in B \Delta A^{-1}].$$

Remark (Failure modes). The reciprocal image is undefined if zero is allowed as an input value. Once zero is excluded, the only definition failures are omitted or spurious reciprocal values.

Remark (Failure mode decomposition). Every element of A^{-1} has the form $1/a$ for a nonzero $a \in A$.

Remark (Dependencies).

[Subset](#); [Multiplication on \$\mathbb{R}\$](#) .

Theorem (Translation Invariance of the Supremum)

Theorem 3.3.34 (Translation Invariance of the Supremum). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded above, and let $c \in \mathbb{R}$. Then*

$$\sup(A + c) = \sup A + c, \quad A + c = \{a + c : a \in A\}.$$

[Go to proof](#).

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } A \subseteq \mathbb{R} \text{ be nonempty and bounded above, and let } c \in \mathbb{R}. \\ &\sup\{a + c : a \in A\} = \sup A + c. \end{aligned}$$

Remark (Predicate reading).

$$\text{Supremum}(s, A) \implies \text{Supremum}(s + c, A + c).$$

Remark (Negated quantified statement).

$$\sup(A + c) \neq \sup A + c$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Supremum}(s, A) \wedge \neg \text{Supremum}(s + c, A + c)$$

is a contradictory configuration.

Remark (Failure modes). The formula could fail only if translation either failed to preserve upper bounds or failed to preserve leastness. Adding the same real number to every comparison preserves the order, so neither failure mode can occur.

Remark (Failure mode decomposition).

$$\begin{aligned} \forall a \in A (a \leq s) &\iff \forall a \in A (a + c \leq s + c), \\ \forall u [\text{UpperBound}(u, A) \implies s \leq u] &\iff \forall v [\text{UpperBound}(v, A + c) \implies s + c \leq v]. \end{aligned}$$

Remark (Interpretation). Translation shifts the entire set and all of its ceilings by the same amount. The least ceiling therefore shifts by exactly that amount.

Remark (Dependencies).

[Upper Bound](#); [Supremum](#).

Theorem (Translation Invariance of the Infimum)

Theorem 3.3.35 (Translation Invariance of the Infimum). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded below, and let $c \in \mathbb{R}$. Then*

$$\inf(A + c) = \inf A + c.$$

[Go to proof](#).

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad A \text{ bounded below} \implies \inf\{a + c : a \in A\} = \inf A + c.$$

Remark (Predicate reading).

$$\text{Infimum}(t, A) \implies \text{Infimum}(t + c, A + c).$$

Remark (Negated quantified statement).

$$\inf(A + c) \neq \inf A + c$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Infimum}(t, A) \wedge \neg \text{Infimum}(t + c, A + c).$$

Remark (Failure modes). The formula could fail only if translation failed to preserve lower bounds or failed to preserve greatestness among lower bounds.

Remark (Failure mode decomposition).

$$(\forall a \in A)(t \leq a) \iff (\forall a \in A)(t + c \leq a + c).$$

Remark (Interpretation). Translation shifts every floor by the same amount, so the greatest floor shifts by that amount.

Remark (Dependencies).

[Lower Bound](#); [Infimum](#); [Set Arithmetic Images](#).

Theorem (Scalar Multiplication by a Positive Scalar)

Theorem 3.3.36 (Scalar Multiplication by a Positive Scalar). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded above, and let $k > 0$. Then*

$$\sup(kA) = k \sup A, \quad kA = \{ka : a \in A\}.$$

[Go to proof](#).

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } A \subseteq \mathbb{R} \text{ be nonempty and bounded above, and let } k > 0. \\ &\sup\{ka : a \in A\} = k \sup A. \end{aligned}$$

Remark (Predicate reading).

$$\text{Supremum}(s, A) \wedge k > 0 \implies \text{Supremum}(ks, kA).$$

Remark (Negated quantified statement).

$$k > 0, \quad \sup(kA) \neq k \sup A$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Supremum}(s, A) \wedge k > 0 \wedge \neg \text{Supremum}(ks, kA)$$

is a contradictory configuration.

Remark (Failure modes). The formula could fail only if multiplying by a positive scalar failed to preserve upper bounds or failed to preserve leastness. Positive multiplication preserves the order, so both properties scale exactly.

Remark (Failure mode decomposition).

$$\begin{aligned} \forall a \in A (a \leq s) &\iff \forall a \in A (ka \leq ks), \\ \text{UpperBound}(u, A) &\iff \text{UpperBound}(ku, kA). \end{aligned}$$

Remark (Interpretation). Positive scaling stretches the set without reversing order. The least ceiling stretches by the same positive factor.

Remark (Dependencies).

[Upper Bound](#); [Supremum](#).

Theorem (Positive Scalar Multiplication of the Infimum)

Theorem 3.3.37 (Positive Scalar Multiplication of the Infimum). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded below, and let $k > 0$. Then*

$$\inf(kA) = k \inf A.$$

[Go to proof](#).

Remark (Standard quantified statement).

$$k > 0, \quad A \neq \emptyset, \quad A \text{ bounded below} \implies \inf\{ka : a \in A\} = k \inf A.$$

Remark (Predicate reading).

$$\text{Infimum}(t, A) \wedge k > 0 \implies \text{Infimum}(kt, kA).$$

Remark (Negated quantified statement).

$$k > 0, \quad \inf(kA) \neq k \inf A$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Infimum}(t, A) \wedge k > 0 \wedge \neg \text{Infimum}(kt, kA).$$

Remark (Failure modes). The formula could fail only if positive multiplication failed to preserve lower bounds or greatestness among lower bounds.

Remark (Failure mode decomposition).

$$(\forall a \in A)(t \leq a) \iff (\forall a \in A)(kt \leq ka) \quad (k > 0).$$

Remark (Interpretation). Positive scaling stretches the floor of the set by the same factor.

Remark (Dependencies).

[Lower Bound](#); [Infimum](#); [Displacement](#), [Dilation](#), and [Reflection](#).

Theorem (Scalar Multiplication by a Negative Scalar)

Theorem 3.3.38 (Scalar Multiplication by a Negative Scalar). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded below, and let $k < 0$. Then*

$$\sup(kA) = k \inf A, \quad kA = \{ka : a \in A\}.$$

[Go to proof](#).

Remark (Standard quantified statement).

Let $A \subseteq \mathbb{R}$ be nonempty and bounded below, and let $k < 0$.
 $\sup\{ka : a \in A\} = k \inf A$.

Remark (Predicate reading).

$$\text{Infimum}(t, A) \wedge k < 0 \implies \text{Supremum}(kt, kA).$$

Remark (Negated quantified statement).

$$k < 0, \quad \sup(kA) \neq k \inf A$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Infimum}(t, A) \wedge k < 0 \wedge \neg \text{Supremum}(kt, kA)$$

is a contradictory configuration.

Remark (Failure modes). The formula could fail only if negative multiplication did not reverse lower bounds into upper bounds, or if the extremal value lost its leastness after reversal. Multiplication by a negative scalar reverses order exactly, so the infimum of A becomes the supremum of kA .

Remark (Failure mode decomposition).

$$\begin{aligned} \forall a \in A (t \leq a) &\iff \forall a \in A (ka \leq kt), \\ \text{LowerBound}(v, A) &\iff \text{UpperBound}(kv, kA) \quad (k < 0). \end{aligned}$$

Remark (Interpretation). Negative scaling reflects the set and changes floors into ceilings. The lowest point threshold of A becomes the highest point threshold of the reflected and scaled set.

Remark (Dependencies).

[Lower Bound](#); [Upper Bound](#); [Infimum](#); [Supremum](#).

Theorem (Negative Scalar Multiplication of the Infimum)

Theorem 3.3.39 (Negative Scalar Multiplication of the Infimum). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded above, and let $k < 0$. Then*

$$\inf(kA) = k \sup A.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$k < 0, \quad A \neq \emptyset, \quad A \text{ bounded above} \implies \inf\{ka : a \in A\} = k \sup A.$$

Remark (Predicate reading).

$$\text{Supremum}(s, A) \wedge k < 0 \implies \text{Infimum}(ks, kA).$$

Remark (Negated quantified statement).

$$k < 0, \quad \inf(kA) \neq k \sup A$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Supremum}(s, A) \wedge k < 0 \wedge \neg \text{Infimum}(ks, kA).$$

Remark (Failure modes). The formula could fail only if negative multiplication failed to reverse upper bounds into lower bounds or failed to preserve the extremal threshold after reversal.

Remark (Failure mode decomposition).

$$(\forall a \in A)(a \leq s) \iff (\forall a \in A)(ks \leq ka) \quad (k < 0).$$

Remark (Interpretation). Negative scaling reflects the set, so the ceiling of A becomes the floor of kA .

Remark (Dependencies).

[Upper Bound](#); [Lower Bound](#); [Supremum](#); [Infimum](#).

Theorem (Negation Exchanges Supremum and Infimum)

Theorem 3.3.40 (Negation Exchanges Supremum and Infimum). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded below. Then $-A = \{-a : a \in A\}$ is bounded above and*

$$\sup(-A) = -\inf A.$$

[Go to proof](#).

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } A \subseteq \mathbb{R} \text{ be nonempty and bounded below.} \\ &\sup\{-a : a \in A\} = -\inf A. \end{aligned}$$

Remark (Predicate reading).

$$\text{Infimum}(t, A) \implies \text{Supremum}(-t, -A).$$

Remark (Negated quantified statement).

$$\sup(-A) \neq -\inf A$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Infimum}(t, A) \wedge \neg \text{Supremum}(-t, -A)$$

is a contradictory configuration.

Remark (Failure modes). The theorem could fail only if negation did not convert lower bounds into upper bounds or did not preserve the extremal threshold after reversing order. The map $x \mapsto -x$ reverses all inequalities, so the exchange is exact.

Remark (Failure mode decomposition).

$$\begin{aligned} \forall a \in A (t \leq a) &\iff \forall a \in A (-a \leq -t), \\ \text{LowerBound}(v, A) &\iff \text{UpperBound}(-v, -A). \end{aligned}$$

Remark (Interpretation). Negation reflects the real line through the origin. Floors become ceilings, so the infimum of the original set becomes the negative of the supremum of the reflected set.

Remark (Dependencies).

[Lower Bound](#); [Upper Bound](#); [Infimum](#); [Supremum](#).

Theorem (Supremum of a Sum Set)

Theorem 3.3.41 (Supremum of a Sum Set). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded above. Then*

$$\sup(A + B) = \sup A + \sup B, \quad A + B = \{a + b : a \in A, b \in B\}.$$

[Go to proof](#).

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } A, B \subseteq \mathbb{R} \text{ be nonempty and bounded above.} \\ &\sup\{a + b : a \in A, b \in B\} = \sup A + \sup B. \end{aligned}$$

Remark (Predicate reading).

$$\text{Supremum}(s_A, A) \wedge \text{Supremum}(s_B, B) \implies \text{Supremum}(s_A + s_B, A + B).$$

Remark (Negated quantified statement).

$$\sup(A + B) \neq \sup A + \sup B$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Supremum}(s_A, A) \wedge \text{Supremum}(s_B, B) \wedge \neg \text{Supremum}(s_A + s_B, A + B)$$

is a contradictory configuration.

Remark (Failure modes). The equality could fail only if the sum of the two suprema did not bound the sum set, or if it were not the least such bound. The first part follows from adding the two upper-bound inequalities; the second follows by approximating each supremum from within its set.

Remark (Failure mode decomposition).

$$\begin{aligned} a \leq \sup A, \quad b \leq \sup B &\implies a + b \leq \sup A + \sup B, \\ \varepsilon > 0 &\implies \exists a \in A, b \in B : \sup A + \sup B - \varepsilon < a + b. \end{aligned}$$

Remark (Interpretation). The highest limiting value of all pairwise sums is obtained by combining values that approach the separate ceilings of A and B .

Remark (Dependencies).

[Supremum](#); [Epsilon Characterization of Supremum](#).

Theorem (Infimum of a Sum Set)

Theorem 3.3.42 (Infimum of a Sum Set). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded below. Then*

$$\inf(A + B) = \inf A + \inf B.$$

[Go to proof](#).

Remark (Standard quantified statement).

$$A, B \neq \emptyset, \quad A, B \text{ bounded below} \implies \inf\{a + b : a \in A, b \in B\} = \inf A + \inf B.$$

Remark (Predicate reading).

$$\text{Infimum}(t_A, A) \wedge \text{Infimum}(t_B, B) \implies \text{Infimum}(t_A + t_B, A + B).$$

Remark (Negated quantified statement).

$$\inf(A + B) \neq \inf A + \inf B$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Infimum}(t_A, A) \wedge \text{Infimum}(t_B, B) \wedge \neg \text{Infimum}(t_A + t_B, A + B).$$

Remark (Failure modes). The equality could fail only if the sum of the two infima failed to be a lower bound of the sum set or failed to be the greatest such lower bound.

Remark (Failure mode decomposition).

$$t_A \leq a, \quad t_B \leq b \implies t_A + t_B \leq a + b,$$

and sharpness follows by choosing elements near each infimum.

Remark (Interpretation). The lowest limiting value of pairwise sums is obtained by combining values near the separate floors of the two sets.

Remark (Dependencies).

[Infimum](#); [Set Arithmetic Images](#).

Theorem (Supremum of a Difference Set)

Theorem 3.3.43 (Supremum of a Difference Set). *Let $A, B \subseteq \mathbb{R}$ be nonempty, with A bounded above and B bounded below. Then*

$$\sup(A - B) = \sup A - \inf B, \quad A - B = \{a - b : a \in A, b \in B\}.$$

[Go to proof](#).

Remark (Standard quantified statement).

Let $A, B \subseteq \mathbb{R}$ be nonempty, A bounded above, and B bounded below.

$$\sup\{a - b : a \in A, b \in B\} = \sup A - \inf B.$$

Remark (Predicate reading).

$$\text{Supremum}(s, A) \wedge \text{Infimum}(t, B) \implies \text{Supremum}(s - t, A - B).$$

Remark (Negated quantified statement).

$$\sup(A - B) \neq \sup A - \inf B$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Supremum}(s, A) \wedge \text{Infimum}(t, B) \wedge \neg \text{Supremum}(s - t, A - B)$$

is a contradictory configuration.

Remark (Failure modes). The formula could fail only if subtracting elements of B did not convert the lower extremal behavior of B into an upper contribution. Since $a - b = a + (-b)$, this is the sum theorem combined with the negation exchange.

Remark (Failure mode decomposition).

$$A - B = A + (-B), \quad \sup(-B) = -\inf B, \quad \sup(A + (-B)) = \sup A + \sup(-B).$$

Remark (Interpretation). The largest possible difference comes from making the first term as large as possible and the subtracted term as small as possible.

Remark (Dependencies).

[Supremum](#); [Infimum](#); [Negation Exchanges Supremum and Infimum](#); [Supremum of a Sum Set](#).

Theorem (Infimum of a Difference Set)

Theorem 3.3.44 (Infimum of a Difference Set). *Let $A, B \subseteq \mathbb{R}$ be nonempty, with A bounded below and B bounded above. Then*

$$\inf(A - B) = \inf A - \sup B.$$

Go to proof.

Remark (Standard quantified statement).

$$A, B \neq \emptyset, \quad A \text{ bounded below}, \quad B \text{ bounded above} \implies \inf\{a-b : a \in A, b \in B\} = \inf A - \sup B. \blacksquare$$

Remark (Predicate reading).

$$\text{Infimum}(t, A) \wedge \text{Supremum}(s, B) \implies \text{Infimum}(t - s, A - B).$$

Remark (Negated quantified statement).

$$\inf(A - B) \neq \inf A - \sup B$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Infimum}(t, A) \wedge \text{Supremum}(s, B) \wedge \neg \text{Infimum}(t - s, A - B).$$

Remark (Failure modes). The formula could fail only if subtracting B failed to convert its upper extremal behavior into a lower contribution.

Remark (Failure mode decomposition).

$$A - B = A + (-B), \quad \inf(-B) = -\sup B, \quad \inf(A + (-B)) = \inf A + \inf(-B).$$

Remark (Interpretation). The smallest possible difference comes from making the first term as small as possible and the subtracted term as large as possible.

Remark (Dependencies).

[Infimum](#); [Supremum](#); [Infimum of a Sum Set](#); [Negative Scalar Multiplication of the Infimum](#).

Theorem (Supremum of a Dilation)

Theorem 3.3.45 (Supremum of a Dilation). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded, and let $\lambda \in \mathbb{R}$. Then*

$$\sup(\lambda A) = \max\{\lambda \sup A, \lambda \inf A\}, \quad \lambda A = \{\lambda a : a \in A\}.$$

Go to proof.

Remark (Standard quantified statement).

Let $A \subseteq \mathbb{R}$ be nonempty and bounded, and let $\lambda \in \mathbb{R}$.
 $\sup\{\lambda a : a \in A\} = \max\{\lambda \sup A, \lambda \inf A\}.$

Remark (Predicate reading).

$\text{Supremum}(s, A) \wedge \text{Infimum}(t, A) \implies \text{Supremum}(\max\{\lambda s, \lambda t\}, \lambda A).$

Remark (Negated quantified statement).

$$\sup(\lambda A) \neq \max\{\lambda \sup A, \lambda \inf A\}$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$\text{Supremum}(s, A) \wedge \text{Infimum}(t, A) \wedge \neg \text{Supremum}(\max\{\lambda s, \lambda t\}, \lambda A)$

is a contradictory configuration.

Remark (Failure modes). The formula separates into the cases $\lambda > 0$, $\lambda = 0$, and $\lambda < 0$. Positive dilation scales the supremum, zero dilation collapses the set to $\{0\}$, and negative dilation turns the infimum into the supremum.

Remark (Failure mode decomposition).

$$\sup(\lambda A) = \begin{cases} \lambda \sup A, & \lambda > 0, \\ 0, & \lambda = 0, \\ \lambda \inf A, & \lambda < 0. \end{cases}$$

Remark (Interpretation). A dilation either preserves the order, collapses all points to zero, or reverses the order. The displayed maximum packages those three cases into one formula.

Remark (Dependencies).

[Supremum](#); [Infimum](#); [Maximum](#); [Scalar Multiplication by a Positive Scalar](#); [Scalar Multiplication by a Negative Scalar](#).

Theorem (Supremum of the Absolute-Value Image)

Theorem 3.3.46 (Supremum of the Absolute-Value Image). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded. Then*

$$\sup |A| = \max\{|\sup A|, |\inf A|\}, \quad |A| = \{|a| : a \in A\}.$$

Go to proof.

Remark (Standard quantified statement).

Let $A \subseteq \mathbb{R}$ be nonempty and bounded.
 $\sup\{|a| : a \in A\} = \max\{|\sup A|, |\inf A|\}.$

Remark (Predicate reading).

$$\text{Supremum}(s, A) \wedge \text{Infimum}(t, A) \implies \text{Supremum}(\max\{|s|, |t|\}, |A|).$$

Remark (Negated quantified statement).

$$\sup |A| \neq \max\{|\sup A|, |\inf A|\}$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$\text{Supremum}(s, A) \wedge \text{Infimum}(t, A) \wedge \neg \text{Supremum}(\max\{|s|, |t|\}, |A|)$$

is a contradictory configuration.

Remark (Failure modes). The formula could fail only if the farthest absolute-value size were not achieved at one of the two extremal thresholds. Since every $a \in A$ lies between $\inf A$ and $\sup A$, its distance from zero is bounded by the larger endpoint magnitude.

Remark (Failure mode decomposition).

$$\inf A \leq a \leq \sup A \implies |a| \leq \max\{|\sup A|, |\inf A|\}.$$

Remark (Interpretation). The absolute-value image measures distance from the origin. For a bounded subset of the real line, the largest possible distance is controlled by whichever extremal endpoint lies farther from zero.

Remark (Dependencies).

[Supremum](#); [Infimum](#); [Maximum](#); [Every Element Lies Below the Supremum](#).

Theorem (Supremum of a Reciprocal Set)

Theorem 3.3.47 (Supremum of a Reciprocal Set). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded, and suppose that $0 < \inf A$ or $\sup A < 0$. Define*

$$A^{-1} := \{1/a : a \in A\}.$$

Then

$$\sup(A^{-1}) = \frac{1}{\inf A}.$$

[Go to proof](#).

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad A \text{ bounded}, \quad 0 \notin [\inf A, \sup A] \implies \sup\{1/a : a \in A\} = (\inf A)^{-1}.$$

Remark (Predicate reading).

$$\text{Infimum}(m, A) \wedge \text{Supremum}(M, A) \wedge 0 \notin [m, M] \implies \text{Supremum}(m^{-1}, A^{-1}).$$

Remark (Negated quantified statement).

$$A \neq \emptyset, \quad A \text{ bounded}, \quad 0 \notin [\inf A, \sup A], \quad \sup(A^{-1}) \neq (\inf A)^{-1}.$$

Remark (Negation predicate reading).

$$\text{Infimum}(m, A) \wedge 0 \notin [m, \sup A] \wedge \neg \text{Supremum}(m^{-1}, A^{-1})$$

is contradictory under the stated hypotheses.

Remark (Failure modes). The formula fails without a nonzero gap from zero: reciprocal images may be unbounded or undefined. With the gap, reciprocal is order-reversing on the whole interval containing A , so the floor of A becomes the ceiling of A^{-1} .

Remark (Failure mode decomposition). Every $a \in A$ lies between $\inf A$ and $\sup A$, and the interval does not cross zero. The reciprocal map reverses comparisons there, so $1/a \leq 1/\inf A$. Leastness follows by applying the same reversal to values of A approaching $\inf A$.

Remark (Interpretation). Taking reciprocals turns small nonzero magnitudes into large magnitudes. The least value threshold of the original set therefore controls the largest value threshold of the reciprocal set.

Remark (Dependencies).

[Bounded Set](#); [Infimum](#); [Supremum](#).

Theorem (Infimum of a Reciprocal Set)

Theorem 3.3.48 (Infimum of a Reciprocal Set). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded, and suppose that $0 < \inf A$ or $\sup A < 0$. Define*

$$A^{-1} := \{1/a : a \in A\}.$$

Then

$$\inf(A^{-1}) = \frac{1}{\sup A}.$$

[Go to proof](#).

Remark (Standard quantified statement).

$$A \neq \emptyset, \quad A \text{ bounded}, \quad 0 \notin [\inf A, \sup A] \implies \inf\{1/a : a \in A\} = (\sup A)^{-1}.$$

Remark (Predicate reading).

$$\text{Infimum}(m, A) \wedge \text{Supremum}(M, A) \wedge 0 \notin [m, M] \implies \text{Infimum}(M^{-1}, A^{-1}).$$

Remark (Negated quantified statement).

$$A \neq \emptyset, \quad A \text{ bounded}, \quad 0 \notin [\inf A, \sup A], \quad \inf(A^{-1}) \neq (\sup A)^{-1}.$$

Remark (Negation predicate reading).

$$\text{Supremum}(M, A) \wedge 0 \notin [\inf A, M] \wedge \neg \text{Infimum}(M^{-1}, A^{-1})$$

is contradictory under the stated hypotheses.

Remark (Failure modes). The formula fails without excluding zero from the closed extremal interval. Once zero is excluded, reciprocal reverses order and the ceiling of A becomes the floor of A^{-1} .

Remark (Failure mode decomposition). Every reciprocal $1/a$ lies above $1/\sup A$ because reciprocal reverses order on the nonzero interval containing A . Greatestness of $\sup A$ gives elements of A arbitrarily close to that ceiling, which makes the lower bound sharp.

Remark (Interpretation). For nonzero bounded sets that stay on one side of zero, reciprocal swaps the two extremal thresholds.

Remark (Dependencies).

[Bounded Set](#); [Infimum](#); [Supremum](#); [Supremum of a Reciprocal Set](#).

Theorem (Supremum of a Product Set)

Theorem 3.3.49 (Supremum of a Product Set). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded. Then*

$$\sup(AB) = \max\{(\inf A)(\inf B), (\inf A)(\sup B), (\sup A)(\inf B), (\sup A)(\sup B)\}.$$

[Go to proof](#).

Remark (Standard quantified statement).

$$A, B \neq \emptyset, \quad A, B \text{ bounded} \quad \implies \quad \sup\{ab : a \in A, b \in B\}$$

equals the maximum of the four endpoint products.

Remark (Predicate reading).

$$\text{Infimum}(\alpha, A) \wedge \text{Supremum}(\beta, A) \wedge \text{Infimum}(\gamma, B) \wedge \text{Supremum}(\delta, B) \implies \text{Supremum}(\max\{\alpha\gamma, \alpha\delta, \beta\gamma, \beta\delta\}, AB)$$

Remark (Negated quantified statement).

$$\sup(AB) \neq \max\{(\inf A)(\inf B), (\inf A)(\sup B), (\sup A)(\inf B), (\sup A)(\sup B)\}$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading). The product endpoint maximum is not the supremum of AB .

Remark (Failure modes). A failure would require a product above all four endpoint products or would require the endpoint maximum not to be least.

Remark (Failure mode decomposition). Every pair (a, b) lies in the rectangle $[\inf A, \sup A] \times [\inf B, \sup B]$. A bilinear product on a rectangle has its extreme values at corners, and elements can approximate the relevant corners from within the two sets.

Remark (Dependencies).

[Set Arithmetic Images](#); [Infimum](#); [Supremum](#); [Maximum](#).

Theorem (Infimum of a Product Set)

Theorem 3.3.50 (Infimum of a Product Set). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded. Then*

$$\inf(AB) = \min\{(\inf A)(\inf B), (\inf A)(\sup B), (\sup A)(\inf B), (\sup A)(\sup B)\}.$$

Go to proof.

Remark (Standard quantified statement).

$$A, B \neq \emptyset, \quad A, B \text{ bounded} \implies \inf\{ab : a \in A, b \in B\}$$

equals the minimum of the four endpoint products.

Remark (Predicate reading).

$$\text{Infimum}(\alpha, A) \wedge \text{Supremum}(\beta, A) \wedge \text{Infimum}(\gamma, B) \wedge \text{Supremum}(\delta, B) \implies \text{Infimum}(\min\{\alpha\gamma, \alpha\delta, \beta\gamma, \beta\delta\}, AB)$$

Remark (Negated quantified statement).

$$\inf(AB) \neq \min\{(\inf A)(\inf B), (\inf A)(\sup B), (\sup A)(\inf B), (\sup A)(\sup B)\}$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading). The product endpoint minimum is not the infimum of AB .

Remark (Failure modes). A failure would require a product below all four endpoint products or would require the endpoint minimum not to be greatest.

Remark (Failure mode decomposition). Reduce to the same endpoint analysis as the supremum product formula, replacing the corner maximum by the corner minimum.

Remark (Dependencies).

[Set Arithmetic Images](#); [Infimum](#); [Supremum](#); [Minimum](#); [Supremum of a Product Set](#).

Theorem (Supremum of a Quotient Set)

Theorem 3.3.51 (Supremum of a Quotient Set). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded, and suppose $0 < \inf B$ or $\sup B < 0$. Then*

$$\sup(A/B) = \max\left\{\frac{\inf A}{\inf B}, \frac{\inf A}{\sup B}, \frac{\sup A}{\inf B}, \frac{\sup A}{\sup B}\right\}.$$

Go to proof.

Remark (Standard quantified statement).

$$A, B \neq \emptyset, \quad A, B \text{ bounded}, \quad 0 \notin [\inf B, \sup B]$$

implies that the supremum of $\{a/b : a \in A, b \in B\}$ is the maximum of the four endpoint quotients.

Remark (Predicate reading).

$$\text{Supremum}\left(\max\left\{\frac{\inf A}{\inf B}, \frac{\inf A}{\sup B}, \frac{\sup A}{\inf B}, \frac{\sup A}{\sup B}\right\}, A/B\right).$$

Remark (Negated quantified statement). The displayed endpoint maximum is not the supremum of A/B .

Remark (Negation predicate reading).

$$\neg \text{Supremum}(\text{endpoint quotient maximum}, A/B).$$

Remark (Failure modes). A failure would require the quotient set to see values not controlled by the reciprocal endpoint interval for B .

Remark (Failure mode decomposition). Use $A/B = AB^{-1}$, the reciprocal endpoint formulas for B^{-1} , and the product endpoint formula.

Remark (Dependencies).

[Set Arithmetic Images](#); [Reciprocal Image](#); [Supremum of a Reciprocal Set](#); [Infimum of a Reciprocal Set](#); [Supremum of a Product Set](#).

Theorem (Infimum of a Quotient Set)

Theorem 3.3.52 (Infimum of a Quotient Set). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded, and suppose $0 < \inf B$ or $\sup B < 0$. Then*

$$\inf(A/B) = \min\left\{\frac{\inf A}{\inf B}, \frac{\inf A}{\sup B}, \frac{\sup A}{\inf B}, \frac{\sup A}{\sup B}\right\}.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$A, B \neq \emptyset, \quad A, B \text{ bounded}, \quad 0 \notin [\inf B, \sup B]$$

implies that the infimum of $\{a/b : a \in A, b \in B\}$ is the minimum of the four endpoint quotients.

Remark (Predicate reading).

$$\text{Infimum}\left(\min\left\{\frac{\inf A}{\inf B}, \frac{\inf A}{\sup B}, \frac{\sup A}{\inf B}, \frac{\sup A}{\sup B}\right\}, A/B\right).$$

Remark (Negated quantified statement). The displayed endpoint minimum is not the infimum of A/B .

Remark (Negation predicate reading).

$$\neg \text{Infimum}(\text{endpoint quotient minimum}, A/B).$$

Remark (Failure modes). A failure would require a quotient below all endpoint quotients or would require the endpoint minimum not to be greatest among lower bounds.

Remark (Failure mode decomposition). Use $A/B = AB^{-1}$ and combine the reciprocal endpoint formulas with the product endpoint infimum formula.

Remark (Dependencies).

[Set Arithmetic Images](#); [Reciprocal Image](#); [Supremum of a Reciprocal Set](#); [Infimum of a Reciprocal Set](#); [Infimum of a Product Set](#).

Proposition (Bounds for Suprema and Infima Under Set Addition)

Proposition 3.3.53 (Bounds for Suprema and Infima Under Set Addition). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded. Then*

$$\inf A + \inf B \leq \inf(A + B) \quad \text{and} \quad \sup(A + B) \leq \sup A + \sup B.$$

[Go to proof.](#)

Remark (Standard quantified statement).

Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded.

$$\inf A + \inf B \leq \inf(A + B) \quad \wedge \quad \sup(A + B) \leq \sup A + \sup B.$$

Remark (Predicate reading).

$$\text{Infimum}(t_A, A) \wedge \text{Infimum}(t_B, B) \wedge \text{Supremum}(s_A, A) \wedge \text{Supremum}(s_B, B) \implies t_A + t_B \leq \inf(A + B) \leq \sup(A + B) \leq s_A + s_B$$

Remark (Negated quantified statement).

$$\inf(A + B) < \inf A + \inf B \quad \text{or} \quad \sup A + \sup B < \sup(A + B)$$

is impossible under the stated hypotheses.

Remark (Negation predicate reading).

$$[\text{Infimum}(t_A, A) \wedge \text{Infimum}(t_B, B) \wedge \inf(A + B) < t_A + t_B] \vee [\text{Supremum}(s_A, A) \wedge \text{Supremum}(s_B, B) \wedge s_A + s_B < \sup(A + B)]$$

is a contradictory configuration.

Remark (Failure modes). The lower inequality could fail only if a sum fell below the sum of the two lower thresholds. The upper inequality could fail only if a sum rose above the sum of the two upper thresholds.

Remark (Failure mode decomposition).

$$\inf A \leq a \leq \sup A, \quad \inf B \leq b \leq \sup B \implies \inf A + \inf B \leq a + b \leq \sup A + \sup B.$$

Remark (Interpretation). Every element of the sum set is built from one element of A and one element of B . Adding the individual lower and upper controls gives immediate lower and upper controls for every such sum.

Remark (Dependencies).

[Infimum](#); [Supremum](#); [Supremum of a Sum Set](#).

Proposition (Upper Bounds Depend on the Ambient Order)

Proposition 3.3.54 (Upper Bounds Depend on the Ambient Order). *Let $(S, \leq_S)$ and $(T, \leq_T)$ be ordered sets, and let $A \subseteq S \cap T$. Whether u is an upper bound of A must be interpreted relative to the chosen ambient order. [Go to proof](#).*

Remark (Standard quantified statement).

$$\begin{aligned} u \text{ is an upper bound of } A \text{ in } S &\iff u \in S \text{ and } \forall x \in A (x \leq_S u), \\ u \text{ is an upper bound of } A \text{ in } T &\iff u \in T \text{ and } \forall x \in A (x \leq_T u). \end{aligned}$$

Remark (Predicate reading).

$$\text{UpperBound}_S(u, A) \iff u \in S \wedge \forall x \in A (x \leq_S u).$$

Remark (Negated quantified statement).

The phrase “ u is an upper bound of A ”

is not well-determined until the ambient ordered set and its order relation are fixed.

Remark (Negation predicate reading).

$$\neg \text{WellDetermined}(\text{UpperBound}(u, A))$$

when the ambient order is omitted.

Remark (Failure modes). Ambiguity occurs when the same set A is considered inside different ordered ambient sets or under different order relations. The candidate bound may belong to one ambient set and not another, or the relevant comparison relation may change.

Remark (Failure mode decomposition).

$$\text{UpperBound}_S(u, A) \text{ and } \text{UpperBound}_T(u, A)$$

are separate assertions unless $S = T$ and $\leq_S = \leq_T$ on the elements being compared.

Remark (Interpretation). An upper bound is not only about the subset; it is also about where the subset lives. Changing the ambient order changes the universe of possible bounds and the comparison used to test them.

Remark (Dependencies).

[Upper Bound](#).

Proposition (Suprema Depend on the Ambient Set)

Proposition 3.3.55 (Suprema Depend on the Ambient Set). *Let $A \subseteq S \subseteq S'$ be subsets of an ordered set. The supremum of A relative to S , if it exists, need not agree with the supremum of A relative to S' . [Go to proof](#).*

Remark (Standard quantified statement).

$$\sup_S A \text{ and } \sup_{S'} A$$

are ambient-relative objects and may differ in existence or value.

Remark (Predicate reading).

$$\text{Supremum}_S(s, A) \text{ and } \text{Supremum}_{S'}(s, A)$$

are different assertions unless the ambient upper-bound sets agree.

Remark (Negated quantified statement).

$$\sup_S A = \sup_{S'} A$$

is not guaranteed by $A \subseteq S \subseteq S'$ alone.

Remark (Negation predicate reading).

$$\text{Supremum}_{S'}(s, A) \wedge \neg \text{Supremum}_S(s, A)$$

is possible when the larger ambient set contains the least upper bound but the smaller one does not.

Remark (Failure modes). The ambient-relative supremum can fail to transfer because the candidate least upper bound may not belong to the smaller ambient set, or because the smaller ambient set may have a different collection of upper bounds.

Remark (Failure mode decomposition).

$$s = \sup_{S'} A, \quad s \notin S \implies s \text{ cannot be the supremum of } A \text{ relative to } S.$$

Remark (Interpretation). Suprema are least upper bounds inside a chosen universe. Enlarging or shrinking that universe can create or remove the point that should serve as the least ceiling.

Remark (Dependencies).

[Upper Bound](#); [Supremum](#).

Theorem (Ambient Existence of Supremum)

Theorem 3.3.56 (Ambient Existence of Supremum). *There are ordered sets $S \subseteq S'$ and a subset $A \subseteq S$ such that A has a supremum relative to S' but has no supremum relative to S . [Go to proof](#).*

Remark (Standard quantified statement).

$$\exists S \subseteq S' \exists A \subseteq S : \sup_{S'} A \text{ exists and } \sup_S A \text{ does not exist.}$$

Remark (Predicate reading).

$$\exists S \subseteq S' \exists A \subseteq S \exists s \in S' : \text{Supremum}_{S'}(s, A) \wedge \neg \exists t \in S \text{Supremum}_S(t, A).$$

Remark (Negated quantified statement).

Every ambient enlargement preserves and reflects existence of suprema.

This assertion is false.

Remark (Negation predicate reading).

$$\forall S \subseteq S' \forall A \subseteq S : [\exists s \in S' \text{Supremum}_{S'}(s, A) \implies \exists t \in S \text{Supremum}_S(t, A)]$$

is false.

Remark (Failure modes). The transfer from S' to S fails when the least upper bound exists in the larger ambient set but is missing from the smaller one. The smaller ambient set may contain upper bounds without containing a least upper bound.

Remark (Failure mode decomposition).

$$s = \sup_{S'} A, \quad s \notin S, \quad \text{and no } t \in S \text{ is least among the upper bounds of } A \text{ in } S.$$

Remark (Interpretation). Completeness is an ambient property. A larger ordered universe can fill a gap that remains absent in a smaller ordered universe.

Remark (Dependencies).

Upper Bound; Supremum; Suprema Depend on the Ambient Set.

Lemma (Rational Gap Example for Suprema)

Lemma 3.3.57 (Rational Gap Example for Suprema). *Let*

$$A = \{q \in \mathbb{Q} : q^2 < 2\}.$$

Then A is nonempty and bounded above in $\mathbb{Q}$, but A has no supremum relative to $\mathbb{Q}$. [Go to proof](#).

Remark (Standard quantified statement).

$$A = \{q \in \mathbb{Q} : q^2 < 2\} \implies \sup_{\mathbb{Q}} A \text{ does not exist.}$$

Remark (Predicate reading).

$$\neg \exists s \in \mathbb{Q} \text{Supremum}_{\mathbb{Q}}(s, A), \quad A = \{q \in \mathbb{Q} : q^2 < 2\}.$$

Remark (Negated quantified statement).

$$\exists s \in \mathbb{Q} (s = \sup_{\mathbb{Q}} A)$$

is false for this set.

Remark (Negation predicate reading).

$$\exists s \in \mathbb{Q} \text{ Supremum}_{\mathbb{Q}}(s, A)$$

is false.

Remark (Failure modes). A rational candidate below $\sqrt{2}$ fails to be an upper bound. A rational candidate above $\sqrt{2}$ is an upper bound but is not least, because another rational upper bound lies between it and $\sqrt{2}$.

Remark (Failure mode decomposition).

$$\begin{cases} s^2 < 2 & \implies s \text{ is not an upper bound of } A, \\ s^2 > 2 & \implies s \text{ is not the least upper bound of } A. \end{cases} \quad (s \in \mathbb{Q})$$

Remark (Interpretation). The set wants $\sqrt{2}$ as its least ceiling, but that number is not rational. The rational ambient set therefore has a visible gap where the supremum should be.

Remark (Dependencies).

Upper Bound; Supremum; Ambient Existence of Supremum.





Figure 3.9: Bound Algebra Cluster Dependency Map. The map originates from the ‘Least Upper Bound Property’ (Axiom) and branches into the dual properties of the continuum. The Negation Exchange serves as the structural bridge between supremum and infimum algebra.



Proofs

Remark (Return). [Return to Theorem](#)

Theorem (Archimedean Property). *For every $x \in \mathbb{R}$, there exists $n \in \mathbb{N}$ such that $n > x$.*

Professional Standard Proof.

Suppose, toward a contradiction, that the conclusion fails. Then there is some $x \in \mathbb{R}$ such that $n \leq x$ for every $n \in \mathbb{N}$; that is, $\mathbb{N}$ is bounded above in $\mathbb{R}$. By completeness, let

$$s = \sup \mathbb{N}.$$

Since $s - 1 < s$, the number $s - 1$ cannot be an upper bound of $\mathbb{N}$; otherwise the leastness of s would give $s \leq s - 1$, impossible. Hence there exists $n \in \mathbb{N}$ such that

$$n > s - 1.$$

Adding 1 gives $n + 1 > s$. But $n + 1 \in \mathbb{N}$, and s is an upper bound of $\mathbb{N}$, so $n + 1 \leq s$. This contradiction shows that $\mathbb{N}$ is not bounded above in $\mathbb{R}$. Therefore for every $x \in \mathbb{R}$ there exists $n \in \mathbb{N}$ such that $n > x$. $\square$

Detailed Learning Proof.

We prove the statement by contradiction. The theorem says that no real number can dominate all natural numbers. Suppose the theorem were false. Then there would be some real number x such that

$$n \leq x \quad \text{for every } n \in \mathbb{N}.$$

Thus $\mathbb{N}$ would be a nonempty subset of $\mathbb{R}$ that is bounded above.

By the least upper bound property of $\mathbb{R}$, the set $\mathbb{N}$ would have a supremum. Let

$$s = \sup \mathbb{N}.$$

So s is an upper bound for $\mathbb{N}$, and it is the least such upper bound.

Now consider the smaller number $s - 1$. Because $s - 1 < s$, it cannot also be an upper bound of $\mathbb{N}$: if it were, then the leastness of s would force $s \leq s - 1$, a contradiction. Since $s - 1$ is not an upper bound, some natural number lies above it. Therefore there exists $n \in \mathbb{N}$ such that

$$n > s - 1.$$

Adding 1 to both sides gives

$$n + 1 > s.$$

But $n + 1$ is again a natural number. Since s is an upper bound of $\mathbb{N}$, every natural number must be at most s , so in particular

$$n + 1 \leq s.$$

This contradicts $n + 1 > s$.

The contradiction came from assuming that some real number bounds all natural numbers above. Hence no such real number exists. Equivalently, for every real number x , there is a natural number n with $n > x$. $\square$

Remark (Proof structure). Assume $\mathbb{N}$ is bounded above and take its supremum s . Since $s - 1$ is smaller than the least upper bound, it fails to be an upper bound, so some $n \in \mathbb{N}$ satisfies $n > s - 1$. Then $n + 1 > s$, contradicting that s is an upper bound for $\mathbb{N}$.

Remark (Dependencies).

- Completeness of the real numbers.
- Supremum.
- Upper bound.

Remark (Return). [Return to Corollary](#)

Corollary (Archimedean Corollary). *For every $\varepsilon \in \mathbb{R}$ such that $\varepsilon > 0$, there exists $n \in \mathbb{N}$ such that $\frac{1}{n} < \varepsilon$.*

Professional Standard Proof.

TODO: professional standard proof for archimedean-reciprocal. □

Detailed Learning Proof.

TODO: detailed learning proof for archimedean-reciprocal. □

Remark (Strategy: The Choice of Leastness Method). Because this corollary is a direct algebraic consequence of the Archimedean Property, we prove it through simple variable transformation ($x = 1/\varepsilon$) rather than using the supremum's ε -characterization directly. In the hierarchy of your repository, the Archimedean Property (which relies on "leastness") does the heavy lifting, allowing the corollary to be a "Side A" algebraic derivation.

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Density of the Rational Numbers in $\mathbb{R}$). *For every $a, b \in \mathbb{R}$ with $a < b$, there exists $q \in \mathbb{Q}$ such that $a < q < b$.*

Professional Standard Proof.

TODO: professional standard proof for density-of-rationals-in-reals. □

Detailed Learning Proof.

TODO: detailed learning proof for density-of-rationals-in-reals. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Density of the Irrational Numbers in $\mathbb{R}$). *For every $a, b \in \mathbb{R}$ with $a < b$, there exists $s \in \mathbb{R} \setminus \mathbb{Q}$ such that $a < s < b$.*

Professional Standard Proof.

TODO: professional standard proof for density-of-irrationals-in-reals. □

Detailed Learning Proof.

TODO: detailed learning proof for density-of-irrationals-in-reals. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Corollary (Rational and Irrational Points in Every Open Interval). *Every open interval in $\mathbb{R}$ contains both a rational number and an irrational number.*

Professional Standard Proof.

TODO: professional standard proof for every-open-interval-contains-rational-and-irrational. □

Detailed Learning Proof.

TODO: detailed learning proof for every-open-interval-contains-rational-and-irrational. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Nested Interval Theorem). *Let $I_n = [a_n, b_n]$ be a sequence of non-empty closed bounded intervals such that $I_{n+1} \subseteq I_n$. Then $\bigcap_{n=1}^{\infty} I_n \neq \emptyset$.*

Professional Standard Proof.

Write $I_n = [a_n, b_n]$. Since the intervals are nonempty, $a_n \leq b_n$ for every n . The set

$$A = \{a_n : n \in \mathbb{N}\}$$

is nonempty and bounded above by b_1 , because each $a_n \in I_n \subseteq I_1$. By completeness, let $x = \sup A$.

Fix $n \in \mathbb{N}$. Since $a_n \in A$, we have $a_n \leq x$. It remains to show $x \leq b_n$. For every $m \in \mathbb{N}$, the nesting of the intervals implies $a_m \leq b_n$: if $m \geq n$, then $I_m \subseteq I_n$, so $a_m \in I_n$ and hence $a_m \leq b_n$; if $m < n$, then $a_m \leq a_n \leq b_n$. Thus b_n is an upper bound of A . Since $x = \sup A$, $x \leq b_n$.

Therefore $a_n \leq x \leq b_n$, so $x \in I_n$. Since n was arbitrary, $x \in \bigcap_{n=1}^{\infty} I_n$, and the intersection is nonempty. $\square$

Detailed Learning Proof.

Each closed interval can be written as

$$I_n = [a_n, b_n].$$

The inclusion $I_{n+1} \subseteq I_n$ says that the intervals are nested: later intervals sit inside earlier ones. In particular, every left endpoint a_n lies inside the first interval $I_1 = [a_1, b_1]$, so

$$a_n \leq b_1 \quad \text{for every } n.$$

Thus the set of all left endpoints,

$$A = \{a_n : n \in \mathbb{N}\},$$

is nonempty and bounded above. By the least upper bound property of $\mathbb{R}$, this set has a supremum. Call it

$$x = \sup A.$$

We now prove that this single number x belongs to every interval I_n . Fix an arbitrary n . Since a_n is one of the elements of A , and x is an upper bound of A , we have

$$a_n \leq x.$$

So x is not to the left of I_n . It remains to prove that x is not to the right of I_n , meaning $x \leq b_n$.

To show $x \leq b_n$, it is enough to show that b_n is an upper bound for the set A of left endpoints. Take any $a_m \in A$. If $m \geq n$, then $I_m \subseteq I_n$, so $a_m \in I_n = [a_n, b_n]$, and therefore $a_m \leq b_n$. If $m < n$, then $I_n \subseteq I_m$, so the left endpoint of the earlier interval lies no farther right than the left endpoint of the later interval: $a_m \leq a_n$. Since $a_n \leq b_n$, this also gives $a_m \leq b_n$.

Thus every element of A is at most b_n , so b_n is an upper bound of A . Because x is the least upper bound of A , it follows that

$$x \leq b_n.$$

Combining the two inequalities gives

$$a_n \leq x \leq b_n,$$

so $x \in I_n$.

The index n was arbitrary, so x belongs to every interval in the nested sequence. Hence

$$x \in \bigcap_{n=1}^{\infty} I_n,$$

and the intersection is nonempty. □

Remark (Proof structure). Use completeness on the set of left endpoints. Nesting makes every right endpoint an upper bound for those left endpoints, so the supremum of the left endpoints lies inside every interval.

Remark (Dependencies).

- Completeness of the real numbers.
- Supremum.
- Upper bound.

Remark (Return). [Return to Theorem](#)

Theorem (Least Upper Bound Property Implies Existence of Suprema). *Let S be a partially ordered set. If S has the Least Upper Bound Property, then every nonempty subset of S that is bounded above in S has a supremum in S .*

Professional Standard Proof.

TODO: professional standard proof for lub-property-implies-existence-of-suprema. □

Detailed Learning Proof.

TODO: detailed learning proof for lub-property-implies-existence-of-suprema. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Upper Bound Property of Supremum). *Let $A \subseteq \mathbb{R}$ and let $s \in \mathbb{R}$. If $s = \sup A$, then*

$$x \leq s \quad \text{for every } x \in A.$$

Professional Standard Proof.

TODO: professional standard proof for upper-bound-property-of-supremum. □

Detailed Learning Proof.

TODO: detailed learning proof for upper-bound-property-of-supremum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Uniqueness of the Minimum). *If a set has a minimum, then that minimum is unique.*

Professional Standard Proof.

TODO: professional standard proof for uniqueness-of-the-minimum. □

Detailed Learning Proof.

TODO: detailed learning proof for uniqueness-of-the-minimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Minimum Implies Infimum). *If a nonempty ordered set has a minimum, then that minimum is its infimum.*

Professional Standard Proof.

TODO: professional standard proof for minimum-implies-infimum. □

Detailed Learning Proof.

TODO: detailed learning proof for minimum-implies-infimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Infimum in the Set is the Minimum). *If the infimum of a nonempty set belongs to the set, then it is the minimum.*

Professional Standard Proof.

TODO: professional standard proof for infimum-in-the-set-is-the-minimum. □

Detailed Learning Proof.

TODO: detailed learning proof for infimum-in-the-set-is-the-minimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Uniqueness of the Infimum). *If a set has an infimum, then that infimum is unique.*

Professional Standard Proof.

TODO: professional standard proof for uniqueness-of-the-infimum. □

Detailed Learning Proof.

TODO: detailed learning proof for uniqueness-of-the-infimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Lower Bound Property of the Infimum). *If i is the infimum of A , then i is a lower bound of A .*

Professional Standard Proof.

TODO: professional standard proof for lower-bound-property-of-infimum. □

Detailed Learning Proof.

TODO: detailed learning proof for lower-bound-property-of-infimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Lemma (Subset Inclusion Preserves Lower Bounds). *If A is a subset of B and ℓ is a lower bound of B , then ℓ is a lower bound of A .*

Professional Standard Proof.

TODO: professional standard proof for subset-inclusion-preserves-lower-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for subset-inclusion-preserves-lower-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Infimum Is Monotone Under Inclusion). *If A is a subset of B and both infima exist, then $\inf B$ is less than or equal to $\inf A$.*

Professional Standard Proof.

TODO: professional standard proof for infimum-is-monotone-under-inclusion. □

Detailed Learning Proof.

TODO: detailed learning proof for infimum-is-monotone-under-inclusion. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Lower Bound Comparison with the Infimum). *For a nonempty set with infimum i , lower bounds are exactly the numbers less than or equal to i .*

Professional Standard Proof.

TODO: professional standard proof for lower-bound-comparison-with-the-infimum. □

Detailed Learning Proof.

TODO: detailed learning proof for lower-bound-comparison-with-the-infimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Every Element Lies Above the Infimum). *If i is the infimum of A , then every element of A lies above i .*

Professional Standard Proof.

TODO: professional standard proof for every-element-lies-above-the-infimum. □

Detailed Learning Proof.

TODO: detailed learning proof for every-element-lies-above-the-infimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Greatest Lower Bound Property Implies Existence of Infima). *Every nonempty subset of R that is bounded below has an infimum in R .*

Professional Standard Proof.

TODO: professional standard proof for glb-property-implies-existence-of-infima. □

Detailed Learning Proof.

TODO: detailed learning proof for glb-property-implies-existence-of-infima. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Lemma](#)

Lemma (Union Preserves Upper Bounds). *If u_A is an upper bound of A and u_B is an upper bound of B , then $\max\{u_A, u_B\}$ is an upper bound of $A \cup B$.*

Professional Standard Proof.

TODO: professional standard proof for union-preserves-upper-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for union-preserves-upper-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Lemma](#)

Lemma (Union Preserves Lower Bounds). *If ℓ_A is a lower bound of A and ℓ_B is a lower bound of B , then $\min\{\ell_A, \ell_B\}$ is a lower bound of $A \cup B$.*

Professional Standard Proof.

TODO: professional standard proof for union-preserves-lower-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for union-preserves-lower-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Proposition](#)

Proposition (Union Is Bounded Above Exactly When Both Pieces Are). *For nonempty $A, B \subseteq \mathbb{R}$, $A \cup B$ is bounded above if and only if A and B are bounded above.*

Professional Standard Proof.

TODO: professional standard proof for union-bounded-above-iff-pieces-bounded-above. $\square$

Detailed Learning Proof.

TODO: detailed learning proof for union-bounded-above-iff-pieces-bounded-above. $\square$

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Proposition](#)

Proposition (Union Is Bounded Below Exactly When Both Pieces Are). *For nonempty $A, B \subseteq \mathbb{R}$, $A \cup B$ is bounded below if and only if A and B are bounded below.*

Professional Standard Proof.

TODO: professional standard proof for union-bounded-below-iff-pieces-bounded-below. $\square$

Detailed Learning Proof.

TODO: detailed learning proof for union-bounded-below-iff-pieces-bounded-below. $\square$

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Corollary](#)

Corollary (Union Is Bounded Exactly When Both Pieces Are). *For nonempty $A, B \subseteq \mathbb{R}$, $A \cup B$ is bounded if and only if A and B are bounded.*

Professional Standard Proof.

TODO: professional standard proof for union-bounded-iff-pieces-bounded. □

Detailed Learning Proof.

TODO: detailed learning proof for union-bounded-iff-pieces-bounded. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Lemma](#)

Lemma (Subsets Preserve Upper Bounds). *If $C \subseteq A \subseteq \mathbb{R}$ and u is an upper bound of A , then u is an upper bound of C .*

Professional Standard Proof.

TODO: professional standard proof for subsets-preserve-upper-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for subsets-preserve-upper-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Lemma](#)

Lemma (Subsets Preserve Lower Bounds). *If $C \subseteq A \subseteq \mathbb{R}$ and ℓ is a lower bound of A , then ℓ is a lower bound of C .*

Professional Standard Proof.

TODO: professional standard proof for subsets-preserve-lower-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for subsets-preserve-lower-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Corollary](#)

Corollary (Intersections Inherit Bounds). *Every upper bound of A is an upper bound of $A \cap B$, and every lower bound of A is a lower bound of $A \cap B$.*

Professional Standard Proof.

TODO: professional standard proof for intersections-inherit-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for intersections-inherit-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Corollary](#)

Corollary (Differences Inherit Bounds). *Every upper bound of A is an upper bound of $A \setminus B$, and every lower bound of A is a lower bound of $A \setminus B$.*

Professional Standard Proof.

TODO: professional standard proof for differences-inherit-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for differences-inherit-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Corollary](#)

Corollary (Complements Inherit Ambient Bounds). *For $A \subseteq E \subseteq \mathbb{R}$, every upper bound of E is an upper bound of $E \setminus A$, and every lower bound of E is a lower bound of $E \setminus A$.*

Professional Standard Proof.

TODO: professional standard proof for complements-inherit-ambient-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for complements-inherit-ambient-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Supremum of the Pointwise Maximum Set). *For nonempty bounded-above $A, B \subseteq \mathbb{R}$, $\sup \max(A, B) = \max\{\sup A, \sup B\}$.*

Professional Standard Proof.

TODO: professional standard proof for supremum-pointwise-maximum-set. □

Detailed Learning Proof.

TODO: detailed learning proof for supremum-pointwise-maximum-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Infimum of the Pointwise Maximum Set). *For nonempty bounded-below $A, B \subseteq \mathbb{R}$, $\inf \max(A, B) = \max\{\inf A, \inf B\}$.*

Professional Standard Proof.

TODO: professional standard proof for infimum-pointwise-maximum-set. □

Detailed Learning Proof.

TODO: detailed learning proof for infimum-pointwise-maximum-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Supremum of the Pointwise Minimum Set). *For nonempty bounded-above $A, B \subseteq \mathbb{R}$, $\sup \min(A, B) = \min\{\sup A, \sup B\}$.*

Professional Standard Proof.

TODO: professional standard proof for supremum-pointwise-minimum-set. □

Detailed Learning Proof.

TODO: detailed learning proof for supremum-pointwise-minimum-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Infimum of the Pointwise Minimum Set). *For nonempty bounded-below $A, B \subseteq \mathbb{R}$, $\inf \min(A, B) = \min\{\inf A, \inf B\}$.*

Professional Standard Proof.

TODO: professional standard proof for infimum-pointwise-minimum-set. □

Detailed Learning Proof.

TODO: detailed learning proof for infimum-pointwise-minimum-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Increasing Image Preserves Suprema). *If f is increasing and continuous at $\sup A$, then $\sup f(A) = f(\sup A)$.*

Professional Standard Proof.

TODO: professional standard proof for increasing-image-preserves-suprema. □

Detailed Learning Proof.

TODO: detailed learning proof for increasing-image-preserves-suprema. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Increasing Image Preserves Infima). *If f is increasing and continuous at $\inf A$, then $\inf f(A) = f(\inf A)$.*

Professional Standard Proof.

TODO: professional standard proof for increasing-image-preserves-infima. □

Detailed Learning Proof.

TODO: detailed learning proof for increasing-image-preserves-infima. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Decreasing Image Sends Infimum to Supremum). *If f is decreasing and continuous at $\inf A$, then $\sup f(A) = f(\inf A)$.*

Professional Standard Proof.

TODO: professional standard proof for decreasing-image-infimum-to-supremum. □

Detailed Learning Proof.

TODO: detailed learning proof for decreasing-image-infimum-to-supremum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Decreasing Image Sends Supremum to Infimum). *If f is decreasing and continuous at $\sup A$, then $\inf f(A) = f(\sup A)$.*

Professional Standard Proof.

TODO: professional standard proof for decreasing-image-supremum-to-infimum. □

Detailed Learning Proof.

TODO: detailed learning proof for decreasing-image-supremum-to-infimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Increasing Inverse Preserves Suprema). *If f^{-1} is increasing and continuous at $\sup B$, then $\sup f^{-1}(B) = f^{-1}(\sup B)$.*

Professional Standard Proof.

TODO: professional standard proof for increasing-inverse-preserves-suprema. □

Detailed Learning Proof.

TODO: detailed learning proof for increasing-inverse-preserves-suprema. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Increasing Inverse Preserves Infima). *If f^{-1} is increasing and continuous at $\inf B$, then $\inf f^{-1}(B) = f^{-1}(\inf B)$.*

Professional Standard Proof.

TODO: professional standard proof for increasing-inverse-preserves-infima. □

Detailed Learning Proof.

TODO: detailed learning proof for increasing-inverse-preserves-infima. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Decreasing Inverse Sends Infimum to Supremum). *If f^{-1} is decreasing and continuous at $\inf B$, then $\sup f^{-1}(B) = f^{-1}(\inf B)$.*

Professional Standard Proof.

TODO: professional standard proof for decreasing-inverse-infimum-to-supremum. □

Detailed Learning Proof.

TODO: detailed learning proof for decreasing-inverse-infimum-to-supremum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Decreasing Inverse Sends Supremum to Infimum). *If f^{-1} is decreasing and continuous at $\sup B$, then $\inf f^{-1}(B) = f^{-1}(\sup B)$.*

Professional Standard Proof.

TODO: professional standard proof for decreasing-inverse-supremum-to-infimum. □

Detailed Learning Proof.

TODO: detailed learning proof for decreasing-inverse-supremum-to-infimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Proposition](#)

Proposition (Translation Preserves Upper Bounds). *If u is an upper bound of A , then $u + c$ is an upper bound of $A + c$.*

Professional Standard Proof.

TODO: professional standard proof for translation-preserves-upper-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for translation-preserves-upper-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Proposition](#)

Proposition (Translation Preserves Lower Bounds). *If ℓ is a lower bound of A , then $\ell + c$ is a lower bound of $A + c$.*

Professional Standard Proof.

TODO: professional standard proof for translation-preserves-lower-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for translation-preserves-lower-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Proposition](#)

Proposition (Positive Dilation Preserves Upper Bounds). *If $\lambda > 0$ and u is an upper bound of A , then λu is an upper bound of λA .*

Professional Standard Proof.

TODO: professional standard proof for positive-dilation-preserves-upper-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for positive-dilation-preserves-upper-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Proposition](#)

Proposition (Positive Dilation Preserves Lower Bounds). *If $\lambda > 0$ and ℓ is a lower bound of A , then $\lambda\ell$ is a lower bound of λA .*

Professional Standard Proof.

TODO: professional standard proof for positive-dilation-preserves-lower-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for positive-dilation-preserves-lower-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Proposition](#)

Proposition (Negative Dilation Sends Lower Bounds to Upper Bounds). *If $\lambda < 0$ and ℓ is a lower bound of A , then $\lambda\ell$ is an upper bound of λA .*

Professional Standard Proof.

TODO: professional standard proof for negative-dilation-sends-lower-to-upper-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for negative-dilation-sends-lower-to-upper-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Proposition](#)

Proposition (Negative Dilation Sends Upper Bounds to Lower Bounds). *If $\lambda < 0$ and u is an upper bound of A , then λu is a lower bound of λA .*

Professional Standard Proof.

TODO: professional standard proof for negative-dilation-sends-upper-to-lower-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for negative-dilation-sends-upper-to-lower-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Corollary](#)

Corollary (Reflection Swaps Upper and Lower Bounds). *If ℓ is a lower bound of A , then $-\ell$ is an upper bound of $-A$. If u is an upper bound of A , then $-u$ is a lower bound of $-A$.*

Professional Standard Proof.

TODO: professional standard proof for reflection-swaps-upper-lower-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for reflection-swaps-upper-lower-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Uniqueness of the Maximum). *If a set $A \subseteq \mathbb{R}$ has a maximum, then that maximum is unique.*

Professional Standard Proof.

TODO: professional standard proof for uniqueness-of-the-maximum. □

Detailed Learning Proof.

TODO: detailed learning proof for uniqueness-of-the-maximum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Uniqueness of the Supremum). *If a set $A \subseteq \mathbb{R}$ has a supremum, then that supremum is unique.*

Professional Standard Proof.

TODO: professional standard proof for uniqueness-of-the-supremum. □

Detailed Learning Proof.

TODO: detailed learning proof for uniqueness-of-the-supremum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Remark (Proof vault). [Handwritten proof record](#)

Proposition (Maximum Implies Supremum). *If a set $A \subseteq \mathbb{R}$ has a maximum m , then m is also the supremum of A .*

Professional Standard Proof.

Suppose m is a maximum of A . Then $m \in A$ and

$$x \leq m$$

for every $x \in A$. Hence m is an upper bound of A .

Now let u be any upper bound of A . Since $m \in A$, the upper-bound property of u gives $m \leq u$. Thus every upper bound u of A satisfies $m \leq u$. Therefore m is the least upper bound of A , so

$$m = \sup A.$$

□

Detailed Learning Proof.

We prove that m satisfies the two defining properties of the supremum of A .

First, since m is a maximum of A , the definition of maximum gives

$$m \in A \quad \text{and} \quad x \leq m \quad \text{for every } x \in A.$$

The second condition says exactly that m is an upper bound of A .

Second, we show that m is the least upper bound. Let u be an arbitrary upper bound of A . Because u is an upper bound, every element of A is less than or equal to u . Since $m \in A$, we may apply this to the element m itself and obtain

$$m \leq u.$$

Thus every upper bound u of A lies above m .

So m is an upper bound of A , and every upper bound of A is at least m . By the definition of supremum,

$$m = \sup A.$$

□

Remark (Proof structure). A maximum is already an upper bound because it dominates every element of the set. It is least among upper bounds because it is itself an element of the set: any upper bound must bound m too.

Remark (Dependencies).

- [Maximum](#)
- [Upper Bound](#)
- [Supremum](#)

Remark (Return). [Return to Proposition](#)

Proposition (Supremum in the Set Is the Maximum). *Let $s = \sup A$. If $s \in A$, then s is the maximum of A .*

Professional Standard Proof.

TODO: professional standard proof for supremum-in-the-set-is-the-maximum. □

Detailed Learning Proof.

TODO: detailed learning proof for supremum-in-the-set-is-the-maximum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Lemma](#)

Lemma (Subset Inclusion Preserves Upper Bounds). *Let $A \subseteq B \subseteq \mathbb{R}$. If u is an upper bound of B , then u is an upper bound of A .*

Professional Standard Proof.

TODO: professional standard proof for subset-inclusion-preserves-upper-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for subset-inclusion-preserves-upper-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Supremum Is Monotone Under Inclusion). *Let A and B be non-empty subsets of $\mathbb{R}$ that are bounded above. If $A \subseteq B$, then $\sup A \leq \sup B$.*

Professional Standard Proof.

TODO: professional standard proof for supremum-is-monotone-under-inclusion. □

Detailed Learning Proof.

TODO: detailed learning proof for supremum-is-monotone-under-inclusion. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Proposition](#)

Proposition (Upper Bound Comparison with the Supremum). *Let $A \subseteq \mathbb{R}$ be non-empty and bounded above, and let $s = \sup A$. For any $u \in \mathbb{R}$, u is an upper bound of A if and only if $s \leq u$.*

Professional Standard Proof.

TODO: professional standard proof for upper-bound-comparison-with-the-supremum. □

Detailed Learning Proof.

TODO: detailed learning proof for upper-bound-comparison-with-the-supremum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Proposition](#)

Remark (Proof vault). [Handwritten proof record](#)

Proposition (Every Element Lies Below the Supremum). *If $s = \sup A$, then for every $x \in A$, $x \leq s$.*

Professional Standard Proof.

Since $s = \sup A$, the definition of supremum says, in particular, that s is an upper bound of A . Hence every element of A lies below s : for every $x \in A$,

$$x \leq s.$$

□

Detailed Learning Proof.

Let $x \in A$ be arbitrary. We need to prove that $x \leq s$.

The hypothesis $s = \sup A$ means that s is the least upper bound of A . In particular, before using the “least” part, we know that s is an upper bound of A . By the definition of upper bound, this means that every element of A is less than or equal to s . Since $x \in A$, it follows that

$$x \leq s.$$

Because $x \in A$ was arbitrary, $x \leq s$ for every $x \in A$.

□

Remark (Proof structure). This proof only unpacks the definition of supremum. The conclusion uses the upper-bound part of $s = \sup A$; the minimality of s among upper bounds is not needed.

Remark (Dependencies).

- [Upper Bound](#)
- [Supremum](#)

Remark (Return). [Return to Theorem](#)

Remark (Proof vault). [Handwritten proof record](#)

Theorem (Infimum-Supremum Comparison). *For any non-empty bounded set $A \subseteq \mathbb{R}$, the infimum of A is less than or equal to the supremum of A .*

Professional Standard Proof.

Because A is nonempty, choose $x \in A$. Since $\inf A$ is a lower bound of A , we have

$$\inf A \leq x.$$

Since $\sup A$ is an upper bound of A , we have

$$x \leq \sup A.$$

By transitivity of the order on $\mathbb{R}$,

$$\inf A \leq \sup A.$$

□

Detailed Learning Proof.

The set A is nonempty, so there exists an element $x \in A$.

By definition, $\inf A$ is the greatest lower bound of A . In particular, it is a lower bound of A . Thus every element of A lies above $\inf A$, so for our chosen $x \in A$,

$$\inf A \leq x.$$

Similarly, $\sup A$ is the least upper bound of A . In particular, it is an upper bound of A . Thus every element of A lies below $\sup A$, so for the same $x \in A$,

$$x \leq \sup A.$$

Combining the two inequalities gives

$$\inf A \leq x \leq \sup A.$$

Therefore

$$\inf A \leq \sup A.$$

□

Remark (Proof structure). Choose one element of the nonempty set. The lower-bound part of the infimum definition places $\inf A$ below that element, and the upper-bound part of the supremum definition places the element below $\sup A$. Transitivity finishes the comparison.

Remark (Dependencies).

- [Lower Bound](#)
- [Upper Bound](#)
- [Infimum](#)
- [Supremum](#)
- [Real Ordered Field](#)

Remark (Return). [Return to Theorem](#)

Proposition (Order Comparison of Supremum). *Let $A, B \subseteq \mathbb{R}$ be non-empty sets that are bounded above. If for every $x \in A$, there exists $y \in B$ such that $x \leq y$, then $\sup A \leq \sup B$.*

Professional Standard Proof.

TODO: professional standard proof for order-comparison-of-suprema. □

Detailed Learning Proof.

TODO: detailed learning proof for order-comparison-of-suprema. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Translation Invariance of the Supremum). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded above, and let $c \in \mathbb{R}$. Then*

$$\sup(A + c) = \sup A + c, \quad A + c = \{a + c : a \in A\}.$$

Professional Standard Proof.

TODO: professional standard proof for translation-invariance-supremum. □

Detailed Learning Proof.

TODO: detailed learning proof for translation-invariance-supremum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Translation Invariance of the Infimum). *For nonempty A bounded below, $\inf(A + c) = \inf A + c$.*

Professional Standard Proof.

TODO: professional standard proof for translation-invariance-infimum. □

Detailed Learning Proof.

TODO: detailed learning proof for translation-invariance-infimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Remark (Proof vault). [Handwritten proof record](#)

Theorem (Scalar Multiplication by a Positive Scalar). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded above, and let $k > 0$. Then*

$$\sup(kA) = k \sup A, \quad kA = \{ka : a \in A\}.$$

Professional Standard Proof.

Let $s = \sup A$. Since s is an upper bound of A , we have $a \leq s$ for every $a \in A$. Because $k > 0$, multiplication by k preserves order, so $ka \leq ks$ for every $a \in A$. Thus ks is an upper bound of kA .

Let $t = \sup(kA)$. Since ks is an upper bound of kA , the least-upper-bound property gives $t \leq ks$. Conversely, since t is an upper bound of kA , we have $ka \leq t$ for every $a \in A$. Dividing by $k > 0$ gives $a \leq t/k$ for every $a \in A$, so t/k is an upper bound of A . Since $s = \sup A$, we get $s \leq t/k$, and multiplying by $k > 0$ gives $ks \leq t$. Hence $t = ks$, so $\sup(kA) = k \sup A$. $\square$

Detailed Learning Proof.

Set $s = \sup A$. We prove that ks is exactly the least upper bound of kA .

First we show that ks is an upper bound of kA . Let $y \in kA$. By the definition of kA , there is some $a \in A$ such that $y = ka$. Since $s = \sup A$, every element of A is at most s , so $a \leq s$. Because $k > 0$, multiplication by k preserves the inequality, and therefore $ka \leq ks$. Hence $y \leq ks$. Since $y \in kA$ was arbitrary, ks is an upper bound of kA .

Now let $t = \sup(kA)$. From the preceding paragraph, ks is an upper bound of kA , so the defining leastness of t gives

$$t \leq ks.$$

It remains to prove the reverse inequality. Since t is an upper bound of kA , for each $a \in A$ the element ka belongs to kA and therefore $ka \leq t$. Because $k > 0$, dividing by k preserves order, so

$$a \leq \frac{t}{k}$$

for every $a \in A$. Thus t/k is an upper bound of A . Since $s = \sup A$, s is less than or equal to every upper bound of A , so

$$s \leq \frac{t}{k}.$$

Multiplying by $k > 0$ gives $ks \leq t$.

Together, $t \leq ks$ and $ks \leq t$ imply $t = ks$. Since $t = \sup(kA)$ and $s = \sup A$, this is precisely

$$\sup(kA) = k \sup A.$$

$\square$

Remark (Proof structure). Show that the proposed value $k \sup A$ is an upper bound of kA , then use the least-upper-bound property in the reverse direction by dividing an arbitrary upper bound of kA by the positive scalar k .

Remark (Dependencies).

- Upper bound.
- Supremum.
- Upper Bound Comparison with the Supremum.
- Multiplication by a positive scalar preserves non-strict order.

Remark (Return). [Return to Theorem](#)

Theorem (Positive Scalar Multiplication of the Infimum). *If $k > 0$ and A is nonempty and bounded below, then $\inf(kA) = k \inf A$.*

Professional Standard Proof.

TODO: professional standard proof for positive-scalar-mult-infimum. □

Detailed Learning Proof.

TODO: detailed learning proof for positive-scalar-mult-infimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Remark (Proof vault). [Handwritten proof record](#)

Theorem (Scalar Multiplication by a Negative Scalar). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded below, and let $k < 0$. Then*

$$\sup(kA) = k \inf A, \quad kA = \{ka : a \in A\}.$$

Professional Standard Proof.

Let $N = -k$. Since $k < 0$, we have $N > 0$. Also

$$kA = \{ka : a \in A\} = \{N(-a) : a \in A\} = N(-A).$$

Because A is bounded below, the set $-A$ is bounded above. By scalar multiplication by a positive scalar,

$$\sup(N(-A)) = N \sup(-A).$$

By the theorem that negation exchanges supremum and infimum,

$$\sup(-A) = -\inf A.$$

Therefore

$$\sup(kA) = \sup(N(-A)) = N \sup(-A) = N(-\inf A) = -N \inf A.$$

Since $N = -k$, we have $-N = k$, and hence $\sup(kA) = k \inf A$. □

Detailed Learning Proof.

We reduce the negative-scalar case to two results that are already known: positive scalar multiplication of suprema, and the exchange of supremum and infimum under negation.

Let $N = -k$. The hypothesis $k < 0$ implies $N > 0$. For every $a \in A$,

$$ka = (-N)a = N(-a).$$

Therefore the image of A under multiplication by k is the same set as the image of $-A$ under multiplication by the positive scalar N :

$$kA = N(-A).$$

Since A is bounded below, negating all of its elements produces a set $-A = \{-a : a \in A\}$ that is bounded above. The positive-scalar theorem applies to the nonempty bounded-above set $-A$ and the positive scalar N , so

$$\sup(N(-A)) = N \sup(-A).$$

The negation theorem gives

$$\sup(-A) = -\inf A.$$

Substituting this into the preceding equality yields

$$\sup(N(-A)) = N(-\inf A).$$

Finally, because $kA = N(-A)$, the two sets have the same supremum. Hence

$$\sup(kA) = \sup(N(-A)) = N(-\inf A) = -N \inf A.$$

Since $N = -k$, we have $-N = k$. Thus

$$\sup(kA) = k \inf A,$$

as required. □

Remark (Proof structure). Convert the negative multiplier k into a positive multiplier $N = -k$ applied to the negated set $-A$. Then combine the positive-scalar supremum formula with the theorem that negation exchanges supremum and infimum.

Remark (Dependencies).

- [Scalar Multiplication by a Positive Scalar](#).
- [Negation Exchanges Supremum and Infimum](#).
- [Supremum](#).
- [Infimum](#).

Remark (Return). [Return to Theorem](#)

Theorem (Negative Scalar Multiplication of the Infimum). *If $k < 0$ and A is nonempty and bounded above, then $\inf(kA) = k \sup A$.*

Professional Standard Proof.

TODO: professional standard proof for negative-scalar-mult-infimum. □

Detailed Learning Proof.

TODO: detailed learning proof for negative-scalar-mult-infimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Remark (Proof vault). [Handwritten proof record](#)

Theorem (Negation Exchanges Supremum and Infimum). *Let $A \subseteq \mathbb{R}$ be a non-empty set. If A is bounded below, then $\sup(-A) = -\inf A$.*

Professional Standard Proof.

Let $i = \inf A$. We prove that $-i$ is the supremum of $-A$.

First let $y \in -A$. Then $y = -x$ for some $x \in A$. Since i is a lower bound of A , we have $i \leq x$. Multiplying by -1 reverses the inequality, so

$$y = -x \leq -i.$$

Thus $-i$ is an upper bound of $-A$. In particular, $-A$ is bounded above.

Now let u be any upper bound of $-A$. For every $x \in A$, the element $-x$ belongs to $-A$, so

$$-x \leq u.$$

Multiplying by -1 gives

$$-u \leq x$$

for every $x \in A$. Hence $-u$ is a lower bound of A . Since $i = \inf A$ is the greatest lower bound of A , we have $-u \leq i$. Multiplying by -1 again gives

$$-i \leq u.$$

Therefore every upper bound u of $-A$ lies above $-i$. Since $-i$ is itself an upper bound of $-A$, it is the least upper bound of $-A$. Hence

$$\sup(-A) = -i = -\inf A.$$

□

Detailed Learning Proof.

Set

$$i = \inf A.$$

We want to show that $-i$ is the least upper bound of the reflected set

$$-A = \{-a : a \in A\}.$$

First we show that $-i$ is an upper bound of $-A$. Take an arbitrary element $y \in -A$. By the definition of $-A$, there is some $x \in A$ such that

$$y = -x.$$

Because $i = \inf A$, the number i is a lower bound of A . Thus

$$i \leq x.$$

After multiplying by -1 , the inequality reverses:

$$-x \leq -i.$$

Since $y = -x$, this gives $y \leq -i$. Therefore every element of $-A$ is less than or equal to $-i$, so $-i$ is an upper bound of $-A$. This also shows that $-A$ is bounded above.

It remains to show that $-i$ is the least upper bound. Let u be any upper bound of $-A$. Then every element of $-A$ is less than or equal to u . In particular, for each $x \in A$, the element $-x \in -A$, so

$$-x \leq u.$$

Multiplying by -1 reverses the inequality:

$$-u \leq x.$$

This holds for every $x \in A$, so $-u$ is a lower bound of A . Since $i = \inf A$ is the greatest lower bound of A , every lower bound of A is less than or equal to i . Hence

$$-u \leq i.$$

Multiplying by -1 one more time gives

$$-i \leq u.$$

Thus every upper bound u of $-A$ lies above $-i$.

We have shown that $-i$ is an upper bound of $-A$ and that every upper bound of $-A$ is at least $-i$. By the definition of supremum,

$$\sup(-A) = -i = -\inf A.$$

□

Remark (Strategy: The Choice of Leastness Method). In this proof, we establish the "leastness" property by comparing $-i$ against an arbitrary upper bound u (the "Side A" method) rather than using the ε -characterization ("Side B"). While the ε -characterization is mathematically equivalent, it often requires "double negation" logic here—showing that $a < i + \varepsilon$ implies $-a > -i - \varepsilon$. When a proof already involves a set transformation like negation, it is often procedurally cleaner to use the direct definition of the supremum to avoid the mental overhead of tracking flipped inequalities across a tolerance variable.

Remark (Proof structure). Negation reverses order. Lower bounds of A become upper bounds of $-A$, and the greatest lower bound becomes the least upper bound after multiplying all inequalities by -1 .

Remark (Dependencies).

- [Lower Bound](#)
- [Upper Bound](#)
- [Infimum](#)
- [Supremum](#)
- [Real Ordered Field](#)

Remark (Return). [Return to Theorem](#)

Theorem (Supremum of a Sum Set). *Let $A, B \subseteq \mathbb{R}$ be non-empty sets bounded above. Define $A + B = \{a + b : a \in A, b \in B\}$. Then $\sup(A + B) = \sup A + \sup B$.*

Professional Standard Proof.

TODO: professional standard proof for supremum-sum-set. □

Detailed Learning Proof.

TODO: detailed learning proof for supremum-sum-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Infimum of a Sum Set). *For nonempty bounded-below sets A, B , $\inf(A + B) = \inf A + \inf B$.*

Professional Standard Proof.

TODO: professional standard proof for infimum-sum-set. □

Detailed Learning Proof.

TODO: detailed learning proof for infimum-sum-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Supremum of a Difference Set). *Let $A, B \subseteq \mathbb{R}$ be nonempty, with A bounded above and B bounded below. Then*

$$\sup(A - B) = \sup A - \inf B, \quad A - B = \{a - b : a \in A, b \in B\}.$$

Professional Standard Proof.

TODO: professional standard proof for supremum-difference-set. □

Detailed Learning Proof.

TODO: detailed learning proof for supremum-difference-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Infimum of a Difference Set). *For nonempty A, B , with A bounded below and B bounded above, $\inf(A - B) = \inf A - \sup B$.*

Professional Standard Proof.

TODO: professional standard proof for infimum-difference-set. □

Detailed Learning Proof.

TODO: detailed learning proof for infimum-difference-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Supremum of a Dilation). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded, and let $\lambda \in \mathbb{R}$. Then*

$$\sup(\lambda A) = \max\{\lambda \sup A, \lambda \inf A\}, \quad \lambda A = \{\lambda a : a \in A\}.$$

Professional Standard Proof.

TODO: professional standard proof for supremum-dilation. □

Detailed Learning Proof.

TODO: detailed learning proof for supremum-dilation. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Supremum of the Absolute-Value Image). *Let $A \subseteq \mathbb{R}$ be nonempty and bounded. Then*

$$\sup |A| = \max\{|\sup A|, |\inf A|\}, \quad |A| = \{|a| : a \in A\}.$$

Professional Standard Proof.

TODO: professional standard proof for supremum-absolute-value-image. □

Detailed Learning Proof.

TODO: detailed learning proof for supremum-absolute-value-image. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Supremum of a Reciprocal Set). *For a nonempty bounded set separated from zero,*

$$\sup(A^{-1}) = (\inf A)^{-1}.$$

Professional Standard Proof.

TODO: professional standard proof for supremum-reciprocal-set. □

Detailed Learning Proof.

TODO: detailed learning proof for supremum-reciprocal-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Infimum of a Reciprocal Set). *For a nonempty bounded set separated from zero,*

$$\inf(A^{-1}) = (\sup A)^{-1}.$$

Professional Standard Proof.

TODO: professional standard proof for infimum-reciprocal-set. □

Detailed Learning Proof.

TODO: detailed learning proof for infimum-reciprocal-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Supremum of a Product Set). *For nonempty bounded A, B , $\sup(AB)$ is the maximum of the four endpoint products.*

Professional Standard Proof.

TODO: professional standard proof for supremum-product-set. □

Detailed Learning Proof.

TODO: detailed learning proof for supremum-product-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Infimum of a Product Set). *For nonempty bounded A, B , $\inf(AB)$ is the minimum of the four endpoint products.*

Professional Standard Proof.

TODO: professional standard proof for infimum-product-set. □

Detailed Learning Proof.

TODO: detailed learning proof for infimum-product-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Supremum of a Quotient Set). *If A, B are nonempty and bounded and B is separated from zero, then $\sup(A/B)$ is the maximum of the four endpoint quotients.*

Professional Standard Proof.

TODO: professional standard proof for supremum-quotient-set. □

Detailed Learning Proof.

TODO: detailed learning proof for supremum-quotient-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Infimum of a Quotient Set). *If A, B are nonempty and bounded and B is separated from zero, then $\inf(A/B)$ is the minimum of the four endpoint quotients.*

Professional Standard Proof.

TODO: professional standard proof for infimum-quotient-set. □

Detailed Learning Proof.

TODO: detailed learning proof for infimum-quotient-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Bounds for Suprema and Infima Under Set Addition). *Let $A, B \subseteq \mathbb{R}$ be nonempty and bounded. Then*

$$\inf A + \inf B \leq \inf(A + B) \quad \text{and} \quad \sup(A + B) \leq \sup A + \sup B.$$

Professional Standard Proof.

TODO: professional standard proof for bounds-sum-set. □

Detailed Learning Proof.

TODO: detailed learning proof for bounds-sum-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Upper Bounds Depend on the Ambient Order). *Let $(S, \leq_S)$ and $(T, \leq_T)$ be ordered sets, and let $A \subseteq S \cap T$. Whether u is an upper bound of A must be interpreted relative to the chosen ambient order.*

Professional Standard Proof.

TODO: professional standard proof for upper-bounds-ambient-order. □

Detailed Learning Proof.

TODO: detailed learning proof for upper-bounds-ambient-order. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Suprema Depend on the Ambient Set). *Let $A \subseteq S \subseteq S'$ be subsets of an ordered set. The supremum of A relative to S , if it exists, need not agree with the supremum of A relative to S' .*

Professional Standard Proof.

TODO: professional standard proof for suprema-ambient-set. □

Detailed Learning Proof.

TODO: detailed learning proof for suprema-ambient-set. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Ambient Existence of Supremum). *There are ordered sets $S \subseteq S'$ and a subset $A \subseteq S$ such that A has a supremum relative to S' but has no supremum relative to S .*

Professional Standard Proof.

TODO: professional standard proof for ambient-existence-supremum. □

Detailed Learning Proof.

TODO: detailed learning proof for ambient-existence-supremum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Lemma (Rational Gap Example for Suprema). *Let*

$$A = \{q \in \mathbb{Q} : q^2 < 2\}.$$

Then A is nonempty and bounded above in $\mathbb{Q}$, but A has no supremum relative to $\mathbb{Q}$.

Professional Standard Proof.

TODO: professional standard proof for rational-gap-suprema. □

Detailed Learning Proof.

TODO: detailed learning proof for rational-gap-suprema. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Capstone

Capstone Theorem: The Minimax Inequality

Theorem. If X and Y are nonempty sets and $\Phi : X \times Y \rightarrow \mathbb{R}$ is bounded, then

$$\sup_{x \in X} \inf_{y \in Y} \Phi(x, y) \leq \inf_{y \in Y} \sup_{x \in X} \Phi(x, y).$$

Remark (Abstract). This capstone proves that a player who must commit first cannot guarantee more than the opposing player can force by committing second. No topology, continuity, compactness, or convexity is involved. The proof uses only the order meaning of supremum and infimum, plus boundedness to ensure that all displayed extrema are real numbers.

Remark (Notation). For each $x \in X$, call

$$r(x) := \inf_{y \in Y} \Phi(x, y)$$

the floor of the row indexed by x . For each $y \in Y$, call

$$c(y) := \sup_{x \in X} \Phi(x, y)$$

the ceiling of the column indexed by y . The lower and upper values are

$$\underline{V} := \sup_{x \in X} r(x), \quad \overline{V} := \inf_{y \in Y} c(y).$$

The theorem is the comparison $\underline{V} \leq \overline{V}$.

Remark (Orientation). The theorem should be read against a simpler warning about coupled and independent indices. If $f, g : X \rightarrow \mathbb{R}$ are bounded, then

$$\sup_{x \in X} (f(x) + g(x)) \leq \sup_{x \in X} f(x) + \sup_{x \in X} g(x),$$

and the inequality can be strict when the two suprema occur at different points. By contrast, the independent grid identity

$$\sup_{(x,y) \in X \times Y} (f(x) + g(y)) = \sup_{x \in X} f(x) + \sup_{y \in Y} g(y)$$

is exactly the result proved in [Supremum of a Sum Set](#). The minimax inequality is another place where the order and independence of indices matter.

Remark (Lemma 1: existence of the displayed extrema). The row floors $r(x)$, column ceilings $c(y)$, lower value $\underline{V}$, and upper value $\overline{V}$ all exist as real numbers.

Proof task. Use the bounds $m \leq \Phi(x, y) \leq M$. Each row image $\{\Phi(x_0, y) : y \in Y\}$ and each column image $\{\Phi(x, y_0) : x \in X\}$ is nonempty and bounded because X and Y are nonempty. Therefore each row has an infimum and each column has a supremum. The family of row floors is bounded by m and M , and the family of column ceilings is also bounded by m and M . Apply the chapter's existence machinery for suprema and infima.

Remark (Lemma 2: the bridge). For every $x_0 \in X$ and $y_0 \in Y$,

$$r(x_0) = \inf_{y \in Y} \Phi(x_0, y) \leq \Phi(x_0, y_0) \leq \sup_{x \in X} \Phi(x, y_0) = c(y_0).$$

Proof task. The left inequality is the lower-bound property of the row infimum. The right inequality is the upper-bound property of the column supremum. The single point-height $\Phi(x_0, y_0)$ is the bridge between the row floor and the column ceiling.

Remark (Proposition 1: every column ceiling dominates the lower value). For every fixed $y_0 \in Y$,

$$\underline{V} \leq c(y_0).$$

Proof task. By Lemma 2, $r(x) \leq c(y_0)$ for every $x \in X$. Thus $c(y_0)$ is an upper bound for the set of row floors $\{r(x) : x \in X\}$. Since $\underline{V}$ is the supremum of that set, the upper-bound comparison property gives $\underline{V} \leq c(y_0)$.

Remark (Proof of the theorem). Proposition 1 holds for every $y_0 \in Y$. Hence $\underline{V}$ is a lower bound for the set of column ceilings $\{c(y) : y \in Y\}$. Since $\overline{V}$ is the infimum of that set,

$$\underline{V} \leq \inf_{y \in Y} c(y) = \overline{V}.$$

Expanding the definitions of r , c , $\underline{V}$, and $\overline{V}$ gives

$$\sup_{x \in X} \inf_{y \in Y} \Phi(x, y) \leq \inf_{y \in Y} \sup_{x \in X} \Phi(x, y).$$

Remark (Proposition 2: a saddle point forces equality). Call (x^*, y^*) a saddle point of Φ if

$$\Phi(x, y^*) \leq \Phi(x^*, y^*) \leq \Phi(x^*, y) \quad \text{for all } x \in X, y \in Y.$$

If a saddle point exists, then

$$\underline{V} = \overline{V} = \Phi(x^*, y^*).$$

Proof task. The left saddle inequality makes $\Phi(x^*, y^*)$ the supremum of the column at y^* . The right saddle inequality makes it the infimum of the row at x^* . Thus

$$\overline{V} \leq \Phi(x^*, y^*) \leq \underline{V}.$$

Combine this with the minimax inequality $\underline{V} \leq \overline{V}$.

Remark (Application: one-step pursuit on the circle). Let $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$, with wrap-around metric

$$d_{\mathbb{S}^1}(a, b) = \min_{k \in \mathbb{Z}} |a - b + k| \in \left[0, \frac{1}{2}\right].$$

Place a pursuer at 0 and an evader at $d_0 \in (0, \frac{1}{2}]$. In one step the pursuer chooses $u \in [-s_P, s_P]$, and the evader chooses $v \in [-s_E, s_E]$, where $s_P, s_E > 0$. Define

$$\Psi(v, u) := d_{\mathbb{S}^1}(u, d_0 + v).$$

The evader maximizes distance and the pursuer minimizes distance, so the one-step lower and upper values are

$$\sup_{v \in [-s_E, s_E]} \inf_{u \in [-s_P, s_P]} \Psi(v, u) \leq \inf_{u \in [-s_P, s_P]} \sup_{v \in [-s_E, s_E]} \Psi(v, u).$$

As a final task, determine whether this circle model has a saddle point, and compare the result with the corresponding one-step model on the line.

Remark (Limits of the theorem). Equality need not hold without extra structure. In particular, $\underline{V} = \bar{V}$ does not by itself imply that a saddle point exists. Existence of saddle points or equality of values usually requires additional hypotheses such as compactness, convexity in one variable, and concavity in the other. Those hypotheses belong to later minimax theorems, not to this bounds-only result.

Remark (Dependencies to state). • [Supremum](#)

- [Infimum](#)

Remark (Dependencies to prove). • [Supremum](#)

- [Infimum](#)
- [Upper-Bound Comparison with the Supremum](#)
- [Infimum Less Than Supremum](#)
- [Supremum of a Sum Set](#)

Remark (Dependency ceiling). This capstone may use only the order, supremum, infimum, and boundedness results routed at or before the Bounds chapter.

Remark (Dependencies).

- [Supremum](#)
- [Infimum](#)
- [Upper-Bound Comparison with the Supremum](#)
- [Infimum Less Than Supremum](#)
- [Supremum of a Sum Set](#)

Chapter 4

Functions



4.1 Algebra of Functions

Algebra of Functions — Quick Reference	
Statement	Role
Injective composition	Injectivity is preserved by composition
Surjective composition	Surjectivity is preserved by composition
Bijective composition	Bijections compose to bijections
Inverse bijection	Bijections have bijective inverses
Preimage algebra	Preimages preserve unions and intersections exactly

Algebra of Functions

Remark (Intervals on $\mathbb{R}$). Every open interval (a, b) consists entirely of cluster points; $[a, b]$ is closed and bounded. Natural domains for functions are typically intervals or unions of intervals.

Proposition

Proposition 4.1.1 (Composition of Injections Is Injective). *Let $f : A \rightarrow B$ and let $g : B \rightarrow C$. If f is injective and g is injective, then $g \circ f : A \rightarrow C$ is injective.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} & \left(\forall a_1, a_2 \in A (f(a_1) = f(a_2) \Rightarrow a_1 = a_2) \right) \wedge \left(\forall b_1, b_2 \in B (g(b_1) = g(b_2) \Rightarrow b_1 = b_2) \right) \\ & \Rightarrow \forall a_1, a_2 \in A (g(f(a_1)) = g(f(a_2)) \Rightarrow a_1 = a_2). \end{aligned}$$

Remark (Definition predicate reading).

$$\text{Injective}(f) \wedge \text{Injective}(g) \implies \text{Injective}(g \circ f).$$

Remark (Negated quantified statement).

$$\exists a_1, a_2 \in A (a_1 \neq a_2 \wedge g(f(a_1)) = g(f(a_2))).$$

Remark (Negation predicate reading).

$$\neg \text{Injective}(g \circ f).$$

Remark (Failure modes). The composite fails to be injective exactly when two distinct inputs of A are sent to the same output in C after applying both functions.

Remark (Failure mode decomposition).

$$\exists a_1, a_2 \in A (a_1 \neq a_2 \wedge g(f(a_1)) = g(f(a_2))).$$

Remark (Interpretation). Injectivity survives composition: equality after $g \circ f$ can be pulled back first through g and then through f .

Remark (Dependencies).

- [Composition](#)
- [Function](#)

Proposition

Proposition 4.1.2 (Composition of Surjections Is Surjective). $\text{Surjective}(f) \wedge \text{Surjective}(g) \implies \text{Surjective}(g \circ f)$.

[Go to proof.](#)

Remark (Standard quantified statement).

$$\text{Surjective}(f) \wedge \text{Surjective}(g) \implies \forall c \in C, \exists a \in A : g(f(a)) = c.$$

Remark (Definition predicate reading).

$$\text{Surjective}(g \circ f, C) \equiv \forall c \in C, \exists a \in A : (g \circ f)(a) = c.$$

Remark (Dependencies).

- [Composition](#)
- [Function](#)

Corollary

Corollary 4.1.3 (Composition of Bijections Is Bijective). *Let $f : A \rightarrow B$ and let $g : B \rightarrow C$. If f is bijective and g is bijective, then $g \circ f : A \rightarrow C$ is bijective.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} & \left[(\forall a_1, a_2 \in A (f(a_1) = f(a_2) \Rightarrow a_1 = a_2)) \wedge (\forall b \in B \exists a \in A (f(a) = b)) \right] \\ & \wedge \left[(\forall b_1, b_2 \in B (g(b_1) = g(b_2) \Rightarrow b_1 = b_2)) \wedge (\forall c \in C \exists b \in B (g(b) = c)) \right] \\ & \Rightarrow \left[(\forall a_1, a_2 \in A (g(f(a_1)) = g(f(a_2)) \Rightarrow a_1 = a_2)) \wedge (\forall c \in C \exists a \in A (g(f(a)) = c)) \right]. \end{aligned}$$

Remark (Definition predicate reading).

$$\text{Bijective}(f) \wedge \text{Bijective}(g) \implies \text{Bijective}(g \circ f).$$

Remark (Negated quantified statement).

$$\begin{aligned} & \exists a_1, a_2 \in A (a_1 \neq a_2 \wedge g(f(a_1)) = g(f(a_2))) \\ & \vee \exists c \in C \forall a \in A (g(f(a)) \neq c). \end{aligned}$$

Remark (Negation predicate reading).

$$\neg \text{Bijective}(g \circ f).$$

Remark (Failure modes). A composite bijection can fail only by losing injectivity or by missing some target value in C .

Remark (Failure mode decomposition).

$$\neg \text{Bijective}(g \circ f) \iff \neg \text{Injective}(g \circ f) \vee \neg \text{Surjective}(g \circ f).$$

Remark (Interpretation). A bijection is an injective and surjective function. Since both of those properties are preserved by composition, bijectivity is preserved too.

Remark (Dependencies).

- [Composition](#)
- [Function](#)

Proposition

Proposition 4.1.4 (Inverse of a Bijection Is a Bijection). $\text{Bijective}(f) \Rightarrow \text{Bijective}(f^{-1})$.

Go to proof.

Remark (Standard quantified statement).

$$\text{Bijective}(f) \Rightarrow \text{InverseFunction}(f, f^{-1}), f^{-1} \circ f = \text{id}_A, f \circ f^{-1} = \text{id}_B.$$

Remark (Definition predicate reading).

$$\text{InverseFunction}(f, f^{-1}) := f^{-1} \circ f = \text{id}_A \wedge f \circ f^{-1} = \text{id}_B.$$

Remark (Dependencies).

- [Composition](#)
- [Function](#)

Proposition

Proposition 4.1.5 (Preimage Distributes over Union and Intersection). *Let $f : A \rightarrow B$, $S, T \subseteq B$. Then:*

$$(i) \quad f^{-1}(S \cup T) = f^{-1}(S) \cup f^{-1}(T).$$

$$(ii) \quad f^{-1}(S \cap T) = f^{-1}(S) \cap f^{-1}(T).$$

$$(iii) \quad f^{-1}(S^c) = (f^{-1}(S))^c.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$(i) \quad x \in f^{-1}(S \cup T) \iff f(x) \in S \vee f(x) \in T \iff x \in f^{-1}(S) \cup f^{-1}(T).$$

$$(ii) \quad x \in f^{-1}(S \cap T) \iff f(x) \in S \wedge f(x) \in T \iff x \in f^{-1}(S) \cap f^{-1}(T).$$

Remark (Negated quantified statement).

$$x \notin f^{-1}(S) \iff f(x) \notin S \iff x \in f^{-1}(S^c).$$

Remark (Interpretation). Preimage is a complete lattice homomorphism: it preserves all set operations including complement. Contrast with image: $f(A \cap B) \subseteq f(A) \cap f(B)$ with strict containment possible when f is not injective. Preimage identities are load-bearing in topology (preimages of open sets are open) and measure theory (preimages of measurable sets are measurable).

Remark (Dependencies).

- [Function](#)

4.2 Real-Valued Functions

Real-Valued Functions — Quick Reference

Concept	Meaning
Pointwise sum	Algebraic addition of function values
Pointwise product	Product fg , distinct from composition
Pointwise quotient	Quotient where denominator is nonzero
Bounded function	A single bound controls all values on A
Supremum on a set	Least upper bound of $f(A)$
Maximum point	Point where the supremum is attained
Monotone function	Increasing or decreasing behavior on a set
Constant function	Same value at every input

Real-Valued Functions

Definition (Pointwise Sum of Functions)

Definition 4.2.1 (Pointwise Sum of Functions). Let $X \subseteq \mathbb{R}$ and let $f, g : X \rightarrow \mathbb{R}$. The pointwise sum $f + g : X \rightarrow \mathbb{R}$ is the function defined by

$$(f + g)(x) = f(x) + g(x) \quad (x \in X).$$

Remark (Standard quantified statement).

$$\forall x \in X ((f + g)(x) = f(x) + g(x)).$$

Remark (Interpretation). The sum is computed output by output; at each input, add the two function values at that same input.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Addition on \$\mathbb{R}\$](#)

Definition (Pointwise Difference of Functions)

Definition 4.2.2 (Pointwise Difference of Functions). Let $X \subseteq \mathbb{R}$ and let $f, g : X \rightarrow \mathbb{R}$. The pointwise difference $f - g : X \rightarrow \mathbb{R}$ is the function defined by

$$(f - g)(x) = f(x) - g(x) \quad (x \in X).$$

Remark (Standard quantified statement).

$$\forall x \in X ((f - g)(x) = f(x) - g(x)).$$

Remark (Interpretation). The difference subtracts the output of g from the output of f at each input separately.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Subtraction on \$\mathbb{R}\$](#)

Definition (Pointwise Product of Functions)

Definition 4.2.3 (Pointwise Product of Functions). Let $X \subseteq \mathbb{R}$ and let $f, g : X \rightarrow \mathbb{R}$. The pointwise product $fg : X \rightarrow \mathbb{R}$ is the function defined by

$$(fg)(x) = f(x)g(x) \quad (x \in X).$$

Remark (Standard quantified statement).

$$\forall x \in X \ ((fg)(x) = f(x)g(x)).$$

Remark (Interpretation). The product is not composition: it multiplies the two real outputs at the same input.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Multiplication on \$\mathbb{R}\$](#)

Definition (Pointwise Scalar Multiple of a Function)

Definition 4.2.4 (Pointwise Scalar Multiple of a Function). Let $X \subseteq \mathbb{R}$, let $f : X \rightarrow \mathbb{R}$, and let $c \in \mathbb{R}$. The pointwise scalar multiple $cf : X \rightarrow \mathbb{R}$ is the function defined by

$$(cf)(x) = cf(x) \quad (x \in X).$$

Remark (Standard quantified statement).

$$\forall x \in X \ ((cf)(x) = cf(x)).$$

Remark (Interpretation). Scalar multiplication stretches every output of the function by the same real scalar.

Remark (Dependencies).

- [Function](#)

- [Subset](#)
- [Multiplication on \$\mathbb{R}\$](#)

Definition (Pointwise Absolute Value of a Function)

Definition 4.2.5 (Pointwise Absolute Value of a Function). Let $X \subseteq \mathbb{R}$ and let $f : X \rightarrow \mathbb{R}$. The pointwise absolute value $|f| : X \rightarrow \mathbb{R}$ is the function defined by

$$|f|(x) = |f(x)| \quad (x \in X).$$

Remark (Standard quantified statement).

$$\forall x \in X \ (|f|(x) = |f(x)|).$$

Remark (Interpretation). The absolute value operation applies the ordinary real absolute value to each output.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Absolute Value on \$\mathbb{R}\$](#)

Definition (Pointwise Maximum of Two Functions)

Definition 4.2.6 (Pointwise Maximum of Two Functions). Let $X \subseteq \mathbb{R}$ and let $f, g : X \rightarrow \mathbb{R}$. The pointwise maximum $\max(f, g) : X \rightarrow \mathbb{R}$ is the function defined by

$$\max(f, g)(x) = \max\{f(x), g(x)\} \quad (x \in X).$$

Remark (Standard quantified statement).

$$\forall x \in X \ (\max(f, g)(x) = \max\{f(x), g(x)\}).$$

Remark (Interpretation). At each input, the new function keeps the larger of the two output values.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Maximum](#)

Definition (Pointwise Quotient of Functions)

Definition 4.2.7 (Pointwise Quotient of Functions). Let $X \subseteq \mathbb{R}$ and let $f, g : X \rightarrow \mathbb{R}$. Suppose $g(x) \neq 0$ for every $x \in X$. The pointwise quotient $f/g : X \rightarrow \mathbb{R}$ is the function defined by

$$(f/g)(x) = \frac{f(x)}{g(x)} \quad (x \in X).$$

Remark (Standard quantified statement).

$$(\forall x \in X (g(x) \neq 0)) \implies \forall x \in X ((f/g)(x) = f(x)/g(x)).$$

Remark (Interpretation). The quotient is defined pointwise only when the denominator function never vanishes on the domain.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Division on \$\mathbb{R}\$](#)

Definition (Linear Combination of Real-Valued Functions)

Definition 4.2.8 (Linear Combination of Real-Valued Functions). Let $X \subseteq \mathbb{R}$, let $f, g : X \rightarrow \mathbb{R}$, and let $\alpha, \beta \in \mathbb{R}$. The real linear combination $\alpha f + \beta g : X \rightarrow \mathbb{R}$ is the function defined by

$$(\alpha f + \beta g)(x) = \alpha f(x) + \beta g(x) \quad (x \in X).$$

Remark (Standard quantified statement).

$$\forall x \in X ((\alpha f + \beta g)(x) = \alpha f(x) + \beta g(x)).$$

Remark (Interpretation). A linear combination is built only from scalar multiples and sums. It is the shared algebraic pattern behind sum rules, constant-multiple rules, and finite linear-combination rules for functions.

Remark (Dependencies).

- [Pointwise Scalar Multiple of a Function](#)
- [Pointwise Sum of Functions](#)

Definition (Real-Linear Rule)

Definition 4.2.9 (Real-Linear Rule). Let $\mathcal{C}$ be a class of real-valued functions closed under real linear combinations. A rule T on $\mathcal{C}$ is real-linear if, whenever $f, g \in \mathcal{C}$ and $\alpha, \beta \in \mathbb{R}$,

$$T(\alpha f + \beta g) = \alpha T(f) + \beta T(g)$$

whenever both sides are defined.

Remark (Standard quantified statement).

$$f, g \in \mathcal{C} \implies T(\alpha f + \beta g) = \alpha T(f) + \beta T(g).$$

Remark (Interpretation). Linearity records exactly the behavior of preserving addition and scalar multiplication. Nonlinear operations such as products, quotients, absolute values, and order comparisons require their own dependency information.

Remark (Dependencies).

- [Function](#)
- [Linear Combination of Real-Valued Functions](#)

Definition (Bounded Above Function)

Definition 4.2.10 (Bounded Above Function). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. The function f is bounded above on A if there exists $M \in \mathbb{R}$ such that

$$f(x) \leq M \quad (x \in A).$$

Remark (Standard quantified statement).

$$\exists M \in \mathbb{R} \forall x \in A (f(x) \leq M).$$

Remark (Negated quantified statement).

$$\forall M \in \mathbb{R} \exists x \in A (f(x) > M).$$

Remark (Failure modes). No proposed real ceiling works: whenever a candidate upper bound is chosen, some input has an output larger than that candidate.

Remark (Failure mode decomposition).

$$\forall M \in \mathbb{R} \exists x \in A (f(x) > M).$$

Remark (Interpretation). Bounded above means all output values lie below one common real ceiling.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Order on \$\mathbb{R}\$](#)
- [Bounded Above](#)

Definition (Bounded Below Function)

Definition 4.2.11 (Bounded Below Function). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. The function f is bounded below on A if there exists $m \in \mathbb{R}$ such that

$$m \leq f(x) \quad (x \in A).$$

Remark (Standard quantified statement).

$$\exists m \in \mathbb{R} \forall x \in A (m \leq f(x)).$$

Remark (Negated quantified statement).

$$\forall m \in \mathbb{R} \exists x \in A (f(x) < m).$$

Remark (Failure modes). No proposed real floor works: whenever a candidate lower bound is chosen, some input has an output smaller than that candidate.

Remark (Failure mode decomposition).

$$\forall m \in \mathbb{R} \exists x \in A (f(x) < m).$$

Remark (Interpretation). Bounded below means all output values lie above one common real floor.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Order on \$\mathbb{R}\$](#)
- [Bounded Below](#)

Definition (Bounded Function)

Definition 4.2.12 (Bounded Function). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. The function f is bounded on A if there exists $B > 0$ such that

$$|f(x)| \leq B \quad (x \in A).$$

Remark (Standard quantified statement).

$$\exists B > 0 \forall x \in A (|f(x)| \leq B).$$

Remark (Definition predicate reading).

$$\text{BoundedOn}(f, A) := \exists B > 0 \forall x \in A (|f(x)| \leq B).$$

Remark (Negated quantified statement).

$$\forall B > 0 \exists x \in A (|f(x)| > B).$$

Remark (Negation predicate reading).

$$\neg \text{BoundedOn}(f, A) \iff \forall B > 0 \exists x \in A (|f(x)| > B).$$

Remark (Failure modes). Every symmetric interval around 0 fails to contain all output values of f on A .

Remark (Failure mode decomposition).

$$\forall B > 0 \exists x \in A (f(x) > B \vee f(x) < -B).$$

Remark (Interpretation). A bounded function has all of its values trapped inside some symmetric interval $[-B, B]$. Being bounded above alone is not enough; the function $f(x) = -x$ on $\mathbb{R}$ is bounded above by 0 but is not bounded.

Remark (Dependencies).

- [Bounded Above Function](#)
- [Bounded Below Function](#)

Definition (Supremum of a Function on a Set)

Definition 4.2.13 (Supremum of a Function on a Set). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. The supremum of f on A is

$$\sup_{x \in A} f(x) = \sup f(A),$$

provided the supremum of $f(A)$ exists.

Remark (Standard quantified statement).

$$s = \sup_{x \in A} f(x) \iff (\forall x \in A (f(x) \leq s)) \wedge (\forall u \in \mathbb{R} ((\forall x \in A (f(x) \leq u)) \Rightarrow s \leq u)).$$

Remark (Interpretation). The supremum is the least real ceiling for the output set $f(A)$; it need not be an output value of f .

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Image Set](#)
- [Supremum](#)

Definition (Infimum of a Function on a Set)

Definition 4.2.14 (Infimum of a Function on a Set). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. The infimum of f on A is

$$\inf_{x \in A} f(x) = \inf f(A),$$

provided the infimum of $f(A)$ exists.

Remark (Standard quantified statement).

$$t = \inf_{x \in A} f(x) \iff (\forall x \in A (t \leq f(x))) \wedge (\forall \ell \in \mathbb{R} ((\forall x \in A (\ell \leq f(x))) \Rightarrow \ell \leq t)).$$

Remark (Interpretation). The infimum is the greatest real floor for the output set $f(A)$; it need not be an output value of f .

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Image Set](#)
- [Infimum](#)

Definition (Maximum Point of a Function)

Definition 4.2.15 (Maximum Point of a Function). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. A point $x^* \in A$ is a maximum point of f on A if

$$f(x) \leq f(x^*) \quad (x \in A).$$

Remark (Standard quantified statement).

$$x^* \in A \wedge \forall x \in A (f(x) \leq f(x^*)).$$

Remark (Negated quantified statement).

$$x^* \notin A \vee \exists x \in A (f(x) > f(x^*)).$$

Remark (Failure modes). A proposed maximum point fails either by not belonging to the domain or by being beaten by another output value.

Remark (Failure mode decomposition).

$$x^* \notin A \vee \exists x \in A (f(x) > f(x^*)).$$

Remark (Interpretation). A maximum point is an input where the largest output value is attained.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Order on \$\mathbb{R}\$](#)
- [Maximum](#)

Definition (Minimum Point of a Function)

Definition 4.2.16 (Minimum Point of a Function). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. A point $x_* \in A$ is a minimum point of f on A if

$$f(x_*) \leq f(x) \quad (x \in A).$$

Remark (Standard quantified statement).

$$x_* \in A \wedge \forall x \in A (f(x_*) \leq f(x)).$$

Remark (Negated quantified statement).

$$x_* \notin A \vee \exists x \in A (f(x) < f(x_*)).$$

Remark (Failure modes). A proposed minimum point fails either by not belonging to the domain or by being undercut by another output value.

Remark (Failure mode decomposition).

$$x_* \notin A \vee \exists x \in A (f(x) < f(x_*)).$$

Remark (Interpretation). A minimum point is an input where the smallest output value is attained.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Order on \$\mathbb{R}\$](#)
- [Minimum](#)

Definition (Increasing Function)

Definition 4.2.17 (Increasing Function). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. The function f is increasing on A if

$$x_1 < x_2 \implies f(x_1) \leq f(x_2) \quad (x_1, x_2 \in A).$$

Remark (Standard quantified statement).

$$\forall x_1, x_2 \in A \ (x_1 < x_2 \Rightarrow f(x_1) \leq f(x_2)).$$

Remark (Negated quantified statement).

$$\exists x_1, x_2 \in A \ (x_1 < x_2 \wedge f(x_1) > f(x_2)).$$

Remark (Failure modes). The function fails to be increasing if some later input has a smaller output than an earlier input.

Remark (Failure mode decomposition).

$$\exists x_1, x_2 \in A \ (x_1 < x_2 \wedge f(x_1) > f(x_2)).$$

Remark (Interpretation). As the input moves to the right in A , the output is not allowed to go down.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Order on \$\mathbb{R}\$](#)

Definition (Strictly Increasing Function)

Definition 4.2.18 (Strictly Increasing Function). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. The function f is strictly increasing on A if

$$x_1 < x_2 \implies f(x_1) < f(x_2) \quad (x_1, x_2 \in A).$$

Remark (Standard quantified statement).

$$\forall x_1, x_2 \in A \ (x_1 < x_2 \Rightarrow f(x_1) < f(x_2)).$$

Remark (Negated quantified statement).

$$\exists x_1, x_2 \in A \ (x_1 < x_2 \wedge f(x_1) \geq f(x_2)).$$

Remark (Failure modes). Strict increase fails if some later input has an output that is no larger than the earlier output.

Remark (Failure mode decomposition).

$$\exists x_1, x_2 \in A \ (x_1 < x_2 \wedge f(x_1) \geq f(x_2)).$$

Remark (Interpretation). Strict increase requires every genuine move to the right in the domain to produce a genuine increase in output.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Strict Order on \$\mathbb{R}\$](#)

Definition (Decreasing Function)

Definition 4.2.19 (Decreasing Function). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. The function f is decreasing on A if

$$x_1 < x_2 \implies f(x_1) \geq f(x_2) \quad (x_1, x_2 \in A).$$

Remark (Standard quantified statement).

$$\forall x_1, x_2 \in A \ (x_1 < x_2 \implies f(x_1) \geq f(x_2)).$$

Remark (Negated quantified statement).

$$\exists x_1, x_2 \in A \ (x_1 < x_2 \wedge f(x_1) < f(x_2)).$$

Remark (Failure modes). The function fails to be decreasing if some later input has a larger output than an earlier input.

Remark (Failure mode decomposition).

$$\exists x_1, x_2 \in A \ (x_1 < x_2 \wedge f(x_1) < f(x_2)).$$

Remark (Interpretation). As the input moves to the right in A , the output is not allowed to go up.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Order on \$\mathbb{R}\$](#)

Definition (Strictly Decreasing Function)

Definition 4.2.20 (Strictly Decreasing Function). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. The function f is strictly decreasing on A if

$$x_1 < x_2 \implies f(x_1) > f(x_2) \quad (x_1, x_2 \in A).$$

Remark (Standard quantified statement).

$$\forall x_1, x_2 \in A \ (x_1 < x_2 \implies f(x_1) > f(x_2)).$$

Remark (Negated quantified statement).

$$\exists x_1, x_2 \in A \ (x_1 < x_2 \wedge f(x_1) \leq f(x_2)).$$

Remark (Failure modes). Strict decrease fails if some later input has an output that is no smaller than the earlier output.

Remark (Failure mode decomposition).

$$\exists x_1, x_2 \in A (x_1 < x_2 \wedge f(x_1) \leq f(x_2)).$$

Remark (Interpretation). Strict decrease requires every genuine move to the right in the domain to produce a genuine decrease in output.

Remark (Dependencies).

- [Function](#)
- [Subset](#)
- [Strict Order on \$\mathbb{R}\$](#)

Definition (Monotone Function)

Definition 4.2.21 (Monotone Function). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. The function f is monotone on A if f is increasing on A or decreasing on A .

Remark (Standard quantified statement).

$$(\forall x_1, x_2 \in A (x_1 < x_2 \Rightarrow f(x_1) \leq f(x_2))) \vee (\forall x_1, x_2 \in A (x_1 < x_2 \Rightarrow f(x_1) \geq f(x_2))).$$

Remark (Definition predicate reading).

$$\text{MonotoneFunction}(f, A) := (\forall x_1, x_2 \in A (x_1 < x_2 \Rightarrow f(x_1) \leq f(x_2))) \vee (\forall x_1, x_2 \in A (x_1 < x_2 \Rightarrow f(x_1) \geq f(x_2))).$$

Remark (Negated quantified statement).

$$(\exists x_1, x_2 \in A (x_1 < x_2 \wedge f(x_1) > f(x_2))) \wedge (\exists y_1, y_2 \in A (y_1 < y_2 \wedge f(y_1) < f(y_2))).$$

Remark (Negation predicate reading).

$$\neg \text{MonotoneFunction}(f, A)$$

Remark (Failure modes). A nonmonotone function has evidence of both kinds of reversal: one pair where the outputs go down as inputs increase, and another pair where the outputs go up as inputs increase.

Remark (Failure mode decomposition).

$$\begin{aligned} &\exists x_1, x_2 \in A (x_1 < x_2 \wedge f(x_1) > f(x_2)) \\ &\wedge \exists y_1, y_2 \in A (y_1 < y_2 \wedge f(y_1) < f(y_2)). \end{aligned}$$

Remark (Interpretation). A monotone function moves consistently in one direction: nondecreasing or nonincreasing.

Remark (Dependencies).

- [Increasing Function](#)
- [Decreasing Function](#)

Definition (Constant Function)

Definition 4.2.22 (Constant Function). Let $A \subseteq \mathbb{R}$ and let $f : A \rightarrow \mathbb{R}$. The function f is constant on A if

$$f(x_1) = f(x_2) \quad (x_1, x_2 \in A).$$

Remark (Standard quantified statement).

$$\forall x_1, x_2 \in A \ (f(x_1) = f(x_2)).$$

Remark (Negated quantified statement).

$$\exists x_1, x_2 \in A \ (f(x_1) \neq f(x_2)).$$

Remark (Failure modes). Constancy fails as soon as two inputs in the domain have different outputs.

Remark (Failure mode decomposition).

$$\exists x_1, x_2 \in A \ (f(x_1) \neq f(x_2)).$$

Remark (Interpretation). A constant function assigns the same real value to every input in its domain.

Remark (Dependencies).

- [Function](#)
- [Subset](#)

4.3 Subsets of $\mathbb{R}$

Subsets of $\mathbb{R}$ — Quick Reference

Concept	Meaning
Epsilon neighbourhood	Unpunctured condition $ x - c < r$
Deleted neighbourhood	Punctured condition $0 < x - c < \delta$
Cluster point	Every neighbourhood contains a distinct point of X
Adherent point	Every neighbourhood contains some point of X
Closure	Points adherent to X
Closed set	Set containing its closure or sequential limits
Bounded set	Set contained in a large interval
Heine–Borel	Closed and bounded iff sequentially compact in $\mathbb{R}$

Subsets of $\mathbb{R}$

Remark (Terminological note). Cluster point = limit point = accumulation point. Bartle and Lebl use *cluster point*; Rudin uses *limit point*; Tao uses both. Adherent point (Tao) is the weaker notion that includes isolated points.

Definition (ε -Neighbourhood)

Definition 4.3.1 (ε -Neighbourhood). Let $x \in \mathbb{R}$ and let $\varepsilon > 0$. The ε -neighbourhood of x is

$$V_\varepsilon(x) := (x - \varepsilon, x + \varepsilon) = \{y \in \mathbb{R} : |y - x| < \varepsilon\}.$$

Remark (Standard quantified statement).

$$y \in V_\varepsilon(x) \iff |y - x| < \varepsilon.$$

Remark (Interpretation). The ε -neighbourhood is the open interval of radius ε centered at x .

Remark (Dependencies).

- [Open Ball on the Real Line](#)

Definition (Deleted ε -Neighbourhood)

Definition 4.3.2 (Deleted ε -Neighbourhood). Let $x \in \mathbb{R}$ and let $\varepsilon > 0$. The **deleted ε -neighbourhood** of x is

$$V_\varepsilon^*(x) := V_\varepsilon(x) \setminus \{x\} = \{y \in \mathbb{R} : 0 < |y - x| < \varepsilon\}.$$

Remark (Standard quantified statement).

$$y \in V_\varepsilon^*(x) \iff 0 < |y - x| < \varepsilon.$$

Remark (Interpretation). The deleted neighbourhood removes the base point. It is the punctured input region used in limit definitions.

Remark (Dependencies).

- [ε-Neighbourhood](#)

Definition (Cluster Point)

Definition 4.3.3 (Cluster Point). Let $X \subseteq \mathbb{R}$ and $x \in \mathbb{R}$. The point x is a **cluster point** of X if

$$\forall \varepsilon > 0, \exists y \in X \setminus \{x\} : |y - x| < \varepsilon.$$

Remark (Standard quantified statement).

$$x \text{ is a cluster point of } X \iff \forall \varepsilon > 0, \exists y \in X \setminus \{x\} : |y - x| < \varepsilon.$$

Remark (Definition predicate reading).

$$x \text{ is a cluster point of } X \equiv \forall \varepsilon > 0, V_\varepsilon^*(x) \cap X \neq \emptyset.$$

Remark (Negated quantified statement).

$$\neg x \text{ is a cluster point of } X \iff \exists \varepsilon > 0, \forall y \in X \setminus \{x\} : \neg |y - x| < \varepsilon.$$

Remark (Negation predicate reading).

$$\neg x \text{ is a cluster point of } X \equiv \exists \varepsilon > 0 : V_\varepsilon^*(x) \cap X = \emptyset.$$

Remark (Interpretation). A cluster point is approached by X via distinct witnesses arbitrarily close. A cluster point need not belong to X : 0 is a cluster point of $(0, 1)$ but $0 \notin (0, 1)$. The condition $\lim_{x \rightarrow c} f(x) = L$ requires $\text{ClusterPoint}(c, \text{dom}(f))$; without it the punctured ball condition is vacuous.

Remark (Dependencies).

- Deleted ε -Neighbourhood

Theorem

Theorem 4.3.4 (Sequential Characterization of Cluster Points). *c is a cluster point of A if and only if there exists $(a_n) \subseteq A \setminus \{c\}$ with $a_n \rightarrow c$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$c \text{ is a cluster point of } A \iff \exists (a_n) \subseteq A \setminus \{c\} : a_n \rightarrow c.$$

Remark (Interpretation). The forward direction: use $\varepsilon = 1/n$ to build the witness sequence. The reverse direction is immediate since the sequence witnesses the ball condition for every ε . This bridges the ε - δ definition with the sequential language.

Remark (Dependencies).

- Cluster Point

Definition (Adherent Point)

Definition 4.3.5 (Adherent Point). Let $X \subseteq \mathbb{R}$ and let $x \in \mathbb{R}$. The point x is an **adherent point** of X if

$$\forall \varepsilon > 0 \exists y \in X (|y - x| < \varepsilon).$$

Remark (Standard quantified statement).

$$\forall \varepsilon > 0 (V_\varepsilon(x) \cap X \neq \emptyset).$$

Remark (Definition predicate reading).

$$\text{AdherentPoint}(x, X) := \forall \varepsilon > 0 \exists y \in X (|y - x| < \varepsilon).$$

Remark (Negated quantified statement).

$$\exists \varepsilon > 0 \forall y \in X (|y - x| \geq \varepsilon).$$

Remark (Interpretation). An adherent point can be witnessed by points of X arbitrarily close to x , including possibly x itself.

Remark (Dependencies).

- [ε-Neighbourhood](#)

Definition (Isolated Point)

Definition 4.3.6 (Isolated Point). Let $X \subseteq \mathbb{R}$ and let $x \in X$. The point x is an **isolated point** of X if

$$\exists \varepsilon > 0 (V_\varepsilon(x) \cap X = \{x\}).$$

Remark (Standard quantified statement).

$$\exists \varepsilon > 0 \forall y \in X (|y - x| < \varepsilon \Rightarrow y = x).$$

Remark (Definition predicate reading).

$$\text{IsolatedPoint}(x, X) := x \in X \wedge \exists \varepsilon > 0 (V_\varepsilon(x) \cap X = \{x\}).$$

Remark (Negated quantified statement).

$$\forall \varepsilon > 0 \exists y \in X \setminus \{x\} (|y - x| < \varepsilon).$$

Remark (Interpretation). An isolated point belongs to the set but has a neighbourhood containing no other points of the set.

Remark (Dependencies).

- [Adherent Point](#)
- [Cluster Point](#)

Definition (Interior Point)

Definition 4.3.7 (Interior Point). Let $X \subseteq \mathbb{R}$ and let $x \in \mathbb{R}$. The point x is an **interior point** of X if

$$\exists \varepsilon > 0 (V_\varepsilon(x) \subseteq X).$$

Remark (Standard quantified statement).

$$\exists \varepsilon > 0 \forall y \in \mathbb{R} (|y - x| < \varepsilon \Rightarrow y \in X).$$

Remark (Definition predicate reading).

$$\text{InteriorPoint}(x, X) := \exists \varepsilon > 0 \ (V_\varepsilon(x) \subseteq X).$$

Remark (Interpretation). An interior point has a full open neighbourhood contained in the set.

Remark (Dependencies).

- ε -Neighbourhood

Definition (Boundary Point)

Definition 4.3.8 (Boundary Point). Let $X \subseteq \mathbb{R}$ and let $x \in \mathbb{R}$. The point x is a **boundary point** of X if

$$\forall \varepsilon > 0 \ (V_\varepsilon(x) \cap X \neq \emptyset) \wedge (V_\varepsilon(x) \cap X^c \neq \emptyset).$$

Remark (Standard quantified statement).

$$\forall \varepsilon > 0 \ (V_\varepsilon(x) \cap X \neq \emptyset) \wedge (V_\varepsilon(x) \cap X^c \neq \emptyset).$$

Remark (Definition predicate reading).

$$\text{BoundaryPoint}(x, X) := \forall \varepsilon > 0 \ (V_\varepsilon(x) \cap X \neq \emptyset) \wedge (V_\varepsilon(x) \cap X^c \neq \emptyset).$$

Remark (Interpretation). A boundary point sees both the set and its complement in every neighbourhood.

Remark (Dependencies).

- ε -Neighbourhood

Definition (Interior)

Definition 4.3.9 (Interior). Let $X \subseteq \mathbb{R}$. The **interior** of X is

$$X^\circ := \{x \in \mathbb{R} : \exists \varepsilon > 0 \ V_\varepsilon(x) \subseteq X\}.$$

Remark (Standard quantified statement).

$$x \in X^\circ \iff \exists \varepsilon > 0 \ V_\varepsilon(x) \subseteq X.$$

Remark (Interpretation). The interior collects exactly the points around which the set contains a whole neighbourhood.

Remark (Dependencies).

- Interior Point

Definition (Boundary)

Definition 4.3.10 (Boundary). Let $X \subseteq \mathbb{R}$. The **boundary** of X is

$$\partial X := \{x \in \mathbb{R} : \forall \varepsilon > 0 (V_\varepsilon(x) \cap X \neq \emptyset \wedge V_\varepsilon(x) \cap X^c \neq \emptyset)\}.$$

Remark (Standard quantified statement).

$$x \in \partial X \iff \forall \varepsilon > 0 (V_\varepsilon(x) \cap X \neq \emptyset \wedge V_\varepsilon(x) \cap X^c \neq \emptyset).$$

Remark (Interpretation). The boundary records the points where every neighbourhood touches both inside and outside of the set.

Remark (Dependencies).

- [Boundary Point](#)

Definition (Closure)

Definition 4.3.11 (Closure).

$$\overline{X} := \{x \in \mathbb{R} : \forall \varepsilon > 0 V_\varepsilon(x) \cap X \neq \emptyset\}.$$

Remark (Standard quantified statement).

$$x \in \overline{X} \iff x \text{ is an adherent point of } X \iff \forall \varepsilon > 0 : V_\varepsilon(x) \cap X \neq \emptyset.$$

Remark (Interpretation). The closure adds all adherent points of X , so it fills in the limiting points that can be reached by points of X .

Remark (Canonical examples). $\overline{(a, b)} = [a, b]$; $\overline{\mathbb{Q}} = \mathbb{R}$; $\overline{\mathbb{Z}} = \mathbb{Z}$; $\overline{\{1/n\}} = \{1/n\} \cup \{0\}$.

Remark (Dependencies).

- [Adherent Point](#)

Lemma

Lemma 4.3.12 (Elementary Properties of Closures). Let $X, Y \subseteq \mathbb{R}$. Then: (i) $X \subseteq \overline{X}$; (ii) $\overline{X \cup Y} = \overline{X} \cup \overline{Y}$; (iii) $\overline{X \cap Y} \subseteq \overline{X} \cap \overline{Y}$; (iv) $X \subseteq Y \Rightarrow \overline{X} \subseteq \overline{Y}$.

Go to proof.

Remark (Standard quantified statement).

$$\overline{X \cup Y} = \overline{X} \cup \overline{Y}; \quad \overline{X \cap Y} \subseteq \overline{X} \cap \overline{Y}.$$

Remark (Failure of equality in (iii)). Take $X = (0, 1)$, $Y = (1, 2)$: $\overline{X \cap Y} = \emptyset$ but $\overline{X} \cap \overline{Y} = \{1\}$.

Remark (Interpretation). Closure preserves inclusions and finite unions, but intersection can lose limit points that are approached separately by the two sets.

Remark (Dependencies).

- [Closure](#)

Definition (Closed Subset of $\mathbb{R}$)

Definition 4.3.13 (Closed Subset of $\mathbb{R}$). E is closed $\iff \overline{E} = E$.

Remark (Standard quantified statement).

$$E \text{ is closed} \iff \forall x \in \mathbb{R} : x \text{ is an adherent point of } E \Rightarrow x \in E.$$

Remark (Definition predicate reading).

$$E \text{ is closed} :\equiv \forall x : x \text{ is an adherent point of } E \Rightarrow x \in E.$$

Remark (Negated quantified statement).

$$\neg E \text{ is closed} \iff \exists x \notin E : x \text{ is an adherent point of } E.$$

Remark (Negation predicate reading).

$$\neg E \text{ is closed} :\equiv \exists x \notin E : \forall \varepsilon > 0 : V_\varepsilon(x) \cap E \neq \emptyset.$$

Remark (Interpretation). E is closed iff it contains its own boundary. Examples: $[a, b]$ closed; (a, b) not closed; $\mathbb{R}$ and $\emptyset$ both closed; $\mathbb{Q}$ not closed.

Remark (Dependencies).

- [Closure](#)

Corollary

Corollary 4.3.14 (Closed iff Sequentially Closed). X is closed $\iff \forall (a_n) \subseteq X, a_n \rightarrow x \Rightarrow x \in X$.

[Go to proof.](#)

Remark (Standard quantified statement).

$$X \text{ is closed} \iff \forall (a_n) \subseteq X, a_n \rightarrow x \Rightarrow x \in X.$$

Remark (Interpretation). A closed set cannot be escaped by convergent sequences. Every adherent point has a witness sequence; closedness forces the limit into the set.

Remark (Dependencies).

- [Closed Subset of \$\mathbb{R}\$](#)
- [Sequential Characterization of Cluster Points](#)

Corollary

Corollary 4.3.15 (Every Point of an Interval Is a Cluster Point). *If I is an interval, then every $x \in I$ is a cluster point of I .*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall x \in I \forall \varepsilon > 0 \exists y \in I \setminus \{x\} (|y - x| < \varepsilon).$$

Remark (Interpretation). This justifies intervals as natural domains for function limits. $\lim_{x \rightarrow c} f(x) = L$ requires $\text{ClusterPoint}(c, \text{dom}(f))$; on any interval this is free.

Remark (Dependencies).

- [Cluster Point](#)

Definition (Bounded Subset of $\mathbb{R}$)

Definition 4.3.16 (Bounded Subset of $\mathbb{R}$). X is bounded $\iff \exists M > 0, \forall x \in X : |x| \leq M$.

Remark (Standard quantified statement).

$$\exists M > 0 \forall x \in X (|x| \leq M).$$

Remark (Negated quantified statement).

$$\forall M > 0 \exists x \in X (|x| > M).$$

Remark (Interpretation). A bounded subset of $\mathbb{R}$ fits inside some symmetric interval $[-M, M]$.

Remark (Dependencies).

- [Subset](#)
- [Absolute Value on \$\mathbb{R}\$](#)
- [Order on \$\mathbb{R}\$](#)

Theorem

Theorem 4.3.17 (Heine–Borel Theorem for $\mathbb{R}$). X is closed $\wedge X$ is bounded $\iff$ every $(a_n) \subseteq X$ has a subsequence converging to a limit in X .

Go to proof.

Remark (Standard quantified statement).

$$X \text{ is closed} \wedge X \text{ is bounded} \iff \forall (a_n) \subseteq X, \exists (a_{n_k}), \exists L \in X : a_{n_k} \rightarrow L.$$

Remark (Definition predicate reading).

$$\text{SequentiallyCompact}(X) :\equiv X \text{ is closed} \wedge X \text{ is bounded}.$$

Remark (Negated quantified statement).

$$\neg(X \text{ is closed} \wedge X \text{ is bounded}) \iff \exists (a_n) \subseteq X : \text{no subsequence converges to a limit in } X. \blacksquare$$

Remark (Failure modes). The compactness conclusion fails when either closedness or boundedness fails.

Remark (Failure mode decomposition). Either X is bounded fails — a sequence escapes to $\pm\infty$ — or X is closed fails — a convergent subsequence exists but its limit lies outside X .

Remark (Interpretation). Closed prevents limits from escaping through boundary gaps; bounded prevents escape to infinity. Both are required. Forward: bounded gives Bolzano–Weierstrass; closed keeps the limit in X . Reverse: negate either hypothesis to construct a sequence with no convergent subsequence in X .

Remark (Dependencies).

- Closed Subset of $\mathbb{R}$
- Bounded Subset of $\mathbb{R}$

Definition (True Near a Point)

Definition 4.3.18 (True Near a Point). Property $P(f(x))$ is **true near** x_0 if $\exists \delta > 0, \forall x : 0 < |x - x_0| < \delta \Rightarrow P(f(x))$.

Remark (Standard quantified statement).

$$P(f(x)) \text{ true near } x_0 \iff \exists \delta > 0, \forall x : 0 < |x - x_0| < \delta \Rightarrow P(f(x)).$$

Remark (Interpretation). $\lim_{x \rightarrow x_0} f(x) = L$ states: for every $\varepsilon > 0$, the property $|f(x) - L| < \varepsilon$ is true near x_0 .

Remark (Dependencies).

- ε -Neighbourhood

Proofs

Remark (Return). [Return to Theorem](#)

Proposition (Composition of Injections Is Injective). *Let $f : A \rightarrow B$ and let $g : B \rightarrow C$. If f is injective and g is injective, then $g \circ f : A \rightarrow C$ is injective.*

Professional Standard Proof.

TODO: professional standard proof for composition-injective. □

Detailed Learning Proof.

TODO: detailed learning proof for composition-injective. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Composition of Surjections Is Surjective). $\text{Surjective}(f) \wedge \text{Surjective}(g) \Rightarrow \text{Surjective}(g \circ f)$.

Professional Standard Proof.

TODO: professional standard proof for composition-surjective. □

Detailed Learning Proof.

TODO: detailed learning proof for composition-surjective. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Corollary (Composition of Bijections Is Bijective). *Let $f : A \rightarrow B$ and let $g : B \rightarrow C$. If f is bijective and g is bijective, then $g \circ f : A \rightarrow C$ is bijective.*

Professional Standard Proof.

TODO: professional standard proof for composition-bijective. □

Detailed Learning Proof.

TODO: detailed learning proof for composition-bijective. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Inverse of a Bijection Is a Bijection). $\text{Bijective}(f) \Rightarrow \text{Bijective}(f^{-1})$.

Professional Standard Proof.

TODO: professional standard proof for inverse-bijection. □

Detailed Learning Proof.

TODO: detailed learning proof for inverse-bijection. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Proposition (Preimage Distributes over Union and Intersection). *Let $f : A \rightarrow B$, $S, T \subseteq B$. Then:*

$$(i) \quad f^{-1}(S \cup T) = f^{-1}(S) \cup f^{-1}(T).$$

$$(ii) \quad f^{-1}(S \cap T) = f^{-1}(S) \cap f^{-1}(T).$$

$$(iii) \quad f^{-1}(S^c) = (f^{-1}(S))^c.$$

Professional Standard Proof.

TODO: professional standard proof for preimage-union-intersection. □

Detailed Learning Proof.

TODO: detailed learning proof for preimage-union-intersection. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Sequential Characterization of Cluster Points). *c is a cluster point of A if and only if there exists $(a_n) \subseteq A \setminus \{c\}$ with $a_n \rightarrow c$.*

Professional Standard Proof.

TODO: professional standard proof for cluster-point-sequential. □

Detailed Learning Proof.

TODO: detailed learning proof for cluster-point-sequential. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Lemma (Elementary Properties of Closures). *Let $X, Y \subseteq \mathbb{R}$. Then: (i) $X \subseteq \overline{X}$; (ii) $\overline{X \cup Y} = \overline{X} \cup \overline{Y}$; (iii) $\overline{X \cap Y} \subseteq \overline{X} \cap \overline{Y}$; (iv) $X \subseteq Y \Rightarrow \overline{X} \subseteq \overline{Y}$.*

Professional Standard Proof.

TODO: professional standard proof for closure-elementary. □

Detailed Learning Proof.

TODO: detailed learning proof for closure-elementary. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Corollary (Closed iff Sequentially Closed). X is closed $\iff \forall(a_n) \subseteq X, a_n \rightarrow x \Rightarrow x \in X$.

Professional Standard Proof.

TODO: professional standard proof for closed-iff-seq-limits. □

Detailed Learning Proof.

TODO: detailed learning proof for closed-iff-seq-limits. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Corollary (Every Point of an Interval Is a Cluster Point). *If I is an interval, then every $x \in I$ is a cluster point of I .*

Professional Standard Proof.

TODO: professional standard proof for interval-all-limit-points. □

Detailed Learning Proof.

TODO: detailed learning proof for interval-all-limit-points. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Heine–Borel Theorem for $\mathbb{R}$). X is closed $\wedge$ X is bounded $\iff$ every $(a_n) \subseteq X$ has a subsequence converging to a limit in X .

Professional Standard Proof.

TODO: professional standard proof for heine-borel-subsets-real-line. □

Detailed Learning Proof.

TODO: detailed learning proof for heine-borel-subsets-real-line. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Chapter 5

Discrete Calculus



5.1 Motivation

Motivation

Classical calculus studies functions defined on continuous domains, typically functions $f : \mathbb{R} \rightarrow \mathbb{R}$. Its central tools are the derivative and the integral:

- The derivative measures *instantaneous change*.
- The integral accumulates *total change*.

However, many important mathematical objects are inherently *discrete*. Examples include

- sequences a_n ,
- recurrence relations,
- sums of integers or combinatorial quantities,
- discrete probability distributions,
- algorithms and computational processes.

In these contexts the domain is typically the natural numbers

$$f : \mathbb{Z} \rightarrow \mathbb{R}$$

rather than the real line.

To study change in such settings, calculus is replaced by its discrete counterpart.

Discrete Analogs of Calculus

The central idea of *discrete calculus* is that the roles of derivative and integral are played by two simple operations.

Continuous vs. Discrete Calculus	
Continuous Calculus	Discrete Calculus
Derivative $\frac{d}{dx}f(x)$	Forward difference $\Delta f(n) = f(n+1) - f(n)$
Integral $\int f(x) dx$	Summation $\sum f(n)$

The forward difference operator measures change between consecutive values of a function.

$$\Delta f(n) = f(n+1) - f(n).$$

This quantity plays the same role for sequences that the derivative plays for functions of a real variable.

Likewise, summation serves as the discrete analogue of integration.

$$\sum_{k=a}^b f(k)$$

accumulates the values of a function across a finite discrete interval.

Telescoping Phenomena

One of the most important identities in discrete mathematics arises when differences are summed.

$$\sum_{k=a}^{b-1} (f(k+1) - f(k)) = f(b) - f(a).$$

The intermediate terms cancel, leaving only the endpoints. Such sums are called *telescoping sums*.

This identity is the discrete counterpart of the *Fundamental Theorem of Calculus*.

Why Discrete Calculus Matters

Discrete calculus appears throughout mathematics:

- in combinatorics and enumerative formulas,
- in recurrence relations and generating functions,
- in probability theory and stochastic processes,

- in numerical analysis and finite difference methods,
- in computer science and algorithm analysis.

For students of analysis, discrete calculus is particularly valuable because it reveals the structural skeleton behind many identities involving sums and sequences.

Many techniques used to estimate limits of sequences or series ultimately rely on discrete analogues of differential identities.

A Structural View


Continuous calculus studies how functions change along the real line.

Discrete calculus studies how functions change along the integers.

Despite the difference in setting, the two theories mirror each other remarkably closely. Understanding this parallel provides powerful intuition and useful techniques for manipulating sums, sequences, and recurrence relations.

In the sections that follow we develop the basic tools of discrete calculus, beginning with the forward difference operator and the algebra of finite differences.

Structural Correspondence Between Continuous and Discrete Calculus



5.2 Foundations

5.2.1 Foundations

Definition (Discrete Function)

A *discrete function* is a function

$$f : \mathbb{Z} \rightarrow \mathbb{R}$$

(or more generally $f : \mathbb{N} \rightarrow \mathbb{R}$). It assigns a real value to each integer input, producing a sequence $f(0), f(1), f(2), \dots$

Remark (Fully quantified form). $\forall n \in \mathbb{Z}, f(n) \in \mathbb{R}$. The domain is the integers (or natural numbers); the codomain is the real line.

Remark (Discrete vs. continuous). In discrete calculus the independent variable moves in *integer steps* rather than continuously. Change is therefore measured between successive integer inputs rather than through infinitesimal limits. The fundamental operator of discrete calculus is the *difference operator*

$$\Delta f(n) = f(n+1) - f(n),$$

which plays a role analogous to the derivative in continuous calculus.

Remark (Sequences as discrete functions). A discrete function is simply another way of viewing a *sequence*. A sequence $a_0, a_1, a_2, \dots$ is identified with

$$f : \mathbb{N} \rightarrow \mathbb{R}, \quad f(n) = a_n.$$

Several equivalent notations appear in the literature: $f(n)$, a_n , $(a_n)_{n \in \mathbb{N}}$, $\{a_n\}_{n=0}^{\infty}$. All describe the same object. The functional notation $f(n)$ is preferred in discrete calculus because the difference operator acts on it much as a derivative acts on functions in continuous calculus.

5.2.2 Summation Notation

Definition (Summation)

Let f be a function defined on the integers. For integers $a \leq b$, the expression

$$\sum_{k=a}^b f(k)$$

denotes the sum $f(a) + f(a+1) + f(a+2) + \dots + f(b)$. The symbol $\sum$ (Greek capital sigma) represents summation; k is the *index of summation*; a and b are the *lower* and *upper limits*; $f(k)$ is the *summand*.

Remark (Standard quantified statement). $\sum_{k=a}^b f(k) := f(a) + f(a+1) + \dots + f(b)$, where $a, b \in \mathbb{Z}$, $a \leq b$, and $f : \mathbb{Z} \rightarrow \mathbb{R}$.

Remark (Interpretation). Sigma notation compresses a finite list of additions into one indexed symbol.

Remark (Dummy variable). The index of summation is a *dummy variable*: $\sum_{k=a}^b f(k) = \sum_{i=a}^b f(i)$. Its name carries no information about the value of the sum.

Basic Properties of Summation

Remark (Basic properties of summation). For functions f and g and constants c ,

$$\begin{aligned}\sum_{k=m}^n (f(k) + g(k)) &= \sum_{k=m}^n f(k) + \sum_{k=m}^n g(k), \\ \sum_{k=m}^n c f(k) &= c \sum_{k=m}^n f(k).\end{aligned}$$

These express the *linearity* of summation.

Splitting Sums

Remark (Splitting sums). A sum over an interval may be divided into smaller pieces:

$$\sum_{k=a}^c f(k) = \sum_{k=a}^b f(k) + \sum_{k=b+1}^c f(k) \quad \text{whenever } a \leq b < c.$$

Changing the Index

Remark (Changing the index). It is often useful to shift the summation index. For example,

$$\sum_{k=0}^n f(k) = \sum_{j=1}^{n+1} f(j-1).$$

Such substitutions frequently simplify expressions or align sums with known formulas.

Empty Sums

Remark (Empty sums). If the upper bound is smaller than the lower bound, the sum contains no terms. By convention,

$$\sum_{k=a}^b f(k) = 0 \quad \text{when } a > b.$$

This convention simplifies many algebraic identities.

Nested Summations

Remark (Nested summations). Summations may be nested:

$$\sum_{i=1}^n \sum_{j=1}^m a_{ij} = \sum_{i=1}^n (a_{i1} + a_{i2} + \cdots + a_{im}).$$

The inner sum is evaluated first. When the limits are independent, the order of summation may be exchanged:

$$\sum_{i=1}^n \sum_{j=1}^m a_{ij} = \sum_{j=1}^m \sum_{i=1}^n a_{ij}.$$

Remark (Analogy with integration). Nested sums behave like iterated integrals: exchanging summation order corresponds to Fubini's theorem in the discrete setting.

Common Examples

Remark (Common examples).

$$\sum_{k=1}^n 1 = n, \quad \sum_{k=1}^n k = \frac{n(n+1)}{2}, \quad \sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}.$$

These formulas appear throughout discrete mathematics and analysis.

5.2.3 The Binomial Theorem

Theorem 5.2.1 (Binomial Theorem). *For any integer $n \geq 0$ and real numbers x, y ,*

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k,$$

where the binomial coefficients are

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}.$$

Remark (Standard quantified statement). $\forall n \in \mathbb{N}, \forall x, y \in \mathbb{R} : (x + y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k$.

Remark (Interpretation). The theorem expands a power of a sum by counting how many terms choose each possible number of y -factors.

Remark (Combinatorial meaning). The binomial coefficients count the number of ways to choose k factors of y from the n factors in the product $(x + y)(x + y) \cdots (x + y)$. Thus $\binom{n}{k}$ is the number of ways to select k objects from a set of n , and the theorem describes how a power of a sum expands into a sum of products.

Basic Properties of Binomial Coefficients

Remark (Basic properties of binomial coefficients).

$$\binom{n}{0} = 1, \quad \binom{n}{n} = 1, \quad \binom{n}{k} = \binom{n}{n-k}.$$

They also satisfy the *Pascal recursion*:

$$\binom{n+1}{k} = \binom{n}{k} + \binom{n}{k-1}.$$

This recursion produces Pascal's triangle, in which each entry is the sum of the two entries directly above it.

$$\begin{array}{ccccccc} & & & & \binom{0}{0} & & \\ & & & & & & \\ & & \binom{1}{0} & & \binom{1}{1} & & \\ & & & & & & \\ & \binom{2}{0} & & \binom{2}{1} & & \binom{2}{2} & \\ & & & & & & \\ & \binom{3}{0} & & \binom{3}{1} & & \binom{3}{2} & & \binom{3}{3} \\ & & & & & & \\ & \binom{4}{0} & & \binom{4}{1} & & \binom{4}{2} & & \binom{4}{3} & & \binom{4}{4} \\ & & & & & & \\ & \binom{5}{0} & & \binom{5}{1} & & \binom{5}{2} & & \binom{5}{3} & & \binom{5}{4} & & \binom{5}{5} \end{array}$$

Remark (Pascal's triangle). Each entry $\binom{n}{k}$ is obtained from the sum of the two entries above it: $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$. These coefficients appear throughout discrete calculus, especially in formulas for higher finite differences and in Newton's forward difference expansion.

Connection with Discrete Calculus

Remark (Connection with discrete calculus). The binomial theorem plays a central role in discrete calculus because difference operators naturally produce binomial coefficients.

Proposition 5.2.2 (Higher difference formula). *For any function f ,*

$$\Delta^n f(x) = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} f(x+k).$$

Go to proof.

Remark (Standard quantified statement). $\forall n \in \mathbb{N}, \forall f : \mathbb{Z} \rightarrow \mathbb{R}, \forall x \in \mathbb{Z} : \Delta^n f(x) = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} f(x+k).$

Remark (Interpretation). The n -th forward difference is a signed binomial combination of $n + 1$ successive function values.

Remark (Proof idea). Applying the difference operator $\Delta = E - I$ (where E is the shift operator) repeatedly gives $\Delta^n = (E - I)^n$. Expanding by the binomial theorem yields the coefficients $(-1)^{n-k} \binom{n}{k}$, and $E^k f(x) = f(x + k)$. This derivation is carried out in full in the Difference Operators section.

Remark (Dependencies).

- [Binomial Theorem](#).
- [Forward Difference Operator](#).
- [Higher Differences](#).

Explicit Examples

Remark (Explicit examples). Applying the difference operator repeatedly to a function f yields

$$\begin{aligned}\Delta f(x) &= f(x + 1) - f(x), \\ \Delta^2 f(x) &= f(x + 2) - 2f(x + 1) + f(x), \\ \Delta^3 f(x) &= f(x + 3) - 3f(x + 2) + 3f(x + 1) - f(x).\end{aligned}$$

The coefficients 1, 2, 1 and 1, 3, 3, 1 are precisely the rows of Pascal's triangle, confirming that binomial coefficients determine the pattern of coefficients in higher-order finite differences.

5.3 Algebra of Summations

Summation Algebra — Quick Reference

Linearity	$\sum (af + bg) = a \sum f + b \sum g$
Splitting	$\sum_a^c = \sum_a^b + \sum_{b+1}^c$
Index shift	$\sum_{k=a}^b f(k) = \sum_{k=a+1}^{b+1} f(k - 1)$
Constant sum	$\sum_{k=a}^b c = (b - a + 1)c$
Empty sum	$\sum_{k=a}^b f(k) = 0$ when $a > b$
Nested sums	$\sum_i \sum_j a_{ij} = \sum_j \sum_i a_{ij}$ (independent limits)

Remark (Orientation). The difference operator and telescoping identity provide the background for the basic algebraic rules used to manipulate summations. These identities allow sums to be combined, split, and reindexed while preserving their value.

Linearity

Remark (Linearity). Summation is a linear operator. For functions f and g and constants a and b ,

$$\sum_{k=m}^n (a f(k) + b g(k)) = a \sum_{k=m}^n f(k) + b \sum_{k=m}^n g(k).$$

This allows constants to be factored out of a sum and sums of expressions to be separated.

Example. $\sum_{k=1}^n (2k + 3) = 2 \sum_{k=1}^n k + 3 \sum_{k=1}^n 1.$

Splitting a Sum

Remark (Splitting a sum). A sum over an interval may be divided into pieces:

$$\sum_{k=a}^c f(k) = \sum_{k=a}^b f(k) + \sum_{k=b+1}^c f(k) \quad \text{whenever } a \leq b < c.$$

Example. $\sum_{k=1}^{10} f(k) = \sum_{k=1}^5 f(k) + \sum_{k=6}^{10} f(k).$

Index Renaming

Remark (Index renaming). The index of summation is a *dummy variable*:

$$\sum_{k=a}^b f(k) = \sum_{i=a}^b f(i).$$

This is useful when manipulating expressions containing several sums, since the index variable can be renamed to avoid conflicts.

Index Shifting

Remark (Index shifting). It is often convenient to shift the summation index:

$$\sum_{k=a}^b f(k) = \sum_{k=a+1}^{b+1} f(k-1).$$

Index shifting allows sums to be aligned so that terms cancel or combine properly. It appears frequently in proofs involving recurrence relations and discrete calculus identities.

Example. $\sum_{k=0}^n f(k+1) = \sum_{k=1}^{n+1} f(k).$

Constant Sum

Remark (Constant sum). If the summand does not depend on the index, the sum counts terms:

$$\sum_{k=a}^b c = (b - a + 1) c.$$

The quantity $b - a + 1$ is the number of integers between a and b , inclusive.

Example. $\sum_{k=3}^7 5 = 5 \cdot 5 = 25.$

Empty Sum

Remark (Empty sum). If the upper limit is smaller than the lower limit, the sum contains no terms. By convention,

$$\sum_{k=a}^b f(k) = 0 \quad \text{whenever } a > b.$$

This convention simplifies many algebraic identities and allows formulas to remain valid without special cases.

Nested Sums

Remark (Nested sums). Summations may be nested:

$$\sum_{i=1}^n \sum_{j=1}^m a_{ij} = \sum_{i=1}^n \left(a_{i1} + a_{i2} + \cdots + a_{im} \right).$$

When the limits are independent, the order may be exchanged:

$$\sum_{i=1}^n \sum_{j=1}^m a_{ij} = \sum_{j=1}^m \sum_{i=1}^n a_{ij}.$$

Remark (Analogy with double integrals). Exchanging summation order is the discrete analogue of Fubini's theorem for iterated integrals.

Common Summation Identities

Remark (Common summation identities). Many sums that appear in discrete mathematics and discrete calculus have closed-form expressions. The following identities are among the most frequently encountered.

Common Summation Identities

$$\begin{aligned} \sum_{k=1}^n 1 &= n & \sum_{k=1}^n k &= \frac{n(n+1)}{2} \\ \sum_{k=1}^n k^2 &= \frac{n(n+1)(2n+1)}{6} & \sum_{k=1}^n k^3 &= \left(\frac{n(n+1)}{2}\right)^2 \\ \sum_{k=0}^n r^k &= \frac{1-r^{n+1}}{1-r} \quad (r \neq 1) & \sum_{k=0}^n \binom{n}{k} &= 2^n \end{aligned}$$

Remark (Role in discrete calculus). Polynomial summation formulas play an important role when working with finite differences and Newton series, while geometric sums arise naturally in recurrence relations and generating functions. Many of these identities can be derived systematically using the falling factorial basis and the Fundamental Theorem of Finite Calculus introduced later in this chapter.

5.4 Difference Operators

5.4.1 Difference Operators

Discrete calculus studies functions defined on the integers and the way their values change from one integer to the next. The central tool for measuring this change is the *difference operator*, which plays a role analogous to the derivative in continuous calculus.

Motivation

In ordinary calculus the derivative measures the rate of change of a function:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}.$$

In the discrete setting the independent variable changes in steps of size 1. Instead of taking a limit, we measure the difference between successive values of the function directly.

Forward Difference

Definition 5.4.1 (Forward Difference Operator). Let $f : \mathbb{Z} \rightarrow \mathbb{R}$. The *forward difference operator* Δ is defined by

$$\Delta f(n) = f(n+1) - f(n).$$

The function Δf measures the change in f when the input increases by one unit.

Remark (Fully quantified form). $\forall f : \mathbb{Z} \rightarrow \mathbb{R}, \forall n \in \mathbb{Z} : \Delta f(n) := f(n+1) - f(n)$.

Remark (Interpretation). The forward difference records the one-step change of a discrete function.

Remark (Analogy with differentiation). The operator Δ is the discrete analogue of the derivative. While the derivative measures infinitesimal change through a limit, the forward difference measures change between successive integer inputs exactly, with no limit required.

Example

Consider the function $f(n) = n^2$. Applying the difference operator gives

$$\begin{aligned}\Delta f(n) &= f(n+1) - f(n) \\ &= (n+1)^2 - n^2 \\ &= 2n+1.\end{aligned}$$

Thus the forward difference of the quadratic function n^2 is the linear function $2n+1$.

Remark (Degree reduction). This mirrors a familiar fact from ordinary calculus: $\frac{d}{dx}x^2 = 2x$. Just as derivatives reduce the degree of a polynomial by one, finite differences also lower the degree of polynomial sequences. Note, however, that $\Delta(n^2) = 2n+1$ rather than $2n$; ordinary powers do not behave as cleanly under Δ as they do under differentiation. This motivates introducing the *falling factorial* basis later in this chapter.

5.4.2 Higher Differences

Definition 5.4.2 (Higher Differences). The *second difference* of f is defined by

$$\Delta^2 f(n) = \Delta(\Delta f(n)).$$

More generally, the k -th difference $\Delta^k f(n)$ denotes the k -fold application of the difference operator, defined recursively by

$$\Delta^0 f = f, \quad \Delta^k f = \Delta(\Delta^{k-1} f).$$

Remark (Fully quantified form). $\forall k \in \mathbb{N}, \forall f : \mathbb{Z} \rightarrow \mathbb{R}, \forall n \in \mathbb{Z} : \Delta^k f(n) := \underbrace{(\Delta \circ \cdots \circ \Delta)}_k f(n)$

Remark (Interpretation). Higher differences measure repeated one-step change, just as higher derivatives measure repeated infinitesimal change.

Remark (Dependencies).

- [Forward Difference Operator](#).

Example

Let $f(n) = n^2$. Then

$$\begin{aligned}\Delta f(n) &= 2n+1, \\ \Delta^2 f(n) &= \Delta(2n+1) = 2(n+1) + 1 - (2n+1) = 2, \\ \Delta^3 f(n) &= \Delta(2) = 0.\end{aligned}$$

Remark (Analogy with higher derivatives). Higher differences behave analogously to higher derivatives. For polynomial functions of degree d , the d -th difference is constant and all higher differences are zero — just as the d -th derivative of a degree- d polynomial is constant and the next derivative vanishes.

5.4.3 Properties of the Difference Operator

The forward difference operator satisfies algebraic properties that closely mirror the familiar rules of differentiation.

Linearity

Proposition 5.4.3 (Linearity of Δ). *For functions $f, g : \mathbb{Z} \rightarrow \mathbb{R}$ and constants $a, b \in \mathbb{R}$,*

$$\Delta(af + bg) = a \Delta f + b \Delta g.$$

Remark (Fully quantified form). $\forall f, g : \mathbb{Z} \rightarrow \mathbb{R}, \forall a, b \in \mathbb{R}, \forall n \in \mathbb{Z} : \Delta(af + bg)(n) = a \Delta f(n) + b \Delta g(n)$.

Remark (Interpretation). Taking a forward difference respects pointwise linear combinations.

Remark (Dependencies).

- [Forward Difference Operator](#).

Proof. By definition of the difference operator,

$$\begin{aligned} \Delta(af + bg)(n) &= (af + bg)(n + 1) - (af + bg)(n) \\ &= af(n + 1) + bg(n + 1) - af(n) - bg(n) \\ &= a(f(n + 1) - f(n)) + b(g(n + 1) - g(n)) \\ &= a \Delta f(n) + b \Delta g(n). \quad \square \end{aligned}$$

Constant Rule

Proposition 5.4.4 (Constant rule). *If c is a constant function, then $\Delta c = 0$.*

Remark (Fully quantified form). $\forall c \in \mathbb{R}, \forall n \in \mathbb{Z} : \Delta c(n) = 0$.

Remark (Interpretation). A constant discrete function has no one-step change.

Remark (Dependencies).

- [Forward Difference Operator](#).

Proof. $\Delta c(n) = c - c = 0$. $\square$

Remark (Analogy). This mirrors $\frac{d}{dx}c = 0$ in continuous calculus.

The Shift Operator and Operator Algebra

Definition 5.4.5 (Shift Operator). The *shift operator* E and the *identity operator* I are defined by

$$Ef(n) = f(n + 1), \quad If(n) = f(n).$$

Remark (Fully quantified form). $\forall f : \mathbb{Z} \rightarrow \mathbb{R}, \forall n \in \mathbb{Z} : Ef(n) = f(n + 1)$ and $If(n) = f(n)$.

Remark (Interpretation). The shift operator advances the input by one step, while the identity operator leaves the input unchanged.

Theorem 5.4.6 ($\Delta = E - I$).

$$\Delta = E - I.$$

Remark (Fully quantified form). $\forall f : \mathbb{Z} \rightarrow \mathbb{R}, \forall n \in \mathbb{Z} : \Delta f(n) = (E - I)f(n) = f(n + 1) - f(n)$.

Remark (Interpretation). The forward difference is the shifted value minus the original value.

Remark (Dependencies).

- [Forward Difference Operator](#).
- [Shift Operator](#).

Proof. Applying the operator $E - I$ to a function f gives

$$(E - I)f(n) = Ef(n) - If(n) = f(n + 1) - f(n) = \Delta f(n). \quad \square$$

Remark (Higher differences via the binomial theorem). Expressing $\Delta = E - I$ reveals that higher differences can be computed using ordinary algebra. Applying the binomial theorem to $\Delta^n = (E - I)^n$ gives

$$\Delta^n = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} E^k.$$

Since $E^k f(n) = f(n + k)$, this yields

$$\Delta^n f(n) = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} f(n + k).$$

This explains why binomial coefficients and Pascal's triangle appear naturally throughout discrete calculus: they are a direct consequence of the algebraic identity $\Delta = E - I$.

5.4.4 Difference Tables and Polynomial Detection

One of the most useful tools in discrete calculus is the *difference table*. By repeatedly applying the difference operator to a sequence, one can detect whether the sequence arises from a polynomial function and determine its degree.

Definition 5.4.7 (Difference Table). Given a sequence $f(0), f(1), f(2), \dots$, the *difference table* is constructed by listing the successive values of the function and then computing their forward differences $\Delta f(n) = f(n + 1) - f(n)$, then $\Delta^2 f$, $\Delta^3 f$, and so on.

Remark (Interpretation). A difference table organizes repeated forward differences into rows so that eventual constant or zero rows become visible.

Remark (Dependencies).


- [Forward Difference Operator](#).
- [Higher Differences](#).

Example

For $f(n) = n^2$:

n	$f(n)$	$\Delta f(n)$	$\Delta^2 f(n)$
0	0	1	
1	1	3	2
2	4	5	2
3	9	7	2
4	16	9	2
5	25		

The second differences are constant (all equal to 2), and the third differences are zero.



Theorem 5.4.8 (Polynomial detection). *Let $f(n)$ be a sequence generated by a polynomial of degree d . Then $\Delta^d f(n)$ is constant, and $\Delta^{d+1} f(n) = 0$.*

Proof. Write $f(n) = a_d n^d + a_{d-1} n^{d-1} + \cdots + a_0$ with $a_d \neq 0$. We show by induction on d that Δf is a polynomial of degree $d - 1$ with leading coefficient $d a_d$.

Base case ($d = 0$): a constant polynomial has $\Delta f = 0$. ✓

Inductive step: Assume the result holds for all polynomials of degree less than d . Then

$$\Delta f(n) = f(n+1) - f(n) = a_d((n+1)^d - n^d) + (\text{lower-degree terms}).$$

Expanding $(n+1)^d - n^d = d n^{d-1} + \binom{d}{2} n^{d-2} + \cdots$, the leading term is $d a_d n^{d-1}$. Thus Δf is a polynomial of degree $d - 1$ with leading coefficient $d a_d \neq 0$.

By induction, applying Δ a further $d - 1$ times yields the constant $\Delta^d f(n) = d! a_d$, and one further application gives 0. $\square$

Remark (Fully quantified form). $\forall d \in \mathbb{N}, \forall f : \mathbb{Z} \rightarrow \mathbb{R}$ polynomial of degree d :
 $\exists c \in \mathbb{R} : \forall n \in \mathbb{Z}, \Delta^d f(n) = c$, and $\forall n \in \mathbb{Z}, \Delta^{d+1} f(n) = 0$.

Remark (Interpretation). Repeated differences reveal polynomial degree: a degree d polynomial has constant d -th differences and vanishing next differences.

Remark (Consequence). Difference tables provide a practical method for identifying polynomial sequences and determining their degree. This observation leads naturally to the *falling factorial basis*: a family of polynomials that interact particularly cleanly with the difference operator, forming the foundation for Newton's discrete analogue of the Taylor expansion.

Remark (Dependencies).

- [Forward Difference Operator](#).
- [Higher Differences](#).
- [Difference Table](#).
- [Binomial Theorem](#).

5.5 Telescoping Sums

5.5.1 Telescoping Sums

One of the most useful patterns in summation is the phenomenon of *telescoping*. In a telescoping sum, consecutive terms cancel with one another, leaving only a small number of terms from the beginning and end of the expression.

A Simple Example

Consider the sum

$$(2 - 1) + (3 - 2) + (4 - 3) + (5 - 4).$$

Expanding gives

$$2 - 1 + 3 - 2 + 4 - 3 + 5 - 4.$$

Most terms cancel in adjacent pairs:

$$\cancel{2} - 1 + \cancel{3} - \cancel{2} + \cancel{4} - \cancel{3} + 5 - \cancel{4}.$$

After cancellation, only the first and last terms remain: $5 - 1$. Thus

$$(2 - 1) + (3 - 2) + (4 - 3) + (5 - 4) = 5 - 1.$$

Definition (Telescoping Sum)

A sum is called a *telescoping sum* if consecutive terms cancel when the sum is expanded, leaving only a few remaining terms — typically the first and last.

General Form

The general telescoping structure is

$$\sum_{k=a}^{b-1} (f(k+1) - f(k)).$$

Writing out the terms explicitly:

$$\begin{aligned} & (f(a+1) - f(a)) \\ & + (f(a+2) - f(a+1)) \\ & + (f(a+3) - f(a+2)) \\ & \vdots \\ & + (f(b) - f(b-1)). \end{aligned}$$

All intermediate terms cancel in pairs, leaving only $f(b) - f(a)$.

Remark (Why “telescoping”). The name comes from an old collapsible telescope: each section slides into the next. The intermediate terms in the sum “slide together” and vanish, collapsing the entire sum down to its endpoints.

The Telescoping Identity

Theorem (Telescoping Identity)

For any function $f : \mathbb{Z} \rightarrow \mathbb{R}$ and integers $a \leq b$,

$$\sum_{k=a}^{b-1} (f(k+1) - f(k)) = f(b) - f(a).$$

Remark (Fully quantified form). $\forall f : \mathbb{Z} \rightarrow \mathbb{R}, \forall a, b \in \mathbb{Z}, a \leq b : \sum_{k=a}^{b-1} (f(k+1) - f(k)) = f(b) - f(a)$.

Proof. Expanding the sum gives

$$\begin{aligned} & (f(a+1) - f(a)) + (f(a+2) - f(a+1)) \\ & \quad + \cdots + (f(b) - f(b-1)). \end{aligned}$$

Every intermediate term $f(a+1), f(a+2), \dots, f(b-1)$ appears once with a positive sign and once with a negative sign, so they all cancel. The remaining terms are $f(b)$ and $-f(a)$, giving $f(b) - f(a)$. $\square$

Cancellation Pattern

The cancellation can be seen explicitly:

$$\begin{array}{ccc}
f(a+1) - f(a) & & -f(a) \\
\\
f(a+2) - f(a+1) & & \cancel{f(a+1)} - \cancel{f(a+1)} \\
\\
f(a+3) - f(a+2) & & \cancel{f(a+2)} - \cancel{f(a+2)} \\
\\
\vdots & & \vdots \\
\\
f(b) - f(b-1) & & f(b)
\end{array}$$

Remark (Intuition). Each intermediate term appears twice in the expanded sum: once with a positive sign and once with a negative sign. These pairs cancel, leaving only the first and last terms.

Operator Form

Using the forward difference operator $\Delta f(n) = f(n+1) - f(n)$, the telescoping identity takes the compact form

$$\sum_{k=a}^{b-1} \Delta f(k) = f(b) - f(a).$$

Remark (Discrete Fundamental Theorem of Calculus). This identity is the discrete analogue of the Fundamental Theorem of Calculus:

$$\int_a^b f'(x) dx = f(b) - f(a).$$

It reveals a deep relationship between the difference operator and summation: summation is the inverse of differencing, just as integration is the inverse of differentiation. This connection is made precise in the next section through the concept of the *discrete antiderivative*.

5.6 Discrete Antiderivatives

5.6.1 Discrete Antiderivatives

The telescoping identity established in the previous section,

$$\sum_{k=a}^{b-1} \Delta f(k) = f(b) - f(a),$$

reveals that summation acts as an inverse operation to the difference operator, much like integration acts as an inverse to differentiation in ordinary calculus. This motivates the following definition.

Definition 5.6.1 (Discrete Antiderivative). Let $f : \mathbb{Z} \rightarrow \mathbb{R}$ be a discrete function. A function $F : \mathbb{Z} \rightarrow \mathbb{R}$ is called a *discrete antiderivative* (or *discrete indefinite sum*) of f if

$$\Delta F(n) = f(n),$$

that is, $F(n+1) - F(n) = f(n)$ for all $n \in \mathbb{Z}$.

Remark (Standard quantified statement). $\forall f, F : \mathbb{Z} \rightarrow \mathbb{R} : F \text{ is a discrete antiderivative of } f \iff \forall n \in \mathbb{Z}, F(n+1) - F(n) = f(n).$ ■

Remark (Interpretation). A discrete antiderivative is a function whose one-step increments reproduce the given summand.

Remark (Analogy). A discrete antiderivative plays the same role in discrete calculus that an antiderivative plays in ordinary calculus: it reverses the fundamental differentiation-like operator of the theory. Instead of reversing differentiation, it reverses the forward difference Δ .

Remark (Dependencies).

- [Forward Difference Operator](#).

Examples

Example 1. Let $f(n) = 1$. Take $F(n) = n$. Then

$$\Delta F(n) = (n+1) - n = 1 = f(n). \checkmark$$

Example 2. Let $f(n) = n$. Take $F(n) = \frac{n(n-1)}{2}$. Then

$$\Delta F(n) = \frac{(n+1)n}{2} - \frac{n(n-1)}{2} = \frac{n(n+1-n+1)}{2} = n = f(n). \checkmark$$

Remark (Connection to falling factorials). Both examples above are instances of a general pattern: the discrete antiderivative of the falling factorial $n^{\underline{k}}$ is $\frac{1}{k+1}n^{\underline{k+1}}$, exactly mirroring the antiderivative rule for ordinary powers. This is developed in the Polynomial Basis section.

Constant of Integration

Discrete antiderivatives are not unique.

Proposition 5.6.2 (Constant of integration). *If F is a discrete antiderivative of f , then so is $F + C$ for any constant $C \in \mathbb{R}$.*

Remark (Fully quantified form). $\forall C \in \mathbb{R} : \Delta F = f \Rightarrow \Delta(F + C) = f.$

Remark (Interpretation). Adding a constant changes the level of a discrete antiderivative but not its one-step increments.

Remark (Dependencies).

- [Discrete Antiderivative](#).
- [Forward Difference Operator](#).

Proof.

$$\Delta(F + C)(n) = (F + C)(n + 1) - (F + C)(n) = F(n + 1) - F(n) = \Delta F(n) = f(n). \quad \square$$

Fundamental Theorem of Finite Calculus

Theorem (Fundamental Theorem of Finite Calculus)

Let F be a discrete antiderivative of f , so that $\Delta F(n) = f(n)$ for all n . Then

$$\sum_{k=a}^{b-1} f(k) = F(b) - F(a).$$

Remark (Fully quantified form). $\forall f, F : \mathbb{Z} \rightarrow \mathbb{R}, \forall a, b \in \mathbb{Z}, a \leq b : (\forall n, \Delta F(n) = f(n)) \Rightarrow \sum_{k=a}^{b-1} f(k) = F(b) - F(a).$

Proof. Since $f(k) = \Delta F(k)$ for all k , substituting gives

$$\sum_{k=a}^{b-1} f(k) = \sum_{k=a}^{b-1} \Delta F(k).$$

By the Telescoping Identity (Theorem [5.5.1](#)),

$$\sum_{k=a}^{b-1} \Delta F(k) = F(b) - F(a). \quad \square$$

Remark (Analogy with continuous calculus). This theorem is the discrete analogue of the Fundamental Theorem of Calculus,

$$\int_a^b f'(x) dx = f(b) - f(a).$$

In continuous calculus, differentiation and integration are inverse operations. In discrete calculus, the analogous inverse pair is the difference operator Δ and summation. Any sum can be evaluated by finding a discrete antiderivative of the summand — just as any definite integral can be evaluated by finding an antiderivative.

Remark (Practical consequence). The theorem reduces the problem of computing $\sum_{k=a}^{b-1} f(k)$ to the algebraic problem of finding F with $\Delta F = f$. Combined with the falling factorial rules developed in the next section, this gives a systematic method for evaluating a wide family of sums in closed form.

5.7 Polynomial Basis

5.7.1 Polynomial Basis: Falling Factorials

In ordinary calculus, polynomial functions are naturally expressed using the monomial powers $1, x, x^2, x^3, \dots$ because the derivative acts simply on them: $\frac{d}{dx}x^n = nx^{n-1}$.

In discrete calculus, however, the difference operator does not behave as cleanly on ordinary powers. For example,

$$\Delta(n^2) = (n+1)^2 - n^2 = 2n + 1.$$

Although the degree decreases by one, the result contains an extra constant term. For this reason it is more natural to work with a different family of polynomials — the *falling factorials* — which interact with Δ exactly as monomials interact with $\frac{d}{dx}$.

Definition (Falling Factorial)

For a nonnegative integer k , the *falling factorial* (or *falling power*) is defined by

$$x^{\underline{k}} = x(x-1)(x-2)\cdots(x-k+1) \quad (k \text{ factors}).$$

By convention, $x^{\underline{0}} = 1$.

Remark (Fully quantified form). $\forall k \in \mathbb{N}, \forall x \in \mathbb{R} : x^{\underline{k}} := \prod_{j=0}^{k-1} (x-j)$, with $x^{\underline{0}} := 1$.

Remark (Relation to factorials). When $x = n \in \mathbb{N}$ and $k \leq n$, the falling factorial equals $\frac{n!}{(n-k)!}$. In particular, $n^{\underline{n}} = n!$. Thus falling factorials generalise factorials to arbitrary inputs.

The first few falling factorials are:

$$\begin{aligned} x^{\underline{0}} &= 1, \\ x^{\underline{1}} &= x, \\ x^{\underline{2}} &= x(x-1), \\ x^{\underline{3}} &= x(x-1)(x-2). \end{aligned}$$

The Difference Rule for Falling Factorials

Theorem (Difference Rule for Falling Factorials)

For $k \geq 1$,

$$\Delta x^{\underline{k}} = k x^{\underline{k-1}}.$$

Remark (Fully quantified form). $\forall k \in \mathbb{N}, k \geq 1, \forall x \in \mathbb{R} : (x+1)^{\underline{k}} - x^{\underline{k}} = k x^{\underline{k-1}}$.

Proof. Observe that

$$(x+1)^{\underline{k}} = (x+1)x(x-1)\cdots(x-k+2).$$

Therefore

$$\Delta x^{\underline{k}} = (x+1)^{\underline{k}} - x^{\underline{k}} = x(x-1)\cdots(x-k+2)[(x+1) - (x-k+1)].$$

The bracket simplifies:

$$(x+1) - (x-k+1) = k.$$

The remaining product $x(x-1)\cdots(x-k+2)$ is exactly $x^{\underline{k-1}}$. Thus

$$\Delta x^{\underline{k}} = k x^{\underline{k-1}}. \quad \square$$

Remark (Mirror of the power rule). This formula exactly mirrors the continuous power rule $\frac{d}{dx}x^k = kx^{k-1}$. Because of this clean behaviour under Δ , falling factorials form the natural polynomial basis for discrete calculus.

Remark (Antiderivative rule). The theorem also yields the discrete antiderivative rule:

$$\Delta\left(\frac{x^{\underline{k+1}}}{k+1}\right) = x^{\underline{k}}.$$

Thus $\frac{x^{\underline{k+1}}}{k+1}$ is a discrete antiderivative of $x^{\underline{k}}$, mirroring the continuous rule $\int x^k dx = \frac{x^{k+1}}{k+1}$.

Summary Table

$f(x)$	$\Delta f(x)$
$x^{\underline{1}} = x$	1
$x^{\underline{2}} = x(x-1)$	$2x$
$x^{\underline{3}} = x(x-1)(x-2)$	$3x(x-1)$
$x^{\underline{4}} = x(x-1)(x-2)(x-3)$	$4x(x-1)(x-2)$

Polynomial Representation in the Falling Factorial Basis

Every polynomial can be written as a linear combination of falling factorials:

$$f(x) = c_0 + c_1 x + c_2 x^{\underline{2}} + c_3 x^{\underline{3}} + \cdots + c_n x^{\underline{n}}.$$

Remark (Role in Newton's expansion). This representation plays the same role in discrete calculus that the Taylor expansion plays in continuous calculus. The coefficients c_k are determined by the successive finite differences of f evaluated at 0: $c_k = \Delta^k f(0)/k!$. This leads directly to Newton's forward difference expansion, developed in the Expansions section.

Consequences of the Difference Rule

The difference rule $\Delta x^{\underline{k}} = k x^{\underline{k-1}}$ (Theorem 5.7.1) has several important consequences.

Higher Differences of Falling Factorials

Applying the rule repeatedly gives:

$$\Delta^m x^{\underline{k}} = \begin{cases} \frac{k!}{(k-m)!} x^{\underline{k-m}} & \text{if } m \leq k, \\ 0 & \text{if } m > k. \end{cases}$$

In particular, evaluating at $x = 0$:

$$\Delta^m x^{\underline{k}}|_{x=0} = \begin{cases} k! & \text{if } m = k, \\ 0 & \text{if } m \neq k. \end{cases}$$

Remark (Basis orthogonality). This vanishing property means the falling factorials behave like a “triangular” basis: Δ^m kills all $x^{\underline{k}}$ with $k < m$, and returns a scalar multiple for $k = m$. This is what makes the coefficient extraction in Newton’s expansion work cleanly.

Connection to Binomial Coefficients

The higher difference formula from the Binomial Theorem section (Proposition 5.2.2),

$$\Delta^n f(x) = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} f(x+k),$$

together with the difference rule, gives an explicit formula for the coefficients of the falling factorial expansion:

$$c_k = \frac{\Delta^k f(0)}{k!}.$$

Remark (Pascal’s triangle revisited). The coefficients appearing in higher finite differences follow exactly the pattern of Pascal’s triangle — a direct consequence of the algebraic identity $\Delta = E - I$ and the binomial theorem.

Worked Example: Expanding n^2

We find the falling factorial expansion of $f(n) = n^2$.

Computing the difference table at $n = 0$:

$$f(0) = 0, \quad \Delta f(0) = 1, \quad \Delta^2 f(0) = 2, \quad \Delta^3 f(0) = 0.$$

The coefficients are $c_k = \Delta^k f(0)/k!$:

$$c_0 = 0, \quad c_1 = 1, \quad c_2 = \frac{2}{2!} = 1, \quad c_3 = 0.$$

Therefore:

$$n^2 = 0 + n^1 + n^2 = n + n(n-1) = n^2 - n + n = n^2. \checkmark$$

Remark (Practical value). Once a function is expressed in the falling factorial basis, both differencing and antidifferencing become purely mechanical, following the rule $\Delta x^{\underline{k}} = k x^{\underline{k-1}}$ and its inverse. This makes the Fundamental Theorem of Finite Calculus immediately applicable to compute sums like $\sum_{k=0}^{n-1} k^2$ without clever tricks.

5.8 Calculus Rules

5.8.1 Calculus Rules

With the difference operator and the polynomial basis in place, we can now develop the full suite of differentiation-like rules for discrete calculus. The linearity of Δ was established in the Difference Operators section (Proposition 5.4.3). Here we add the *product rule* — the discrete counterpart of the Leibniz rule $(fg)' = f'g + fg'$.

Product Rule

Theorem (Discrete Product Rule)

For functions $f, g : \mathbb{Z} \rightarrow \mathbb{R}$,

$$\Delta(fg)(n) = f(n) \Delta g(n) + g(n+1) \Delta f(n).$$

Remark (Fully quantified form). $\forall f, g : \mathbb{Z} \rightarrow \mathbb{R}, \forall n \in \mathbb{Z} : (f(n+1)g(n+1)) - (f(n)g(n)) = f(n)(g(n+1) - g(n)) + g(n+1)(f(n+1) - f(n))$.

Proof. By definition of Δ ,

$$\Delta(fg)(n) = f(n+1)g(n+1) - f(n)g(n).$$

Add and subtract $f(n)g(n+1)$:

$$= f(n+1)g(n+1) - f(n)g(n+1) + f(n)g(n+1) - f(n)g(n).$$

Grouping the first two and last two terms:

$$= g(n+1)(f(n+1) - f(n)) + f(n)(g(n+1) - g(n)) = g(n+1) \Delta f(n) + f(n) \Delta g(n). \quad \square$$

Remark (Comparison with the continuous product rule). In continuous calculus, $(fg)' = f'g + fg'$ is symmetric in f and g . In discrete calculus the rule is slightly asymmetric: one factor is evaluated at n and the other at $n+1$. This asymmetry is unavoidable and is a characteristic feature of the discrete setting.

Remark (Consequence — summation by parts). The product rule leads directly to the summation by parts formula: summing both sides of $\Delta(fg) = f \Delta g + g(E) \Delta f$ over $k = a$ to $b-1$ and applying the Fundamental Theorem gives the formula in the next section.

Summation by Parts

Summation by parts is the discrete analogue of integration by parts. It is derived from the product rule exactly as the continuous version is derived from the Leibniz rule.

Theorem (Summation by Parts)

For functions $f, g : \mathbb{Z} \rightarrow \mathbb{R}$ and integers $a \leq b$,

$$\sum_{k=a}^{b-1} f(k) \Delta g(k) = \left[f(k)g(k) \right]_{k=a}^{k=b} - \sum_{k=a}^{b-1} g(k+1) \Delta f(k).$$

That is,

$$\sum_{k=a}^{b-1} f(k) \Delta g(k) = f(b)g(b) - f(a)g(a) - \sum_{k=a}^{b-1} g(k+1) \Delta f(k).$$

Remark (Fully quantified form). $\forall f, g : \mathbb{Z} \rightarrow \mathbb{R}, \forall a, b \in \mathbb{Z}, a \leq b : \sum_{k=a}^{b-1} f(k) \Delta g(k) =$

$$f(b)g(b) - f(a)g(a) - \sum_{k=a}^{b-1} g(k+1) \Delta f(k).$$

Proof. By the Discrete Product Rule (Theorem 5.8.1),

$$\Delta(fg)(k) = f(k) \Delta g(k) + g(k+1) \Delta f(k).$$

Rearranging,

$$f(k) \Delta g(k) = \Delta(fg)(k) - g(k+1) \Delta f(k).$$

Sum both sides from $k = a$ to $k = b - 1$:

$$\sum_{k=a}^{b-1} f(k) \Delta g(k) = \sum_{k=a}^{b-1} \Delta(fg)(k) - \sum_{k=a}^{b-1} g(k+1) \Delta f(k).$$

Applying the Fundamental Theorem of Finite Calculus (Theorem 5.6.1) to the first sum on the right gives

$$\sum_{k=a}^{b-1} \Delta(fg)(k) = f(b)g(b) - f(a)g(a),$$

which completes the proof. □

Remark (Analogy with integration by parts). The continuous integration by parts formula,

$$\int_a^b f dg = [fg]_a^b - \int_a^b g df,$$

is recovered from summation by parts by replacing $\sum$ with $\int$ and Δ with d . The asymmetry in the discrete formula — $g(k+1)$ rather than $g(k)$ in the remaining sum — reflects the shift inherent in the product rule.

Remark (Applications). Summation by parts is useful for evaluating sums involving products, particularly when one factor is easy to sum (as a falling factorial) and the other is easy to difference. Classic applications include evaluating $\sum kr^k$ (where $f(k) = k$, $\Delta g(k) = r^k$ leads to a geometric-type sum) and establishing Abel's summation formula.

5.9 Expansions

5.9.1 Newton's Forward Difference Expansion

In continuous calculus, smooth functions can be expanded as power series called Taylor series:

$$f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{k!} x^k = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \cdots.$$

The coefficients are determined by successive derivatives evaluated at 0, and the basis functions are ordinary powers x^k .

Discrete calculus has a precise analogue: the *Newton forward difference expansion*, in which derivatives are replaced by finite differences and ordinary powers are replaced by falling factorials.

Theorem (Newton Forward Difference Expansion)

Let $f : \mathbb{Z} \rightarrow \mathbb{R}$. Then f can be expressed as

$$f(x) = \sum_{k=0}^n \frac{\Delta^k f(0)}{k!} x^{\underline{k}} = f(0) + \Delta f(0)x + \frac{\Delta^2 f(0)}{2!}x^{\underline{2}} + \frac{\Delta^3 f(0)}{3!}x^{\underline{3}} + \cdots,$$

where $x^{\underline{k}} = x(x-1)\cdots(x-k+1)$ is the falling factorial. For a polynomial of degree n , the expansion is exact (the series terminates at $k = n$).

Remark (Fully quantified form). $\forall f : \mathbb{Z} \rightarrow \mathbb{R}$ polynomial of degree $\leq n$, $\forall x \in \mathbb{Z} : f(x) = \sum_{k=0}^n \binom{x}{k} \Delta^k f(0)$, where $\binom{x}{k} := x^{\underline{k}}/k!$.

Structural Parallel with the Taylor Series

The Newton expansion mirrors the Taylor series at every point. The correspondence is exact:

Taylor Series (continuous)	Newton Expansion (discrete)
Coefficients $f^{(k)}(0)/k!$	Coefficients $\Delta^k f(0)/k!$
Basis functions x^k	Basis functions $x^{\underline{k}}$
Derivative $f^{(k)}$	Difference Δ^k
Convergence required in general	Exact for polynomials

Remark (Why falling factorials play the role of powers). In the Taylor series, powers x^k are the natural basis because the derivative satisfies $\frac{d}{dx}x^k = kx^{k-1}$ and $\frac{d^m}{dx^m}x^k|_{x=0} = k!$ when $m = k$, and 0 otherwise. The falling factorials satisfy the same two properties under Δ : $\Delta x^{\underline{k}} = kx^{\underline{k-1}}$ and $\Delta^m x^{\underline{k}}|_{x=0} = k!$ when $m = k$, else 0. This “triangular” property is exactly what allows the coefficients $c_k = \Delta^k f(0)/k!$ to be read off independently, level by level.

Proof Sketch

We construct the expansion term by term, matching the values of f at successive integers.

The constant term $f(0)$ matches f at $x = 0$.


Adding the linear term $\Delta f(0) \cdot x$ matches f at $x = 0$ and $x = 1$.

Adding the quadratic term $\frac{\Delta^2 f(0)}{2!} x^2$ corrects the value at $x = 2$ while leaving earlier matches intact, because $x^2 = x(x-1)$ vanishes at $x = 0$ and $x = 1$.

At each stage, x^k vanishes at $x = 0, 1, \dots, k-1$, so the k -th term corrects only the value at $x = k$. The coefficient $\Delta^k f(0)/k!$ achieves this correction by the triangular property noted above.

Reading Coefficients from the Difference Table

The coefficients in the Newton expansion are read from the *left edge* of the difference table — the “Newton diagonal”:



Each level of the difference table contributes exactly one term to the falling-factorial expansion.

Worked Example: Expanding $f(n) = n^3$

We compute the difference table at $n = 0$:

n	$f(n)$	$\Delta f(n)$	$\Delta^2 f(n)$	$\Delta^3 f(n)$
0	0	1	6	6
1	1	7	12	
2	8	19		
3	27			

The coefficients are $c_k = \Delta^k f(0)/k!$:

$$c_0 = 0, \quad c_1 = 1, \quad c_2 = \frac{6}{2} = 3, \quad c_3 = \frac{6}{6} = 1.$$

Therefore the Newton expansion is:

$$n^3 = n + 3n^2 + n^3 = n + 3n(n-1) + n(n-1)(n-2).$$

Verification: $n + 3n^2 - 3n + n^3 - 3n^2 + 2n = n^3$. ✓

Application: Computing $\sum_{k=0}^{n-1} k^2$

Using the expansion $k^2 = k^1 + k^2$ and the Fundamental Theorem of Finite Calculus (Theorem 5.6.1),

$$\sum_{k=0}^{n-1} k^2 = \sum_{k=0}^{n-1} k^1 + \sum_{k=0}^{n-1} k^2.$$

Since $\Delta\left(\frac{k^2}{2}\right) = k^1$ and $\Delta\left(\frac{k^3}{3}\right) = k^2$, the Fundamental Theorem gives:

$$\sum_{k=0}^{n-1} k^1 = \frac{n^2}{2} - \frac{0^2}{2} = \frac{n(n-1)}{2},$$

$$\sum_{k=0}^{n-1} k^2 = \frac{n^3}{3} - \frac{0^3}{3} = \frac{n(n-1)(n-2)}{3}.$$

Adding:

$$\sum_{k=0}^{n-1} k^2 = \frac{n(n-1)}{2} + \frac{n(n-1)(n-2)}{3} = \frac{n(n-1)}{6} [3 + 2(n-2)] = \frac{n(n-1)(2n-1)}{6}.$$

Remark (Check against the standard formula). The standard formula is $\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$.

The computed result $\sum_{k=0}^{n-1} k^2 = \frac{(n-1)n(2n-1)}{6}$ matches: substitute $n \mapsto n-1$ in the standard formula to get $\frac{(n-1)n(2n-1)}{6}$. ✓

This example illustrates the power of the method: the sum was evaluated by pure algebraic antidifferentiation, with no need for induction or clever manipulation.

5.10 Special Functions

5.10.1 Special Functions in Discrete Calculus

Just as continuous calculus has its distinguished special functions — the exponential e^{ax} , the trigonometric functions, and the logarithm — discrete calculus has analogous functions that behave simply under the difference operator.

The Discrete Exponential

In continuous calculus, the exponential function e^{ax} is its own derivative (up to a constant):

$$\frac{d}{dx} e^{ax} = a e^{ax}.$$

The analogous function in discrete calculus is the *discrete exponential* $(1+a)^n$.

Definition 5.10.1 (Discrete Exponential). For a constant $a \neq -1$ and $n \in \mathbb{Z}$, the *discrete exponential* with base $(1 + a)$ is the function

$$E_a(n) = (1 + a)^n.$$

Remark (Fully quantified form). $\forall a \in \mathbb{R} \setminus \{-1\}, \forall n \in \mathbb{Z} : E_a(n) := (1 + a)^n$.

Remark (Interpretation). The discrete exponential is the geometric function whose ratio between successive inputs is $1 + a$.

Proposition 5.10.2 (Discrete exponential rule).

$$\Delta(1 + a)^n = a(1 + a)^n.$$

Remark (Fully quantified form). $\forall a \in \mathbb{R} \setminus \{-1\}, \forall n \in \mathbb{Z} : (1 + a)^{n+1} - (1 + a)^n = a(1 + a)^n$.

Remark (Interpretation). The forward difference of a discrete exponential is a scalar multiple of the same discrete exponential.

Proof.

$$\Delta(1 + a)^n = (1 + a)^{n+1} - (1 + a)^n = (1 + a)^n[(1 + a) - 1] = a(1 + a)^n. \quad \square$$

Remark (Analogy). The continuous exponential satisfies $\frac{d}{dx}e^{ax} = ae^{ax}$. The discrete exponential satisfies $\Delta(1 + a)^n = a(1 + a)^n$. The base e in continuous calculus is replaced by $(1 + a)$ in discrete calculus. When a is small, $(1 + a) \approx e^a$, recovering the continuous limit.

Remark (Antiderivative). From the proposition, a discrete antiderivative of $a(1 + a)^n$ is $(1 + a)^n$ itself. Equivalently, a discrete antiderivative of $(1 + a)^n$ is $\frac{(1+a)^n}{a}$. Combined with the Fundamental Theorem of Finite Calculus, this gives the geometric series formula:

$$\sum_{k=0}^{n-1} (1 + a)^k = \frac{(1 + a)^n - 1}{a} \quad (a \neq 0).$$

Remark (Dependencies).

- [Discrete Exponential](#).
- [Forward Difference Operator](#).

5.10.2 The Function 2^n in Discrete Calculus

Among all discrete exponentials, the case $a = 1$ — giving $(1 + 1)^n = 2^n$ — is perhaps the most important. It appears in combinatorics as the count of subsets, in probability as the sample space of fair coin sequences, in computer science as the natural measure of binary information, and in discrete calculus as a canonical test function. This section develops its properties systematically.

Differencing 2^n

Setting $a = 1$ in the discrete exponential rule (Proposition 5.10.2) gives immediately:

Proposition ($\Delta 2^n = 2^n$)

$$\Delta 2^n = 2^n.$$

Remark (Standard quantified statement). $\forall n \in \mathbb{Z} : 2^{n+1} - 2^n = 2^n$.

Proof.

$$\Delta 2^n = 2^{n+1} - 2^n = 2 \cdot 2^n - 2^n = (2 - 1) \cdot 2^n = 2^n. \quad \square$$

Remark (Self-similarity). This says that 2^n is its own forward difference. In continuous calculus, the analogous property is $\frac{d}{dx} e^x = e^x$: the exponential is its own derivative. The function 2^n is the discrete analogue of e^x in the sense that $\Delta(2^n) = 2^n$. More precisely, $2^n = e^{n \ln 2}$, and the relationship $\Delta(2^n) = 2^n$ corresponds to the special case $a = 1$ of $\Delta(1 + a)^n = a(1 + a)^n$.

Antiderivative and Geometric Sum

Since $\Delta(2^n) = 2^n$, a discrete antiderivative of 2^n is 2^n itself (up to a constant).

Proposition (Antiderivative of 2^n)

A discrete antiderivative of 2^n is 2^n itself:

$$\Delta(2^n) = 2^n \quad \text{so} \quad \sum_{k=0}^{n-1} 2^k = 2^n - 1.$$

Remark (Standard quantified statement). $\forall n \in \mathbb{N} : \sum_{k=0}^{n-1} 2^k = 2^n - 1$.

Proof. Applying the Fundamental Theorem of Finite Calculus (Theorem 5.6.1) with $F(k) = 2^k$ and $f(k) = \Delta F(k) = 2^k$:

$$\sum_{k=0}^{n-1} 2^k = F(n) - F(0) = 2^n - 2^0 = 2^n - 1. \quad \square$$

Remark (Verification by expansion). Expanding: $1 + 2 + 4 + 8 + \cdots + 2^{n-1}$. Multiply the partial sum $S = \sum_{k=0}^{n-1} 2^k$ by 2 to get $2S = \sum_{k=1}^n 2^k$. Subtracting: $2S - S = 2^n - 1$, so $S = 2^n - 1$. This is the classical derivation of the geometric series formula; the discrete calculus approach via antidifferentiation gives the same result mechanically.

Higher Differences of 2^n

Proposition 5.10.3 (Higher differences of 2^n). *For all $k \geq 0$,*

$$\Delta^k 2^n = 2^n.$$

Remark (Standard quantified statement). $\forall k \in \mathbb{N}, \forall n \in \mathbb{Z} : \Delta^k(2^n) = 2^n$.

Remark (Interpretation). Every forward-difference row for 2^n reproduces the same function.

Proof. By induction. The base case $k = 0$ is $\Delta^0(2^n) = 2^n$ by definition. For the inductive step, assume $\Delta^k(2^n) = 2^n$. Then

$$\Delta^{k+1}(2^n) = \Delta(\Delta^k(2^n)) = \Delta(2^n) = 2^n. \quad \square$$

Remark (Consequence for difference tables). Since every finite difference of 2^n is 2^n again, the difference table of 2^n consists entirely of the sequence 2^n at every level. This is the discrete analogue of the fact that the Taylor series of e^x has the same coefficient pattern at every derivative level. In particular, 2^n is *not* a polynomial — a polynomial of degree d would reach 0 at the $(d + 1)$ -th difference row. The non-termination of the difference table is the signature of a genuinely transcendental discrete function.

Remark (Dependencies).

- [Higher Differences](#).
- [Discrete Exponential Rule](#).

2^n in Combinatorics and the Newton Expansion

The Binomial Theorem (Theorem 5.2.1) gives a direct connection: setting $x = y = 1$ yields

$$\sum_{k=0}^n \binom{n}{k} = 2^n.$$

This counts the total number of subsets of an n -element set: there are $\binom{n}{k}$ subsets of size k , and the sum over all k gives 2^n .

More deeply, the Newton expansion (Theorem 5.9.1) of 2^n in the falling factorial basis is:

$$2^n = \sum_{k=0}^n \binom{n}{k} = \sum_{k=0}^n \frac{n^{\underline{k}}}{k!},$$

since $\Delta^k(2^n)|_{n=0} = 2^0 = 1$ for every k . This expresses 2^n as the sum of all normalised falling factorials up to degree n , and reveals why binomial coefficients count subsets: they are the coefficients of 2^n in its own Newton expansion.

Remark (Analogy with e^x). In continuous calculus, $e^x = \sum_{k=0}^{\infty} x^k/k!$. The Newton expansion gives $2^n = \sum_{k=0}^n n^{\underline{k}}/k!$. Both express a canonical exponential as a sum of normalised basis functions, with all coefficients equal to 1. The analogy is exact: ordinary powers replace falling factorials, and the infinite series terminates at degree n in the discrete case because 2^n has degree exactly n in the falling factorial basis.

Discrete Differential Equation

In continuous calculus, the equation $f' = f$ (with $f(0) = 1$) has the unique solution $f(x) = e^x$. In discrete calculus, the analogous *difference equation* is:

$$\Delta F(n) = F(n), \quad F(0) = 1.$$

That is, $F(n+1) - F(n) = F(n)$, so $F(n+1) = 2F(n)$. The unique solution to this initial value problem is $F(n) = 2^n$.

Proposition 5.10.4 (Discrete IVP). *The unique function $F : \mathbb{N} \rightarrow \mathbb{R}$ satisfying*

$$F(n+1) = 2F(n), \quad F(0) = 1$$

is $F(n) = 2^n$.

Remark (Standard quantified statement). $\forall n \in \mathbb{N} : (F(n+1) = 2F(n) \text{ and } F(0) = 1) \Rightarrow F(n) = 2^n$.

Remark (Interpretation). The recurrence $F(n+1) = 2F(n)$ with initial value 1 determines the discrete exponential 2^n uniquely.

Proof. Uniqueness: if G is another solution, then $G(0) = 1 = F(0)$, and inductively $G(n+1) = 2G(n) = 2F(n) = F(n+1)$, so $G = F$. Existence: verify directly that $2^{n+1} = 2 \cdot 2^n$ and $2^0 = 1$. $\square$

Remark (Significance). This characterises 2^n as the unique fixed point of “multiply by 2” starting from 1. The analogous characterisation of e^x is: unique solution to $f' = f$, $f(0) = 1$. Both are initial-value problems in their respective calculi, and both have closed-form solutions that serve as the canonical exponential of that calculus.

Remark (Dependencies).

- [Discrete Exponential](#).

Difference Formulas for Special Functions

The table below collects the difference formulas for the principal special functions of discrete calculus.

Function	Forward Difference
$x^{\underline{k}}$ (falling factorial)	$k x^{\underline{k-1}}$
$(1+a)^n$ (discrete exponential)	$a(1+a)^n$
<i>Remark</i> (Difference formulas). $\binom{x}{k} = x^{\underline{k}}/k!$	$\binom{x}{k-1}$
c (constant)	0
n	1
$n^2 = n(n-1)$	$2n$

Remark (Binomial coefficient as a special case). The formula $\Delta \binom{x}{k} = \binom{x}{k-1}$ follows immediately from the falling factorial rule $\Delta x^{\underline{k}} = k x^{\underline{k-1}}$ by dividing through by $k!$. It says that the binomial coefficient $\binom{x}{k}$ is its own natural “unit” in discrete calculus, analogous to the monomial $x^k/k!$ in the Taylor expansion.

Antiderivative Table

By reversing the difference formulas, we obtain the corresponding discrete antiderivatives:

Function $f(n)$	Antiderivative $F(n)$ with $\Delta F = f$
$x^{\underline{k}}$	$\frac{x^{\underline{k+1}}}{k+1}$
<i>Remark</i> (Discrete antiderivatives). $(1+a)^n$ ($a \neq 0$)	$\frac{(1+a)^n}{a}$
$\binom{x}{k}$	$\binom{x}{k+1}$
c (constant)	cn

Remark (Using the Fundamental Theorem). Each antiderivative in the table, combined with the Fundamental Theorem of Finite Calculus (Theorem 5.6.1), yields an immediate summation formula. For instance, from $\Delta \binom{x^{\underline{k+1}}}{k+1} = x^{\underline{k}}$:

$$\sum_{j=0}^{n-1} j^{\underline{k}} = \frac{n^{\underline{k+1}}}{k+1}.$$

This is the discrete analogue of $\int_0^n x^k dx = \frac{n^{k+1}}{k+1}$.

Proofs

Remark (Return). [Return to Theorem](#)

Theorem (Binomial Theorem). *For any integer $n \geq 0$ and real numbers x, y ,*

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k.$$

Professional Standard Proof.

TODO: professional standard proof for binomial. □

Detailed Learning Proof.

TODO: detailed learning proof for binomial. □

Remark (Proof structure).

Remark (Return). [Return to Theorem](#)

Proposition (Higher difference formula). *For any function f ,*

$$\Delta^n f(x) = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} f(x+k).$$

Professional Standard Proof.

TODO: professional standard proof for higher-diff-formula. □

Detailed Learning Proof.

TODO: detailed learning proof for higher-diff-formula. □

Remark (Proof structure).

Remark (Dependencies).

- [Binomial Theorem](#).
- [Forward Difference Operator](#).
- [Higher Differences](#).

Remark (Return). [Return to Theorem](#)

Proposition (Linearity of Δ). *The forward difference operator is linear.*

Professional Standard Proof.

TODO: professional standard proof for delta-linearity. □

Detailed Learning Proof.

TODO: detailed learning proof for delta-linearity. □

Remark (Proof structure).

Remark (Dependencies).

- [Forward Difference Operator](#).

Remark (Return). [Return to Theorem](#)

Proposition (Constant rule). *The forward difference of a constant sequence is zero.*

Professional Standard Proof.

TODO: professional standard proof for delta-constant. □

Detailed Learning Proof.

TODO: detailed learning proof for delta-constant. □

Remark (Proof structure).

Remark (Dependencies).

- [Forward Difference Operator](#).

Remark (Return). [Return to Theorem](#)

Theorem ($\Delta = E - I$). *The forward difference operator is the shift operator minus the identity operator.*

Professional Standard Proof.

TODO: professional standard proof for delta-shift-minus-identity. □

Detailed Learning Proof.

TODO: detailed learning proof for delta-shift-minus-identity. □

Remark (Proof structure).

Remark (Dependencies).

- [Forward Difference Operator](#).
- [Shift Operator](#).

Remark (Return). [Return to Theorem](#)

Theorem (Polynomial detection). *Finite differences detect polynomial degree.*

Professional Standard Proof.

TODO: professional standard proof for polynomial-detection. □

Detailed Learning Proof.

TODO: detailed learning proof for polynomial-detection. □

Remark (Proof structure).

Remark (Dependencies).

- [Forward Difference Operator](#).
- [Higher Differences](#).
- [Difference Table](#).
- [Binomial Theorem](#).

Remark (Return). [Return to Theorem](#)

Proposition (Constant of integration). *Discrete antiderivatives are unique up to an additive constant.*

Professional Standard Proof.

TODO: professional standard proof for constant-of-integration. □

Detailed Learning Proof.

TODO: detailed learning proof for constant-of-integration. □

Remark (Proof structure).

Remark (Dependencies).

- [Discrete Antiderivative](#).
- [Forward Difference Operator](#).

Remark (Return). [Return to Theorem](#)

Proposition (Discrete exponential rule). *The discrete exponential has geometric forward differences.*

Professional Standard Proof.

TODO: professional standard proof for discrete-exponential-rule. □

Detailed Learning Proof.

TODO: detailed learning proof for discrete-exponential-rule. □

Remark (Proof structure).

Remark (Dependencies).

- [Discrete Exponential](#).
- [Forward Difference Operator](#).

Remark (Return). [Return to Theorem](#)

Proposition (Higher differences of 2^n). *All higher forward differences of 2^n are again 2^n .*

Professional Standard Proof.

TODO: professional standard proof for higher-diff-two-n. □

Detailed Learning Proof.

TODO: detailed learning proof for higher-diff-two-n. □

Remark (Proof structure).

Remark (Dependencies).

- [Higher Differences](#).
- [Discrete Exponential Rule](#).

Remark (Return). [Return to Theorem](#)

Proposition (Discrete IVP). *A first-order discrete initial value problem has the stated solution.*

Professional Standard Proof.

TODO: professional standard proof for discrete-ivp. □

Detailed Learning Proof.

TODO: detailed learning proof for discrete-ivp. □

Remark (Proof structure).

Remark (Dependencies).

- [Discrete Exponential](#).

Chapter 6

Sequences



6.1 Sequence Definitions

6.1.1 Definitions

Definition (Sequence)

Definition 6.1.1 (Sequence). Let X be a nonempty set. A **sequence** in X is a function

$$x : \mathbb{N} \rightarrow X.$$

The value $x(n)$ is written x_n and called the **n -th term** of the sequence. The sequence itself is denoted $(x_n)_{n \in \mathbb{N}}$, or simply (x_n) when the index set is clear.

Remark (Standard quantified statement).

$$\forall X \neq \emptyset \forall x : \mathbb{N} \rightarrow X \text{ (Sequence}(x, X) \iff x : \mathbb{N} \rightarrow X).$$

Remark (Definition predicate reading).

$$\text{Sequence}(x_n, X) := x : \mathbb{N} \rightarrow X.$$

Remark (Interpretation). A sequence is not a set of terms but a function: the index carries independent weight. Two sequences (x_n) and (y_n) in X are equal precisely when $x_n = y_n$ for every $n \in \mathbb{N}$, so order and repetition are intrinsic to the object. The notational collapse from $x(n)$ to x_n is conventional suppression of function-application syntax; the underlying object remains a function $\mathbb{N} \rightarrow X$. When $X = \mathbb{R}$, the sequence is called a **real sequence**, which is the default ambient context for this chapter.

Remark (Dependencies).

- [Function](#)
- [Natural Numbers](#)

Definition (Real Sequence)

Definition 6.1.2 (Real Sequence). A **real sequence** is a sequence whose image is contained in $\mathbb{R}$. Equivalently, a real sequence is a function

$$x : \mathbb{N} \rightarrow \mathbb{R}.$$

Remark (Standard quantified statement).

$$\text{RealSequence}(x_n) \iff x : \mathbb{N} \rightarrow \mathbb{R}.$$

Remark (Definition predicate reading).

$$\text{RealSequence}(x_n) := \text{Sequence}(x_n, \mathbb{R}).$$

Remark (Interpretation). A real sequence is not a new kind of indexing object; it is the specialization of a sequence to the ambient space $\mathbb{R}$. Thus the generic function structure is inherited from sequences, while order, absolute value, and arithmetic enter through the real codomain.

Remark (Dependencies).

- [Sequence](#)
- [Real Numbers](#)

Example (Constant Sequence). Let $c \in \mathbb{R}$. The **constant sequence** is defined by

$$x_n := c \quad \forall n \in \mathbb{N}.$$

The constant sequence converges to c : given any $\varepsilon > 0$, take $N = 1$, and observe $|x_n - c| = 0 < \varepsilon$ for all $n \geq 1$. It is the degenerate case of convergence in which the tail is not merely eventually close to L but identically equal to it from the first term.

Example (The Sequence $(1/n)$). The sequence (x_n) defined by

$$x_n := \frac{1}{n} \quad \forall n \in \mathbb{N}$$

converges to 0. Given $\varepsilon > 0$, the Archimedean property supplies $N \in \mathbb{N}$ with $N > 1/\varepsilon$, and then for all $n \geq N$,

$$\left| \frac{1}{n} - 0 \right| = \frac{1}{n} \leq \frac{1}{N} < \varepsilon.$$

This is the canonical example of a sequence that approaches its limit monotonically from above, never reaching it. It also serves as the standard witness in squeeze arguments and in the proof that $\inf\{1/n : n \in \mathbb{N}\} = 0$.

Example (The Sequence (n)). The sequence (x_n) defined by

$$x_n := n \quad \forall n \in \mathbb{N}$$

is divergent. For any proposed limit $L \in \mathbb{R}$, take $\varepsilon = 1$; the Archimedean property supplies $n \in \mathbb{N}$ with $n > L + 1$, so $|x_n - L| = |n - L| > 1 = \varepsilon$ for infinitely many n . The sequence is unbounded, and every convergent sequence is bounded, so divergence also follows from that theorem once it is established. This is the canonical example of divergence to $+\infty$: the terms are not oscillating outside a neighborhood of L but growing without bound in a single direction.

6.1.2 Null and Constant Sequences

Null and Constant Sequences

Definition (Constant Sequence)

Definition 6.1.3 (Constant Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **constant** exactly when there exists $c \in \mathbb{R}$ such that $x_n = c$ for every $n \in \mathbb{N}$.

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence.} \\ &(x_n) \text{ is constant} \iff \exists c \in \mathbb{R} \, \forall n \in \mathbb{N} \, (x_n = c). \end{aligned}$$

Remark (Definition predicate reading).

$$\text{ConstantSequence}(x_n, c) := \forall n \in \mathbb{N} \, (x_n = c).$$

Remark (Negated quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence.} \\ &(x_n) \text{ is not constant} \iff \forall c \in \mathbb{R} \, \exists n \in \mathbb{N} \, (x_n \neq c). \end{aligned}$$

Remark (Negation predicate reading).

$$\neg \exists c \in \mathbb{R} \, \text{ConstantSequence}(x_n, c).$$

Remark (Failure modes). Every proposed constant value fails at some index.

Remark (Failure mode decomposition).

$$\forall c \in \mathbb{R} \quad \underbrace{\exists n \in \mathbb{N} \, (x_n \neq c)}_{\text{a term differs from the proposed constant value}}.$$

Remark (Interpretation). A constant sequence has no genuine dependence on the index. The indexing is still part of the object, but every index is assigned the same real value.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Null Sequence)

Definition 6.1.4 (Null Sequence). Let (x_n) be a real sequence. The sequence (x_n) is a **null sequence** exactly when for every $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that $|x_n| < \varepsilon$ for every $n \geq N$.

Remark (Standard quantified statement).

Let (x_n) be a real sequence.

(x_n) is a null sequence $\iff \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n| < \varepsilon)$.

Remark (Definition predicate reading).

$\text{NullSequence}(x_n) := \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n| < \varepsilon)$.

Remark (Negated quantified statement).

Let (x_n) be a real sequence.

(x_n) is not a null sequence $\iff \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N (|x_n| \geq \varepsilon)$.

Remark (Negation predicate reading).

$\neg \text{NullSequence}(x_n)$.

Remark (Failure modes). Some positive tolerance is missed infinitely far out in the sequence.

Remark (Failure mode decomposition).

$$\exists \varepsilon > 0 \forall N \in \mathbb{N} \underbrace{\exists n \geq N (|x_n| \geq \varepsilon)}_{\text{the tail is not trapped inside the } \varepsilon\text{-band}}.$$

Remark (Interpretation). A null sequence is a sequence whose tail is eventually contained in every symmetric neighborhood of zero. The first finitely many terms are irrelevant; only the eventual behavior matters.

Remark (Dependencies).

- [Real Sequence](#)

Theorem (Constant Sequence Convergence)

Theorem 6.1.5 (Constant Sequence Convergence). Let (x_n) be a real sequence, and let $c \in \mathbb{R}$. If $x_n = c$ for every $n \in \mathbb{N}$, then (x_n) converges to c .

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence and $c \in \mathbb{R}$.

$(\forall n \in \mathbb{N} (x_n = c)) \implies \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - c| < \varepsilon)$.

Remark (Predicate reading).

$$\text{ConstantSequence}(x_n, c) \implies \text{ConvergentSequence}(x_n, c).$$

Remark (Interpretation). A constant sequence is already equal to its proposed limit at every index. Every positive tolerance is therefore satisfied by any tail of the sequence.

Remark (Dependencies).

- [Constant Sequence](#)
- [Convergent Sequence](#)

Theorem (Zero Sequence is Null)

Theorem 6.1.6 (Zero Sequence is Null). *Let (x_n) be a real sequence. If $x_n = 0$ for every $n \in \mathbb{N}$, then (x_n) is a null sequence.*

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence.

$$(\forall n \in \mathbb{N} (x_n = 0)) \implies \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n| < \varepsilon).$$

Remark (Predicate reading).

$$\text{ConstantSequence}(x_n, 0) \implies \text{NullSequence}(x_n).$$

Remark (Interpretation). The zero sequence satisfies every null-sequence tolerance immediately because the distance from each term to zero is exactly zero.

Remark (Dependencies).

- [Null Sequence](#)
- [Constant Sequence](#)

Theorem (Constant Null Sequence)

Theorem 6.1.7 (Constant Null Sequence). *Let (x_n) be a real sequence, and let $c \in \mathbb{R}$. If $x_n = c$ for every $n \in \mathbb{N}$, then (x_n) is a null sequence if and only if $c = 0$.*

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence and $c \in \mathbb{R}$.

$$(\forall n \in \mathbb{N} (x_n = c)) \implies \left([\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n| < \varepsilon)] \iff c = 0 \right).$$

Remark (Predicate reading).

$$\text{ConstantSequence}(x_n, c) \implies (\text{NullSequence}(x_n) \iff c = 0).$$

Remark (Interpretation). A sequence that never changes can approach zero only when its fixed value is already zero. Conversely, the fixed value zero gives the zero sequence.

Remark (Dependencies).

- [Constant Sequence](#)
- [Null Sequence](#)

Definition (Ultimately Constant Sequence)

Definition 6.1.8 (Ultimately Constant Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **ultimately constant** exactly when there exist $N \in \mathbb{N}$ and $c \in \mathbb{R}$ such that $x_n = c$ for every $n \geq N$.

Remark (Standard quantified statement).

Let (x_n) be a real sequence.

$$(x_n) \text{ is ultimately constant} \iff \exists N \in \mathbb{N} \exists c \in \mathbb{R} \forall n \geq N (x_n = c).$$

Remark (Definition predicate reading).

$$\text{UltimatelyConstantSequence}(x_n, c) := \exists N \in \mathbb{N} \forall n \geq N (x_n = c).$$

Remark (Negated quantified statement).

Let (x_n) be a real sequence.

$$(x_n) \text{ is not ultimately constant} \iff \forall N \in \mathbb{N} \forall c \in \mathbb{R} \exists n \geq N (x_n \neq c).$$

Remark (Negation predicate reading).

$$\neg \exists c \in \mathbb{R} \text{ UltimatelyConstantSequence}(x_n, c).$$

Remark (Failure modes). No tail of the sequence becomes permanently equal to a single real value.

Remark (Failure mode decomposition).

$$\forall N \in \mathbb{N} \forall c \in \mathbb{R} \underbrace{\exists n \geq N (x_n \neq c)}_{\text{the proposed tail value fails later in the sequence}}.$$

Remark (Interpretation). An ultimately constant sequence may vary on an initial finite segment, but its tail becomes fixed. The concept isolates eventual behavior from finite initial behavior.

Remark (Dependencies).

- [Real Sequence](#)

Theorem (Difference from the Limit is Null)

Theorem 6.1.9 (Difference from the Limit is Null). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. The sequence (x_n) converges to L if and only if the sequence $(x_n - L)$ is a null sequence.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence and } L \in \mathbb{R}. \\ &[\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - L| < \varepsilon)] \\ &\iff [\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - L| < \varepsilon)]. \end{aligned}$$

Remark (Predicate reading).

$$\text{ConvergentSequence}(x_n, L) \iff \text{NullSequence}(x_n - L).$$

Remark (Interpretation). Subtracting the proposed limit translates the convergence problem to the special case of convergence to zero. Null sequences are therefore the local normal form for sequence convergence.

Remark (Dependencies).

- [Convergent Sequence](#)
- [Null Sequence](#)
- [Pointwise Difference of Sequences](#)

Theorem (Ultimately Constant Sequence Convergence)

Theorem 6.1.10 (Ultimately Constant Sequence Convergence). *Let (x_n) be a real sequence. If there exist $N_0 \in \mathbb{N}$ and $c \in \mathbb{R}$ such that $x_n = c$ for every $n \geq N_0$, then (x_n) converges to c .*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence.} \\ &\forall c \in \mathbb{R} \left([\exists N_0 \in \mathbb{N} \forall n \geq N_0 (x_n = c)] \right. \\ &\quad \implies [\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - c| < \varepsilon)] \Big). \end{aligned}$$

Remark (Predicate reading).

$$\text{UltimatelyConstantSequence}(x_n, c) \implies \text{ConvergentSequence}(x_n, c).$$

Remark (Interpretation). Once the tail is equal to a fixed value, all sufficiently late terms are already exactly at the proposed limit.

Remark (Dependencies).

- [Ultimately Constant Sequence](#)
- [Convergent Sequence](#)

Theorem (Constant Implies Ultimately Constant)

Theorem 6.1.11 (Constant Implies Ultimately Constant). *Every constant real sequence is ultimately constant.*

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence.} \\ &[\exists c \in \mathbb{R} \forall n \in \mathbb{N} (x_n = c)] \\ &\implies [\exists N \in \mathbb{N} \exists c \in \mathbb{R} \forall n \geq N (x_n = c)]. \end{aligned}$$

Remark (Predicate reading).

$$\text{ConstantSequence}(x_n, c) \implies \text{UltimatelyConstantSequence}(x_n, c).$$

Remark (Interpretation). A constant sequence has already stabilized from the first index, so it is automatically stabilized on some tail.

Remark (Dependencies).

- [Constant Sequence](#)
- [Ultimately Constant Sequence](#)

Theorem (Ultimately Zero Sequence is Null)

Theorem 6.1.12 (Ultimately Zero Sequence is Null). *Let (x_n) be a real sequence. If there exists $N_0 \in \mathbb{N}$ such that $x_n = 0$ for every $n \geq N_0$, then (x_n) is a null sequence.*

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence.} \\ &[\exists N_0 \in \mathbb{N} \forall n \geq N_0 (x_n = 0)] \\ &\implies \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n| < \varepsilon). \end{aligned}$$

Remark (Predicate reading).

$$\text{UltimatelyConstantSequence}(x_n, 0) \implies \text{NullSequence}(x_n).$$

Remark (Interpretation). If the tail is identically zero, then every sufficiently late term lies inside every positive tolerance around zero.

Remark (Dependencies).

- [Ultimately Constant Sequence](#)
- [Null Sequence](#)

Theorem (Ultimately Constant Null Sequence)

Theorem 6.1.13 (Ultimately Constant Null Sequence). *Let (x_n) be a real sequence. Suppose there exist $N_0 \in \mathbb{N}$ and $c \in \mathbb{R}$ such that $x_n = c$ for every $n \geq N_0$. Then (x_n) is a null sequence if and only if $c = 0$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence.} \\ &\forall c \in \mathbb{R} \left(\left[\exists N_0 \in \mathbb{N} \forall n \geq N_0 (x_n = c) \right] \right. \\ &\quad \left. \implies \left(\left[\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n| < \varepsilon) \right] \iff c = 0 \right) \right). \end{aligned}$$

Remark (Predicate reading).

$$\text{UltimatelyConstantSequence}(x_n, c) \implies (\text{NullSequence}(x_n) \iff c = 0).$$

Remark (Interpretation). Only the eventual value controls whether an ultimately constant sequence is null. A finite initial segment cannot prevent or create null behavior.

Remark (Dependencies).

- [Ultimately Constant Sequence](#)
- [Null Sequence](#)

Theorem (Tail Equality Preserves Convergence)

Theorem 6.1.14 (Tail Equality Preserves Convergence). *Let (x_n) and (y_n) be real sequences. If there exists $N_0 \in \mathbb{N}$ such that $x_n = y_n$ for every $n \geq N_0$, then for every $L \in \mathbb{R}$, (x_n) converges to L if and only if (y_n) converges to L .*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ and } (y_n) \text{ be real sequences.} \\ &\left[\exists N_0 \in \mathbb{N} \forall n \geq N_0 (x_n = y_n) \right] \\ &\implies \forall L \in \mathbb{R} \left(\left[\forall \varepsilon > 0 \exists N_x \in \mathbb{N} \forall n \geq N_x (|x_n - L| < \varepsilon) \right] \right. \\ &\quad \left. \iff \left[\forall \varepsilon > 0 \exists N_y \in \mathbb{N} \forall n \geq N_y (|y_n - L| < \varepsilon) \right] \right). \end{aligned}$$

Remark (Predicate reading).

$\text{EventuallyEqual}(x_n, y_n) \implies \forall L \in \mathbb{R} \left(\text{ConvergentSequence}(x_n, L) \iff \text{ConvergentSequence}(y_n, L) \right).$ ■

Remark (Interpretation). Convergence is a tail property. Changing finitely many initial terms cannot change whether a sequence converges to a specified real number.

Remark (Dependencies).

- [Convergent Sequence](#)

Definition (Sequence Bounded Above)

Definition 6.1.15 (Sequence Bounded Above). Let (x_n) be a real sequence. The sequence (x_n) is **bounded above** exactly when there exists $M \in \mathbb{R}$ such that $x_n \leq M$ for every $n \in \mathbb{N}$.

Remark (Standard quantified statement).

Let (x_n) be a real sequence.
 (x_n) is bounded above $\iff \exists M \in \mathbb{R} \forall n \in \mathbb{N} (x_n \leq M)$.

Remark (Definition predicate reading).

$\text{BoundedAboveSequence}(x_n) := \exists M \in \mathbb{R} \forall n \in \mathbb{N} (x_n \leq M)$.

Remark (Negated quantified statement).

Let (x_n) be a real sequence.
 (x_n) is not bounded above $\iff \forall M \in \mathbb{R} \exists n \in \mathbb{N} (M < x_n)$.

Remark (Negation predicate reading).

$\neg \text{BoundedAboveSequence}(x_n)$.

Remark (Failure modes). Every proposed upper bound is exceeded by some term of the sequence.

Remark (Failure mode decomposition).

$$\forall M \in \mathbb{R} \underbrace{\exists n \in \mathbb{N} (M < x_n)}_{\text{a term rises above the proposed ceiling}}.$$

Remark (Interpretation). A bounded-above sequence has a finite ceiling. Failure means no real number serves as a ceiling for all terms.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Sequence Bounded Below)

Definition 6.1.16 (Sequence Bounded Below). Let (x_n) be a real sequence. The sequence (x_n) is **bounded below** exactly when there exists $m \in \mathbb{R}$ such that $m \leq x_n$ for every $n \in \mathbb{N}$.

Remark (Standard quantified statement).

Let (x_n) be a real sequence.

(x_n) is bounded below $\iff \exists m \in \mathbb{R} \forall n \in \mathbb{N} (m \leq x_n)$.

Remark (Definition predicate reading).

$\text{BoundedBelowSequence}(x_n) := \exists m \in \mathbb{R} \forall n \in \mathbb{N} (m \leq x_n)$.

Remark (Negated quantified statement).

Let (x_n) be a real sequence.

(x_n) is not bounded below $\iff \forall m \in \mathbb{R} \exists n \in \mathbb{N} (x_n < m)$.

Remark (Negation predicate reading).

$\neg \text{BoundedBelowSequence}(x_n)$.

Remark (Failure modes). Every proposed lower bound is undercut by some term of the sequence.

Remark (Failure mode decomposition).

$$\forall m \in \mathbb{R} \quad \underbrace{\exists n \in \mathbb{N} (x_n < m)}_{\text{a term falls below the proposed floor}} .$$

Remark (Interpretation). A bounded-below sequence has a finite floor. Failure means no real number serves as a floor for all terms.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Bounded Sequence)

Definition 6.1.17 (Bounded Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **bounded** exactly when there exists $M > 0$ such that $|x_n| \leq M$ for every $n \in \mathbb{N}$.

Remark (Standard quantified statement).

Let (x_n) be a real sequence.

(x_n) is bounded $\iff \exists M > 0 \forall n \in \mathbb{N} (|x_n| \leq M)$.

Remark (Definition predicate reading).

$$\text{BoundedSequence}(x_n) := \exists M > 0 \forall n \in \mathbb{N} (|x_n| \leq M).$$

Remark (Negated quantified statement).

Let (x_n) be a real sequence.

$$(x_n) \text{ is not bounded} \iff \forall M > 0 \exists n \in \mathbb{N} (|x_n| > M).$$

Remark (Negation predicate reading).

$$\neg \text{BoundedSequence}(x_n).$$

Remark (Failure modes). Every symmetric bound around zero is escaped by some term of the sequence.

Remark (Failure mode decomposition).

$$\forall M > 0 \quad \underbrace{\exists n \in \mathbb{N} (|x_n| > M)}_{\text{a term escapes the symmetric band of radius } M}.$$

Remark (Interpretation). A bounded sequence is trapped in a finite symmetric band around zero. This is equivalent to having both a finite ceiling and a finite floor.

Remark (Dependencies).

- [Real Sequence](#)

Theorem (Eventually Bounded Above Tail Formulation)

Theorem 6.1.18 (Eventually Bounded Above Tail Formulation). *Let (x_n) be a real sequence. The sequence (x_n) is bounded above if and only if there exist $N_0 \in \mathbb{N}$ and $M \in \mathbb{R}$ such that $x_n \leq M$ for every $n \geq N_0$.*

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence.

$$\begin{aligned} & [\exists M \in \mathbb{R} \forall n \in \mathbb{N} (x_n \leq M)] \\ & \iff [\exists N_0 \in \mathbb{N} \exists M \in \mathbb{R} \forall n \geq N_0 (x_n \leq M)]. \end{aligned}$$

Remark (Predicate reading).

$$\text{BoundedAboveSequence}(x_n) \iff \exists M \in \mathbb{R} \text{ Eventually}(x_n, x_n \leq M).$$

Remark (Interpretation). One-sided boundedness is unchanged by ignoring finitely many initial terms, because finitely many exceptional values can be absorbed into a larger upper bound.

Remark (Dependencies).

- [Sequence Bounded Above](#)

Theorem (Eventually Bounded Below Tail Formulation)

Theorem 6.1.19 (Eventually Bounded Below Tail Formulation). *Let (x_n) be a real sequence. The sequence (x_n) is bounded below if and only if there exist $N_0 \in \mathbb{N}$ and $m \in \mathbb{R}$ such that $m \leq x_n$ for every $n \geq N_0$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence.} \\ &[\exists m \in \mathbb{R} \forall n \in \mathbb{N} (m \leq x_n)] \\ &\iff [\exists N_0 \in \mathbb{N} \exists m \in \mathbb{R} \forall n \geq N_0 (m \leq x_n)]. \end{aligned}$$

Remark (Predicate reading).

$$\text{BoundedBelowSequence}(x_n) \iff \exists m \in \mathbb{R} \text{ Eventually}(x_n, m \leq x_n).$$

Remark (Interpretation). One-sided lower boundedness is unchanged by ignoring finitely many initial terms, because finitely many exceptional values can be absorbed into a lower bound.

Remark (Dependencies).

- [Sequence Bounded Below](#)

Theorem (Eventually Bounded Tail Formulation)

Theorem 6.1.20 (Eventually Bounded Tail Formulation). *Let (x_n) be a real sequence. The sequence (x_n) is bounded if and only if there exist $N_0 \in \mathbb{N}$ and $M > 0$ such that $|x_n| \leq M$ for every $n \geq N_0$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence.} \\ &[\exists M > 0 \forall n \in \mathbb{N} (|x_n| \leq M)] \\ &\iff [\exists N_0 \in \mathbb{N} \exists M > 0 \forall n \geq N_0 (|x_n| \leq M)]. \end{aligned}$$

Remark (Predicate reading).

$$\text{BoundedSequence}(x_n) \iff \exists M > 0 \text{ Eventually}(x_n, |x_n| \leq M).$$

Remark (Interpretation). Boundedness is a tail property up to finitely many exceptions. A finite initial segment can be enclosed by increasing the bound.

Remark (Dependencies).

- [Bounded Sequence](#)

Theorem (Bounded Sequence If and Only If Bounded Above and Below)

Theorem 6.1.21 (Bounded Sequence If and Only If Bounded Above and Below). *Let (x_n) be a real sequence. The sequence (x_n) is bounded if and only if it is both bounded above and bounded below.*

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence.

$$\begin{aligned} & [\exists K > 0 \forall n \in \mathbb{N} (|x_n| \leq K)] \\ & \iff [\exists m \in \mathbb{R} \exists M \in \mathbb{R} \forall n \in \mathbb{N} (m \leq x_n \wedge x_n \leq M)]. \end{aligned}$$

Remark (Predicate reading).

$$\text{BoundedSequence}(x_n) \iff (\text{BoundedBelowSequence}(x_n) \wedge \text{BoundedAboveSequence}(x_n)).$$

Remark (Interpretation). An absolute-value bound gives a symmetric interval around zero. Having both a floor and a ceiling gives an interval that may not be symmetric, but it can be enlarged to a symmetric one.

Remark (Dependencies).

- [Bounded Sequence](#)
- [Sequence Bounded Above](#)
- [Sequence Bounded Below](#)

Theorem (Absolute Bound Gives Upper and Lower Bounds)

Theorem 6.1.22 (Absolute Bound Gives Upper and Lower Bounds). *Let (x_n) be a real sequence, and let $K > 0$. If $|x_n| \leq K$ for every $n \in \mathbb{N}$, then $-K \leq x_n \leq K$ for every $n \in \mathbb{N}$.*

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence.

$$\begin{aligned} & \forall K > 0 [\forall n \in \mathbb{N} (|x_n| \leq K)] \\ & \implies \forall n \in \mathbb{N} (-K \leq x_n \wedge x_n \leq K). \end{aligned}$$

Remark (Interpretation). An absolute-value inequality simultaneously controls the positive and negative directions. The upper barrier is K , and the lower barrier is $-K$.

Remark (Dependencies).

- [Bounded Sequence](#)
- [Sequence Bounded Above](#)
- [Sequence Bounded Below](#)

Theorem (Upper and Lower Bounds Give an Absolute Bound)

Theorem 6.1.23 (Upper and Lower Bounds Give an Absolute Bound). *Let (x_n) be a real sequence. If there exist $m, M \in \mathbb{R}$ such that $m \leq x_n \leq M$ for every $n \in \mathbb{N}$, then there exists $K > 0$ such that $|x_n| \leq K$ for every $n \in \mathbb{N}$.*

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence.

$$\begin{aligned} & [\exists m \in \mathbb{R} \exists M \in \mathbb{R} \forall n \in \mathbb{N} (m \leq x_n \wedge x_n \leq M)] \\ & \implies [\exists K > 0 \forall n \in \mathbb{N} (|x_n| \leq K)]. \end{aligned}$$

Remark (Interpretation). Two one-sided barriers can always be replaced by one symmetric barrier around zero. Any positive K at least as large as both $|m|$ and $|M|$ works.

Remark (Dependencies).

- [Sequence Bounded Above](#)
- [Sequence Bounded Below](#)
- [Bounded Sequence](#)

6.2 Examples and Counterexamples

Sequence Examples — Quick Reference

Example	Role
Constant sequence	Basic convergent model
Reciprocal sequence	Canonical null sequence
Alternating null sequence	Sign oscillation with convergence to 0
Oscillating sequence	Bounded nonconvergent model
Geometric sequence	Ratio-driven convergence and divergence
Bounded not convergent	Boundedness alone is insufficient
Small differences not Cauchy	Vanishing increments do not ensure Cauchy

Examples and Counterexamples

Examples provide test objects for the definitions. A good sequence example usually isolates one behavior: settling to a limit, oscillating between several values, escaping to infinity, or satisfying a tempting condition that is still too weak to force convergence.

Example (Constant Sequence). Let $c \in \mathbb{R}$ and define $x_n = c$ for every $n \in \mathbb{N}$. Then (x_n) converges to c .

Remark (Interpretation). A constant sequence is the baseline example of convergence: every term is already equal to the proposed limit.

Example (Reciprocal Sequence). Define $x_n = 1/n$ for every $n \in \mathbb{N}$. Then (x_n) is decreasing, bounded below by 0, and converges to 0.

Remark (Interpretation). The reciprocal sequence is the standard positive null sequence. It is also the model for estimates of the form “eventually smaller than any fixed positive tolerance.”

Example (Alternating Null Sequence). Define

$$x_n = \frac{(-1)^n}{n} \quad \text{for every } n \in \mathbb{N}.$$

Then (x_n) is not monotone, but it converges to 0.

Remark (Interpretation). Oscillation in sign does not by itself prevent convergence. The amplitudes must be watched: here the oscillation collapses to zero.

Example (Oscillating Sequence). Define

$$x_n = (-1)^n \quad \text{for every } n \in \mathbb{N}.$$

Then (x_n) is bounded and divergent. Its even-indexed subsequence converges to 1, and its odd-indexed subsequence converges to -1 .

Remark (Interpretation). This is the canonical oscillator. The sequence never settles because two subsequences settle to different limits.

Example (Geometric Sequence). Let $r \in \mathbb{R}$ and define $x_n = r^n$ for every $n \in \mathbb{N}$. Then:

- if $|r| < 1$, then $x_n \rightarrow 0$;
- if $r = 1$, then $x_n \rightarrow 1$;
- if $r = -1$, then (x_n) oscillates and diverges;
- if $|r| > 1$, then (x_n) is unbounded.

Remark (Interpretation). The geometric sequence is the standard laboratory for rate, sign, and magnitude. The threshold $|r| = 1$ separates decay from growth.

Example (Bounded Does Not Imply Convergent). The sequence $x_n = (-1)^n$ is bounded, since $|x_n| \leq 1$ for every $n \in \mathbb{N}$, but it does not converge.

Remark (Interpretation). Boundedness prevents escape to infinity, but it does not prevent persistent oscillation.

Example (Vanishing Successive Differences Do Not Imply Cauchy). Let

$$x_n = 1 + \frac{1}{2} + \cdots + \frac{1}{n} \quad \text{for every } n \in \mathbb{N}.$$

Then

$$x_{n+1} - x_n = \frac{1}{n+1} \rightarrow 0,$$

but (x_n) is not Cauchy.

Remark (Interpretation). Cauchy control requires all sufficiently late terms to be mutually close. Controlling only adjacent jumps is weaker.

6.3 Convergence

Definition (Convergence of a Sequence)

Definition 6.3.1 (Convergence of a Sequence). Let $(x_n) \subseteq \mathbb{R}$. The sequence (x_n) **converges** to $L \in \mathbb{R}$ if

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - L| < \varepsilon).$$

The value L is called the **limit** of (x_n) , written $x_n \rightarrow L$ or $\lim_{n \rightarrow \infty} x_n = L$. A sequence that converges to some $L \in \mathbb{R}$ is called **convergent**; otherwise it is called **divergent**.

Remark (Standard quantified statement).

$$\begin{aligned} & \forall (x_n) \subseteq \mathbb{R} \forall L \in \mathbb{R} \\ & \left(x_n \rightarrow L \iff \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - L| < \varepsilon) \right). \end{aligned}$$

Remark (Definition predicate reading).

$$\text{ConvergentSequence}(x_n, L) := \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N \text{OutputTolerance}(x_n, L, \varepsilon).$$

Remark (Negated quantified statement).

$$\begin{aligned} x_n \not\rightarrow L & \iff \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N \\ & (|x_n - L| \geq \varepsilon). \end{aligned}$$

Remark (Negation predicate reading).

$$\text{NotConvergentTo}(x_n, L) := \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N (|x_n - L| \geq \varepsilon).$$

Remark (Failure modes). Failure of convergence to a fixed L means that some positive tolerance cannot be controlled on any tail. The escaped terms may oscillate, drift monotonically away, or grow without bound, but all such behavior is captured by the same persistent-tolerance obstruction.

Remark (Failure mode decomposition). The negation decomposes into a single structural obstruction:

$$\underbrace{\exists \varepsilon > 0}_{\text{a fixed tolerance}} \quad \underbrace{\forall N \in \mathbb{N}}_{\text{no tail suffices}} \quad \underbrace{\exists n \geq N}_{\text{terms escape infinitely often}} \quad \underbrace{|x_n - L| \geq \varepsilon}_{\text{outside the target interval}}.$$

The sequence fails to converge to L not by drifting away once, but by returning outside the ε -interval infinitely often regardless of how far along the tail one looks.

Remark (Interpretation). The quantifier order $\forall \varepsilon \exists N \forall n \geq N$ encodes a strict dependency: N is permitted to depend on ε , but not on n . This asymmetry is load-bearing. Reversing the outer quantifiers to $\exists N \forall \varepsilon$ would assert that a single index N controls all tolerances simultaneously, which is a strictly stronger and generally false condition. The condition $|x_n - L| < \varepsilon$ is the membership assertion $x_n \in (L - \varepsilon, L + \varepsilon)$; convergence is therefore the statement that the tail of the sequence is eventually contained in every ε -neighborhood of L .

Remark (Dependencies).

- [Real Sequence](#)

Definition (Convergence — Neighborhood Formulation)

Definition 6.3.2 (Convergence — Neighborhood Formulation). Let $(x_n) \subseteq \mathbb{R}$ and let $L \in \mathbb{R}$. The sequence (x_n) **converges** to L if for every $\varepsilon > 0$ the ε -neighborhood

$$V_\varepsilon(L) := (L - \varepsilon, L + \varepsilon)$$

contains all but finitely many terms of (x_n) ; that is,

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (x_n \in V_\varepsilon(L)).$$

Remark (Standard quantified statement).

$$\begin{aligned} & \forall (x_n) \subseteq \mathbb{R} \forall L \in \mathbb{R} \\ & \left(x_n \rightarrow L \iff \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (x_n \in V_\varepsilon(L)) \right). \end{aligned}$$

Remark (Definition predicate reading).

$$\text{ConvergentSequence}(x_n, L) := \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (x_n \in V_\varepsilon(L)).$$

Remark (Negated quantified statement).

$$\begin{aligned} x_n \not\rightarrow L & \iff \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N \\ & (x_n \notin V_\varepsilon(L)). \end{aligned}$$

Remark (Negation predicate reading).

$$\text{NotConvergentTo}(x_n, L) := \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N (x_n \notin V_\varepsilon(L)).$$

Remark (Failure modes). Failure of the neighborhood formulation is exactly failure to trap a tail in every neighborhood of L . Different examples may escape by oscillation, by monotone drift, or by unbounded growth, but the formal failure is always the existence of one neighborhood that every tail misses at least once.

Remark (Failure mode decomposition).

$$\underbrace{\exists \varepsilon > 0}_{\text{a fixed neighborhood}} \quad \underbrace{\forall N \in \mathbb{N}}_{\text{no tail suffices}} \quad \underbrace{\exists n \geq N}_{\text{terms escape infinitely often}} \quad \underbrace{x_n \notin V_\varepsilon(L)}_{\text{outside every candidate neighborhood}} .$$

The sequence fails to converge to L precisely when some neighborhood of L contains only finitely many terms: infinitely many terms lie permanently outside it.

Remark (Interpretation). The neighborhood formulation reframes convergence as a set-containment condition rather than an inequality condition. The assertion $x_n \in V_\varepsilon(L)$ is identical to $|x_n - L| < \varepsilon$; the two definitions are logically equivalent and the choice between them is one of language, not content. The neighborhood language is the natural bridge to the topological formulation: in a general topological space, open sets replace ε -neighborhoods and the definition reads identically with $V_\varepsilon(L)$ replaced by any open set U containing L . The phrase *all but finitely many terms* is the canonical prose compression of the quantifier block $\exists N \in \mathbb{N} \forall n \geq N$, and is standard in the literature.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [\$\varepsilon\$ -neighbourhood](#)

Theorem (Equivalence of Convergence Formulations)

Theorem 6.3.3 (Equivalence of Convergence Formulations). *Let $(x_n) \subseteq \mathbb{R}$ and let $L \in \mathbb{R}$. The following statements are equivalent.*

- (a) $x_n \rightarrow L$.
- (b) $\forall \varepsilon > 0 \exists K \in \mathbb{N} \forall n \geq K (|x_n - L| < \varepsilon)$.
- (c) $\forall \varepsilon > 0 \exists K \in \mathbb{N} \forall n \geq K (L - \varepsilon < x_n < L + \varepsilon)$.
- (d) $\forall \varepsilon > 0 \exists K \in \mathbb{N} \forall n \geq K (x_n \in V_\varepsilon(L))$.

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \forall L \in \mathbb{R} \left((a) \iff (b) \iff (c) \iff (d) \right).$$

Remark (Theorem predicate reading).

$\text{EquivalentConvergenceForms}(x_n, L) := \text{ConvergentSequence}(x_n, L) \iff \text{NeighborhoodConvergentSequence}(x_n, L)$

Remark (Interpretation). The four conditions are the same geometric fact written in four notational registers. Condition (a) is the informal assertion. Condition (b) is the canonical ε - K definition: the absolute value formulation. Condition (c) makes the interval structure explicit by expanding $|x_n - L| < \varepsilon$ into its equivalent double inequality $L - \varepsilon < x_n < L + \varepsilon$; this is useful in proofs where upper and lower bounds are handled separately. Condition (d) is the neighborhood formulation: $x_n \in V_\varepsilon(L)$ is identical to conditions (b) and (c) but frames convergence as set membership, which is the natural language for the topological generalization. The index K plays the same role as N in the base definition [Definition 6.3.1](#); the letter changes here to signal that K is the tail offset in the sense of [Definition 6.4.1](#).

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Convergence — Neighborhood Formulation](#)

6.3.1 Limits

Theorem (Uniqueness of Limits)

Theorem 6.3.4 (Uniqueness of Limits). *Let $(x_n) \subseteq \mathbb{R}$. If $x_n \rightarrow L$ and $x_n \rightarrow K$, then $L = K$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \, \forall L \in \mathbb{R} \, \forall K \in \mathbb{R} \\ \left((x_n \rightarrow L \wedge x_n \rightarrow K) \implies L = K \right).$$

Remark (Theorem predicate reading).

$$\text{UniqueSequentialLimit}(x_n, L, K) := (\text{ConvergentSequence}(x_n, L) \wedge \text{ConvergentSequence}(x_n, K)) \implies L = K.$$

Remark (Negated quantified statement).

$$\exists (x_n) \subseteq \mathbb{R} \, \exists L \in \mathbb{R} \, \exists K \in \mathbb{R} \\ \left((x_n \rightarrow L \wedge x_n \rightarrow K) \wedge L \neq K \right).$$

Remark (Negation predicate reading).

$$\neg \text{UniqueSequentialLimit}(x_n, L, K) := \text{ConvergentSequence}(x_n, L) \wedge \text{ConvergentSequence}(x_n, K) \wedge L \neq K.$$

Remark (Failure modes). The only failure mode is the existence of one real sequence with two distinct real limits. In $\mathbb{R}$ this is impossible because distinct real numbers can be separated by disjoint neighborhoods.

Remark (Failure mode decomposition).

$$\underbrace{\exists(x_n) \subseteq \mathbb{R}}_{\text{some sequence}} \underbrace{x_n \rightarrow L \wedge x_n \rightarrow K}_{\text{converges to two values}} \underbrace{L \neq K}_{\text{distinct limits}}.$$

The theorem asserts this configuration is impossible in $\mathbb{R}$. The failure mode is not merely exotic: it occurs in non-Hausdorff topological spaces where distinct points cannot be separated by disjoint open sets. The proof that it cannot occur in $\mathbb{R}$ is precisely an argument that $\mathbb{R}$ has enough room to separate any two distinct points by disjoint ε -neighborhoods.

Remark (Interpretation). The theorem licenses the notation $\lim_{n \rightarrow \infty} x_n = L$: the definite article is justified precisely because the limit, when it exists, is unique. Without uniqueness the limit would be a relation rather than a function, and the notation would be ambiguous. The proof strategy is a separation argument: if $L \neq K$ then $|L - K| > 0$, and choosing $\varepsilon = |L - K|/2$ produces two disjoint neighborhoods that the tail of (x_n) cannot simultaneously occupy. The Hausdorff remark in the failure mode block is the forward pointer: when sequences are studied in general topological spaces, uniqueness of limits is equivalent to the Hausdorff separation axiom.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Triangle inequality](#)

Theorem (Limit Preserves Eventual Order)

Theorem 6.3.5 (Limit Preserves Eventual Order). *Let (x_n) and (y_n) be real sequences, and let $L, M \in \mathbb{R}$. Suppose that $x_n \rightarrow L$ and $y_n \rightarrow M$. If there exists $N_0 \in \mathbb{N}$ such that $x_n \leq y_n$ for every $n \geq N_0$, then $L \leq M$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} & \forall (x_n), (y_n) \subseteq \mathbb{R} \, \forall L, M \in \mathbb{R} \\ & \left(x_n \rightarrow L \wedge y_n \rightarrow M \wedge \exists N_0 \in \mathbb{N} \, \forall n \geq N_0 \, (x_n \leq y_n) \right) \\ & \implies L \leq M. \end{aligned}$$

Remark (Theorem predicate reading).

$$(\text{ConvergentSequence}(x_n, L) \wedge \text{ConvergentSequence}(y_n, M) \wedge \text{Eventually}(x_n, x_n \leq y_n)) \implies L \leq M. \blacksquare$$

Remark (Interpretation). An inequality that holds on a tail survives passage to the limit, but only as a weak inequality. Strict inequalities on the terms may collapse at the limit, as in $0 < x_n$ with $x_n \rightarrow 0$.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Uniqueness of Limits](#)

Theorem (Strict Limit Separation Gives Eventual Order)

Theorem 6.3.6 (Strict Limit Separation Gives Eventual Order). *Let (x_n) and (y_n) be real sequences, and let $A, B \in \mathbb{R}$. Suppose that $x_n \rightarrow A$ and $y_n \rightarrow B$. If $A < B$, then there exists $N \in \mathbb{N}$ such that $x_n < y_n$ for every $n \geq N$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \forall (x_n), (y_n) \subseteq \mathbb{R} \forall A, B \in \mathbb{R} \\ \left(x_n \rightarrow A \wedge y_n \rightarrow B \wedge A < B \right) \\ \implies \exists N \in \mathbb{N} \forall n \geq N (x_n < y_n). \end{aligned}$$

Remark (Theorem predicate reading).

$$(\text{ConvergentSequence}(x_n, A) \wedge \text{ConvergentSequence}(y_n, B) \wedge A < B) \implies \text{Eventually}(x_n, x_n < y_n). \blacksquare$$

Remark (Interpretation). If the limiting values have a positive gap, then sufficiently late terms inherit the same order. The proof uses the open interval between A and B : eventually (x_n) lies near A and (y_n) lies near B , with room left between the two tails.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Uniqueness of Limits](#)

Theorem (Eventual Strict Comparison Preserves Weak Limit Order)

Theorem 6.3.7 (Eventual Strict Comparison Preserves Weak Limit Order). *Let (x_n) and (y_n) be real sequences, and let $A, B \in \mathbb{R}$. Suppose that $x_n \rightarrow A$ and $y_n \rightarrow B$.*

(a) *If there exists $N \in \mathbb{N}$ such that $x_n < y_n$ for every $n \geq N$, then $A \leq B$.*

(b) *If there exists $N \in \mathbb{N}$ such that $x_n > y_n$ for every $n \geq N$, then $A \geq B$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \forall (x_n), (y_n) \subseteq \mathbb{R} \forall A, B \in \mathbb{R} \\ \left(x_n \rightarrow A \wedge y_n \rightarrow B \wedge \exists N \in \mathbb{N} \forall n \geq N (x_n < y_n) \right) \implies A \leq B, \\ \left(x_n \rightarrow A \wedge y_n \rightarrow B \wedge \exists N \in \mathbb{N} \forall n \geq N (x_n > y_n) \right) \implies A \geq B. \end{aligned}$$

Remark (Theorem predicate reading).

$$\begin{aligned} &(\text{ConvergentSequence}(x_n, A) \wedge \text{ConvergentSequence}(y_n, B) \wedge \text{Eventually}(x_n, x_n < y_n)) \implies A \leq B, \\ &(\text{ConvergentSequence}(x_n, A) \wedge \text{ConvergentSequence}(y_n, B) \wedge \text{Eventually}(x_n, x_n > y_n)) \implies A \geq B. \end{aligned}$$

Remark (Interpretation). Strict comparison of terms does not force strict comparison of limits. The strict inequality can collapse in the limit, so the conclusion is weak: for example $1/n > 0$ for every $n \in \mathbb{N}$, but both sides have limit 0.

Remark (Dependencies).

- [Limit Preserves Eventual Order](#)

Theorem (Constant Comparison for Sequence Limits)

Theorem 6.3.8 (Constant Comparison for Sequence Limits). *Let (x_n) be a real sequence, and let $A, B \in \mathbb{R}$. Suppose that $x_n \rightarrow A$.*

- (a) *If there exists $N \in \mathbb{N}$ such that $x_n \leq B$ for every $n \geq N$, then $A \leq B$.*
- (b) *If there exists $N \in \mathbb{N}$ such that $x_n < B$ for every $n \geq N$, then $A \leq B$.*
- (c) *If there exists $N \in \mathbb{N}$ such that $x_n \geq B$ for every $n \geq N$, then $A \geq B$.*
- (d) *If there exists $N \in \mathbb{N}$ such that $x_n > B$ for every $n \geq N$, then $A \geq B$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} &\forall (x_n) \subseteq \mathbb{R} \forall A, B \in \mathbb{R} \\ &\left(x_n \rightarrow A \wedge \exists N \in \mathbb{N} \forall n \geq N (x_n \leq B) \right) \implies A \leq B, \\ &\left(x_n \rightarrow A \wedge \exists N \in \mathbb{N} \forall n \geq N (x_n < B) \right) \implies A \leq B, \\ &\left(x_n \rightarrow A \wedge \exists N \in \mathbb{N} \forall n \geq N (x_n \geq B) \right) \implies A \geq B, \\ &\left(x_n \rightarrow A \wedge \exists N \in \mathbb{N} \forall n \geq N (x_n > B) \right) \implies A \geq B. \end{aligned}$$

Remark (Theorem predicate reading).

$$\begin{aligned} &\text{ConvergentSequence}(x_n, A) \wedge \text{Eventually}(x_n, x_n \leq B) \implies A \leq B, \\ &\text{ConvergentSequence}(x_n, A) \wedge \text{Eventually}(x_n, x_n < B) \implies A \leq B, \\ &\text{ConvergentSequence}(x_n, A) \wedge \text{Eventually}(x_n, x_n \geq B) \implies A \geq B, \\ &\text{ConvergentSequence}(x_n, A) \wedge \text{Eventually}(x_n, x_n > B) \implies A \geq B. \end{aligned}$$

Remark (Interpretation). This is the constant version of the comparison theorem. A fixed real number B may be treated as the constant sequence with value B , so each conclusion is a direct application of eventual order preservation. Again, strict term inequalities produce only weak inequalities between limits.

Remark (Dependencies).

- [Constant Sequence](#)
- [Limit Preserves Eventual Order](#)
- [Eventual Strict Comparison Preserves Weak Limit Order](#)

Theorem (Constant Squeeze Theorem)

Theorem 6.3.9 (Constant Squeeze Theorem). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. If there exists $N_0 \in \mathbb{N}$ such that $L \leq x_n \leq L$ for every $n \geq N_0$, then $x_n \rightarrow L$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \forall (x_n) \subseteq \mathbb{R} \forall L \in \mathbb{R} \\ \left(\exists N_0 \in \mathbb{N} \forall n \geq N_0 (L \leq x_n \leq L) \right) \\ \implies \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - L| < \varepsilon). \end{aligned}$$

Remark (Theorem predicate reading).

$$\text{Eventually}(x_n, L \leq x_n \leq L) \implies \text{ConvergentSequence}(x_n, L).$$

Remark (Interpretation). This is the degenerate squeeze theorem: the lower and upper barriers are the same constant sequence. It says that eventual equality to a real number is enough for convergence to that real number.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Constant Sequence](#)

Theorem (Sequence Squeeze Theorem)

Theorem 6.3.10 (Sequence Squeeze Theorem). *Let (a_n) , (x_n) , and (b_n) be real sequences, and let $L \in \mathbb{R}$. Suppose that $a_n \rightarrow L$ and $b_n \rightarrow L$. If there exists $N_0 \in \mathbb{N}$ such that $a_n \leq x_n \leq b_n$ for every $n \geq N_0$, then $x_n \rightarrow L$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \forall (a_n), (x_n), (b_n) \subseteq \mathbb{R} \forall L \in \mathbb{R} \\ \left(a_n \rightarrow L \wedge b_n \rightarrow L \wedge \exists N_0 \in \mathbb{N} \forall n \geq N_0 (a_n \leq x_n \leq b_n) \right) \\ \implies \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - L| < \varepsilon). \end{aligned}$$

Remark (Theorem predicate reading).

$(\text{ConvergentSequence}(a_n, L) \wedge \text{ConvergentSequence}(b_n, L) \wedge \text{Eventually}(x_n, a_n \leq x_n \leq b_n)) \implies \text{ConvergentSequence}(x_n, L)$

Remark (Interpretation). The middle sequence is trapped between two tails that approach the same number. Once both barriers lie inside an ε -neighborhood of L , the trapped sequence must also lie inside that neighborhood.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Limit Preserves Eventual Order](#)

Theorem (Absolute-Value Squeeze Theorem)

Theorem 6.3.11 (Absolute-Value Squeeze Theorem). *Let (x_n) and (u_n) be real sequences, and let $L \in \mathbb{R}$. Suppose that $u_n \rightarrow 0$. If there exists $N_0 \in \mathbb{N}$ such that $|x_n - L| \leq u_n$ for every $n \geq N_0$, then $x_n \rightarrow L$.*

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned} \forall (x_n), (u_n) \subseteq \mathbb{R} \forall L \in \mathbb{R} \\ \left(u_n \rightarrow 0 \wedge \exists N_0 \in \mathbb{N} \forall n \geq N_0 (|x_n - L| \leq u_n) \right) \\ \implies \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - L| < \varepsilon). \end{aligned}$$

Remark (Theorem predicate reading).

$(\text{ConvergentSequence}(u_n, 0) \wedge \text{Eventually}(x_n, |x_n - L| \leq u_n)) \implies \text{ConvergentSequence}(x_n, L).$ ■

Remark (Interpretation). This is the estimate form of the squeeze theorem. Instead of constructing two explicit bounding sequences, it bounds the error $|x_n - L|$ by a null sequence.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Null Sequence](#)
- [Sequence Squeeze Theorem](#)

6.3.2 Algebra of Limits

Algebra of Convergent Sequences

The algebra of convergent sequences records that limits are compatible with the ordinary algebraic operations on real numbers. The pointwise operations below turn real sequences

into new real sequences, and the theorem blocks that follow state how convergence passes through those operations. The reciprocal and quotient laws require nonzero denominator hypotheses, while the square-root law requires nonnegative terms.

Definition (Pointwise Sum of Sequences)

Definition 6.3.12 (Pointwise Sum of Sequences). Let (x_n) and (y_n) be real sequences. The **pointwise sum** of (x_n) and (y_n) is the real sequence $(x_n + y_n)$ defined by

$$(x + y)_n := x_n + y_n \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

Let (x_n) and (y_n) be real sequences.

$$\forall n \in \mathbb{N} \ ((x + y)_n = x_n + y_n).$$

Remark (Interpretation). The sum sequence is formed term by term. No limiting process is involved in the definition; convergence enters only when the behavior of the resulting sequence is studied.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Pointwise Difference of Sequences)

Definition 6.3.13 (Pointwise Difference of Sequences). Let (x_n) and (y_n) be real sequences. The **pointwise difference** of (x_n) and (y_n) is the real sequence $(x_n - y_n)$ defined by

$$(x - y)_n := x_n - y_n \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

Let (x_n) and (y_n) be real sequences.

$$\forall n \in \mathbb{N} \ ((x - y)_n = x_n - y_n).$$

Remark (Interpretation). The difference sequence subtracts corresponding terms. It is equivalently the sum of (x_n) with the pointwise negation of (y_n) .

Remark (Dependencies).

- [Real Sequence](#)
- [Pointwise Sum of Sequences](#)

Definition (Scalar Multiple of a Sequence)

Definition 6.3.14 (Scalar Multiple of a Sequence). Let (x_n) be a real sequence, and let $\alpha \in \mathbb{R}$. The **scalar multiple** $\alpha(x_n)$ is the real sequence (αx_n) defined by

$$(\alpha x)_n := \alpha x_n \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

Let (x_n) be a real sequence and $\alpha \in \mathbb{R}$.
 $\forall n \in \mathbb{N} \ ((\alpha x)_n = \alpha x_n)$.

Remark (Interpretation). Scalar multiplication rescales every term by the same real number. It changes the vertical scale of the sequence but not its indexing.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Linear Combination of Real Sequences)

Definition 6.3.15 (Linear Combination of Real Sequences). Let (x_n) and (y_n) be real sequences, and let $\alpha, \beta \in \mathbb{R}$. The real linear combination $\alpha(x_n) + \beta(y_n)$ is the real sequence $(\alpha x_n + \beta y_n)$ defined by

$$(\alpha x + \beta y)_n := \alpha x_n + \beta y_n \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

Let $(x_n), (y_n)$ be real sequences and $\alpha, \beta \in \mathbb{R}$.
 $\forall n \in \mathbb{N} \ ((\alpha x + \beta y)_n = \alpha x_n + \beta y_n)$.

Remark (Interpretation). A linear combination of sequences is built term by term from scalar multiples and sums. It is the common algebraic shape behind sequence limit rules and Cauchy closure under addition and scalar multiplication.

Remark (Dependencies).

- [Scalar Multiple of a Sequence](#)
- [Pointwise Sum of Sequences](#)

Definition (Pointwise Negation of a Sequence)

Definition 6.3.16 (Pointwise Negation of a Sequence). Let (x_n) be a real sequence. The **pointwise negation** of (x_n) is the real sequence $(-x_n)$ defined by

$$(-x)_n := -x_n \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

Let (x_n) be a real sequence.
 $\forall n \in \mathbb{N} \ ((-x)_n = -x_n)$.

Remark (Interpretation). Pointwise negation reflects the sequence across zero. It is the special scalar multiple corresponding to the scalar -1 .

Remark (Dependencies).

- [Scalar Multiple of a Sequence](#)

Definition (Pointwise Product of Sequences)

Definition 6.3.17 (Pointwise Product of Sequences). Let (x_n) and (y_n) be real sequences. The **pointwise product** of (x_n) and (y_n) is the real sequence $(x_n y_n)$ defined by

$$(xy)_n := x_n y_n \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

Let (x_n) and (y_n) be real sequences.

$$\forall n \in \mathbb{N} \left((xy)_n = x_n y_n \right).$$

Remark (Interpretation). The product sequence multiplies corresponding terms. The product law for limits is subtler than the sum law because one factor must be controlled while the other is approximated.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Reciprocal Sequence)

Definition 6.3.18 (Reciprocal Sequence). Let (x_n) be a real sequence such that $x_n \neq 0$ for every $n \in \mathbb{N}$. The **reciprocal sequence** of (x_n) is the real sequence $(1/x_n)$ defined by

$$\left(\frac{1}{x} \right)_n := \frac{1}{x_n} \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

Let (x_n) be a real sequence with $\forall n \in \mathbb{N} (x_n \neq 0)$.

$$\forall n \in \mathbb{N} \left(\left(\frac{1}{x} \right)_n = \frac{1}{x_n} \right).$$

Remark (Interpretation). The reciprocal sequence is defined only when division by each term is legal. For limit theorems, a nonzero limit guarantees eventual nonzero behavior, but the sequence-level construction itself requires nonzero terms wherever it is defined.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Pointwise Quotient of Sequences)

Definition 6.3.19 (Pointwise Quotient of Sequences). Let (x_n) and (y_n) be real sequences such that $y_n \neq 0$ for every $n \in \mathbb{N}$. The **pointwise quotient** of (x_n) by (y_n) is the real sequence (x_n/y_n) defined by

$$\left(\frac{x}{y}\right)_n := \frac{x_n}{y_n} \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n), (y_n) \text{ be real sequences with } \forall n \in \mathbb{N} (y_n \neq 0). \\ &\forall n \in \mathbb{N} \left(\left(\frac{x}{y}\right)_n = \frac{x_n}{y_n} \right). \end{aligned}$$

Remark (Interpretation). The quotient sequence is the product of (x_n) with the reciprocal of (y_n) . Its convergence law is therefore a product law plus a reciprocal law.

Remark (Dependencies).

- [Pointwise Product of Sequences](#)
- [Reciprocal Sequence](#)

Definition (Square Sequence)

Definition 6.3.20 (Square Sequence). Let (x_n) be a real sequence. The **square sequence** of (x_n) is the real sequence (x_n^2) defined by

$$(x^2)_n := x_n^2 \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence.} \\ &\forall n \in \mathbb{N} ((x^2)_n = x_n^2). \end{aligned}$$

Remark (Interpretation). The square sequence is the product of a sequence with itself. Its limit law is therefore a direct specialization of the product law.

Remark (Dependencies).

- [Pointwise Product of Sequences](#)

Definition (Absolute Value Sequence)

Definition 6.3.21 (Absolute Value Sequence). Let (x_n) be a real sequence. The **absolute value sequence** of (x_n) is the real sequence $(|x_n|)$ defined by

$$(|x|)_n := |x_n| \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

Let (x_n) be a real sequence.
 $\forall n \in \mathbb{N} \left((|x|)_n = |x_n| \right).$

Remark (Interpretation). The absolute value sequence records the magnitude of each term. It removes sign information while preserving distance from zero.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Square Root Sequence)

Definition 6.3.22 (Square Root Sequence). Let (x_n) be a real sequence such that $0 \leq x_n$ for every $n \in \mathbb{N}$. The **square root sequence** of (x_n) is the real sequence $(\sqrt{x_n})$ defined by

$$(\sqrt{x})_n := \sqrt{x_n} \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

Let (x_n) be a real sequence with $\forall n \in \mathbb{N} (0 \leq x_n)$.
 $\forall n \in \mathbb{N} \left((\sqrt{x})_n = \sqrt{x_n} \right).$

Remark (Interpretation). The square root sequence is defined term by term on the nonnegative real terms. The nonnegativity hypothesis is part of the construction, not a proof artifact.

Remark (Dependencies).

- [Real Sequence](#)

Theorem (Limit of a Scalar Multiple)

Theorem 6.3.23 (Limit of a Scalar Multiple). Let (x_n) be a real sequence, let $L \in \mathbb{R}$, and let $\alpha \in \mathbb{R}$. If $x_n \rightarrow L$, then $\alpha x_n \rightarrow \alpha L$.

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence and let $L, \alpha \in \mathbb{R}$.
$$\left[\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - L| < \varepsilon) \right]$$
$$\implies \left[\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|\alpha x_n - \alpha L| < \varepsilon) \right].$$

Remark (Predicate reading).

$$\text{ConvergentSequence}(x_n, L) \implies \text{ConvergentSequence}(\alpha x_n, \alpha L).$$

Remark (Interpretation). Multiplying every term by a fixed scalar multiplies the limiting value by the same scalar. The case $\alpha = 0$ collapses the sequence to the constant zero sequence.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Scalar Multiple of a Sequence](#)

Theorem (Limit of a Sum)

Theorem 6.3.24 (Limit of a Sum). *Let (x_n) and (y_n) be real sequences, and let $L, M \in \mathbb{R}$. If $x_n \rightarrow L$ and $y_n \rightarrow M$, then $x_n + y_n \rightarrow L + M$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n), (y_n) \text{ be real sequences and let } L, M \in \mathbb{R}. \\ &[x_n \rightarrow L \wedge y_n \rightarrow M] \implies [x_n + y_n \rightarrow L + M]. \end{aligned}$$

Remark (Predicate reading).

$$(\text{ConvergentSequence}(x_n, L) \wedge \text{ConvergentSequence}(y_n, M)) \implies \text{ConvergentSequence}(x_n + y_n, L + M). \blacksquare$$

Remark (Interpretation). The sum law says that convergent sequences may be added before or after taking limits. The proof splits a target tolerance into two smaller tolerances, one for each summand.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Pointwise Sum of Sequences](#)
- [Triangle Inequality](#)

Theorem (Limit of a Negation)

Theorem 6.3.25 (Limit of a Negation). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. If $x_n \rightarrow L$, then $-x_n \rightarrow -L$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence and let } L \in \mathbb{R}. \\ &[x_n \rightarrow L] \implies [-x_n \rightarrow -L]. \end{aligned}$$

Remark (Predicate reading).

$$\text{ConvergentSequence}(x_n, L) \implies \text{ConvergentSequence}(-x_n, -L).$$

Remark (Interpretation). Negation is scalar multiplication by -1 , so this theorem is the scalar multiple law in its most common special case.

Remark (Dependencies).

- [Limit of a Scalar Multiple](#)
- [Pointwise Negation of a Sequence](#)

Theorem (Limit of a Difference)

Theorem 6.3.26 (Limit of a Difference). *Let (x_n) and (y_n) be real sequences, and let $L, M \in \mathbb{R}$. If $x_n \rightarrow L$ and $y_n \rightarrow M$, then $x_n - y_n \rightarrow L - M$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\text{Let } (x_n), (y_n) \text{ be real sequences and let } L, M \in \mathbb{R}.$$

$$[x_n \rightarrow L \wedge y_n \rightarrow M] \implies [x_n - y_n \rightarrow L - M].$$

Remark (Predicate reading).

$$(\text{ConvergentSequence}(x_n, L) \wedge \text{ConvergentSequence}(y_n, M)) \implies \text{ConvergentSequence}(x_n - y_n, L - M). \blacksquare$$

Remark (Interpretation). The difference law follows by adding (x_n) to the negation of (y_n) . It is therefore not a new analytic mechanism, but a useful named rule.

Remark (Dependencies).

- [Limit of a Sum](#)
- [Limit of a Negation](#)
- [Pointwise Difference of Sequences](#)

Theorem (Limit of a Product)

Theorem 6.3.27 (Limit of a Product). *Let (x_n) and (y_n) be real sequences, and let $L, M \in \mathbb{R}$. If $x_n \rightarrow L$ and $y_n \rightarrow M$, then $x_n y_n \rightarrow LM$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\text{Let } (x_n), (y_n) \text{ be real sequences and let } L, M \in \mathbb{R}.$$

$$[x_n \rightarrow L \wedge y_n \rightarrow M] \implies [x_n y_n \rightarrow LM].$$

Remark (Predicate reading).

$$(\text{ConvergentSequence}(x_n, L) \wedge \text{ConvergentSequence}(y_n, M)) \implies \text{ConvergentSequence}(x_n y_n, LM). \blacksquare$$

Remark (Interpretation). The product law uses both approximation and boundedness. A convergent sequence is eventually bounded, so one factor can be controlled while the other is made small.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Pointwise Product of Sequences](#)
- [Bounded Sequence](#)

Theorem (Nonzero Limit is Eventually Nonzero)

Theorem 6.3.28 (Nonzero Limit is Eventually Nonzero). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$ with $L \neq 0$. If $x_n \rightarrow L$, then there exists $N \in \mathbb{N}$ such that $x_n \neq 0$ for every $n \geq N$.*

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence and let $L \in \mathbb{R}$.

$$[L \neq 0 \wedge x_n \rightarrow L] \implies \exists N \in \mathbb{N} \forall n \geq N (x_n \neq 0).$$

Remark (Interpretation). A sequence converging to a nonzero number eventually stays away from zero. This is the denominator-control result needed for reciprocal and quotient rules.

Remark (Dependencies).

- [Convergence of a Sequence](#)

Theorem (Limit of a Reciprocal)

Theorem 6.3.29 (Limit of a Reciprocal). *Let (x_n) be a real sequence such that $x_n \neq 0$ for every $n \in \mathbb{N}$, and let $L \in \mathbb{R}$ with $L \neq 0$. If $x_n \rightarrow L$, then $1/x_n \rightarrow 1/L$.*

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence with $\forall n \in \mathbb{N} (x_n \neq 0)$, and let $L \neq 0$.

$$[x_n \rightarrow L] \implies \left[\frac{1}{x_n} \rightarrow \frac{1}{L} \right].$$

Remark (Predicate reading).

$$\text{ConvergentSequence}(x_n, L) \implies \text{ConvergentSequence}\left(\frac{1}{x_n}, \frac{1}{L}\right).$$

Remark (Interpretation). Reciprocation is continuous away from zero. The nonzero limit supplies a tail on which the denominators stay uniformly separated from zero.

Remark (Dependencies).

- [Reciprocal Sequence](#)
- [Nonzero Limit is Eventually Nonzero](#)

Theorem (Limit of a Quotient)

Theorem 6.3.30 (Limit of a Quotient). *Let (x_n) and (y_n) be real sequences such that $y_n \neq 0$ for every $n \in \mathbb{N}$, and let $L, M \in \mathbb{R}$ with $M \neq 0$. If $x_n \rightarrow L$ and $y_n \rightarrow M$, then $x_n/y_n \rightarrow L/M$.*

[Go to proof.](#)

Remark (Standard quantified statement).

Let $(x_n), (y_n)$ be real sequences with $\forall n \in \mathbb{N} (y_n \neq 0)$, and let $M \neq 0$.

$$[x_n \rightarrow L \wedge y_n \rightarrow M] \implies \left[\frac{x_n}{y_n} \rightarrow \frac{L}{M}\right].$$

Remark (Predicate reading).

$$(\text{ConvergentSequence}(x_n, L) \wedge \text{ConvergentSequence}(y_n, M)) \implies \text{ConvergentSequence}\left(\frac{x_n}{y_n}, \frac{L}{M}\right).$$

Remark (Interpretation). The quotient law is the product law applied to (x_n) and the reciprocal of (y_n) . The condition $M \neq 0$ prevents the limiting denominator from collapsing.

Remark (Dependencies).

- [Pointwise Quotient of Sequences](#)
- [Limit of a Product](#)
- [Limit of a Reciprocal](#)

Theorem (Limit of a Square)

Theorem 6.3.31 (Limit of a Square). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. If $x_n \rightarrow L$, then $x_n^2 \rightarrow L^2$.*

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence and let $L \in \mathbb{R}$.

$$[x_n \rightarrow L] \implies [x_n^2 \rightarrow L^2].$$

Remark (Predicate reading).

$$\text{ConvergentSequence}(x_n, L) \implies \text{ConvergentSequence}(x_n^2, L^2).$$

Remark (Interpretation). Squaring preserves limits because it is multiplication of a sequence by itself.

Remark (Dependencies).

- [Square Sequence](#)
- [Limit of a Product](#)

Theorem (Limit of an Absolute Value)

Theorem 6.3.32 (Limit of an Absolute Value). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. If $x_n \rightarrow L$, then $|x_n| \rightarrow |L|$.*

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence and let } L \in \mathbb{R}. \\ &[x_n \rightarrow L] \implies [|x_n| \rightarrow |L|]. \end{aligned}$$

Remark (Predicate reading).

$$\text{ConvergentSequence}(x_n, L) \implies \text{ConvergentSequence}(|x_n|, |L|).$$

Remark (Interpretation). Taking absolute values cannot increase distance by more than the original distance: the magnitude function is continuous on $\mathbb{R}$.

Remark (Dependencies).

- [Absolute Value Sequence](#)
- [Triangle Inequality](#)

Theorem (Positive Limit is Eventually Positive)

Theorem 6.3.33 (Positive Limit is Eventually Positive). *Let (x_n) be a real sequence, and let $L > 0$. If $x_n \rightarrow L$, then there exists $N \in \mathbb{N}$ such that $0 < x_n$ for every $n \geq N$.*

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence and let } L > 0. \\ &[x_n \rightarrow L] \implies \exists N \in \mathbb{N} \forall n \geq N (0 < x_n). \end{aligned}$$

Remark (Interpretation). Convergence to a positive limit eventually traps the sequence inside the positive half-line. This result is the sign-control counterpart of the eventual nonzero theorem.

Remark (Dependencies).

- [Convergence of a Sequence](#)

Theorem (Limit of a Square Root)

Theorem 6.3.34 (Limit of a Square Root). *Let (x_n) be a real sequence such that $0 \leq x_n$ for every $n \in \mathbb{N}$, and let $L \in \mathbb{R}$. If $x_n \rightarrow L$, then $0 \leq L$ and $\sqrt{x_n} \rightarrow \sqrt{L}$.*

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence with $\forall n \in \mathbb{N} (0 \leq x_n)$, and let $L \in \mathbb{R}$.

$$[x_n \rightarrow L] \implies [0 \leq L \wedge \sqrt{x_n} \rightarrow \sqrt{L}].$$

Remark (Predicate reading).

$$\text{ConvergentSequence}(x_n, L) \implies \text{ConvergentSequence}(\sqrt{x_n}, \sqrt{L}).$$

Remark (Interpretation). The square-root function is continuous on the nonnegative real line. Near zero the proof uses the order relation; away from zero it can be rationalized by multiplying by the conjugate.

Remark (Dependencies).

- [Square Root Sequence](#)
- [Convergence of a Sequence](#)

Theorem (Polynomial Sequence Limit)

Theorem 6.3.35 (Polynomial Sequence Limit). *Let (x_n) be a real sequence, let $L \in \mathbb{R}$, and let $p \in \mathbb{R}[t]$ be a polynomial. If $x_n \rightarrow L$, then $p(x_n) \rightarrow p(L)$.*

[Go to proof.](#)

Remark (Standard quantified statement).

Let (x_n) be a real sequence, $L \in \mathbb{R}$, and $p \in \mathbb{R}[t]$.

$$[x_n \rightarrow L] \implies [p(x_n) \rightarrow p(L)].$$

Remark (Predicate reading).

$$\text{ConvergentSequence}(x_n, L) \implies \text{ConvergentSequence}(p(x_n), p(L)).$$

Remark (Interpretation). Polynomial limits are built from finitely many additions, scalar multiples, and products. The theorem packages those algebra laws into one reusable composition principle.

Remark (Dependencies).

- [Limit of a Sum](#)

- [Limit of a Scalar Multiple](#)
- [Limit of a Product](#)
- [Limit of a Square](#)

Theorem (Rational Sequence Limit)

Theorem 6.3.36 (Rational Sequence Limit). *Let (x_n) be a real sequence, let $L \in \mathbb{R}$, and let $p, q \in \mathbb{R}[t]$ be polynomials. Suppose $q(L) \neq 0$ and $q(x_n) \neq 0$ for every $n \in \mathbb{N}$. If $x_n \rightarrow L$, then*

$$\frac{p(x_n)}{q(x_n)} \rightarrow \frac{p(L)}{q(L)}.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} &\text{Let } (x_n) \text{ be a real sequence, } L \in \mathbb{R}, \text{ and } p, q \in \mathbb{R}[t]. \\ &[q(L) \neq 0 \wedge \forall n \in \mathbb{N} (q(x_n) \neq 0) \wedge x_n \rightarrow L] \\ &\implies \left[\frac{p(x_n)}{q(x_n)} \rightarrow \frac{p(L)}{q(L)} \right]. \end{aligned}$$

Remark (Predicate reading).

$$\text{ConvergentSequence}(x_n, L) \implies \text{ConvergentSequence}\left(\frac{p(x_n)}{q(x_n)}, \frac{p(L)}{q(L)}\right).$$

Remark (Interpretation). A rational expression is a quotient of two polynomial expressions. The condition $q(L) \neq 0$ keeps the limiting denominator legitimate, and the pointwise condition $q(x_n) \neq 0$ keeps every term of the quotient sequence defined.

Remark (Dependencies).

- [Polynomial Sequence Limit](#)
- [Limit of a Quotient](#)
- [Nonzero Limit is Eventually Nonzero](#)

6.4 Tails

Definition (M -Tail of a Sequence)

Definition 6.4.1 (M -Tail of a Sequence). Let $(x_n) \subseteq \mathbb{R}$ and let $M \in \mathbb{N}$. The **M -tail** of (x_n) is the sequence

$$(x_n)_{n \geq M} := (x_M, x_{M+1}, x_{M+2}, \dots).$$

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \forall M \in \mathbb{N} \left((x_n)_{n \geq M} = (x_M, x_{M+1}, x_{M+2}, \dots) \right).$$

Remark (Definition predicate reading).

$$\text{TailSequence}(x_n, M, y_k) := \forall k \in \mathbb{N} (y_k = x_{M+k-1}).$$

Remark (Interpretation). The M -tail discards the first $M - 1$ terms and retains everything from index M onward. It is the formal object underlying the phrase *all but finitely many terms*: to say that all but finitely many terms of (x_n) satisfy a property P is precisely to say that some M -tail of (x_n) satisfies P termwise. The index M is an offset into the sequence, not a bound on the terms themselves; two sequences with identical M -tails share all asymptotic properties, differing only in finitely many initial values which convergence arguments discard entirely.

Remark (Dependencies).

- [Real Sequence](#)

Theorem (Convergence of a Tail)

Theorem 6.4.2 (Convergence of a Tail). *Let $(x_n) \subseteq \mathbb{R}$ and let $m \in \mathbb{N}$. The m -tail $(x_{m+n})_{n \in \mathbb{N}}$ converges if and only if (x_n) converges. Moreover, in this case,*

$$\lim_{n \rightarrow \infty} x_{m+n} = \lim_{n \rightarrow \infty} x_n.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} & \forall (x_n) \subseteq \mathbb{R} \forall m \in \mathbb{N} \\ & \left((x_{m+n})_{n \in \mathbb{N}} \text{ converges} \iff (x_n) \text{ converges} \right) \\ & \wedge \left((x_n) \text{ converges} \implies \lim_{n \rightarrow \infty} x_{m+n} = \lim_{n \rightarrow \infty} x_n \right). \end{aligned}$$

Remark (Theorem predicate reading).

$$\text{TailConvergenceInvariant}(x_n, m) := \left(\exists L \in \mathbb{R} \text{ ConvergentSequence}(x_n, L) \right) \iff \left(\exists L \in \mathbb{R} \text{ ConvergentSequence}(x_{m+n}, L) \right)$$

Remark (Contrapositive quantified statement).

$$\begin{aligned} & \forall (x_n) \subseteq \mathbb{R} \forall m \in \mathbb{N} \\ & \left((x_{m+n})_{n \in \mathbb{N}} \text{ diverges} \iff (x_n) \text{ diverges} \right). \end{aligned}$$

Remark (Contrapositive predicate reading).

$$\neg \text{TailConvergenceInvariant}(x_n, m) := \left((x_n) \text{ diverges} \iff (x_{m+n})_{n \in \mathbb{N}} \text{ diverges} \right).$$

Remark (Interpretation). Convergence is a *tail property*: it depends only on the eventual behavior of the sequence, not on any finite initial segment. Discarding the first m terms neither creates nor destroys convergence, and when the limit exists it is preserved exactly. This theorem formalizes the informal principle that finitely many terms are irrelevant to asymptotic behavior. It is the rigorous justification for the common proof move of assuming without loss of generality that n is large enough for some auxiliary condition to hold: one simply passes to a suitable m -tail, applies the argument there, and concludes for the full sequence by this theorem. The contrapositive is equally useful: to show (x_n) diverges it suffices to show that some m -tail diverges.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [M-Tail of a Sequence](#)
- [Uniqueness of Limits](#)

Theorem (Convergence by Domination)

Theorem 6.4.3 (Convergence by Domination). *Let $(x_n) \subseteq \mathbb{R}$ and let $L \in \mathbb{R}$. Let (a_n) be a sequence of positive real numbers with $\lim_{n \rightarrow \infty} a_n = 0$. If there exist a constant $c > 0$ and an index $m \in \mathbb{N}$ such that*

$$|x_n - L| \leq c a_n \quad \forall n \geq m,$$

then $\lim_{n \rightarrow \infty} x_n = L$.

Go to proof.

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \forall L \in \mathbb{R} \forall (a_n) \subseteq (0, \infty) \left(\left(\lim_{n \rightarrow \infty} a_n = 0 \wedge \exists c > 0 \exists m \in \mathbb{N} \forall n \geq m (|x_n - L| \leq c a_n) \right) \implies \lim_{n \rightarrow \infty} x_n = L \right).$$

Remark (Contrapositive quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \forall L \in \mathbb{R} \forall (a_n) \subseteq (0, \infty) \left(\lim_{n \rightarrow \infty} x_n \neq L \implies \lim_{n \rightarrow \infty} a_n \neq 0 \vee \forall c > 0 \forall m \in \mathbb{N} \exists n \geq m (|x_n - L| > c a_n) \right).$$

Remark (Contrapositive predicate reading).

$$\text{NotDominatedByNullError}(x_n, L, a_n) := \lim_{n \rightarrow \infty} a_n \neq 0 \vee \forall c > 0 \forall m \in \mathbb{N} \exists n \geq m (|x_n - L| > c a_n).$$

Remark (Interpretation). The theorem is a *domination principle*: it is not necessary to verify the ε - N condition directly for (x_n) . It suffices to find a null sequence (a_n) – one converging to zero – that dominates the error $|x_n - L|$ on some tail, up to a fixed scalar c . The scalar c

is irrelevant to the conclusion: since $a_n \rightarrow 0$, the product $c a_n \rightarrow 0$ regardless of the size of c , and any $\varepsilon > 0$ is eventually beaten by $c a_n$. This theorem is the rigorous engine behind the common proof pattern of bounding $|x_n - L|$ by a simpler expression and observing that the simpler expression tends to zero. The canonical instance is $a_n = 1/n$, where domination by c/n is sufficient to conclude convergence to L .

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [M-Tail of a Sequence](#)

Theorem (Ratio Limit Less Than One Implies Null)

Theorem 6.4.4 (Ratio Limit Less Than One Implies Null). *Let (x_n) be a sequence of positive real numbers. Suppose that the limit*

$$L := \lim_{n \rightarrow \infty} \frac{x_{n+1}}{x_n}$$

exists. If $L < 1$, then (x_n) converges and

$$\lim_{n \rightarrow \infty} x_n = 0.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} & \forall (x_n) \subseteq (0, \infty) \forall L \in \mathbb{R} \\ & \left(\lim_{n \rightarrow \infty} \frac{x_{n+1}}{x_n} = L \wedge L < 1 \right) \implies \lim_{n \rightarrow \infty} x_n = 0. \end{aligned}$$

Remark (Theorem predicate reading).

$$(\text{ConvergentSequence}(x_{n+1}/x_n, L) \wedge L < 1) \implies \text{NullSequence}(x_n).$$

Remark (Negated quantified statement).

$$\begin{aligned} & \exists (x_n) \subseteq (0, \infty) \exists L \in \mathbb{R} \\ & \left(\lim_{n \rightarrow \infty} \frac{x_{n+1}}{x_n} = L \wedge L < 1 \wedge \lim_{n \rightarrow \infty} x_n \neq 0 \right). \end{aligned}$$

Remark (Negation predicate reading).

$$\text{ConvergentSequence}(x_{n+1}/x_n, L) \wedge L < 1 \wedge \text{NotConvergentTo}(x_n, 0).$$

Remark (Failure modes). A failure would be a positive sequence whose consecutive ratios settle below one in the limit but whose terms do not tend to zero. The theorem says this cannot happen: once the ratio limit is less than one, some tail has ratios bounded by a fixed number $r < 1$, forcing geometric decay.

Remark (Failure mode decomposition).

$$\underbrace{(x_n) \subseteq (0, \infty)}_{\text{positive terms}} \quad \underbrace{\lim_{n \rightarrow \infty} \frac{x_{n+1}}{x_n} = L < 1}_{\text{eventual ratio gap}} \quad \underbrace{\lim_{n \rightarrow \infty} x_n \neq 0}_{\text{fails to be null}}.$$

The positive-term hypothesis makes each ratio meaningful. The strict gap $L < 1$ gives a tail on which $x_{n+1} \leq rx_n$ for some $r < 1$, and repeated iteration then gives domination by a null geometric sequence.

Remark (Interpretation). This is the sequence form of the ratio test. The theorem does not merely say that the terms decrease eventually; it says they decrease at an eventually geometric rate. The strict inequality $L < 1$ is essential. When the ratio limit equals 1, the conclusion may hold, as with $1/n$, or fail, as with the constant sequence 1.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Null Sequence](#)
- [M-Tail of a Sequence](#)
- [Convergence by Domination](#)

6.5 Monotonicity

Monotonicity of Sequences

Monotonicity is an order condition on successive terms of a sequence. It is independent of convergence by itself, but when combined with boundedness it becomes one of the fundamental convergence mechanisms in real analysis.

Definition (Increasing Sequence)

Definition 6.5.1 (Increasing Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **increasing** if

$$x_n \leq x_{n+1} \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$(x_n) \text{ is increasing} \iff \forall n \in \mathbb{N} (x_n \leq x_{n+1}).$$

Remark (Definition predicate reading).

$$\text{MonotoneIncreasingSequence}(x_n) := \forall n \in \mathbb{N} (x_n \leq x_{n+1}).$$

Remark (Negated quantified statement).

$$(x_n) \text{ is not increasing} \iff \exists n \in \mathbb{N} (x_{n+1} < x_n).$$

Remark (Negation predicate reading).

$$\neg \text{MonotoneIncreasingSequence}(x_n) \iff \exists n \in \mathbb{N} (x_{n+1} < x_n).$$

Remark (Interpretation). An increasing sequence may stay flat. The term “increasing” is therefore used here in the non-strict sense: each term is at least as large as the one before it.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Decreasing Sequence)

Definition 6.5.2 (Decreasing Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **decreasing** if

$$x_{n+1} \leq x_n \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$(x_n) \text{ is decreasing} \iff \forall n \in \mathbb{N} (x_{n+1} \leq x_n).$$

Remark (Definition predicate reading).

$$\text{MonotoneDecreasingSequence}(x_n) := \forall n \in \mathbb{N} (x_{n+1} \leq x_n).$$

Remark (Negated quantified statement).

$$(x_n) \text{ is not decreasing} \iff \exists n \in \mathbb{N} (x_n < x_{n+1}).$$

Remark (Negation predicate reading).

$$\neg \text{MonotoneDecreasingSequence}(x_n) \iff \exists n \in \mathbb{N} (x_n < x_{n+1}).$$

Remark (Interpretation). A decreasing sequence may stay flat. The order condition says the sequence never rises as the index advances.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Strictly Increasing Sequence)

Definition 6.5.3 (Strictly Increasing Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **strictly increasing** if

$$x_n < x_{n+1} \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$(x_n) \text{ is strictly increasing} \iff \forall n \in \mathbb{N} (x_n < x_{n+1}).$$

Remark (Definition predicate reading).

$$\text{StrictlyIncreasingSequence}(x_n) := \forall n \in \mathbb{N} (x_n < x_{n+1}).$$

Remark (Negated quantified statement).

$$(x_n) \text{ is not strictly increasing} \iff \exists n \in \mathbb{N} (x_{n+1} \leq x_n).$$

Remark (Negation predicate reading).

$$\neg \text{StrictlyIncreasingSequence}(x_n) \iff \exists n \in \mathbb{N} (x_{n+1} \leq x_n).$$

Remark (Interpretation). Strict increase excludes equality between consecutive terms. Every strictly increasing sequence is increasing, but the converse need not hold.

Remark (Dependencies).

- [Real Sequence](#)
- [Increasing Sequence](#)

Definition (Strictly Decreasing Sequence)

Definition 6.5.4 (Strictly Decreasing Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **strictly decreasing** if

$$x_{n+1} < x_n \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$(x_n) \text{ is strictly decreasing} \iff \forall n \in \mathbb{N} (x_{n+1} < x_n).$$

Remark (Definition predicate reading).

$$\text{StrictlyDecreasingSequence}(x_n) := \forall n \in \mathbb{N} (x_{n+1} < x_n).$$

Remark (Negated quantified statement).

$$(x_n) \text{ is not strictly decreasing} \iff \exists n \in \mathbb{N} (x_n \leq x_{n+1}).$$

Remark (Negation predicate reading).

$$\neg \text{StrictlyDecreasingSequence}(x_n) \iff \exists n \in \mathbb{N} (x_n \leq x_{n+1}).$$

Remark (Interpretation). Strict decrease excludes equality between consecutive terms. Every strictly decreasing sequence is decreasing, but the converse need not hold.

Remark (Dependencies).

- [Real Sequence](#)

- [Decreasing Sequence](#)

Definition (Monotone Sequence)

Definition 6.5.5 (Monotone Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **monotone** if it is increasing or decreasing.

Remark (Standard quantified statement).

$$(x_n) \text{ is monotone} \\ \iff [\forall n \in \mathbb{N} (x_n \leq x_{n+1})] \vee [\forall n \in \mathbb{N} (x_{n+1} \leq x_n)].$$

Remark (Definition predicate reading).

$\text{MonotoneSequence}(x_n) := \text{MonotoneIncreasingSequence}(x_n) \vee \text{MonotoneDecreasingSequence}(x_n).$ ■

Remark (Negated quantified statement).

$$(x_n) \text{ is not monotone} \\ \iff [\exists n \in \mathbb{N} (x_{n+1} < x_n)] \wedge [\exists m \in \mathbb{N} (x_m < x_{m+1})].$$

Remark (Negation predicate reading).

$\neg \text{MonotoneSequence}(x_n) \iff \neg \text{MonotoneIncreasingSequence}(x_n) \wedge \neg \text{MonotoneDecreasingSequence}(x_n).$ ■

Remark (Interpretation). Monotonicity means the sequence has one global direction. It may move upward, move downward, or stay constant, but it cannot reverse direction.

Remark (Dependencies).

- [Increasing Sequence](#)
- [Decreasing Sequence](#)

Definition (Eventually Increasing Sequence)

Definition 6.5.6 (Eventually Increasing Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **eventually increasing** if there exists $N \in \mathbb{N}$ such that

$$x_n \leq x_{n+1} \quad \text{for every } n \geq N.$$

Remark (Standard quantified statement).

$$(x_n) \text{ is eventually increasing} \iff \exists N \in \mathbb{N} \forall n \geq N (x_n \leq x_{n+1}).$$

Remark (Definition predicate reading).

$\text{EventuallyIncreasingSequence}(x_n) := \exists N \in \mathbb{N} \forall n \geq N (x_n \leq x_{n+1}).$

Remark (Negated quantified statement).

$$(x_n) \text{ is not eventually increasing} \iff \forall N \in \mathbb{N} \exists n \geq N (x_{n+1} < x_n).$$

Remark (Negation predicate reading).

$$\neg \text{EventuallyIncreasingSequence}(x_n) \iff \forall N \in \mathbb{N} \exists n \geq N (x_{n+1} < x_n).$$

Remark (Interpretation). Eventual increase allows finitely many early reversals. Only the tail of the sequence must move in a nondecreasing direction.

Remark (Dependencies).

- [M-Tail of a Sequence](#)
- [Increasing Sequence](#)

Definition (Eventually Decreasing Sequence)

Definition 6.5.7 (Eventually Decreasing Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **eventually decreasing** if there exists $N \in \mathbb{N}$ such that

$$x_{n+1} \leq x_n \quad \text{for every } n \geq N.$$

Remark (Standard quantified statement).

$$(x_n) \text{ is eventually decreasing} \iff \exists N \in \mathbb{N} \forall n \geq N (x_{n+1} \leq x_n).$$

Remark (Definition predicate reading).

$$\text{EventuallyDecreasingSequence}(x_n) := \exists N \in \mathbb{N} \forall n \geq N (x_{n+1} \leq x_n).$$

Remark (Negated quantified statement).

$$(x_n) \text{ is not eventually decreasing} \iff \forall N \in \mathbb{N} \exists n \geq N (x_n < x_{n+1}).$$

Remark (Negation predicate reading).

$$\neg \text{EventuallyDecreasingSequence}(x_n) \iff \forall N \in \mathbb{N} \exists n \geq N (x_n < x_{n+1}).$$

Remark (Interpretation). Eventual decrease allows the initial terms to behave irregularly. The order condition begins only after some finite threshold.

Remark (Dependencies).

- [M-Tail of a Sequence](#)
- [Decreasing Sequence](#)

Definition (Eventually Monotone Sequence)

Definition 6.5.8 (Eventually Monotone Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **eventually monotone** if it is eventually increasing or eventually decreasing.

Remark (Standard quantified statement).

$$\begin{aligned} (x_n) \text{ is eventually monotone} \\ \iff [\exists N \in \mathbb{N} \forall n \geq N (x_n \leq x_{n+1})] \vee [\exists N \in \mathbb{N} \forall n \geq N (x_{n+1} \leq x_n)]. \end{aligned}$$

Remark (Definition predicate reading).

$\text{EventuallyMonotoneSequence}(x_n) := \text{EventuallyIncreasingSequence}(x_n) \vee \text{EventuallyDecreasingSequence}(x_n)$

Remark (Negated quantified statement).

$$\begin{aligned} (x_n) \text{ is not eventually monotone} \\ \iff [\forall N \in \mathbb{N} \exists n \geq N (x_{n+1} < x_n)] \wedge [\forall N \in \mathbb{N} \exists n \geq N (x_n < x_{n+1})]. \end{aligned}$$

Remark (Negation predicate reading).

$\neg \text{EventuallyMonotoneSequence}(x_n) \iff \neg \text{EventuallyIncreasingSequence}(x_n) \wedge \neg \text{EventuallyDecreasingSequence}(x_n)$

Remark (Interpretation). Eventual monotonicity is the tail version of monotonicity. It is often the right hypothesis in applications, because finitely many early terms do not affect convergence.

Remark (Dependencies).

- [Eventually Increasing Sequence](#)
- [Eventually Decreasing Sequence](#)
- [Convergence of a Tail](#)

Theorem (Monotone Convergence Theorem)

Theorem 6.5.9 (Monotone Convergence Theorem). Let (x_n) be a real sequence.

(a) If (x_n) is increasing and bounded above, then (x_n) converges and

$$\lim_{n \rightarrow \infty} x_n = \sup\{x_n : n \in \mathbb{N}\}.$$

(b) If (x_n) is decreasing and bounded below, then (x_n) converges and

$$\lim_{n \rightarrow \infty} x_n = \inf\{x_n : n \in \mathbb{N}\}.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned}
& \forall (x_n) \subseteq \mathbb{R} \\
& \left([\forall n \in \mathbb{N} (x_n \leq x_{n+1})] \wedge [\exists M \in \mathbb{R} \forall n \in \mathbb{N} (x_n \leq M)] \right) \\
& \implies \exists L \in \mathbb{R} \left(\lim_{n \rightarrow \infty} x_n = L \wedge L = \sup\{x_n : n \in \mathbb{N}\} \right), \\
& \left([\forall n \in \mathbb{N} (x_{n+1} \leq x_n)] \wedge [\exists m \in \mathbb{R} \forall n \in \mathbb{N} (m \leq x_n)] \right) \\
& \implies \exists L \in \mathbb{R} \left(\lim_{n \rightarrow \infty} x_n = L \wedge L = \inf\{x_n : n \in \mathbb{N}\} \right).
\end{aligned}$$

Remark (Theorem predicate reading).

$$\begin{aligned}
& \text{MonotoneIncreasingSequence}(x_n) \wedge \text{BoundedAboveSequence}(x_n) \implies \text{ConvergentSequence}(x_n, \sup\{x_n : n \in \mathbb{N}\}) \\
& \text{MonotoneDecreasingSequence}(x_n) \wedge \text{BoundedBelowSequence}(x_n) \implies \text{ConvergentSequence}(x_n, \inf\{x_n : n \in \mathbb{N}\})
\end{aligned}$$

Remark (Negated quantified statement).

$$\begin{aligned}
& \exists (x_n) \subseteq \mathbb{R} \left([\forall n \in \mathbb{N} (x_n \leq x_{n+1})] \wedge [\exists M \in \mathbb{R} \forall n \in \mathbb{N} (x_n \leq M)] \right. \\
& \quad \left. \wedge (x_n) \text{ does not converge} \right), \\
& \exists (x_n) \subseteq \mathbb{R} \left([\forall n \in \mathbb{N} (x_{n+1} \leq x_n)] \wedge [\exists m \in \mathbb{R} \forall n \in \mathbb{N} (m \leq x_n)] \right. \\
& \quad \left. \wedge (x_n) \text{ does not converge} \right).
\end{aligned}$$

Remark (Negation predicate reading).

$$\begin{aligned}
& \text{MonotoneIncreasingSequence}(x_n) \wedge \text{BoundedAboveSequence}(x_n) \wedge \neg \exists L \in \mathbb{R} \text{ ConvergentSequence}(x_n, L), \\
& \text{MonotoneDecreasingSequence}(x_n) \wedge \text{BoundedBelowSequence}(x_n) \wedge \neg \exists L \in \mathbb{R} \text{ ConvergentSequence}(x_n, L).
\end{aligned}$$

Remark (Failure modes). The theorem would fail only if a one-directional bounded sequence could avoid settling to a real limit. Completeness of $\mathbb{R}$ rules this out: an increasing bounded-above sequence is forced toward the least upper bound of its range, while a decreasing bounded-below sequence is forced toward the greatest lower bound of its range.

Remark (Failure mode decomposition).

$$\underbrace{\forall n \in \mathbb{N} (x_n \leq x_{n+1})}_{\text{one direction}} \quad \underbrace{\exists M \in \mathbb{R} \forall n \in \mathbb{N} (x_n \leq M)}_{\text{ceiling}} \quad \underbrace{(x_n) \text{ does not converge}}_{\text{forbidden by completeness}}.$$

The decreasing case is dual: reverse the inequalities and replace the ceiling by a floor.

Remark (Interpretation). Boundedness prevents escape, and monotonicity prevents oscillation. Together they force convergence. The theorem is one of the standard forms in which the completeness of the real numbers becomes a sequence theorem.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Increasing Sequence](#)
- [Decreasing Sequence](#)
- [Sequence Bounded Above](#)
- [Sequence Bounded Below](#)
- [Supremum](#)
- [Infimum](#)

Theorem (Strict Monotonicity Implies Monotonicity)

Theorem 6.5.10 (Strict Monotonicity Implies Monotonicity). *Let (x_n) be a real sequence.*

- (a) *If (x_n) is strictly increasing, then (x_n) is increasing.*
- (b) *If (x_n) is strictly decreasing, then (x_n) is decreasing.*

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned} \forall(x_n) \subseteq \mathbb{R} \\ \left([\forall n \in \mathbb{N} (x_n < x_{n+1})] \implies [\forall n \in \mathbb{N} (x_n \leq x_{n+1})] \right), \\ \left([\forall n \in \mathbb{N} (x_{n+1} < x_n)] \implies [\forall n \in \mathbb{N} (x_{n+1} \leq x_n)] \right). \end{aligned}$$

Remark (Theorem predicate reading).

$$\begin{aligned} \text{StrictlyIncreasingSequence}(x_n) &\implies \text{MonotoneIncreasingSequence}(x_n), \\ \text{StrictlyDecreasingSequence}(x_n) &\implies \text{MonotoneDecreasingSequence}(x_n). \end{aligned}$$

Remark (Interpretation). Strict monotonicity is stronger than non-strict monotonicity because every strict inequality is also a weak inequality.

Remark (Dependencies).

- [Strictly Increasing Sequence](#)
- [Strictly Decreasing Sequence](#)
- [Increasing Sequence](#)
- [Decreasing Sequence](#)

Theorem (Bounded Monotone Sequence Equivalences)

Theorem 6.5.11 (Bounded Monotone Sequence Equivalences). *Let (x_n) be a real sequence.*

- (a) *If (x_n) is increasing, then (x_n) converges if and only if (x_n) is bounded above.*
- (b) *If (x_n) is decreasing, then (x_n) converges if and only if (x_n) is bounded below.*
- (c) *If (x_n) is monotone, then (x_n) converges if and only if (x_n) is bounded.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned}\forall (x_n) \subseteq \mathbb{R} \\ [\forall n \in \mathbb{N} (x_n \leq x_{n+1})] &\implies \left(\exists L \in \mathbb{R} (x_n \rightarrow L) \iff \exists M \in \mathbb{R} \forall n \in \mathbb{N} (x_n \leq M) \right), \\ [\forall n \in \mathbb{N} (x_{n+1} \leq x_n)] &\implies \left(\exists L \in \mathbb{R} (x_n \rightarrow L) \iff \exists m \in \mathbb{R} \forall n \in \mathbb{N} (m \leq x_n) \right).\end{aligned}$$

Remark (Theorem predicate reading).

$$\begin{aligned}\text{MonotoneIncreasingSequence}(x_n) &\implies \left(\exists L \in \mathbb{R} \text{ConvergentSequence}(x_n, L) \iff \text{BoundedAboveSequence}(x_n) \right), \\ \text{MonotoneDecreasingSequence}(x_n) &\implies \left(\exists L \in \mathbb{R} \text{ConvergentSequence}(x_n, L) \iff \text{BoundedBelowSequence}(x_n) \right).\end{aligned}$$

Remark (Interpretation). For monotone sequences, the only obstruction to real convergence is escape in the monotone direction. Boundedness removes that obstruction.

Remark (Dependencies).

- [Monotone Convergence Theorem](#)
- [Bounded Sequence](#)
- [Convergence of a Sequence](#)

Theorem (Eventually Monotone Convergence Theorem)

Theorem 6.5.12 (Eventually Monotone Convergence Theorem). *Let (x_n) be a real sequence.*

- (a) *If (x_n) is eventually increasing and bounded above, then (x_n) converges.*
- (b) *If (x_n) is eventually decreasing and bounded below, then (x_n) converges.*
- (c) *If (x_n) is eventually monotone and bounded, then (x_n) converges.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned}
& \forall (x_n) \subseteq \mathbb{R} \\
& \left(\exists N \in \mathbb{N} \forall n \geq N (x_n \leq x_{n+1}) \wedge \exists M \in \mathbb{R} \forall n \in \mathbb{N} (x_n \leq M) \right) \\
& \implies \exists L \in \mathbb{R} (x_n \rightarrow L), \\
& \left(\exists N \in \mathbb{N} \forall n \geq N (x_{n+1} \leq x_n) \wedge \exists m \in \mathbb{R} \forall n \in \mathbb{N} (m \leq x_n) \right) \\
& \implies \exists L \in \mathbb{R} (x_n \rightarrow L).
\end{aligned}$$

Remark (Theorem predicate reading).

$$\begin{aligned}
& \text{EventuallyIncreasingSequence}(x_n) \wedge \text{BoundedAboveSequence}(x_n) \implies \exists L \in \mathbb{R} \text{ConvergentSequence}(x_n, L) \\
& \text{EventuallyDecreasingSequence}(x_n) \wedge \text{BoundedBelowSequence}(x_n) \implies \exists L \in \mathbb{R} \text{ConvergentSequence}(x_n, L)
\end{aligned}$$

Remark (Interpretation). This is the tail version of the Monotone Convergence Theorem. Once the tail is monotone and bounded in the right direction, finite initial behavior is irrelevant.

Remark (Dependencies).

- [Eventually Increasing Sequence](#)
- [Eventually Decreasing Sequence](#)
- [Convergence of a Tail](#)
- [Monotone Convergence Theorem](#)

Theorem (Unbounded Monotone Divergence)

Theorem 6.5.13 (Unbounded Monotone Divergence). *Let (x_n) be a real sequence.*

(a) *If (x_n) is increasing and is not bounded above, then $x_n \rightarrow +\infty$.*

(b) *If (x_n) is decreasing and is not bounded below, then $x_n \rightarrow -\infty$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned}
& \forall (x_n) \subseteq \mathbb{R} \\
& \left([\forall n \in \mathbb{N} (x_n \leq x_{n+1})] \wedge [\forall M \in \mathbb{R} \exists n \in \mathbb{N} (M < x_n)] \right) \\
& \implies \forall M \in \mathbb{R} \exists N \in \mathbb{N} \forall n \geq N (M < x_n), \\
& \left([\forall n \in \mathbb{N} (x_{n+1} \leq x_n)] \wedge [\forall M \in \mathbb{R} \exists n \in \mathbb{N} (x_n < M)] \right) \\
& \implies \forall M \in \mathbb{R} \exists N \in \mathbb{N} \forall n \geq N (x_n < M).
\end{aligned}$$

Remark (Theorem predicate reading).

$\text{MonotoneIncreasingSequence}(x_n) \wedge \neg \text{BoundedAboveSequence}(x_n) \implies \text{DivergesToPositiveInfinity}(x_n),$
 $\text{MonotoneDecreasingSequence}(x_n) \wedge \neg \text{BoundedBelowSequence}(x_n) \implies \text{DivergesToNegativeInfinity}(x_n).$

Remark (Interpretation). For a monotone sequence, unboundedness has a direction. An increasing sequence that has no upper bound eventually exceeds every real number; the decreasing case is the order-dual statement.

Remark (Dependencies).

- [Divergence to Positive Infinity](#)
- [Divergence to Negative Infinity](#)
- [Increasing Sequence](#)
- [Decreasing Sequence](#)
- [Sequence Bounded Above](#)
- [Sequence Bounded Below](#)

Theorem (Algebraic Transformations Preserve Monotonicity)

Theorem 6.5.14 (Algebraic Transformations Preserve Monotonicity). *Let (x_n) be a real sequence and let $c, \alpha \in \mathbb{R}$.*

- (a) *If (x_n) is increasing, then $(x_n + c)$ is increasing.*
- (b) *If (x_n) is decreasing, then $(x_n + c)$ is decreasing.*
- (c) *If (x_n) is increasing and $\alpha > 0$, then (αx_n) is increasing.*
- (d) *If (x_n) is increasing and $\alpha < 0$, then (αx_n) is decreasing.*
- (e) *If (x_n) is increasing, then $(-x_n)$ is decreasing.*
- (f) *If (x_n) is decreasing, then $(-x_n)$ is increasing.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \forall (x_n) \subseteq \mathbb{R} \, \forall c, \alpha \in \mathbb{R} \\ [\forall n \in \mathbb{N} (x_n \leq x_{n+1})] &\implies [\forall n \in \mathbb{N} (x_n + c \leq x_{n+1} + c)], \\ (\alpha > 0 \wedge \forall n \in \mathbb{N} (x_n \leq x_{n+1})) &\implies [\forall n \in \mathbb{N} (\alpha x_n \leq \alpha x_{n+1})], \\ (\alpha < 0 \wedge \forall n \in \mathbb{N} (x_n \leq x_{n+1})) &\implies [\forall n \in \mathbb{N} (\alpha x_{n+1} \leq \alpha x_n)]. \end{aligned}$$

Remark (Theorem predicate reading).

$$\begin{aligned} \text{MonotoneIncreasingSequence}(x_n) &\implies \text{MonotoneIncreasingSequence}(x_n + c), \\ \text{MonotoneIncreasingSequence}(x_n) \wedge \alpha > 0 &\implies \text{MonotoneIncreasingSequence}(\alpha x_n), \\ \text{MonotoneIncreasingSequence}(x_n) \wedge \alpha < 0 &\implies \text{MonotoneDecreasingSequence}(\alpha x_n). \end{aligned}$$

Remark (Interpretation). Translation preserves order, multiplication by a positive scalar preserves order, and multiplication by a negative scalar reverses order. Negation is the special case $\alpha = -1$.

Remark (Dependencies).

- [Pointwise Sum of Sequences](#)
- [Scalar Multiple of a Sequence](#)
- [Pointwise Negation of a Sequence](#)
- [Increasing Sequence](#)
- [Decreasing Sequence](#)

6.6 Subsequences

Subsequences

Subsequences record what remains after passing to an infinite ordered selection of indices. They are the basic tool for extracting structure from a sequence: limits, oscillation, boundedness, compactness, and failure of convergence are all detected by suitable subsequences.

Definition (Strictly Increasing Index Map)

Definition 6.6.1 (Strictly Increasing Index Map). Let $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ be a function. The function σ is a **strictly increasing index map** if

$$k < \ell \implies \sigma(k) < \sigma(\ell) \quad \text{for all } k, \ell \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$\sigma : \mathbb{N} \rightarrow \mathbb{N} \text{ is strictly increasing} \iff \forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)).$$

Remark (Definition predicate reading).

$$\text{StrictlyIncreasing}(\sigma) := \forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)).$$

Remark (Negated quantified statement).

$$\sigma \text{ is not strictly increasing} \iff \exists k, \ell \in \mathbb{N} (k < \ell \wedge \sigma(\ell) \leq \sigma(k)).$$

Remark (Negation predicate reading).

$$\neg \text{StrictlyIncreasing}(\sigma) \iff \exists k, \ell \in \mathbb{N} (k < \ell \wedge \sigma(\ell) \leq \sigma(k)).$$

Remark (Interpretation). The index map selects terms without repetition and without changing their order. This is the structural difference between a subsequence and an arbitrary rearrangement.

Remark (Dependencies).

- [Function](#)
- [Natural Numbers](#)
- [Strict Order Successor Characterization](#)

Definition (Subsequence)

Definition 6.6.2 (Subsequence). Let (x_n) be a real sequence. A **subsequence** of (x_n) is a sequence $(x_{\sigma(k)})_{k \in \mathbb{N}}$, where $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ is a strictly increasing index map.

Remark (Standard quantified statement).

$$\begin{aligned} (y_k) \text{ is a subsequence of } (x_n) \\ \iff \exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \left(\forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \wedge \forall k \in \mathbb{N} (y_k = x_{\sigma(k)}) \right). \end{aligned}$$

Remark (Definition predicate reading).

$$\text{Subsequence}(y_k, x_n) := \exists \sigma \left(\text{StrictlyIncreasing}(\sigma) \wedge \forall k \in \mathbb{N} (y_k = x_{\sigma(k)}) \right).$$

Remark (Negated quantified statement).

$$\begin{aligned} (y_k) \text{ is not a subsequence of } (x_n) \\ \iff \forall \sigma : \mathbb{N} \rightarrow \mathbb{N} \left(\exists k, \ell \in \mathbb{N} (k < \ell \wedge \sigma(\ell) \leq \sigma(k)) \vee \exists k \in \mathbb{N} (y_k \neq x_{\sigma(k)}) \right). \end{aligned}$$

Remark (Negation predicate reading).

$$\neg \text{Subsequence}(y_k, x_n).$$

Remark (Interpretation). A subsequence keeps the original chronological order but may skip terms. The new index k counts the selected terms, while $\sigma(k)$ records where each selected term came from in the original sequence.

Remark (Dependencies).

- [Real Sequence](#)
- [Strictly Increasing Index Map](#)

Definition (Subsequential Limit)

Definition 6.6.3 (Subsequential Limit). Let (x_n) be a real sequence and let $L \in \mathbb{R}$. The real number L is a **subsequential limit** of (x_n) if there exists a subsequence $(x_{\sigma(k)})$ such that

$$\lim_{k \rightarrow \infty} x_{\sigma(k)} = L.$$

Remark (Standard quantified statement).

L is a subsequential limit of (x_n)

$$\iff \exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \left(\forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \wedge \lim_{k \rightarrow \infty} x_{\sigma(k)} = L \right).$$

Remark (Definition predicate reading).

$\text{SubsequentialLimit}(x_n, L) := \exists \sigma \left(\text{StrictlyIncreasing}(\sigma) \wedge \text{ConvergentSequence}(x_{\sigma(k)}, L) \right).$

Remark (Interpretation). Subsequential limits detect limiting behavior along an infinite selection of terms. A sequence may have no ordinary limit but still have one or more subsequential limits.

Remark (Dependencies).

- [Subsequence](#)
- [Convergence of a Sequence](#)

Definition (Has Convergent Subsequence)

Definition 6.6.4 (Has Convergent Subsequence). Let (x_n) be a real sequence. The sequence (x_n) **has a convergent subsequence** if there exist a strictly increasing index map $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ and a real number L such that

$$\lim_{k \rightarrow \infty} x_{\sigma(k)} = L.$$

Remark (Standard quantified statement).

(x_n) has a convergent subsequence

$$\iff \exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \exists L \in \mathbb{R} \left(\forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \wedge \lim_{k \rightarrow \infty} x_{\sigma(k)} = L \right).$$

Remark (Definition predicate reading).

$\text{HasConvergentSubsequence}(x_n) := \exists L \in \mathbb{R} \text{ SubsequentialLimit}(x_n, L).$

Remark (Interpretation). This property asks for at least one convergent infinite selection. It does not assert convergence of the whole sequence and does not require uniqueness of the subsequential limit.

Remark (Dependencies).

- [Subsequential Limit](#)

Theorem (Subsequence Indices Dominate the Identity)

Theorem 6.6.5 (Subsequence Indices Dominate the Identity). *Let $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ be a strictly increasing index map. Then*

$$k \leq \sigma(k) \quad \text{for every } k \in \mathbb{N}.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall \sigma : \mathbb{N} \rightarrow \mathbb{N} \left([\forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell))] \implies [\forall k \in \mathbb{N} (k \leq \sigma(k))] \right).$$

Remark (Theorem predicate reading).

$$\text{StrictlyIncreasing}(\sigma) \implies \forall k \in \mathbb{N} (k \leq \sigma(k)).$$

Remark (Interpretation). A subsequence cannot outrun the original indexing backwards. By the time the subsequence has selected its k th term, it must have reached at least the k th original position.

Remark (Dependencies).

- [Strictly Increasing Index Map](#)
- [Well-Ordering Principle](#)

Theorem (Subsequences Preserve Limits)

Theorem 6.6.6 (Subsequences Preserve Limits). *Let (x_n) be a real sequence and let $L \in \mathbb{R}$. If $x_n \rightarrow L$, then every subsequence of (x_n) converges to L .*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \forall L \in \mathbb{R} \forall \sigma : \mathbb{N} \rightarrow \mathbb{N} \left(\left(x_n \rightarrow L \wedge \forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \right) \implies \lim_{k \rightarrow \infty} x_{\sigma(k)} = L \right).$$

Remark (Theorem predicate reading).

$$\text{ConvergentSequence}(x_n, L) \wedge \text{StrictlyIncreasing}(\sigma) \implies \text{ConvergentSequence}(x_{\sigma(k)}, L).$$

Remark (Interpretation). Subsequences cannot escape a limit already possessed by the whole sequence. The domination of subsequence indices by the identity transfers every tail estimate for (x_n) to a tail estimate for $(x_{\sigma(k)})$.

Remark (Dependencies).

- [Subsequence](#)
- [Convergence of a Sequence](#)
- [Subsequence Indices Dominate the Identity](#)

Theorem (Subsequential Limit of a Convergent Sequence)

Theorem 6.6.7 (Subsequential Limit of a Convergent Sequence). *Let (x_n) be a real sequence and let $L \in \mathbb{R}$. If $x_n \rightarrow L$, then L is the only possible subsequential limit of (x_n) .*

Go to proof.

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \forall L, K \in \mathbb{R} \\ \left(x_n \rightarrow L \wedge K \text{ is a subsequential limit of } (x_n) \right) \implies K = L.$$

Remark (Theorem predicate reading).

$$\text{ConvergentSequence}(x_n, L) \wedge \text{SubsequentialLimit}(x_n, K) \implies K = L.$$

Remark (Interpretation). A convergent sequence has no hidden alternative limiting behavior. Every convergent subsequence inherits the same limit, and uniqueness of limits then forces equality.

Remark (Dependencies).

- [Subsequences Preserve Limits](#)
- [Uniqueness of Limits](#)

Theorem (Divergence by Two Subsequential Limits)

Theorem 6.6.8 (Divergence by Two Subsequential Limits). *Let (x_n) be a real sequence. If (x_n) has two distinct subsequential limits, then (x_n) does not converge.*

Go to proof.

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \forall L, K \in \mathbb{R} \\ \left(L \text{ is a subsequential limit of } (x_n) \wedge K \text{ is a subsequential limit of } (x_n) \wedge L \neq K \right) \\ \implies (x_n) \text{ does not converge.}$$

Remark (Theorem predicate reading).

$$\text{SubsequentialLimit}(x_n, L) \wedge \text{SubsequentialLimit}(x_n, K) \wedge L \neq K \implies \neg \exists A \in \mathbb{R} \text{ ConvergentSequence}(x_n, A).$$

Remark (Interpretation). Two different subsequential limits reveal persistent oscillation or splitting. If the whole sequence had a limit, every subsequence would have to inherit that same limit.

Remark (Dependencies).

- [Subsequential Limit](#)
- [Subsequential Limit of a Convergent Sequence](#)

Theorem (Boundedness Passes to Subsequences)

Theorem 6.6.9 (Boundedness Passes to Subsequences). *Every subsequence of a bounded real sequence is bounded.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \forall (x_n) \subseteq \mathbb{R} \ \forall \sigma : \mathbb{N} \rightarrow \mathbb{N} \\ \left(\exists M > 0 \ \forall n \in \mathbb{N} (|x_n| \leq M) \wedge \forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \right) \\ \implies \exists M > 0 \ \forall k \in \mathbb{N} (|x_{\sigma(k)}| \leq M). \end{aligned}$$

Remark (Theorem predicate reading).

$$\text{BoundedSequence}(x_n) \wedge \text{StrictlyIncreasing}(\sigma) \implies \text{BoundedSequence}(x_{\sigma(k)}).$$

Remark (Interpretation). A subsequence cannot create new values; it only selects from the original sequence. Any global bound for the original sequence is automatically a bound for the selection.

Remark (Dependencies).

- [Bounded Sequence](#)
- [Subsequence](#)

Theorem (Monotonicity Passes to Subsequences)

Theorem 6.6.10 (Monotonicity Passes to Subsequences). *Every subsequence of an increasing real sequence is increasing, and every subsequence of a decreasing real sequence is decreasing.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \forall (x_n) \subseteq \mathbb{R} \ \forall \sigma : \mathbb{N} \rightarrow \mathbb{N} \\ \left(\left[\forall n \in \mathbb{N} (x_n \leq x_{n+1}) \right] \wedge \left[\forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \right] \right) \\ \implies \forall k \in \mathbb{N} (x_{\sigma(k)} \leq x_{\sigma(k+1)}). \end{aligned}$$

Remark (Theorem predicate reading).

$\text{MonotoneIncreasingSequence}(x_n) \wedge \text{StrictlyIncreasing}(\sigma) \implies \text{MonotoneIncreasingSequence}(x_{\sigma(k)})$. ■

Remark (Interpretation). Selecting terms in the original order preserves the order pattern already present in the sequence. This is why the index map must be strictly increasing.

Remark (Dependencies).

- [Increasing Sequence](#)
- [Decreasing Sequence](#)
- [Subsequence](#)

Theorem (Subsequence of a Subsequence)

Theorem 6.6.11 (Subsequence of a Subsequence). *Every subsequence of a subsequence of (x_n) is a subsequence of (x_n) .*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} & \forall (x_n) \subseteq \mathbb{R} \, \forall \sigma, \tau : \mathbb{N} \rightarrow \mathbb{N} \\ & \left(\left[\forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \right] \wedge \left[\forall k, \ell \in \mathbb{N} (k < \ell \implies \tau(k) < \tau(\ell)) \right] \right) \\ & \implies (x_{\sigma(\tau(k))})_{k \in \mathbb{N}} \text{ is a subsequence of } (x_n). \end{aligned}$$

Remark (Theorem predicate reading).

$$\text{StrictlyIncreasing}(\sigma) \wedge \text{StrictlyIncreasing}(\tau) \implies \text{Subsequence}(x_{\sigma(\tau(k))}, x_n).$$

Remark (Interpretation). Passing to subsequences is closed under iteration. The composed index map $\sigma \circ \tau$ is still strictly increasing.

Remark (Dependencies).

- [Subsequence](#)
- [Composition](#)

Theorem (Eventual Properties Pass to Subsequences)

Theorem 6.6.12 (Eventual Properties Pass to Subsequences). *Let $P(n)$ be a property of indices. If there exists $N \in \mathbb{N}$ such that $P(n)$ holds for every $n \geq N$, then for every strictly increasing index map σ there exists $K \in \mathbb{N}$ such that $P(\sigma(k))$ holds for every $k \geq K$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} & \left(\exists N \in \mathbb{N} \forall n \geq N P(n) \wedge \forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \right) \\ & \implies \exists K \in \mathbb{N} \forall k \geq K P(\sigma(k)). \end{aligned}$$

Remark (Theorem predicate reading).

$$\text{Eventually}(x_n, P) \wedge \text{StrictlyIncreasing}(\sigma) \implies \text{Eventually}(x_{\sigma(k)}, P).$$

Remark (Interpretation). Subsequences eventually enter every tail of the original sequence. Any property that holds on a tail of the original therefore holds on a tail of the subsequence.

Remark (Dependencies).

- [Subsequence Indices Dominate the Identity](#)
- [Subsequence](#)

Theorem (Frequent Properties Yield Subsequences)

Theorem 6.6.13 (Frequent Properties Yield Subsequences). *Let $P(n)$ be a property of indices. If for every $N \in \mathbb{N}$ there exists $n \geq N$ such that $P(n)$ holds, then there exists a strictly increasing index map $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ such that $P(\sigma(k))$ holds for every $k \in \mathbb{N}$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\left[\forall N \in \mathbb{N} \exists n \geq N P(n) \right] \implies \exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \left(\forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \wedge \forall k \in \mathbb{N} P(\sigma(k)) \right).$$

Remark (Theorem predicate reading).

$$\text{Frequently}(x_n, P) \implies \exists \sigma \left(\text{StrictlyIncreasing}(\sigma) \wedge \forall k \in \mathbb{N} P(\sigma(k)) \right).$$

Remark (Interpretation). An infinitely recurring property can be threaded into a subsequence. The construction chooses one later witness at each step.

Remark (Dependencies).

- [Well-Ordering Principle](#)
- [Subsequence](#)

Theorem (Subsequential Limits Respect Bounds)

Theorem 6.6.14 (Subsequential Limits Respect Bounds). *Let (x_n) be a real sequence and let L be a subsequential limit of (x_n) . If $m \leq x_n \leq M$ for every $n \in \mathbb{N}$, then $m \leq L \leq M$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \forall L, m, M \in \mathbb{R} \\ \left(L \text{ is a subsequential limit of } (x_n) \wedge \forall n \in \mathbb{N} (m \leq x_n \leq M) \right) \implies m \leq L \leq M.$$

Remark (Theorem predicate reading).

$$\text{SubsequentialLimit}(x_n, L) \wedge \forall n \in \mathbb{N} (m \leq x_n \leq M) \implies m \leq L \leq M.$$

Remark (Interpretation). Bounds pass to subsequences, and limits preserve eventual inequalities. Hence subsequential limits remain inside any closed interval that contains all terms.

Remark (Dependencies).

- [Boundedness Passes to Subsequences](#)
- [Constant Comparison for Sequence Limits](#)

Theorem (Squeeze Passes to Subsequences)

Theorem 6.6.15 (Squeeze Passes to Subsequences). *Let (a_n) , (x_n) , and (b_n) be real sequences. If $a_n \leq x_n \leq b_n$ eventually, then for every subsequence $(x_{\sigma(k)})$ there exists $K \in \mathbb{N}$ such that*

$$a_{\sigma(k)} \leq x_{\sigma(k)} \leq b_{\sigma(k)} \quad \text{for every } k \geq K.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\left(\exists N \in \mathbb{N} \forall n \geq N (a_n \leq x_n \leq b_n) \wedge \forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \right) \\ \implies \exists K \in \mathbb{N} \forall k \geq K (a_{\sigma(k)} \leq x_{\sigma(k)} \leq b_{\sigma(k)}).$$

Remark (Theorem predicate reading).

$$\text{Eventually}(x_n, a_n \leq x_n \leq b_n) \wedge \text{StrictlyIncreasing}(\sigma) \implies \text{Eventually}(x_{\sigma(k)}, a_{\sigma(k)} \leq x_{\sigma(k)} \leq b_{\sigma(k)}). \blacksquare$$

Remark (Interpretation). Squeeze hypotheses are tail hypotheses, so they survive passage to subsequences.

Remark (Dependencies).

- [Eventual Properties Pass to Subsequences](#)
- [Sequence Squeeze Theorem](#)

Theorem (Monotone Subsequence Theorem)

Theorem 6.6.16 (Monotone Subsequence Theorem). *Every real sequence has a monotone subsequence.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall(x_n) \subseteq \mathbb{R} \exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \left(\forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \wedge \left[\forall k \in \mathbb{N} (x_{\sigma(k)} \leq x_{\sigma(k+1)}) \vee \forall k \in \mathbb{N} (x_{\sigma(k+1)} \leq x_{\sigma(k)}) \right] \right)$$

Remark (Theorem predicate reading).

$$\exists \sigma \left(\text{StrictlyIncreasing}(\sigma) \wedge \text{MonotoneSequence}(x_{\sigma(k)}) \right).$$

Remark (Interpretation). Every sequence contains an infinite ordered pattern. The proof uses the peak dichotomy: either there are infinitely many terms larger than all later terms, or there is an infinite rising chain.

Remark (Dependencies).

- [Monotone Sequence](#)
- [Subsequence](#)
- [Well-Ordering Principle](#)

Theorem (Bolzano-Weierstrass Theorem for Sequences)

Theorem 6.6.17 (Bolzano-Weierstrass Theorem for Sequences). *Every bounded real sequence has a convergent subsequence.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall(x_n) \subseteq \mathbb{R} \left(\exists M > 0 \forall n \in \mathbb{N} (|x_n| \leq M) \implies \exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \exists L \in \mathbb{R} \left[\forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \wedge \lim_{k \rightarrow \infty} x_{\sigma(k)} = L \right] \right)$$

Remark (Theorem predicate reading).

$$\text{BoundedSequence}(x_n) \implies \text{HasConvergentSubsequence}(x_n).$$

Remark (Interpretation). Bolzano-Weierstrass is the compactness principle for bounded real sequences. Boundedness prevents escape, and the monotone subsequence theorem supplies an ordered subsequence to which the Monotone Convergence Theorem applies.

Remark (Dependencies).

- [Bounded Sequence](#)
- [Monotone Subsequence Theorem](#)
- [Boundedness Passes to Subsequences](#)
- [Bounded Monotone Sequence Equivalences](#)

Theorem (Sequential Compactness of Closed Bounded Intervals)

Theorem 6.6.18 (Sequential Compactness of Closed Bounded Intervals). *Let $a, b \in \mathbb{R}$ with $a \leq b$. Every sequence in $[a, b]$ has a convergent subsequence whose limit belongs to $[a, b]$.*

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned} & \forall a, b \in \mathbb{R} \, \forall (x_n) \subseteq \mathbb{R} \\ & \left(a \leq b \wedge \forall n \in \mathbb{N} (a \leq x_n \leq b) \right) \\ & \implies \exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \, \exists L \in \mathbb{R} \left(\forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \wedge \lim_{k \rightarrow \infty} x_{\sigma(k)} = L \wedge a \leq L \leq b \right). \end{aligned}$$

Remark (Theorem predicate reading).

$$[\forall n \in \mathbb{N} (a \leq x_n \leq b)] \implies \exists L \in [a, b] \, \text{SubsequentialLimit}(x_n, L).$$

Remark (Interpretation). Closed bounded intervals are sequentially compact: a sequence trapped in the interval has a convergent subsequence, and closedness keeps the subsequential limit inside the interval.

Remark (Dependencies).

- [Bolzano-Weierstrass Theorem for Sequences](#)
- [Subsequential Limits Respect Bounds](#)

6.7 Divergence

Divergence — Quick Reference

Concept	Meaning
Divergent sequence	Sequence that does not converge in $\mathbb{R}$
Diverges to $+\infty$	Eventually exceeds every real bound
Diverges to $-\infty$	Eventually lies below every real bound
Oscillatory sequence	Divergence by persistent oscillation
Infinite divergence	Divergence to infinity implies real divergence
Two subsequential limits	Distinct subsequential limits force divergence
Bounded divergence	Bounded divergence yields multiple subsequential limits

Divergence

Divergence is not a single behavior. A sequence can fail to converge because it escapes upward, escapes downward, oscillates between incompatible limiting patterns, or remains

bounded while refusing to settle. The purpose of this section is to separate these failure modes and connect them to subsequences.

Definition (Divergent Sequence)

Definition 6.7.1 (Divergent Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **divergent** if there is no $L \in \mathbb{R}$ such that $x_n \rightarrow L$.

Remark (Standard quantified statement).

$$(x_n) \text{ is divergent} \\ \iff \forall L \in \mathbb{R} \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N (|x_n - L| \geq \varepsilon).$$

Remark (Definition predicate reading).

$$\text{DivergentSequence}(x_n) := \forall L \in \mathbb{R} \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N (|x_n - L| \geq \varepsilon).$$

Remark (Negated quantified statement).

$$(x_n) \text{ is not divergent} \iff \exists L \in \mathbb{R} \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - L| < \varepsilon).$$

Remark (Negation predicate reading).

$$\neg \text{DivergentSequence}(x_n) \iff \exists L \in \mathbb{R} \text{ConvergentSequence}(x_n, L).$$

Remark (Interpretation). Divergence is the formal negation of real convergence. It says that every candidate limit fails at some fixed tolerance on every tail.

Remark (Dependencies).

- [Real Sequence](#)
- [Convergence of a Sequence](#)

Definition (Divergence to Positive Infinity)

Definition 6.7.2 (Divergence to Positive Infinity). Let (x_n) be a real sequence. The sequence (x_n) **diverges to** $+\infty$ if

$$\forall M \in \mathbb{R} \exists N \in \mathbb{N} \forall n \geq N (M < x_n).$$

Remark (Standard quantified statement).

$$x_n \rightarrow +\infty \iff \forall M \in \mathbb{R} \exists N \in \mathbb{N} \forall n \geq N (M < x_n).$$

Remark (Definition predicate reading).

$$\text{DivergesToPositiveInfinity}(x_n) := \forall M \in \mathbb{R} \exists N \in \mathbb{N} \forall n \geq N (M < x_n).$$

Remark (Negated quantified statement).

$$x_n \not\rightarrow +\infty \iff \exists M \in \mathbb{R} \forall N \in \mathbb{N} \exists n \geq N (x_n \leq M).$$

Remark (Interpretation). Divergence to $+\infty$ means every horizontal threshold is eventually left below the sequence. It is not convergence to a real number; it is escape from the real line in the positive direction.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Divergence to Negative Infinity)

Definition 6.7.3 (Divergence to Negative Infinity). Let (x_n) be a real sequence. The sequence (x_n) **diverges to** $-\infty$ if

$$\forall M \in \mathbb{R} \exists N \in \mathbb{N} \forall n \geq N (x_n < M).$$

Remark (Standard quantified statement).

$$x_n \rightarrow -\infty \iff \forall M \in \mathbb{R} \exists N \in \mathbb{N} \forall n \geq N (x_n < M).$$

Remark (Definition predicate reading).

$$\text{DivergesToNegativeInfinity}(x_n) := \forall M \in \mathbb{R} \exists N \in \mathbb{N} \forall n \geq N (x_n < M).$$

Remark (Negated quantified statement).

$$x_n \not\rightarrow -\infty \iff \exists M \in \mathbb{R} \forall N \in \mathbb{N} \exists n \geq N (M \leq x_n).$$

Remark (Interpretation). Divergence to $-\infty$ is escape past every lower threshold. It is the order-dual of divergence to $+\infty$.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Oscillatory Sequence)

Definition 6.7.4 (Oscillatory Sequence). Let (x_n) be a real sequence. The sequence (x_n) is **oscillatory** if it is divergent, does not diverge to $+\infty$, and does not diverge to $-\infty$.

Remark (Standard quantified statement).

$$\begin{aligned} (x_n) \text{ is oscillatory} \\ \iff & \left[\forall L \in \mathbb{R} \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N (|x_n - L| \geq \varepsilon) \right] \\ & \wedge \left[\exists M \in \mathbb{R} \forall N \in \mathbb{N} \exists n \geq N (x_n \leq M) \right] \\ & \wedge \left[\exists M \in \mathbb{R} \forall N \in \mathbb{N} \exists n \geq N (M \leq x_n) \right]. \end{aligned}$$

Remark (Definition predicate reading).

$$\begin{aligned}\text{OscillatorySequence}(x_n) &:= \text{DivergentSequence}(x_n) \\ &\quad \wedge \neg \text{DivergesToPositiveInfinity}(x_n) \\ &\quad \wedge \neg \text{DivergesToNegativeInfinity}(x_n).\end{aligned}$$

Remark (Interpretation). Oscillation means failure to settle without one-directional escape. The sequence may remain bounded, as $(-1)^n$ does, or it may be unbounded while still repeatedly returning in incompatible directions.

Remark (Dependencies).

- [Divergent Sequence](#)
- [Divergence to Positive Infinity](#)
- [Divergence to Negative Infinity](#)

Theorem (Divergence to Infinity Implies Real Divergence)

Theorem 6.7.5 (Divergence to Infinity Implies Real Divergence). *Let (x_n) be a real sequence. If $x_n \rightarrow +\infty$ or $x_n \rightarrow -\infty$, then (x_n) is divergent.*

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned}&\left[\forall M \in \mathbb{R} \exists N \in \mathbb{N} \forall n \geq N (M < x_n) \right] \\ &\quad \vee \left[\forall M \in \mathbb{R} \exists N \in \mathbb{N} \forall n \geq N (x_n < M) \right] \\ &\implies \forall L \in \mathbb{R} \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N (|x_n - L| \geq \varepsilon).\end{aligned}$$

Remark (Definition predicate reading).

$$\text{DivergesToPositiveInfinity}(x_n) \vee \text{DivergesToNegativeInfinity}(x_n) \implies \text{DivergentSequence}(x_n). \blacksquare$$

Remark (Interpretation). Infinity divergence is a stronger, directional failure of real convergence. A sequence escaping past every real threshold cannot approach a finite real limit.

Remark (Dependencies).

- [Divergent Sequence](#)
- [Divergence to Positive Infinity](#)
- [Divergence to Negative Infinity](#)

Theorem (Two Subsequential Limits Force Divergence)

Theorem 6.7.6 (Two Subsequential Limits Force Divergence). *Let (x_n) be a real sequence. If (x_n) has two subsequential limits $L, K \in \mathbb{R}$ with $L \neq K$, then (x_n) is divergent.*

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned} \exists L, K \in \mathbb{R} \left(L \neq K \wedge L \text{ is a subsequential limit of } (x_n) \wedge K \text{ is a subsequential limit of } (x_n) \right) \\ \implies \forall A \in \mathbb{R} \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N (|x_n - A| \geq \varepsilon). \end{aligned}$$

Remark (Definition predicate reading).

$$\begin{aligned} \exists L, K \in \mathbb{R} (L \neq K \wedge \text{SubsequentialLimit}(x_n, L) \wedge \text{SubsequentialLimit}(x_n, K)) \\ \implies \text{DivergentSequence}(x_n). \end{aligned}$$

Remark (Interpretation). Convergence forces all subsequences to inherit the same limit. Two different subsequential limits therefore witness an irreparable conflict in the tail behavior of the original sequence.

Remark (Dependencies).

- [Subsequential Limit](#)
- [Subsequences Preserve Limits](#)
- [Uniqueness of Limits](#)
- [Divergence by Two Subsequential Limits](#)

Theorem (Unbounded Above Sequence Has a Positive-Infinity Subsequence)

Theorem 6.7.7 (Unbounded Above Sequence Has a Positive-Infinity Subsequence). *Let (x_n) be a real sequence. If (x_n) is not bounded above, then there exists a subsequence $(x_{\sigma(k)})$ such that $x_{\sigma(k)} \rightarrow +\infty$.*

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned} \left[\forall M \in \mathbb{R} \exists n \in \mathbb{N} (M < x_n) \right] \\ \implies \exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \left(\forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \wedge \forall M \in \mathbb{R} \exists K \in \mathbb{N} \forall k \geq K (M < x_{\sigma(k)}) \right). \end{aligned}$$

Remark (Definition predicate reading).

$$\neg \text{BoundedAboveSequence}(x_n) \implies \exists \sigma (\text{StrictlyIncreasing}(\sigma) \wedge \text{DivergesToPositiveInfinity}(x_{\sigma(k)})).$$

Remark (Interpretation). Unboundedness above may be sparse. This theorem says the high values can still be selected in increasing index order to form a subsequence that escapes to $+\infty$.

Remark (Dependencies).

- [Sequence Bounded Above](#)
- [Subsequence](#)
- [Divergence to Positive Infinity](#)

Theorem (Unbounded Below Sequence Has a Negative-Infinity Subsequence)

Theorem 6.7.8 (Unbounded Below Sequence Has a Negative-Infinity Subsequence). *Let (x_n) be a real sequence. If (x_n) is not bounded below, then there exists a subsequence $(x_{\sigma(k)})$ such that $x_{\sigma(k)} \rightarrow -\infty$.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\left[\forall M \in \mathbb{R} \exists n \in \mathbb{N} (x_n < M) \right] \\ \implies \exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \left(\forall k, \ell \in \mathbb{N} (k < \ell \implies \sigma(k) < \sigma(\ell)) \wedge \forall M \in \mathbb{R} \exists K \in \mathbb{N} \forall k \geq K (x_{\sigma(k)} < M) \right).$$

Remark (Definition predicate reading).

$$\neg \text{BoundedBelowSequence}(x_n) \implies \exists \sigma (\text{StrictlyIncreasing}(\sigma) \wedge \text{DivergesToNegativeInfinity}(x_{\sigma(k)})).$$

Remark (Interpretation). Unboundedness below can be converted into a selected tail that escapes past every lower threshold. The proof is the order-reversed version of the positive-infinity subsequence construction.

Remark (Dependencies).

- [Sequence Bounded Below](#)
- [Subsequence](#)
- [Divergence to Negative Infinity](#)

Theorem (Bounded Divergence Produces Two Subsequential Limits)

Theorem 6.7.9 (Bounded Divergence Produces Two Subsequential Limits). *Let (x_n) be a bounded real sequence. If (x_n) is divergent, then there exist $L, K \in \mathbb{R}$ with $L \neq K$ such that L and K are subsequential limits of (x_n) .*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\left[\exists M > 0 \forall n \in \mathbb{N} (|x_n| \leq M) \right] \wedge \left[\forall A \in \mathbb{R} \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N (|x_n - A| \geq \varepsilon) \right] \\ \implies \exists L, K \in \mathbb{R} (L \neq K \wedge L \text{ is a subsequential limit of } (x_n) \wedge K \text{ is a subsequential limit of } (x_n)).$$

Remark (Definition predicate reading).

$$\text{BoundedSequence}(x_n) \wedge \text{DivergentSequence}(x_n) \implies \exists L, K \in \mathbb{R} (L \neq K \wedge \text{SubsequentialLimit}(x_n, L) \wedge \text{SubsequentialLimit}(x_n, K)).$$

Remark (Interpretation). Bounded divergence cannot be explained by escape to infinity. In the real line, compactness forces convergent subsequences, and divergence forces at least two incompatible subsequential limits.

Remark (Dependencies).

- [Bounded Sequence](#)
- [Divergent Sequence](#)
- [Subsequential Limit](#)
- [Bolzano-Weierstrass Theorem for Sequences](#)
- [Subsequences Preserve Limits](#)

6.8 Liminf and Limsup

Liminf–Limsup — Quick Reference

Concept	Meaning
Tail supremum sequence	Supremum of each tail
Tail infimum sequence	Infimum of each tail
Limit superior	Limit of tail suprema
Limit inferior	Limit of tail infima
Liminf below limsup	Lower envelope never exceeds upper envelope
Convergence criterion	Convergence iff liminf equals limsup
Largest subsequential limit	Limsup is the top cluster value
Smallest subsequential limit	Liminf is the bottom cluster value
Oscillation criterion	Gap detects nonconvergence
Eventual-order squeeze	Squeeze estimates transfer to limsup and liminf

Liminf and Limsup

The limit superior and limit inferior measure the eventual upper and lower envelopes of a bounded sequence. Instead of asking whether all late terms settle to one number, they

ask how high and how low the tails continue to reach. This makes them natural tools for detecting convergence, oscillation, and subsequential limits.

Definition (Tail Supremum Sequence)

Definition 6.8.1 (Tail Supremum Sequence). Let (x_n) be a bounded real sequence. The **tail supremum sequence** of (x_n) is the real sequence (s_n) defined by

$$s_n := \sup\{x_k : k \geq n\} \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$\forall n \in \mathbb{N} \left(s_n = \sup\{x_k : k \geq n\} \right).$$

Remark (Interpretation). The term s_n records the least ceiling of the n -th tail. As n grows, earlier terms are discarded, so the tail has fewer opportunities to reach large values.

Remark (Dependencies).

- [Bounded Sequence](#)
- [Supremum](#)
- [M-Tail of a Sequence](#)

Definition (Tail Infimum Sequence)

Definition 6.8.2 (Tail Infimum Sequence). Let (x_n) be a bounded real sequence. The **tail infimum sequence** of (x_n) is the real sequence (i_n) defined by

$$i_n := \inf\{x_k : k \geq n\} \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$\forall n \in \mathbb{N} \left(i_n = \inf\{x_k : k \geq n\} \right).$$

Remark (Interpretation). The term i_n records the greatest floor of the n -th tail. As earlier terms are discarded, the tail loses opportunities to reach small values.

Remark (Dependencies).

- [Bounded Sequence](#)
- [Infimum](#)
- [M-Tail of a Sequence](#)

Definition (Limit Superior of a Sequence)

Definition 6.8.3 (Limit Superior of a Sequence). Let (x_n) be a bounded real sequence, and let (s_n) be its tail supremum sequence. The **limit superior** of (x_n) is

$$\limsup_{n \rightarrow \infty} x_n := \lim_{n \rightarrow \infty} s_n.$$

Remark (Standard quantified statement).

$$\limsup_{n \rightarrow \infty} x_n = L \iff \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|s_n - L| < \varepsilon).$$

Remark (Definition predicate reading).

$$\text{LimsupSequence}(x_n, L) := \text{ConvergentSequence}(s_n, L).$$

Remark (Interpretation). The limit superior is the eventual upper envelope of the sequence. It is the number to which the ceilings of the tails descend.

Remark (Dependencies).

- [Tail Supremum Sequence](#)
- [Convergence of a Sequence](#)

Definition (Limit Inferior of a Sequence)

Definition 6.8.4 (Limit Inferior of a Sequence). Let (x_n) be a bounded real sequence, and let (i_n) be its tail infimum sequence. The **limit inferior** of (x_n) is

$$\liminf_{n \rightarrow \infty} x_n := \lim_{n \rightarrow \infty} i_n.$$

Remark (Standard quantified statement).

$$\liminf_{n \rightarrow \infty} x_n = L \iff \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|i_n - L| < \varepsilon).$$

Remark (Definition predicate reading).

$$\text{LiminfSequence}(x_n, L) := \text{ConvergentSequence}(i_n, L).$$

Remark (Interpretation). The limit inferior is the eventual lower envelope of the sequence. It is the number to which the floors of the tails ascend.

Remark (Dependencies).

- [Tail Infimum Sequence](#)
- [Convergence of a Sequence](#)

Theorem (Tail Suprema Are Decreasing)

Theorem 6.8.5 (Tail Suprema Are Decreasing). *Let (x_n) be a bounded real sequence, and let (s_n) be its tail supremum sequence. Then (s_n) is decreasing.*

Go to proof.

Remark (Standard quantified statement).

$$\forall n \in \mathbb{N} (s_{n+1} \leq s_n).$$

Remark (Definition predicate reading).

$$\text{MonotoneDecreasingSequence}(s_n).$$

Remark (Interpretation). The $(n + 1)$ -st tail is contained in the n -th tail. Passing from a set to a smaller subset cannot raise its supremum.

Remark (Dependencies).

- [Tail Supremum Sequence](#)
- [Supremum Is Monotone Under Inclusion](#)

Theorem (Tail Infima Are Increasing)

Theorem 6.8.6 (Tail Infima Are Increasing). *Let (x_n) be a bounded real sequence, and let (i_n) be its tail infimum sequence. Then (i_n) is increasing.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall n \in \mathbb{N} \ (i_n \leq i_{n+1}).$$

Remark (Definition predicate reading).

$$\text{MonotoneIncreasingSequence}(i_n).$$

Remark (Interpretation). The $(n + 1)$ -st tail is contained in the n -th tail. Passing from a set to a smaller subset cannot lower its infimum.

Remark (Dependencies).

- [Tail Infimum Sequence](#)
- [Infimum](#)

Theorem (Liminf Is Below Limsup)

Theorem 6.8.7 (Liminf Is Below Limsup). *Let (x_n) be a bounded real sequence. Then*

$$\liminf_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} x_n.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\liminf_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} x_n.$$

Remark (Interpretation). Every tail infimum lies below every corresponding tail supremum. Passing to the limits preserves this order.

Remark (Dependencies).

- [Limit Inferior of a Sequence](#)
- [Limit Superior of a Sequence](#)
- [Constant Comparison for Sequence Limits](#)

Theorem (Convergence iff Liminf Equals Limsup)

Theorem 6.8.8 (Convergence iff Liminf Equals Limsup). *Let (x_n) be a bounded real sequence, and let $L \in \mathbb{R}$. Then $x_n \rightarrow L$ if and only if*

$$\liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n = L.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$x_n \rightarrow L \iff \liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n = L.$$

Remark (Definition predicate reading).

$$\text{ConvergentSequence}(x_n, L) \iff (\text{LiminfSequence}(x_n, L) \wedge \text{LimsupSequence}(x_n, L)).$$

Remark (Interpretation). Convergence occurs exactly when the eventual lower envelope and eventual upper envelope collapse to the same number.

Remark (Dependencies).

- [Limit Inferior of a Sequence](#)
- [Limit Superior of a Sequence](#)
- [Convergence of a Sequence](#)
- [Sequence Squeeze Theorem](#)

Theorem (Limsup Is the Largest Subsequential Limit)

Theorem 6.8.9 (Limsup Is the Largest Subsequential Limit). *Let (x_n) be a bounded real sequence, and let $S = \limsup_{n \rightarrow \infty} x_n$. Then S is a subsequential limit of (x_n) , and every subsequential limit of (x_n) is less than or equal to S .*

[Go to proof.](#)

Remark (Standard quantified statement).

$$S = \limsup_{n \rightarrow \infty} x_n \implies \left[S \text{ is a subsequential limit of } (x_n) \wedge \forall L \in \mathbb{R} \left(L \text{ is a subsequential limit of } (x_n) \implies L \leq S \right) \right].$$

Remark (Definition predicate reading).

$\text{LimsupSequence}(x_n, S) \implies (\text{SubsequentialLimit}(x_n, S) \wedge \forall L \in \mathbb{R} (\text{SubsequentialLimit}(x_n, L) \implies L \leq S)).$

Remark (Interpretation). The limit superior is not merely an upper bound on subsequential behavior. It is actually attained as a subsequential limit, and no subsequential limit lies above it.

Remark (Dependencies).

- [Limit Superior of a Sequence](#)
- [Subsequential Limit](#)
- [Bolzano-Weierstrass Theorem for Sequences](#)

Theorem (Liminf Is the Smallest Subsequential Limit)

Theorem 6.8.10 (Liminf Is the Smallest Subsequential Limit). *Let (x_n) be a bounded real sequence, and let $I = \liminf_{n \rightarrow \infty} x_n$. Then I is a subsequential limit of (x_n) , and every subsequential limit of (x_n) is greater than or equal to I .*

[Go to proof.](#)

Remark (Standard quantified statement).

$$I = \liminf_{n \rightarrow \infty} x_n \implies \left[I \text{ is a subsequential limit of } (x_n) \wedge \forall L \in \mathbb{R} (L \text{ is a subsequential limit of } (x_n) \implies I \leq L) \right].$$

Remark (Definition predicate reading).

$\text{LiminfSequence}(x_n, I) \implies (\text{SubsequentialLimit}(x_n, I) \wedge \forall L \in \mathbb{R} (\text{SubsequentialLimit}(x_n, L) \implies I \leq L)).$

Remark (Interpretation). The limit inferior is attained as a subsequential limit and no subsequential limit lies below it.

Remark (Dependencies).

- [Limit Inferior of a Sequence](#)
- [Subsequential Limit](#)
- [Bolzano-Weierstrass Theorem for Sequences](#)

Theorem (Oscillation Criterion via Liminf and Limsup)

Theorem 6.8.11 (Oscillation Criterion via Liminf and Limsup). *Let (x_n) be a bounded real sequence. Then*

$$\liminf_{n \rightarrow \infty} x_n < \limsup_{n \rightarrow \infty} x_n$$

if and only if (x_n) has two distinct subsequential limits.

[Go to proof.](#)

Remark (Standard quantified statement).

$$\liminf_{n \rightarrow \infty} x_n < \limsup_{n \rightarrow \infty} x_n \\ \iff \exists L, K \in \mathbb{R} (L \neq K \wedge L \text{ is a subsequential limit of } (x_n) \wedge K \text{ is a subsequential limit of } (x_n)).$$

Remark (Definition predicate reading).

$$\exists I, S \in \mathbb{R} (\text{LiminfSequence}(x_n, I) \wedge \text{LimsupSequence}(x_n, S) \wedge I < S) \\ \iff \exists L, K \in \mathbb{R} (L \neq K \wedge \text{SubsequentialLimit}(x_n, L) \wedge \text{SubsequentialLimit}(x_n, K)).$$

Remark (Interpretation). A strict gap between the lower and upper envelopes is exactly persistent oscillation in bounded sequences. The sequence keeps returning near at least two incompatible limiting levels.

Remark (Dependencies).

- [Limsup Is the Largest Subsequential Limit](#)
- [Liminf Is the Smallest Subsequential Limit](#)
- [Two Subsequential Limits Force Divergence](#)

Theorem (Limsup Comparison Under Eventual Order)

Theorem 6.8.12 (Limsup Comparison Under Eventual Order). *Let (x_n) and (y_n) be bounded real sequences. If there exists $N \in \mathbb{N}$ such that $x_n \leq y_n$ for every $n \geq N$, then*

$$\limsup_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} y_n.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\exists N \in \mathbb{N} \forall n \geq N (x_n \leq y_n) \\ \implies \limsup_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} y_n.$$

Remark (Interpretation). Eventual order of sequences passes to eventual order of their upper tail envelopes. Taking the limiting ceiling preserves that comparison.

Remark (Dependencies).

- [Limit Superior of a Sequence](#)
- [Limit Preserves Eventual Order](#)

Theorem (Liminf Comparison Under Eventual Order)

Theorem 6.8.13 (Liminf Comparison Under Eventual Order). *Let (x_n) and (y_n) be bounded real sequences. If there exists $N \in \mathbb{N}$ such that $x_n \leq y_n$ for every $n \geq N$, then*

$$\liminf_{n \rightarrow \infty} x_n \leq \liminf_{n \rightarrow \infty} y_n.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \exists N \in \mathbb{N} \forall n \geq N (x_n \leq y_n) \\ \implies \liminf_{n \rightarrow \infty} x_n \leq \liminf_{n \rightarrow \infty} y_n. \end{aligned}$$

Remark (Interpretation). Eventual order also passes to the lower tail envelopes. If one sequence is eventually below another, its eventual floor cannot sit above the other's eventual floor.

Remark (Dependencies).

- [Limit Inferior of a Sequence](#)
- [Limit Preserves Eventual Order](#)

Theorem (Limsup Squeeze Under Eventual Order)

Theorem 6.8.14 (Limsup Squeeze Under Eventual Order). *Let (a_n) , (x_n) , and (b_n) be bounded real sequences. If there exists $N \in \mathbb{N}$ such that*

$$a_n \leq x_n \leq b_n \quad \text{for every } n \geq N,$$

then

$$\limsup_{n \rightarrow \infty} a_n \leq \limsup_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} b_n.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \exists N \in \mathbb{N} \forall n \geq N (a_n \leq x_n \leq b_n) \\ \implies \limsup_{n \rightarrow \infty} a_n \leq \limsup_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} b_n. \end{aligned}$$

Remark (Interpretation). If one sequence is eventually trapped between two others, then its eventual upper envelope is trapped between their eventual upper envelopes.

Remark (Dependencies).

- [Limsup Comparison Under Eventual Order](#)

Theorem (Liminf Squeeze Under Eventual Order)

Theorem 6.8.15 (Liminf Squeeze Under Eventual Order). *Let (a_n) , (x_n) , and (b_n) be bounded real sequences. If there exists $N \in \mathbb{N}$ such that*

$$a_n \leq x_n \leq b_n \quad \text{for every } n \geq N,$$

then

$$\liminf_{n \rightarrow \infty} a_n \leq \liminf_{n \rightarrow \infty} x_n \leq \liminf_{n \rightarrow \infty} b_n.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \exists N \in \mathbb{N} \forall n \geq N (a_n \leq x_n \leq b_n) \\ \implies \liminf_{n \rightarrow \infty} a_n \leq \liminf_{n \rightarrow \infty} x_n \leq \liminf_{n \rightarrow \infty} b_n. \end{aligned}$$

Remark (Interpretation). The lower-envelope version of the squeeze principle says that eventual two-sided order traps the eventual floors as well as the eventual ceilings.

Remark (Dependencies).

- [Liminf Comparison Under Eventual Order](#)

Remark (Illustrative cases). For the alternating sequence $x_n = (-1)^n$, the tail suprema are constantly 1 and the tail infima are constantly -1 , so

$$\limsup_{n \rightarrow \infty} (-1)^n = 1, \quad \liminf_{n \rightarrow \infty} (-1)^n = -1.$$

For any convergent sequence $x_n \rightarrow L$, both envelopes collapse to L .

6.9 Order-Theoretic Limit Constructions

Order-Theoretic Limits — Quick Reference

Statement	Role
Increasing sequence limit	Limit realized as a supremum
Decreasing sequence limit	Limit realized as an infimum
Tail extremals converge	Tail suprema and infima are monotone
Bounded sequence limsup/liminf	Bounded sequences have asymptotic envelopes

Order-Theoretic Limit Constructions

Completeness lets order data produce limits. An increasing sequence bounded above converges to the least ceiling of its range; a decreasing sequence bounded below converges to the greatest

floor of its range. Applying these same ideas to tail suprema and tail infima produces limsup and liminf for every bounded sequence.

Theorem (Increasing Sequence Limit as Supremum)

Theorem 6.9.1 (Increasing Sequence Limit as Supremum). *Let (x_n) be an increasing real sequence that is bounded above. Then*

$$\lim_{n \rightarrow \infty} x_n = \sup\{x_n : n \in \mathbb{N}\}.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \forall(x_n) \subseteq \mathbb{R} \left(\left[\forall n \in \mathbb{N} (x_n \leq x_{n+1}) \right] \wedge \left[\exists M \in \mathbb{R} \forall n \in \mathbb{N} (x_n \leq M) \right] \right) \\ \implies \lim_{n \rightarrow \infty} x_n = \sup\{x_n : n \in \mathbb{N}\}. \end{aligned}$$

Remark (Theorem predicate reading).

$\text{MonotoneIncreasingSequence}(x_n) \wedge \text{BoundedAboveSequence}(x_n) \implies \text{ConvergentSequence}(x_n, \sup\{x_n : n \in \mathbb{N}\})$

Remark (Interpretation). The supremum is the only possible destination: all terms lie below it, and the defining approximation property of the supremum forces the sequence eventually above every lower tolerance.

Remark (Dependencies).

- [Increasing Sequence](#)
- [Sequence Bounded Above](#)
- [Supremum](#)
- [Monotone Convergence Theorem](#)

Theorem (Decreasing Sequence Limit as Infimum)

Theorem 6.9.2 (Decreasing Sequence Limit as Infimum). *Let (x_n) be a decreasing real sequence that is bounded below. Then*

$$\lim_{n \rightarrow \infty} x_n = \inf\{x_n : n \in \mathbb{N}\}.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \forall(x_n) \subseteq \mathbb{R} \left(\left[\forall n \in \mathbb{N} (x_{n+1} \leq x_n) \right] \wedge \left[\exists m \in \mathbb{R} \forall n \in \mathbb{N} (m \leq x_n) \right] \right) \\ \implies \lim_{n \rightarrow \infty} x_n = \inf\{x_n : n \in \mathbb{N}\}. \end{aligned}$$

Remark (Theorem predicate reading).

$\text{MonotoneDecreasingSequence}(x_n) \wedge \text{BoundedBelowSequence}(x_n) \implies \text{ConvergentSequence}(x_n, \inf\{x_n : n \in \mathbb{N}\})$

Remark (Interpretation). The infimum is the decreasing analogue of the supremum target: the sequence falls toward its greatest lower barrier.

Remark (Dependencies).

- [Decreasing Sequence](#)
- [Sequence Bounded Below](#)
- [Infimum](#)
- [Monotone Convergence Theorem](#)

Theorem (Tail Suprema and Infima Converge)

Theorem 6.9.3 (Tail Suprema and Infima Converge). *Let (x_n) be a bounded real sequence. Define*

$$s_n = \sup\{x_k : k \geq n\}, \quad i_n = \inf\{x_k : k \geq n\}.$$

Then (s_n) and (i_n) converge.

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \left((x_n) \text{ is bounded} \implies \exists S, I \in \mathbb{R} \left[\lim_{n \rightarrow \infty} \sup\{x_k : k \geq n\} = S \wedge \lim_{n \rightarrow \infty} \inf\{x_k : k \geq n\} = I \right] \right).$$

Remark (Theorem predicate reading).

$\text{BoundedSequence}(x_n) \implies \exists S, I \in \mathbb{R} \left(\text{ConvergentSequence}(\sup\{x_k : k \geq n\}, S) \wedge \text{ConvergentSequence}(\inf\{x_k : k \geq n\}, I) \right)$

Remark (Interpretation). Tail suprema decrease because later tails contain fewer terms; tail infima increase for the same reason. Boundedness supplies the opposite bound needed for monotone convergence.

Remark (Dependencies).

- [Tail Supremum Sequence](#)
- [Tail Infimum Sequence](#)
- [Tail Suprema Are Decreasing](#)
- [Tail Infima Are Increasing](#)
- [Monotone Convergence Theorem](#)

Theorem (Bounded Sequence Has Limsup and Liminf)

Theorem 6.9.4 (Bounded Sequence Has Limsup and Liminf). *Every bounded real sequence has a limit superior and a limit inferior.*

Go to proof.

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \left((x_n) \text{ is bounded} \implies \exists S, I \in \mathbb{R} \left(S = \limsup_{n \rightarrow \infty} x_n \wedge I = \liminf_{n \rightarrow \infty} x_n \right) \right).$$

Remark (Theorem predicate reading).

$$\text{BoundedSequence}(x_n) \implies \exists S, I \in \mathbb{R} \left(\text{LimsupSequence}(x_n, S) \wedge \text{LiminfSequence}(x_n, I) \right).$$

Remark (Interpretation). For bounded sequences, limsup and liminf are not extra assumptions. They are guaranteed order-theoretic limits of the tail envelopes.

Remark (Dependencies).

- [Limit Superior of a Sequence](#)
- [Limit Inferior of a Sequence](#)
- [Tail Suprema and Infima Converge](#)

6.10 Cluster Values of Sequences

Cluster Values — Quick Reference

Concept	Meaning
Cluster value	Value approached along a subsequence
Cluster equals subsequential limit	Converts frequent approach to subsequences
Bounded sequences have cluster values	Compactness produces accumulation
Extremal cluster values	Limsup and liminf are the outer cluster values

Cluster Values of Sequences

A convergent sequence has only one limiting value. A bounded divergent sequence may still have values that it approaches along selected subsequences. Cluster values name this repeated limiting behavior without requiring the entire sequence to converge.

Definition (Cluster Value of a Sequence)

Definition 6.10.1 (Cluster Value of a Sequence). Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. The number L is a **cluster value** of (x_n) if for every $\varepsilon > 0$ and every $N \in \mathbb{N}$ there exists $n \geq N$ such that

$$|x_n - L| < \varepsilon.$$

Remark (Standard quantified statement).

L is a cluster value of $(x_n) \iff \forall \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N (|x_n - L| < \varepsilon)$.

Remark (Definition predicate reading).

$\text{SequenceClusterValue}(x_n, L) := \forall \varepsilon > 0 \forall N \in \mathbb{N} \exists n \geq N (|x_n - L| < \varepsilon)$.

Remark (Negated quantified statement).

L is not a cluster value of $(x_n) \iff \exists \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - L| \geq \varepsilon)$.

Remark (Interpretation). No matter how far out in the sequence one looks, some later term returns inside every neighborhood of L .

Remark (Dependencies).

- [Real Sequence](#)

Theorem (Cluster Values Are Subsequential Limits)

Theorem 6.10.2 (Cluster Values Are Subsequential Limits). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. Then L is a cluster value of (x_n) if and only if L is a subsequential limit of (x_n) .*

[Go to proof.](#)

Remark (Standard quantified statement).

$\forall (x_n) \subseteq \mathbb{R} \forall L \in \mathbb{R} \left(L \text{ is a cluster value of } (x_n) \iff L \text{ is a subsequential limit of } (x_n) \right)$.

Remark (Theorem predicate reading).

$\text{SequenceClusterValue}(x_n, L) \iff \text{SubsequentialLimit}(x_n, L)$.

Remark (Interpretation). The neighborhood formulation and the subsequence formulation describe the same phenomenon: infinitely many opportunities to get arbitrarily close to L .

Remark (Dependencies).

- [Cluster Value of a Sequence](#)
- [Subsequential Limit](#)
- [Frequent Properties Yield Subsequences](#)

Theorem (Bounded Sequences Have Cluster Values)

Theorem 6.10.3 (Bounded Sequences Have Cluster Values). *Every bounded real sequence has at least one cluster value.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall(x_n) \subseteq \mathbb{R} \left([\exists M > 0 \forall n \in \mathbb{N} (|x_n| \leq M)] \implies \exists L \in \mathbb{R} (L \text{ is a cluster value of } (x_n)) \right).$$

Remark (Theorem predicate reading).

$$\text{BoundedSequence}(x_n) \implies \exists L \in \mathbb{R} \text{ SequenceClusterValue}(x_n, L).$$

Remark (Interpretation). Boundedness traps the sequence in a finite interval, and Bolzano-Weierstrass extracts a convergent subsequence from that trap.

Remark (Dependencies).

- [Bounded Sequence](#)
- [Bolzano-Weierstrass Theorem for Sequences](#)
- [Cluster Values Are Subsequential Limits](#)

Theorem (Limsup and Liminf Are Extremal Cluster Values)

Theorem 6.10.4 (Limsup and Liminf Are Extremal Cluster Values). *Let (x_n) be a bounded real sequence. Then $\limsup_{n \rightarrow \infty} x_n$ is the largest cluster value of (x_n) , and $\liminf_{n \rightarrow \infty} x_n$ is the smallest cluster value of (x_n) .*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall(x_n) \subseteq \mathbb{R} \left((x_n) \text{ is bounded} \implies \left[\limsup_{n \rightarrow \infty} x_n \text{ is a cluster value of } (x_n) \wedge \forall L \in \mathbb{R} (L \text{ is a cluster value of } (x_n) \implies L \leq \limsup_{n \rightarrow \infty} x_n) \right] \wedge \left[\liminf_{n \rightarrow \infty} x_n \text{ is a cluster value of } (x_n) \wedge \forall L \in \mathbb{R} (L \text{ is a cluster value of } (x_n) \implies \liminf_{n \rightarrow \infty} x_n \leq L) \right] \right).$$

Remark (Theorem predicate reading).

$$\text{BoundedSequence}(x_n) \implies$$

$$\text{SequenceClusterValue}(x_n, \limsup_{n \rightarrow \infty} x_n) \wedge \forall L \in \mathbb{R} (\text{SequenceClusterValue}(x_n, L) \implies L \leq \limsup_{n \rightarrow \infty} x_n),$$

$$\text{BoundedSequence}(x_n) \implies$$

$$\text{SequenceClusterValue}(x_n, \liminf_{n \rightarrow \infty} x_n) \wedge \forall L \in \mathbb{R} (\text{SequenceClusterValue}(x_n, L) \implies \liminf_{n \rightarrow \infty} x_n \leq L).$$

Remark (Interpretation). Limsup and liminf describe the top and bottom edges of the sequence's subsequential behavior.

Remark (Dependencies).

- [Limit Superior of a Sequence](#)

- [Limit Inferior of a Sequence](#)
- [Limsup Is the Largest Subsequential Limit](#)
- [Liminf Is the Smallest Subsequential Limit](#)
- [Cluster Values Are Subsequential Limits](#)

6.11 Cauchy

Cauchy Sequences

Cauchy sequences measure internal stabilization. Instead of comparing the terms to an already named limit, the Cauchy condition compares sufficiently late terms to each other. Over the real numbers this internal criterion is equivalent to convergence, and that equivalence is one of the central sequence forms of completeness.

Definition (Cauchy Sequence)

Definition 6.11.1 (Cauchy Sequence). Let (x_n) be a real sequence. The sequence (x_n) is a **Cauchy sequence** if

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N (|x_m - x_n| < \varepsilon).$$

Remark (Standard quantified statement).

$$\begin{aligned} (x_n) \text{ is Cauchy} \\ \iff \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N (|x_m - x_n| < \varepsilon). \end{aligned}$$

Remark (Definition predicate reading).

$$\text{CauchySequence}(x_n) := \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N (|x_m - x_n| < \varepsilon).$$

Remark (Negated quantified statement).

$$\begin{aligned} (x_n) \text{ is not Cauchy} \\ \iff \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists m, n \geq N (|x_m - x_n| \geq \varepsilon). \end{aligned}$$

Remark (Negation predicate reading).

$$\neg \text{CauchySequence}(x_n) \iff \exists \varepsilon > 0 \forall N \in \mathbb{N} \exists m, n \geq N (|x_m - x_n| \geq \varepsilon).$$

Remark (Interpretation). The Cauchy condition says that the tails of the sequence eventually have arbitrarily small internal spread. It does not name a candidate limit; it only asserts that late terms become mutually close.

Remark (Dependencies).

- [Real Sequence](#)

Theorem (Convergent Sequences Are Cauchy)

Theorem 6.11.2 (Convergent Sequences Are Cauchy). *Let (x_n) be a real sequence. If (x_n) converges, then (x_n) is Cauchy.*

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned} \forall(x_n) \subseteq \mathbb{R} \\ \left((\exists L \in \mathbb{R} \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - L| < \varepsilon)) \right. \\ \left. \implies \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N (|x_m - x_n| < \varepsilon) \right). \end{aligned}$$

Remark (Definition predicate reading).

$$\forall(x_n) \subseteq \mathbb{R} \left(\exists L \in \mathbb{R} \text{ConvergentSequence}(x_n, L) \implies \text{CauchySequence}(x_n) \right).$$

Remark (Interpretation). Once every sufficiently late term is close to the same limit, any two sufficiently late terms are close to each other. This theorem is the easy half of the Cauchy criterion.

Remark (Dependencies).

- [Convergence of a Sequence](#)
- [Cauchy Sequence](#)

Theorem (Cauchy Sequences Are Bounded)

Theorem 6.11.3 (Cauchy Sequences Are Bounded). *Every Cauchy real sequence is bounded.*

Go to proof.

Remark (Standard quantified statement).

$$\begin{aligned} \forall(x_n) \subseteq \mathbb{R} \\ \left((\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N (|x_m - x_n| < \varepsilon)) \right. \\ \left. \implies \exists M > 0 \forall n \in \mathbb{N} (|x_n| \leq M) \right). \end{aligned}$$

Remark (Definition predicate reading).

$$\forall(x_n) \subseteq \mathbb{R} \left(\text{CauchySequence}(x_n) \implies \text{BoundedSequence}(x_n) \right).$$

Remark (Interpretation). A Cauchy sequence has one tightly controlled tail. The finitely many terms before that tail can be absorbed into a single larger bound.

Remark (Dependencies).

- [Cauchy Sequence](#)
- [Bounded Sequence](#)

Theorem (Cauchy Sequence with a Convergent Subsequence)

Theorem 6.11.4 (Cauchy Sequence with a Convergent Subsequence). *Let (x_n) be a Cauchy real sequence. If a subsequence of (x_n) converges to $L \in \mathbb{R}$, then (x_n) converges to L .*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} & \forall (x_n) \subseteq \mathbb{R} \, \forall L \in \mathbb{R} \\ & \left(\left[\forall \varepsilon > 0 \, \exists N \in \mathbb{N} \, \forall m, n \geq N \, (|x_m - x_n| < \varepsilon) \right] \right. \\ & \quad \wedge \left[\exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \, (\forall k, \ell \in \mathbb{N} \, (k < \ell \implies \sigma(k) < \sigma(\ell))) \wedge \forall \varepsilon > 0 \, \exists K \in \mathbb{N} \, \forall k \geq K \, (|x_{\sigma(k)} - L| < \varepsilon) \right] \\ & \quad \implies \forall \varepsilon > 0 \, \exists N \in \mathbb{N} \, \forall n \geq N \, (|x_n - L| < \varepsilon) \Big). \end{aligned}$$

Remark (Definition predicate reading).

$$\begin{aligned} & \forall (x_n) \subseteq \mathbb{R} \, \forall L \in \mathbb{R} \\ & \left(\text{CauchySequence}(x_n) \wedge \text{SubsequentialLimit}(x_n, L) \implies \text{ConvergentSequence}(x_n, L) \right). \end{aligned}$$

Remark (Interpretation). The Cauchy condition prevents the rest of the tail from wandering away from any convergent subsequence. A single subsequential limit therefore determines the limit of the whole sequence.

Remark (Dependencies).

- [Cauchy Sequence](#)
- [Subsequence](#)
- [Subsequential Limit](#)
- [Convergence of a Sequence](#)

Theorem (Cauchy Criterion for Real Sequences)

Theorem 6.11.5 (Cauchy Criterion for Real Sequences). *Let (x_n) be a real sequence. Then (x_n) converges if and only if (x_n) is Cauchy.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \forall (x_n) \subseteq \mathbb{R} \\ \left(\exists L \in \mathbb{R} \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - L| < \varepsilon) \right. \\ \left. \iff \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N (|x_m - x_n| < \varepsilon) \right). \end{aligned}$$

Remark (Definition predicate reading).

$$\forall (x_n) \subseteq \mathbb{R} \left(\exists L \in \mathbb{R} \text{ConvergentSequence}(x_n, L) \iff \text{CauchySequence}(x_n) \right).$$

Remark (Interpretation). This theorem is the sequence form of completeness of $\mathbb{R}$. In an incomplete ambient space, a sequence may be internally stable without converging inside that space.

Remark (Dependencies).

- [Convergent Sequences Are Cauchy](#)
- [Cauchy Sequences Are Bounded](#)
- [Bolzano-Weierstrass Theorem for Sequences](#)
- [Cauchy Sequence with a Convergent Subsequence](#)

Theorem (Cauchy Criterion via Tails)

Theorem 6.11.6 (Cauchy Criterion via Tails). *Let (x_n) be a real sequence. Then (x_n) is Cauchy if and only if for every $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that the N -tail has pairwise term distances less than ε .*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} (x_n) \text{ is Cauchy} \\ \iff \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall p, q \in \mathbb{N} (|x_{N+p} - x_{N+q}| < \varepsilon). \end{aligned}$$

Remark (Definition predicate reading).

$$\text{CauchySequence}(x_n) \iff \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall p, q \in \mathbb{N} (|x_{N+p} - x_{N+q}| < \varepsilon).$$

Remark (Interpretation). The Cauchy condition is a tail property. Reindexing a sufficiently late tail does not change the substance of the condition; it only changes the names of the indices.

Remark (Dependencies).

- [Cauchy Sequence](#)

- [M-Tail of a Sequence](#)

Theorem (Cauchy Tail Diameter Criterion)

Theorem 6.11.7 (Cauchy Tail Diameter Criterion). *Let (x_n) be a real sequence. Then (x_n) is Cauchy if and only if every sufficiently late tail has arbitrarily small pairwise diameter.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} (x_n) \text{ is Cauchy} \\ \iff \forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall N \geq N_0 \\ \forall m, n \geq N (|x_m - x_n| < \varepsilon). \end{aligned}$$

Remark (Definition predicate reading).

$$\text{CauchySequence}(x_n) \iff \forall \varepsilon > 0 \exists N_0 \in \mathbb{N} \forall N \geq N_0 \forall m, n \geq N (|x_m - x_n| < \varepsilon).$$

Remark (Interpretation). The tail-diameter form says that once one tail is small, every later tail is small as well. It records the monotone nature of tail control without introducing additional notation for diameters.

Remark (Dependencies).

- [Cauchy Sequence](#)
- [M-Tail of a Sequence](#)

Theorem (Cauchy Sequences Have Vanishing Successive Differences)

Theorem 6.11.8 (Cauchy Sequences Have Vanishing Successive Differences). *If (x_n) is a Cauchy real sequence, then*

$$|x_{n+1} - x_n| \rightarrow 0.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} \forall (x_n) \subseteq \mathbb{R} \\ \left(\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N (|x_m - x_n| < \varepsilon) \right) \\ \implies \forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_{n+1} - x_n| < \varepsilon). \end{aligned}$$

Remark (Definition predicate reading).

$$\text{CauchySequence}(x_n) \implies \text{NullSequence}(|x_{n+1} - x_n|).$$

Remark (Interpretation). Cauchy control applies to every pair of late indices, so it applies in particular to consecutive late indices. The converse is false in general: small successive steps do not prevent accumulated drift.

Remark (Dependencies).

- [Cauchy Sequence](#)
- [Null Sequence](#)

Theorem (Scalar Multiples of Cauchy Sequences)

Theorem 6.11.9 (Scalar Multiples of Cauchy Sequences). *Let (x_n) be a Cauchy real sequence, and let $\alpha \in \mathbb{R}$. Then (αx_n) is Cauchy.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \, \forall \alpha \in \mathbb{R} \left((x_n) \text{ is Cauchy} \implies (\alpha x_n) \text{ is Cauchy} \right).$$

Remark (Definition predicate reading).

$$\text{CauchySequence}(x_n) \implies \text{CauchySequence}(\alpha x_n).$$

Remark (Interpretation). Multiplying all terms by one fixed real number rescales all pairwise tail distances by the same fixed factor.

Remark (Dependencies).

- [Cauchy Sequence](#)
- [Scalar Multiple of a Sequence](#)

Theorem (Sums of Cauchy Sequences)

Theorem 6.11.10 (Sums of Cauchy Sequences). *Let (x_n) and (y_n) be Cauchy real sequences. Then $(x_n + y_n)$ is Cauchy.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \, \forall (y_n) \subseteq \mathbb{R} \left((x_n) \text{ is Cauchy} \wedge (y_n) \text{ is Cauchy} \implies (x_n + y_n) \text{ is Cauchy} \right).$$

Remark (Definition predicate reading).

$$\text{CauchySequence}(x_n) \wedge \text{CauchySequence}(y_n) \implies \text{CauchySequence}(x_n + y_n).$$

Remark (Interpretation). The triangle inequality splits the tail distance of the sum into the two tail distances already controlled by the original sequences.

Remark (Dependencies).

- [Cauchy Sequence](#)
- [Pointwise Sum of Sequences](#)

Theorem (Differences of Cauchy Sequences)

Theorem 6.11.11 (Differences of Cauchy Sequences). *Let (x_n) and (y_n) be Cauchy real sequences. Then $(x_n - y_n)$ is Cauchy.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall(x_n) \subseteq \mathbb{R} \forall(y_n) \subseteq \mathbb{R} \\ \left((x_n) \text{ is Cauchy} \wedge (y_n) \text{ is Cauchy} \implies (x_n - y_n) \text{ is Cauchy} \right).$$

Remark (Definition predicate reading).

$$\text{CauchySequence}(x_n) \wedge \text{CauchySequence}(y_n) \implies \text{CauchySequence}(x_n - y_n).$$

Remark (Interpretation). The difference theorem is the sum theorem applied after negating one sequence. It is often the form used to compare two approximating sequences.

Remark (Dependencies).

- [Sums of Cauchy Sequences](#)
- [Scalar Multiples of Cauchy Sequences](#)
- [Pointwise Difference of Sequences](#)

Theorem (Linear Combinations of Cauchy Sequences)

Theorem 6.11.12 (Linear Combinations of Cauchy Sequences). *Let (x_n) and (y_n) be Cauchy real sequences, and let $\alpha, \beta \in \mathbb{R}$. Then $(\alpha x_n + \beta y_n)$ is Cauchy.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall(x_n) \subseteq \mathbb{R} \forall(y_n) \subseteq \mathbb{R} \forall \alpha \in \mathbb{R} \forall \beta \in \mathbb{R} \\ \left((x_n) \text{ is Cauchy} \wedge (y_n) \text{ is Cauchy} \implies (\alpha x_n + \beta y_n) \text{ is Cauchy} \right).$$

Remark (Definition predicate reading).

$$\text{CauchySequence}(x_n) \wedge \text{CauchySequence}(y_n) \implies \text{CauchySequence}(\alpha x_n + \beta y_n).$$

Remark (Interpretation). The Cauchy real sequences are closed under finite real linear combinations. This is the additive vector-space part of Cauchy algebra.

Remark (Dependencies).

- [Linear Combination of Real Sequences](#)
- [Scalar Multiples of Cauchy Sequences](#)
- [Sums of Cauchy Sequences](#)

Theorem (Products of Cauchy Sequences)

Theorem 6.11.13 (Products of Cauchy Sequences). *Let (x_n) and (y_n) be Cauchy real sequences. Then $(x_n y_n)$ is Cauchy.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} & \forall (x_n) \subseteq \mathbb{R} \, \forall (y_n) \subseteq \mathbb{R} \\ & \left((x_n) \text{ is Cauchy} \wedge (y_n) \text{ is Cauchy} \implies (x_n y_n) \text{ is Cauchy} \right). \end{aligned}$$

Remark (Definition predicate reading).

$$\text{CauchySequence}(x_n) \wedge \text{CauchySequence}(y_n) \implies \text{CauchySequence}(x_n y_n).$$

Remark (Interpretation). Products require boundedness in addition to tail closeness. The boundedness is supplied automatically because every Cauchy real sequence is bounded.

Remark (Dependencies).

- [Cauchy Sequences Are Bounded](#)
- [Pointwise Product of Sequences](#)

Theorem (Reciprocals of Cauchy Sequences)

Theorem 6.11.14 (Reciprocals of Cauchy Sequences). *Let (x_n) be a Cauchy real sequence. If there exist $c > 0$ and $N_0 \in \mathbb{N}$ such that $|x_n| \geq c$ for every $n \geq N_0$, then $(1/x_n)$ is Cauchy.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} & \forall (x_n) \subseteq \mathbb{R} \\ & \left((x_n) \text{ is Cauchy} \wedge \exists c > 0 \, \exists N_0 \in \mathbb{N} \, \forall n \geq N_0 \, (|x_n| \geq c) \right. \\ & \quad \left. \implies (1/x_n) \text{ is Cauchy} \right). \end{aligned}$$

Remark (Definition predicate reading).

$$\text{CauchySequence}(x_n) \wedge \exists c > 0 \exists N_0 \in \mathbb{N} \forall n \geq N_0 (|x_n| \geq c) \implies \text{CauchySequence}(1/x_n).$$

Remark (Interpretation). The lower bound away from zero prevents division from amplifying small tail differences without control. Without it, reciprocals need not preserve Cauchy behavior.

Remark (Dependencies).

- [Cauchy Sequence](#)
- [Reciprocal Sequence](#)

Theorem (Quotients of Cauchy Sequences)

Theorem 6.11.15 (Quotients of Cauchy Sequences). *Let (x_n) and (y_n) be Cauchy real sequences. If there exist $c > 0$ and $N_0 \in \mathbb{N}$ such that $|y_n| \geq c$ for every $n \geq N_0$, then (x_n/y_n) is Cauchy.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\begin{aligned} & \forall (x_n) \subseteq \mathbb{R} \forall (y_n) \subseteq \mathbb{R} \\ & \left((x_n) \text{ is Cauchy} \wedge (y_n) \text{ is Cauchy} \wedge \exists c > 0 \exists N_0 \in \mathbb{N} \forall n \geq N_0 (|y_n| \geq c) \right. \\ & \quad \left. \implies (x_n/y_n) \text{ is Cauchy} \right). \end{aligned}$$

Remark (Definition predicate reading).

$$\text{CauchySequence}(x_n) \wedge \text{CauchySequence}(y_n) \wedge \exists c > 0 \exists N_0 \in \mathbb{N} \forall n \geq N_0 (|y_n| \geq c) \implies \text{CauchySequence}(x_n/y_n).$$

Remark (Interpretation). The quotient theorem is the product theorem combined with the reciprocal theorem for a denominator eventually bounded away from zero.

Remark (Dependencies).

- [Products of Cauchy Sequences](#)
- [Reciprocals of Cauchy Sequences](#)
- [Pointwise Quotient of Sequences](#)

Theorem (Absolute Values of Cauchy Sequences)

Theorem 6.11.16 (Absolute Values of Cauchy Sequences). *Let (x_n) be a Cauchy real sequence. Then $(|x_n|)$ is Cauchy.*

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall (x_n) \subseteq \mathbb{R} \left((x_n) \text{ is Cauchy} \implies (|x_n|) \text{ is Cauchy} \right).$$

Remark (Definition predicate reading).

$$\text{CauchySequence}(x_n) \implies \text{CauchySequence}(|x_n|).$$

Remark (Interpretation). The reverse triangle inequality bounds the distance between absolute values by the original distance between terms.

Remark (Dependencies).

- [Cauchy Sequence](#)
- [Absolute Value of a Sequence](#)

6.12 Applications of Sequences

Sequence Applications — Quick Reference

Construction	Role
Newton sequence for $\sqrt{2}$	Iterative approximation model
Newton approximation	Converges to $\sqrt{2}$
Factorial partial sums	Series approximants for e
Approximation of e	Factorial sums converge to e
Compound-interest sequence	Classical exponential approximants
Compound-interest approximation	Compound interest converges to e
Decimal truncation sequence	Decimal approximants from below
Decimal truncations	Truncations converge to the target real

Applications of Sequences

The convergence theorems for sequences become numerical tools once the sequence is computable. Newton iteration turns an equation into a convergent recursive approximation; factorial partial sums and compound-interest terms approximate the Euler number; decimal truncations approximate arbitrary real numbers by finite decimals.

Definition (Newton Sequence for $\sqrt{2}$)

Definition 6.12.1 (Newton Sequence for $\sqrt{2}$). The **Newton sequence for $\sqrt{2}$** is the real sequence (x_n) defined by

$$x_1 := \frac{3}{2}, \quad x_{n+1} := \frac{1}{2} \left(x_n + \frac{2}{x_n} \right) \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$x_1 = \frac{3}{2} \\ \wedge \quad \forall n \in \mathbb{N} \left(x_{n+1} = \frac{1}{2} \left(x_n + \frac{2}{x_n} \right) \right).$$

Remark (Interpretation). The recursion is Newton's method applied to the equation $x^2 - 2 = 0$. Each new term averages the current estimate with the correction $2/x_n$.

Remark (Dependencies).

- [Real Sequence](#)

Theorem (Newton Approximation of $\sqrt{2}$)

Theorem 6.12.2 (Newton Approximation of $\sqrt{2}$). *Let (x_n) be the Newton sequence for $\sqrt{2}$. Then (x_n) converges and*

$$\lim_{n \rightarrow \infty} x_n = \sqrt{2}.$$

Go to proof.

Remark (Standard quantified statement).

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|x_n - \sqrt{2}| < \varepsilon).$$

Remark (Definition predicate reading).

$$\text{ConvergentSequence}(x_n, \sqrt{2}).$$

Remark (Interpretation). The Newton sequence is decreasing after the first term and bounded below by $\sqrt{2}$, so monotone convergence supplies a limit. Passing the recursion to the limit identifies the limit as the positive solution of $L^2 = 2$.

Remark (Dependencies).

- [Newton Sequence for \$\sqrt{2}\$](#)
- [Monotone Convergence Theorem](#)
- [Rational Sequence Limit](#)
- [Square Root Sequence](#)

Definition (Factorial Partial Sums)

Definition 6.12.3 (Factorial Partial Sums). The **factorial partial-sum sequence** is the real sequence (s_n) defined by

$$s_n := \sum_{k=0}^n \frac{1}{k!} \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$\forall n \in \mathbb{N} \left(s_n = \sum_{k=0}^n \frac{1}{k!} \right).$$

Remark (Interpretation). The sequence adds the reciprocal factorial terms one at a time. The terms decrease very rapidly, so the partial sums stabilize quickly.

Remark (Dependencies).

- [Real Sequence](#)

Theorem (Factorial Partial Sums Approximate e)

Theorem 6.12.4 (Factorial Partial Sums Approximate e). *Let (s_n) be the factorial partial-sum sequence. Then (s_n) converges to the Euler number e :*

$$\lim_{n \rightarrow \infty} s_n = e.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|s_n - e| < \varepsilon).$$

Remark (Definition predicate reading).

$$\text{ConvergentSequence}(s_n, e).$$

Remark (Interpretation). This is the sequence form of the factorial-series definition of e . In practice the approximation is efficient because the error after n terms is controlled by a rapidly shrinking factorial tail.

Remark (Dependencies).

- [Factorial Partial Sums](#)
- [The Number \$e\$](#)
- [Cauchy Criterion for Real Sequences](#)

Definition (Compound-Interest Sequence)

Definition 6.12.5 (Compound-Interest Sequence). The **compound-interest sequence** is the real sequence (c_n) defined by

$$c_n := \left(1 + \frac{1}{n}\right)^n \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$\forall n \in \mathbb{N} \left(c_n = \left(1 + \frac{1}{n} \right)^n \right).$$

Remark (Interpretation). The term c_n is the value of one unit of principal after one unit of time when interest is compounded n times at nominal rate 1.

Remark (Dependencies).

- [Real Sequence](#)

Theorem (Compound-Interest Approximation of e)

Theorem 6.12.6 (Compound-Interest Approximation of e). *Let (c_n) be the compound-interest sequence. Then*

$$\lim_{n \rightarrow \infty} c_n = e.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|c_n - e| < \varepsilon).$$

Remark (Definition predicate reading).

$$\text{ConvergentSequence}(c_n, e).$$

Remark (Interpretation). The compound-interest approximation gives a second natural sequence converging to the same constant as the factorial partial sums. It is slower numerically, but it explains why e appears in continuous growth.

Remark (Dependencies).

- [Compound-Interest Sequence](#)
- [The Number \$e\$](#)
- [Sequence Squeeze Theorem](#)

Definition (Decimal Truncation Sequence)

Definition 6.12.7 (Decimal Truncation Sequence). Let $\alpha \in \mathbb{R}$. The **decimal truncation sequence** of α is the real sequence (d_n) defined by

$$d_n := \frac{\lfloor 10^n \alpha \rfloor}{10^n} \quad \text{for every } n \in \mathbb{N}.$$

Remark (Standard quantified statement).

$$\forall n \in \mathbb{N} \left(d_n = \frac{\lfloor 10^n \alpha \rfloor}{10^n} \right).$$

Remark (Interpretation). The number d_n is obtained by cutting α off after n decimal places. Truncation is one-sided: it does not round to the nearest decimal.

Remark (Dependencies).

- [Real Sequence](#)

Theorem (Decimal Truncations Converge)

Theorem 6.12.8 (Decimal Truncations Converge). *Let $\alpha \in \mathbb{R}$, and let (d_n) be the decimal truncation sequence of α . Then*

$$\lim_{n \rightarrow \infty} d_n = \alpha.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n \geq N (|d_n - \alpha| < \varepsilon).$$

Remark (Definition predicate reading).

$$\text{ConvergentSequence}(d_n, \alpha).$$

Remark (Interpretation). Decimal truncations converge because the truncation error is always less than 10^{-n} . This is one of the most concrete examples of convergence: finite decimal data can approximate a real number to arbitrary accuracy.

Remark (Dependencies).

- [Decimal Truncation Sequence](#)
- [Sequence Squeeze Theorem](#)

Proofs

Remark (Return). [Return to Theorem](#)

Theorem (Constant Sequence Convergence). *Let (x_n) be a real sequence, and let $c \in \mathbb{R}$. If $x_n = c$ for every $n \in \mathbb{N}$, then (x_n) converges to c .*

Professional Standard Proof.

TODO: professional standard proof for constant-sequence-convergence. □

Detailed Learning Proof.

TODO: detailed learning proof for constant-sequence-convergence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Zero Sequence is Null). *Let (x_n) be a real sequence. If $x_n = 0$ for every $n \in \mathbb{N}$, then (x_n) is a null sequence.*

Professional Standard Proof.

TODO: professional standard proof for zero-sequence-is-null.

□

Detailed Learning Proof.

TODO: detailed learning proof for zero-sequence-is-null.

□

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Constant Null Sequence). *Let (x_n) be a real sequence, and let $c \in \mathbb{R}$. If $x_n = c$ for every $n \in \mathbb{N}$, then (x_n) is a null sequence if and only if $c = 0$.*

Professional Standard Proof.

TODO: professional standard proof for constant-null-sequence. □

Detailed Learning Proof.

TODO: detailed learning proof for constant-null-sequence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Difference from the Limit is Null). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. The sequence (x_n) converges to L if and only if the sequence $(x_n - L)$ is a null sequence.*

Professional Standard Proof.

TODO: professional standard proof for difference-from-limit-is-null. □

Detailed Learning Proof.

TODO: detailed learning proof for difference-from-limit-is-null. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Ultimately Constant Sequence Convergence). *Let (x_n) be a real sequence. If there exist $N_0 \in \mathbb{N}$ and $c \in \mathbb{R}$ such that $x_n = c$ for every $n \geq N_0$, then (x_n) converges to c .*

Professional Standard Proof.

TODO: professional standard proof for ultimately-constant-sequence-convergence. □

Detailed Learning Proof.

TODO: detailed learning proof for ultimately-constant-sequence-convergence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Constant Implies Ultimately Constant). *Every constant real sequence is ultimately constant.*

Professional Standard Proof.

TODO: professional standard proof for constant-implies-ultimately-constant. □

Detailed Learning Proof.

TODO: detailed learning proof for constant-implies-ultimately-constant. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Ultimately Zero Sequence is Null). *Let (x_n) be a real sequence. If there exists $N_0 \in \mathbb{N}$ such that $x_n = 0$ for every $n \geq N_0$, then (x_n) is a null sequence.*

Professional Standard Proof.

TODO: professional standard proof for ultimately-zero-sequence-is-null. □

Detailed Learning Proof.

TODO: detailed learning proof for ultimately-zero-sequence-is-null. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Ultimately Constant Null Sequence). *Let (x_n) be a real sequence. Suppose there exist $N_0 \in \mathbb{N}$ and $c \in \mathbb{R}$ such that $x_n = c$ for every $n \geq N_0$. Then (x_n) is a null sequence if and only if $c = 0$.*

Professional Standard Proof.

TODO: professional standard proof for ultimately-constant-null-sequence. □

Detailed Learning Proof.

TODO: detailed learning proof for ultimately-constant-null-sequence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Tail Equality Preserves Convergence). *Let (x_n) and (y_n) be real sequences. If there exists $N_0 \in \mathbb{N}$ such that $x_n = y_n$ for every $n \geq N_0$, then for every $L \in \mathbb{R}$, (x_n) converges to L if and only if (y_n) converges to L .*

Professional Standard Proof.

TODO: professional standard proof for tail-equality-preserves-convergence. □

Detailed Learning Proof.

TODO: detailed learning proof for tail-equality-preserves-convergence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Eventually Bounded Above Tail Formulation). *Let (x_n) be a real sequence. The sequence (x_n) is bounded above if and only if there exist $N_0 \in \mathbb{N}$ and $M \in \mathbb{R}$ such that $x_n \leq M$ for every $n \geq N_0$.*

Professional Standard Proof.

TODO: professional standard proof for eventually-bounded-above-tail-formulation. □

Detailed Learning Proof.

TODO: detailed learning proof for eventually-bounded-above-tail-formulation. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Eventually Bounded Below Tail Formulation). *Let (x_n) be a real sequence. The sequence (x_n) is bounded below if and only if there exist $N_0 \in \mathbb{N}$ and $m \in \mathbb{R}$ such that $m \leq x_n$ for every $n \geq N_0$.*

Professional Standard Proof.

TODO: professional standard proof for eventually-bounded-below-tail-formulation. □

Detailed Learning Proof.

TODO: detailed learning proof for eventually-bounded-below-tail-formulation. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Eventually Bounded Tail Formulation). *Let (x_n) be a real sequence. The sequence (x_n) is bounded if and only if there exist $N_0 \in \mathbb{N}$ and $M > 0$ such that $|x_n| \leq M$ for every $n \geq N_0$.*

Professional Standard Proof.

TODO: professional standard proof for eventually-bounded-tail-formulation. □

Detailed Learning Proof.

TODO: detailed learning proof for eventually-bounded-tail-formulation. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Bounded Sequence If and Only If Bounded Above and Below). *Let (x_n) be a real sequence. The sequence (x_n) is bounded if and only if it is both bounded above and bounded below.*

Professional Standard Proof.

TODO: professional standard proof for bounded-sequence-iff-bounded-above-and-below. $\square$

Detailed Learning Proof.

TODO: detailed learning proof for bounded-sequence-iff-bounded-above-and-below. $\square$

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Absolute Bound Gives Upper and Lower Bounds). *Let (x_n) be a real sequence, and let $K > 0$. If $|x_n| \leq K$ for every $n \in \mathbb{N}$, then $-K \leq x_n \leq K$ for every $n \in \mathbb{N}$.*

Professional Standard Proof.

TODO: professional standard proof for absolute-bound-gives-upper-and-lower-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for absolute-bound-gives-upper-and-lower-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Upper and Lower Bounds Give an Absolute Bound). *Let (x_n) be a real sequence. If there exist $m, M \in \mathbb{R}$ such that $m \leq x_n \leq M$ for every $n \in \mathbb{N}$, then there exists $K > 0$ such that $|x_n| \leq K$ for every $n \in \mathbb{N}$.*

Professional Standard Proof.

TODO: professional standard proof for upper-and-lower-bounds-give-absolute-bound. □

Detailed Learning Proof.

TODO: detailed learning proof for upper-and-lower-bounds-give-absolute-bound. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limit of a Scalar Multiple). *Let (x_n) be a real sequence, let $L \in \mathbb{R}$, and let $\alpha \in \mathbb{R}$. If $x_n \rightarrow L$, then $\alpha x_n \rightarrow \alpha L$.*

Professional Standard Proof.

TODO: professional standard proof for limit-of-scalar-multiple. □

Detailed Learning Proof.

TODO: detailed learning proof for limit-of-scalar-multiple. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limit of a Sum). *Let (x_n) and (y_n) be real sequences, and let $L, M \in \mathbb{R}$. If $x_n \rightarrow L$ and $y_n \rightarrow M$, then $x_n + y_n \rightarrow L + M$.*

Professional Standard Proof.

TODO: professional standard proof for limit-of-a-sum. □

Detailed Learning Proof.

TODO: detailed learning proof for limit-of-a-sum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limit of a Negation). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. If $x_n \rightarrow L$, then $-x_n \rightarrow -L$.*

Professional Standard Proof.

TODO: professional standard proof for limit-of-a-negation. □

Detailed Learning Proof.

TODO: detailed learning proof for limit-of-a-negation. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limit of a Difference). *Let (x_n) and (y_n) be real sequences, and let $L, M \in \mathbb{R}$. If $x_n \rightarrow L$ and $y_n \rightarrow M$, then $x_n - y_n \rightarrow L - M$.*

Professional Standard Proof.

TODO: professional standard proof for limit-of-a-difference. □

Detailed Learning Proof.

TODO: detailed learning proof for limit-of-a-difference. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limit of a Product). *Let (x_n) and (y_n) be real sequences, and let $L, M \in \mathbb{R}$. If $x_n \rightarrow L$ and $y_n \rightarrow M$, then $x_n y_n \rightarrow LM$.*

Professional Standard Proof.

TODO: professional standard proof for limit-of-a-product. □

Detailed Learning Proof.

TODO: detailed learning proof for limit-of-a-product. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Nonzero Limit is Eventually Nonzero). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$ with $L \neq 0$. If $x_n \rightarrow L$, then there exists $N \in \mathbb{N}$ such that $x_n \neq 0$ for every $n \geq N$.*

Professional Standard Proof.

TODO: professional standard proof for nonzero-limit-eventually-nonzero. □

Detailed Learning Proof.

TODO: detailed learning proof for nonzero-limit-eventually-nonzero. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limit of a Reciprocal). *Let (x_n) be a real sequence such that $x_n \neq 0$ for every $n \in \mathbb{N}$, and let $L \in \mathbb{R}$ with $L \neq 0$. If $x_n \rightarrow L$, then $1/x_n \rightarrow 1/L$.*

Professional Standard Proof.

TODO: professional standard proof for limit-of-a-reciprocal. □

Detailed Learning Proof.

TODO: detailed learning proof for limit-of-a-reciprocal. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limit of a Quotient). *Let (x_n) and (y_n) be real sequences such that $y_n \neq 0$ for every $n \in \mathbb{N}$, and let $L, M \in \mathbb{R}$ with $M \neq 0$. If $x_n \rightarrow L$ and $y_n \rightarrow M$, then $x_n/y_n \rightarrow L/M$.*

Professional Standard Proof.

TODO: professional standard proof for limit-of-a-quotient. □

Detailed Learning Proof.

TODO: detailed learning proof for limit-of-a-quotient. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limit of a Square). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. If $x_n \rightarrow L$, then $x_n^2 \rightarrow L^2$.*

Professional Standard Proof.

TODO: professional standard proof for limit-of-a-square. □

Detailed Learning Proof.

TODO: detailed learning proof for limit-of-a-square. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limit of an Absolute Value). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. If $x_n \rightarrow L$, then $|x_n| \rightarrow |L|$.*

Professional Standard Proof.

TODO: professional standard proof for limit-of-an-absolute-value. □

Detailed Learning Proof.

TODO: detailed learning proof for limit-of-an-absolute-value. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Positive Limit is Eventually Positive). *Let (x_n) be a real sequence, and let $L > 0$. If $x_n \rightarrow L$, then there exists $N \in \mathbb{N}$ such that $0 < x_n$ for every $n \geq N$.*

Professional Standard Proof.

TODO: professional standard proof for positive-limit-eventually-positive. □

Detailed Learning Proof.

TODO: detailed learning proof for positive-limit-eventually-positive. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limit of a Square Root). *Let (x_n) be a real sequence such that $0 \leq x_n$ for every $n \in \mathbb{N}$, and let $L \in \mathbb{R}$. If $x_n \rightarrow L$, then $0 \leq L$ and $\sqrt{x_n} \rightarrow \sqrt{L}$.*

Professional Standard Proof.

TODO: professional standard proof for limit-of-a-square-root.

□

Detailed Learning Proof.

TODO: detailed learning proof for limit-of-a-square-root.

□

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Polynomial Sequence Limit). *Let (x_n) be a real sequence, let $L \in \mathbb{R}$, and let $p \in \mathbb{R}[t]$ be a polynomial. If $x_n \rightarrow L$, then $p(x_n) \rightarrow p(L)$.*

Professional Standard Proof.

TODO: professional standard proof for polynomial-sequence-limit. □

Detailed Learning Proof.

TODO: detailed learning proof for polynomial-sequence-limit. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Rational Sequence Limit). *Let (x_n) be a real sequence, let $L \in \mathbb{R}$, and let $p, q \in \mathbb{R}[t]$ be polynomials. Suppose $q(L) \neq 0$ and $q(x_n) \neq 0$ for every $n \in \mathbb{N}$. If $x_n \rightarrow L$, then*

$$\frac{p(x_n)}{q(x_n)} \rightarrow \frac{p(L)}{q(L)}.$$

Professional Standard Proof.

TODO: professional standard proof for rational-sequence-limit. □

Detailed Learning Proof.

TODO: detailed learning proof for rational-sequence-limit. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Equivalence of Convergence Formulations). *Let $(x_n) \subseteq \mathbb{R}$ and let $L \in \mathbb{R}$. The following statements are equivalent.*

- (a) $x_n \rightarrow L$.
- (b) $\forall \varepsilon > 0 \exists K \in \mathbb{N} \forall n \geq K (|x_n - L| < \varepsilon)$.
- (c) $\forall \varepsilon > 0 \exists K \in \mathbb{N} \forall n \geq K (L - \varepsilon < x_n < L + \varepsilon)$.
- (d) $\forall \varepsilon > 0 \exists K \in \mathbb{N} \forall n \geq K (x_n \in V_\varepsilon(L))$.

Professional Standard Proof.

TODO: professional standard proof for equivalence-of-convergence-formulations. □

Detailed Learning Proof.

TODO: detailed learning proof for equivalence-of-convergence-formulations. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Uniqueness of Limits). *Let $(x_n) \subseteq \mathbb{R}$. If $x_n \rightarrow L$ and $x_n \rightarrow K$, then $L = K$.*

Professional Standard Proof.

TODO: professional standard proof for uniqueness-of-limits. □

Detailed Learning Proof.

TODO: detailed learning proof for uniqueness-of-limits. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limit Preserves Eventual Order). *Let (x_n) and (y_n) be real sequences, and let $L, M \in \mathbb{R}$. Suppose that $x_n \rightarrow L$ and $y_n \rightarrow M$. If there exists $N_0 \in \mathbb{N}$ such that $x_n \leq y_n$ for every $n \geq N_0$, then $L \leq M$.*

Professional Standard Proof.

TODO: professional standard proof for limit-preserves-eventual-order. □

Detailed Learning Proof.

TODO: detailed learning proof for limit-preserves-eventual-order. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Strict Limit Separation Gives Eventual Order). *Let (x_n) and (y_n) be real sequences, and let $A, B \in \mathbb{R}$. Suppose that $x_n \rightarrow A$ and $y_n \rightarrow B$. If $A < B$, then there exists $N \in \mathbb{N}$ such that $x_n < y_n$ for every $n \geq N$.*

Professional Standard Proof.

TODO: professional standard proof for strict-limit-separation-gives-eventual-order. □

Detailed Learning Proof.

TODO: detailed learning proof for strict-limit-separation-gives-eventual-order. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Eventual Strict Comparison Preserves Weak Limit Order). *Let (x_n) and (y_n) be real sequences, and let $A, B \in \mathbb{R}$. Suppose that $x_n \rightarrow A$ and $y_n \rightarrow B$.*

(a) If there exists $N \in \mathbb{N}$ such that $x_n < y_n$ for every $n \geq N$, then $A \leq B$.

(b) If there exists $N \in \mathbb{N}$ such that $x_n > y_n$ for every $n \geq N$, then $A \geq B$.

Professional Standard Proof.

TODO: professional standard proof for eventual-strict-comparison-preserves-weak-limit-order. □

Detailed Learning Proof.

TODO: detailed learning proof for eventual-strict-comparison-preserves-weak-limit-order. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Constant Comparison for Sequence Limits). *Let (x_n) be a real sequence, and let $A, B \in \mathbb{R}$. Suppose that $x_n \rightarrow A$.*

- (a) *If there exists $N \in \mathbb{N}$ such that $x_n \leq B$ for every $n \geq N$, then $A \leq B$.*
- (b) *If there exists $N \in \mathbb{N}$ such that $x_n < B$ for every $n \geq N$, then $A \leq B$.*
- (c) *If there exists $N \in \mathbb{N}$ such that $x_n \geq B$ for every $n \geq N$, then $A \geq B$.*
- (d) *If there exists $N \in \mathbb{N}$ such that $x_n > B$ for every $n \geq N$, then $A \geq B$.*

Professional Standard Proof.

TODO: professional standard proof for constant-comparison-for-sequence-limits. □

Detailed Learning Proof.

TODO: detailed learning proof for constant-comparison-for-sequence-limits. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Constant Squeeze Theorem). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. If there exists $N_0 \in \mathbb{N}$ such that $L \leq x_n \leq L$ for every $n \geq N_0$, then $x_n \rightarrow L$.*

Professional Standard Proof.

TODO: professional standard proof for constant-squeeze-theorem. □

Detailed Learning Proof.

TODO: detailed learning proof for constant-squeeze-theorem. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Sequence Squeeze Theorem). *Let (a_n) , (x_n) , and (b_n) be real sequences, and let $L \in \mathbb{R}$. Suppose that $a_n \rightarrow L$ and $b_n \rightarrow L$. If there exists $N_0 \in \mathbb{N}$ such that $a_n \leq x_n \leq b_n$ for every $n \geq N_0$, then $x_n \rightarrow L$.*

Professional Standard Proof.

TODO: professional standard proof for sequence-squeeze-theorem. □

Detailed Learning Proof.

TODO: detailed learning proof for sequence-squeeze-theorem. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Absolute-Value Squeeze Theorem). *Let (x_n) and (u_n) be real sequences, and let $L \in \mathbb{R}$. Suppose that $u_n \rightarrow 0$. If there exists $N_0 \in \mathbb{N}$ such that $|x_n - L| \leq u_n$ for every $n \geq N_0$, then $x_n \rightarrow L$.*

Professional Standard Proof.

TODO: professional standard proof for absolute-value-squeeze-theorem. □

Detailed Learning Proof.

TODO: detailed learning proof for absolute-value-squeeze-theorem. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Convergence of a Tail). *Let $(x_n) \subseteq \mathbb{R}$ and let $m \in \mathbb{N}$. The m -tail $(x_{m+n})_{n \in \mathbb{N}}$ converges if and only if (x_n) converges. Moreover, in this case,*

$$\lim_{n \rightarrow \infty} x_{m+n} = \lim_{n \rightarrow \infty} x_n.$$

Professional Standard Proof.

TODO: professional standard proof for convergence-of-tail. □

Detailed Learning Proof.

TODO: detailed learning proof for convergence-of-tail. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Convergence by Domination). *Let $(x_n) \subseteq \mathbb{R}$ and let $L \in \mathbb{R}$. Let (a_n) be a sequence of positive real numbers with $\lim_{n \rightarrow \infty} a_n = 0$. If there exist a constant $c > 0$ and an index $m \in \mathbb{N}$ such that*

$$|x_n - L| \leq c a_n \quad \forall n \geq m,$$

then $\lim_{n \rightarrow \infty} x_n = L$.

Professional Standard Proof.

TODO: professional standard proof for convergence-by-domination. □

Detailed Learning Proof.

TODO: detailed learning proof for convergence-by-domination. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Ratio Limit Less Than One Implies Null). *Let (x_n) be a sequence of positive real numbers. Suppose that the limit*

$$L := \lim_{n \rightarrow \infty} \frac{x_{n+1}}{x_n}$$

exists. If $L < 1$, then (x_n) converges and

$$\lim_{n \rightarrow \infty} x_n = 0.$$

Professional Standard Proof.

TODO: professional standard proof for ratio-limit-less-than-one-implies-null. □

Detailed Learning Proof.

TODO: detailed learning proof for ratio-limit-less-than-one-implies-null. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Monotone Convergence Theorem). *Let (x_n) be a real sequence.*

(a) *If (x_n) is increasing and bounded above, then (x_n) converges and*

$$\lim_{n \rightarrow \infty} x_n = \sup\{x_n : n \in \mathbb{N}\}.$$

(b) *If (x_n) is decreasing and bounded below, then (x_n) converges and*

$$\lim_{n \rightarrow \infty} x_n = \inf\{x_n : n \in \mathbb{N}\}.$$

Professional Standard Proof.

TODO: professional standard proof for monotone-convergence-theorem. □

Detailed Learning Proof.

TODO: detailed learning proof for monotone-convergence-theorem. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Strict Monotonicity Implies Monotonicity). *Let (x_n) be a real sequence.*

(a) If (x_n) is strictly increasing, then (x_n) is increasing.

(b) If (x_n) is strictly decreasing, then (x_n) is decreasing.

Professional Standard Proof.

TODO: professional standard proof for strict-monotonicity-implies-monotonicity. □

Detailed Learning Proof.

TODO: detailed learning proof for strict-monotonicity-implies-monotonicity. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Bounded Monotone Sequence Equivalences). *Let (x_n) be a real sequence.*

- (a) If (x_n) is increasing, then (x_n) converges if and only if (x_n) is bounded above.*
- (b) If (x_n) is decreasing, then (x_n) converges if and only if (x_n) is bounded below.*
- (c) If (x_n) is monotone, then (x_n) converges if and only if (x_n) is bounded.*

Professional Standard Proof.

TODO: professional standard proof for bounded-monotone-sequence-equivalences. □

Detailed Learning Proof.

TODO: detailed learning proof for bounded-monotone-sequence-equivalences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Eventually Monotone Convergence Theorem). *Let (x_n) be a real sequence.*

- (a) If (x_n) is eventually increasing and bounded above, then (x_n) converges.*
- (b) If (x_n) is eventually decreasing and bounded below, then (x_n) converges.*
- (c) If (x_n) is eventually monotone and bounded, then (x_n) converges.*

Professional Standard Proof.

TODO: professional standard proof for eventually-monotone-convergence-theorem. □

Detailed Learning Proof.

TODO: detailed learning proof for eventually-monotone-convergence-theorem. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Unbounded Monotone Divergence). *Let (x_n) be a real sequence.*

(a) If (x_n) is increasing and is not bounded above, then $x_n \rightarrow +\infty$.

(b) If (x_n) is decreasing and is not bounded below, then $x_n \rightarrow -\infty$.

Professional Standard Proof.

TODO: professional standard proof for unbounded-monotone-divergence. □

Detailed Learning Proof.

TODO: detailed learning proof for unbounded-monotone-divergence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Algebraic Transformations Preserve Monotonicity). *Let (x_n) be a real sequence and let $c, \alpha \in \mathbb{R}$.*

- (a) *If (x_n) is increasing, then $(x_n + c)$ is increasing.*
- (b) *If (x_n) is decreasing, then $(x_n + c)$ is decreasing.*
- (c) *If (x_n) is increasing and $\alpha > 0$, then (αx_n) is increasing.*
- (d) *If (x_n) is increasing and $\alpha < 0$, then (αx_n) is decreasing.*
- (e) *If (x_n) is increasing, then $(-x_n)$ is decreasing.*
- (f) *If (x_n) is decreasing, then $(-x_n)$ is increasing.*

Professional Standard Proof.

TODO: professional standard proof for algebraic-transformations-preserve-monotonicity. □

Detailed Learning Proof.

TODO: detailed learning proof for algebraic-transformations-preserve-monotonicity. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Subsequence Indices Dominate the Identity). *Let $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ be a strictly increasing index map. Then*

$$k \leq \sigma(k) \quad \text{for every } k \in \mathbb{N}.$$

Professional Standard Proof.

TODO: professional standard proof for subsequence-indices-dominate-identity. □

Detailed Learning Proof.

TODO: detailed learning proof for subsequence-indices-dominate-identity. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Subsequences Preserve Limits). *Let (x_n) be a real sequence and let $L \in \mathbb{R}$. If $x_n \rightarrow L$, then every subsequence of (x_n) converges to L .*

Professional Standard Proof.

TODO: professional standard proof for subsequences-preserve-limits. □

Detailed Learning Proof.

TODO: detailed learning proof for subsequences-preserve-limits. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Subsequential Limit of a Convergent Sequence). *Let (x_n) be a real sequence and let $L \in \mathbb{R}$. If $x_n \rightarrow L$, then L is the only possible subsequential limit of (x_n) .*

Professional Standard Proof.

TODO: professional standard proof for subsequential-limit-of-convergent-sequence. □

Detailed Learning Proof.

TODO: detailed learning proof for subsequential-limit-of-convergent-sequence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Divergence by Two Subsequential Limits). *Let (x_n) be a real sequence. If (x_n) has two distinct subsequential limits, then (x_n) does not converge.*

Professional Standard Proof.

TODO: professional standard proof for divergence-by-two-subsequential-limits. □

Detailed Learning Proof.

TODO: detailed learning proof for divergence-by-two-subsequential-limits. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Boundedness Passes to Subsequences). *Every subsequence of a bounded real sequence is bounded.*

Professional Standard Proof.

TODO: professional standard proof for boundedness-passes-to-subsequences. □

Detailed Learning Proof.

TODO: detailed learning proof for boundedness-passes-to-subsequences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Monotonicity Passes to Subsequences). *Every subsequence of an increasing real sequence is increasing, and every subsequence of a decreasing real sequence is decreasing.*

Professional Standard Proof.

TODO: professional standard proof for monotonicity-passes-to-subsequences. □

Detailed Learning Proof.

TODO: detailed learning proof for monotonicity-passes-to-subsequences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Subsequence of a Subsequence). *Every subsequence of a subsequence of (x_n) is a subsequence of (x_n) .*

Professional Standard Proof.

TODO: professional standard proof for subsequence-of-subsequence. □

Detailed Learning Proof.

TODO: detailed learning proof for subsequence-of-subsequence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Eventual Properties Pass to Subsequences). *Let $P(n)$ be a property of indices. If there exists $N \in \mathbb{N}$ such that $P(n)$ holds for every $n \geq N$, then for every strictly increasing index map σ there exists $K \in \mathbb{N}$ such that $P(\sigma(k))$ holds for every $k \geq K$.*

Professional Standard Proof.

TODO: professional standard proof for eventual-properties-pass-to-subsequences. □

Detailed Learning Proof.

TODO: detailed learning proof for eventual-properties-pass-to-subsequences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Frequent Properties Yield Subsequences). *Let $P(n)$ be a property of indices. If for every $N \in \mathbb{N}$ there exists $n \geq N$ such that $P(n)$ holds, then there exists a strictly increasing index map $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ such that $P(\sigma(k))$ holds for every $k \in \mathbb{N}$.*

Professional Standard Proof.

TODO: professional standard proof for frequent-properties-*yield*-subsequences. □

Detailed Learning Proof.

TODO: detailed learning proof for frequent-properties-*yield*-subsequences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Subsequential Limits Respect Bounds). *Let (x_n) be a real sequence and let L be a subsequential limit of (x_n) . If $m \leq x_n \leq M$ for every $n \in \mathbb{N}$, then $m \leq L \leq M$.*

Professional Standard Proof.

TODO: professional standard proof for subsequential-limits-respect-bounds. □

Detailed Learning Proof.

TODO: detailed learning proof for subsequential-limits-respect-bounds. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Squeeze Passes to Subsequences). *Let (a_n) , (x_n) , and (b_n) be real sequences. If $a_n \leq x_n \leq b_n$ eventually, then for every subsequence $(x_{\sigma(k)})$ there exists $K \in \mathbb{N}$ such that*

$$a_{\sigma(k)} \leq x_{\sigma(k)} \leq b_{\sigma(k)} \quad \text{for every } k \geq K.$$

Professional Standard Proof.

TODO: professional standard proof for squeeze-passes-to-subsequences. □

Detailed Learning Proof.

TODO: detailed learning proof for squeeze-passes-to-subsequences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Monotone Subsequence Theorem). *Every real sequence has a monotone subsequence.*

Professional Standard Proof.

TODO: professional standard proof for monotone-subsequence-theorem. □

Detailed Learning Proof.

TODO: detailed learning proof for monotone-subsequence-theorem. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Bolzano-Weierstrass Theorem for Sequences). *Every bounded real sequence has a convergent subsequence.*

Professional Standard Proof.

TODO: professional standard proof for bolzano-weierstrass-sequences. □

Detailed Learning Proof.

TODO: detailed learning proof for bolzano-weierstrass-sequences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Sequential Compactness of Closed Bounded Intervals). *Let $a, b \in \mathbb{R}$ with $a \leq b$. Every sequence in $[a, b]$ has a convergent subsequence whose limit belongs to $[a, b]$.*

Professional Standard Proof.

TODO: professional standard proof for sequential-compactness-closed-bounded-interval. $\square$

Detailed Learning Proof.

TODO: detailed learning proof for sequential-compactness-closed-bounded-interval. $\square$

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Divergence to Infinity Implies Real Divergence). *Let (x_n) be a real sequence. If $x_n \rightarrow +\infty$ or $x_n \rightarrow -\infty$, then (x_n) is divergent.*

Professional Standard Proof.

TODO: professional standard proof for divergence-to-infinity-implies-real-divergence. □

Detailed Learning Proof.

TODO: detailed learning proof for divergence-to-infinity-implies-real-divergence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Two Subsequential Limits Force Divergence). *Let (x_n) be a real sequence. If (x_n) has two subsequential limits $L, K \in \mathbb{R}$ with $L \neq K$, then (x_n) is divergent.*

Professional Standard Proof.

TODO: professional standard proof for two-subsequential-limits-force-divergence. □

Detailed Learning Proof.

TODO: detailed learning proof for two-subsequential-limits-force-divergence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Unbounded Above Sequence Has a Positive-Infinity Subsequence). *Let (x_n) be a real sequence. If (x_n) is not bounded above, then there exists a subsequence $(x_{\sigma(k)})$ such that $x_{\sigma(k)} \rightarrow +\infty$.*

Professional Standard Proof.

TODO: professional standard proof for unbounded-above-has-positive-infinity-subsequence. □

Detailed Learning Proof.

TODO: detailed learning proof for unbounded-above-has-positive-infinity-subsequence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Unbounded Below Sequence Has a Negative-Infinity Subsequence). *Let (x_n) be a real sequence. If (x_n) is not bounded below, then there exists a subsequence $(x_{\sigma(k)})$ such that $x_{\sigma(k)} \rightarrow -\infty$.*

Professional Standard Proof.

TODO: professional standard proof for unbounded-below-has-negative-infinity-subsequence. □

Detailed Learning Proof.

TODO: detailed learning proof for unbounded-below-has-negative-infinity-subsequence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Bounded Divergence Produces Two Subsequential Limits). *Let (x_n) be a bounded real sequence. If (x_n) is divergent, then there exist $L, K \in \mathbb{R}$ with $L \neq K$ such that L and K are subsequential limits of (x_n) .*

Professional Standard Proof.

TODO: professional standard proof for bounded-divergence-produces-two-subsequential-limits. □

Detailed Learning Proof.

TODO: detailed learning proof for bounded-divergence-produces-two-subsequential-limits. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Tail Suprema Are Decreasing). *Let (x_n) be a bounded real sequence, and let (s_n) be its tail supremum sequence. Then (s_n) is decreasing.*

Professional Standard Proof.

TODO: professional standard proof for tail-suprema-are-decreasing. □

Detailed Learning Proof.

TODO: detailed learning proof for tail-suprema-are-decreasing. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Tail Infima Are Increasing). *Let (x_n) be a bounded real sequence, and let (i_n) be its tail infimum sequence. Then (i_n) is increasing.*

Professional Standard Proof.

TODO: professional standard proof for tail-infima-are-increasing. □

Detailed Learning Proof.

TODO: detailed learning proof for tail-infima-are-increasing. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Liminf Is Below Limsup). *Let (x_n) be a bounded real sequence. Then*

$$\liminf_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} x_n.$$

Professional Standard Proof.

TODO: professional standard proof for liminf-below-limsup. □

Detailed Learning Proof.

TODO: detailed learning proof for liminf-below-limsup. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Convergence iff Liminf Equals Limsup). *Let (x_n) be a bounded real sequence, and let $L \in \mathbb{R}$. Then $x_n \rightarrow L$ if and only if*

$$\liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n = L.$$

Professional Standard Proof.

TODO: professional standard proof for convergence-iff-liminf-equals-limsup. □

Detailed Learning Proof.

TODO: detailed learning proof for convergence-iff-liminf-equals-limsup. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limsup Is the Largest Subsequential Limit). *Let (x_n) be a bounded real sequence, and let $S = \limsup_{n \rightarrow \infty} x_n$. Then S is a subsequential limit of (x_n) , and every subsequential limit of (x_n) is less than or equal to S .*

Professional Standard Proof.

TODO: professional standard proof for limsup-largest-subsequential-limit. □

Detailed Learning Proof.

TODO: detailed learning proof for limsup-largest-subsequential-limit. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Liminf Is the Smallest Subsequential Limit). *Let (x_n) be a bounded real sequence, and let $I = \liminf_{n \rightarrow \infty} x_n$. Then I is a subsequential limit of (x_n) , and every subsequential limit of (x_n) is greater than or equal to I .*

Professional Standard Proof.

TODO: professional standard proof for liminf-smallest-subsequential-limit. □

Detailed Learning Proof.

TODO: detailed learning proof for liminf-smallest-subsequential-limit. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Oscillation Criterion via Liminf and Limsup). *Let (x_n) be a bounded real sequence. Then*

$$\liminf_{n \rightarrow \infty} x_n < \limsup_{n \rightarrow \infty} x_n$$

if and only if (x_n) has two distinct subsequential limits.

Professional Standard Proof.

TODO: professional standard proof for oscillation-criterion-via-liminf-limsup. □

Detailed Learning Proof.

TODO: detailed learning proof for oscillation-criterion-via-liminf-limsup. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limsup Comparison Under Eventual Order). *Let (x_n) and (y_n) be bounded real sequences. If there exists $N \in \mathbb{N}$ such that $x_n \leq y_n$ for every $n \geq N$, then*

$$\limsup_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} y_n.$$

Professional Standard Proof.

TODO: professional standard proof for limsup-comparison-under-eventual-order. □

Detailed Learning Proof.

TODO: detailed learning proof for limsup-comparison-under-eventual-order. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Liminf Comparison Under Eventual Order). *Let (x_n) and (y_n) be bounded real sequences. If there exists $N \in \mathbb{N}$ such that $x_n \leq y_n$ for every $n \geq N$, then*

$$\liminf_{n \rightarrow \infty} x_n \leq \liminf_{n \rightarrow \infty} y_n.$$

Professional Standard Proof.

TODO: professional standard proof for liminf-comparison-under-eventual-order. □

Detailed Learning Proof.

TODO: detailed learning proof for liminf-comparison-under-eventual-order. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limsup Squeeze Under Eventual Order). *Let (a_n) , (x_n) , and (b_n) be bounded real sequences. If there exists $N \in \mathbb{N}$ such that*

$$a_n \leq x_n \leq b_n \quad \text{for every } n \geq N,$$

then

$$\limsup_{n \rightarrow \infty} a_n \leq \limsup_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} b_n.$$

Professional Standard Proof.

TODO: professional standard proof for limsup-squeeze-under-eventual-order. □

Detailed Learning Proof.

TODO: detailed learning proof for limsup-squeeze-under-eventual-order. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Liminf Squeeze Under Eventual Order). *Let (a_n) , (x_n) , and (b_n) be bounded real sequences. If there exists $N \in \mathbb{N}$ such that*

$$a_n \leq x_n \leq b_n \quad \text{for every } n \geq N,$$

then

$$\liminf_{n \rightarrow \infty} a_n \leq \liminf_{n \rightarrow \infty} x_n \leq \liminf_{n \rightarrow \infty} b_n.$$

Professional Standard Proof.

TODO: professional standard proof for liminf-squeeze-under-eventual-order. □

Detailed Learning Proof.

TODO: detailed learning proof for liminf-squeeze-under-eventual-order. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Increasing Sequence Limit as Supremum). *Let (x_n) be an increasing real sequence that is bounded above. Then*

$$\lim_{n \rightarrow \infty} x_n = \sup\{x_n : n \in \mathbb{N}\}.$$

Professional Standard Proof.

TODO: professional standard proof for increasing-sequence-limit-as-supremum. □

Detailed Learning Proof.

TODO: detailed learning proof for increasing-sequence-limit-as-supremum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Decreasing Sequence Limit as Infimum). *Let (x_n) be a decreasing real sequence that is bounded below. Then*

$$\lim_{n \rightarrow \infty} x_n = \inf\{x_n : n \in \mathbb{N}\}.$$

Professional Standard Proof.

TODO: professional standard proof for decreasing-sequence-limit-as-infimum. □

Detailed Learning Proof.

TODO: detailed learning proof for decreasing-sequence-limit-as-infimum. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Tail Suprema and Infima Converge). *Let (x_n) be a bounded real sequence. Define*

$$s_n = \sup\{x_k : k \geq n\}, \quad i_n = \inf\{x_k : k \geq n\}.$$

Then (s_n) and (i_n) converge.

Professional Standard Proof.

TODO: professional standard proof for tail-suprema-and-infima-converge. □

Detailed Learning Proof.

TODO: detailed learning proof for tail-suprema-and-infima-converge. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Bounded Sequence Has Limsup and Liminf). *Every bounded real sequence has a limit superior and a limit inferior.*

Professional Standard Proof.

TODO: professional standard proof for bounded-sequence-has-limsup-and-liminf. □

Detailed Learning Proof.

TODO: detailed learning proof for bounded-sequence-has-limsup-and-liminf. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Cluster Values Are Subsequential Limits). *Let (x_n) be a real sequence, and let $L \in \mathbb{R}$. Then L is a cluster value of (x_n) if and only if L is a subsequential limit of (x_n) .*

Professional Standard Proof.

TODO: professional standard proof for cluster-values-are-subsequential-limits. □

Detailed Learning Proof.

TODO: detailed learning proof for cluster-values-are-subsequential-limits. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Bounded Sequences Have Cluster Values). *Every bounded real sequence has at least one cluster value.*

Professional Standard Proof.

TODO: professional standard proof for bounded-sequences-have-cluster-values. □

Detailed Learning Proof.

TODO: detailed learning proof for bounded-sequences-have-cluster-values. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Limsup and Liminf Are Extremal Cluster Values). *Let (x_n) be a bounded real sequence. Then $\limsup_{n \rightarrow \infty} x_n$ is the largest cluster value of (x_n) , and $\liminf_{n \rightarrow \infty} x_n$ is the smallest cluster value of (x_n) .*

Professional Standard Proof.

TODO: professional standard proof for limsup-liminf-extremal-cluster-values. □

Detailed Learning Proof.

TODO: detailed learning proof for limsup-liminf-extremal-cluster-values. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Convergent Sequences Are Cauchy). *Let (x_n) be a real sequence. If (x_n) converges, then (x_n) is Cauchy.*

Professional Standard Proof.

TODO: professional standard proof for convergent-sequences-are-cauchy. □

Detailed Learning Proof.

TODO: detailed learning proof for convergent-sequences-are-cauchy. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Cauchy Sequences Are Bounded). *Every Cauchy real sequence is bounded.*

Professional Standard Proof.

TODO: professional standard proof for cauchy-sequences-are-bounded.

□

Detailed Learning Proof.

TODO: detailed learning proof for cauchy-sequences-are-bounded.

□

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Cauchy Sequence with a Convergent Subsequence). *Let (x_n) be a Cauchy real sequence. If a subsequence of (x_n) converges to $L \in \mathbb{R}$, then (x_n) converges to L .*

Professional Standard Proof.

TODO: professional standard proof for cauchy-sequence-with-convergent-subsequence. □

Detailed Learning Proof.

TODO: detailed learning proof for cauchy-sequence-with-convergent-subsequence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Cauchy Criterion for Real Sequences). *Let (x_n) be a real sequence. Then (x_n) converges if and only if (x_n) is Cauchy.*

Professional Standard Proof.

TODO: professional standard proof for cauchy-criterion-real-sequences. □

Detailed Learning Proof.

TODO: detailed learning proof for cauchy-criterion-real-sequences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Cauchy Criterion via Tails). *Let (x_n) be a real sequence. Then (x_n) is Cauchy if and only if for every $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that the N -tail has pairwise term distances less than ε .*

Professional Standard Proof.

TODO: professional standard proof for cauchy-criterion-via-tails. □

Detailed Learning Proof.

TODO: detailed learning proof for cauchy-criterion-via-tails. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Cauchy Tail Diameter Criterion). *Let (x_n) be a real sequence. Then (x_n) is Cauchy if and only if every sufficiently late tail has arbitrarily small pairwise diameter.*

Professional Standard Proof.

TODO: professional standard proof for cauchy-tail-diameter-criterion. □

Detailed Learning Proof.

TODO: detailed learning proof for cauchy-tail-diameter-criterion. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Cauchy Sequences Have Vanishing Successive Differences). *If (x_n) is a Cauchy real sequence, then*

$$|x_{n+1} - x_n| \rightarrow 0.$$

Professional Standard Proof.

TODO: professional standard proof for cauchy-successive-differences-vanish. □

Detailed Learning Proof.

TODO: detailed learning proof for cauchy-successive-differences-vanish. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Scalar Multiples of Cauchy Sequences). *Let (x_n) be a Cauchy real sequence, and let $\alpha \in \mathbb{R}$. Then (αx_n) is Cauchy.*

Professional Standard Proof.

TODO: professional standard proof for scalar-multiple-cauchy-sequence. □

Detailed Learning Proof.

TODO: detailed learning proof for scalar-multiple-cauchy-sequence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Sums of Cauchy Sequences). *Let (x_n) and (y_n) be Cauchy real sequences. Then $(x_n + y_n)$ is Cauchy.*

Professional Standard Proof.

TODO: professional standard proof for sum-cauchy-sequences. □

Detailed Learning Proof.

TODO: detailed learning proof for sum-cauchy-sequences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Differences of Cauchy Sequences). *Let (x_n) and (y_n) be Cauchy real sequences. Then $(x_n - y_n)$ is Cauchy.*

Professional Standard Proof.

TODO: professional standard proof for difference-cauchy-sequences. □

Detailed Learning Proof.

TODO: detailed learning proof for difference-cauchy-sequences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Linear Combinations of Cauchy Sequences). *Let (x_n) and (y_n) be Cauchy real sequences, and let $\alpha, \beta \in \mathbb{R}$. Then $(\alpha x_n + \beta y_n)$ is Cauchy.*

Professional Standard Proof.

TODO: professional standard proof for linear-combination-cauchy-sequences. □

Detailed Learning Proof.

TODO: detailed learning proof for linear-combination-cauchy-sequences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Products of Cauchy Sequences). *Let (x_n) and (y_n) be Cauchy real sequences. Then $(x_n y_n)$ is Cauchy.*

Professional Standard Proof.

TODO: professional standard proof for product-cauchy-sequences. □

Detailed Learning Proof.

TODO: detailed learning proof for product-cauchy-sequences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Reciprocals of Cauchy Sequences). *Let (x_n) be a Cauchy real sequence. If there exist $c > 0$ and $N_0 \in \mathbb{N}$ such that $|x_n| \geq c$ for every $n \geq N_0$, then $(1/x_n)$ is Cauchy.*

Professional Standard Proof.

TODO: professional standard proof for reciprocal-cauchy-sequence. □

Detailed Learning Proof.

TODO: detailed learning proof for reciprocal-cauchy-sequence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Quotients of Cauchy Sequences). *Let (x_n) and (y_n) be Cauchy real sequences. If there exist $c > 0$ and $N_0 \in \mathbb{N}$ such that $|y_n| \geq c$ for every $n \geq N_0$, then (x_n/y_n) is Cauchy.*

Professional Standard Proof.

TODO: professional standard proof for quotient-cauchy-sequences. □

Detailed Learning Proof.

TODO: detailed learning proof for quotient-cauchy-sequences. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Absolute Values of Cauchy Sequences). *Let (x_n) be a Cauchy real sequence. Then $(|x_n|)$ is Cauchy.*

Professional Standard Proof.

TODO: professional standard proof for absolute-value-cauchy-sequence. □

Detailed Learning Proof.

TODO: detailed learning proof for absolute-value-cauchy-sequence. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Newton Approximation of 2). *Let (x_n) be the Newton sequence for $\sqrt{2}$. Then (x_n) converges and*

$$\lim_{n \rightarrow \infty} x_n = \sqrt{2}.$$

Professional Standard Proof.

TODO: professional standard proof for newton-approximation-sqrt-two. □

Detailed Learning Proof.

TODO: detailed learning proof for newton-approximation-sqrt-two. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Factorial Partial Sums Approximate e). *Let (s_n) be the factorial partial-sum sequence. Then (s_n) converges to the Euler number e :*

$$\lim_{n \rightarrow \infty} s_n = e.$$

Professional Standard Proof.

TODO: professional standard proof for factorial-partial-sums-approximate-e. □

Detailed Learning Proof.

TODO: detailed learning proof for factorial-partial-sums-approximate-e. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Compound-Interest Approximation of e). *Let (c_n) be the compound-interest sequence. Then*

$$\lim_{n \rightarrow \infty} c_n = e.$$

Professional Standard Proof.

TODO: professional standard proof for compound-interest-approximation-e. □

Detailed Learning Proof.

TODO: detailed learning proof for compound-interest-approximation-e. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Remark (Return). [Return to Theorem](#)

Theorem (Decimal Truncations Converge). *Let $\alpha \in \mathbb{R}$, and let (d_n) be the decimal truncation sequence of α . Then*

$$\lim_{n \rightarrow \infty} d_n = \alpha.$$

Professional Standard Proof.

TODO: professional standard proof for decimal-truncations-converge. □

Detailed Learning Proof.

TODO: detailed learning proof for decimal-truncations-converge. □

Remark (Proof structure).

Remark (Dependencies).

- TODO: list mathematical dependencies.

Capstone

Capstone Theorem: The Contraction Mapping Principle on $\mathbb{R}$

Theorem. Let $f : \mathbb{R} \rightarrow \mathbb{R}$. Suppose there is a constant L with $0 \leq L < 1$ such that

$$|f(x) - f(y)| \leq L|x - y| \quad \text{for all } x, y \in \mathbb{R}.$$

Fix $x_0 \in \mathbb{R}$, and define $x_{n+1} = f(x_n)$. Then (x_n) converges to a point $x^* \in \mathbb{R}$,

$$f(x^*) = x^*,$$

this fixed point is unique, every starting point converges to it, and

$$|x_n - x^*| \leq L^n |x_0 - x^*|, \quad |x_n - x^*| \leq \frac{L^n}{1 - L} |x_1 - x_0|.$$

Remark (Abstract). This capstone is the chapter's completeness payoff. The hypothesis is only a distance-shrinking inequality; no continuity theorem is used. The orbit first proves itself Cauchy by geometric decay of its successive steps. Completeness then supplies a limit, and only afterward does the contraction inequality identify that limit as the fixed point.

Remark (Orientation). The orbit is the sequence

$$x_0, \quad f(x_0), \quad f(f(x_0)), \quad \dots$$

A fixed point is a point x^* satisfying $f(x^*) = x^*$. The contraction hypothesis says that applying f pulls every pair of points at least a factor $L < 1$ closer. The proof cannot aim at x^* , because no fixed point has been produced yet. Instead it shows that the orbit crowds together internally, invokes the Cauchy criterion, and then checks what the limit must be.

Remark (Step 1: one step is controlled by the previous step). For every $n \geq 1$,

$$|x_{n+1} - x_n| = |f(x_n) - f(x_{n-1})| \leq L|x_n - x_{n-1}|.$$

This is the contraction hypothesis applied to the pair (x_n, x_{n-1}) .

Remark (Step 2: every step is controlled by the first step). For every $n \geq 0$,

$$|x_{n+1} - x_n| \leq L^n |x_1 - x_0|.$$

The case $n = 0$ is equality. If the estimate holds at $n - 1$, then Step 1 gives

$$|x_{n+1} - x_n| \leq L|x_n - x_{n-1}| \leq L \cdot L^{n-1} |x_1 - x_0| = L^n |x_1 - x_0|.$$

Induction proves the estimate.

Remark (Step 3: the orbit is Cauchy). Let $m > n$. Telescoping and applying the triangle inequality gives

$$|x_m - x_n| = \left| \sum_{k=n}^{m-1} (x_{k+1} - x_k) \right| \leq \sum_{k=n}^{m-1} |x_{k+1} - x_k| \leq \sum_{k=n}^{m-1} L^k |x_1 - x_0|.$$

Since $0 \leq L < 1$,

$$\sum_{k=n}^{m-1} L^k = L^n \frac{1 - L^{m-n}}{1 - L} \leq \frac{L^n}{1 - L}.$$

Hence

$$|x_m - x_n| \leq \frac{L^n}{1 - L} |x_1 - x_0|. \quad (*)$$

Because $L^n \rightarrow 0$, the right-hand side tends to 0. Thus for every $\varepsilon > 0$ there is N such that $m > n \geq N$ implies $|x_m - x_n| < \varepsilon$. The orbit is Cauchy.

Remark (Step 4: completeness produces the limit). By the Cauchy criterion for real sequences, every Cauchy sequence of real numbers converges. Therefore there is $x^* \in \mathbb{R}$ such that

$$x_n \rightarrow x^*.$$

This is the only step in which a limit not given by a formula is brought into existence.

Remark (Step 5: the limit is a fixed point). For every n ,

$$\begin{aligned} |x^* - f(x^*)| &\leq |x^* - x_{n+1}| + |x_{n+1} - f(x^*)| \\ &= |x^* - x_{n+1}| + |f(x_n) - f(x^*)| \\ &\leq |x^* - x_{n+1}| + L|x_n - x^*|. \end{aligned}$$

The shifted sequence x_{n+1} also converges to x^* , and $|x_n - x^*| \rightarrow 0$. The right-hand side therefore tends to 0. Since $|x^* - f(x^*)|$ is a fixed nonnegative number bounded above by a null sequence, it is 0. Hence $f(x^*) = x^*$.

Remark (Step 6: uniqueness and convergence from every start). If p and q are fixed points, then

$$|p - q| = |f(p) - f(q)| \leq L|p - q|.$$

Thus

$$0 \leq (1 - L)|p - q| \leq 0.$$

Since $1 - L > 0$, we have $|p - q| = 0$, so $p = q$. The construction above started from an arbitrary x_0 , and uniqueness forces every such orbit to converge to the same fixed point.

Remark (Step 7: rates). Once x^* is known to be fixed,

$$|x_n - x^*| = |f(x_{n-1}) - f(x^*)| \leq L|x_{n-1} - x^*|.$$

Induction gives

$$|x_n - x^*| \leq L^n |x_0 - x^*|.$$

For the computable estimate, fix n in $(*)$ and let $m \rightarrow \infty$. Using $x_m \rightarrow x^*$ and preservation of eventual order under limits gives

$$|x_n - x^*| \leq \frac{L^n}{1 - L} |x_1 - x_0|.$$

Remark (Limits of the theorem). The strict inequality $L < 1$ is essential. At $L = 1$, a map may preserve distances without having any fixed point; for example $f(x) = x + 1$. The general theorem replaces $\mathbb{R}$ by any complete metric space and $|x - y|$ by the metric $d(x, y)$. That version is the Banach fixed-point theorem and belongs to the metric-spaces layer.

Remark (Forward connection). Later, the continuity chapter names the hypothesis as a Lipschitz condition with constant $L < 1$. Here the name is deliberately avoided: the proof uses only the inequality, geometric decay, and Cauchy completeness. The same iteration idea becomes Picard iteration when differential equations are recast as fixed-point problems for integral operators.

Remark (Dependencies to state). • [Sequence](#)

- [Convergence of a Sequence](#)
- [Cauchy Sequence](#)

Remark (Dependencies to prove). • [Cauchy Criterion for Real Sequences](#)

- [Ratio Limit Less Than One Implies Null](#)
- [Subsequences Preserve Limits](#)
- [Constant Comparison for Sequence Limits](#)
- [Limit Preserves Eventual Order](#)

Remark (Dependency ceiling). This capstone may use only real-number arithmetic, absolute value and the triangle inequality, finite geometric sums, and sequence results routed at or before the Sequences chapter. It must not use continuity, Lipschitz maps as a named concept, metric-space completeness, or the Banach fixed-point theorem.

Remark (Dependencies).

- [Sequence](#)
- [Convergence of a Sequence](#)
- [Cauchy Sequence](#)
- [Cauchy Criterion for Real Sequences](#)
- [Ratio Limit Less Than One Implies Null](#)
- [Subsequences Preserve Limits](#)
- [Constant Comparison for Sequence Limits](#)
- [Limit Preserves Eventual Order](#)

Chapter 7

Series



7.1 Finite Series

Remark (Exposition). Tao introduces finite series by first fixing a finite real sequence indexed by consecutive integers and then defining its sum recursively [Tao \[2006\]](#). The source bundles these conventions into one definition; here they are split into atomic definitions.

Definition (Finite Real Sequence on an Integer Interval)

Definition 7.1.1 (Finite Real Sequence on an Integer Interval). Let $m, n \in \mathbb{Z}$ with $m \leq n$. A finite real sequence indexed from m to n is an assignment of a real number a_i to each integer i satisfying $m \leq i \leq n$. We denote such a sequence by $(a_i)_{i=m}^n$.

Remark (Interpretation). The indices form the integer interval $\{m, m+1, \dots, n\}$. The notation $(a_i)_{i=m}^n$ records both the terms and the finite range of indices on which the terms are defined.

Remark (Dependencies).

- [Real Sequence](#)

Definition (Empty Finite Sum)

Definition 7.1.2 (Empty Finite Sum). Let $m, n \in \mathbb{Z}$. If $n < m$, the finite sum from m to n is defined by

$$\sum_{i=m}^n a_i := 0.$$

Remark (Interpretation). When the upper index lies below the lower index, there are no terms to add. The sum is therefore assigned the additive identity 0.

Remark (Dependencies).

- [Finite Real Sequence on an Integer Interval](#)

Definition (Finite Sum)

Definition 7.1.3 (Finite Sum). Let $m, n \in \mathbb{Z}$, and let real numbers a_i be assigned for the relevant integer indices. The notation

$$\sum_{i=m}^n a_i$$

denotes the finite sum, or finite series, of the terms indexed from m to n .

Remark (Standard quantified statement).

$$\sum_{i=m}^n a_i \quad \text{denotes the finite sum of the terms } a_i \text{ indexed by } m \leq i \leq n.$$

Remark (Interpretation). This definition fixes the notation and name of the object being defined. The value of the object is then determined by the empty-sum convention and the recursive rule below.

Remark (Dependencies).

- [Finite Real Sequence on an Integer Interval](#)

Definition (Recursive Finite Sum)

Definition 7.1.4 (Recursive Finite Sum). Let $m \in \mathbb{Z}$, and let real numbers a_i be assigned for the relevant indices. The finite sum, also called a finite series, is defined recursively from the empty finite sum by the rule

$$\sum_{i=m}^{n+1} a_i := \left(\sum_{i=m}^n a_i \right) + a_{n+1} \quad \text{whenever } n \geq m - 1.$$

Remark (Standard quantified statement).

$$\forall m, n \in \mathbb{Z}, \quad n \geq m - 1 \implies \sum_{i=m}^{n+1} a_i := \left(\sum_{i=m}^n a_i \right) + a_{n+1}.$$

Remark (Interpretation). The recursive rule says that extending the upper endpoint by one index extends the sum by adding exactly the new last term. Together with the empty-sum convention, this determines every finite sum.

Remark (Examples). For a sequence indexed from m , the first few cases are

$$\sum_{i=m}^{m-2} a_i = 0, \quad \sum_{i=m}^{m-1} a_i = 0, \quad \sum_{i=m}^m a_i = a_m,$$

and

$$\sum_{i=m}^{m+1} a_i = a_m + a_{m+1}, \quad \sum_{i=m}^{m+2} a_i = a_m + a_{m+1} + a_{m+2}.$$

Remark (Dependencies).

- [Empty Finite Sum](#)
- [Finite Sum](#)
- [Finite Real Sequence on an Integer Interval](#)

Lemma (Splitting a Finite Sum)

Lemma 7.1.5 (Splitting a Finite Sum). *Let $m \leq n < p$ be integers, and let a_i be a real number assigned to each integer i satisfying $m \leq i \leq p$. Then*

$$\sum_{i=m}^n a_i + \sum_{i=n+1}^p a_i = \sum_{i=m}^p a_i.$$

Go to proof.

Remark (Interpretation). A finite sum over one integer interval may be split into two adjacent finite sums without changing its value.

Remark (Dependencies).

- [Recursive Finite Sum](#)

Definition (Summation over a Finite Set)

Definition 7.1.6 (Summation over a Finite Set). Let X be a finite set with n elements, where $n \in \mathbb{N}$, and let $f : X \rightarrow \mathbb{R}$. Choose a bijection

$$g : \{i \in \mathbb{N} : 1 \leq i \leq n\} \rightarrow X.$$

The finite sum of f over X is defined by

$$\sum_{x \in X} f(x) := \sum_{i=1}^n f(g(i)).$$

Remark (Standard quantified statement).

$$X = \{g(1), \dots, g(n)\} \implies \sum_{x \in X} f(x) := \sum_{i=1}^n f(g(i)).$$

Remark (Interpretation). The definition computes a sum over an unordered finite set by first listing its elements through a bijection from $\{1, \dots, n\}$. The notation $\sum_{x \in X} f(x)$ records the set being summed over; the chosen listing is auxiliary.

Remark (Dependencies).

- [Finite Sum](#)

Proposition (Finite Summations Are Well-Defined)

Proposition 7.1.7 (Finite Summations Are Well-Defined). *Let X be a finite set with n elements, where $n \in \mathbb{N}$, let $f : X \rightarrow \mathbb{R}$, and let*

$$g : \{i \in \mathbb{N} : 1 \leq i \leq n\} \rightarrow X \quad \text{and} \quad h : \{i \in \mathbb{N} : 1 \leq i \leq n\} \rightarrow X$$

be bijections. Then

$$\sum_{i=1}^n f(g(i)) = \sum_{i=1}^n f(h(i)).$$

[Go to proof.](#)

Remark (Interpretation). The value of a finite summation over a finite set does not depend on which bijection is used to list the elements of that set. Tao notes that the corresponding issue for summations over infinite sets is more complicated.

Remark (Dependencies).

- [Summation over a Finite Set](#)

Proposition (Empty Finite Set Summation)

Proposition 7.1.8 (Empty Finite Set Summation). *If X is empty, and $f : X \rightarrow \mathbb{R}$ is a function, then*

$$\sum_{x \in X} f(x) = 0.$$

[Go to proof.](#)

Remark (Interpretation). The finite summation over the empty set has value 0.

Remark (Dependencies).

- [Summation over a Finite Set](#)

Proposition (Singleton Finite Set Summation)

Proposition 7.1.9 (Singleton Finite Set Summation). *If X consists of a single element, $X = \{x_0\}$, and $f : X \rightarrow \mathbb{R}$ is a function, then*

$$\sum_{x \in X} f(x) = f(x_0).$$

[Go to proof.](#)

Remark (Interpretation). Summing a function over a one-element set gives the value of the function at that element.

Remark (Dependencies).

- [Summation over a Finite Set](#)

Proposition (Substitution for Finite Set Summation)

Proposition 7.1.10 (Substitution for Finite Set Summation). *If X is a finite set, $f : X \rightarrow \mathbb{R}$ is a function, and $g : Y \rightarrow X$ is a bijection, then*

$$\sum_{x \in X} f(x) = \sum_{y \in Y} f(g(y)).$$

[Go to proof.](#)

Remark (Interpretation). A finite summation over X may be computed by summing over any set bijective with X , after composing the summand with the bijection.

Remark (Dependencies).

- [Summation over a Finite Set](#)
- [Finite Summations Are Well-Defined](#)

Proposition (Integer Interval as a Finite Set)

Proposition 7.1.11 (Integer Interval as a Finite Set). *Let $n \leq m$ be integers, and let*

$$X := \{i \in \mathbb{Z} : n \leq i \leq m\}.$$

If a_i is a real number assigned to each integer $i \in X$, then

$$\sum_{i=n}^m a_i = \sum_{i \in X} a_i.$$

[Go to proof.](#)

Remark (Interpretation). Summing over a consecutive integer interval agrees with summing over that same interval regarded as a finite set.

Remark (Dependencies).

- [Finite Sum](#)
- [Summation over a Finite Set](#)

Proposition (Disjoint Union Finite Set Summation)

Proposition 7.1.12 (Disjoint Union Finite Set Summation). *Let X and Y be disjoint finite sets, so $X \cap Y = \emptyset$, and let $f : X \cup Y \rightarrow \mathbb{R}$ be a function. Then*

$$\sum_{z \in X \cup Y} f(z) = \left(\sum_{x \in X} f(x) \right) + \left(\sum_{y \in Y} f(y) \right).$$

[Go to proof.](#)

Remark (Interpretation). A sum over a disjoint union splits into the sum over each component set.

Remark (Dependencies).

- [Summation over a Finite Set](#)

Proposition (Addition Rule for Finite Set Summation)

Proposition 7.1.13 (Addition Rule for Finite Set Summation). *Let X be a finite set, and let $f : X \rightarrow \mathbb{R}$ and $g : X \rightarrow \mathbb{R}$ be functions. Then*

$$\sum_{x \in X} (f(x) + g(x)) = \sum_{x \in X} f(x) + \sum_{x \in X} g(x).$$

[Go to proof.](#)

Remark (Interpretation). Finite set summation distributes over pointwise addition.

Remark (Dependencies).

- [Summation over a Finite Set](#)

Proposition (Scalar Multiplication Rule for Finite Set Summation)

Proposition 7.1.14 (Scalar Multiplication Rule for Finite Set Summation). *Let X be a finite set, let $f : X \rightarrow \mathbb{R}$ be a function, and let c be a real number. Then*

$$\sum_{x \in X} cf(x) = c \sum_{x \in X} f(x).$$

[Go to proof.](#)

Remark (Interpretation). A constant scalar may be pulled outside a finite set summation.

Remark (Dependencies).

- [Summation over a Finite Set](#)

Proposition (Monotonicity for Finite Set Summation)

Proposition 7.1.15 (Monotonicity for Finite Set Summation). *Let X be a finite set, and let $f : X \rightarrow \mathbb{R}$ and $g : X \rightarrow \mathbb{R}$ be functions such that $f(x) \leq g(x)$ for all $x \in X$. Then*

$$\sum_{x \in X} f(x) \leq \sum_{x \in X} g(x).$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$(\forall x \in X, f(x) \leq g(x)) \implies \sum_{x \in X} f(x) \leq \sum_{x \in X} g(x).$$

Remark (Interpretation). Pointwise order between two functions on a finite set gives the same order between their finite sums.

Remark (Dependencies).

- [Summation over a Finite Set](#)

Proposition (Triangle Inequality for Finite Set Summation)

Proposition 7.1.16 (Triangle Inequality for Finite Set Summation). *Let X be a finite set, and let $f : X \rightarrow \mathbb{R}$ be a function. Then*

$$\left| \sum_{x \in X} f(x) \right| \leq \sum_{x \in X} |f(x)|.$$

[Go to proof.](#)

Remark (Interpretation). The absolute value of a finite set summation is bounded by the corresponding finite set summation of absolute values.

Remark (Dependencies).

- [Summation over a Finite Set](#)

Lemma (Iterated Summation over a Product)

Lemma 7.1.17 (Iterated Summation over a Product). *Let X and Y be finite sets, and let $f : X \times Y \rightarrow \mathbb{R}$ be a function. Then*

$$\sum_{x \in X} \left(\sum_{y \in Y} f(x, y) \right) = \sum_{(x, y) \in X \times Y} f(x, y).$$

[Go to proof.](#)

Remark (Interpretation). Summing first over Y for each $x \in X$, and then summing those values over X , gives the same value as summing over the product set $X \times Y$.

Remark (Dependencies).

- [Summation over a Finite Set](#)

Corollary (Fubini's Theorem for Finite Series)

Corollary 7.1.18 (Fubini's Theorem for Finite Series). *Let X and Y be finite sets, and let $f : X \times Y \rightarrow \mathbb{R}$ be a function. Then*

$$\sum_{x \in X} \left(\sum_{y \in Y} f(x, y) \right) = \sum_{(x, y) \in X \times Y} f(x, y) = \sum_{(y, x) \in Y \times X} f(x, y) = \sum_{y \in Y} \left(\sum_{x \in X} f(x, y) \right).$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\sum_{x \in X} \left(\sum_{y \in Y} f(x, y) \right) = \sum_{y \in Y} \left(\sum_{x \in X} f(x, y) \right).$$

Remark (Interpretation). For a function on a finite product set, the order of summation over the two finite factors may be interchanged.

Remark (Dependencies).

- [Iterated Summation over a Product](#)
- [Summation over a Finite Set](#)

Lemma (Reindexing a Finite Sum)

Lemma 7.1.19 (Reindexing a Finite Sum). *Let $m \leq n$ be integers, let k be an integer, and let a_i be a real number assigned to each integer i satisfying $m \leq i \leq n$. Then*

$$\sum_{i=m}^n a_i = \sum_{j=m+k}^{n+k} a_{j-k}.$$

[Go to proof.](#)

Remark (Interpretation). Shifting every index by the same integer changes the labels of the summation but not the ordered list of terms being summed.

Remark (Dependencies).

- [Recursive Finite Sum](#)

Lemma (Finite Sum of Sums)

Lemma 7.1.20 (Finite Sum of Sums). *Let $m \leq n$ be integers, and let a_i and b_i be real numbers assigned to each integer i satisfying $m \leq i \leq n$. Then*

$$\sum_{i=m}^n (a_i + b_i) = \left(\sum_{i=m}^n a_i \right) + \left(\sum_{i=m}^n b_i \right).$$

[Go to proof.](#)

Remark (Interpretation). Finite summation distributes over termwise addition.

Remark (Dependencies).

- [Recursive Finite Sum](#)

Lemma (Finite Sum of Scalar Multiples)

Lemma 7.1.21 (Finite Sum of Scalar Multiples). *Let $m \leq n$ be integers, let a_i be a real number assigned to each integer i satisfying $m \leq i \leq n$, and let $c \in \mathbb{R}$. Then*

$$\sum_{i=m}^n (ca_i) = c \left(\sum_{i=m}^n a_i \right).$$

[Go to proof.](#)

Remark (Interpretation). A common scalar factor may be pulled outside a finite sum.

Remark (Dependencies).

- [Recursive Finite Sum](#)

Lemma (Triangle Inequality for Finite Series)

Lemma 7.1.22 (Triangle Inequality for Finite Series). *Let $m \leq n$ be integers, and let a_i be a real number assigned to each integer i satisfying $m \leq i \leq n$. Then*

$$\left| \sum_{i=m}^n a_i \right| \leq \sum_{i=m}^n |a_i|.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall m, n \in \mathbb{Z}, \quad m \leq n \implies \left| \sum_{i=m}^n a_i \right| \leq \sum_{i=m}^n |a_i|.$$

Remark (Interpretation). The absolute value of a finite series is bounded by the corresponding finite series of absolute values.

Remark (Dependencies).

- [Recursive Finite Sum](#)

Lemma (Comparison Test for Finite Series)

Lemma 7.1.23 (Comparison Test for Finite Series). *Let $m \leq n$ be integers, and let a_i and b_i be real numbers assigned to each integer i satisfying $m \leq i \leq n$. Suppose that $a_i \leq b_i$ for all i satisfying $m \leq i \leq n$. Then*

$$\sum_{i=m}^n a_i \leq \sum_{i=m}^n b_i.$$

[Go to proof.](#)

Remark (Standard quantified statement).

$$\forall i (m \leq i \leq n \implies a_i \leq b_i) \implies \sum_{i=m}^n a_i \leq \sum_{i=m}^n b_i.$$

Remark (Interpretation). Termwise comparison of two finite series gives comparison of their finite sums.

Remark (Dependencies).

- [Recursive Finite Sum](#)

7.2 Infinite Series

Definition (Formal Infinite Series)

Definition 7.2.1 (Formal Infinite Series). A formal infinite series is an expression of the form

$$\sum_{n=m}^{\infty} a_n,$$

where m is an integer, and a_n is a real number for every integer $n \geq m$. This series is sometimes written as

$$a_m + a_{m+1} + a_{m+2} + \cdots.$$

Remark (Standard quantified statement).

$\sum_{n=m}^{\infty} a_n$ denotes a formal infinite series with $m \in \mathbb{Z}$ and $a_n \in \mathbb{R}$ for every $n \geq m$.

Remark (Interpretation). The notation records an infinite list of real terms beginning at the integer index m . At this stage it is a formal expression rather than a claim that a value has already been assigned to the series.

Remark (Dependencies).

- [Real Sequence](#)

Proofs

Remark (Return). [Return to Lemma](#)

Lemma (Splitting a Finite Sum). *Let $m \leq n < p$ be integers, and let a_i be a real number assigned to each integer i satisfying $m \leq i \leq p$. Then*

$$\sum_{i=m}^n a_i + \sum_{i=n+1}^p a_i = \sum_{i=m}^p a_i.$$

Professional Standard Proof.

TODO: professional standard proof for finite-sum-splitting. □

Detailed Learning Proof.

TODO: detailed learning proof for finite-sum-splitting. □

Remark (Proof structure).

Remark (Dependencies).

- [Recursive Finite Sum](#)

Remark (Return). [Return to Proposition](#)

Proposition (Finite Summations Are Well-Defined). *Let X be a finite set with n elements, where $n \in \mathbb{N}$, let $f : X \rightarrow \mathbb{R}$, and let*

$$g : \{i \in \mathbb{N} : 1 \leq i \leq n\} \rightarrow X \quad \text{and} \quad h : \{i \in \mathbb{N} : 1 \leq i \leq n\} \rightarrow X$$

be bijections. Then

$$\sum_{i=1}^n f(g(i)) = \sum_{i=1}^n f(h(i)).$$

Professional Standard Proof.

TODO: professional standard proof for finite-summations-well-defined. □

Detailed Learning Proof.

TODO: detailed learning proof for finite-summations-well-defined. □

Remark (Proof structure).

Remark (Dependencies).

- [Summation over a Finite Set](#)

Remark (Return). [Return to Proposition](#)

Proposition (Empty Finite Set Summation). *If X is empty, and $f : X \rightarrow \mathbb{R}$ is a function, then*

$$\sum_{x \in X} f(x) = 0.$$

Professional Standard Proof.

TODO: professional standard proof for empty-finite-set-summation. □

Detailed Learning Proof.

TODO: detailed learning proof for empty-finite-set-summation. □

Remark (Proof structure).

Remark (Dependencies).

- [Summation over a Finite Set](#)

Remark (Return). [Return to Proposition](#)

Proposition (Singleton Finite Set Summation). *If X consists of a single element, $X = \{x_0\}$, and $f : X \rightarrow \mathbb{R}$ is a function, then*

$$\sum_{x \in X} f(x) = f(x_0).$$

Professional Standard Proof.

TODO: professional standard proof for singleton-finite-set-summation. □

Detailed Learning Proof.

TODO: detailed learning proof for singleton-finite-set-summation. □

Remark (Proof structure).

Remark (Dependencies).

- [Summation over a Finite Set](#)

Remark (Return). [Return to Proposition](#)

Proposition (Substitution for Finite Set Summation). *If X is a finite set, $f : X \rightarrow \mathbb{R}$ is a function, and $g : Y \rightarrow X$ is a bijection, then*

$$\sum_{x \in X} f(x) = \sum_{y \in Y} f(g(y)).$$

Professional Standard Proof.

TODO: professional standard proof for substitution-finite-set-summation. □

Detailed Learning Proof.

TODO: detailed learning proof for substitution-finite-set-summation. □

Remark (Proof structure).

Remark (Dependencies).

- [Summation over a Finite Set](#)
- [Finite Summations Are Well-Defined](#)

Remark (Return). [Return to Proposition](#)

Proposition (Integer Interval as a Finite Set). *Let $n \leq m$ be integers, and let*

$$X := \{i \in \mathbb{Z} : n \leq i \leq m\}.$$

If a_i is a real number assigned to each integer $i \in X$, then

$$\sum_{i=n}^m a_i = \sum_{i \in X} a_i.$$

Professional Standard Proof.

TODO: professional standard proof for integer-interval-finite-set-summation. □

Detailed Learning Proof.

TODO: detailed learning proof for integer-interval-finite-set-summation. □

Remark (Proof structure).

Remark (Dependencies).

- [Finite Sum](#)
- [Summation over a Finite Set](#)

Remark (Return). [Return to Proposition](#)

Proposition (Disjoint Union Finite Set Summation). *Let X and Y be disjoint finite sets, so $X \cap Y = \emptyset$, and let $f : X \cup Y \rightarrow \mathbb{R}$ be a function. Then*

$$\sum_{z \in X \cup Y} f(z) = \left(\sum_{x \in X} f(x) \right) + \left(\sum_{y \in Y} f(y) \right).$$

Professional Standard Proof.

TODO: professional standard proof for disjoint-union-finite-set-summation. □

Detailed Learning Proof.

TODO: detailed learning proof for disjoint-union-finite-set-summation. □

Remark (Proof structure).

Remark (Dependencies).

- [Summation over a Finite Set](#)

Remark (Return). [Return to Proposition](#)

Proposition (Addition Rule for Finite Set Summation). *Let X be a finite set, and let $f : X \rightarrow \mathbb{R}$ and $g : X \rightarrow \mathbb{R}$ be functions. Then*

$$\sum_{x \in X} (f(x) + g(x)) = \sum_{x \in X} f(x) + \sum_{x \in X} g(x).$$

Professional Standard Proof.

TODO: professional standard proof for addition-rule-finite-set-summation. □

Detailed Learning Proof.

TODO: detailed learning proof for addition-rule-finite-set-summation. □

Remark (Proof structure).

Remark (Dependencies).

- [Summation over a Finite Set](#)

Remark (Return). [Return to Proposition](#)

Proposition (Scalar Multiplication Rule for Finite Set Summation). *Let X be a finite set, let $f : X \rightarrow \mathbb{R}$ be a function, and let c be a real number. Then*

$$\sum_{x \in X} cf(x) = c \sum_{x \in X} f(x).$$

Professional Standard Proof.

TODO: professional standard proof for scalar-multiplication-rule-finite-set-summation. □

Detailed Learning Proof.

TODO: detailed learning proof for scalar-multiplication-rule-finite-set-summation. □

Remark (Proof structure).

Remark (Dependencies).

- [Summation over a Finite Set](#)

Remark (Return). [Return to Proposition](#)

Proposition (Monotonicity for Finite Set Summation). *Let X be a finite set, and let $f : X \rightarrow \mathbb{R}$ and $g : X \rightarrow \mathbb{R}$ be functions such that $f(x) \leq g(x)$ for all $x \in X$. Then*

$$\sum_{x \in X} f(x) \leq \sum_{x \in X} g(x).$$

Professional Standard Proof.

TODO: professional standard proof for monotonicity-finite-set-summation. □

Detailed Learning Proof.

TODO: detailed learning proof for monotonicity-finite-set-summation. □

Remark (Proof structure).

Remark (Dependencies).

- [Summation over a Finite Set](#)

Remark (Return). [Return to Proposition](#)

Proposition (Triangle Inequality for Finite Set Summation). *Let X be a finite set, and let $f : X \rightarrow \mathbb{R}$ be a function. Then*

$$\left| \sum_{x \in X} f(x) \right| \leq \sum_{x \in X} |f(x)|.$$

Professional Standard Proof.

TODO: professional standard proof for triangle-inequality-finite-set-summation. □

Detailed Learning Proof.

TODO: detailed learning proof for triangle-inequality-finite-set-summation. □

Remark (Proof structure).

Remark (Dependencies).

- [Summation over a Finite Set](#)

Remark (Return). [Return to Lemma](#)

Lemma (Iterated Summation over a Product). *Let X and Y be finite sets, and let $f : X \times Y \rightarrow \mathbb{R}$ be a function. Then*

$$\sum_{x \in X} \left(\sum_{y \in Y} f(x, y) \right) = \sum_{(x, y) \in X \times Y} f(x, y).$$

Professional Standard Proof.

TODO: professional standard proof for iterated-summation-over-product. □

Detailed Learning Proof.

TODO: detailed learning proof for iterated-summation-over-product. □

Remark (Proof structure).

Remark (Dependencies).

- [Summation over a Finite Set](#)

Remark (Return). [Return to Corollary](#)

Corollary (Fubini's Theorem for Finite Series). *Let X and Y be finite sets, and let $f : X \times Y \rightarrow \mathbb{R}$ be a function. Then*

$$\sum_{x \in X} \left(\sum_{y \in Y} f(x, y) \right) = \sum_{(x, y) \in X \times Y} f(x, y) = \sum_{(y, x) \in Y \times X} f(x, y) = \sum_{y \in Y} \left(\sum_{x \in X} f(x, y) \right).$$

Professional Standard Proof.

TODO: professional standard proof for fubini-theorem-finite-series. □

Detailed Learning Proof.

TODO: detailed learning proof for fubini-theorem-finite-series. □

Remark (Proof structure).

Remark (Dependencies).

- [Iterated Summation over a Product](#)
- [Summation over a Finite Set](#)

Remark (Return). [Return to Lemma](#)

Lemma (Reindexing a Finite Sum). *Let $m \leq n$ be integers, let k be an integer, and let a_i be a real number assigned to each integer i satisfying $m \leq i \leq n$. Then*

$$\sum_{i=m}^n a_i = \sum_{j=m+k}^{n+k} a_{j-k}.$$

Professional Standard Proof.

TODO: professional standard proof for finite-sum-reindexing. □

Detailed Learning Proof.

TODO: detailed learning proof for finite-sum-reindexing. □

Remark (Proof structure).

Remark (Dependencies).

- [Recursive Finite Sum](#)

Remark (Return). [Return to Lemma](#)

Lemma (Finite Sum of Sums). *Let $m \leq n$ be integers, and let a_i and b_i be real numbers assigned to each integer i satisfying $m \leq i \leq n$. Then*

$$\sum_{i=m}^n (a_i + b_i) = \left(\sum_{i=m}^n a_i \right) + \left(\sum_{i=m}^n b_i \right).$$

Professional Standard Proof.

TODO: professional standard proof for finite-sum-of-sums. □

Detailed Learning Proof.

TODO: detailed learning proof for finite-sum-of-sums. □

Remark (Proof structure).

Remark (Dependencies).

- [Recursive Finite Sum](#)

Remark (Return). [Return to Lemma](#)

Lemma (Finite Sum of Scalar Multiples). *Let $m \leq n$ be integers, let a_i be a real number assigned to each integer i satisfying $m \leq i \leq n$, and let $c \in \mathbb{R}$. Then*

$$\sum_{i=m}^n (ca_i) = c \left(\sum_{i=m}^n a_i \right).$$

Professional Standard Proof.

TODO: professional standard proof for finite-sum-scalar-multiple. □

Detailed Learning Proof.

TODO: detailed learning proof for finite-sum-scalar-multiple. □

Remark (Proof structure).

Remark (Dependencies).

- [Recursive Finite Sum](#)

Remark (Return). [Return to Lemma](#)

Lemma (Triangle Inequality for Finite Series). *Let $m \leq n$ be integers, and let a_i be a real number assigned to each integer i satisfying $m \leq i \leq n$. Then*

$$\left| \sum_{i=m}^n a_i \right| \leq \sum_{i=m}^n |a_i|.$$

Professional Standard Proof.

TODO: professional standard proof for triangle-inequality-finite-series. □

Detailed Learning Proof.

TODO: detailed learning proof for triangle-inequality-finite-series. □

Remark (Proof structure).

Remark (Dependencies).

- [Recursive Finite Sum](#)

Remark (Return). [Return to Lemma](#)

Lemma (Comparison Test for Finite Series). *Let $m \leq n$ be integers, and let a_i and b_i be real numbers assigned to each integer i satisfying $m \leq i \leq n$. Suppose that $a_i \leq b_i$ for all i satisfying $m \leq i \leq n$. Then*

$$\sum_{i=m}^n a_i \leq \sum_{i=m}^n b_i.$$

Professional Standard Proof.

TODO: professional standard proof for comparison-test-finite-series. □

Detailed Learning Proof.

TODO: detailed learning proof for comparison-test-finite-series. □

Remark (Proof structure).

Remark (Dependencies).

- [Recursive Finite Sum](#)

Chapter 8

Function Sequences



Proofs

Bibliography

Terence Tao. *Analysis I*, volume 37 of *Texts and Readings in Mathematics*. Hindustan Book Agency, New Delhi, 2006. ISBN 978-8185931751.