

RICE UNIVERSITY

**Impacts of Two-fluid Effects on Emission from
Impulsively Heated Solar Coronal Loops**

by

Will T. Barnes

A THESIS SUBMITTED
IN PARTIAL FULFILLMENT OF THE
REQUIREMENTS FOR THE DEGREE

Master of Science

APPROVED, THESIS COMMITTEE:

Stephen J. Bradshaw, Chair
Assistant Professor of Physics and
Astronomy

David Alexander
Professor of Physics and Astronomy

Marjorie Corcoran
Professor of Physics and Astronomy

Houston, Texas

December, 2015

ABSTRACT

Impacts of Two-fluid Effects on Emission from Impulsively Heated Solar Coronal
Loops

by

Will T. Barnes

The coronal heating problem has puzzled physicists and astronomers for nearly a century. While the origin of these surprisingly high temperatures in the upper solar atmosphere is most certainly magnetic,

Contents

Abstract	ii
List of Illustrations	iv
List of Tables	v
1 Introduction	1
1.1 Structure of the Solar Atmosphere	3
1.2 The Solar Magnetic Field	6
1.3 The Coronal Heating Problem	8
1.4 Summary	9
2 Coronal Loops	10
2.1 Observations	10
2.2 Modeling	10
3 Emission Diagnostics	11
4 Numerical Modeling	12
4.1 One-dimensional Hydrodynamics	12
4.2 The EBTEL Model	13
4.3 The Two-fluid EBTEL Model	14
5 Results	15
6 Conclusions	16

Illustrations

1.1	Total eclipse as seen from Svalbard, Norway on March 20, 2015. Open and closed loops in the highly-structured solar corona are clearly visible. Photo courtesy of Miloslav Druckmüller.	2
1.2	Layers of the solar atmosphere (a) Courtesy of NASA (b) Taken from Gary et al. (2007)	5
1.3	Taken from Gary (2001).	7

Tables

Chapter 1

Introduction

Intro to the first chapter Give a nice history of solar observations and discuss current observing efforts as well as modeling efforts, but briefly; quickly move into atmosphere, corona

The Sun is the most important celestial body to life on Earth. For the last five billion years, it has provided the light by which humans observe the world around them and the heat to save the planet from the frigid temperatures of interplanetary space. Because of its proximity, the Sun provides astronomers an exclusive and unique look into how stars behave. By observing and understanding the Sun, one can make conclusions about other types of stars in our galaxy and the universe.

Perhaps no other consistent celestial event has attracted as much attention as the solar eclipse. Solar eclipses have been observed and recorded for thousands of years, with some reports dating back to the fourteenth century BC (Golub & Pasachoff, 2010). The recordings of ancient eclipses have been heavily studied. Chinese rock drawings dating back to the Han dynasty (approximately 1900 years ago) appear to show the moon completely obscuring the Sun. Additionally, some have even suggested the Aubrey holes that surround the Stonehenge site were used to track and predict both solar and lunar eclipses (Golub & Pasachoff, 2010). Though some claims of



Figure 1.1 : Total eclipse as seen from Svalbard, Norway on March 20, 2015. Open and closed loops in the highly-structured solar corona are clearly visible. Photo courtesy of Miloslav Druckmüller.

ancient eclipse studies are controversial, it suffices to say that humans have long sought to study and explain the behavior of the nearest star to Earth.

In particular, solar eclipses have captured the attention of artists for centuries. Cosmas Damian Asam, a Bavarian painter and architect active in the early eighteenth century, used images of solar eclipses in many of his works, including several frescoes and an altarpiece. Olson & Pasachoff (2007) discuss how Asam, a deeply religious artist, was commissioned several times to depict the vision of St. Gregory the Great, a Benedictine monk, as described in his work *Dialogues*.

One of these depictions, an altarpiece at a Benedictine monastery in Kladruby, Czech Republic (see Fig. 7 of Olson & Pasachoff (2007)), shows “the visionary globe surrounded by a glowing halo of yellow light that more closely resembles the solar

corona” (Olson & Pasachoff, 2007). An additional altarpiece at another monastery in Weltenburg, Germany shows a perhaps even more pronounced depiction of the solar corona during a total solar eclipse. Olson & Pasachoff (2007) note that, given his detailed depictions, Asam must have observed several solar eclipses as well as the solar corona, with these astronomical events profoundly impacting his depictions of supernatural events in his works. This is but one example of how solar eclipses and their consequential insight into the highly structured solar atmosphere, have shaped scientific and artistic discourse throughout history.

1.1 Structure of the Solar Atmosphere

This section will discuss the structure of the solar atmosphere including the different layers of the Sun and how they are connected. This will help to introduce the solar corona

Though the Sun can be easily seen from Earth, its dynamic and highly structured atmosphere is not observable with the naked eye, with the one exception being brief glimpses of the corona during an eclipse. The interior of the Sun is of course very complex and constitutes a very different regime of physics than that seen in the solar atmosphere. Thus, this work will be primarily limited to the upper solar atmosphere with some discussion of the lower layers.

The solar atmosphere is often divided up into four separate regions: the photosphere, the chromosphere, the transition region, and the corona. Fig. 1.2(a) shows

a cartoon of the different layers while Fig. 1.2(b) shows the density and temperature profiles of the atmosphere with each region labeled. The *photosphere* is what we typically refer to as the solar surface, with the actual surface located where the optical depth, τ , is equal to 1. This region also contains the lowest temperature on the Sun, approximately 4400 K, located about 525 km above the surface (Carroll & Ostlie, 2007). The photosphere is where the majority of the visibly (with the naked eye) observable photons originate.

The temperature minimum defines the top of the photosphere above which lies the *chromosphere*. As can be seen from Fig. 1.2(b), the density in the chromosphere is many orders of magnitude less than that of the photosphere and the temperature has increased from the minimum up to about 1×10^4 K. Though not visible with the naked eye, the chromosphere is highly structured. Structures such as spicules, tall columns of gas that extend high into the solar atmosphere thought to heavily impact the behavior of plasma in the corona (De Pontieu et al., 2011), originate in the photosphere as well as filaments, essentially spicules observed on-disk and plage, bright regions surrounding sunspots.

Next is the *transition region*, so called because of the steep temperature and density gradients (see Fig. 1.2(b)) that mark the transition between the chromosphere and the corona. The transition region is extremely thin, only a few hundred kilometers as compared to the chromosphere which extends over many thousands of kilometers. However, in this very short change in altitude, the temperature in the




Figure 1.2 : Layers of the solar atmosphere (a)Courtesy of NASA (b) Taken from Gary et al. (2007)

solar atmosphere jumps from $\approx 1 \times 10^4$ K to temperatures exceeding 1×10^5 K.

Finally, the solar *corona*, or “crown”, is the highly-dynamic uppermost layer of the Sun’s atmosphere. Visible with the naked eye only during a total solar eclipse (see Fig. 1.1), the corona is highly-structured and diffuse. Here, the temperature continues to increase, with typical coronal temperatures exceeding 1×10^6 K. Particularly high temperatures ($\approx 1 \times 10^7$ K) have also been observed in *active regions*, sites of intense magnetic activity associated with sunspots. These active regions can contain plasma as cool as 1×10^4 K as well and represent some of the most dynamic portions of the solar corona.

The work presented here will focus primarily on the plasma dynamics in the corona, in particular, in the cores of active regions where the intense magnetic field drives the motion of the plasma. In the following sections, we will discuss the origin

of the solar magnetic field, its topology, how it impacts plasma in the solar corona, and finally how it connects to the anomalously high temperatures seen in the upper solar atmosphere.

1.2 The Solar Magnetic Field

touch on field origin (dynamo theory), discuss how the field gets tangled, formation of loops, drives behavior of the atmosphere, lead into discussion of coronal heating, low beta versus high beta

Because of the high temperatures that characterize both the solar atmosphere and interior, much of the gas that makes up the Sun is ionized; that is, each atom has been stripped of at least one of its electrons. This means that the Sun is filled by a sea of charged particles, or plasma. In the corona where temperatures are often higher than 1×10^6 K, even heavy elements such as Mg, Ca, and Fe are stripped of their electrons. Because these particles are charged, their motion is strongly dictated by both the electric and magnetic fields via the Lorentz force law. Thus, to understand the dynamics of the coronal plasma, one must first try to understand the magnetic field that ultimately controls it.

The Sun, like the Earth, possesses an intrinsic magnetic field, though its origin and dynamics constitute a largely unsolved problem. Additionally, unlike Earth, the solar magnetic field is highly dynamic and comparatively quite strong (up to 1000 G in sunspots as compared to the < 1 G field on Earth) (Aschwanden, 2006). From the




Figure 1.3 : Taken from Gary (2001).

earliest studies of magnetism, it has been known that a conductor moving through a magnetic field produces a current. This current then in turn produces an additional magnetic field. Treating the convection zone (below the photosphere) plasma as the conductor and assuming some preexisting, but small magnetic field leads to what is commonly known as *dynamo theory* (Golub & Pasachoff, 2010).

Dynamo theory seeks to show self-consistently how the interaction between the Sun's hot plasma and some small initial field leads to the observed complicated topologies and field strengths. Modeling such a system is no easy task as the interactions between the field and the plasma are highly non-linear, meaning solving the so-called “dynamo equations” requires a significant amount of computational effort. something here about how the field pops and creates loop things, Reale loop paper

Fortunately, we can still learn something about the corona without completely

solving the problem of where the driving magnetic field comes from. The corona constitutes what is commonly referred to as a “low-beta plasma” such that $\beta = p_{\text{thermal}}/(B^2 8\pi)$, the ratio between the gas pressure and magnetic pressure, is $\beta < 1$ (see Fig. 1.3). This means that coronal magnetic field acts as a guide for the plasma, with the flow directed primarily in the field direction with negligible cross-field motion. A useful analogy is that of water flowing through a pipe: the rigid pipe directs the flow of the water, but has no affect the underlying fluid dynamics. Provided the field is reasonably stable and the relevant fluid equations are parameterized in terms of the field-aligned coordinate, our equations governing the plasma dynamics need not include the field at all!

Talk about drawbacks, mention MHD and hydrodynamic and hydrostatic; lead into coronal heating conversation

1.3 The Coronal Heating Problem

Here discuss coronal heating broadly, first evidence for high coronal temperatures, some proposed heating mechanisms, magnetic reconnection, AC versus DC heating, wave heating versus braiding etc. Use Klimchuk review of coronal heating for this section + Cargill nanoflare paper(s)

1.4 Summary

End with outline of the rest of the thesis: in ch. such and such we will discuss such and such

Chapter 2

Coronal Loops

This chapter will discuss the discrete nature of corona in terms of coronal loop structures. Need a section on general plasma dynamics of loops to discuss energy transfer/loss/gain through heating/enthalpy/radiation/draining/filling. Also discuss general structure and how they are formed. Give some general characteristics about them like length, temperature, density, through what layers they extend etc. Show nice schematic. The solar corona is typically described in terms of discrete structures known as coronal loops. Heated

2.1 Observations

Discuss some observations of loops and what has been learned about them, what constraints, multi-stranded versus single stranded. Show some pretty pictures

2.2 Modeling

Discuss modeling approaches, hydrodynamics versus magnetohydrodynamics, etc.

Chapter 3

Emission Diagnostics

This chapter will discuss emission measure (EM), differential EM (DEM), line intensities etc. and how they are interpreted in observational and modeling contexts. Discuss how this is how we know anything about plasma in the solar atmosphere. Can discuss forward modeling as well; this sets us up for the rest of the thesis.

Chapter 4

Numerical Modeling

This chapter will outline our numerical approach to studying the solar corona

Often, the full 3D magnetohydrodynamic (MHD) equations are used when modeling loops in the solar corona. However, such treatments are computationally expensive and are prohibitive when studies of large parameter spaces are needed. Instead, the one-dimensional (1D) hydrodynamic equations are used to model plasma dynamics in coronal loops. These hydrodynamic equations do not include the magnetic field B . This is because the corona These equations are parameterized in terms of time, t , and the field-aligned coordinate, s .

4.1 One-dimensional Hydrodynamics

List the 1D hydrodynamic equations and briefly discuss their origin (i.e. how they were derived) and how they differ from MHD approach Include plasma β justification of why we don't include the magnetic field

The single-fluid 1D hydrodynamic equations are given by,

$$\frac{\partial \rho}{\partial t} = -\frac{\partial(\rho v)}{\partial s} \quad (4.1)$$

$$\frac{\partial(\rho v)}{\partial t} = -\frac{\partial(\rho v^2)}{\partial s} - \frac{\partial(p)}{\partial s} + \frac{\partial}{\partial s} \left(\frac{4}{3} \mu \frac{\partial v}{\partial s} \right) + \rho g_{\parallel} \quad (4.2)$$

$$\frac{\partial E}{\partial t} = -\frac{\partial}{\partial s} ((E + p)v) - \frac{\partial F}{\partial s} + E_H - E_R + \rho g_{\parallel} v, \quad (4.3)$$

where

$$E = \frac{1}{\gamma - 1} p + \frac{1}{2} \rho v^2. \quad (4.4)$$

s and t are the field-aligned spatial coordinate and time, respectively, $\rho = mn$ is the mass density, v is the bulk velocity, μ is the viscosity coefficient and g_{\parallel} is the component of gravitational acceleration parallel to s , E is the energy density and $p = 2nk_B T$ is the pressure subject to closure by the ideal gas law.

Additionally, we also have three terms that are specific to energy balance in a coronal loop: the heat flux $F = \kappa_0 T^{5/2} \partial T / \partial s$, typically given by the classical Spitzer-Harm expression, the ad-hoc heating term E_H , and the volumetric radiative loss term $E_R = n^2 \Lambda(T)$, where $\Lambda(T)$ is the radiative loss function. Emphasize terms that are “non-standard” hydrodynamic terms (e.g. radiation, heat flux, ad-hoc heating, enthalpy transfer) Bring up HYDRAD model and discuss speed limitations

4.2 The EBTEL Model

When using hydrodynamic simulations to study coronal loops, it is often best to compare computed observables with observations in order to make constraints on

loop properties. To do this, one must explore a very large parameter space of initial conditions (e.g. loop length, heating frequency, power-law index, etc.). Though the 1D hydrodynamic equations have been shown to accurately model coronal loops,

Derive the EBTEL equations from the 1D hydrodynamic equations and discuss the physics behind the EBTEL model; what physics is left out? what physical insight can we still gain from this model? Say why EBTEL is important (i.e. large parameter sweeps that can't be done with HYDRAD) Discuss 0D nature, maybe include discussion of previous 0D models and why EBTEL is better Show comparisons between EBTEL and HYDRAD

4.3 The Two-fluid EBTEL Model

Discuss importance of two-fluid effects in hydrodynamic models of the solar corona; include some quick calculations to show how electron and ion fluids can become de-coupled Derive two-fluid EBTEL equations from 1D hydrodynamic equations Show several comparisons between HYDRAD and two-fluid EBTEL to justify its use in this study remember to show EM calculation, justification, how EBTEL treats DEM as opposed to HYDRAD Briefly detail solver(s) used, adaptive timestep, why this is necessary, etc. If there is time, discuss speed comparison between EBTEL and HYDRAD

Chapter 5

Results

This chapter will include the results of our numerical study. Here we will also describe how the study was performed and what tools were used to perform the study. It is best not to introduce any new tools here; just pool from those that have already been discussed and show how they were applied. Show lots of plots and tables.

Chapter 6

Conclusions

This chapter will discuss the conclusions that we can draw based on the results in the results section. What do these results mean? What are the implications in the context of loops in active region cores? May include some topics for future work in this section as well or just wait and put it in a different chapter

Bibliography

- Aschwanden, M. J. 2006, Physics of the Solar Corona: An Introduction with Problems and Solutions, 2nd edn., Springer-Praxis books in geophysical sciences (Berlin ; New York: Springer)
- Carroll, B. W., & Ostlie, D. A. 2007, An Introduction to Modern Astrophysics, second edition edn. (2007: Pearson)
- De Pontieu, B., McIntosh, S. W., Carlsson, M., et al. 2011, Science, 331, 55
- Gary, G. A. 2001, Solar Physics, 203, 71
- Gary, G. A., West, E. A., Rees, D., et al. 2007, Astronomy and Astrophysics, 461, 707
- Golub, L., & Pasachoff, J. M. 2010, The Solar Corona (Cambridge, UK ; New York : Cambridge University Press, 2010.)
- Olson, R. J. M., & Pasachoff, J. M. 2007, Religion & the Arts, 11, 299