

$$\vec{a}\times\vec{b}=|a|\cdot|b|\cdot\sin\langle\vec{a},\vec{b}\rangle$$

$$\mathrm{e}^{\mathrm{i}\pi}+1=0$$

$$\mathrm{e}^{\mathrm{i}\theta}=\cos\theta+\mathrm{i}\sin\theta$$

$$a^2+b^2=c^2$$

$$\alpha\beta\gamma\delta\epsilon\zeta\eta\theta\iota\kappa\lambda\mu\nu\xi\omicron\pi\rho\sigma\tau\nu\phi\psi\chi\omega$$

$$\varphi\varepsilon\vartheta$$

$$\frac{\mathrm{d} x}{\mathrm{d} y}$$

$$f'(x), \quad f^n(x)$$

$$\frac{\partial f}{\partial x}$$

$$\int f(x)\,\mathrm{d} x$$

$$\int_a^b f(x)\,\mathrm{d} x = F(b)-F(a)$$

$$\iint_D f(x,y)\,\mathrm{d} x\,\mathrm{d} y$$

$$\iint_D f(x,y)\,\mathrm{d} \sigma$$

$$\oint_L P\,\mathrm{d} x+Q\,\mathrm{d} y=\iint_D\left(\frac{\partial P}{\partial x}-\frac{\partial Q}{\partial y}\right)\,\mathrm{d} \sigma$$

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}$$

$$\nabla \cdot \vec{E} = \frac{\rho}{\varepsilon_0}$$

$$\nabla \cdot \vec{B} = 0$$

$$\nabla \times \vec{E} = -\frac{\partial \vec{B}}{\partial t}$$

$$\nabla \times \vec{B} = \mu_0 \vec{J} + \mu_0 \varepsilon_0 \frac{\partial \vec{E}}{\partial t}$$

$$\nabla f, \quad \nabla \cdot \vec{F}, \quad \nabla \times \vec{F}$$

$$\vec{F}(x,y,z) \quad \nabla^2 f$$

