

Algebra

Serge Lang

August 4, 2021

Contents

1	Groups	2
1.1	Monoids	2
1.2	Groups	2
1.3	Normal Subgroups	2
1.4	Direct Sums and Free Abelian Groups	2
2	Rings	5
2.1	Rings and Homomorphisms	5
2.2	Commutative Rings	11
2.3	Polynomials and Group Rings	15
2.4	Localization	19
3	Modules	20
3.1	Basic Definitions	20
3.2	The Group of Homomorphisms	22
3.3	Direct Products and Sums of Modules	24
4	Algebraic Extensions	27
4.1	Finite and Algebraic Extensions	27
5	Real Fields	27
5.1	Ordered Fields	27

1 Groups

1.1 Monoids

1.2 Groups

1.3 Normal Subgroups

Let $f : G \rightarrow G'$ be a group homomorphism, and let H be its kernel. If x is an element of G , then $xH = Hx$, because both are equal to $f^{-1}(f(x))$. We can also rewrite this relation as $xHx^{-1} = H$

Conversely, let G be a group and let H be a subgroup. Assume that for all elements $x \in G$, we have $xH \subset Hx$ (or equivalently, $xHx^{-1} \subset H$), which implies $H \subset xHx^{-1}$. Thus our condition is equivalent to the condition $xHx^{-1} = H$ for all $x \in G$. A subgroup H satisfying this condition will be called **normal**

Let G' be the set of cosets of H . (A left coset is equal to a right coset). If xH and yH are cosets, then their product

$$xHyH = xyHH = xyH$$

is also a coset. Hence G' is a group.

Let $f : G \rightarrow G'$ be the mapping s.t. $f(x)$ is the coset xH . Then f is clearly a homomorphism and H is equal to the kernel.

The group of cosets of a normal subgroup H is denoted by G/H (which we read G modulo H , or $G \bmod H$). The map f of G onto G/H constructed above is called the **canonical map**, and G/H is called the **factor group** of G by H

1.4 Direct Sums and Free Abelian Groups

Let $\{A_i\}_{i \in I}$ be a family of abelian groups. We define their **direct sum**

$$A = \bigoplus_{i \in I} A_i$$

to be the subset of the direct product $\prod A_i$ consisting of all families $(x_i)_{i \in I}$ with $x_i \in A_i$ s.t. $x_i = 0$ for all but a finite number of indices i . For each index $j \in I$, we map

$$\lambda_j : A_j \rightarrow A$$

by letting $\lambda_j(x)$ be the element whose j -th component is x , and having all other components equal to 0. Then λ_j is an injective homomorphism

Proposition 1.1. Let $\{f_i : A_i \rightarrow B\}$ be a family of homomorphisms into an abelian group B . Let $A = \bigoplus A_i$. There exists a unique homomorphism

$$f : A \rightarrow B$$

s.t. $f \circ \lambda_j = f_j$ for all j

Proof. Define

$$f((x_i)_{i \in I}) = \sum_{i \in I} f_i(x_i)$$

□

The property in Proposition 1.1 is called the **universal property** of the direct sum.

Let A be an abelian group and B, C subgroups. If $B + C = A$ and $B \cap C = \{0\}$ then the map

$$B \times C \rightarrow A$$

given by $(x, y) \mapsto x + y$ is an isomorphism. Instead of writing $A = B \times C$ we shall write $A = B \oplus C$ and say that A is the **direct sum** of B and C . We use a similar notation for the direct sum of a finite number of subgroups B_1, \dots, B_n s.t.

$$B_1 + \dots + B_n = A$$

and

$$B_{i+1} \cap (B_1 + \dots + B_i) = 0$$

In that case, we write

$$A = B_1 \oplus B_2 \oplus \dots \oplus B_n$$

Let A be an abelian group. Let $\{e_i\}_{i \in I}$ be a family of elements of A . We say that this family is a **basis** of A if the family is not empty, and if every element of A has a unique expression as a linear combination

$$x = \sum x_i e_i$$

with $x_i \in \mathbb{Z}$ and almost all $x_i = 0$. Thus the sum is actually a finite sum. An abelian group is **free** if it has a basis. If that is the case, then if we let $Z_i = \mathbb{Z}$ for all i , then A is isomorphic to the direct sum

$$A \cong \bigoplus_{i \in I} Z_i$$

Now let S be a set. Let $\mathbb{Z}\langle S \rangle$ be the set of all maps $\varphi : S \rightarrow \mathbb{Z}$ s.t. $\varphi(x) = 0$ for almost all $x \in S$. Then $\mathbb{Z}\langle S \rangle$ is an abelian group. if k is an integer and

$x \in S$, we denote by $k \cdot x$ the map φ s.t. $\varphi(x) = k$ and $\varphi(y) = 0$ if $y \neq x$. Then every element φ of $\mathbb{Z}\langle S \rangle$ can be written in the form

$$\varphi = k_1 \cdot x_1 + \cdots + k_n \cdot x_n$$

for $k_i \in \mathbb{Z}$ and $x_i \in S$, all the x_i being distinct. Furthermore, φ **admits a unique such expression**, because if we have

$$\varphi = \sum_{x \in S} k_x \cdot x = \sum_{x \in S} k'_x \cdot x$$

then

$$0 = \sum_{x \in S} (k_x - k'_x) \cdot x$$

whence $k'_x = k_x$ for all $x \in S$

We map S into $\mathbb{Z}\langle S \rangle$ by the map $f_S = f$ s.t. $f(x) = 1 \cdot x$. $f(S)$ generates $\mathbb{Z}\langle S \rangle$. If $g : S \rightarrow B$ is a mapping of S into some abelian group B , then we define a map

$$g_* : \mathbb{Z}\langle S \rangle \rightarrow B$$

s.t.

$$g_* \left(\sum_{x \in S} k_x \cdot x \right) = \sum_{x \in S} k_x g(x)$$

It's unique for any such homomorphism g_* must be s.t. $g_*(1 \cdot x) = g(x)$

Proposition 1.2. *if $\lambda : S \rightarrow S'$ is a mapping of sets, there is a unique homomorphism $\bar{\lambda}$ making the following diagram commutative*

$$\begin{array}{ccc} S & \xrightarrow{f_S} & \mathbb{Z}\langle S \rangle \\ \downarrow \lambda & & \downarrow \bar{\lambda} \\ S' & \xrightarrow{f_{S'}} & \mathbb{Z}\langle S' \rangle \end{array}$$

In fact, $\bar{\lambda}$ is none other than $(f_S \circ \lambda)_$*

We shall denote $\mathbb{Z}\langle S \rangle$ also $F_{ab}(S)$ and call $F_{ab}(S)$ the **free abelian group generated by S** . We call elements of S its **free generators**

2 Rings

2.1 Rings and Homomorphisms

A **ring** A is a set

1. w.r.t. addition, A is a commutative group
2. the multiplication is associative, and has a unit element
3. for all $x, y, z \in A$ we have

$$(x + y)z = xz + yz \quad \text{and} \quad z(x + y) = zx + zy$$

(called **distributivity**)

We denote the unit element for addition by 0, and the unit element for multiplication by 1. Observe that $0x = 0$ for all $x \in A$. *Proof:* $0x + x = (0 + 1)x = x$

For any $x, y \in A$ we have $(-x)y = -(xy)$

Let A be a ring, and let U be the set of elements of A which have both a right and left inverse. Then U is a multiplicative group. Indeed, if a has a right inverse b , so that $ab = 1$, and a left inverse c , so that $ca = 1$, then $cab = b$, whence $c = b$, and we see that c is a two-sided inverse, and that c itself has a two-sided inverse, namely a . Therefore U satisfies all the axioms of a multiplicative group, and is called the group of **units** of A . It is sometimes denoted by A^* , and is also called the group of **invertible** elements of A . A ring A s.t. $1 \neq 0$ and s.t. every non-zero element is invertible is called a **division ring**.

Example 2.1 (The Shift Operator). Let E be the set of all sequences

$$a = (a_1, a_2, a_3, \dots)$$

of integers. One can define addition componentwise. Let R be the set of all mappings $f : E \rightarrow E$ of E into itself s.t. $f(a + b) = f(a) + f(b)$. Then R is a ring. Let

$$T(a_1, a_2, a_3, \dots) = (0, a_1, a_2, a_3, \dots)$$

Verify that T is left invertible but not right invertible

A ring A is said to be **commutative** if $xy = yx$ for all $x, y \in A$. A commutative division ring is called a **field**. By definition, a field contains at least two elements, namely 0 and 1.

A subset B of ring A is called a **subring** if it is an additive subgroup, if it contains the multiplicative unit, and if $x, y \in B$ implies $xy \in B$. If that is the case, then B is in itself a ring, the laws of operation in B being the same as the laws of operation in A .

For example, the **center** of a ring A is the subset of A consisting of all elements $a \in A$ s.t. $ax = xa$ for all $x \in A$. The center of A is a subring.

If x, y_1, \dots, y_n are elements of a ring, then by induction one sees that

$$x(y_1 + \dots + y_n) = xy_1 + \dots + xy_n$$

If $x_i (i = 1, \dots, n)$ and $y_j (j = 1, \dots, m)$ are elements of A , then it is also easily proved that

$$\left(\sum_{i=1}^n x_i \right) \left(\sum_{j=1}^m y_j \right) = \sum_{i=1}^n \sum_{j=1}^m x_i y_j$$

Furthermore, distributivity holds for subtraction, e.g.

$$x(y_1 - y_2) = xy_1 - xy_2$$

Example 2.2. Let S be a set and A a ring. Let $\text{Map}(S, A)$ be the set of mappings of S into A . Then $\text{Map}(S, A)$ is a ring if for $f, g \in \text{Map}(S, A)$ we define

$$(fg)(x) = f(x)g(x) \quad \text{and} \quad (f + g)(x) = f(x) + g(x)$$

for all $x \in S$.

Let M be an additive abelian group, and let A be the set $\text{End}(M)$ of group-homomorphisms of M into itself. We define addition in A to be the addition of mappings, and we define multiplication to be **composition** of mappings

Example 2.3 (The convolution product). Let G be a group and let K be a field. Denote by $K[G]$ the set of all formal linear combinations $\alpha = \sum a_x x$ with $x \in G$ and $a_x \in K$, s.t. all but finite number of a_x are equal to 0. If $\beta = \sum b_x x \in K[G]$, then one can define the product

$$\alpha\beta = \sum_{x \in G} \sum_{y \in G} a_x b_y xy = \sum_{z \in G} \left(\sum_{xy=z} a_x b_y \right) z$$

With this product, the **group ring** $K[G]$ is a ring. $K[G]$ is commutative iff G is commutative. The second sum on the right defines what is called a **convolution product**. If f, g are functions on a group G , we define their **convolution** $f * g$ by

$$(f * g)(z) = \sum_{xy=z} f(x)g(y)$$

A **left ideal** \mathfrak{a} in a ring A is a subset of A which is a subgroup of the additive group of A , s.t. $A\mathfrak{a} \subset \mathfrak{a}$ (and hence $A\mathfrak{a} = \mathfrak{a}$ since A contains 1). To define a right ideal, we require $\mathfrak{a}A = \mathfrak{a}$, and a **two-sided ideal** is a subset which is both a left and right ideal. A two-sided ideal is called an **ideal** in this section.

If A is a ring and $a \in A$, then Aa is a left ideal, called **principal**. We say that a is a generator of \mathfrak{a} (over A). AaA is a principal two-sided ideal if $AaA = \{\sum x_i a y_i \mid x_i, y_i \in A\}$. More generally, let $a_1, \dots, a_n \in A$. We denote by (a_1, \dots, a_n) the set of elements of A which can be written in the form

$$x_1 a_1 + \dots + x_n a_n \quad \text{with } x_i \in A$$

Then this set of elements is immediately verified to be a left ideal, and a_1, \dots, a_n are called **generators** of the left ideal.

If $\{\mathfrak{a}_i\}_{i \in I}$ is a family of ideals, then their intersection

$$\bigcap_{i \in I} \mathfrak{a}_i$$

is also an ideal

A **commutative** ring s.t. every ideal is principal and s.t. $1 \neq 0$ is called a **principal ring**

Example 2.4. The integers \mathbb{Z} form a ring, which is commutative. Let \mathfrak{a} be an ideal $\neq \mathbb{Z}$ and $\neq 0$. If $n \in \mathfrak{a}$ then $-n \in \mathfrak{a}$. Let d be the smallest integer > 0 lying in \mathfrak{a} . If $n \in \mathfrak{a}$ then there exists integers q, r with $0 \leq r < d$ s.t.

$$n = dq + r$$

Since \mathfrak{a} is an ideal, it follows that r lies in \mathfrak{a} , hence $r = 0$. Hence \mathfrak{a} consists of all multiples qd of d , which $q \in \mathbb{Z}$, and \mathbb{Z} is a principal ring.

Let $\mathfrak{a}, \mathfrak{b}$ be ideals of A . We define $\mathfrak{a}\mathfrak{b}$ to be the set of all sums

$$x_1 y_1 + \dots + x_n y_n$$

with $x_i \in \mathfrak{a}$ and $y_i \in \mathfrak{b}$. $\mathfrak{a}\mathfrak{b}$ is an ideal, and that the set of ideals forms a multiplicative monoid, the unit element being the ring itself. This unit element is called the **unit ideal** and is often written (1).

If $\mathfrak{a}, \mathfrak{b}$ are left ideals of A , then $\mathfrak{a} + \mathfrak{b}$ (the sum being taken as additive subgroup of A) is obviously a left ideal. Thus ideals also form a monoid under addition. We also have distributivity: if $\mathfrak{a}_1, \dots, \mathfrak{a}_n, \mathfrak{b}$ are ideals of A , then

$$\mathfrak{b}(\mathfrak{a}_1 + \dots + \mathfrak{a}_n) = \mathfrak{b}\mathfrak{a}_1 + \dots + \mathfrak{b}\mathfrak{a}_n$$

Let \mathfrak{a} be a left ideal. Define $\mathfrak{a}A$ to be the set of all sums $a_1x_1 + \cdots + a_nx_n$ with $a_i \in \mathfrak{a}$ and $x_i \in A$. Then $\mathfrak{a}A$ is an ideal.

Suppose that A is commutative. Let $\mathfrak{a}, \mathfrak{b}$ be ideals. Then trivially

$$\mathfrak{a}\mathfrak{b} \subset \mathfrak{a} \cap \mathfrak{b}$$

If $\mathfrak{a} + \mathfrak{b} = A$ then $\mathfrak{a}\mathfrak{b} = \mathfrak{a} \cap \mathfrak{b}$. Suppose $x \in \mathfrak{a} \cap \mathfrak{b}$ and $x = a_x + b_x$, where $a_x \in \mathfrak{a}$ and $b_x \in \mathfrak{b}$. Then $a_x \in \mathfrak{b}$ and $b_x \in \mathfrak{a}$. If $1 = a_1 + b_1$ then $x \cdot 1 = (a_x + b_x)(a_1 + b_1) \in \mathfrak{a}\mathfrak{b}$

By a **ring homomorphism** one means a mapping $f : A \rightarrow B$ where A, B are rings, and s.t. f is a monoid-homomorphism for the multiplicative structures on A and B , and also a monoid homomorphism for the additive structure. In other words

$$\begin{aligned} f(a + a') &= f(a) + f(a') & f(aa') &= f(a)f(a') \\ f(1) &= 1 & f(0) &= 0 \end{aligned}$$

for all $a, a' \in A$.

The kernel of a ring homomorphism $f : A \rightarrow B$ is an ideal of A .

Conversely, let \mathfrak{a} be an ideal of the ring A . We can construct the **factor ring** A/\mathfrak{a} as follows. Viewing A and \mathfrak{a} as additive groups, let A/\mathfrak{a} be the factor group. If $x + \mathfrak{a}$ and $y + \mathfrak{a}$ are two cosets of \mathfrak{a} , we define $(x + \mathfrak{a})(y + \mathfrak{a})$ to be the coset $xy + \mathfrak{a}$. This coset is well-defined, for if x_1, y_1 are in the same coset as x, y respectively, then one verifies that x_1y_1 is in the same coset as xy . Unit element is $1 + \mathfrak{a}$.

We therefore defined a ring structure on A/\mathfrak{a} and the canonical map

$$f : A \rightarrow A/\mathfrak{a}$$

is then clearly a ring homomorphism

Proposition 2.1. *If $g : A \rightarrow A'$ is a ring homomorphism whose kernel contains \mathfrak{a} , then there exists a unique ring homomorphism $g_* : A/\mathfrak{a} \rightarrow A'$ making the following diagram commutative*

$$\begin{array}{ccc} A & \xrightarrow{g} & A' \\ & \searrow f & \nearrow g_* \\ & A/\mathfrak{a} & \end{array}$$

Indeed, viewing f, g as group homomorphisms, there is a unique group homomorphism g_* making our diagram commutative

Proof. If $x \in A$ then $g(x) = g_*f(x)$. Hence for $x, y \in A$

$$\begin{aligned} g_*(f(x)f(y)) &= g_*(f(xy)) = g(xy) = g(x)g(y) \\ &= g_*f(x)g_*f(y) \end{aligned}$$

Given $\xi, \eta \in A/\mathfrak{a}$, there exists $x, y \in A$ s.t. $f(x) = \xi$ and $f(y) = \eta$. Since $f(1) = 1$, we get $g_*f(1) = g(1) = 1$ and hence the two conditions that g_* be a multiplicative monoid-homomorphism are satisfied \square

Let A be a ring, and denote its unit element by e for the moment. The map

$$\lambda : \mathbb{Z} \rightarrow A$$

s.t. $\lambda(n) = ne$ is a ring homomorphism, and its kernel is an ideal (n) , generated by an integer $n \geq 0$. We have a canonical injective homomorphism $\mathbb{Z}/n\mathbb{Z} \rightarrow A$ which is a (ring) isomorphism between $\mathbb{Z}/n\mathbb{Z}$ and a subring of A . If $n\mathbb{Z}$ is a prime ideal, then $n = 0$ or $n = p$ for some prime number p . In the first place, A contains as a subring a ring which is isomorphic to \mathbb{Z} , and which is often identified with \mathbb{Z} . In that case, we say that A has **characteristic 0**. if on the other hand $n = p$ then we say that A has **characteristic p** , and A contains (an isomorphic image of) $\mathbb{Z}/p\mathbb{Z}$ as a subring. We abbreviate $\mathbb{Z}/p\mathbb{Z}$ by \mathbb{F}_p .

If K is a field, then K has characteristic 0 or $p > 0$. (if its characteristic is $a \cdot b$, then $a \cdot b \cdot 1 = 0$ but field is an integral domain). In the first case, K contains as a subfield an isomorphic image of the rational numbers, and in the second case, it contains an isomorphic image of \mathbb{F}_p . In either case, this subfield will be called the **prime field** (contained in K). Since this prime field is the smallest subfield of K containing 1 and has no automorphism except the identity, it is customary to identify it with \mathbb{Q} or \mathbb{F}_p as the case may be. By the **prime ring** (in K) we shall mean either the integers \mathbb{Z} if K has characteristic 0 or \mathbb{F}_p if K has characteristic p .

Let A be a subring of a ring B . Let S be a subset of B commuting with A . We denote by $A[S]$ the set of all elements

$$\sum a_{i_1 \dots i_n} s_1^{i_1} \dots s_n^{i_n}$$

the sum ranging over a finite number of n -tuples (i_1, \dots, i_n) of integers ≥ 0 , and $a_{i_1, \dots, i_n} \in A$, $s_1, \dots, s_n \in S$. If $B = A[S]$, we say that S is a set of **generators** (or **ring generators**) for B over A , or that B is **generated** by S over A . If S is finite, B is **finitely generated as a ring over A** . Note that S is not commutative.

Let A be a ring, \mathfrak{a} an ideal, and S a subset of A . We write

$$S \equiv 0 \pmod{\mathfrak{a}}$$

if $S \subset \mathfrak{a}$. If $x, y \in A$ we write

$$x \equiv y \pmod{\mathfrak{a}}$$

if $x - y \in \mathfrak{a}$. If \mathfrak{a} is principal, equal to (a) , then we also write

$$x \equiv y \pmod{a}$$

If $f : A \rightarrow A/\mathfrak{a}$ is the canonical homomorphism, then $x \equiv y \pmod{\mathfrak{a}}$ means that $f(x) = f(y)$

The factor ring A/\mathfrak{a} is also called a **residue class ring**. Cosets of \mathfrak{a} in A are called **residue classes** modulo \mathfrak{a} , and if $x \in A$, then the coset $x + \mathfrak{a}$ is called the **residue class of x modulo \mathfrak{a}**

An injective ring homomorphism $f : A \rightarrow B$ establishes a ring isomorphism between A and its image. Such a homomorphism will be called an **embedding**

Let $f : A \rightarrow A'$ be a ring homomorphism, and let \mathfrak{a}' be an ideal of A' . Then $f^{-1}(\mathfrak{a}')$ is an ideal \mathfrak{a} in A , and we have an induced injective homomorphism

$$A/\mathfrak{a} \rightarrow A'/\mathfrak{a}'$$

Proposition 2.2. *Products exist in the category of rings*

Let A be a ring. Elements $x, y \in A$ are said to be **zero divisors** if $x \neq 0$, $y \neq 0$ and $xy = 0$. A ring A is **entire** if $1 \neq 0$, if A is commutative and if there are no zero divisors in the ring. (Entire rings are also called **integral domains**)

Let m be a positive integer $\neq 1$. The ring $\mathbb{Z}/m\mathbb{Z}$ has zero divisors iff m is not prime.

Proposition 2.3. *Let A be an entire ring, and let a, b be non-zero elements of A . Then a, b generate the same ideal iff there exists a unit u of A s.t. $b = au$.*

Proof. Assume $Aa = Ab$. Then $a = bc$ and $b = ad$ for some $c, d \in A$. Hence $a = adc$ whence $a(1 - dc) = 0$ and therefore $dc = 1$. Hence c is a unit \square

2.2 Commutative Rings

Assume A is commutative

A **prime** ideal in A is an ideal $\mathfrak{p} \neq A$ s.t. A/\mathfrak{p} is entire. Equivalently, we could say that it is an ideal $\mathfrak{p} \neq A$ s.t. whenever $x, y \in A$ and $xy \in \mathfrak{p}$ then $x \in \mathfrak{p}$ or $y \in \mathfrak{p}$. A prime ideal is often called simply a **prime**

Proposition 2.4. *Every maximal ideal is prime*

Proof. Let \mathfrak{m} be maximal and let $x, y \in A$ s.t. $xy \in \mathfrak{m}$. Suppose $x \notin \mathfrak{m}$, then $\mathfrak{m} + Ax$ is an ideal properly containing \mathfrak{m} , hence equal to A . Hence we can write

$$1 = u + ax$$

with $u \in \mathfrak{m}$ and $a \in A$. Multiplying by y we find

$$y = yu + axy$$

whence $y \in \mathfrak{m}$. □

Proposition 2.5. *Let \mathfrak{a} be an ideal $\neq A$. Then \mathfrak{a} is contained in some maximal ideal \mathfrak{m}*

Proposition 2.6. *The ideal $\{0\}$ is a prime ideal of A iff A is entire*

The only ideals of a field are itself and the zero ideal

Proposition 2.7. *If \mathfrak{m} is a maximal ideal of A , then A/\mathfrak{m} is a field*

Proof. If $x \in A$, we denote by \bar{x} its residue class mod \mathfrak{m} . Since $\mathfrak{m} \neq A$ we note that A/\mathfrak{m} has a unit element $\neq 0$. Any non-zero element of A/\mathfrak{m} can be written as \bar{x} for some $x \in A, x \notin \mathfrak{m}$. To find its inverse, note that $\mathfrak{m} + Ax$ is an ideal of $A \neq \mathfrak{m}$ and hence equal to A . Hence we can write

$$1 = u + yx$$

with $u \in \mathfrak{m}$ and $y \in A$. This means that $\bar{y}\bar{x} = 1 = \bar{1}$ and hence that \bar{x} has an inverse. □

Proposition 2.8. *Let $f : A \rightarrow A'$ be a homomorphism of commutative rings. Let \mathfrak{p}' be a prime ideal of A' and let $\mathfrak{p} = f^{-1}\mathfrak{p}'$. Then \mathfrak{p} is prime*

Example 2.5. Let \mathbb{Z} be the ring of integers. Since an ideal is also an additive subgroup of \mathbb{Z} , every ideal $\neq \{0\}$ is principal, of the form $n\mathbb{Z}$ for some integer $n > 0$. (proof)

Let \mathfrak{p} be a prime ideal $\neq \{0\}$, $\mathfrak{p} = n\mathbb{Z}$. Then n must be a prime number. Conversely, if p is a prime number, then $p\mathbb{Z}$ is a prime ideal. Furthermore, $p\mathbb{Z}$ is a maximal ideal. Suppose $p\mathbb{Z}$ is contained in some ideal $n\mathbb{Z}$, then $p = nm$ for some integer m , whence $n = p$ or $n = 1$, thereby proving $p\mathbb{Z}$ maximal

if n is an integer, the factor ring $\mathbb{Z}/n\mathbb{Z}$ is called the ring of **integers modulo n** . We also denote

$$\mathbb{Z}/n\mathbb{Z} = \mathbb{Z}(n)$$

If n is a prime number p , then the ring of integers modulo p is in fact a field, denoted by \mathbb{F}_p . In particular, the multiplicative group of \mathbb{F}_p is called the group of non-zero integers modulo p . From the elementary properties of groups, we get a standard fact of elementary number theory: if x is an integer $\neq 0 \pmod p$, then $x^{p-1} \equiv 1 \pmod p$ (Fermat's Theorem). Similarly given an integer $n > 1$, the units in the ring $\mathbb{Z}/n\mathbb{Z}$ consist of those residue class mod $n\mathbb{Z}$ which are represented by integers $m \neq 0$ and prime to n . The order of the group of units in $\mathbb{Z}/n\mathbb{Z}$ is called by definition $\varphi(n)$ (where φ is known as the **Euler phi-function**). Consequently, if x is an integer prime to n , then $x^{\varphi(n)} \equiv 1 \pmod n$

Theorem 2.9 (Chinese Remainder Theorem). *Let $\mathfrak{a}_1, \dots, \mathfrak{a}_n$ be ideals of A s.t. $\mathfrak{a}_i + \mathfrak{a}_j = A$ for all $i \neq j$. Given elements $x_1, \dots, x_n \in A$, there exists $x \in A$ s.t. $x \equiv x_i \pmod{\mathfrak{a}_i}$ for all i*

Proof. For $n = 2$ we have an expression

$$1 = a_1 + a_2$$

for some $a_i \in \mathfrak{a}_i$, and we let $x = x_2 a_1 + x_1 a_2$

For each $i \geq 2$ we can find elements $a_i \in \mathfrak{a}_1$ and $b_i \in \mathfrak{a}_i$ s.t.

$$a_i + b_i = 1, \quad i \geq 2$$

The products $\prod_{i=2}^n (a_i + b_i)$ is equal to 1, and lies in

$$\mathfrak{a}_1 + \prod_{i=2}^n \mathfrak{a}_i$$

Hence

$$\mathfrak{a}_1 + \prod_{i=2}^n \mathfrak{a}_i = A$$

By theorem for $n = 2$, we can find an element $y_1 \in A$ s.t.

$$\begin{aligned} y_1 &\equiv 1 \pmod{\mathfrak{a}_1} \\ y_1 &\equiv 0 \pmod{\prod_{i=2}^n \mathfrak{a}_i} \end{aligned}$$

We find similarly elements y_2, \dots, y_n s.t.

$$y_j \equiv 1 \pmod{\mathfrak{a}_j} \quad \text{and} \quad y_j \equiv 0 \pmod{\mathfrak{a}_i \text{ for } i \neq j}$$

Then $x = x_1 y_1 + \dots + x_n y_n$ satisfies our requirements □

In the same vein as above, we observe that if $\mathfrak{a}_1, \dots, \mathfrak{a}_n$ are ideals of a ring A s.t.

$$\mathfrak{a}_1 + \dots + \mathfrak{a}_n = A$$

and if v_1, \dots, v_n are positive integers, then

$$\mathfrak{a}_1^{v_1} + \dots + \mathfrak{a}_n^{v_n} = A$$

Corollary 2.10. *Let $\mathfrak{a}_1, \dots, \mathfrak{a}_n$ be ideals of A . Assume that $\mathfrak{a}_i + \mathfrak{a}_j = A$ for $i \neq j$. Let*

$$f : A \rightarrow \prod_{i=1}^n A/\mathfrak{a}_i = (A/\mathfrak{a}_1) \times \dots \times (A/\mathfrak{a}_n)$$

be the map of A into the product induced by the canonical map of A onto A/\mathfrak{a}_i for each factor. Then the kernel of f is $\bigcap_{i=1}^n \mathfrak{a}_i$ and f is surjective, thus giving an isomorphism

$$A/\bigcap_{i=1}^n \mathfrak{a}_i \cong \prod_{i=1}^n A/\mathfrak{a}_i$$

Proof. Surjectivity follows from the theorem □

Let m be an integer > 1 , and let

$$m = \prod_i p_i^{r_i}$$

be a factorization of m into primes, with exponents $r_i \geq 1$. Then we have a ring isomorphism

$$\mathbb{Z}/m\mathbb{Z} \cong \prod_i \mathbb{Z}/p_i^{r_i}\mathbb{Z}$$

If A is a ring, we denote as usual by A^* the multiplicative group of invertible elements of A

Proposition 2.11. *The preceding ring isomorphism of $\mathbb{Z}/m\mathbb{Z}$ onto the product induces a group isomorphism*

$$(\mathbb{Z}/m\mathbb{Z})^* \cong \prod_i (\mathbb{Z}/p_i^{r_i}\mathbb{Z})^*$$

In view of our isomorphism, we have

$$\varphi(m) = \prod_i \varphi(p_i^{r_i})$$

If p is a prime number and r an integer ≥ 1 , then

$$\varphi(p^r) = (p-1)p^{r-1}$$

If $r = 1$, then $\mathbb{Z}/p\mathbb{Z}$ is a field, and the multiplicative group of that field has order $p-1$. Let r be ≥ 1 , and consider the canonical ring homomorphism

$$\mathbb{Z}/p^{r+1}\mathbb{Z} \rightarrow \mathbb{Z}/p^r\mathbb{Z}$$

arising from the inclusion of ideals $(p^{r+1}) \subset (p^r)$. We get an induced group homomorphism

$$\lambda : (\mathbb{Z}/p^{r+1}\mathbb{Z})^* \rightarrow (\mathbb{Z}/p^r\mathbb{Z})^*$$

which is surjective because any integer a which represents an element of $\mathbb{Z}/p^r\mathbb{Z}$ and is prime to p will represent an element of $(\mathbb{Z}/p^{r+1}\mathbb{Z})^*$. Let a be an integer representing an element of $(\mathbb{Z}/p^{r+1}\mathbb{Z})^*$ s.t. $\lambda(a) = 1$. Then

$$a \equiv 1 \pmod{p^r\mathbb{Z}}$$

P96

Application: The ring of endomorphisms of a cyclic group.

Theorem 2.12. *Let A be a cyclic group of order n . For each $k \in \mathbb{Z}$ let $f_k : A \rightarrow A$ be the endomorphism $x \mapsto kx$ (writing A additively). Then $k \mapsto f_k$ induces a ring homomorphism $\mathbb{Z}/n\mathbb{Z} \cong \text{End}(A)$, and a group isomorphism $(\mathbb{Z}/n\mathbb{Z})^* \cong \text{Aut}(A)$*

Proof. The fact that $k \mapsto f_k$ is ring homomorphism is a restatement of the formulas

$$1a = a, \quad (k + k')a = ka + k'a, \quad (kk')a = k(k'a)$$

□

2.3 Polynomials and Group Rings

Consider an infinite cyclic group generated by an element X . We let S be the subset consisting of powers X^r with $r \geq 0$. Then S is a monoid. We define the set of **polynomials** $A[X]$ to be the set of functions $S \rightarrow A$ which are equal to 0 except for a finite number of elements of S . For each element $a \in A$ we denote by aX^n the function which has the value a on X^n and the value 0 for all other elements of S . Then it is immediate that a polynomial can be written uniquely as a finite sum

$$a_0X^0 + \cdots + a_nX^n$$

for some integer $n \in \mathbb{N}$ and $a_i \in A$. Such a polynomial is denoted by $f(X)$. The elements $a_i \in A$ are called the **coefficients** of f . We define the product according to the convolution rule. Thus, given polynomials

$$f(X) = \sum_{i=0}^n a_i X^i \quad \text{and} \quad g(X) = \sum_{j=0}^m b_j X^j$$

we define the product to be

$$f(X)g(X) = \sum_{k=0}^{m+n} \left(\sum_{i+j=k} a_i b_j \right) X^k$$

This product is associative and distributive. $1X^0$ is the unit element. There is also an embedding

$$\begin{aligned} A &\rightarrow A[X] \\ a &\mapsto aX^0 \end{aligned}$$

Let A be a subring of a commutative ring B . Let $x \in B$. If $f \in A[X]$ is a polynomial, we define the associated **polynomial function**

$$f_B : B \rightarrow B$$

by letting

$$f_B(x) = f(x) = a_0 + a_1x + \cdots + a_nx^n$$

Given an element $b \in B$, directly from the definition of multiplication of polynomials, we find

Proposition 2.13. *The association*

$$ev_b : f \mapsto f(b)$$

is a ring homomorphism of $A[X]$ into B

This homomorphism is called the **evaluation homomorphism**, and is also said to be obtained by **substituting** b for X in the polynomial

Let $x \in B$. We see that the subring $A[x]$ of B generated by x over A is a ring of all polynomial values $f(x)$ for $f \in A[X]$. If the evaluation map $f \mapsto f(x)$ gives an isomorphism of $A[X]$ with $A[x]$, then we say that x is **transcendental** over A , or that x is a **variable** over A . In particular, X is a variable over A

Example 2.6. Let $\alpha = \sqrt{2}$. Then the set of all real numbers of the form $a + b\alpha$, with $a, b \in \mathbb{Z}$ is a subring of the real numbers, generated by $\sqrt{2}$. α is not transcendental over \mathbb{Z} , because the polynomial $X^2 - 2$ lies in the kernel of the evaluation map $f \mapsto f(\sqrt{2})$. On the other hand, it can be shown that e and π are transcendental over \mathbb{Q}

Example 2.7. Let p be a prime number and let $K = \mathbb{Z}/p\mathbb{Z}$. Then K is a field. Let $f(X) = X^p - X \in K[X]$. Then f is not the zero polynomial. But f_K is the zero function. Indeed, $f_K(0) = 0$. If $x \in K$, $x \neq 0$, then since the multiplicative group of K has order $p - 1$, it follows that $x^{p-1} = 1$, whence $x^p = x$, so $f(x) = 0$. Thus a non-zero polynomial gives rise to the zero function on K

Let

$$\varphi : A \rightarrow B$$

be a homomorphism of commutative rings. Then there is an associated homomorphism of the polynomial rings $A[X] \rightarrow B[X]$ s.t.

$$f(X) = \sum a_i X^i \mapsto \sum \varphi(a_i) X^i = (\varphi f)(X)$$

We call $f \mapsto \varphi f$ the **reduction map**

Let \mathfrak{p} be a prime ideal of A . Let $\varphi : A \rightarrow A/\mathfrak{p}$ be the canonical homomorphism of A onto A/\mathfrak{p} . If $f(X)$ is a polynomial in $A[X]$, then φf will sometimes be called the **reduction of f modulo \mathfrak{p}** .

For example, taking $A = \mathbb{Z}$ and $\mathfrak{p} = (p)$ for some prime number p , we can speak of the polynomial $3X^4 - X + 2$ as a polynomial mod 5, viewing the coefficients as elements of $\mathbb{Z}/5\mathbb{Z}$

Proposition 2.14. Let $\varphi : A \rightarrow B$ be a homomorphism of commutative rings. Let $x \in B$. There is a unique homomorphism extending φ

$$A[X] \rightarrow B \quad \text{s.t.} \quad X \mapsto x$$

and for this homomorphism $\sum a_i X^i \mapsto \sum \varphi(a_i) x^i$

The homomorphism of the above statement may be viewed as the composite

$$A[X] \longrightarrow B[X] \xrightarrow{\text{ev}_x} B$$

When writing a polynomial $f(X) = \sum_{i=1}^n a_i X^i$, if $a_n \neq 0$ then we define n to be the **degree** of f . Thus the degree of f is the smallest integer n s.t. $a_r = 0$ for $r > n$. If $f = 0$ (i.e. f is the zero polynomial), then by convention, we define the degree of f to be $-\infty$. We agree to the convention that

$$-\infty + -\infty = -\infty, \quad -\infty + n = -\infty, \quad -\infty < n$$

for all $n \in \mathbb{Z}$, and no other operation with $-\infty$ is defined. A polynomial of degree 1 is also called a **linear** polynomial. If $f \neq 0$ and $\deg f = n$ then we call a_n the **leading coefficient** of f . We call a_0 its **constant term**

Let

$$g(X) = b_0 + \cdots + b_m X^m$$

be a polynomial in $A[X]$, of degree m , and assume $g \neq 0$. Then

$$f(X)g(X) = a_0 b_0 + \cdots + a_n b_m X^{m+n}$$

Therefore

Proposition 2.15. If we assume that at least one of the leading coefficients a_n or b_m is not a divisor of 0 in A , then

$$\deg(fg) = \deg f + \deg g$$

and the leading coefficient of fg is $a_n b_m$. This holds in particular when a_n or b_m is a unit in A , or when A is entire. Consequently, when A is entire, $A[X]$ is also entire

If $f = 0$ or $g = 0$ we still have

$$\deg(fg) = \deg f + \deg g$$

if we agree that $-\infty + m = -\infty$ for any integer m

Let A be a subring of a commutative ring B . Let $x_1, \dots, x_n \in B$. For each n -tuple of integers $(v_1, \dots, v_n) = \mathbf{v} \in \mathbb{N}^n$, let $\mathbf{x} = (x_1, \dots, x_n)$, and

$$M_{\mathbf{v}}(\mathbf{x}) = x_1^{v_1} \dots x_n^{v_n}$$

The set of such elements forms a monoid under multiplication. Let $A[x] = A[x_1, \dots, x_n]$ be the subring of B generated by x_1, \dots, x_n over A . Then every element of $A[x]$ can be written as a finite sum

$$\sum a_{\mathbf{v}} M_{\mathbf{v}}(\mathbf{x}) \quad \text{and} \quad a_{\mathbf{v}} \in A$$

Using the construction of polynomials in one variable repeatedly, we may form the ring

$$A[X_1, \dots, X_n] = A[X_1][X_2] \dots [X_n]$$

selecting X_n to be variable over $A[X_1, \dots, X_{n-1}]$. Then every element f of $A[X_1, \dots, X_n] = A[X]$ has a *unique* expression as a finite sum

$$f = \sum_{j=0}^{d_n} f_j(X_1, \dots, X_{n-1}) X_n^j \quad \text{with} \quad f_j \in A[X_1, \dots, X_{n-1}]$$

Therefore by induction we can write f uniquely as a sum

$$\begin{aligned} f &= \sum_{v_n=0}^{d_n} \left(\sum_{v_1, \dots, v_{n-1}} a_{v_1 \dots v_n} X_1^{v_1} \dots X_{n-1}^{v_{n-1}} \right) X_n^{v_n} \\ &= \sum a_{\mathbf{v}} M_{\mathbf{v}}(X) = \sum a_{\mathbf{v}} X_1^{v_1} \dots X_n^{v_n} \end{aligned}$$

with elements $a_{\mathbf{v} \in A}$, which are called the **coefficients** of f . The products

$$M_{\mathbf{v}}(X) = X_1^{v_1} \dots X_n^{v_n}$$

will be called **primitive monomials**. Elements of $A[X]$ are called **polynomials** (in n variables). We call $a_{\mathbf{v}}$ its **coefficients**

Given $\mathbf{x} = (x_1, \dots, x_n)$ and f , we define

$$f(\mathbf{x}) = \sum a_{\mathbf{v}} M_{\mathbf{v}}(\mathbf{x}) = \sum a_{\mathbf{v}} x_1^{v_1} \dots x_n^{v_n}$$

Then the **evaluation map**

$$\text{ev}_{\mathbf{x}} : A[X] \rightarrow B \quad \text{with} \quad f \mapsto f(\mathbf{x})$$

is a ring homomorphism

Elements $x_1, \dots, x_n \in B$ are called **algebraically independent** over A if the evaluation map

$$f \mapsto f(x)$$

is injective. Equivalently, we could say that if $f \in A[X]$ is a polynomial and $f(x) = 0$ then $f = 0$; in other words, there are no non-trivial polynomial relations among x_1, \dots, x_n over A .

By the **degree** of a primitive monomial

$$M_v(X) = X_1^{v_1} \dots X_n^{v_n}$$

we shall mean the integer $|v| = v_1 + \dots + v_n$

A polynomial

$$aX_1^{v_1} \dots X_n^{v_n} \quad (a \in A)$$

will be called a **monomial**

If $f(X)$ is a polynomial in $A[X]$ written as

$$f(X) = \sum a_v X_1^{v_1} \dots X_n^{v_n}$$

we define the **degree** of f to be the maximum of the degrees of the monomials $M_v(X)$ s.t. $a_v \neq 0$. (Such monomials are said to **occur** in the polynomial)

For each integer $d \geq 0$, given a polynomial f , let $f^{(d)}$ be the sum of all monomials occurring in f and having degree d . Then

$$f = \sum_d f^{(d)}$$

Suppose $f \neq 0$, we say that f is **homogeneous** of degree d if $f = f^{(d)}$

Algebraically independent elements will also be called **variables**

2.4 Localization

A a commutative ring

By a **multiplicative subset** of A we shall mean a submonoid of A

We shall now construct the **quotient ring of A by S** , also known as the **ring of fractions of A by S**

We consider pairs (a, s) with $a \in A$ and $s \in S$. We define a relation

$$(a, s) \sim (a', s')$$

if there exists $s_1 \in S$ s.t.

$$s_1(s'a - sa') = 0$$

The equivalence class containing a pair (a, s) is denoted by a/s . The set of equivalence classes is denoted by $S^{-1}A$

if $0 \in S$, then $S^{-1}A$ has precisely one element $0/1$

$$(a/s)(a'/s') = aa'/ss'$$

$$\frac{a}{s} + \frac{a'}{s'} = \frac{s'a + sa'}{ss'}$$

Let $\varphi_S : A \rightarrow S^{-1}A$ be the s.t. $\varphi_S(a) = a/1$. Every element of $\varphi_S(S)$ is invertible in $S^{-1}(A)$ (the inverse of $s/1$ is $1/s$)

Let \mathcal{C} be the category whose objects are ring homomorphism

$$f : A \rightarrow B$$

s.t. for every $s \in S$ the elements $f(s)$ is invertible in B . If $f : A \rightarrow B$ and

Proposition 2.16. *Let A be an entire ring, and let S be a multiplicative subset which does not contain 0. Then*

$$\varphi_S : A \rightarrow S^{-1}A$$

is injective

Let A be an entire ring, and let S be the set of non-zero elements of A . Then S is a multiplicative set, and $S^{-1}A$ is then a field, called the **quotient field** or the *field of fractions of A .

3 Modules

3.1 Basic Definitions

Let A be a ring. A **left module** over A , or a left A -module M is an abelian group, together with an operation of A on M , s.t. for all $a, b \in A$ and $x, y \in M$

$$(a + b)x = ax + bx \quad \text{and} \quad a(x + y) = ax + ay$$

Let A be an entire ring and let M be an A -module. We define the **torsion submodule** M_{tor} to be the subset of elements $x \in M$ s.t. there exists $a \in A$, $a \neq 0$ s.t. $ax = 0$.

By a **module homomorphism** we means a map

$$f : M \rightarrow M'$$

which is an additive group homomorphism and s.t.

$$f(ax) = af(x)$$

for all $a \in A$ and $x \in M$. If we wish to refer to the ring A , we also say that f is an **A -homomorphism**, or also that it is an **A -linear map**

For any module M and M' , the map $\zeta : M \rightarrow M'$ s.t. $\zeta(x) = 0$ for all $x \in M$ is a homomorphism, called **zero**

Let $f : M \rightarrow M'$ be a homomorphism. By the **cokernel** of f we mean the factor module $M' / \text{im } f = M' / f(M)$.

Like groups

Proposition 3.1. *Let N, N' be two submodules of a module of M . Then $N + N'$ is also a submodule, and we have an isomorphism*

$$N / (N \cap N') \cong (N + N') / N'$$

If $M \supset M' \supset M''$ are modules, then

$$(M / M'') / (M' / M'') \cong M / M'$$

If $f : M \rightarrow M'$ is a module homomorphism, and N' is a submodule of M' , then $f^{-1}(N')$ is a submodule of M and we have a canonical injective homomorphism

$$\bar{f} : M / f^{-1}(N') \rightarrow M' / N'$$

If f is surjective, then \bar{f} is a module isomorphism

A sequence of module homomorphisms

$$M' \xrightarrow{f} M \xrightarrow{g} M''$$

is **exact** if $\text{im } f = \ker g$. If N is a submodule of M , then

$$0 \longrightarrow N \longrightarrow M \longrightarrow M/N \longrightarrow 0$$

If a homomorphism $u : N \rightarrow M$ is s.t.

$$0 \longrightarrow N \xrightarrow{u} M$$

is exact, then we also say that u is a **monomorphism** or an **embedding**. Dually if

$$N \xrightarrow{u} M \longrightarrow 0$$

is exact, we say that u is an **epimorphism**

Let A be a commutative ring. Let E, F be modules. By a **bilinear map**

$$g : E \times E \rightarrow F$$

we mean a map s.t. given $x \in E$ the map $y \mapsto g(x, y)$ is A -linear and given $y \in E$, the map $x \mapsto g(x, y)$ is A -linear. By an **A -algebra** we mean a module together with a bilinear map $g : E \times E \rightarrow E$. We view such a map as a law of composition on E .

3.2 The Group of Homomorphisms

Let A be a ring, and let X, X' be A -modules. We denote by $\text{Hom}_A(X', X)$ the set of A -homomorphisms of X' into X . Then $\text{Hom}_A(X', X)$ is an abelian group, the law of addition being that of addition for mappings into an abelian group.

If A is *commutative* then we can make $\text{Hom}_A(X', X)$ into an A -module by defining af for $a \in A$ and $f \in \text{Hom}_A(X', X)$ to be the map s.t.

$$(af)(x) = af(x)$$

Let Y be an A -module, and let

$$X' \xrightarrow{f} X$$

be an A -homomorphism. Then we get an induced homomorphism

$$\text{Hom}_A(f, Y) : \text{Hom}_A(X, Y) \rightarrow \text{Hom}_A(X', Y)$$

given by $g \mapsto g \circ f$. The fact that $\text{Hom}_A(f, Y)$ is a homomorphism is a rephrasing of the $(g_1 + g_2) \circ f = g_1 \circ f + g_2 \circ f$

If we have a sequence of A -homomorphisms

$$X' \longrightarrow X \longrightarrow X''$$

then we get an induced sequence

$$\text{Hom}_A(X', Y) \longleftarrow \text{Hom}_A(X, Y) \longleftarrow \text{Hom}_A(X'', Y)$$

Proposition 3.2. *A sequence*

$$X' \xrightarrow{\lambda} X \longrightarrow X'' \longrightarrow 0$$

is exact iff the sequence

$$\mathrm{Hom}_A(X', Y) \longleftarrow \mathrm{Hom}_A(X, Y) \longleftarrow \mathrm{Hom}_A(X'', Y) \longleftarrow 0$$

is exact for all Y

Proof. Suppose the first sequence is exact. If $g : X'' \rightarrow Y$ is an A -homomorphism, its image in $\mathrm{Hom}_A(X, Y)$ is obtained by composing g with the surjective map of X on X'' . If this composition is 0, it follows that $g = 0$. Consider a homomorphism $g : X \rightarrow Y$ s.t. the composition

$$X' \xrightarrow{\lambda} X \xrightarrow{g} Y$$

is 0. Then g vanishes on the image of λ . Hence we can factor g through the factor module

$$\begin{array}{ccc} & X/\mathrm{im} \lambda & \\ \nearrow & & \searrow \\ X & \xrightarrow{g} & Y \end{array}$$

Since $X \rightarrow X''$ is surjective, we have an isomorphism

$$X/\mathrm{im} \lambda \cong X''$$

Hence we can factor g through X'' , thereby showing that the kernel of

$$\mathrm{Hom}_A(X', Y) \longleftarrow \mathrm{Hom}_A(X, Y)$$

is contained in the image of

$$\mathrm{Hom}_A(X, Y) \longleftarrow \mathrm{Hom}_A(X'', Y)$$

□

similarly, we have

Proposition 3.3. *A sequence*

$$0 \longrightarrow Y' \longrightarrow Y \longrightarrow Y''$$

is exact iff

$$0 \longrightarrow \mathrm{Hom}_A(X, Y') \longrightarrow \mathrm{Hom}_A(X, Y) \longrightarrow \mathrm{Hom}_A(X, Y'')$$

is exact for all X

Let $\text{Mod}(A)$ and $\text{Mod}(B)$ be the categories of modules over rings A and B , and let $F : \text{Mod}(A) \rightarrow \text{Mod}(B)$ be a functor. One says that F is **exact** if F transforms exact sequences into exact sequences.

let M be an A -module. From the relations

$$\begin{aligned}(g_1 + g_2) \circ f &= g_1 \circ f + g_2 \circ f \\ g \circ (f_1 + f_2) &= g \circ f_1 + g \circ f_2\end{aligned}$$

and the fact that there is an identity for composition, namely id_M , we conclude that $\text{Hom}_A(M, M)$ is a ring. We call $\text{End}_A(M) = \text{Hom}_A(M, M)$ the ring of **endomorphisms**

3.3 Direct Products and Sums of Modules

Proposition 3.4. *Let M be an A -module and n an integer ≥ 1 . For each $i = 1, \dots, n$ let $\varphi_i : M \rightarrow M$ be an A -homomorphism s.t.*

$$\sum_{i=1}^n \varphi_i = \text{id} \quad \text{and} \quad \varphi_i \circ \varphi_j = 0 \quad \text{if } i \neq j$$

Then $\varphi_i^2 = \varphi_i$ for all i . Let $M_i = \varphi_i(M)$, and let $\varphi : M \rightarrow \prod M_i$ be s.t.

$$\varphi(x) = (\varphi_1(x), \dots, \varphi_n(x))$$

Then φ is an A -isomorphism of M onto the direct product $\prod M_i$

Proof. for each j , we have

$$\varphi_j = \varphi_j \circ \text{id} = \varphi_j \circ \sum_{i=1}^n \varphi_i = \varphi_j \circ \varphi_j = \varphi_j^2$$

thereby proving the first assertion. It is clear that φ is an A -homomorphism. Let $x \in \ker \varphi$. Since

$$x = \text{id}(x) = \sum_{i=1}^n \varphi_i(x)$$

we conclude that $x = 0$, so φ is injective. □

Let M be a module over a ring A and let S be a subset of M . By a **linear combination** of elements of S (with coefficients in A) one means a sum

$$\sum_{x \in S} a_x x$$

where $\{a_x\}$ is a set of elements of A , almost all of which are equal to 0. Let N be the set of all linear combinations of elements of S . Then N is a submodule of M , for if

$$\sum_{x \in S} a_x x \quad \text{and} \quad \sum_{x \in S} b_x x$$

are two linear combinations, then their sum is equal to

$$\sum_{x \in S} (a_x + b_x) x$$

and if $c \in A$, then

$$c \left(\sum_{x \in S} a_x x \right) = \sum_{x \in S} c a_x x$$

We shall call N the submodule **generated** by S , and we call S a set of **generators** for N . We sometimes write $N = A\langle S \rangle$. If S consists of one element x , the module generated by x is also written Ax , or simply (x) , and sometimes we say that (x) is a **principal module**

A module M is said to be **finitely generated**, or of **finite type** or **finite** over A , if it has a finite number of generators

A subset S of a module M is said to be **linearly independent** (over A) if whenever we have a linear combination

$$\sum_{x \in S} a_x x$$

which is equal to 0, then $a_x = 0$ for all $x \in S$. If S is linearly independent and if two linear combinations

$$\sum a_x x \quad \text{and} \quad \sum b_x x$$

are equal, then $a_x = b_x$ for all $x \in S$.

Let M be an A -module, and let $\{M_i\}_{i \in I}$ be a family of submodules. Since we have inclusion-homomorphism

$$\lambda_i : M_i \rightarrow M$$

we have an induced homomorphism

$$\lambda_* : \bigoplus M_i \rightarrow M$$

which is s.t. for any family of elements $(x_i)_{i \in I}$ all but a finite number of which are 0, we have

$$\lambda_*((x_i)) = \sum_{i \in I} x_i$$

if λ_* is an isomorphism, then we say that $\{M_i\}_{i \in I}$ is a **direct sum decomposition** of M . This is equivalent to saying that every element of M has a unique expression as a sum

$$\sum x_i$$

with $x_i \in M$ and almost all $x_i = 0$. By abuse of notation, we also write

$$M = \bigoplus_i M_i$$

in this case

If M is a module and N, N' are two submodules s.t. $N + N' = M$ and $N \cap N' = 0$, then we have a module isomorphism

$$M \cong N \oplus N'$$

Proposition 3.5. *Let M, M', N be modules. Then we have an isomorphism of abelian groups*

$$\text{Hom}_A(M \oplus M', N) \cong \text{Hom}_A(M, N) \times \text{Hom}_A(M', N)$$

and

$$\text{Hom}_A(N, M \times M') \cong \text{Hom}_A(N, M) \times \text{Hom}_A(N, M')$$

Proof. if $f : M \oplus M' \rightarrow N$ is a homomorphism, then f induces a homomorphism $f_1 : M \rightarrow N$ and a homomorphism $f_2 : M' \rightarrow N$ by composing injections

$$\begin{aligned} M &\rightarrow M \oplus \{0\} \subset M \oplus M' \xrightarrow{f} N \\ M' &\rightarrow \{0\} \oplus M' \subset M \oplus M' \xrightarrow{f} N \end{aligned}$$

Then

$$f \mapsto (f_1, f_2)$$

is an isomorphism □

Proposition 3.6. *Let $0 \rightarrow M' \xrightarrow{f} M \xrightarrow{g} M'' \rightarrow 0$ be an exact sequence of modules. The following are equivalent*

1. *there exists a homomorphism $\varphi : M' \rightarrow M$ s.t. $g \circ \varphi = \text{id}$*
2. *there exists a homomorphism $\psi : M \rightarrow M'$ s.t. $\psi \circ f = \text{id}$*

if these conditions are satisfied, then we have isomorphisms

$$\begin{aligned} M &= \text{im } f \oplus \ker \psi, & M &= \ker g \oplus \text{im } \varphi \\ M &\cong M' \oplus M'' \end{aligned}$$

Proof. Let $x \in M$, then $x - \varphi(g(x)) \in \ker g$, and hence $M = \ker g + \text{im } \varphi$. If $x \in \ker g \cap \text{im } \varphi$, then $x = \varphi(w)$ and $g(x) = g(\varphi(w)) = w = 0$, thus $\ker g \cap \text{im } \varphi = \{0\}$ \square

4 Algebraic Extensions

4.1 Finite and Algebraic Extensions

Let F be a field. If F is a subfield of a field E , then we also say that E is an **extension field** of F . We may view E as a vector space over F , and we say E is **finite** or **infinite** extension of F according as the dimension of this vector space is finite or infinite.

Let F be a subfield of a field E . An element α of E is said to be **algebraic** over F if there exists elements $a_0, \dots, a_n \in F$, not all equal to 0, s.t.

$$a_0 + a_1\alpha + \dots + a_n\alpha^n = 0$$

If $\alpha \neq 0$, and α is algebraic, then we can always find elements a_i as above s.t. $a_0 \neq 0$

Let X be a variable over F . We can also say that α is algebraic over F if the homomorphism

$$F[X] \rightarrow E$$

which is the identity on F and maps X on α has a non-zero kernel. In that case the kernel is an ideal which is principal, generated by a single polynomial $p(X)$, which we may assume has leading coefficient 1. We then have an isomorphism

$$F[X]/(p(X)) \cong F[\alpha]$$

5 Real Fields

5.1 Ordered Fields

Let K be a field. An **ordering** of K is a subset P of K having the following properties

ORD 1. Given $x \in K$, we have either $x \in P$, or $x = 0$ or $-x \in P$, and these three possibilities are mutually exclusive

ORD 2. If $x, y \in P$, then $x + y, xy \in P$

K is **ordered by P** , and we call P the set of **positive elements**

Suppose K is ordered by P . Since $1 \neq 0$ and $1 = 1^2 = (-1)^2$, we see that $1 \in P$. By **ORD 2**, it follows that $1 + \dots + 1 \in P$, whence K has characteristic 0. If $x \in P$ and $x \neq 0$, then $xx^{-1} = 1 \in P$ implies that $x^{-1} \in P$

Let E be a field. Then a product of sums of squares in E is a sum of squares.

If $a, b \in E$ are sum of squares and $b \neq 0$, then a/b is a sum of squares

Consider complex number:)

Let $x, y \in K$. We define $x < y$ to mean that $y - x \in P$. If $x < 0$ we say that x is **negative**.

If K is ordered and $x \in K, x \neq 0$, then x^2 is positive

If E has characteristic $\neq 2$, and -1 is a sum of squares in E , then every element $a \in E$ is a sum of squares, because $4a = (1 + a)^2 - (1 - a)^2$

If K is a field with an ordering P , and F is a subfield, then obviously, $P \cap F$ defines an ordering of F , which is called the **induced** ordering

Let K be an ordered field and let F be a subfield with the induced ordering. We put $|x| = x$ if $x > 0$ and $|x| = -x$ if $x < 0$. An element $\alpha \in K$ is **infinitely large** over F if $|\alpha| \geq x$ for all $x \in F$. It is **infinitely small** over F if $0 \leq |\alpha| \leq |x|$ for all $x \in F, x \neq 0$. α is infinitely large if and only if α^{-1} is infinitely small. K is **archimedean** over F if K has no elements which are infinitely large over F . An intermediate field $F_1, K \supset F_1 \supset F$ is **maximal archimedean over F** in K if it is archimedean over F and no other intermediate field containing F_1 is archimedean over F . We say that F is **maximal archimedean in K** if it is maximal archimedean over itself in K

Let K be an ordered field and F a subfield. Let \mathfrak{o} be the set of elements of K which are not infinitely large over F . Then \mathfrak{o} is a ring and that for any $\alpha \in K$, we have α or $\alpha^{-1} \in \mathfrak{o}$. Hence \mathfrak{o} is what is called a valuation ring, containing F . Let \mathfrak{m} be the ideal of all $\alpha \in K$ which are infinitely small over F . Then \mathfrak{m} is the unique maximal ideal of \mathfrak{o} , because any element in \mathfrak{o} which is not in \mathfrak{m} has an inverse in \mathfrak{o} . We call \mathfrak{o} the **valuation ring determined by the ordering of K/F**

Proposition 5.1. *Let K be an ordered field and F a subfield. Let \mathfrak{o} be the valuation ring determined by the ordering of K/F , and let \mathfrak{m} be its maximal ideal. Then $\mathfrak{o}/\mathfrak{m}$ is a real field.*

Proof. Otherwise, we could write

$$-1 = \sum \alpha_i^2 + a$$

with $\alpha_i \in \mathfrak{o}$ and $a \in \mathfrak{m}$. Since $\sum \alpha_i^2$ is positive and a is infinitely small, such a relation is clearly impossible \square