

Introductory model theory
with elements of universal algebra
(version 5)

Will Johnson

Autumn 2022

Preface

This is a provisional draft of the course notes for Introductory Model Theory at Fudan University in Autumn 2022. The goal is to present the basic topics of model theory without assuming much background in abstract algebra. Please send any comments or corrections to willjohnson@fudan.edu.cn.

The current draft has no examples, motivations, or explanations after Chapter 2.

The approach of these notes

Many textbooks in model theory assume a large amount of mathematical background knowledge, especially in abstract algebra. For example, Poizat’s *Course in Model Theory*—which is better than most textbooks in this regard—assumes you know the definition of fields, rings, homomorphisms, and ideals.

There is a good reason for this approach: model theory as a subject is mostly about the applications of mathematical logic to other branches of mathematics, especially abstract algebra. However, this works poorly for our class, which is not in the mathematics department and not intended solely for mathematics majors.

In contrast, these notes will try to minimize the required mathematical background. Model theory is a branch of mathematics, so it would be impossible to explain the subject without *any* mathematical background. I assume you know what sets, functions, and ordered pairs are. (Much of this material is reviewed in Appendix A.) But I will not assume any background in abstract algebra. In theory, mathematical logic is a prerequisite for this course, so these notes will assume a little bit of background from mathematical logic, mostly set theory.

One approach would be to avoid the concepts from abstract algebra

(fields, rings, etc.) entirely, instead focusing on combinatorial examples like the random graph. Model theory *can* be presented in this way, but one loses much of the motivation, examples, intuition, and applications of the subject.

Instead, we will build up elements of abstract algebra from scratch, defining concepts like fields, rings, homomorphisms, and ideals and proving their basic properties. This will allow us to work through the model theory of algebraically closed fields, which has served as an important example in the history of the subject.

Our approach towards abstract algebra will emphasize ideas from *universal algebra*. Universal algebra is a subject which finds the parallels between ring theory, group theory, lattice theory, and other topics in algebra. For example, “ideals” in ring theory and “normal subgroups” in group theory are both instances of the more fundamental concept of “congruences” in universal algebra.

There are two reasons to use universal algebra in these notes. On a basic level, it allows us to efficiently explain concepts in ring theory and group theory simultaneously. More importantly, universal algebra is tightly connected to model theory. One could say that universal algebra is a special case of model theory, or model theory is a generalization of universal algebra. For example, Chang and Keisler describe model theory as “universal algebra plus logic” in their textbook. Consequently, one of the goals of these notes will be to emphasize the parallels between universal algebra and model theory.

Notation and conventions

We generally follow standard mathematical notation, some of which is reviewed in Appendix A. Here are some points to be aware of:

- $A \subseteq B$ means that A is a subset of B . $A \subsetneq B$ means that A is a proper subset of B . We don’t use the symbol $A \subset B$. $A \not\subseteq B$ means that A is not a subset of B .
- $A \subseteq_f B$ means that A is a finite subset of B .
- The powerset of S is written $\mathfrak{P}(S)$.
- The empty set is written \emptyset .

- \bar{x} means (x_1, x_2, \dots, x_n) , where n is determined by the context. The symbols $\bar{a}, \bar{b}, \bar{c}, \bar{d}, \bar{y}, \bar{z}, \dots$ work similarly.
- $f : A \rightarrow B$ means that f is a function from A to B . If we are talking about some function f , then $x \mapsto y$ means $f(x) = y$. If $X \subseteq A$, then $f(X)$ denotes the direct image, and if $X \subseteq B$ then $f^{-1}(X)$ denotes the inverse image. The domain and range of f are written $\text{dom}(f)$ and $\text{im}(f)$.
- $x := \dots$ means “[by the way,] let x be \dots ”. The notation $\dots =: x$ means the same thing.
- \mathbb{N} means the natural numbers, including 0:

$$\mathbb{N} = \{0, 1, 2, 3, \dots\}.$$

We also denote this set by ω or \aleph_0 .

- $\mathbb{R}, \mathbb{Q}, \mathbb{Z}$, and \mathbb{C} denote the real numbers, the rational numbers, the integers, and the complex numbers, respectively.
- $\forall x$ means “for every x ”, and $\exists x$ mean “there is an x such that”.
- If E is an equivalence relation on some set X and a is an element, then $[a]$ or $[a]_E$ denotes the equivalence class of a , and X/E denotes the set of all equivalence classes.
- If A is a set, we write the cardinality (size) of A as $|A|$.

Introduction

The story of model theory is like a novel that unexpectedly changes genres halfway through. What began as a branch of mathematical logic later became “universal algebra plus logic” in Chang and Keisler’s formulation, and then “algebraic geometry minus fields” in Hodges’ formulation. This makes the subject a little hard to precisely define.

At the end of the day, model theory is a network of closely connected definitions and tools, all related to the notion of “model.” This introduction is a survey of these definitions and tools. The goal of model theory is not so much to understand “models”, but instead to build on this network and apply it to other branches of mathematics.

0.1 Models

Mathematical knowledge is obtained through deductive reasoning—through *proofs*. Proofs must begin somewhere, with a set of *axioms*—statements we assume without proof. When reasoning about numbers, one might take the following set of statements as axioms:

1. If x and y are numbers, then $x + y$ and $x \cdot y$ are numbers.
2. If x and y are numbers, then $x + y = y + x$ and $x \cdot y = y \cdot x$.
3. If x , y , and z are numbers, then $x + (y + z) = (x + y) + z$ and $x \cdot (y \cdot z) = (x \cdot y) \cdot z$.
4. If x , y , and z are numbers, then $x \cdot (y + z) = (x \cdot y) + (x \cdot z)$.
5. There is a number 0 such that $x + 0 = x$ for any number x .
6. There is a number 1 such that $x \cdot 1 = x$ for any number x .

7. For any number x , there is a number called $-x$ such that $x + (-x) = 0$.
8. For any number x other than 0, there is a number called x^{-1} such that $x \cdot (x^{-1}) = 1$.

We can use these axioms to deduce other facts about numbers. Here is an example:

Theorem 0.1.1. *For any number x , $x \cdot 0 = 0$.*

Proof. If $y = x \cdot 0$, then

$$\begin{aligned}
 y &\stackrel{(5)}{=} y + 0 \stackrel{(7)}{=} y + (y + (-y)) \stackrel{(3)}{=} (y + y) + (-y) \\
 &= ((x \cdot 0) + (x \cdot 0)) + (-y) \stackrel{(4)}{=} x \cdot (0 + 0) + (-y) \\
 &\stackrel{(5)}{=} x \cdot 0 + (-y) = y + (-y) \stackrel{(7)}{=} 0. \quad \square
 \end{aligned}$$

With more work (Corollary 1.4.14, Theorem 1.4.16), one can prove other algebraic facts, such as

$$\begin{aligned}
 x \cdot y = 0 &\iff (x = 0 \text{ or } y = 0) \\
 (-x) \cdot (-y) &= x \cdot y.
 \end{aligned}$$

How about the following?

$$x + x = y + y \stackrel{?}{\implies} x = y. \quad (*)$$

This is certainly true for real numbers, but can we prove $(*)$ from the axioms? Surprisingly, **we cannot**. To see this, suppose we that 0 and 1 are the only “numbers”, and that “+” and “ \cdot ” are defined as follows:

+	0	1
0	0	1
1	1	0

\cdot	0	1
0	0	0
1	0	1

For example, “ $1 + 1 = 0$ ” under this interpretation. By checking all the cases, one can verify that all of Axioms (1)–(8) hold. Anything which is provable from the axioms, such as Theorem 0.1.1, must be true in this interpretation. However, $(*)$ fails:

$$1 + 1 = 0 + 0 \text{ but } 1 \neq 0.$$

Therefore $(*)$ cannot be proven from Axioms (1)–(8).

Our strange way of interpreting “number”, “0”, and “1” is an example of a **model**—in this case a model of Axioms (1)–(8).

Remark 0.1.2. Axioms (1)–(8) are called the *field axioms*, and their models are called *fields*. Fields are one of the central objects of study in the branch of mathematics called *abstract algebra*.

Some history

The toy example above is parallel to an important saga in the history of mathematics. In ancient times, the Hellenistic mathematician Euclid wrote a book called the *Elements* in which he developed geometry from a set of five axioms. Four of these axioms were simple, intuitive statements, like “any two points are on a line” or “any two right angles are congruent.” In contrast, the fifth axiom was a complicated geometric statement about parallel lines. This was the so-called *parallel postulate*.

Many people were unhappy with the parallel postulate, because it seemed less obvious and more arbitrary than the other four axioms. It felt like something that should be a theorem or a lemma, not an axiom. Euclid himself avoided using the parallel postulate until absolutely necessary. *If* one could find a proof of the parallel postulate from the other four axioms, the problem would go away: the parallel postulate could be demoted to a theorem and geometry could be developed on the firm foundation of the four intuitive axioms.

In the 2000 years after Euclid, there were many unsuccessful attempts to find such a proof. Eventually the matter was settled in 1868 when the mathematician Beltrami constructed a model of Euclid’s first four axioms in which the parallel postulate is false. This model is called *hyperbolic geometry*, and is the precursor to the sort of “curved” geometries used in Einstein’s theory of general relativity.

0.2 Complete theories

A *theory* is a set of axioms. If T is a theory, we write $M \models T$ to indicate that M is a model of T , and $M \models \varphi$ to indicate that M satisfies a sentence φ . For example, if T is the field axioms (1)–(8) of the previous section and φ is Statement (*), then the standard real numbers \mathbb{R} satisfy both T and φ

$$\mathbb{R} \models T \text{ and } \mathbb{R} \models \varphi,$$

but we constructed a different model M such that

$$M \models T \text{ and } M \not\models \varphi.$$

A theory T is *complete* if for every sentence φ , either T proves φ or T disproves φ . If we can completely axiomatize a structure, then we can determine all its logical properties:

Fact 0.2.1. *Let T be a complete theory.*

1. *If $M \models T$, then for any sentence φ ,*

$$M \models \varphi \iff (\varphi \text{ is provable from } T)$$

2. *If T is finite or computable, then there is an algorithm which takes a sentence φ as input and outputs whether or not φ is provable from T .*

The problem we ran into in the previous section is that the field axioms are not complete—they neither prove nor disprove the statement (*). Can we fix the problem and write down a complete axiomatization for numbers?

To begin with, let's add the following axioms:

$$\begin{aligned} 1 + 1 &\neq 0 \\ 1 + 1 + 1 &\neq 0 \\ 1 + 1 + 1 + 1 &\neq 0 \\ &\dots \end{aligned}$$

The first of these axioms gets rid of the bad model M of the previous section, which is a good start. Models of the resulting theory are called *fields of characteristic 0*.

Sadly, the resulting theory is still not complete. For example, consider the statement

$$\text{For every number } x, \text{ there is a number } y \text{ with } y^2 = x.$$

This is true if “number” means “complex number”, and false if “number” means “real number.” In other words, the two models \mathbb{C} and \mathbb{R} show incompleteness.

We should probably decide whether we are trying to axiomatize the complex numbers \mathbb{C} or the real numbers \mathbb{R} . It turns out to be easier to axiomatize \mathbb{C} .

Definition 0.2.2. A field is *algebraically closed* if it satisfies the following axiom: for any numbers a_0, a_1, \dots, a_{n-1} , there is a number x such that

$$x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0 = 0.$$

In other words, every polynomial equation has a solution.

The field of complex numbers \mathbb{C} is algebraically closed, a fact known as the *fundamental theorem of algebra*. In contrast, the field of real numbers \mathbb{R} is *not* algebraically closed, as $x^2 + 1 = 0$ has no solutions.

If we combine the axioms defining fields of characteristic zero and the axioms defining algebraically closed fields, we get a theory called ACF_0 , whose models are algebraically closed fields of characteristic 0. The complex numbers are a model. In Section 9.4 we will prove the following:

Fact 0.2.3. *The theory ACF_0 is complete.*

Consequently:

1. A sentence φ is true in \mathbb{C} if and only if it is provable from the ACF_0 axioms.
2. There is an algorithm which takes φ as input and determines whether φ is true in \mathbb{C} .

Remark 0.2.4. There is also a theory called RCF which completely axiomatizes \mathbb{R} . Models of RCF are called *real closed fields*.

0.3 Categoricity

Model theory provides a number of methods to prove that a theory T is complete by analyzing the structure of the models of T . One such method is *categoricity*.

Definition 0.3.1. Let κ be an infinite cardinal number like \aleph_0 or \aleph_1 . A theory T is κ -categorical if T has exactly one model of size κ .

Fact 0.3.2 (Łoś-Vaught criterion). *If a theory T is κ -categorical and all models of T are infinite, then T is complete.*

Conveniently, it is a theorem in abstract algebra that ACF_0 is κ -categorical for all uncountable κ :

Fact 0.3.3 (Steinitz). *If κ is an uncountable cardinal, then there is exactly one algebraically closed field of characteristic 0 and size κ .*

Algebraically closed fields are infinite (see Theorem 9.2.3), and so ACF_0 is complete by the Łoś-Vaught criterion.

Remark 0.3.4. This technique is less flexible than one might hope for. In fact, algebraically closed fields are the *only* fields which can be completely axiomatized by κ -categorical theories. We will see a little bit of the proof in Section 13.4, but the bulk of the proof requires stability theory and is beyond the scope of this course.

0.4 Definable sets

If M is a model, a subset $D \subseteq M$ is *definable* if $D = \{x \in M : \varphi(x)\}$ where $\varphi(x)$ is a logical statement about x . For example, the set

$$\{x \in \mathbb{R} : x \text{ has a square root}\} = \{x \in \mathbb{R} : x \geq 0\}$$

is definable in the field \mathbb{R} . More generally, a set $D \subseteq M^n$ in n variables is definable if

$$D = \{(x_1, \dots, x_n) \in M^n : \varphi(x_1, \dots, x_n)\}$$

where $\varphi(x_1, \dots, x_n)$ is a statement about the variables x_1, \dots, x_n . For example, the unit circle is definable in the field \mathbb{R}

$$\{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}. \quad (\dagger)$$

In some models, there are very many definable sets:

Fact 0.4.1 (Robinson). *In the field \mathbb{Q} , every computable set $S \subseteq \mathbb{Q}^n$ is definable.*

For example, the set of Fibonacci numbers $\{1, 2, 3, 5, 8, 13, \dots\}$ is definable. We will see a variant of Fact 0.4.1 in Section 3.9.

In other models, there are very few definable sets:

Definition 0.4.2. In a field K , a set is *constructible* if it is a finite union of sets of the form

$$\begin{aligned} \{(x_1, \dots, x_n) \in K^n : P_1(x_1, \dots, x_n) = P_2(x_1, \dots, x_n) = \dots \\ = P_m(x_1, \dots, x_n) = 0 \neq Q(x_1, \dots, x_n)\} \end{aligned}$$

where P_1, \dots, P_m, Q are polynomials.

For example, the unit circle (\dagger) is constructible.

Fact 0.4.3. *In algebraically closed fields such as \mathbb{C} , the definable sets are exactly the constructible sets.*

Constructible sets are a slight generalization of what algebraic geometers call *varieties*, and Fact 0.4.3 is equivalent to a fact called Chevalley's Theorem in algebraic geometry. We will prove Fact 0.4.3 in Section 9.4 as a consequence of *quantifier elimination*.

One consequence of Fact 0.4.3 is that definable sets in one variable are very simple:

Corollary 0.4.4. *If M is an algebraically closed field and $D \subseteq M$ is definable, then D is finite or the complement of a finite set.*

One says that a theory T is *strongly minimal* if the models of T have this property. It turns out that strong minimality is closely connected to categoricity.

Fact 0.4.5 (= Corollary 14.2.7). *If T is strongly minimal and complete, then T is κ -categorical for all $\kappa > \aleph_0$.*

Remark 0.4.6. In more advanced model theory, one can show that if T is κ -categorical for at least one $\kappa > \aleph_0$, then T is tightly connected to a strongly minimal theory in a certain sense. As a consequence, T is κ -categorical for all $\kappa > \aleph_0$, a fact known as *Morley's Theorem*.

Strong minimality also has an interesting consequence for definable sets in more than one variable:

Fact 0.4.7. *Let M be a model of a strongly minimal theory. To each definable set $D \subseteq M^n$, we can associate a natural number called the dimension of D , written $\dim(D)$, with the following properties, among others:*

1. $\dim(X) > 0$ if and only if D is infinite.
2. $\dim(M^n) = n$.
3. $\dim(X \cup Y) = \max(\dim(X), \dim(Y))$.
4. $\dim(X \times Y) = \dim(X) + \dim(Y)$.
5. If $f : X \rightarrow Y$ is a definable bijection, then $\dim(X) = \dim(Y)$.

The intuition is that $\dim(D)$ measures the number of “degrees of freedom” of a value in D . For example, the unit circle $x^2 + y^2 = 1$ has one degree of freedom, because the value x almost determines the pair (x, y) . In contrast, the plane \mathbb{C}^2 has two degrees of freedom, because x and y can vary freely. We will verify Fact 0.4.7 in Sections 14.5–14.6.

In the case of algebraically closed fields, $\dim(D)$ is something that algebraic geometers call *Krull dimension*. More generally, model theorists like to take concepts from algebraic geometry and extend them from algebraically closed fields to other strongly minimal theories. This is perhaps why Hodges characterizes model theory as “algebraic geometry minus fields” in his textbook.

0.5 An outline of this book

Chapters 1 and 2 are about *universal algebra*, which is the special case of model theory where axioms must be *equations* like the distributive law or associative law

$$\begin{aligned}x \cdot (y + z) &= (x \cdot y) + (x \cdot z) \\x + (y + z) &= (x + y) + z\end{aligned}$$

rather than more complicated sentences like

For every x , there is a y such that $y \cdot y = x$ or $y \cdot y = -x$.

An *equational theory* is a set of equations. Many important classes of structures from abstract algebra, such as groups and rings, are defined by equational theories, as we will see in Chapter 1. Moreover, certain algebraic constructions like products and quotients make sense in any equational theory, as we will see in Chapter 2. In fact, these constructions precisely characterize which classes of structures are defined by equational theories, a fact known as *Birkhoff’s HSP theorem* (Theorem 2.9.2). Along the way, we will develop basic ring theory, using it as a running example of universal algebra. This ring theory will be used later in Chapter 9 when we axiomatize the complex numbers.

We begin the study of model theory proper in Chapter 3, where we define the basic notions of languages, structures, formulas, satisfaction, theories, models, elementary classes, elementary equivalence, elementary maps,

and definable sets. Chapter 4 is about the *compactness theorem*, the most important fundamental tool of model theory. We apply the compactness theorem in Chapter 5 on categoricity, where we prove the Łoś-Vaught criterion (Theorem 5.6.2) mentioned above.

Chapter 6 is about *ultraproducts*, a mysterious construction which plays the same role in general model theory that products, quotients, etc. play in universal algebra. We use ultraproducts to give another proof of the compactness theorem (Theorem 6.2.6), and then discuss some analogues of Birkhoff’s HSP theorem using ultraproducts in Sections 6.3–6.4.

In Chapter 7, we review basic pointset topology, and apply it to build the topological space S of complete theories. The compactness theorem can be understood as the statement that S is compact (Theorem 7.3.4), and ultraproducts correspond to a certain kind of limit in this topological space (Theorem 7.3.5). The construction of S serves as a prototype of several constructions appearing later in the book.

Chapter 8 is about types and quantifier elimination, two topics which happen to be interconnected. A *type* is a description of a potential object which may not exist in the current model, but exists in some other model. Most applications of the compactness theorem boil down to *realizing* types—finding the element described by a type. Meanwhile, *quantifier elimination* is an important technical tool that allows us to model-theoretically analyze good theories and structures. For example, quantifier elimination gives control over definable sets, and can be used to detect whether a theory is complete.

In Chapter 9, we begin studying the model theory of algebraically closed fields, leading to the complete axiomatization of the complex numbers in Corollary 9.4.7. We continue to develop the model theory of algebraically closed fields in later chapters, using it as a running example.

The next two chapters are about models where “anything which can happen does happen”, in two slightly different senses. Chapter 10 is about *existentially closed models*. Existentially closed models are a useful way of constructing new models of a given theory—for example, we will use them to show that algebraically closed fields *exist*. Existentially closed models can also be used to construct new theories and prove quantifier elimination.

Meanwhile, Chapter 11 is about *monster models*—models where every “small” type is realized. Monster models exist (Theorem 11.4.8), and possess a number of nice properties with respect to definable sets, types, and automorphisms (Sections 11.2–11.3). In advanced model theory, it is common to fix a monster model \mathbb{M} and always work inside \mathbb{M} . Monster models

also provide a nice way of thinking about quantifier elimination (see Theorem 11.5.4).

The final three chapters return to the theme of categoricity. Recall that a theory is κ -categorical if it has exactly one model of size κ . Chapter 12 is about *countable categoricity*, the case where $\kappa = \aleph_0$. It turns out that a structure M can be axiomatized by an \aleph_0 -categorical theory if and only if M satisfies certain structural properties with respect to types, definable sets, and automorphisms (Theorems 12.3.5 and 12.3.6). Chapter 13 is a digression on abstract *closure operations* and the special family of closure operations known as *pregeometries* or *matroids*. We use pregeometries to classify vector spaces, getting more examples of κ -categorical theories (Corollary 13.7.6).

Finally, Chapter 14 is about *strongly minimal theories*, a special class of uncountably categorical theories including algebraically closed fields. Using pregeometries, we prove that strongly minimal theories are uncountably categorical (Corollary 14.2.7), and we develop the dimension theory described in Section 0.4 above. In fact, this dimension theory works in *any* uncountably categorical theory, but the proof is beyond the scope of this book.

Contents

0.1	Models	7
0.2	Complete theories	9
0.3	Categoricity	11
0.4	Definable sets	12
0.5	An outline of this book	14
1	Algebras and equational classes	21
1.1	Monoids and groups	22
1.2	Rings and fields	25
1.3	Languages, algebras, and equational classes	26
1.4	Homomorphisms and isomorphisms	32
2	New algebras from old	39
2.1	Subalgebras and generators	40
2.2	Products	43
2.3	Descending functions along surjections	49
2.4	Congruences and quotients	50
2.5	Application: finite fields	57
2.6	The fundamental theorem on homomorphisms	60
2.7	Application: characteristic of fields	63
2.8	Term algebras and free algebras	66
2.9	Birkhoff's HSP theorem	70
3	First-order logic	75
3.1	Languages, structures, formulas, and satisfaction	76
3.2	Theories and elementary classes	79
3.3	Common abbreviations	80
3.4	Elementary equivalence and embeddings	80
3.5	Substructures and extensions	82

3.6	Complete theories	83
3.7	Completeness via back-and-forth systems	85
3.8	Definable sets and functions	87
3.9	Incompleteness of Peano Arithmetic	88
4	The compactness theorem	93
4.1	A temporary convention	94
4.2	Premodels	94
4.3	The witness property	96
4.4	Compactness via Henkin's method	98
5	Categoricity	101
5.1	A useful fact around finite satisfiability	102
5.2	Diagrams and elementary amalgamation	102
5.3	Tarski-Vaught	104
5.4	The Löwenheim-Skolem theorem	105
5.5	Absolute categoricity	107
5.6	κ -categoricity	108
6	Ultraproducts	111
6.1	Ultrafilters	112
6.2	Ultraproducts	114
6.3	Characterizations of elementary classes	117
6.4	Universal classes	118
7	Topologies	121
7.1	Topological spaces	121
7.2	Ultralimits	124
7.3	The space of complete theories	127
7.4	Stone spaces	129
8	Types and quantifier elimination	131
8.1	Types	132
8.2	The topology on type space	136
8.3	Quantifier elimination	138
8.4	Quantifier elimination in DLO	141

9	Algebraically closed fields	143
9.1	Polynomial rings	144
9.2	The theory ACF	147
9.3	Algebraic and transcendental elements	147
9.4	Quantifier elimination in ACF	148
10	Existentially closed models	153
10.1	Unions of chains of structures	153
10.2	Existentially closed models	154
10.3	Existentially closed fields	156
11	Monster models	159
11.1	Pushing types along partial elementary maps	159
11.2	κ -saturated models	161
11.3	Strongly κ -homogeneous models	163
11.4	Construction of monster models	165
11.5	Saturation and back-and-forth	169
11.6	Application: discrete linear orders with endpoints	170
12	Countable categoricity	175
12.1	Baire category theorem	176
12.2	The omitting types theorem	177
12.3	Countably categorical theories	179
13	Closure operators and pregeometries	183
13.1	Closure operators	184
13.2	Algebraic and definable closure	186
13.3	Definable closure, algebraic closure, and substructures	187
13.4	Countably categorical fields	188
13.5	Algebraic closure in ACF	189
13.6	Pregeometries and vector spaces	191
13.7	The classification of vector spaces	197
14	Strong minimality and geometric theories	201
14.1	Strong minimality	202
14.2	Uncountable categoricity	203
14.3	Uniform finiteness	205
14.4	Pregeometric theories	207

14.5	Dimension theory in the monster	209
14.6	Dimension theory in small models	213
A	Mathematical background	217
A.1	Logical notation	217
A.2	Sets	218
A.3	$\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$	219
A.4	Tuples, products, and relations	219
A.5	Functions	221
A.6	Cardinalities	223
B	Set theory	225
B.1	Ordinals	225
B.2	Induction and recursion	226
B.3	Cardinals	227
B.4	Zorn's lemma	228

Chapter 1

Algebras and equational classes

In this chapter, we introduce several important classes of structures from abstract algebra, including *monoids*, *groups*, *rings*, and *fields*. The definitions of these concepts look very similar to each other on an abstract level. Each definition has the form

A FOO is a set together with some operations, satisfying a certain list of equations,

where FOO is “monoid”, “group”, “ring”, and so on.

This suggests the abstract notion of an *equational class*—a class of structures defined by a set of equations. The class of monoids is an equational class, and the same holds for groups and rings. We precisely define equational classes in Section 1.3. Equational classes allow us to uniformly treat certain concepts in algebra. For example, in Section 1.4, we define the important algebraic concepts of *homomorphisms* and *isomorphisms* in a uniform way across all equational classes.

The study of equational classes is called *universal algebra*, and is the subject of this chapter and Chapter 2. Model theory proper, which begins in Chapter 3, can be seen as a generalization of universal algebra. We will use universal algebra as both a source of inspiration for model theory, and as a technical tool to develop the ring theory we will need later when we axiomatize the complex numbers in Chapter 9.

1.1 Monoids and groups

Let A be a set. For $k \geq 0$, a k -ary operation or k -ary function on A is a function $f : A^k \rightarrow A$, that is, a function which takes k inputs from A and outputs one value in A . *Binary*, *unary*, and *nullary* mean 2-ary, 1-ary, and 0-ary, respectively.

For example, the function

$$\max(x, y) = \begin{cases} x & \text{if } x \geq y \\ y & \text{if } x \leq y \end{cases}$$

is a function $\mathbb{R}^2 \rightarrow \mathbb{R}$, so it is a binary (2-ary) operation on \mathbb{R} . Other binary operations on \mathbb{R} include $+$, $-$, and \times . Most binary operations are written in infix notation, like $2 + 3$, rather than prefix notation, like $+(2, 3)$.

Note that A^0 has one element $()$. Therefore, we can identify nullary functions $f : A^0 \rightarrow A$ with elements of A via the correspondence

$$f \mapsto f().$$

For example, we identify the element $8 \in \mathbb{R}$ with the function $f : \mathbb{R}^0 \rightarrow \mathbb{R}$ sending the one element $()$ in \mathbb{R}^0 to the element $8 \in \mathbb{R}^1$.

A *magma* is a pair (A, \star) where A is a set and \star is a binary operation on A . For example, $(\mathbb{R}, +)$ and $(\mathbb{R}, -)$ are magmas. A magma (A, \star) is a semigroup if \star is associative, meaning that:

$$x \star (y \star z) = (x \star y) \star z$$

for all $x, y, z \in A$. For example, $(\mathbb{R}, +)$ and (\mathbb{R}, \max) are semigroups, because

$$\begin{aligned} x + (y + z) &= (x + y) + z \\ \max(x, \max(y, z)) &= \max(x, y, z) = \max(\max(x, y), z). \end{aligned}$$

On the other hand, $(\mathbb{R}, -)$ is not a semigroup, because

$$5 - (3 - 2) = 4 \neq (5 - 3) - 2 = 0.$$

Let (A, \star) be a semigroup. An element $e \in A$ is an *identity element* if

$$x \star e = x = e \star x$$

holds for all $x \in A$. For example, 0 is an identity element in the semigroup $(\mathbb{R}, +)$ because $x + 0 = 0 + x = x$. Similarly, 1 is an identity element in the semigroup (\mathbb{R}, \cdot) , because $x \cdot 1 = 1 \cdot x = x$. In contrast, (\mathbb{R}, \max) has no identity element.

Theorem 1.1.1. *If (A, \star) is a semigroup, there is at most one identity element.*

Proof. Suppose e_1, e_2 are identity elements. Then $e_1 \star e_2 = e_1$ (because e_2 is an identity element), and $e_1 \star e_2 = e_2$ (because e_1 is an identity element). Therefore $e_1 = e_2$. \square

Consequently, we can talk about “the” identity element, when it exists.

Definition 1.1.2. A *monoid* is a triple (A, \star, e) where (A, \star) is a semigroup and e is an identity element.

For example, $(\mathbb{R}, +, 0)$ and $(\mathbb{R}, \cdot, 1)$ are monoids.

Remark 1.1.3. The traditional definition of “monoid” is “a semigroup in which an identity element exists.” With the traditional definition, $(\mathbb{R}, +)$ is a monoid, rather than $(\mathbb{R}, +, 0)$. However, there is a clear correspondence between the traditional definition and Definition 1.1.2, thanks to Theorem 1.1.1. Definition 1.1.2 works slightly better for the purposes of universal algebra.

Let (A, \star, e) be a monoid. If $x \in A$, then an *inverse* of x is an element $x' \in A$ such that $x \star x' = x' \star x = e$.

Example 1.1.4. In the monoid $(\mathbb{R}, \cdot, 1)$, the element $2/3$ is an inverse of $3/2$, while the element 0 has no inverse.

Theorem 1.1.5. *Let (A, \star, e) be a monoid. If $x \in A$, then x has at most one inverse.*

Proof. Suppose y, z are both inverses of x . Then

$$y = y \star e = y \star (x \star z) = (y \star x) \star z = e \star z = z. \quad \square$$

Consequently, we can talk about “the” inverse of x , when it exists.

Definition 1.1.6. A *group* is a 4-tuple $(A, \star, e, (-)')$ where (A, \star, e) is a monoid and $(-)'$ is a unary function $A \rightarrow A$ such that x' is an inverse of x for every $x \in A$.

Equivalently, a group is a 4-tuple $(A, \star, e, (-)')$ where A is a set, \star is a binary operation on A , $e \in A$ is a nullary operation (an element), and $(-)'$ is a unary operation such that the following equations hold for all $x, y, z \in A$.

$$\begin{aligned} x \star (y \star z) &= (x \star y) \star z \\ x \star e &= e \star x = x \\ x \star x' &= x' \star x = e. \end{aligned}$$

Example 1.1.7. $(\mathbb{R}, +, 0, -)$ is a group, where $-$ is the unary negation function $-x$.

Again, the traditional definition of “group” is slightly different—a group is a monoid (A, \star) in which every element has an inverse. With this definition, $(\mathbb{R}, +)$ is a group rather than $(\mathbb{R}, +, 0, -)$. Again, there is a clear correspondence between the traditional definition and Definition 1.1.6, and Definition 1.1.6 is better for universal algebra.

A semigroup, monoid, or group is *commutative* if the equation

$$x \star y = y \star x$$

holds for any x, y . Commutative groups are usually called *abelian groups*. So far, all our examples have been commutative. Here is an important example of a non-abelian group.

Example 1.1.8. Let S be a set. Let $\text{Perm}(S)$ be the set of bijections $f : S \rightarrow S$. Then $\text{Perm}(S)$ is a group $(\text{Perm}(S), \circ, \text{id}, (-)^{-1})$, where

1. $f \circ g$ is the function composition $(f \circ g)(x) := f(g(x))$.
2. $\text{id} : S \rightarrow S$ is the identity function $\text{id}(x) = x$.
3. f^{-1} is the inverse function of f .

$\text{Perm}(S)$ is usually non-abelian, since function composition is non-abelian in general. For example, if $f, g \in \text{Perm}(\mathbb{R})$ are

$$\begin{aligned} f(x) &= x + 1 \\ g(x) &= 2x \end{aligned}$$

then $(f \circ g)(x) = 2x + 1$ and $(g \circ f)(x) = 2(x + 1) = 2x + 2$, so $f \circ g \neq g \circ f$.

When S is $\{1, 2, \dots, n\}$, the group $\text{Perm}(S)$ is called the *n th symmetric group*.

Example 1.1.9. If you know linear algebra, the group of $n \times n$ invertible matrices is a group with respect to matrix multiplication. This group is called the *n th general linear group*, usually written $GL(n)$ or $GL_n(\mathbb{R})$.

Remark 1.1.10. Groups are usually written using multiplicative or additive notation.

	Multiplicative notation	Additive notation
$x \star y$	$x \cdot y$	$x + y$
e	1	0
x'	x^{-1}	$-x$

When using multiplicative notation, one uses the usual abbreviations for multiplication, writing $xy^{-1}z$ instead of $x \cdot (y^{-1}) \cdot z$. Here are the group axioms in multiplicative notation:

$$\begin{aligned} x(yz) &= (xy)z \\ x \cdot 1 &= 1 \cdot x = x \\ xx^{-1} &= x^{-1}x = 1 \end{aligned}$$

Additive notation is traditionally reserved for abelian groups. Here are the abelian group axioms in additive notation:

$$\begin{aligned} x + (y + z) &= (x + y) + z \\ x + 0 &= 0 + x = x \\ x + (-x) &= (-x) + x = 0 \\ x + y &= y + x. \end{aligned}$$

1.2 Rings and fields

Definition 1.2.1. A *ring* is a 6-tuple $(A, +, \cdot, -, 0, 1)$ such that

1. $(A, +, 0, -)$ is an abelian group.
2. $(A, \cdot, 1)$ is a commutative monoid.
3. The distributive law holds for $x, y, z \in A$:

$$x \cdot (y + z) = (x \cdot y) + (x \cdot z).$$

For example, $(\mathbb{R}, +, \cdot, -, 0, 1)$ is a ring, where $+, \cdot, -, 0, 1$ have their usual meanings. Similarly, \mathbb{Z}, \mathbb{Q} , and \mathbb{C} are rings with respect to the usual operations.

Definition 1.2.2. A *field* is a ring $(K, +, \cdot, -, 0, 1)$ such that $0 \neq 1$ and any $x \neq 0$ has an inverse x^{-1} in the monoid $(K, \cdot, 1)$.

For example, the rings \mathbb{R} , \mathbb{Q} , and \mathbb{C} are fields. On the other hand, \mathbb{Z} is not, because, for example, $2 \in \mathbb{Z}$ does not have a multiplicative inverse.

Remark 1.2.3. 1. Again, the traditional definition of “ring” is slightly different, so that $(\mathbb{R}, +, \cdot)$ is a ring rather than $(\mathbb{R}, +, \cdot, -, 0, 1)$, but the two definitions are equivalent and Definition 1.2.1 is better for universal algebra.

2. What we are calling “rings” should really be called “commutative rings.” We will follow the conventions of commutative algebra and assume all rings are commutative. In noncommutative rings, $(A, +)$ is an abelian group but (A, \cdot) is a monoid, not necessarily commutative. One needs *two* distributive laws

$$\begin{aligned}x \cdot (y + z) &= (x \cdot y) + (x \cdot z) \\(x + y) \cdot z &= (x \cdot z) + (y \cdot z),\end{aligned}$$

as these are no longer equivalent without commutativity. A typical example of a noncommutative ring is the ring $M_n(\mathbb{R})$ of $n \times n$ matrices of real numbers, with $+$ and \cdot given by matrix addition and matrix multiplication.

3. In some fields of mathematics, rings are not assumed to have a multiplicative identity element $1 \in A$, so that (A, \cdot) is a semigroup rather than a monoid. In this case, what we are calling “rings” should be called “commutative unital rings.”

When working with rings, we use the usual notational abbreviations from algebra. For example, $x^2y - yz$ means $((x \cdot x) \cdot y) + (-(y \cdot z))$.

1.3 Languages, algebras, and equational classes

Each of the concepts *magma*, *semigroup*, *monoid*, *group*, *abelian group*, and *ring* is defined by a list of operations and a set of equations. More generally, an *equational class* is a class of “algebras” defined by a set of “equations”. In this section, we make these notions precise.

Before we can define equations and algebras, we need to deal with the fact that there are different kinds of algebras with different signatures. It doesn’t make sense to talk about a semigroup (G, \cdot) satisfying the distributive law

$x \cdot (y + z) = (x \cdot y) + (x \cdot z)$, because semigroups don't have an addition operation $x + y$. The concept of *languages* allows us to distinguish between these different types of algebras.

Definition 1.3.1. A (*functional*) *language* \mathcal{L} is a set of *function symbols* and a map assigning to each function symbol f an integer $n_f \in \mathbb{N}$ called the *arity* of f . An *n -ary function symbol* is a function symbol of arity n . Nullary function symbols are usually called *constant symbols*.

In Chapters 1–2, all “languages” will be functional languages. Later, we will see a more general notion of “language” (Definition 3.1.1). Languages are also called *signatures*, which is probably a better name. But at least in model theory, the term “language” is more common than “signature,” for historical reasons.

Example 1.3.2. The *language of abelian groups* has one binary function symbol $+$, one constant symbol 0 , and one unary function symbol $-$.

Definition 1.3.3. Given a language \mathcal{L} , an \mathcal{L} -*algebra* \mathcal{A} is a set A and a map assigning to each n -ary function symbol f in \mathcal{L} a corresponding n -ary function $f^{\mathcal{A}} : A^n \rightarrow A$. The set A is called the *underlying set* of \mathcal{A} , and $f^{\mathcal{A}}$ is called the *interpretation of f in \mathcal{A}* .

Usually we don't distinguish between an algebra \mathcal{A} and its underlying set A , writing both as A .

Example 1.3.4. If \mathcal{L} is the language of abelian groups, then an \mathcal{L} -algebra is essentially a 4-tuple $(A, +^A, 0^A, -^A)$ where A is a set, $+^A$ is a binary operation on A , $0^A \in A$, and $-^A$ is a unary operation on A .

Fix some infinite set $\mathcal{V} = \{x, y, z, \dots\}$ of “variable symbols.”

Definition 1.3.5. An \mathcal{L} -*term* is a string generated by the following rules:

- If x is a variable symbol, then x is a term.
- If f is an n -ary function symbol in \mathcal{L} , and t_1, \dots, t_n are \mathcal{L} -terms, then $f(t_1, \dots, t_n)$ is an \mathcal{L} -term.

Example 1.3.6. These are terms in the language of abelian groups:

$$x + (-y), \quad 0 + (x + 0), \quad z, \quad -(-(0 + x)), \quad 0.$$

When we say “let $t(x_1, \dots, x_n)$ be a term,” we mean that $t(x_1, \dots, x_n)$ is a term and the variables occurring in $t(x_1, \dots, x_n)$ are contained in $\{x_1, \dots, x_n\}$. If s_1, \dots, s_n are terms and $t(x_1, \dots, x_n)$ is a term, then $t(s_1, \dots, s_n)$ denotes the result of replacing x_i with s_i in $t(x_1, \dots, x_n)$.

A *closed term* is a term with no variables. If t is a closed term and A is an \mathcal{L} -algebra, we define the *interpretation of t in A* , written t^A , recursively as follows:

$$f(t_1, \dots, t_k)^A = f^A(t_1^A, \dots, t_k^A).$$

Example 1.3.7. If \mathcal{L} is the language of abelian groups, then the interpretation of the closed \mathcal{L} -term $(0 + (-0)) + 0$ in an \mathcal{L} -algebra $(A, +^A, 0^A, -^A)$ is the value $(0^A +^A (-^A 0^A)) +^A 0^A$.

The language $\mathcal{L}(A)$ consists of \mathcal{L} with each element of A added as a new constant symbol. We regard A as an $\mathcal{L}(A)$ -algebra by interpreting each new constant symbol as the corresponding element of A , so that $c^A = c$ for $c \in A$. If $t(x_1, \dots, x_n)$ is an \mathcal{L} -term and $a_1, \dots, a_n \in A$, then $t(a_1, \dots, a_n)$ is a closed $\mathcal{L}(A)$ -term. The *interpretation of t in A* , written t^A , is the function $t^A : A^n \rightarrow A$ defined by

$$t^A(a_1, \dots, a_n) = (t(a_1, \dots, a_n))^A.$$

Example 1.3.8. If \mathcal{L} is the language of abelian groups and $t(x, y) = (-x) + (0 + y)$, then the interpretation of t in an \mathcal{L} -algebra $(A, +^A, 0^A, -^A)$ is the function

$$t^A(a, b) = ((-a) + (0 + b))^A = (-^A a) +^A (0^A +^A b).$$

Definition 1.3.9. An \mathcal{L} -*equation* is a formal expression of the form

$$t(x_1, \dots, x_n) = s(x_1, \dots, x_n)$$

for two \mathcal{L} -terms $t(\bar{x})$ and $s(\bar{x})$. An \mathcal{L} -algebra A *satisfies* an equation $(t(\bar{x}) = s(\bar{x}))$ if for any $\bar{a} \in A^n$,

$$t^A(\bar{a}) = s^A(\bar{a}).$$

The notation $A \models \varphi$ means that A satisfies φ .

The group $(\mathbb{R}, +)$ satisfies the equation $x \cdot y = y \cdot x$, but the non-abelian group $(\text{Perm}(S), \circ)$ does not:

$$\begin{aligned} (\mathbb{R}, +) &\models x \cdot y = y \cdot x \\ (\text{Perm}(S), \circ) &\not\models x \cdot y = y \cdot x. \end{aligned}$$

Definition 1.3.10. An *equational \mathcal{L} -theory* is a set Σ of \mathcal{L} -equations. Elements of Σ are called *axioms* of Σ . If Σ is an equational \mathcal{L} -theory, and A is an \mathcal{L} -algebra, then A is a *model* of Σ , written $A \models \Sigma$, if $A \models \varphi$ for every $\varphi \in \Sigma$. The class of models of Σ is written $\text{Mod}(\Sigma)$. An *equational class* is a class of the form $\text{Mod}(\Sigma)$ for some Σ .

Example 1.3.11. Let \mathcal{L} be the language of abelian groups. The *theory of abelian groups* consists of the equations

$$\begin{aligned} x + (y + z) &= (x + y) + z \\ x + 0 &= x \\ x + (-x) &= 0 \\ x + y &= y + x. \end{aligned}$$

Models are abelian groups.

Example 1.3.12. The classes of rings, groups, monoids, semigroups, and magmas are equational classes.

Warning 1.3.13. The class of fields is *not* an equational class. We will see a couple proofs of this in the next chapter (Examples 2.1.6 and 2.2.6). Nevertheless, we can still learn useful facts about fields by applying the tools of universal algebra to rings.

Example 1.3.14. An *idempotent monoid* is a monoid (A, \star, e) in which the equation $x \star x = x$ holds. Idempotent monoids form an equational class.

Example 1.3.15. A *boolean algebra* is a 6-tuple $(B, \wedge, \vee, \neg, 0, 1)$ where $(B, \vee, 0)$ is an idempotent commutative monoid, $(B, \wedge, 1)$ is an idempotent

commutative monoid, and $\neg : B \rightarrow B$ is a unary operation such that the following equations hold:

$$\begin{aligned} x \wedge \neg x &= 0 \\ x \vee \neg x &= 1 \\ x \wedge (y \vee z) &= (x \wedge y) \vee (x \wedge z) \\ x \vee (y \wedge z) &= (x \vee y) \wedge (x \vee z) \end{aligned}$$

Boolean algebras form an equational class. If S is a set, then the powerset $\mathfrak{P}(S)$ is a boolean algebra with

$$\begin{aligned} x \wedge y &= x \cap y \\ x \vee y &= x \cup y \\ 0 &= \emptyset \\ 1 &= S \\ \neg x &= S \setminus x \end{aligned}$$

Another important boolean algebra is the set of truth values $\{\text{FALSE}, \text{TRUE}\}$, with \wedge, \vee, \neg interpreted as the basic logical operations:

$$\begin{aligned} x \wedge y &= x \text{ AND } y \\ x \vee y &= x \text{ OR } y \\ \neg x &= \text{NOT } x \\ 0 &= \text{FALSE} \\ 1 &= \text{TRUE}. \end{aligned}$$

The fact that these two examples satisfy the axioms of boolean algebras can be checked directly. We will see later in Example 2.2.14 that there is a reason why any equation satisfied by the boolean algebra of truth values must also be satisfied by the boolean algebra $\mathfrak{P}(S)$.

Warning 1.3.16. In universal algebra, equational classes are usually called *varieties*. We will avoid this terminology because “variety” tends to mean something completely different in model theory. Confusingly, “equational theory” also has a technical meaning in model theory, though it is much less common.

Note that languages are allowed to have infinitely many symbols, and theories are allowed to have infinitely many axioms. Here is an important example:

Definition 1.3.17. Let (G, \cdot) be a group and S be a set. An *action* of G on S is a map

$$\star : G \times S \rightarrow S$$

satisfying the following axioms for $g, h \in G$ and $x \in S$:

$$\begin{aligned} (g \cdot h) \star x &= g \star (h \star x) \\ 1 \star x &= x \end{aligned}$$

Group actions are usually written using \cdot rather than \star .

Example 1.3.18. Suppose S is a set and $\text{Perm}(S)$ is the group of permutations on S (Example 1.1.8). There is a natural action of $\text{Perm}(S)$ on S given by $f \cdot x = f(x)$. The axioms of group actions hold as follows:

$$\begin{aligned} (f \cdot g) \cdot x &= (f \circ g)(x) = f(g(x)) = f \cdot (g \cdot x) \\ 1 \cdot x &= \text{id}(x) = x. \end{aligned}$$

Example 1.3.19. If you know linear algebra, another example is the group of invertible $n \times n$ matrices, which acts by matrix multiplication on the space of $n \times 1$ column vectors.

Fix a specific group G .

Definition 1.3.20. A G -set is a pair (S, \cdot) where S and \cdot is an action of G on S .

This doesn't immediately fit into the framework of universal algebra, because the binary operator \cdot involves two different sets, one of which is fixed. However, the following equivalent definition works:

Definition 1.3.21. A G -set is a set S and a unary operator μ_g for each $g \in G$, satisfying the following identities:

$$\begin{aligned} \mu_{g \cdot h}(x) &= \mu_g(\mu_h(x)) \text{ for each } g, h \in G \\ \mu_1(x) &= x \end{aligned}$$

The translation between Definitions 1.3.20 and 1.3.21 is

$$\mu_g(x) = g \cdot x.$$

Definition 1.3.21 fits into the framework of universal algebra. Note that the language of G -sets has one symbol for each element of G . When G is an infinite group, we need infinitely many symbols. Likewise, the theory of G -sets needs infinitely many equations (when G is infinite).

Remark 1.3.22. If you know what vector spaces are, the same thing happens: for any fixed field K , the class of K -vector spaces is an equational class. We will review vector spaces later, in Section 13.6.

1.4 Homomorphisms and isomorphisms

Fix a language \mathcal{L} .

Definition 1.4.1. Let A, B be \mathcal{L} -algebras. A *homomorphism* from A to B is a function $\alpha : A \rightarrow B$ such that for any k -ary function symbol $f \in \mathcal{L}$, and any $a_1, \dots, a_k \in A$,

$$\alpha(f^A(a_1, \dots, a_k)) = f^B(\alpha(a_1), \alpha(a_2), \dots, \alpha(a_k)). \quad (*)$$

An *isomorphism* is a bijective homomorphism. Two \mathcal{L} -algebras A, B are *isomorphic*, written $A \cong B$, if there is an isomorphism $\alpha : A \rightarrow B$.

In what follows, we extend functions to tuples componentwise, so that $\alpha(a_1, \dots, a_k) := (\alpha(a_1), \dots, \alpha(a_k))$. Then $(*)$ can be rewritten as $\alpha(f^A(\bar{a})) = f^B(\alpha(\bar{a}))$.

Remark 1.4.2. The idea of $(*)$ is that α “preserves” the function symbol f . It may help to rewrite $(*)$ as the following logically equivalent condition:

$$f^A(a_1, \dots, a_k) = b \implies f^B(\alpha(a_1), \dots, \alpha(a_k)) = \alpha(b).$$

Example 1.4.3. Let $\mathcal{L} = \{+, 0, -\}$ be the language of abelian groups. Let $h : \mathbb{Z} \rightarrow \mathbb{Z}$ be $h(x) = 2x$. Then h is a homomorphism from \mathbb{Z} to \mathbb{Z} , because

$$\begin{aligned} 2(x + y) &= (2x) + (2y) \\ 2(0) &= 0 \\ 2(-x) &= -(2x). \end{aligned}$$

However, h is not an isomorphism, because h is not surjective.

Here is another example which is slightly more subtle.

Example 1.4.4. Let $\mathcal{L} = \{\cdot, 1\}$ be the language of monoids. The exponential map $\exp(x) = e^x$ is a homomorphism from the monoid $(\mathbb{R}, +, 0)$ to $(\mathbb{R}, \cdot, 1)$, because

$$\begin{aligned}\exp(x + y) &= \exp(x) \exp(y) \\ \exp(0) &= 1\end{aligned}$$

If this looks confusing, remember that the interpretation of “ \cdot ” in the monoid $(\mathbb{R}, +, 0)$ is $+$, and the interpretation of “ 1 ” is 0 .

The exponential map is not an isomorphism, as it is not surjective. On the other hand, if $\mathbb{R}_{>0}$ denotes the set of positive real numbers, then \exp is an isomorphism from the monoid $(\mathbb{R}, +, 0)$ to the monoid $(\mathbb{R}_{>0}, \cdot, 1)$.

Example 1.4.5. An action of G on a set S (Definition 1.3.17) can be thought of as a map assigning to each element $g \in G$ a permutation $\mu_g \in \text{Perm}(S)$ satisfying the conditions

$$\begin{aligned}\mu_{g \cdot h} &= \mu_g \circ \mu_h \\ \mu_1 &= \text{id}_S\end{aligned}$$

(See Definition 1.3.21.) These conditions precisely say that $g \mapsto \mu_g$ is a monoid homomorphism $(G, \cdot, 1) \rightarrow (\text{Perm}(S), \circ, \text{id}_S)$. We will see later that such a homomorphism must actually be a group homomorphism (Theorem 1.4.13). Therefore, an action of G on S is equivalent to a group homomorphism from G to $\text{Perm}(S)$.

The next theorem expresses some basic facts about homomorphisms and isomorphisms: we can compose homomorphisms and take inverses of isomorphisms.

Theorem 1.4.6. *Let \mathcal{L} be a language and A, B, C be \mathcal{L} -algebras.*

1. *Let $\alpha : A \rightarrow B$ and $\beta : B \rightarrow C$ be homomorphisms. Then $\beta \circ \alpha : A \rightarrow C$ is a homomorphism.*
2. *The identity map $\text{id}_A : A \rightarrow A$ is an isomorphism.*
3. *If $\alpha : A \rightarrow B$ is an isomorphism, then $\alpha^{-1} : B \rightarrow A$ is an isomorphism.*

Proof. 1. If $f \in \mathcal{L}$ is a k -ary function symbol, then

$$\beta(\alpha(f^A(\bar{a}))) = \beta(f^B(\alpha(\bar{a}))) = f^C(\beta(\alpha(\bar{a})))$$

because β and α are homomorphisms.

2. If $f \in \mathcal{L}$ is a k -ary relation symbol, then

$$\text{id}(f(\bar{a})) = f(\bar{a}) = f(\text{id}(\bar{a})).$$

3. Suppose $f \in \mathcal{L}$ is a k -ary function symbol and $b_1, \dots, b_k \in B$. Let $a_i = \alpha^{-1}(b_i) \in A$. Then $b_i = \alpha(a_i)$. As α is a homomorphism,

$$\alpha(f^A(\bar{a})) = f^B(\alpha(\bar{a})) = f^B(\bar{b}).$$

Applying α^{-1} to both sides, we see that

$$\alpha^{-1}(f^B(\bar{b})) = f^A(\bar{a}) = f^A(\alpha^{-1}(\bar{b})). \quad \square$$

Corollary 1.4.7. *The relation of isomorphism is an equivalence relation on \mathcal{L} -algebras:*

1. $A \cong A$ for any \mathcal{L} -algebra A .
2. If $A \cong B$, then $B \cong A$.
3. If $A \cong B$ and $B \cong C$, then $A \cong C$.

Proof. 1. $\text{id}_A : A \rightarrow A$ is an isomorphism.

2. If $\alpha : A \rightarrow B$ is an isomorphism, then $\alpha^{-1} : B \rightarrow A$ is an isomorphism.

3. If $\alpha : A \rightarrow B$ and $\beta : B \rightarrow C$ are isomorphisms, then $\beta \circ \alpha : A \rightarrow C$ are isomorphisms. \square

Example 1.4.8. Here is an example of an isomorphism. Let S be $\{\{x\} : x \in \mathbb{R}\}$, the set of one element subsets of \mathbb{R} . Define operations on S as follows:

$$\begin{aligned} \{x\} + \{y\} &:= \{x + y\} \\ \{x\} \cdot \{y\} &:= \{x \cdot y\} \\ -\{x\} &= \{-x\} \\ 1 &:= \{1\} \\ 0 &:= \{0\}. \end{aligned}$$

For example, $\{2\} + \{3\} \cdot \{5\} = \{17\}$. This makes $(S, +, \cdot, -, 0, 1)$ into a ring. Then there is an isomorphism from the ring \mathbb{R} to the ring S given by

$$\begin{aligned} f : \mathbb{R} &\rightarrow S \\ f(x) &= \{x\}. \end{aligned}$$

The intuition you should have is that the ring S is a poorly disguised copy of \mathbb{R} . On some level, S is “the same thing” as \mathbb{R} .

This is the intuition we have for isomorphisms in general—if two algebras A and B are isomorphic, then we think of them as being two “copies” of the same algebra. In algebra and model theory, the objects of study are not so much algebras as isomorphism classes of algebras.

This mindset has the following corollary: if two algebras A and B are isomorphic, then they should have identical properties. For example, A and B will have the same cardinality because the isomorphism $f : A \rightarrow B$ is a bijection, showing that $|A| = |B|$.

Of course, not all properties respect isomorphisms. In our example above, the property “contains 7” is true for \mathbb{R} but false for S . What we should conclude from this is that the property “contains 7” is a stupid property, or an *evil property* as they would say on the website nLab. The “meaningful” or “good” properties of algebras are the ones that respect isomorphisms. In the remainder of this section, we show that the properties defined by equations are good properties. In other words, we will show that isomorphic algebras satisfy the same equations (Theorem 1.4.11).

The next theorem shows that homomorphisms preserve not only the basic function symbols, but also all terms. This is intuitively reasonable, since terms are expressions built up from function symbols.

Theorem 1.4.9. *Let $\alpha : A \rightarrow B$ be a homomorphism of \mathcal{L} -algebras. Let $t(x_1, \dots, x_n)$ be an \mathcal{L} -term. Then for any $a_1, \dots, a_n \in A$, we have*

$$\alpha(t^A(\bar{a})) = t^B(\alpha(\bar{a})). \quad (\dagger)$$

Proof. Proceed by induction on the complexity of t .

- If $t(\bar{x}) = x_i$, then both sides of (\dagger) are $\alpha(a_i)$.
- Suppose $t(\bar{x}) = f(s_1(\bar{x}), \dots, s_k(\bar{x}))$ for some k -ary function symbol f and some simpler \mathcal{L} -terms s_1, \dots, s_k . By definition of t^A ,

$$\alpha(t^A(\bar{a})) = \alpha(f^A(s_1^A(\bar{a}), \dots, s_k^A(\bar{a}))).$$

As α is a homomorphism,

$$\alpha(f^A(s_1^A(\bar{a}), \dots, s_k^A(\bar{a}))) = f^B(\alpha(s_1^A(\bar{a})), \dots, \alpha(s_k^A(\bar{a}))).$$

By induction,

$$\alpha(s_i^A(\bar{a})) = s_i^B(\alpha(\bar{a})).$$

for each i . Therefore,

$$f^B(\alpha(s_1^A(\bar{a})), \dots, \alpha(s_k^A(\bar{a}))) = f^B(s_1^B(\alpha(\bar{a})), \dots, s_k^B(\alpha(\bar{a}))).$$

By definition, the right hand side is $t^B(\alpha(\bar{a}))$. \square

Lemma 1.4.10. *Let $\alpha : A \rightarrow B$ be a surjective homomorphism of \mathcal{L} -algebras, and let φ be an \mathcal{L} -equation. Then $A \models \varphi \implies B \models \varphi$.*

Proof. Suppose φ is $t(x_1, \dots, x_n) = s(x_1, \dots, x_n)$. Suppose $b_1, \dots, b_n \in B$. By surjectivity, we can write b_i as $\alpha(a_i)$ for some $a_i \in A$. Then

$$\begin{aligned} t^B(\bar{b}) &= t^B(\alpha(\bar{a})) = \alpha(t^A(\bar{a})) \\ s^B(\bar{b}) &= s^B(\alpha(\bar{a})) = \alpha(s^A(\bar{a})) \end{aligned}$$

by Theorem 1.4.9, as α is a homomorphism. Because $A \models \varphi$, the right hand sides are equal. Therefore the left hand sides are equal, meaning

$$t^B(\bar{b}) = s^B(\bar{b}).$$
 \square

This allows us to complete the proof that isomorphic algebras satisfy the same equations.

Theorem 1.4.11. *Suppose $A \cong B$.*

1. $A \models s = t \iff B \models s = t$.
2. $A \models \Sigma \iff B \models \Sigma$
3. $A \in \text{Mod}(\Sigma) \iff B \in \text{Mod}(\Sigma)$.
4. If \mathcal{K} is an equational class, then $A \in \mathcal{K} \iff B \in \mathcal{K}$.

Group homomorphisms

In this section, we study homomorphisms of groups, learning something useful about rings and fields along the way.

Lemma 1.4.12 (Cancellation). *Let G be a group. If $ax = ay$, then $x = y$. Similarly, if $xa = ya$, then $x = y$.*

Proof. If $ax = ay$, then

$$x = 1x = (a^{-1}a)x = a^{-1}(ax) = a^{-1}(ay) = (a^{-1}a)y = 1y = y.$$

The other case is similar. □

Theorem 1.4.13. *Let G, H be groups. If $f : G \rightarrow H$ is a semigroup homomorphism, meaning that f preserves multiplication*

$$f(xy) = f(x)f(y),$$

then f also preserves 1 and $(-)^{-1}$:

$$\begin{aligned} f(1) &= 1 \\ f(x^{-1}) &= f(x)^{-1}, \end{aligned}$$

and so f is a group homomorphism.

Proof. First note that

$$f(1) \cdot f(1) = f(1 \cdot 1) = f(1) = f(1) \cdot 1.$$

Canceling $f(1)$ from both sides, we see $f(1) = 1$. Next,

$$f(x)f(x^{-1}) = f(xx^{-1}) = f(1) = 1 = f(x)f(x)^{-1}.$$

Canceling $f(x)$ from both sides, we see $f(x^{-1}) = f(x)^{-1}$. □

Corollary 1.4.14. *Let R be a ring. Then the following equations hold, for $a, x \in R$:*

$$\begin{aligned} 0a &= 0 \\ (-x)a &= -(xa) \\ (-1)a &= -a \end{aligned}$$

Proof. Let $\mu_a(x) = ax$. Then μ_a is a semigroup homomorphism $(R, +) \rightarrow (R, +)$, by the distributive law:

$$a(x + y) = ax + ay.$$

Thus μ_a preserves zero and negation:

$$\begin{aligned}\mu_a(0) &= 0 \\ \mu_a(-x) &= -\mu_a(x),\end{aligned}$$

which are the first two equations. The third equation holds by taking $x = 1$ in the second equation. \square

Lemma 1.4.15. *The following holds in any field K :*

$$xy = xz \implies y = z \text{ when } x \neq 0.$$

Proof. Multiply both sides of $xy = xz$ by x^{-1} , as in Lemma 1.4.12. \square

Theorem 1.4.16. *If K is a field and $x, y \in K$, then*

$$xy = 0 \iff (x = 0 \text{ or } y = 0)$$

Proof. If $x = 0$ or $y = 0$, then $xy = 0$ by the zero law (Corollary 1.4.14). Conversely, suppose $xy = 0$, but $x \neq 0$ and $y \neq 0$. Then

$$xy = 0 = x0$$

by the zero law. Cancelling x from both sides, $y = 0$, a contradiction. \square

Chapter 2

New algebras from old

Let Σ be an equational theory, like the theory of rings or the theory of groups. In this chapter, we will meet three ways to build new models of Σ from old models: *subalgebras*, *products*, and *quotients*, in Sections 2.1, 2.2, and 2.4, respectively. These constructions can roughly be described as follows:

- A *subalgebra* of A is an algebra that sits inside A , the way that the ring \mathbb{Z} sits inside \mathbb{R} and \mathbb{R} sits inside \mathbb{C} .
- A *product* of two algebras A and B is the algebra where we carry out operations from A and B in parallel, a bit like vector addition in linear algebra. One can also form products of more than two algebras, even taking the product of infinitely many algebras.
- A *quotient* of an algebra A is an algebra obtained by “collapsing” A , forcing certain elements of A to become equal. For example, there is a way to collapse the ring \mathbb{Z} making all the even numbers be equal to 0 and all the odd numbers be equal to 1. This gives a ring called $\mathbb{Z}/2\mathbb{Z}$ with two elements.

Each of these constructions respects equational theories: if we start with models of Σ , the result will be a model of Σ . For example, a product of models is a model, and a quotient of a model is a model. This provides a new source of models. For example, in Section 2.5, we use quotients to construct new examples of rings, including finite fields.

At the same time, this shows the limits of what can be expressed by equational theories. For example, a product of two fields is not a field, so

the class of fields cannot be defined by an equational theory (Example 2.2.6). On the other hand, we will see in Theorem 2.9.2 that if a class of structures \mathcal{K} is closed under products, subalgebras, and quotients, then \mathcal{K} is defined by an equational theory, a fact known as *Birkhoff's HSP Theorem*. This gives a structural characterization of equational classes.

A further significance of these constructions is that they reveal the inner structure of homomorphisms. The *fundamental theorem of homomorphisms* (Theorem 2.6.7) shows that if $f : A \rightarrow B$ is a homomorphism, then the image of f , which is a subalgebra of B , is isomorphic to a quotient of A . We apply this to rings in Section 2.7, to define *characteristic*, an important invariant of rings and fields. We will see further applications of these ideas when we study algebraically closed fields in Chapter 9.

2.1 Subalgebras and generators

Definition 2.1.1. Let A be an \mathcal{L} -algebra. A *subalgebra* is a subset $B \subseteq A$ such that for any k -ary relation symbol $f \in \mathcal{L}$,

$$b_1, \dots, b_k \in B \implies f^A(b_1, \dots, b_k) \in B.$$

Example 2.1.2. \mathbb{Z} is a subalgebra of the ring $(\mathbb{R}, +, \cdot, -, 0, 1)$, because

$$x, y \in \mathbb{Z} \implies x + y \in \mathbb{Z}$$

$$x, y \in \mathbb{Z} \implies xy \in \mathbb{Z}$$

$$x \in \mathbb{Z} \implies -x \in \mathbb{Z}$$

$$0 \in \mathbb{Z}$$

$$1 \in \mathbb{Z}.$$

If B is a subalgebra of an \mathcal{L} -algebra A , then we can make B into an \mathcal{L} -algebra by defining f^B to be the restriction of f^A to B , for each function symbol f :

$$f^B(b_1, \dots, b_k) := f^A(b_1, \dots, b_k) \in B.$$

In this way, we regard subalgebras as algebras, not just sets.

Theorem 2.1.3. Suppose B is a subalgebra of A .

1. The inclusion $B \rightarrow A$ is a homomorphism.

2. If $A \models s = t$, then $B \models s = t$.

3. If $\mathcal{K} = \text{Mod}(\Sigma)$ is an equational class, then $A \in \mathcal{K} \implies B \in \mathcal{K}$.

Proof. (1) holds by choice of the structure on B . For (2), note that

$$\begin{aligned} t^B(\bar{b}) &= t^A(\bar{b}) \\ s^B(\bar{b}) &= s^A(\bar{b}) \end{aligned}$$

for \bar{b} in B by Theorem 1.4.9 and (1). If $t^A = s^A$, then $t^B = s^B$. Finally, (3) follows directly from (2). \square

Example 2.1.4. A subalgebra of a monoid is a monoid, and a subalgebra of a group is a group.

Remark 2.1.5. Note that this *wouldn't work* if we had used the traditional definitions of monoids and groups (see Remark 1.1.3). For example, if we use the traditional definition then the algebra $(\mathbb{R}, +)$ is a group, but the subalgebra $(\mathbb{N}, +)$ is not, and the subalgebra $(\{1, 2, 3, \dots\}, +)$ is not even a monoid. Including the identity element 0 and the inverse map $-x$ as part of the structure makes things work.

Because a subalgebra of a *foo* is usually a *foo*, we often say *subfoo* instead of subalgebra. For example, we say “subgroup” when working with groups, and “subring” when working with rings.

Example 2.1.6. Theorem 2.1.3 can be used to show that certain classes are not equational classes. For example, the ring \mathbb{R} is a field, but the subring \mathbb{Z} is not. Therefore, fields are not an equational class. Of course, maybe this problem could be fixed the same way we fixed monoids and groups, by adding the division operation \div as part of the structure. In Example 2.2.6, we will see that this doesn't work, and there is no sensible way to make fields into an equational class.

Definition 2.1.7. If S is a subset of an algebra A , then $\langle S \rangle$ or $\langle S \rangle_A$ denotes the set

$$\{t^A(\bar{b}) : t(x_1, \dots, x_n) \text{ is an } \mathcal{L}\text{-term and } \bar{b} \in S^n\} \quad (*)$$

We often omit brackets $\{, \}$ inside \langle, \rangle , using abbreviations like

$$\begin{aligned} \langle a_1, \dots, a_n \rangle &= \langle \{a_1, \dots, a_n\} \rangle \\ \langle A, b \rangle &= \langle A \cup \{b\} \rangle. \end{aligned}$$

Remark 2.1.8. $\langle S \rangle$ is the smallest subalgebra of A containing S . To see this, first note that by taking $t(x) = x$, we get $b = t(b) \in \langle S \rangle$ for any $b \in S$, and so $S \subseteq \langle S \rangle$. If A' is a subalgebra of A containing S , then $\langle S \rangle \subseteq A'$ because if $t(\bar{x})$ is a term and $\bar{b} \in S^n$, then $t^A(\bar{b}) = t^{A'}(\bar{b}) \in A'$. The final step is to show that $\langle S \rangle$ is itself a subalgebra. This is a little confusing to formally prove; here is one approach. For $\bar{b} \in S^n$, let

$$A_{\bar{b}} = \{t^A(\bar{b}) : t(x_1, \dots, x_n) \text{ is a term}\} \subseteq \langle \bar{b} \rangle.$$

It is easy to see that $A_{\bar{b}}$ is a subalgebra containing b_1, \dots, b_n . Then $\langle \bar{b} \rangle \subseteq A_{\bar{b}}$, equality holds, and $\langle \bar{b} \rangle$ equals the subalgebra $A_{\bar{b}}$. This shows that $\langle S \rangle$ is a subalgebra when S is finite.

For the case where S is infinite, take a k -ary function symbol f and elements $a_1, \dots, a_k \in \langle S \rangle$. Each a_i is in $\langle S_i \rangle$ for some finite $S_i \subseteq_f S$. Let $S' = \bigcup_{i=1}^n S_i$. Then $a_1, \dots, a_k \in \langle S' \rangle$, and $\langle S' \rangle$ is a subalgebra (by the finite case), and so $f^A(a_1, \dots, a_k) \in \langle S' \rangle \subseteq \langle S \rangle$.

Definition 2.1.9. The subalgebra $\langle S \rangle$ is called the subalgebra *generated* by S . We say that A is *finitely generated* if $A = \langle S \rangle$ for some finite $S \subseteq A$.

Example 2.1.10. The subring of \mathbb{R} generated by $\{\sqrt{2}, \sqrt{3}\}$ contains all the expressions built up from $\sqrt{2}$ and $\sqrt{3}$ using the ring operations, such as

$$\sqrt{2} + \sqrt{3}, \quad -\sqrt{2}, 1 + \sqrt{3}, 1 + 1, \sqrt{3} \cdot (\sqrt{2} + 1).$$

In fact, one can show that

$$\langle \sqrt{2}, \sqrt{3} \rangle = \{w + x\sqrt{2} + y\sqrt{3} + z\sqrt{6} : w, x, y, z \in \mathbb{Z}\}. \quad (\dagger)$$

To prove this, let A be the right hand side of (\dagger) . One proves the following:

1. A is a subring of \mathbb{R}
2. Any subalgebra of \mathbb{R} containing $\sqrt{2}$ and $\sqrt{3}$ must contain A .

Both of these are straightforward algebraic exercises.

Example 2.1.11. The subgroup of $(\mathbb{R}, +, 0, -)$ generated by π is $\{n\pi : n \in \mathbb{Z}\}$.

Example 2.1.12. The subring of \mathbb{R} generated by \emptyset is the smallest subring of \mathbb{R} , which is \mathbb{Z} . Indeed, \mathbb{Z} is a subring of \mathbb{R} , and if A is any other subring of \mathbb{R} , one can show that

$$n \in \mathbb{Z} \implies \pm n \in A$$

by induction on n .

Equivalently, $\mathbb{Z} = \{t^{\mathbb{R}} : t \text{ is a closed term}\}$. That is \mathbb{Z} is the set of real numbers built up from the ring operations with no parameters, like so:

There are some close links between generators and homomorphisms:

Theorem 2.1.13. *Let A be generated by $S \subseteq A$. Suppose two homomorphisms $\alpha_1, \alpha_2 : A \rightarrow B$ have the same restriction to S . Then $\alpha_1 = \alpha_2$.*

Proof. Any element of A has the form $t(\bar{a})$ for some term $t(x_1, \dots, x_n)$ and some $\bar{a} \in S^n$. As homomorphisms preserve terms (Theorem 1.4.9),

$$\alpha_1(t(\bar{a})) = t(\alpha_1(\bar{a})) = t(\alpha_2(\bar{a})) = \alpha_2(t(\bar{a})). \quad \square$$

Theorem 2.1.14. *Let A be generated by $S \subseteq A$. Let $\alpha : A \rightarrow B$ be a homomorphism. Then the image $\alpha(A)$ is $\langle \alpha(S) \rangle_B$.*

Proof.

$$\begin{aligned} \alpha(A) &= \alpha(\langle S \rangle) = \{\alpha(t^A(\bar{b})) : t \text{ is a term and } \bar{b} \in S^n\} \\ &= \{t^B(\alpha(\bar{b})) : t \text{ is a term and } \bar{b} \in S^n\} \\ &= \{t^B(\bar{b}) : t \text{ is a term and } \bar{b} \in \alpha(S)^n\} \\ &= \langle \alpha(S) \rangle_B. \end{aligned} \quad \square$$

2.2 Products

Binary products

Recall that if A and B are sets, the *Cartesian* or *direct product* is the set

$$A \times B = \{(x, y) : x \in A \text{ and } y \in B\}.$$

The terminology “product” reflects the fact that $|A \times B| = |A| \cdot |B|$. For example, if A has 3 elements and B has 5 elements, then $A \times B$ has 15 elements, the product of 3 and 5.

Definition 2.2.1. Let A, B be two \mathcal{L} -algebras. The *product algebra* $A \times B$ is the \mathcal{L} -algebra with underlying set $A \times B$ and

$$f^{A \times B}((a_1, b_1), \dots, (a_k, b_k)) := (f^A(a_1, \dots, a_k), f^B(b_1, \dots, b_k)).$$

for each k -ary function symbol $f \in \mathcal{L}$.

The products in Definition 2.2.1 are called *binary products* because there are 2 factors A and B . Later we will consider k -ary products for other values of k , including infinite k .

Example 2.2.2. The direct product of the rings \mathbb{R} and \mathbb{Z} is the structure $(\mathbb{R} \times \mathbb{Z}, +, \cdot, -, 0, 1)$, where the operations are defined as follows:

$$\begin{aligned} (x_1, y_1) + (x_2, y_2) &:= (x_1 + x_2, y_1 + y_2) \\ (x_1, y_1) \cdot (x_2, y_2) &:= (x_1 x_2, y_1 y_2) \\ -(x, y) &:= (-x, -y) \\ 1 &:= (1, 1) \\ 0 &:= (0, 0). \end{aligned}$$

In other words, all the operations are componentwise.¹

Any equational class is closed under binary products. For example, the product of two rings is a ring. This will take a little work to show.

Remark 2.2.3. Let A_1, A_2 be two \mathcal{L} -algebras. For $i = 1, 2$, let $\pi_i : A_1 \times A_2 \rightarrow A_i$ be the projection map $\pi_i(x_1, x_2) = x_i$. Then each π_i is a homomorphism. For example,

$$\begin{aligned} \pi_1(f^{A_1 \times A_2}((a_1, b_1), \dots, (a_k, b_k))) &= f^{A_1}(a_1, \dots, a_k) \\ &= f^{A_1}(\pi_1(a_1, b_1), \dots, \pi_1(a_k, b_k)). \end{aligned}$$

Theorem 2.2.4. Let A_1, A_2 be \mathcal{L} -algebras.

1. If φ is an equation and $A_i \models \varphi$ for $i = 1, 2$, then $A_1 \times A_2 \models \varphi$.
2. If \mathcal{K} is an equational class and $A_1, A_2 \in \mathcal{K}$, then $A_1 \times A_2 \in \mathcal{K}$.

¹The operation $+$ is essentially vector addition. In contrast, \cdot is not a natural operation on vectors, except in settings like machine learning and NumPy.

Proof. Part (2) follows formally from (1). For (1), suppose $A_i \models t = s$ for $i = 1, 2$. Let $P = A_1 \times A_2$. We claim $P \models t = s$. Otherwise there are $a_1, \dots, a_k \in P$ such that $t^P(\bar{a}) \neq s^P(\bar{a})$. Then there is $i \in \{1, 2\}$ such that

$$\pi_i(t^P(\bar{a})) \neq \pi_i(s^P(\bar{a})).$$

As π_i is a homomorphism, we can rewrite the two sides as follows:

$$t^{A_i}(\pi_i(\bar{a})) \neq s^{A_i}(\pi_i(\bar{a})).$$

Then $A_i \not\models t = s$, a contradiction. \square

Example 2.2.5. The product of two groups is a group. The product of two rings is a ring.

Again, Theorem 2.2.4 can be used to see that certain classes are not varieties.

Example 2.2.6. The ring \mathbb{R} is a field, but $\mathbb{R} \times \mathbb{R}$ is not, because the zero law (Theorem 1.4.16) fails:

$$\begin{aligned} (0, 1) &\neq (0, 0) \\ (1, 0) &\neq (0, 0) \\ (0, 1) \cdot (1, 0) &= (0, 0). \end{aligned}$$

Therefore fields are not an equational class. Unlike the argument in Example 2.1.6, this argument is resilient under adding new symbols to the language, like how we made groups into an equational class by adding a symbol for the inverse map.

One could still try to salvage fields as an equational class by using a completely different set of operations. This doesn't work either. We will see later that if n is finite, then there is a field of size n if and only if n is a prime power. In particular, there are fields of size 2 and 3, but no field of size 6. Regardless of which operations we use, products cannot possibly work.

Perhaps we need to add some extra elements to fields, such as the element ∞ . This *still* doesn't work. Suppose we had some way of representing a field with n elements as an algebra with $n + 1$ elements. The field of size 2 would give an algebra of size 3. Taking a product of this algebra with itself

three times, we would get an algebra of size $3^3 = 27$, which would need to correspond to a field of size 26. But there is no field of size 26.²

Infinite products

Let I be a set.

Definition 2.2.7. An I -tuple is a function with domain I . If a is an I -tuple, we write $a(i)$ as $\pi_i(a)$. The notation $(a_i : i \in I)$ means the function $i \mapsto a_i$.

For example, $(2n : n \in \omega)$ is the ω -tuple corresponding to the function $f(n) = 2n$. Although I -tuples are officially functions, we think of them as distinct kinds of object, and use notation to hide the identification.

When $I = \{1, 2, \dots, n\}$, we identify n -tuples and I -tuples, so that

$$(a_1, a_2, \dots, a_n) = (a_i : i \in \{1, \dots, n\}).$$

Likewise, we think of ω -tuples as tuples of length ω , i.e., sequences, so that

$$(a_1, a_2, a_3, \dots) = (a_i : i \in \omega).$$

Definition 2.2.8. Let $\{A_i\}_{i \in I}$ be a family of sets. The *direct product* $\prod_{i \in I} A_i$ is the set of I -tuples $(a_i : i \in I)$ such that $a_i \in A_i$ for each $i \in I$.

For example, when $I = \{1, 2\}$,

$$\prod_{i \in \{1, 2\}} A_i = \{(a_1, a_2) : a_1 \in A_1, a_2 \in A_2\} = A_1 \times A_2.$$

Similarly,

$$\prod_{i \in \{1, 2, \dots, n\}} A_i = A_1 \times A_2 \times \dots \times A_n.$$

Consequently, $\prod_{i \in I} A_i$ generalizes binary direct products $A_1 \times A_2$.

Remark 2.2.9. Suppose A_i doesn't depend on i , so that $A_i = A$ for some fixed set A as i varies. Then $\prod_{i \in I} A$ is the set of functions from I to A . This set is also written A^I , and is called a *power* of A .

²Likewise, supposed we tried adding *two* elements, for ∞ and \mathbf{NaN} . The field of size 2 would give an algebra of size 4. Taking the product of this algebra with itself three times we would get an algebra of size $4^3 = 64$, corresponding to a non-existent field of size 62.

Definition 2.2.10. Let I be a set and let A_i be an \mathcal{L} -algebra for each $i \in I$. We make $\prod_{i \in I} A_i$ into an \mathcal{L} -algebra by interpreting each k -ary function symbol f as

$$f(a_1, \dots, a_k) = (f^{A_i}(\pi_i(a_1), \dots, \pi_i(a_k)) : i \in I).$$

For example, if each A_i is a ring, then addition is defined by

$$a + b := (\pi_i(a) + \pi_i(b) : i \in I)$$

or equivalently

$$(a_i : i \in I) + (b_i : i \in I) = (a_i + b_i : i \in I).$$

To be even more specific, when $I = \omega$ this says

$$\begin{aligned} & (a_1, a_2, a_3, \dots) \\ & + (b_1, b_2, b_3, \dots) \\ & = (a_1 + b_1, a_2 + b_2, a_3 + b_3, \dots) \end{aligned}$$

As in the case of binary products, the operations are carried out componentwise. In fact Definition 2.2.10 generalizes Definition 2.2.1, which is the special case when $I = \{1, 2\}$.

Remark 2.2.11. The structure on $\prod_{i \in I} A_i$ is chosen to make π_i a homomorphism:

$$\pi_i(f(a_1, \dots, a_k)) = f^{A_i}(\pi_i(a_1), \dots, \pi_i(a_k)).$$

Theorem 2.2.12. Let A_i be an \mathcal{L} -algebra for each $i \in I$.

1. If φ is an equation and $A_i \models \varphi$ for each $i \in I$, then $\prod_{i \in I} A_i \models \varphi$.
2. If \mathcal{K} is an equational class and $A_i \in \mathcal{K}$ for all $i \in I$, then $\prod_{i \in I} A_i \in \mathcal{K}$.

Proof. The proof of Theorem 2.2.4 works here. □

Example 2.2.13 (Powers). If I is a set and A is an \mathcal{L} -algebra, the power $A^I := \prod_{i \in I} A$ is the set of functions $I \rightarrow A$, with all the operations defined

pointwise. For example, the ring $\mathbb{R}^{\mathbb{R}}$ is the set of functions $f : \mathbb{R} \rightarrow \mathbb{R}$, with ring operations defined like so:

$$\begin{aligned}(f + g)(x) &:= f(x) + g(x) \\ (f \cdot g)(x) &:= f(x)g(x) \\ (-f)(x) &:= -f(x) \\ 0(x) &:= 0 \\ 1(x) &:= 1\end{aligned}$$

We can get more examples of rings by taking certain subrings of $\mathbb{R}^{\mathbb{R}}$. For example, the set of differential functions (in calculus) is a subring. One subring of interest to us is the ring $\mathbb{R}[x]$ of *polynomials*, functions of the form

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$$

for constants $a_0, \dots, a_n \in \mathbb{R}$. We will say more about the ring $\mathbb{R}[x]$ later.

Example 2.2.14. Recall the boolean algebras $\{\text{FALSE}, \text{TRUE}\}$ and $\mathfrak{P}(S)$ from Example 1.3.15. It turns out that

$$\mathfrak{P}(S) \cong \{\text{FALSE}, \text{TRUE}\}^S. \quad (*)$$

The right hand side $\{\text{FALSE}, \text{TRUE}\}^S$ is the set of truth-valued functions on S , with operations defined pointwise. For example,

$$(f \wedge g)(x) = f(x) \wedge g(x) = (f(x) \text{ AND } g(x)).$$

We can identify a truth-valued function $f : S \rightarrow \{\text{FALSE}, \text{TRUE}\}$ with the set $\{x \in S : f(x)\}$, and then the pointwise AND corresponds to intersections:

$$\{x \in S : f(x) \text{ and } g(x)\} = \{x \in S : f(x)\} \cap \{x \in S : g(x)\}.$$

Similarly, pointwise OR corresponds to unions, and so on, giving the isomorphism of (*).

Applying Theorem 2.2.12 to (*), we see that any equation satisfied by the algebra $\{\text{FALSE}, \text{TRUE}\}$ must also be satisfied by $\mathfrak{P}(S)$. For example, the logical identity

$$x \text{ and } (y \text{ or } z) \iff (x \text{ and } y) \text{ or } (x \text{ and } z)$$

corresponds to the set-theoretic fact

$$X \cap (Y \cup Z) = (X \cap Y) \cup (X \cap Z).$$

If I recall correctly, the observation that truth values and sets are governed by the same equations goes back to Boole, which is why boolean algebras are named in his honor.

2.3 Descending functions along surjections

Often in mathematics we want to define a function $f : A \rightarrow B$, where $A = \{\pi(x) : x \in A'\}$ for some function $\pi : A' \rightarrow A$. Since every element of A has the form $\pi(x)$, to specify f we only need to specify $f(\pi(x))$ for $x \in A'$. The following theorem gives the precise criterion that $f(\pi(x))$ needs to satisfy in order for $f(-)$ to be well-defined.

Theorem 2.3.1. *Let $\pi : A' \rightarrow A$ be a surjection. Let $f' : A' \rightarrow B$ be a function such that*

$$\pi(x) = \pi(y) \implies f'(x) = f'(y). \quad (*)$$

Then there is a unique function $f : A \rightarrow B$ such that $f(\pi(x)) = f'(x)$, i.e., the following diagram commutes:

$$\begin{array}{ccc} A' & & \\ \pi \downarrow & \searrow f' & \\ A & \xrightarrow{f} & B. \end{array}$$

Proof. For $z \in A$, let $S_z = \{f'(x) : x \in A', \pi(x) = z\}$. There is at least one $x \in A'$ with $\pi(x) = z$ because π is surjective, so S_z is non-empty. If $x_1, x_2 \in A'$ satisfy $\pi(x_i) = z$ for $i = 1, 2$, then $f(x_1) = f(x_2)$ by (*). Therefore S_z has a unique element. Let $f(z)$ be the unique element of S_z .

If $x \in A'$ and $z = \pi(x)$, then $f'(x) \in S_z = \{f(z)\}$, so $f'(x) = f(z) = f(\pi(x))$. This proves existence of f . For uniqueness, suppose $f'' : A \rightarrow B$ is another function satisfying $f''(\pi(x)) = f'(x)$. For any $z \in A$, there is $x \in A'$ with $\pi(x) = z$, and then

$$f''(z) = f''(\pi(x)) = f'(x) = f(\pi(x)) = f(z),$$

so $f'' = f$. □

Example 2.3.2. Let \mathcal{K} be the class of \mathcal{L} -algebras. If $X, Y \in \mathcal{K}$ are isomorphic, then $|X| = |Y|$. Therefore, there is a map from the class \mathcal{K}/\cong of isomorphism classes to the class Card of cardinals sending the isomorphism class of X to the cardinality $|X|$. Ignoring the difference between sets and classes, this is an instance of Theorem 2.3.1:

$$\begin{array}{ccc} \mathcal{K} & & \\ \downarrow & \searrow & \\ \mathcal{K}/\cong & \dashrightarrow & \text{Card}. \end{array}$$

The surjectivity of $\mathcal{K} \rightarrow \mathcal{K}/\cong$ holds because every isomorphism class is the isomorphism class of some algebra. The condition $(*)$ in Theorem 2.3.1 holds because

$$[X]_{\cong} = [Y]_{\cong} \iff X \cong Y \implies |X| = |Y|.$$

The conclusion of Theorem 2.3.1 says that the map $f : \mathcal{K}/\cong \rightarrow \text{Card}$ satisfies

$$f([X]_{\cong}) = |X|,$$

so that f sends the isomorphism class of X to the cardinality of X .

2.4 Congruences and quotients

Congruences

Recall that equivalence relations on A are subsets of $A \times A$.

Definition 2.4.1. Let A be an \mathcal{L} -algebra. A *congruence* on A is an equivalence relation E on A that is also a subalgebra of $A \times A$.

Unwinding the definition, an equivalence relation \sim on A is a congruence iff

$$(a_1 \sim b_1 \text{ and } a_2 \sim b_2 \text{ and } \dots \text{ and } a_k \sim b_k) \implies f(a_1, \dots, a_k) \sim f(b_1, \dots, b_k)$$

for any k -ary function symbol $f \in \mathcal{L}$. For example, an equivalence relation \sim on a group $(G, \cdot, 1, (-)^{-1})$ is a congruence if

$$\begin{aligned} (a_1 \sim b_1 \text{ and } a_2 \sim b_2) &\implies a_1 a_2 \sim b_1 b_2 \\ a \sim b &\implies a^{-1} \sim b^{-1} \\ 1 &\sim 1. \end{aligned}$$

The next lemma gives a slightly different criterion for being a congruence, which is sometimes easier to check.

Theorem 2.4.2. *Let \sim be an equivalence relation on an \mathcal{L} -algebra A . Then \sim is a congruence iff the following holds: for any k -ary function symbol $f \in \mathcal{L}$ and any $1 \leq i \leq k$,*

$$a_i \sim a'_i \implies f(a_1, \dots, a_k) \sim f(a_1, \dots, a_{i-1}, a'_i, a_{i+1}, \dots, a_k). \quad (*)$$

Proof. First suppose \sim is a congruence, and $a_i \sim a'_i$. For $j \neq i$ define $a'_j := a_j$. Then $a_j \sim a'_j$ because \sim is reflexive. As $a_j \sim a'_j$ for all $j \leq k$, we see that

$$f(a_1, \dots, a_k) \sim f(a'_1, \dots, a'_k) = f(a_1, \dots, a_{i-1}, a'_i, a_{i+1}, \dots, a_k).$$

Conversely, suppose $(*)$ holds. Suppose $a_i \sim b_i$ for $i = 1, \dots, k$. Then $(*)$ gives

$$\begin{aligned} & f(a_1, a_2, a_3, \dots, a_k) \\ & \sim f(b_1, a_2, a_3, \dots, a_k) \\ & \sim f(b_1, b_2, a_3, \dots, a_k) \\ & \sim \dots \\ & \sim f(b_1, b_2, \dots, b_k). \end{aligned} \quad \square$$

For example, a relation \sim on a group G is a congruence iff

$$\begin{aligned} a \sim a' & \implies ab \sim a'b \\ b \sim b' & \implies ab \sim ab' \\ a \sim b & \implies a^{-1} \sim b^{-1}. \end{aligned}$$

Congruences are a little abstract, but can be understood in terms of more concrete objects in the case of rings and groups.

Definition 2.4.3. Let R be a ring. An *ideal* is a subset $I \subseteq R$ such that

1. $0 \in I$.
2. $x, y \in I \implies x + y \in I$.
3. $(x \in R \text{ and } y \in I) \implies xy \in I$.

Remark 2.4.4. The even numbers $2\mathbb{Z}$ are an ideal in the ring \mathbb{Z} . More generally, if R is any ring and $a \in R$, then the set $aR := \{ax : x \in R\}$ is an ideal. Such ideals are called *principal ideals*.

Theorem 2.4.5. *Let R be a ring.*

1. *If I is an ideal, define $x \equiv_I y$ to mean $x - y \in I$. Then \equiv_I is a congruence.*
2. *This gives a bijection between congruences on R and ideals on R .*

Proof. First, note that if \sim is a congruence, then

$$x \sim y \iff x - y \sim 0$$

for any $x, y \in R$. Indeed,

$$\begin{aligned} x \sim y &\implies x - y \sim y - y = 0 \\ x - y \sim 0 &\implies x = (x - y) + y \sim 0 + y = y. \end{aligned}$$

Therefore \sim must have the form

$$x \sim y \iff x - y \in I$$

for some set $I \subseteq R$, namely $I = \{z \in R : z \sim 0\}$. It remains to characterize which sets I yield congruences.

1. Reflexivity says that $x - x \in I$ for any x . This holds iff $0 \in I$.
2. Symmetry says that $x - y \in I \iff y - x \in I$. This holds iff I is closed under negation.
3. Transitivity says that if $x - y \in I$ and $y - z \in I$, then $x - z \in I$. This holds iff I is closed under addition.
4. Compatibility with $+$ says that if $x - y \in I$, then $(x + a) - (y + a) \in I$. This condition holds for any I .
5. Compatibility with \cdot says that if $x - y \in I$, then $(ax) - (ay) \in I$. This condition holds iff I is closed under multiplication by R .

In summary, I yields a congruence if and only if the following four properties hold:

- $0 \in I$
- $x \in I \implies -x \in I$
- $x, y \in I \implies x + y \in I$
- $a \in R, x \in I \implies ax \in I.$

The second condition is an instance of the fourth (take $a = -1$), so it can be removed. The remaining three conditions are the definition of “ideal.” \square

Remark 2.4.6. The relation $x \equiv_I y$ is called “congruence modulo I ”, and is usually written like $x \equiv y \pmod{I}$. When I is a principal ideal aR , it is usually written $x \equiv y \pmod{a}$.

Example 2.4.7. In the ring \mathbb{Z} , $x \equiv y \pmod{2}$ holds iff $x - y$ is even. There are two equivalence classes, the even numbers and odd numbers.

Definition 2.4.8. Let G be a group. A *normal subgroup* is a subgroup $N \subseteq G$ such that

$$(x \in N \text{ and } y \in G) \implies yxy^{-1} \in N. \quad (\dagger)$$

When G is abelian, $yxy^{-1} = xyy^{-1} = x$, so (\dagger) says

$$(x \in N \text{ and } y \in G) \implies x \in N,$$

which is trivial. Therefore for abelian groups, a normal subgroup is the same thing as a subgroup. But for non-abelian groups, the two concepts are usually different.

Lemma 2.4.9. *Let G be a group.*

1. *If N is a normal subgroup, define $x \equiv_N y$ to mean $xy^{-1} \in N$. Then \equiv_N is a congruence.*
2. *This gives a bijection between congruences on G and normal subgroups of G .*

Proof. Similar to Lemma 2.4.5. Compatibility with $(-)^{-1}$ can be ignored thanks to the following:

Claim. If \approx is a monoid congruence (on $(G, \cdot, 1)$), then \approx is a group congruence.

Indeed, if $x \approx y$ then

$$x^{-1} = x^{-1}yy^{-1} \approx x^{-1}xy^{-1} = y^{-1}. \quad \square$$

Remark 2.4.10. 1. The relation $x \equiv_N y$ is called “congruence modulo N ”, and is usually written like $x \equiv y \pmod{N}$.

2. In group theory, the kernel of a homomorphism $f : G \rightarrow H$ is the *subgroup* $\{x \in G : f(x) = 1\}$. This is the normal subgroup corresponding to the kernel in the sense of Definition 2.6.2.

Quotients

If X is a set and E is an equivalence relation on X , let X/E denote the quotient, the set $\{[a]_E : a \in X\}$, where $[a]_E$ is the E -equivalence class $[a]_E = \{b \in X : a E b\}$. Recall that $[a]_E = [b]_E \iff a E b$. We omit the subscript when E is clear from context.

Theorem 2.4.11 (Quotients). *Let A be an \mathcal{L} -algebra and E be a congruence on A . Then there is a unique \mathcal{L} -algebra with underlying set A/E such that $A \rightarrow A/E$ is a homomorphism, meaning that*

$$f^{A/E}([a_1], \dots, [a_k]) = [f^A(a_1, \dots, a_k)] \quad (\dagger)$$

for any k -ary function symbol in \mathcal{L} .

Proof. The equation (\dagger) is essentially an instance of Theorem 2.3.1:

$$\begin{array}{ccc} A^k & & \\ \downarrow & \searrow & \\ (A/E)^k & \dashrightarrow & A/E, \end{array}$$

where the vertical arrow is $(x_1, \dots, x_k) \mapsto ([x_1], \dots, [x_k])$, the diagonal arrow is $\bar{x} \mapsto [f^A(\bar{x})]$, and the dashed arrow is the map $f^{A/E} : (A/E)^k \rightarrow A/E$ we are trying to define.

By Theorem 2.3.1, the function $f^{A/E} : (A/E)^k \rightarrow A/E$ is uniquely determined by (\dagger) , as long as

$$([a_1], \dots, [a_k]) = ([b_1], \dots, [b_k]) \implies [f^A(\bar{a})] = [f^A(\bar{b})],$$

or equivalently,

$$(a_i E b_i \text{ for } i = 1, \dots, k) \implies (f^A(\bar{a}) E f^A(\bar{b})).$$

This holds because E is a congruence. □

Definition 2.4.12. If A is an \mathcal{L} -algebra and E is a congruence, then A/E is called the *quotient algebra* of A by E .

Remark 2.4.13. In group theory and ring theory, if A is a group or ring, and B is a normal subgroup or ideal, then A/B means A/\equiv_B where \equiv_B is the congruence associated with B . In these cases, one can show that $|A| = |A/B| \cdot |B|$. In particular, when A is finite, one can show that $|A/B| = |A|/|B|$, which explains the term “quotient.”

Example 2.4.14. Let A be $\mathbb{Z}/2\mathbb{Z}$, that is, the quotient of the ring \mathbb{Z} by the equivalence relation $x \equiv_2 y \iff x - y \in 2\mathbb{Z}$. There are two equivalence classes, EVEN and ODD, with $[x] = \text{EVEN}$ for $x \in 2\mathbb{Z}$, and $[x] = \text{ODD}$ for $x \notin 2\mathbb{Z}$. Addition on A is defined so that

$$[x] +^A [y] = [x + y].$$

If x and y are even, then $x + y$ is even, so $\text{EVEN} + \text{EVEN} = \text{EVEN}$. Similarly, we can complete the tables of addition and multiplication as follows:

+	EVEN	ODD
EVEN	EVEN	ODD
ODD	ODD	EVEN

·	EVEN	ODD
EVEN	EVEN	EVEN
ODD	EVEN	ODD

Note that this isn't just a ring—it's a field.

Like subalgebras and products, quotients preserve equations:

Theorem 2.4.15. *Let A be an \mathcal{L} -algebra and E be a congruence on A .*

1. *If A satisfies an equation φ , then A/E satisfies φ .*

2. If $\mathcal{K} = \text{Mod}(\Sigma)$ is an equational class, then $A \in \mathcal{K} \implies A/E \in \mathcal{K}$.

Proof. By construction, there is a surjective homomorphism $A \rightarrow A/E$. Then $A \models \varphi \implies A/E \models \varphi$ (Lemma 1.4.10). \square

If E is an equivalence relation on A , a *set of representatives* for E is a set $S \subseteq A$ containing exactly one element from each E -equivalence class. The next theorem gives a more concrete way to understand quotient structures.

Theorem 2.4.16. *Let A be an \mathcal{L} -algebra and E be a congruence. Let S be a set of representatives for E . Let $\rho : A \rightarrow S$ be the map sending $x \in A$ to the unique $y \in S \cap [x]_E$. For each k -ary function symbol $f \in \mathcal{L}$, define*

$$f^S(x_1, \dots, x_k) = \rho(f^A(x_1, \dots, x_k)).$$

This makes S into an \mathcal{L} -algebra isomorphic to A/E .

Proof. The construction certainly makes S into an \mathcal{L} -algebra. Define $\alpha : S \rightarrow A/E$ by $\alpha(x) = [x]$. We claim that α is an isomorphism.

1. α is a homomorphism: if f is a k -ary function symbol, then

$$\alpha(f^S(x_1, \dots, x_k)) = [\rho(f^A(x_1, \dots, x_k))]$$

But $\rho(y) E y$ for any $y \in A$, so $[\rho(y)] = [y]$. Therefore

$$[\rho(f^A(x_1, \dots, x_k))] = [f^A(x_1, \dots, x_k)].$$

By construction of the quotient,

$$[f^A(x_1, \dots, x_k)] = f^{A/E}([x_1], \dots, [x_k]) = f^{A/E}(\alpha(x_1), \dots, \alpha(x_k)).$$

Putting everything together,

$$\alpha(f^S(x_1, \dots, x_k)) = f^{A/E}(\alpha(x_1), \dots, \alpha(x_k)).$$

2. α is a bijection: clear by choice of S . \square

Example 2.4.17. In the ring \mathbb{Z} , $\{0, 1\}$ is a set of representatives for \equiv_2 , and so the quotient ring \mathbb{Z}/\equiv_2 is isomorphic to the ring

+	0	1
0	0	1
1	1	0

·	0	1
0	0	0
1	0	1

2.5 Application: finite fields

The rings we have seen so far are subrings of \mathbb{C} , like \mathbb{Z} , \mathbb{Q} , \mathbb{R} , and \mathbb{C} . All of them are infinite. In this section, we use the machinery of quotients to construct examples of finite rings and fields. These finite rings and fields play important roles in number theory, cryptography, and error correcting codes.

Work in the ring of integers \mathbb{Z} . Recall the relation $x \equiv_n y$ or $x \equiv y \pmod{n}$, meaning that $x - y \in n\mathbb{Z}$. This relation is called *congruence modulo n* .

Lemma 2.5.1. *If $n > 0$, then $\{0, 1, \dots, n-1\}$ is a system of representatives for \equiv_n . That is, for every $x \in \mathbb{Z}$ there is a unique $y \in \{0, \dots, n-1\}$ with $x \equiv y \pmod{n}$.*

Proof. An exercise, by induction on x . □

For example, every integer x is congruent modulo 10 to exactly one value y in the set $\{0, 1, 2, \dots, 9\}$. If x is positive, then y is the final digit in the base 10 representation of x . In general, if x and n are positive integers, then the unique $y \in \{0, 1, \dots, n-1\}$ such that $x \equiv y \pmod{n}$ is the remainder of dividing x by n . In computer science, this operation is often written $x \bmod n$. We will encounter it later in Section 3.9.

Lemma 2.5.2. *If $0 \neq n \in \mathbb{Z}$, and $x \in \mathbb{Z}$, there is $y \in \mathbb{Z}$ with*

$$\begin{aligned} x &\equiv y \pmod{n} \\ |y| &< |n|. \end{aligned}$$

Proof. Since $n\mathbb{Z} = (-n)\mathbb{Z}$, we may assume $n > 0$ by replacing n with $-n$ if necessary. Then there is some $y \in \{0, 1, 2, \dots, n-1\}$ with $x \equiv y \pmod{n}$ by Lemma 2.5.1. □

Recall that a *principal ideal* in a ring R is an ideal of the form $aR = \{ax : x \in R\}$.

Theorem 2.5.3. *Every ideal $I \subseteq \mathbb{Z}$ is a principal ideal $I = n\mathbb{Z}$ for some $n \geq 0$.*

Proof. Note that $\{0\} \subseteq I$. If $I = \{0\}$, take $n = 0$. Otherwise, take $n \in I \setminus \{0\}$ minimizing $|n|$. Multiplying n by -1 if necessary, we may assume $n \geq 0$.

Then $n\mathbb{Z} \subseteq I$ because I is an ideal. We claim $n\mathbb{Z} = I$. Otherwise, take $a \in I \setminus n\mathbb{Z}$. By Lemma 2.5.2 there is $b \in \mathbb{Z}$ with

$$\begin{aligned} b &\equiv a \pmod{n} \\ |b| &< |n|. \end{aligned}$$

Then $b - a \in n\mathbb{Z} \subseteq I$, and $a \in I$, so $b \in I$ because I is closed under addition. If $b \neq 0$ then b contradicts the choice of n . If $b = 0$, then $a \equiv b = 0 \pmod{n}$, so $a \in n\mathbb{Z}$, contradicting the choice of a . \square

Remark 2.5.4. If $n > 0$, then $\mathbb{Z}/n\mathbb{Z}$ is isomorphic to $R = (\{0, 1, \dots, n-1\}, +^R, \cdot^R)$, where $x +^R y$ is the unique $z \in R$ with $x + y \equiv z \pmod{n}$, and $x \cdot^R y$ is the unique $w \in R$ with $xy \equiv w \pmod{n}$. This follows by Theorem 2.4.16 applied to the system of representatives in Lemma 2.5.1. In particular, $\mathbb{Z}/n\mathbb{Z}$ is finite, of size n .

Using the notation from computer science,

$$\begin{aligned} x +^R y &= (x + y) \bmod n \\ x \cdot^R y &= (xy) \bmod n. \end{aligned}$$

For example, here is the table of multiplication and addition for $\mathbb{Z}/5\mathbb{Z}$:

+	0	1	2	3	4
0	0	1	2	3	4
1	1	2	3	4	0
2	2	3	4	0	1
3	3	4	0	1	2
4	4	0	1	2	3

+	0	1	2	3	4
0	0	0	0	0	0
1	0	1	2	3	4
2	0	2	4	1	3
3	0	3	1	4	2
4	0	4	3	2	1

Note that this is not just a ring but a field ($1^{-1} = 1$, $2^{-1} = 3$, $3^{-1} = 2$, $4^{-1} = 4$). We will see shortly that $\mathbb{Z}/n\mathbb{Z}$ is a field whenever n is a prime number. We first prove a couple general facts about rings.

Lemma 2.5.5. *Let I, J be ideals in a ring R . Let $I + J := \{x + y : x \in I, y \in J\}$. Then $I + J$ is an ideal containing I and J .*

Proof. To show that $I + J$ is an ideal, there are three things to check:

1. $0 \in I + J$: take $x = 0 \in I$ and $y = 0 \in J$.

2. $I + J$ is closed under addition: if $x, x' \in I$ and $y, y' \in J$, then $(x + y) + (x' + y') = (x + x') + (y + y') \in I + J$.
3. $I + J$ is closed under multiplication by R : if $x \in I, y \in J$, and $a \in R$, then $a(x + y) = (ax) + (ay) \in I + J$.

Finally, if $x \in I$ then $x + 0 \in I + J$ because $0 \in J$. This shows $I \subseteq I + J$, and $J \subseteq I + J$ follows similarly. \square

An ideal $I \subseteq R$ is *proper* if $1 \notin I$. If $1 \in I$, then $x = x1 \in I$ for every $x \in R$, and so $I = R$. Thus there is exactly one improper ideal, namely R itself. A *maximal ideal* is a maximal proper ideal. Using Zorn's lemma, one can show that every proper ideal is contained in a maximal proper ideal, though we will not use this fact.

Theorem 2.5.6. *If I is a maximal ideal, then R/I is a field.*

Proof. Let $[a] \in R/I$ denote the image of $a \in R$. First we show that $1 \neq 0$ in R/I . The fact that $1 \notin I$ means that $1 \not\equiv 0 \pmod{I}$, so $[1] \neq [0]$. Next we show that any $[a] \neq 0$ has a multiplicative inverse. By Lemma 2.5.5, $aR + I$ is an ideal containing aR and I . The fact that $[a] \neq 0$ means that $a = a - 0 \notin I$. Therefore $aR + I \supsetneq I$, as $a \in aR \subseteq aR + I$. By maximality of I , $aR + I$ is improper, so $1 \in aR + I$. Therefore there are $x \in R$ and $y \in I$ with $1 = ax + y$. Then $ax - 1 = y \in I$, so that $ax \equiv 1 \pmod{I}$ and $[a][x] = [ax] = [1]$. Then $[x]$ is the multiplicative inverse of $[a]$. \square

Theorem 2.5.7. *If p is prime, then $\mathbb{Z}/p\mathbb{Z}$ is a field.*

Proof. It suffices to show that the ideal $p\mathbb{Z}$ is maximal. If not, take a larger proper ideal $I \supsetneq p\mathbb{Z}$. By Theorem 2.5.3, $I = n\mathbb{Z}$ for some $n \in \mathbb{Z}$. Then $p \in p\mathbb{Z} \subseteq n\mathbb{Z}$, so p is a multiple of n . In other words, $p = nm$ for some $m \in \mathbb{Z}$. As p is prime, one of n or m is ± 1 .

- If $n = \pm 1$, then $n\mathbb{Z}$ is improper, a contradiction.
- If $m = \pm 1$, then $n = \pm p$, and $n\mathbb{Z} = p\mathbb{Z}$, a contradiction. \square

On the other hand, if n is a composite number ab , then $\mathbb{Z}/n\mathbb{Z}$ is not a field. To see this, note that a and b are not multiples of n , but $ab = n$ is a multiple of n . Thus

$$\begin{aligned} a &\not\equiv 0 \pmod{n} \\ b &\not\equiv 0 \pmod{n} \\ ab &\equiv 0 \pmod{n}. \end{aligned}$$

In the quotient $\mathbb{Z}/n\mathbb{Z}$, it follows that $[a] \neq 0$ and $[b] \neq 0$, but $[a][b] = [ab] = 0$, contradicting Theorem 1.4.16 if $\mathbb{Z}/n\mathbb{Z}$ is a field.

As for $n = 1$, the quotient $\mathbb{Z}/n\mathbb{Z}$ is the *trivial ring* with only one element, which fails to be a field because of the requirement that $0 \neq 1$ in fields. We will see later in Lemma 2.7.8 that $\mathbb{Z}/0\mathbb{Z}$ is isomorphic to \mathbb{Z} , so it is not a field. In summary, the only quotients of \mathbb{Z} that are fields are $\mathbb{Z}/p\mathbb{Z}$ for prime p .

2.6 The fundamental theorem on homomorphisms

In this section, we prove the *fundamental theorem of homomorphisms* (Theorem 2.6.7), which analyzes the structure of homomorphisms in terms of quotients and subalgebras. This theorem shows that any homomorphism is built out of three components:

1. A homomorphism of the form $A \rightarrow A/E$ where A is an algebra and A/E is a quotient.
2. An isomorphism.
3. An inclusion homomorphism $A \rightarrow B$ where A is a subalgebra of B .

We will see some applications of this theorem in Section 2.7 and much later in Chapter 9.

Theorem 2.6.1 (Images). *Let $\alpha : A \rightarrow B$ be a homomorphism of \mathcal{L} -algebras. Then the image $\text{im}(\alpha) = \alpha(A) = \{\alpha(x) : x \in A\}$ is a subalgebra of B .*

Proof. Suppose $f \in \mathcal{L}$ is a k -ary function symbol, and $b_1, \dots, b_k \in \text{im}(\alpha)$. Each b_i can be written as $\alpha(a_i)$ for some $a_i \in A$. Then

$$f(b_1, \dots, b_k) = f(\alpha(a_1), \dots, \alpha(a_k)) = \alpha(f(a_1, \dots, a_k)) \in \text{im}(\alpha). \quad \square$$

Conversely, if A is a subalgebra of B , then A is the image of a homomorphism, namely the inclusion $A \hookrightarrow B$ which is a homomorphism by Theorem 2.1.3. Thus the subalgebras of B are precisely those sets which are images of homomorphisms into B . Next, we do something similar with congruences.

Definition 2.6.2. Let $\alpha : A \rightarrow B$ be a homomorphism of \mathcal{L} -algebras. The *kernel* of α is the equivalence relation

$$a \sim b \iff \alpha(a) = \alpha(b).$$

We write the kernel as $\ker(\alpha)$.

Theorem 2.6.3. If $\alpha : A \rightarrow B$ is a homomorphism of \mathcal{L} -algebras, then the kernel is a congruence on A .

Proof. Let $E = \ker(\alpha)$. Let $f \in \mathcal{L}$ be a k -ary function symbol. If $a_i E b_i$ for $i = 1, \dots, k$, then $\alpha(a_i) = \alpha(b_i)$ for each i . As α is a homomorphism,

$$\alpha(f(\bar{a})) = f(\alpha(\bar{a})) = f(\alpha(\bar{b})) = \alpha(f(\bar{b})).$$

Therefore $f(\bar{a}) E f(\bar{b})$. □

Conversely, every congruence E on A is the kernel of some homomorphism out of A , namely the quotient homomorphism $A \rightarrow A/E$. Thus the congruences on A are precisely those relations which are kernels of homomorphisms out of A .

Remark 2.6.4. In ring theory, the kernel of a homomorphism $f : R \rightarrow S$ is the *ideal* $\{x \in R : f(x) = 0\}$. This is the ideal corresponding to the kernel in the sense of Definition 2.6.2.

Likewise, in group theory the kernel of a homomorphism $f : G \rightarrow H$ is $\{x \in G : f(x) = 1\}$, which is the normal subgroup corresponding to the kernel congruence.

Lemma 2.6.5. Let A, B, C be algebras. Let $\alpha : A \rightarrow B$ be a surjective homomorphism and $\beta : B \rightarrow C$ be a function such that $\beta \circ \alpha : A \rightarrow C$ is a homomorphism. Then β is a homomorphism.

Proof. Let f be a k -ary function symbol. If $\bar{b} \in B^k$, then $\bar{b} = \alpha(\bar{a})$ for some $\bar{a} \in A^k$. Then

$$\beta(f(\bar{b})) = \beta(f(\alpha(\bar{a}))) = \beta(\alpha(f(\bar{a}))) = f(\beta(\alpha(\bar{a}))) = f(\beta(\bar{b}))$$

because α and $\beta \circ \alpha$ are homomorphisms. Therefore β is a homomorphism. □

Theorem 2.6.6 (Universal property of quotients). *Let A be an algebra and E be a congruence on A . If $\alpha : A \rightarrow B$ is a homomorphism and $E \subseteq \ker(\alpha)$, then there is a unique homomorphism $\beta : A/E \rightarrow B$ such that $\alpha(x) = \beta([x])$, or equivalently, the following diagram commutes:*

$$\begin{array}{ccc} A & & \\ \downarrow & \searrow \alpha & \\ A/E & \xrightarrow{\beta} & B \end{array}$$

Proof. The condition $E \subseteq \ker(\alpha)$ means that

$$[x]_E = [y]_E \implies \alpha(x) = \alpha(y).$$

By Theorem 2.3.1 there is a unique function $\beta : A/E \rightarrow B$ such that $\alpha(x) = \beta([x])$. We need β to be a homomorphism. The composition $A \rightarrow A/E \rightarrow B$ is the homomorphism α , and $A \rightarrow A/E$ is a surjective homomorphism, so $\beta : A/E \rightarrow B$ is a homomorphism by Lemma 2.6.5. \square

Theorem 2.6.7 (Fundamental theorem on homomorphisms). *Let $\alpha : A \rightarrow B$ be a homomorphism of \mathcal{L} -algebras. Let E be the kernel (a congruence on A) and let $\text{im}(\alpha)$ be the image (a subalgebra of B). There is an isomorphism $\beta : A/E \rightarrow \text{im}(\alpha)$, and α is the composition of the following three homomorphisms:*

$$A \rightarrow A/E \xrightarrow{\beta} \text{im}(\alpha) \xrightarrow{\subseteq} B$$

Proof. We can regard $\alpha : A \rightarrow B$ as a surjective homomorphism $\alpha' : A \rightarrow \text{im}(\alpha)$. Then $\ker(\alpha') = \ker(\alpha) \supseteq E$, so Theorem 2.6.6 gives a homomorphism $\beta : A/E \rightarrow \text{im}(\alpha)$ making the diagram commute:

$$\begin{array}{ccc} A & \xrightarrow{\alpha} & B \\ & \searrow \alpha' & \nearrow \\ & A/E \xrightarrow{\beta} \text{im}(\alpha) & \end{array}$$

Then β is an isomorphism:

- β is surjective because any element of $\text{im}(\alpha)$ has the form $\alpha(a) = \beta([a])$ for some $a \in A$.

- β is injective because for $a, b \in A$,

$$\beta([a]) = \beta([b]) \iff \alpha(a) = \alpha(b) \iff a E b \iff [a] = [b]. \quad \square$$

In the case of surjective homomorphisms, Theorem 2.6.7 says the following:

Corollary 2.6.8. *If $\alpha : A \rightarrow B$ is a surjective homomorphism, then there is an isomorphism $A/\ker(\alpha) \rightarrow B$.*

This gives some abstract intuition or motivation for quotients: the quotient A/E is the unique algebra, up to isomorphism, such that there is a surjective homomorphism $\alpha : A \rightarrow A/E$ with kernel E .

2.7 Application: characteristic of fields

Using the fundamental theorem on homomorphisms, we can define the *characteristic* of a ring or field. This is an important invariant, especially for fields, as we will see later in Chapter 9 and Theorem 14.2.10.

If R is a ring and $n \in \mathbb{Z}$, let n^R be the interpretation of n in R , that is,

$$n^R = \begin{cases} \underbrace{(1 + \cdots + 1)}_{n \text{ times}}^R & \text{if } n > 0 \\ 0^R & \text{if } n = 0 \\ \underbrace{(-(1 + \cdots + 1))}_{n \text{ times}}^R & \text{if } n < 0 \end{cases}$$

For example, $n^{\mathbb{Z}} = n$.

Lemma 2.7.1. *The map $\alpha : n \mapsto n^R$ is the unique homomorphism from \mathbb{Z} to R .*

Proof. The fact that α is a homomorphism is an exercise in induction and the ring axioms. If $\beta : \mathbb{Z} \rightarrow R$ is another homomorphism, then $\beta(n) = \alpha(n)$ by Theorem 1.4.9. For example,

$$\beta(3) = \beta(1 + 1 + 1) = (1 + 1 + 1)^R = \alpha(3).$$

Alternatively, uniqueness follows because a homomorphism is determined by what it does on a set of generators (Theorem 2.1.13), and \mathbb{Z} is generated by the empty set. \square

The subring of R generated by the empty set, written $\langle \emptyset \rangle_R$, can be described abstractly as the smallest subring of R , or explicitly as the set of things of the form t^R for t a closed term.

Lemma 2.7.2. *The image $\text{im}(\alpha)$ equals $\langle \emptyset \rangle_R$.*

This follows from Theorem 2.1.14, but we give another proof.

Proof. The image $\text{im}(\alpha)$ is a subring, so $\text{im}(\alpha) \supseteq \langle \emptyset \rangle_R$. On the other hand, $\alpha(n) = n^R$ is clearly t^R for some closed term t , so $\alpha(n) \in \langle \emptyset \rangle_R$. Thus $\text{im}(\alpha) \subseteq \langle \emptyset \rangle_R$. \square

We summarize the situation below:

Theorem 2.7.3. *If R is a ring, then there is a unique homomorphism $\alpha : \mathbb{Z} \rightarrow R$ given by $\alpha(n) = n^R$, and the image $\text{im}(\alpha)$ is the minimal subring $\langle \emptyset \rangle_R$.*

Definition 2.7.4. The *characteristic* of R , written $\text{char}(R)$, is the unique $n \in \mathbb{N}$ such that the kernel of $\mathbb{Z} \rightarrow R$ is the principal ideal $n\mathbb{Z}$.

Here, we are using the kernel ideal $\ker(\alpha) = \{x : \alpha(x) = 0\}$, rather than the kernel congruence $\ker(\alpha) = \{(x, y) : \alpha(x) = \alpha(y)\}$; see Remark 2.6.4. The fact that the kernel is a principal ideal comes from Theorem 2.5.3.

Theorem 2.7.5. *If R has characteristic n , then the minimal subring $\langle \emptyset \rangle_R$ is isomorphic to $\mathbb{Z}/n\mathbb{Z}$.*

Proof. By Theorem 2.7.3, $\langle \emptyset \rangle_R$ is the image of the unique homomorphism $\alpha : \mathbb{Z} \rightarrow R$. By the fundamental theorem on homomorphisms, $\langle \emptyset \rangle_R = \text{im}(\alpha) \cong \mathbb{Z}/\ker(\alpha) = \mathbb{Z}/n\mathbb{Z}$, where $n = \text{char}(R)$. \square

Theorem 2.7.6. *If K is a field, then $\text{char}(K) \in \{0, 2, 3, 5, 7, 11, \dots\}$.*

Proof. If $n = \text{char}(K)$, and $\alpha : \mathbb{Z} \rightarrow R$ is the unique homomorphism, then $\ker(\alpha) = n\mathbb{Z}$. We must rule out the following cases:

- $n = 1$. Then $1 \in 1\mathbb{Z} = \ker(\alpha)$, so $1^K = \alpha(1) = 0^K$, and K is not a field.

- n is a composite number ab , for some integers $a, b > 1$. Then $a, b \notin n\mathbb{Z}$ and $n \in n\mathbb{Z}$, so $a, b \notin \ker(\alpha)$ but $ab = n \in \ker(\alpha)$. This means that

$$\begin{aligned}\alpha(a) &\neq 0 \\ \alpha(b) &\neq 0 \\ \alpha(a)\alpha(b) &= \alpha(ab) = 0,\end{aligned}$$

contradicting the zero law (Theorem 1.4.16). \square

Conversely, each of the values $0, 2, 3, 5, 7, \dots$ occurs as the characteristic of some field:

Theorem 2.7.7. *If $p \in \{0, 2, 3, 5, 7, \dots\}$, then there is a field of characteristic p .*

Proof. The field \mathbb{R} has characteristic 0. If $p > 0$, the kernel of $\mathbb{Z} \rightarrow \mathbb{Z}/p\mathbb{Z}$ is $p\mathbb{Z}$, so the field $\mathbb{Z}/p\mathbb{Z}$ has characteristic p . \square

Lastly, let's spell out what the minimal subring of a field is in terms of the characteristic.

Lemma 2.7.8. *Let R be a ring. Then $R/0R \cong R$.*

Proof. By the fundamental theorem on homomorphisms (Corollary 2.6.8) applied to the surjective homomorphism $\text{id}_R : R \rightarrow R$, we have $R \cong R/I$, where $I = \ker(\text{id}_R) = \{x \in R : \text{id}(x) = 0\} = \{0\} = 0R$. \square

(We saw an instance of Lemma 2.7.8 earlier—the ring S in Example 1.4.8 was exactly $\mathbb{R}/0\mathbb{R}$.)

Theorem 2.7.9. *If K is a field, then the minimal subring $\langle \emptyset \rangle_K$ is isomorphic to \mathbb{Z} if $\text{char}(K) = 0$, and $\mathbb{Z}/p\mathbb{Z}$ if $\text{char}(K) = p > 0$.*

Theorem 2.7.10. *Let $L \supseteq K$ be an extension of fields. Then $\text{char}(L) = \text{char}(K)$.*

Proof. Let $\alpha : \mathbb{Z} \rightarrow K$ be the unique homomorphism from \mathbb{Z} to K . Then α is also the unique homomorphism from \mathbb{Z} to L . The kernel of α is the same whether we regard α as a homomorphism to K or to L . \square

2.8 Term algebras and free algebras

Let Σ be an equational theory. We have seen a number of ways to construct new models of Σ , such as taking subalgebras, products, and quotients of existing models. Another important source of models is *free algebras*, which are in some sense the “most general” or “generic” models of Σ . Free algebras are built out of the more primitive notion of *term algebras*.

Let \bar{x} be a tuple of variables, possibly infinite. Let $T(\bar{x})$ be the set of terms in the variables \bar{x} . We make $T(\bar{x})$ into an algebra by setting

$$f^{T(\bar{x})}(t_1, \dots, t_k) = f(t_1, \dots, t_k).$$

The resulting algebra is called the *term algebra*.

Example 2.8.1. In the language of rings, the term algebra $T(x, y, z)$ has elements like

$$x + (y + z), 0, 0 \cdot y, (-y) + y.$$

If we “add” the first two of these elements, we get the element $(x + (y + z)) + 0$. Note that the term algebra is not itself a ring. For example, $x + (y + z)$ and $(x + y) + z$ are distinct elements of $T(x, y, z)$, because they are not equal as terms (i.e., as strings of symbols). Thus $T(x, y, z)$ fails to satisfy the associative law.

Theorem 2.8.2. *If A is an algebra and \bar{a} is a tuple in A of the same length as \bar{x} , then the evaluation map*

$$\begin{aligned} \eta : T(\bar{x}) &\rightarrow A \\ \eta(t(\bar{x})) &= t^A(\bar{a}) \end{aligned}$$

is a homomorphism $T(\bar{x}) \rightarrow A$.

Proof. Suppose f is a k -ary function symbol, and $t_1(\bar{x}), \dots, t_k(\bar{x}) \in T(\bar{x})$. Then

$$\begin{aligned} \eta(f^{T(\bar{x})}(t_1(\bar{x}), \dots, t_k(\bar{x}))) &= \eta(f(t_1(\bar{x}), \dots, t_k(\bar{x}))) \\ &= f^A(t_1^A(\bar{a}), \dots, t_k^A(\bar{a})) = f^A(\eta(t_1(\bar{x})), \dots, \eta(t_k(\bar{x}))). \end{aligned} \quad \square$$

Fix an equational class \mathcal{K} . Let $\text{Eq}(\mathcal{K})$ denote the set of equations φ such that every $A \in \mathcal{K}$ satisfies φ .

Definition 2.8.3. If t, s are two terms, then $t \equiv_{\mathcal{K}} s$ if the equation $t = s$ holds for all $A \in \mathcal{K}$. In other words, $t \equiv_{\mathcal{K}} s$ means that $(t = s) \in \text{Eq}(\mathcal{K})$.

Example 2.8.4. For example, if \mathcal{K} is the class of rings, then $0 \cdot x \equiv_{\mathcal{K}} 0$ because the equation $0x = 0$ holds in any ring (Corollary 1.4.14).

Theorem 2.8.5. If \bar{x} is a tuple of variables, then $\equiv_{\mathcal{K}}$ is a congruence on the term algebra $T(\bar{x})$.

Proof. It is easy to see that $\equiv_{\mathcal{K}}$ is an equivalence relation. We show that $\equiv_{\mathcal{K}}$ is a congruence. Suppose t_1, \dots, t_k and s_1, \dots, s_k are terms in $T(\bar{x})$ with $t_i \equiv_{\mathcal{K}} s_i$ for each i , and f is a k -ary function symbol. Then for any $A \in \mathcal{K}$ and tuple \bar{a} in A ,

$$t_i^A(\bar{a}) = s_i^A(\bar{a}) \text{ for } i = 1, \dots, k,$$

and so

$$f(t_1(\bar{a}), \dots, t_k(\bar{a}))^A = f(s_1(\bar{a}), \dots, s_k(\bar{a}))^A.$$

It follows that

$$f(t_1(\bar{x}), \dots, t_k(\bar{x})) \equiv_{\mathcal{K}} f(s_1(\bar{x}), \dots, s_k(\bar{x})). \quad \square$$

Definition 2.8.6. The (\mathcal{K}) -free algebra on the variables \bar{x} , written $F_{\mathcal{K}}(\bar{x})$ or $F(\bar{x})$, is the quotient $T(\bar{x})/\equiv_{\mathcal{K}}$.

Theorem 2.8.7. $F_{\mathcal{K}}(\bar{x}) \in \mathcal{K}$.

Proof. Let $t(y_1, \dots, y_k) = s(y_1, \dots, y_k)$ be any of the axioms defining \mathcal{K} . For any terms $u_1, \dots, u_k \in T(\bar{x})$, any $A \in \mathcal{K}$, and any \bar{a} in A , we have

$$t(u_1(\bar{a}), \dots, u_k(\bar{a})) = s(u_1(\bar{a}), \dots, u_k(\bar{a}))$$

because $A \models t = s$. Therefore

$$\begin{aligned} t(u_1(\bar{x}), \dots, u_k(\bar{x})) &\equiv_{\mathcal{K}} s(u_1(\bar{x}), \dots, u_k(\bar{x})) \\ [t(u_1(\bar{x}), \dots, u_k(\bar{x}))] &= [s(u_1(\bar{x}), \dots, u_k(\bar{x}))] \text{ in } F(\bar{x}) \\ t([u_1(\bar{x})], \dots, [u_k(\bar{x})]) &= s([u_1(\bar{x})], \dots, [u_k(\bar{x})]) \text{ in } F(\bar{x}) \end{aligned}$$

Thus $F(\bar{x}) \models t = s$. \square

Example 2.8.8. Suppose \mathcal{K} is the class of monoids. The free monoid $F(x, y, z)$ has elements like $(xy)z$, $x(yz)$, and xy . The first two are equal, because $(xy)z \equiv_{\mathcal{K}} x(yz)$. The third element is not equal to the first two, because the equation $(xy)z = xy$ fails in at least one monoid, such as $(\mathbb{R}, \cdot, 1)$.

Because $F(x, y, z)$ is a monoid, we can forget the parentheses, and write each element (other than 1) as a string of x , y , and z , like

$$xyz, xyzzy, yyyxyyy, z, \dots$$

If $\{x, y, z\}^*$ denotes the set of finite strings in the alphabet $\{x, y, z\}$, then one can make $\{x, y, z\}^*$ into a monoid by letting $w \cdot v$ be the concatenation of w and v , and 1 be the empty string. One can show that $\{x, y, z\}^* \cong F(x, y, z)$.

Example 2.8.9. Suppose \mathcal{K} is the class of rings. The free ring $F(x)$ has elements like $x + 0$, $x^2 + x$, and so on. One can show that each element of $F(x)$ can be written in a unique way as a polynomial

$$a_0 + a_1x + a_2x^2 + \dots + a_nx^n$$

with integer coefficients a_0, \dots, a_n . Consequently, $F(x)$ is isomorphic to the ring $\mathbb{Z}[x]$ of polynomials with integer coefficients. We will say more about rings of polynomials in Section 9.1.

More generally, the free ring $F(x_1, \dots, x_n)$ is isomorphic to the ring $\mathbb{Z}[x_1, \dots, x_n]$ of polynomials in the variables x_1, \dots, x_n with coefficients in \mathbb{Z} . Going in the other direction, the free ring $F()$ turns out to be isomorphic to \mathbb{Z} .

Fact 2.8.10. *If \mathcal{K} is the class of boolean algebras (Example 1.3.15), then the free boolean algebra $F(x_1, \dots, x_n)$ is isomorphic to the set of all n -ary boolean functions*

$$\{\text{FALSE}, \text{TRUE}\}^n \rightarrow \{\text{FALSE}, \text{TRUE}\}$$

with the operations \wedge, \vee, \dots defined pointwise, like

$$(f \wedge g)(\bar{x}) = f(\bar{x}) \wedge g(\bar{x}).$$

In other words, $F(x_1, \dots, x_n)$ is isomorphic to the power $\{\text{FALSE}, \text{TRUE}\}^I$ where $I = \{\text{FALSE}, \text{TRUE}\}^n$. (See examples 2.2.13 and 2.2.14 for powers.)

The next theorem is an analogue of Theorem 2.8.2 for free algebras instead of term algebras.

Theorem 2.8.11 (Universal mapping property). *For any $A \in \mathcal{K}$ and tuple \bar{a} in A (of the same length as \bar{x}), there is a homomorphism $\alpha : F(\bar{x}) \rightarrow A$ sending $[t(\bar{x})]$ to $t(\bar{a})$.*

Proof. Let $\beta : T(\bar{x}) \rightarrow A$ be the evaluation map $t(\bar{x}) \mapsto t(\bar{a})$. Note that

$$t \equiv_{\mathcal{K}} s \implies t(\bar{a}) = s(\bar{a}) \iff \beta(t) = \beta(s).$$

Therefore $(\equiv_{\mathcal{K}}) \subseteq \ker(\beta)$. By the universal property of quotients (Theorem 2.6.6), there is a homomorphism $\alpha : F(\bar{x}) \rightarrow A$ making the diagram commute

$$\begin{array}{ccc} T(\bar{x}) & & \\ \downarrow & \searrow \beta & \\ F(\bar{x}) & \xrightarrow{\alpha} & A. \end{array}$$

Thus $\alpha([t(\bar{x})]) = \beta(t(\bar{x})) = t(\bar{a})$.

□

Example 2.8.12. For example, if we are working with rings, then $F(x)$ is the polynomial ring $\mathbb{Z}[x]$, and Theorem 2.8.11 says that for any ring A and element $a \in A$, the evaluation map

$$\begin{aligned} \mathbb{Z}[x] &\rightarrow A \\ P(x) &\mapsto P(a) \end{aligned}$$

is a homomorphism.

Similarly, there is a homomorphism from $F()$ to A sending t to t^A . In fact, $F()$ is \mathbb{Z} and this is merely the homomorphism $\mathbb{Z} \rightarrow A$ of Theorem 2.7.3.

Corollary 2.8.13. *If $A \in \mathcal{K}$, then there is a surjective homomorphism $\alpha : F(\bar{x}) \rightarrow A$ for some \bar{x} .*

Proof. Let $\bar{a} = (a_i : i \in I)$ be a tuple (probably infinite) enumerating all of A . Let $\bar{x} = (x_i : i \in I)$ be a tuple of variables of the same length. Let $\alpha : F(\bar{x}) \rightarrow A$ be the homomorphism sending $[t(\bar{x})]$ to $t^A(\bar{a})$. Letting $t(\bar{x}) = x_i$, we see that $\alpha([x_i]) = a_i$, so α is surjective. □

In light of Corollary 2.6.8, this means that any algebra in \mathcal{K} is a quotient of a free algebra, up to isomorphism. For example, any ring is a quotient of a free ring. This will play an important role in the next section.

2.9 Birkhoff's HSP theorem

If \mathcal{K} is an equational class, and $A \in \mathcal{K}$, then any subalgebra or quotient of A is in \mathcal{K} , and anything isomorphic to A is in \mathcal{K} . Moreover, any product of algebras in \mathcal{K} is in \mathcal{K} . (See Theorems 2.1.3, 2.4.15, 1.4.11, and 2.2.12.) In Examples 2.1.6 and 2.2.6 we used these facts to see that the class of fields is *not* an equational class. Birkhoff's HSP theorem—the goal of this section—says that this is the *only* obstruction to being an equational class: if a class of algebras \mathcal{K} respects subalgebras, quotients, products, and isomorphisms, then \mathcal{K} is an equational class, axiomatized by some set of equations.

Lemma 2.9.1. *Let $\alpha_i : A \rightarrow B_i$ be a homomorphism of \mathcal{L} -algebras for $i \in I$. Let $\alpha : A \rightarrow \prod_{i \in I} B_i$ be the map $\alpha(a) = (\alpha_i(a) : i \in I)$. Then α is a homomorphism.*

Proof. Let π_j be the j th coordinate projection from $\prod_{i \in I} B_i$ to B_j . If f is a k -ary function symbol and $a_1, \dots, a_k \in A$, we must show

$$\alpha(f(\bar{a})) \stackrel{?}{=} f(\alpha(\bar{a})).$$

Both sides are in $\prod_{i \in I} B_i$. If the two sides disagree, then there is $i \in I$ such that

$$\pi_i(\alpha(f(\bar{a}))) \neq \pi_i(f(\alpha(\bar{a}))).$$

As π_i is a homomorphism (Remark 2.2.11), we can change the right hand side:

$$\pi_i(\alpha(f(\bar{a}))) \neq f(\pi_i(\alpha(\bar{a}))).$$

Now $\pi_i \circ \alpha = \alpha_i$ by definition of α , so

$$\alpha_i(f(\bar{a})) \neq f(\alpha_i(\bar{a})).$$

This contradicts the fact that α_i is a homomorphism. □

Let \mathcal{K} be a class of \mathcal{L} -algebras.

1. \mathcal{K} is *closed under isomorphisms* if for any $A \in \mathcal{K}$ and $B \cong A$, $B \in \mathcal{K}$.
2. \mathcal{K} is *closed under subalgebras* if for any $A \in \mathcal{K}$ and subalgebra $B \subseteq A$, $B \in \mathcal{K}$.
3. \mathcal{K} is *closed under products* if for any family $\{A_i\}_{i \in I}$ with $A_i \in \mathcal{K}$, the product $\prod_{i \in I} A_i$ is in \mathcal{K} .

4. \mathcal{K} is closed under quotients if for any $A \in \mathcal{K}$ and congruence E on A , $A/E \in \mathcal{K}$.

Theorem 2.9.2 (Birkhoff's HSP theorem). *Let \mathcal{K} be a class of algebras. Then \mathcal{K} is an equational class if and only if \mathcal{K} is closed under isomorphisms, subalgebras, products, and quotients.*

Proof. Equational classes are closed under isomorphisms (Theorem 1.4.11), subalgebras (Theorem 2.1.3), products (Theorem 2.2.12), and quotients (Theorem 2.4.15).

Conversely, suppose \mathcal{K} is closed under isomorphisms, subalgebras, products, and quotients. Let Σ be $\text{Eq}(\mathcal{K})$, the set of equations holding on \mathcal{K} . If $A \in \mathcal{K}$, then $A \models \Sigma$, and so $\mathcal{K} \subseteq \text{Mod}(\Sigma) =: \bar{\mathcal{K}}$.

Claim. If \bar{x} is a tuple of variables, then the $\bar{\mathcal{K}}$ -free algebra $F_{\bar{\mathcal{K}}}(\bar{x})$ is in \mathcal{K} .

Proof. Let $\{(a_i, b_i)\}_{i \in I}$ enumerate all the pairs of distinct elements of $F = F_{\bar{\mathcal{K}}}(\bar{x})$. For each i , we can write $a_i = [t_i(\bar{x})]$ and $b_i = [s_i(\bar{x})]$ for terms $t_i, s_i \in T(\bar{x})$. The fact that $a_i \neq b_i$ means that $t_i(\bar{x}) \not\equiv_{\bar{\mathcal{K}}} s_i(\bar{x})$. Thus $(t_i = s_i) \notin \Sigma = \text{Eq}(\mathcal{K})$, so there is $A_i \in \mathcal{K}$ with $A_i \not\models t_i = s_i$. Then there is $\bar{a}_i \in A_i$ with $t_i^{A_i}(\bar{a}_i) \neq s_i^{A_i}(\bar{a}_i)$. As $A_i \in \mathcal{K} \subseteq \bar{\mathcal{K}}$, there is a homomorphism $\beta_i : F(\bar{x}) \rightarrow A_i$ sending $[t(\bar{x})]$ to $t^{A_i}(\bar{a}_i)$ (see Theorem 2.8.11). Then

$$\beta_i(a_i) = \beta_i([t_i(\bar{x})]) = t_i^{A_i}(\bar{a}_i) \neq s_i^{A_i}(\bar{a}_i) = \beta_i([s_i(\bar{x})]) = \beta_i(b_i).$$

Thus, for every $i \in I$,

$$\beta_i(a_i) \neq \beta_i(b_i). \quad (*)$$

By Lemma 2.9.1, there is a homomorphism $\beta : F(\bar{x}) \rightarrow \prod_{i \in I} A_i$ with $\beta(x) = (\beta_i(x) : i \in I)$. Note that $\prod_{i \in I} A_i \in \mathcal{K}$ because \mathcal{K} is closed under products. We claim that β is injective. Indeed, if $a, b \in F(\bar{x})$ and $a \neq b$, then $(a, b) = (a_i, b_i)$ for some $i \in I$. Then $\beta_i(a) = \beta_i(a_i) \neq \beta_i(b_i) = \beta_i(b)$, so $\beta(a)$ and $\beta(b)$ differ at the i th coordinate.

Then β is an isomorphism from $F(\bar{x})$ to $\text{im}(\beta)$. As $\text{im}(\beta)$ is a subalgebra of $\prod_{i \in I} A_i \in \mathcal{K}$, we have $\text{im}(\beta) \in \mathcal{K}$ and then $F(\bar{x}) \in \mathcal{K}$. \square_{Claim}

Now if $A \in \bar{\mathcal{K}}$ then there is a surjective homomorphism $F(\bar{x}) \rightarrow A$ for some $\bar{\mathcal{K}}$ -free algebra $F(\bar{x})$ (Corollary 2.8.13). By the fundamental theorem on homomorphisms (Corollary 2.6.8), A is isomorphic to a quotient of $F(\bar{x})$. By the claim, $F(\bar{x}) \in \mathcal{K}$, and thus $A \in \mathcal{K}$.

This shows that $\bar{\mathcal{K}} \subseteq \mathcal{K}$. As $\mathcal{K} \subseteq \bar{\mathcal{K}}$, the class \mathcal{K} equals the equational class $\bar{\mathcal{K}}$. \square

Using the HSP theorem, one can prove the following:

Theorem 2.9.3. *If \mathcal{K} is any class of \mathcal{L} -algebras, then there is a smallest equational class $\overline{\mathcal{K}}$ containing \mathcal{K} .*

In fact, $\overline{\mathcal{K}}$ is just the intersection of all equational classes containing \mathcal{K} . The class $\overline{\mathcal{K}}$ is called the equational class *generated by* \mathcal{K} . Then, it turns out that one can characterize groups, rings, and boolean algebras as follows.

Fact 2.9.4. *The equational class of rings is generated by the ring \mathbb{R} .*

Fact 2.9.5. *The equational class of boolean algebras is generated by the two-element boolean algebra $\{\text{FALSE}, \text{TRUE}\}$.*

Recall the group $\text{Perm}(S)$ of permutations on a set S from Example 1.1.8.

Fact 2.9.6. *The equational class of groups is generated by the class of groups of the form $\text{Perm}(S)$.*

These facts help motivate the definitions of groups, rings, and boolean algebras, which may seem arbitrary at first glance.

Facts 2.9.4–2.9.6 can also be used to show that the axioms of groups, rings, and boolean algebras are “complete” with respect to equations. For example, suppose Σ is the set of ring axioms, and φ is some equation in the language of rings. Then one of two things happens:

1. The equational class $\text{Mod}(\Sigma \cup \{\varphi\})$ equals $\text{Mod}(\Sigma)$, and so any ring satisfies φ . Then φ is a redundant axiom, a logical consequence of Σ .
2. The equational class $\text{Mod}(\Sigma \cup \{\varphi\})$ is strictly smaller than the class of rings $\text{Mod}(\Sigma)$, so it does *not* contain \mathbb{R} by Fact 2.9.4. Then the equation φ is not satisfied by \mathbb{R} .

Therefore, every equation satisfied by \mathbb{R} is a logical consequence of the ring axioms. Similarly, every equation satisfied by $\{\text{FALSE}, \text{TRUE}\}$ is a logical consequence of the boolean algebra axioms.

Remark 2.9.7. The letters “HSP” stand for Homomorphic image, Subalgebra, and Product. A class \mathcal{K} is *closed under homomorphic images* if whenever $\alpha : A \rightarrow B$ is a homomorphism of \mathcal{L} -algebras (not necessarily in \mathcal{K}) and $A \in \mathcal{K}$, then $\text{im}(\alpha) \in \mathcal{K}$. Using the fundamental homomorphism theorem (Theorem 2.6.7), one can see that \mathcal{K} is closed under homomorphic images if

and only if \mathcal{K} is closed under quotients and closed under isomorphisms. Thus Theorem 2.9.2 says that \mathcal{K} is an equational class if and only if \mathcal{K} is closed under **h**omomorphic images, **s**ubalgebras, and **p**roducts.

Remark 2.9.8. Let \mathcal{K} be any class of \mathcal{L} -algebras. Define the operations H, S, P as follows:

1. $H(\mathcal{K})$ is the class of homomorphic images of algebras in \mathcal{K} .
2. $S(\mathcal{K})$ is the class of subalgebras of algebras in \mathcal{K} .
3. $P(\mathcal{K})$ is the class of products of algebras in \mathcal{K} .

Theorem 2.9.2 is actually a corollary of the full HSP theorem, which says the following

Fact 2.9.9. *If \mathcal{K} is any class of algebras, then the smallest equational class containing \mathcal{K} is $H(S(P(\mathcal{K})))$.*

The proof is similar to the proof of Theorem 2.9.2, but slightly more complicated.

Chapter 3

First-order logic

We now begin the study of model theory proper. Model theory can be seen as a generalization of universal algebra, where equations are replaced by *first-order sentences*. First-order sentences are more general than equations in two ways. First, we can use logical operations like \wedge (and), \vee (or), \neg (not), \exists (there exists), and \forall (for all). This lets us express things like the last axiom of fields:

$$(\neg(0 = 1)) \wedge \forall x (x = 0 \vee \exists y : x \cdot y = 1)$$

Second, first-order logic allows not just function symbols, but also *relation symbols* like \leq . This allows for things like the axioms of partial orders:

$$\begin{aligned} &\forall x (x \leq x) \\ &\forall x \forall y (x \leq y \wedge y \leq x \rightarrow x = y) \\ &\forall x \forall y \forall z (x \leq y \wedge y \leq z \rightarrow x \leq z). \end{aligned}$$

We can also mix function symbols and relation symbols, as in the ordered field $(\mathbb{R}, +, \cdot, -, 0, 1, \leq)$.

The following analogies hold between universal algebra and model theory:

Universal algebra	Model theory
Equations	(First-order) sentences
Algebras	Structures
Equational theories	Theories
Equational classes	Elementary classes

For example, a theory is a set of sentences, and an elementary class is a class of structures defined by a theory.

While first-order logic is much more expressive than the equational logic of universal algebra, there are still limits to its expressive power. We will see many of these in future chapters, especially Chapter 5. In the present chapter we will already see that the linear orders (\mathbb{R}, \leq) and (\mathbb{Q}, \leq) satisfy exactly the same first-order sentences (Corollary 3.7.11). One says that (\mathbb{R}, \leq) and (\mathbb{Q}, \leq) are *elementarily equivalent*.

The greater expressive power of first-order logic comes at a cost. Tools like subalgebras, products, and quotients no longer work properly.¹ Additionally, the notion of homomorphism is not very useful. Consequently, the basic tools of model theory are fairly different from universal algebra, in spite of the formal analogies between the two subjects.

This chapter is a review of these basic tools. In Sections 3.1–3.3, we define the basic concepts of model theory, namely languages, structures, terms, formulas, satisfaction, theories, models, and elementary classes. In Sections 3.4–3.5 and Section 3.8, we define some of the fundamental tools of model theory—the notions of elementary equivalence, (elementary) embeddings, (elementary) substructures, definable sets and definable functions.

Section 3.6 is about *complete theories*. As discussed in the introduction, complete theories are an important part of model theory’s motivations in mathematical logic. We give an example of a complete theory in Section 3.7, and an example of an incomplete theory in Section 3.9.

3.1 Languages, structures, formulas, and satisfaction

Definition 3.1.1. A *language* \mathcal{L} consists of a set of *function symbols*, a set of *relation symbols*, and a function assigning to each function or relation symbol X an integer $n_X \in \mathbb{N}$ called the *arity* of X . A symbol X is said to be *k-ary* if $n_X = k$. Nullary function symbols are called *constant symbols*.

Example 3.1.2. The *language of posets* contains one binary relation symbol \leq . The *language of monoids* contains one binary function symbol \cdot and one constant symbol 1 .

Let M be a set. For $n \geq 0$, an *n-ary relation* on M is a subset $R \subseteq M^n$.

¹The closest thing to these operations is *ultraproducts*, which will be discussed in Chapter 6.

If $a_1, \dots, a_n \in M$, then “ $R(a_1, \dots, a_n)$ ” means “ $(a_1, \dots, a_n) \in R$ ”. We think of R as an n -ary function $M^n \rightarrow \{\text{FALSE}, \text{TRUE}\}$.

Definition 3.1.3. Let \mathcal{L} be a language. An \mathcal{L} -structure \mathcal{M} consists of the following:

1. A set M , called the *underlying set* of \mathcal{M} .
2. A map assigning an n -ary operation $f^{\mathcal{M}} : M^n \rightarrow M$ to each n -ary function symbol f .
3. A map assigning an n -ary relation $R^{\mathcal{M}} \subseteq M^n$ to each n -ary relation symbol R .

Usually we don’t distinguish between a structure \mathcal{M} and its underlying set M , writing both as M .

Example 3.1.4. If \mathcal{L} is the language of posets, then an \mathcal{L} -structure is essentially a pair (M, \leq^M) where M is a set and \leq^M is a binary relation on M .

Definition 3.1.5. Let \mathcal{L}^+ be a language and \mathcal{L}^- be a sublanguage. For any \mathcal{L}^+ -structure M , we can form an \mathcal{L}^- -structure from M by forgetting about the symbols in $\mathcal{L}^+ \setminus \mathcal{L}^-$. The resulting structure $M \upharpoonright \mathcal{L}^-$ is called a *reduct* of M . Conversely, M is an *expansion* of $M \upharpoonright \mathcal{L}^-$.

Fix a language \mathcal{L} and a class $\mathcal{V} = \{x, y, z, \dots\}$ of *variable symbols* disjoint from the symbols in \mathcal{L} .

Definition 3.1.6. An \mathcal{L} -term is a string generated by the following rules:

- If x is a variable symbol, then x is a term.
- If f is an n -ary function symbol in \mathcal{L} , and t_1, \dots, t_n are \mathcal{L} -terms, then $f(t_1, \dots, t_n)$ is an \mathcal{L} -term.

Definition 3.1.7. An \mathcal{L} -formula is a string generated by the following rules:

1. If R is a k -ary relation symbol in \mathcal{L} and t_1, \dots, t_k are \mathcal{L} -terms, then $R(t_1, \dots, t_k)$ is an \mathcal{L} -formula.
2. If t and s are \mathcal{L} -terms, then $(t = s)$ is an \mathcal{L} -formula.

3. If φ and ψ are \mathcal{L} -formulas, then so are $\varphi \wedge \psi$, $\varphi \vee \psi$, and $\neg\varphi$.
4. \top and \perp are \mathcal{L} -formulas.
5. If φ is an \mathcal{L} -formula and x is a variable symbol, then the following are \mathcal{L} -formulas:

$$\exists x (\varphi) \in \mathcal{F}$$

$$\forall x (\varphi) \in \mathcal{F}.$$

The set of *atomic \mathcal{L} -formulas* is the subset generated by (1)–(2), while the set of *quantifier-free \mathcal{L} -formulas* is generated by (1)–(4). If x is a variable symbol and α is a term or formula, an occurrence of x in α is *bound* if it occurs inside a quantifier $\exists x (\dots)$ or $\forall x (\dots)$, and *free* otherwise. The *free variables* of α are the variables with free occurrences in α . A *closed term* is a term with no free variables (i.e., no variables). A *sentence* is a formula with no free variables.

We write “let $\varphi(x_1, \dots, x_n)$ be a formula” to mean that $\varphi(x_1, \dots, x_n)$ is a formula whose free variables are contained in $\{x_1, \dots, x_n\}$. We use the same convention for terms. If $\varphi(x_1, \dots, x_n)$ is a formula and t_1, \dots, t_n are terms, then $\varphi(t_1, \dots, t_n)$ denotes the result of replacing the free occurrences of x_i with t_i in φ .

Let M be an \mathcal{L} -structure and A be a subset of M . The language $\mathcal{L}(A)$ is obtained by adding each element of A as a new constant symbol. We regard M as an $\mathcal{L}(A)$ -structure by interpreting each new constant symbol as the corresponding element of A :

$$a^M = a.$$

If t is a closed $\mathcal{L}(M)$ -term, define the interpretation t^M recursively as follows:

$$f(t_1, \dots, t_k)^M = f^M(t_1^M, \dots, t_k^M),$$

If $t(x_1, \dots, x_n)$ is an \mathcal{L} -term or $\mathcal{L}(M)$ -term, then the interpretation t^M is the function $M^n \rightarrow M$ defined by

$$t^M(a_1, \dots, a_n) = (t(a_1, \dots, a_n))^M.$$

If φ is an $\mathcal{L}(M)$ -sentence, we define $M \models \varphi$ recursively:

$$\begin{aligned}
 M \models t = s &\iff t^M = s^M \\
 M \models R(t_1, \dots, t_k) &\iff (t_1^M, \dots, t_k^M) \in R^M \\
 M \models \varphi \vee \psi &\iff (M \models \varphi \text{ or } M \models \psi) \\
 M \models \varphi \wedge \psi &\iff (M \models \varphi \text{ and } M \models \psi) \\
 M \models \neg \varphi &\iff M \not\models \varphi \\
 M \models \top &\text{ is always true} \\
 M \models \perp &\text{ is always false} \\
 M \models \exists x \varphi(x) &\iff \exists a \in M : M \models \varphi(a) \\
 M \models \forall x \varphi(x) &\iff \forall a \in M : M \models \varphi(a).
 \end{aligned}$$

If $M \models \varphi$, we say that M *satisfies* φ , or φ is *true* in M .

3.2 Theories and elementary classes

Fix a language \mathcal{L} . An \mathcal{L} -*theory* is a set T of \mathcal{L} -sentences. Elements of T are called *axioms*. An \mathcal{L} -structure M *satisfies* T , or is a *model* of T , written $M \models T$, if

$$M \models \varphi \text{ for every } \varphi \in T.$$

The set of models of T is written $\text{Mod}(T)$. An *elementary class* is a class of the form $\text{Mod}(T)$.

Example 3.2.1. Let \mathcal{L} be the language of rings. The *theory of rings* T_{rings} has the following axioms:

$$\begin{aligned}
 &\forall x \forall y (x + y = y + x) \\
 &\forall x \forall y \forall z (x + (y + z) = (x + y) + z) \\
 &\dots
 \end{aligned}$$

Models of T_{rings} are exactly rings. The *theory of fields* T_{fields} is T_{rings} plus the following two axioms:

$$\begin{aligned}
 &\neg(0 = 1) \\
 &\forall x (x = 0 \vee \exists y (x \cdot y = 1)).
 \end{aligned}$$

Models of T_{fields} are fields.

3.3 Common abbreviations

The following abbreviations are standard:

Abbreviation	Meaning
$\varphi \rightarrow \psi$	$\neg\varphi \vee \psi$
$\varphi \leftarrow \psi$	$\psi \rightarrow \varphi$
$\varphi \leftrightarrow \psi$	$(\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi)$
$t \neq s$	$\neg t = s$
$\bigwedge_{i=1}^n \varphi_i$	$\varphi_1 \wedge \cdots \wedge \varphi_n$ [or \top if $n = 0$]
$\bigvee_{i=1}^n \varphi_i$	$\varphi_1 \vee \cdots \vee \varphi_n$ [or \perp if $n = 0$]
$\exists^{\geq n} \varphi(x)$	$\exists y_1, \dots, y_n \left(\bigwedge_{i=1}^n \varphi(y_i) \wedge \bigwedge_{1 \leq i < j \leq n} (y_i \neq y_j) \right)$
$\exists^= \varphi(x)$	$\exists^{\geq n} \varphi(x) \wedge \neg \exists^{\geq n+1} \varphi(x)$
$\exists! x \varphi(x)$	$\exists^=1 x \varphi(x)$.

Additionally, if \mathcal{L} contains a symbol \leq , then

$$\begin{aligned} t \geq s &\text{ means } s \leq t \\ t < s &\text{ means } t \leq s \wedge t \neq s \\ t > s &\text{ means } s < t. \end{aligned}$$

Example 3.3.1. Let $\mathcal{L} = \{\leq\}$ be the language of posets. The *theory of posets*, whose models are posets, has the following axioms:

$$\begin{aligned} &\forall x (x \leq x) \\ &\forall x, y, z (x \leq y \wedge y \leq z \rightarrow x \leq z) \\ &\forall x, y (x \leq y \wedge y \leq x \rightarrow x = y). \end{aligned}$$

The *theory of linear orders* adds one more axiom:

$$\forall x, y (x \leq y \vee y \leq x).$$

3.4 Elementary equivalence and embeddings

The *complete theory* of an \mathcal{L} -structure M is the set

$$\text{Th}(M) := \{\varphi : \varphi \text{ is an } \mathcal{L}\text{-sentence and } M \models \varphi\}.$$

Two \mathcal{L} -structures M and N are *elementarily equivalent*, written $M \equiv N$, if $\text{Th}(M) = \text{Th}(N)$, meaning that $M \models \varphi \iff N \models \varphi$ for every \mathcal{L} -sentence φ .

Theorem 3.4.1. *A structure N is a model of $\text{Th}(M)$ if and only if $N \equiv M$.*

Proof. Note that $N \models \text{Th}(M)$ iff $\text{Th}(N) \supseteq \text{Th}(M)$. (More generally, $N \models T$ iff $\text{Th}(N) \supseteq T$.) So we must rule out the case that $\text{Th}(N) \supsetneq \text{Th}(M)$. Suppose $\varphi \in \text{Th}(N) \setminus \text{Th}(M)$. Then $N \models \varphi$ and $M \not\models \varphi$, implying that $N \not\models \neg\varphi$ and $M \models \neg\varphi$. Then $(\neg\varphi) \in \text{Th}(M) \setminus \text{Th}(N)$, contradicting the fact that $\text{Th}(N) \supseteq \text{Th}(M)$. \square

Definition 3.4.2. Let M, N be \mathcal{L} -structures. A function $\alpha : M \rightarrow N$ is an *embedding* if α is injective, and α strictly preserves the function and relation symbols:

$$\begin{aligned} \alpha(f^M(b_1, \dots, b_n)) &= f^N(\alpha(b_1), \dots, \alpha(b_n)) \\ R^M(b_1, \dots, b_n) &\iff R^N(\alpha(b_1), \dots, \alpha(b_n)). \end{aligned}$$

An *isomorphism* is a bijective embedding. Two structures M and N are *isomorphic*, written $M \cong N$, if there is an isomorphism from M to N .

The analogue of Theorem 1.4.6 holds, and so \cong is an equivalence relation.

Theorem 3.4.3. *Let $f : M \rightarrow N$ be an embedding and a_1, \dots, a_n be in M .*

1. *If $t(x_1, \dots, x_n)$ is a term, then*

$$f(t^M(a_1, \dots, a_n)) = t^N(f(a_1), \dots, f(a_n)).$$

2. *If $\varphi(x_1, \dots, x_n)$ is a quantifier-free formula, then*

$$M \models \varphi(a_1, \dots, a_n) \iff N \models \varphi(f(a_1), \dots, f(a_n)).$$

3. *If f is an isomorphism and $\varphi(x_1, \dots, x_n)$ is any formula, then*

$$M \models \varphi(a_1, \dots, a_n) \iff N \models \varphi(f(a_1), \dots, f(a_n)).$$

Proof. By induction on the complexity of t or φ , like in Theorem 1.4.9. (In fact, part (1) is an instance of Theorem 1.4.9.) \square

Definition 3.4.4. A function $\alpha : M \rightarrow N$ is an *elementary embedding* if α preserves all \mathcal{L} -formulas $\varphi(x_1, \dots, x_n)$, in the sense that

$$M \models \varphi(b_1, \dots, b_n) \iff N \models \varphi(\alpha(b_1), \dots, \alpha(b_n)). \quad (*)$$

Remark 3.4.5. 1. Any elementary embedding is an embedding: apply (*) to the following formulas:

$$\begin{aligned} x_1 &= x_2 \\ R(x_1, \dots, x_k) \\ f(x_1, \dots, x_k) &= x_{k+1} \end{aligned}$$

2. Theorem 3.4.3(3) says that isomorphisms are elementary embeddings.
3. If $\alpha : M \rightarrow N$ is an elementary embedding, then $M \equiv N$. To see this, take $n = 0$ in (*), so that φ is a sentence. Then (*) says that $M \models \varphi \iff N \models \varphi$.

Corollary 3.4.6. *If $M \cong N$, then $M \equiv N$.*

3.5 Substructures and extensions

Let M be an \mathcal{L} -structure.

Definition 3.5.1. A *substructure* of M is a subset $A \subseteq M$ such that for any k -ary function symbol f in \mathcal{L} ,

$$a_1, \dots, a_k \in A \implies f^M(a_1, \dots, a_k) \in A.$$

If A is a substructure of M , we regard A as an \mathcal{L} -structure by defining f^A and R^A to be the restrictions of f^M and R^M to A . In this way, substructures are structures, not just sets.

If M is an \mathcal{L} -structure, an *extension* of M is an \mathcal{L} -structure N such that M is a substructure of N .

Definition 3.5.2. If A is a subset of M , then $\langle A \rangle_M$ denotes the smallest substructure of M containing A , which is

$$\{t^M(\bar{b}) : t(x_1, \dots, x_n) \text{ is an } \mathcal{L}\text{-term and } \bar{b} \in A^n\}$$

The substructure $\langle A \rangle_M$ is called the substructure *generated* by A . We say that M is *finitely generated* if $M = \langle A \rangle$ for some finite $A \subseteq M$.

Remark 3.5.3. If the language \mathcal{L} has no function symbols or constant symbols, then every subset of M is a substructure, and so $\langle A \rangle_M = A$.

Remark 3.5.4. If A is a substructure of M , then the inclusion $A \hookrightarrow M$ is an embedding. Therefore

$$A \models \varphi(\bar{a}) \iff M \models \varphi(\bar{a})$$

for *quantifier-free* formulas φ and tuples \bar{a} in A .

Definition 3.5.5. Let M and N be \mathcal{L} -structures. We say that M is an *elementary substructure* of N , written $M \preceq N$, or that N is an *elementary extension* of M , written $N \succeq M$, if M is a substructure of N and the inclusion $M \rightarrow N$ is an elementary embedding, meaning that

$$M \models \varphi(a_1, \dots, a_n) \iff N \models \varphi(a_1, \dots, a_n) \quad (*)$$

for \mathcal{L} -formulas φ and $\bar{a} \in M^n$.

Equation $(*)$ says that M and N satisfy the same $\mathcal{L}(M)$ -sentences.

Theorem 3.5.6. *Let f be an embedding from M to N .*

1. *The image $\text{im}(f)$ is a substructure of N , and $M \rightarrow \text{im}(f)$ is an isomorphism.*
2. *If f is an elementary embedding, then $\text{im}(f) \preceq N$.*

Proof. 1. Straightforward. The fact that $\text{im}(f)$ is a substructure is like the analogous fact for homomorphisms (Theorem 2.6.1).

2. If $M' = \text{im}(f)$, then the inclusion $M' \hookrightarrow N$ is the composition of two elementary maps:

$$M' \xrightarrow{\cong} M \xrightarrow{f} N. \quad \square$$

3.6 Complete theories

We write $T \vdash \varphi$ if φ is provable from T , and $T \models \varphi$ if every model of T satisfies φ .

Fact 3.6.1 (Soundness and completeness theorem). *If T is a theory and φ is a sentence, then*

$$T \vdash \varphi \iff T \models \varphi.$$

A theory T is *inconsistent* if the following equivalent conditions hold:

1. $T \vdash \perp$.
2. $T \vdash \varphi$ for all φ .
3. $T \vdash \varphi$ and $T \vdash \neg\varphi$ for some φ .

Otherwise, T is *consistent*. Note that $T \models \perp$ if and only if T has no models, because no structure M satisfies \perp . Therefore the completeness theorem implies the following:

Theorem 3.6.2. *T is consistent if and only if T has a model.*

A theory T is *complete* if T is consistent and for any sentence φ ,

$$T \vdash \varphi \text{ or } T \vdash \neg\varphi.$$

Theorem 3.6.3. *If T is complete and $M \models T$, then for any sentence φ ,*

$$M \models \varphi \iff T \vdash \varphi.$$

Proof. As $M \models T$ and T is complete,

$$\begin{aligned} T \vdash \varphi &\implies M \models \varphi \\ T \not\vdash \varphi &\implies T \vdash \neg\varphi \implies M \models \neg\varphi \implies M \not\models \varphi. \end{aligned} \quad \square$$

Theorem 3.6.4. *Let T be a theory. Then T is complete if and only if any two models $M_1, M_2 \models T$ are elementarily equivalent.*

Proof. First suppose T is complete. If M_1, M_2 are two models, then $\text{Th}(M_1) = \{\varphi : T \vdash \varphi\} = \text{Th}(M_2)$ by Theorem 3.6.3, and so $M_1 \equiv M_2$.

Conversely suppose T is incomplete, with $T \not\vdash \varphi$ and $T \not\vdash \neg\varphi$. By the completeness theorem, there are models $M_1, M_2 \models T$ with $M_1 \models \neg\varphi$ and $M_2 \models \varphi$. Then $M_1 \not\equiv M_2$. \square

Fact 3.6.5. *Let \mathcal{L} be a countable language. If T is finite or more generally if T is computably enumerable, then the set $\bar{T} = \{\varphi : T \vdash \varphi\}$ is computably enumerable.*

Theorem 3.6.6. *If \mathcal{L} is countable, T is complete and computably enumerable, and M is a model of T , then $\text{Th}(M)$ is computable—there is an algorithm which takes an \mathcal{L} -sentence φ and determines whether $M \models \varphi$.*

Proof. By Theorem 3.6.3, $\text{Th}(M) = \overline{T}$, which is computably enumerable by Fact 3.6.5. However,

$$\varphi \notin \text{Th}(M) \iff \neg\varphi \in \text{Th}(M),$$

and so $\text{Th}(M)$ is also co-c.e. As $\text{Th}(M)$ is both c.e. and co-c.e., it is computable. \square

3.7 Completeness via back-and-forth systems

Let M, N be two \mathcal{L} -structures.

Definition 3.7.1. A *partial elementary map* from M to N is a bijection $f : A \rightarrow B$ where A and B are subsets of M and N , respectively, and f preserves all \mathcal{L} -formulas, in the sense that

$$M \models \varphi(a_1, \dots, a_n) \iff N \models \varphi(f(a_1), \dots, f(a_n))$$

for $n \geq 0$, $\varphi(x_1, \dots, x_n)$ an \mathcal{L} -formula, and $a_1, \dots, a_n \in A$.

Remark 3.7.2. Taking $n = 0$, we see that if f is a partial elementary map from M to N , then $M \equiv N$.

Conversely, $\emptyset : \emptyset \rightarrow \emptyset$ is a partial elementary map from M to N if and only if $M \equiv N$.

Remark 3.7.3. An elementary embedding from M to N is the same thing as a partial elementary map f from M to N with $\text{dom}(f) = M$.

Definition 3.7.4. A *partial isomorphism* from M to N is an isomorphism $f : A \rightarrow B$ where A is a substructure of M and B is a substructure of N .

By Theorem 3.4.3, partial isomorphisms preserve quantifier-free formulas: if $f : A \rightarrow B$ is a partial isomorphism and $\bar{a} \in A^n$ and $\varphi(\bar{x})$ is quantifier-free, then

$$M \models \varphi(\bar{a}) \iff A \models \varphi(\bar{a}) \iff B \models \varphi(f(\bar{a})) \iff N \models \varphi(f(\bar{a})).$$

Definition 3.7.5. A *back-and-forth system* between M and N is a family \mathcal{F} of partial isomorphisms satisfying the following conditions:

1. Forward: if $f : A \rightarrow B$ is in \mathcal{F} , and $a' \in M$, then there is $f' : A' \rightarrow B'$ in \mathcal{F} such that f' extends f and $a' \in A'$.
2. Backward: if $f : A \rightarrow B$ is in \mathcal{F} , and $b' \in N$, then there is $f' : A' \rightarrow B'$ in \mathcal{F} such that f' extends f and $b' \in B'$.

Theorem 3.7.6. *If \mathcal{F} is a back-and-forth system, then every $f \in \mathcal{F}$ is a partial elementary map from M to N .*

Proof. We show by induction on the complexity of $\varphi(x_1, \dots, x_n)$ that if $f : A \rightarrow B$ is in \mathcal{F} and $a_1, \dots, a_n \in A$, then

$$M \models \varphi(a_1, \dots, a_n) \iff N \models \varphi(f(a_1), \dots, f(a_n)). \quad (\dagger)$$

The case of atomic formulas holds because f is a partial isomorphism. The case of boolean operations is straightforward. It remains to consider the case where $\varphi(\bar{x})$ is $\exists y : \psi(\bar{x}, y)$. By symmetry, we only need to prove the (\Rightarrow) direction of (\dagger) . Suppose $M \models \exists y : \psi(\bar{a}, y)$. Take some $a' \in M$ such that $M \models \psi(\bar{a}, a')$. By the “forward” condition, there is $g \in \mathcal{F}$ extending f with $a' \in \text{dom}(g)$. By induction, $M \models \psi(\bar{a}, a') \implies N \models \psi(g(\bar{a}), g(a'))$, and then $N \models \varphi(g(\bar{a}))$. But $g(\bar{a}) = f(\bar{a})$, so $N \models \varphi(f(\bar{a}))$ as desired. \square

Corollary 3.7.7. *Let M, N be two structures. If there is a non-empty back-and-forth system between M and N , then $M \equiv N$.*

Definition 3.7.8. A linear order (M, \leq) is *dense* if for any $x < y$ there is z with $x < z < y$.

Definition 3.7.9. DLO is the theory of non-empty dense linear orders without endpoints. That is, $(M, \leq) \models \text{DLO}$ if (M, \leq) is a non-empty dense linear order without a greatest or least element.

Theorem 3.7.10. *Let M and N be models of DLO and let \mathcal{F} be the class of finite partial isomorphisms $f : A \rightarrow B$ between M and N . Then \mathcal{F} is a back-and-forth system.*

Proof. We prove the forward condition; the backward condition follows by symmetry. Fix a finite partial isomorphism $f : A \rightarrow B$ and an element $a' \in M$. We must extend f to a larger finite partial isomorphism f' containing a' . Enumerate the elements of A :

$$\begin{aligned} A &= \{a_1, \dots, a_n\} \\ a_1 &< a_2 < \dots < a_n \end{aligned}$$

Let $b_i = f(a_i)$. Then

$$B = \{b_1, \dots, b_n\}$$

$$b_1 < b_2 < \dots < b_n.$$

There are five cases.

1. $n = 0$, so that A, B are empty. Take any $b' \in N$. (N is non-empty.) Then $f' : \{a'\} \rightarrow \{b'\}$ extends f and is defined at a' .
2. $a' = a_i \in A$. Then we can take $f' = f$.
3. $a' < a_1$. Take $b' \in N$ with $b' < b_1$. (N has no minimum.) Then there is a partial isomorphism

$$f' : \{a', a_1, \dots, a_n\} \rightarrow \{b', b_1, \dots, b_n\}$$

extending f and sending a' to b' .

4. $a' > a_n$. This case is similar to the previous case.
5. $a_i < a' < a_{i+1}$ for some i . Take $b' \in N$ with $b_i < b' < b_{i+1}$. (N is densely ordered.) Then there is a partial isomorphism

$$f' : \{a_1, \dots, a_i, a', a_{i+1}, \dots, a_n\} \rightarrow \{b_1, \dots, b_i, b', b_{i+1}, \dots, b_n\}$$

extending f and sending a' to b' . □

Corollary 3.7.11. *1. The theory DLO is complete—any two models M, N are elementarily equivalent.*

2. *If $M \models \text{DLO}$, then $\text{Th}(M)$ is decidable.*

For example, the complete theory of (\mathbb{Q}, \leq) is decidable.

3.8 Definable sets and functions

Fix a language \mathcal{L} and an \mathcal{L} -structure M . If $\varphi(x_1, \dots, x_n)$ is an \mathcal{L} -formula, then $\varphi(M^n)$ or $\varphi(M)$ denotes the set *defined by* φ :

$$\varphi(M^n) := \{\bar{a} \in M^n : M \models \varphi(\bar{a})\}.$$

More generally, if $\varphi(x_1, \dots, x_n; y_1, \dots, y_m)$ is an \mathcal{L} -formula and $\bar{b} \in M^m$, then $\varphi(M^n, \bar{b})$ or $\varphi(M, \bar{b})$ denotes the set defined by $\varphi(\bar{x}, \bar{b})$:

$$\varphi(M^n, \bar{b}) := \{\bar{a} \in M^n : M \models \varphi(\bar{a}, \bar{b})\}.$$

Sets of this form are called *definable sets*. A set $D \subseteq M^n$ is *A-definable* or *definable over A* if $D = \varphi(M^n, \bar{b})$ with $\bar{b} \in A^m$. That is, an *A-definable set* is a set defined by an $\mathcal{L}(A)$ -formula. Definable sets are the same thing as *M-definable sets*. A *0-definable set* is a \emptyset -definable set. If D, E are definable sets, a function $f : D \rightarrow E$ is *definable* if the graph $\Gamma(f) = \{(x, y) \in D \times E : f(x) = y\}$ is a definable set. *A-definable functions* are defined similarly.

Remark 3.8.1. If $t(x_1, \dots, x_n)$ is an $\mathcal{L}(M)$ -term, then the function $t^M : M^n \rightarrow M$ is definable, defined by the formula $t(\bar{x}) = y$.

Theorem 3.8.2. *If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are definable, then $g \circ f : X \rightarrow Z$ is definable.*

Proof. Let $\varphi(\bar{x}, \bar{y})$ be an $\mathcal{L}(M)$ -formula defining f , meaning that

$$f(\bar{a}) = \bar{b} \iff M \models \varphi(\bar{a}, \bar{b}).$$

Similarly, let $\psi(\bar{y}, \bar{z})$ define g . Let $\theta(\bar{x}, \bar{z})$ be

$$\exists \bar{y} (\varphi(\bar{x}, \bar{y}) \wedge \psi(\bar{y}, \bar{z})).$$

Then θ defines $g \circ f$:

$$\begin{aligned} M \models \theta(\bar{a}, \bar{c}) &\iff \exists \bar{b} (M \models \varphi(\bar{a}, \bar{b}) \text{ and } M \models \psi(\bar{b}, \bar{c})) \\ &\iff \exists \bar{b} (f(\bar{a}) = \bar{b} \text{ and } g(\bar{b}) = \bar{c}) \\ &\iff g(f(\bar{a})) = \bar{c}. \end{aligned}$$

□

3.9 Incompleteness of Peano Arithmetic

Consider \mathbb{N} as a structure $(\mathbb{N}, +, \cdot, 0, 1)$.

Remark 3.9.1. The relation $x \leq y$ is definable, defined by

$$\exists z : z + x = y.$$

Remark 3.9.2. The function $|x - y|$ is definable.

Proof. $|x - y| = z$ if and only if $x = y + z \vee y = x + z$. \square

Definition 3.9.3. If $n > 0$ and $a \in \mathbb{N}$, then $a \bmod n$ denotes the unique $x \in \{0, 1, \dots, n-1\}$ such that $x \equiv a \pmod{n}$.

Lemma 3.9.4. *The function $x \bmod y$ is definable.*

Proof. $(x \bmod y) = z$ if and only if

$$z + 1 \leq y \wedge (\exists k : k \cdot y + z = x). \quad \square$$

Definition 3.9.5. For $a, b, x \in \mathbb{N}$, $\gamma(a, b, x) = (a \bmod ((x+1)b+1))$.

Lemma 3.9.6. *For any finite sequence $c_0, c_1, \dots, c_n \in \mathbb{N}$, there are $a, b \in \mathbb{N}$ such that $c_i = \gamma(a, b, i)$ for $0 \leq i \leq n$.*

Proof. Take $m > \max(c_0, \dots, c_n, n)$, and let $b = m! = 1 \cdot 2 \cdot 3 \cdot 4 \cdots m$.

Claim. The numbers $\{b+1, 2b+1, 3b+1, \dots, (n+1)b+1\}$ are pairwise coprime.

Proof. Suppose a prime number p divides both $ib+1$ and $jb+1$ for some $1 \leq i < j \leq n+1$. Then p divides the difference $(j-i)b$, and one of two things happens:

- p divides b .
- p divides $j-i$. Then $p \leq j-i \leq n \leq m$, so p divides $m! = b$.

Either way, p divides b , and so

$$0 \equiv ib+1 \equiv i0+1 = 1 \pmod{p},$$

a contradiction. \square_{Claim}

By the Chinese remainder theorem, there is an a such that

$$a \equiv c_i \pmod{(i+1)b+1}$$

for each $0 \leq i \leq n$. But $c_i \leq m \leq b < (i+1)b+1$, and so

$$c_i = (a \bmod ((i+1)b+1)) = \gamma(a, b, i). \quad \square$$

Theorem 3.9.7. *Let $f(n)$ be the n th Fibonacci number $(1, 1, 2, 3, 5, 8, 13, \dots)$*

$$\begin{aligned} f(0) &= f(1) = 1 \\ f(n+2) &= f(n) + f(n+1) \end{aligned}$$

Then the function $f : \mathbb{N} \rightarrow \mathbb{N}$ is definable.

Proof. For fixed n and k , the following are equivalent:

1. $f(n) = k$.
2. There are c_0, \dots, c_n such that the following hold:

$$\begin{aligned} c_0 &= 1 \\ c_1 &= 1 \\ c_i &= c_{i-1} + c_{i-2} \text{ for all } 2 \leq i \leq n. \end{aligned}$$

3. There are a and b such that the following hold:

$$\begin{aligned} \gamma(a, b, 0) &= 1 \\ \gamma(a, b, 1) &= 1 \\ \gamma(a, b, i) &= \gamma(a, b, i-1) + \gamma(a, b, i-2) \text{ for all } 2 \leq i \leq n. \end{aligned}$$

Indeed, (2) and (3) are equivalent by Lemma 3.9.6. Condition (3) is easily expressed by a formula. \square

Theorem 3.9.8. *If $f : \mathbb{N}^k \rightarrow \mathbb{N}$ is computable, then f is definable.*

Proof sketch. The only hard thing to check is primitive recursion, which is handled by the method of Theorem 3.9.7. \square

Corollary 3.9.9. *If $S \subseteq \mathbb{N}^k$ is computable, then S is definable.*

Theorem 3.9.10. *$\text{Th}(\mathbb{N})$ is incomputable.*

Proof. Fix a reasonable enumeration of Turing machines. Let $S \subseteq \mathbb{N}^2$ be the set of pairs (n, k) such that the n th Turing machine halts within k steps. By Corollary 3.9.9, S is definable, defined by an $\mathcal{L}(\mathbb{N})$ -formula $\varphi(x, y, n_1, n_2, \dots)$. Replacing n_i with $\underbrace{1 + 1 + \dots + 1}_{n_i \text{ times}}$, we see that S is \emptyset -definable, defined by

an \mathcal{L} -formula $\varphi(x, y)$. Then for any $n \in \mathbb{N}$,

$$\mathbb{N} \models \exists y \varphi(\underbrace{1 + 1 + \dots + 1}_{n \text{ times}}, y) \iff (\text{the } n\text{th Turing machine halts}).$$

Therefore the halting problem reduces to $\text{Th}(\mathbb{N})$. \square

Let \mathcal{L} be the language $\{+, \cdot, 0, 1\}$. *Peano Arithmetic* (PA) consists of the following axioms:

$$\begin{aligned}
& \forall x : x + 1 \neq 0 \\
& \forall x : (x = 0 \vee (\exists y : x = y + 1)) \\
& \forall x, y : (x + 1 = y + 1 \rightarrow x = y) \\
& \forall x : x + 0 = x \\
& \forall x, y : x + (y + 1) = (x + y) + 1 \\
& \forall x : x \cdot 0 = 0 \\
& \forall x, y : x \cdot (y + 1) = (x \cdot y) + x
\end{aligned}$$

as well as the infinite set of axioms $\{Ind_\varphi : \varphi(x, y_1, \dots, y_n) \text{ an } \mathcal{L}\text{-formula}\}$, where Ind_φ is the induction axiom

$$\forall \bar{y} (\varphi(0, \bar{y}) \wedge (\forall x \varphi(x, \bar{y}) \rightarrow \varphi(x + 1, \bar{y})) \rightarrow \forall x \varphi(x, \bar{y})).$$

For example, $(\mathbb{N}, +, \cdot, 0, 1) \models \text{PA}$.

Remark 3.9.11. The induction schema says that if $D \subseteq M^1$ is a definable set such that $0 \in D$ and $x \in D \implies x + 1 \in D$, then D contains every $x \in M$.

PA is computably axiomatized, so Theorems 3.6.6 and 3.9.10 yield the following:

Corollary 3.9.12. *Peano arithmetic is incomplete.*

Then Theorem 3.6.4 gives the following:

Corollary 3.9.13. *There is a model $M \models \text{PA}$ with $M \not\cong \mathbb{N}$.*

Chapter 4

The compactness theorem

The *compactness theorem* is the fundamental tool of model theory. It says the following:

Theorem (Compactness theorem). *Let T be a theory. Suppose every finite subset $T_0 \subseteq_f T$ has a model. Then T_0 has a model.*

The reason for the name “compactness” will be made clear in Chapter 7 when we discuss topologies.

As we will see, the compactness theorem is a powerful method for constructing models. In the next chapter, we will use it to prove the Löwenheim-Skolem theorem (Theorem 5.4.4), and in Chapter 8 we will use it to realize types. For now, we focus on proving the compactness theorem. In the current chapter we give the proof using *Henkin’s method*, and we will later see a different proof using *ultraproducts* (Theorem 6.2.6).

The core idea of Henkin’s method is similar to the construction of free algebras in Section 2.8: given a theory T we want to take the “free” model of T , the model in which equations and relations only hold if T forces them to hold, and every element is named by a term. Unfortunately, this “free” model of T usually doesn’t exist, *unless* T has two unusual properties—*completeness* and the *witness property* (Section 4.3). However, it turns out that if T is a theory which “should” have a model, then we can modify T to get a theory with the two desired properties. All we need to do is apply the following operations ad nauseam:

1. If T doesn’t disprove φ , add φ to T .

2. If T proves $\exists x \varphi(x)$, add a new constant symbol c to the language and add the sentence $\varphi(c)$ into T .

This process is discussed in Section 4.4.

4.1 A temporary convention

When we defined formulas (Definition 3.1.7), we chose the following logical operators to be fundamental:

$$\wedge, \vee, \top, \perp, \exists, \forall, \neg.$$

In contrast, notions like \leftrightarrow and $\exists!$ are defined in terms the fundamental operators (see Section 3.3). For example, $\varphi \rightarrow \psi$ is an abbreviation for $\neg\varphi \vee \psi$.

In this chapter, we temporarily switch conventions, regarding

$$\wedge, \top, \exists, \neg,$$

as fundamental logical operators, and \vee, \forall, \perp as derived notions like \leftrightarrow or $\exists!$. Specifically:

$$\begin{aligned}\varphi \vee \psi &:= \neg(\neg\varphi \wedge \neg\psi) \\ \perp &:= \neg\top \\ \exists x \varphi(x) &:= \neg\forall x \neg\varphi(x).\end{aligned}$$

This convention simplifies the proofs of the compactness and completeness theorems, without impacting the conclusions of these theorems.

4.2 Premodels

A \mathcal{L} -prestructure is a pair (M, \approx) where M is an \mathcal{L} -structure and \approx is an equivalence relation on M such that the following hold:

1. If f is a k -ary function symbol and $a_i \approx b_i$ for $i = 1, \dots, k$, then

$$f(a_1, \dots, a_k) \approx f(b_1, \dots, b_k).$$

2. If R is a k -ary relation symbol and $a_i \approx b_i$ for $i = 1, \dots, k$, then

$$R(a_1, \dots, a_k) \iff R(b_1, \dots, b_k).$$

The first condition says that \approx is a congruence with respect to the function symbols.

Definition 4.2.1. If (M, \approx) is an \mathcal{L} -prestructure, then M/\approx is the following \mathcal{L} -structure:

1. The underlying set is $M/\approx = \{[a] : a \in M\}$, where $[a] = \{b \in M : b \approx a\}$.
2. If f is a k -ary function symbol, then

$$f([a_1], \dots, [a_k]) = [f(a_1, \dots, a_k)]$$

3. If R is a k -ary relation symbol, then

$$R([a_1], \dots, [a_n]) \iff R(a_1, \dots, a_n).$$

This is well-defined by definition of prestructure (using Theorem 2.3.1).

If M is an \mathcal{L} -prestructure, if $\varphi(x_1, \dots, x_n)$ is an \mathcal{L} -sentence, and $a_1, \dots, a_n \in M$, then we define $M \models \varphi(a_1, \dots, a_n)$ exactly as for ordinary structures, except that

$$M \models t(\bar{a}) = s(\bar{a}) \iff t^M(\bar{a}) \approx s^M(\bar{a}).$$

In other words, we treat the symbol “=” as a non-logical symbol whose interpretation is \approx .

Definition 4.2.2. A *premodel* of T is an \mathcal{L} -prestructure satisfying T .

Theorem 4.2.3. If (M, \approx) is an \mathcal{L} -prestructure, then

$$M \models \varphi(a_1, \dots, a_n) \iff (M/\approx) \models \varphi([a_1], \dots, [a_n]).$$

Proof. By induction on the complexity of φ . For example, if $\varphi(x_1, \dots, x_n)$ is $\exists y : \psi(\bar{x}, y)$, then

$$\begin{aligned} M \models \varphi(a_1, \dots, a_n) &\iff \exists b \in M : M \models \psi(\bar{a}, b) \\ &\iff \exists b \in M : (M/\approx) \models \psi([a_1], \dots, [a_n], [b]) \\ &\iff \exists c \in (M/\approx) : (M/\approx) \models \psi([a_1], \dots, [a_n], c) \\ &\iff (M/\approx) \models \varphi([a_1], \dots, [a_n]), \end{aligned}$$

as every element of M/\approx has the form $[b]$ for some $b \in M$. \square

Corollary 4.2.4. If (M, \approx) is a premodel of T , then M/\approx is a model of T .

4.3 The witness property

Definition 4.3.1. Let T be an \mathcal{L} -theory.

- T is *satisfiable* if T has a model.
- T is *finitely satisfiable* if every $T_0 \subseteq_f T$ is satisfiable.
- T is *complete* if for every \mathcal{L} -sentence φ , $\varphi \in T$ or $\neg\varphi \in T$.
- T has the *witness property* if whenever $(\exists x \varphi(x)) \in T$, there is a closed term t such that $\varphi(t) \in T$.

Recall from §3.6 that the notation $T \models \varphi$ means that every model of T satisfies φ .

Lemma 4.3.2. Suppose an \mathcal{L} -theory T is finitely satisfiable and complete, $T_0 \subseteq_f T$, and $T_0 \models \varphi$. Then $\varphi \in T$.

Proof. Otherwise, $\neg\varphi \in T$. Then $T_0 \cup \{\neg\varphi\}$ shows T is *not* finitely satisfiable. \square

Lemma 4.3.3. Suppose an \mathcal{L} -theory T is finitely satisfiable and complete and has the witness property. Let φ, ψ be sentences and $\theta(x)$ be a formula.

1. $\neg\varphi \in T \iff \varphi \notin T$.
2. $\varphi \wedge \psi \in T \iff (\varphi \in T \text{ and } \psi \in T)$
3. $\top \in T$.
4. $(\exists x \theta(x)) \in T$ if and only if there is a closed term t such that $\theta(t) \in T$.

Proof. 1. Either $\varphi \in T$ or $\neg\varphi \in T$ by completeness. If both φ and $\neg\varphi$ are in T , then T is not finitely satisfiable.

2. Both directions hold by Lemma 4.3.2, because

$$\begin{aligned} \{\varphi, \psi\} &\models \varphi \wedge \psi \\ \varphi \wedge \psi &\models \varphi \text{ and } \varphi \wedge \psi \models \psi. \end{aligned}$$

3. Lemma 4.3.2 again.

4. The \Leftarrow direction holds by Lemma 4.3.2, and the \Rightarrow direction holds because T has the witness property. \square

Theorem 4.3.4. *Let T be finitely satisfiable and complete, with the witness property. Then T has a model.*

Proof. By Corollary 4.2.4, it suffices to build a premodel $(M, \approx) \models T$. The underlying set of M will be the set of closed \mathcal{L} -terms. If f is a k -ary function symbol and $t_1, \dots, t_k \in M$, let $f^M(t_1, \dots, t_k) = f(t_1, \dots, t_k)$. If R is a k -ary function symbol and $t_1, \dots, t_k \in M$, let $R^M(t_1, \dots, t_k)$ hold iff $R(t_1, \dots, t_k) \in T$. Finally, let $t \approx s$ hold iff $(t = s) \in T$.

Claim. The relation \approx is transitive.

Proof. Suppose $\{s = t, t = u\} \subseteq T$. Note that $\{s = t, t = u\} \models \{s = u\}$. (Any structure which satisfies $s = t$ and $t = u$ must also satisfy $s = u$.) By Lemma 4.3.2, $(s = u) \in T$. \square_{Claim}

Similar arguments show that \approx is symmetric, reflexive, and respects the function and relation symbols. Thus M is an \mathcal{L} -prestructure.

Claim. If φ is an \mathcal{L} -sentence, then $M \models \varphi \iff \varphi \in T$.

Proof. Proceed by induction on φ :

1. If φ is an atomic formula, then $M \models \varphi \iff \varphi \in T$ by choice of the \mathcal{L} -structure:
 - (a) $M \models R(t_1, \dots, t_k) \iff R(t_1, \dots, t_k) \in T$ by choice of R^M .
 - (b) $M \models t = s \iff (t = s) \in T$ by choice of \approx .
 - (c) $M \models f(t_1, \dots, t_k) = s \iff (f(t_1, \dots, t_k) = s) \in T$ by choice of f^M .
2. The logical operators $\neg, \top, \wedge, \exists$ work correctly by Lemma 4.3.3 and induction. For example,

$$\begin{aligned}
 M \models \exists x \varphi(x) &\iff \exists t \in M : M \models \varphi(t) \\
 &\iff \exists t \in M : \varphi(t) \in T && \text{by induction} \\
 &\iff (\exists x \varphi(x)) \in T && \text{by Lemma 4.3.3(4).} \quad \square_{\text{Claim}}
 \end{aligned}$$

By the Claim, M is a premodel of T . By Corollary 4.2.4, the quotient M/\approx is a model of T . \square

Definition 4.3.5. Suppose T is finitely satisfiable and complete and has the witness property. The *canonical model* of T is the model constructed in the proof of Theorem 4.3.4.

The canonical model of T is characterized up to isomorphism by the fact that every element $a \in M$ is named by a closed term t , in the sense that $a = t^M$.

Definition 4.3.6. Let \mathcal{C} be a collection of closed \mathcal{L} -terms. An \mathcal{L} -theory has the *witness property over \mathcal{C}* if whenever $(\exists x \varphi(x)) \in T$, there is a term $t \in \mathcal{C}$ such that $\varphi(t) \in T$.

Theorem 4.3.7. Let T be a finitely satisfiable, complete theory with the witness property over \mathcal{C} , and let M be the canonical model of T . Then every element of M is named by a term in \mathcal{C} .

Proof. Fix $a \in M$. Then $a = t^M$ for some term t . The \mathcal{L} -sentence

$$\exists x x = t$$

holds in any \mathcal{L} -structure, so it must be in T by Lemma 4.3.2. By the witness property over \mathcal{C} , there is $t' \in \mathcal{C}$ such that $(t' = t) \in T$. Then $M \models t' = t$, so $a = t^M = (t')^M$, and a is named by a term in \mathcal{C} . \square

4.4 Compactness via Henkin's method

Theorem 4.4.1. Let I be a linear order and $\{T_i\}_{i \in I}$ be a chain of finitely satisfiable theories, meaning that $i < i' \implies T_i \subseteq T_{i'}$. Then the union $T = \bigcup_{i \in I} T_i$ is finitely satisfiable.

Proof. Suppose $T_0 \subseteq_f T$. Let $T_0 = \{\varphi_1, \dots, \varphi_n\}$. For each $j \leq n$, we have $\varphi_j \in T = \bigcup_i T_i$, so there is some $i_j \in I$ with $\varphi_j \in T_{i_j}$. Let $\ell = \max(i_1, \dots, i_n) \in I$. Then for each $j \leq n$, we have $\varphi_j \in T_{i_j} \subseteq T_\ell$. Thus $T_0 \subseteq_f T_\ell$, and T_0 is satisfiable because T_ℓ is finitely satisfiable. \square

Lemma 4.4.2. If T is a finitely satisfiable \mathcal{L} -theory, then there is a complete, finitely satisfiable \mathcal{L} -theory $T' \supseteq T$.

Proof. By Zorn's lemma and Theorem 4.4.1 there is a $T' \supseteq T$ which is maximal among finitely satisfiable \mathcal{L} -theories. We claim T' is complete. Otherwise

there is a sentence φ with $\varphi \notin T'$ and $\neg\varphi \notin T'$. By maximality, $T' \cup \{\varphi\}$ is not finitely satisfiable, so there is $T_1 \subseteq_f T$ with $T_1 \models \neg\varphi$. Similarly, $\neg\varphi \notin T'$ implies that there is $T_2 \subseteq_f T$ with $T_2 \models \varphi$. Then $T_1 \cup T_2$ is not satisfiable, a contradiction. \square

Lemma 4.4.3. *Let T be a finitely satisfiable \mathcal{L} -theory containing a sentence $\exists x \varphi(x)$. Let $\mathcal{L}' = \mathcal{L} \cup \{c\}$ where c is a new constant symbol. Then $T \cup \{\varphi(c)\}$ is a finitely satisfiable \mathcal{L}' -theory.*

Proof. Otherwise, $T_0 \cup \{\varphi(c)\}$ is unsatisfiable for some $T_0 \subseteq_f T$. Take $M \models T_0 \cup \{\exists x \varphi(x)\}$ and take $b \in M$ such that $M \models \varphi(b)$. Expand the \mathcal{L} -structure M to an \mathcal{L}' -structure by interpreting c as b . Then $M \models T_0 \cup \{\varphi(c)\}$, contradicting the choice of T_0 . \square

Lemma 4.4.4. *Let T be a finitely satisfiable \mathcal{L} -theory. There is a language $\mathcal{L}' \supseteq \mathcal{L}$ and an \mathcal{L}' -theory $T' \supseteq T$ that is finitely satisfiable and complete, and has the witness property.*

Proof. Build increasing chains

$$\begin{aligned} \mathcal{L}_0 &\subseteq \mathcal{L}_1 \subseteq \cdots \\ T_0 &\subseteq T_1 \subseteq \cdots \end{aligned}$$

where T_i is a finitely satisfiable \mathcal{L}_i -theory as follows:

1. $\mathcal{L}_0 = \mathcal{L}$ and $T_0 = T$.
2. If $n > 0$ and n is odd, then $\mathcal{L}_n = \mathcal{L}_{n-1}$ and T_n is a completion of T_{n-1} from Lemma 4.4.2.
3. If $n > 0$ and n is even, let $\{\varphi_i(x) : i \in I\}$ enumerate the \mathcal{L}_{n-1} -formulas such that $(\exists x \varphi(x)) \in T_{n-1}$. Let $\mathcal{L}_n = \mathcal{L}_{n-1} \cup \{c_i : i \in I\}$ where the c_i are new constant symbols. Let $T_n = T_{n-1} \cup \{\varphi_i(c_i) : i \in I\}$. Then T_n is finitely satisfiable by Lemma 4.4.3.

Finally, take $\mathcal{L}' = \bigcup_n \mathcal{L}_n$ and $T' = \bigcup_n T_n$. Then T' is finitely satisfiable because each T_i is, T' is complete because of the odd-numbered steps, and T' has the witness property because of the even-numbered steps. \square

Theorem 4.4.5 (Compactness). *If T is finitely satisfiable, then T has a model.*

Proof. By Lemma 4.4.4, there is a language $\mathcal{L}' \supseteq \mathcal{L}$ and an \mathcal{L}' -theory $T' \supseteq T$ such that T' is finitely satisfiable and complete and has the witness property. By Theorem 4.3.4 there is an \mathcal{L}' -structure M satisfying T' , and therefore T . \square

Chapter 5

Categoricity

With the compactness theorem proved, we now turn towards applications. The key result of this chapter is the *Löwenheim-Skolem theorem*, which in one form says the following:

Theorem. *Let T be a theory in a countable language. Suppose T has an infinite model M . Then for any infinite cardinal κ , T has a model M_κ of size κ .*

On one hand, this gives us a source of new models. For example, it tells us that there are fields of size κ for any infinite κ , because there is at least one infinite field (\mathbb{R}).

On the other hand, the Löwenheim-Skolem theorem shows the limits of what can be expressed using first-order logic. In particular, first-order sentences have no way of saying anything about infinite cardinalities—there is no way to say “ M is countable” or “ M is uncountable,” because any theory with a countable infinite model will have uncountable models, any theory with an uncountable model will have a countable infinite model.

We apply the Löwenheim-Skolem theorem to the notion of categoricity. A theory T is *absolutely categorical* if it has exactly one model, up to isomorphism. Absolutely categorical theories are complete, because any two models are elementarily equivalent. Unfortunately, the Löwenheim-Skolem theorem makes it nearly impossible to be absolutely categorical (see Theorem 5.5.2).

However, the more refined notion of κ -*categoricity* works well. If κ is an infinite cardinal, then a theory T is κ -*categorical* if it has exactly one model of size κ . Using the Löwenheim-Skolem theorem, one can show that

κ -categorical theories are complete, modulo a few small issues (see Theorem 5.6.2). Several important theories turn out to be κ -categorical for various values of κ .

In the process of studying categoricity, we will also meet some important technical tools. The *diagram* of a structure M is a theory whose models are essentially the extensions of M (the structures extending M). There is also a related theory called the *elementary diagram* of M , whose models are essentially the elementary extensions of M . Diagrams are a common tool in applications of the compactness theorem. Using elementary diagrams, we also prove a fact called *elementary amalgamation*, which says that if $M_1 \equiv M_2$, then up to isomorphism the two structures M_1 and M_2 are both elementary substructures of a third structure M_3 . We will see later in Chapter 11 that for any structure M one can in fact make a big elementary extension $\mathbb{M} \succeq M$ such that \mathbb{M} is an elementary extension of every “small” $N \equiv M$.

5.1 A useful fact around finite satisfiability

Let T be a set of sentences. Suppose T is closed under finite conjunctions, in the sense that

$$\varphi_1, \dots, \varphi_n \in T \implies \bigwedge_{i=1}^n \varphi_i \in T.$$

Then T is finitely satisfiable if and only if every $\varphi \in T$ is satisfiable, because any finite subset $\{\varphi_1, \dots, \varphi_n\} \subseteq T$ is equivalent to a single formula $\bigwedge_{i=1}^n \varphi_i$ in T .

More generally, suppose T_1, T_2 are two sets of sentences, and T_1 is closed under finite conjunctions. Then the following are equivalent:

- $T_1 \cup T_2$ is finitely satisfiable.
- For every $\varphi \in T_1$ and finite subset $T'_2 \subseteq_f T_2$, the theory $\{\varphi\} \cup \{T'_2\}$ is satisfiable.

We will use this fact implicitly in what follows.

5.2 Diagrams and elementary amalgamation

Let M be an \mathcal{L} -structure and A be a subset of M . The language $\mathcal{L}(A)$ is obtained by adding each element of A as a new constant symbol. We regard

M as an $\mathcal{L}(A)$ -structure by interpreting each new constant symbol as the corresponding element of A :

$$a^M = a.$$

Definition 5.2.1. The *elementary diagram* of M , written $\text{eldiag}(M)$, is the set of $\mathcal{L}(M)$ -sentences φ such that $M \models \varphi$. The *diagram* of M is the set of atomic and negated atomic formulas in $\text{eldiag}(M)$.

A model of $\text{diag}(M)$ consists of an \mathcal{L} -structure N and a map

$$\begin{aligned} M &\rightarrow N \\ a &\mapsto a^N \end{aligned}$$

such that for any atomic \mathcal{L} -formula $\varphi(x_1, \dots, x_n)$ and any $\bar{a} \in M^n$,

$$\begin{aligned} M \models \varphi(a_1, \dots, a_n) &\implies N \models \varphi(a_1^N, \dots, a_n^N) \\ M \models \neg\varphi(a_1, \dots, a_n) &\implies N \models \neg\varphi(a_1^N, \dots, a_n^N). \end{aligned}$$

This means that $a \mapsto a^N$ is an embedding.

Therefore, a model of $\text{diag}(M)$ is the same thing as an \mathcal{L} -structure N with an embedding $f : M \rightarrow N$. In particular, if N is an extension of M , then N is a model of $\text{diag}(M)$ in a natural way, interpreting $a \in M$ as $a \in N$. Every model of $\text{diag}(M)$ is isomorphic as an $\mathcal{L}(M)$ -structure to an extension of M .

Similarly, a model of $\text{eldiag}(M)$ is the same thing as an \mathcal{L} -structure N with an elementary embedding $f : M \rightarrow N$. Any elementary extension of M is a model of $\text{eldiag}(M)$, and every model of $\text{eldiag}(M)$ is isomorphic as an $\mathcal{L}(M)$ -structure to an elementary extension of M .

Theorem 5.2.2 (Elementary amalgamation). *If M_1 and M_2 are elementarily equivalent, then there is a structure N and elementary embeddings $M_1 \rightarrow N$ and $M_2 \rightarrow N$.*

We first prove a lemma. Recall the notion of *expansion* from Definition 3.1.5.

Lemma 5.2.3. *If $M_1 \equiv M_2$ and $\varphi(\bar{c}) \in \text{eldiag}(M_2)$, then there is an $\mathcal{L}(M_2)$ -structure N expanding the \mathcal{L} -structure M_1 such that $N \models \varphi$.*

Proof. The fact that $M_2 \models \varphi(\bar{c})$ implies that $M_2 \models \exists \bar{x} \varphi(\bar{x})$ and then $M_1 \models \exists \bar{x} \varphi(\bar{x})$. Take \bar{a} in M_1 such that $M_1 \models \varphi(\bar{a})$. Expand M_1 to an $\mathcal{L}(M_2)$ -structure N by interpreting the symbols c_1, \dots, c_n as a_1, \dots, a_n , respectively. Then $N \models \varphi(\bar{c})$ because $\bar{c}^N = \bar{a}$ and $M_1 \models \varphi(\bar{a})$. \square

Now we prove Theorem 5.2.2.

Proof. It suffices to show that $\text{eldiag}(M_1) \cup \text{eldiag}(M_2)$ is consistent. Otherwise, by compactness, there is a tuple \bar{c} in M_2 and formula $\varphi(\bar{c}) \in \text{eldiag}(M_2)$ such that $\text{eldiag}(M_1) \cup \{\varphi(\bar{c})\}$ is inconsistent. But $M_1 \models \text{eldiag}(M_1)$, and Lemma 5.2.3 gives an expansion of M_1 satisfying $\{\varphi(\bar{c})\}$, and so $\text{eldiag}(M_1) \cup \{\varphi(\bar{c})\}$ is consistent. \square

5.3 Tarski-Vaught

Theorem 5.3.1. *Let M be a structure and A be a subset. Then $A \preceq M$ iff the following holds: for every non-empty A -definable $D \subseteq M$, we have $D \cap A \neq \emptyset$.*

This condition is called the *Tarski-Vaught criterion*.

Proof. First suppose $A \preceq M$. Suppose D is A -definable and non-empty. Write D as $\varphi(M, \bar{a})$ for some $\mathcal{L}(A)$ -formula $\varphi(x, \bar{a})$. Then

$$M \models \exists x : \varphi(x, \bar{a}) \implies A \models \exists x : \varphi(x, \bar{a})$$

so there is $b \in A$ with $A \models \varphi(b, \bar{a})$, which implies $M \models \varphi(b, \bar{a})$ as $A \preceq M$. Then $b \in A$ and $b \in D = \varphi(M, \bar{a})$.

Conversely, suppose the Tarski-Vaught criterion holds. First we show that A is a substructure. Suppose f is a k -ary function symbol and $a_1, \dots, a_k \in A$. The set $\{f(\bar{a})\}$ is non-empty and A -definable, so A intersects it, meaning that $f(\bar{a}) \in A$.

Next we show that for any \mathcal{L} -formula $\varphi(\bar{x})$ and any $\bar{a} \in A$,

$$M \models \varphi(\bar{a}) \iff A \models \varphi(\bar{a}). \quad (*)$$

We may assume φ makes no use of $\wedge, \forall, \top, \perp$. Proceed by induction on the complexity of φ . If φ is atomic or more generally quantifier-free, then $(*)$ holds because A is a substructure.

If φ is $\psi \vee \theta$, then

$$\begin{aligned} M \models \varphi(\bar{a}) &\iff M \models \psi(\bar{a}) \text{ or } M \models \theta(\bar{a}) \\ &\iff A \models \psi(\bar{a}) \text{ or } A \models \theta(\bar{a}) \\ &\iff A \models \varphi(\bar{a}) \end{aligned}$$

by induction. Finally, if $\varphi(\bar{x})$ is $\exists y : \psi(\bar{x}, y)$, then

$$\begin{aligned} M \models \varphi(\bar{a}) &\iff \exists b \in M : M \models \psi(\bar{a}, b) \\ &\stackrel{\text{TV}}{\iff} \exists b \in A : M \models \psi(\bar{a}, b) \\ &\stackrel{\text{ind}}{\iff} \exists b \in A : A \models \psi(\bar{a}, b) \\ &\iff A \models \varphi(\bar{a}). \end{aligned}$$

The second line uses the Tarski-Vaught criterion, and the third line uses induction. \square

5.4 The Löwenheim-Skolem theorem

Definition 5.4.1. If \mathcal{L} is a language, the *size* of \mathcal{L} , written $|\mathcal{L}|$, is \aleph_0 plus the number of symbols in \mathcal{L} .

If \bar{x} is a finite tuple of variables, then $|\mathcal{L}|$ is the number of \mathcal{L} -formulas $\varphi(\bar{x})$.

Remark 5.4.2. If M is a \mathcal{L} -structure and $A \subseteq M$, then the number of A -definable sets in M is at most $|A| + |\mathcal{L}|$.

Theorem 5.4.3 (Downward Löwenheim-Skolem theorem). *Let M be an \mathcal{L} -structure.*

1. *If $A \subseteq M$, there is an elementary substructure $N \preceq M$ with $N \supseteq A$ and $|N| \leq |A| + |\mathcal{L}|$.*
2. *For any κ with $|\mathcal{L}| \leq \kappa \leq |M|$, there is $N \preceq M$ with $|N| = \kappa$.*

Proof. 1. Let $F : \mathfrak{P}(M) \setminus \{\emptyset\} \rightarrow M$ be a function such that $F(X) \in X$. Recursively define $A_0 \subseteq A_1 \subseteq \dots$ by letting $A_0 = A$ and

$$A_{n+1} = A_n \cup \{F(X) : X \subseteq M \text{ is } A_n\text{-definable and non-empty}\}.$$

Let $N = \bigcup_{n=0}^{\infty} A_n$. Then $A = A_0 \subseteq N$. By induction on n , each A_n has size at most $|A| + |\mathcal{L}|$, and therefore $|N| \leq |A| + |\mathcal{L}|$. If X is N -definable and non-empty, then X is A_n -definable for some n , so $F(X) \in A_{n+1} \subseteq N$. Thus N intersects every N -definable set, and $N \preceq M$ by Tarski-Vaught.

2. Take a subset $A \subseteq M$ with $|A| = \kappa$, and take $N \preceq M$ as in part (1). Then $|N| \leq |A| + |\mathcal{L}| = \kappa + |\mathcal{L}| = \kappa$. On the other hand, $|N| \geq \kappa$ because $N \supseteq A$. \square

Theorem 5.4.4 (Löwenheim-Skolem theorem). *Let T be an \mathcal{L} -theory. Suppose T has an infinite model, or more generally that for every $n < \omega$, T has a model of size at least n . Then for any $\kappa \geq |\mathcal{L}|$, T has a model of size κ .*

Proof. Let $\mathcal{L}' = \mathcal{L} \cup \{c_\alpha : \alpha < \kappa\}$, where the c_α are new constant symbols. Let $\Sigma = \{(c_\alpha \neq c_\beta) : \alpha < \beta < \kappa\}$.

Claim. $T \cup \Sigma$ is finitely satisfiable.

Proof. Suppose $\Sigma_0 \subseteq_f \Sigma$. Let S be the finite set of $\alpha \in \kappa$ such that c_α appears in Σ_0 . Take a model $M \models T$ with $|M| > |S|$. Expand M to an \mathcal{L}' -structure by interpreting the c_α for $\alpha \in S$ as distinct elements of M , and interpreting c_α for $\alpha \in \kappa \setminus S$ arbitrarily. Then $M \models T \cup \Sigma_0$. \square_{Claim}

By compactness, $T \cup \Sigma$ has a model M . Then the c_α^M are pairwise distinct, so $|M| \geq \kappa$. By downward Löwenheim-Skolem (Theorem 5.4.3), there is an elementary substructure $N \preceq M$ with $|N| = \kappa$. Then $N \equiv M$ and $M \models T$, so $N \models T$. \square

Corollary 5.4.5 (Löwenheim-Skolem theorem, second version). *Let M be an infinite \mathcal{L} -structure. For any $\kappa \geq |\mathcal{L}|$, there is $N \equiv M$ with $|N| = \kappa$.*

Proof. Apply Theorem 5.4.4 to the \mathcal{L} -theory $\text{Th}(M)$. Models of $\text{Th}(M)$ are elementarily equivalent to M (Theorem 3.4.1). \square

Example 5.4.6. Earlier we saw that Peano Arithmetic has models which are not elementarily equivalent to \mathbb{N} (Corollary 3.9.13). Using the Löwenheim-Skolem theorem, we can also produce models that are elementarily equivalent to \mathbb{N} but not isomorphic to it. For example, Corollary 5.4.5 gives $M \equiv \mathbb{N}$ with $|M| > \aleph_0$. Then $M \models \text{PA}$ but $M \not\cong \mathbb{N}$.

Theorem 5.4.7 (Upward Löwenheim-Skolem theorem). *If M is an infinite \mathcal{L} -structure and $\kappa \geq |M| + |\mathcal{L}|$, then there is an elementary extension $N \succeq M$ with $|N| = \kappa$.*

Proof. As $\kappa \geq |M| + |\mathcal{L}| = |\mathcal{L}(M)|$, we can apply the Löwenheim-Skolem theorem (Theorem 5.4.4) to the $\mathcal{L}(M)$ -theory $\text{eldiag}(M)$ to get a model $N \models \text{eldiag}(M)$ with $|N| = \kappa$. Moving N by an isomorphism, we may assume $N \succeq M$. \square

5.5 Absolute categoricity

Definition 5.5.1. A theory T is *absolutely categorical* if there is a unique model of T , up to isomorphism.

Theorem 5.5.2. *If T is absolutely categorical, then the unique model M is finite.*

Proof. Take $\kappa > |M| + |\mathcal{L}|$. If M is infinite, then there is a model $N \models T$ with $|N| = \kappa$ by Theorem 5.4.4. Then M and N are two non-isomorphic models, contradicting categoricity. \square

Conversely, if M is finite, then $\text{Th}(M)$ is absolutely categorical. Recall that models of $\text{Th}(M)$ are the same thing as structures elementarily equivalent to M (Theorem 3.4.1).

Theorem 5.5.3. *If M is finite and $M \equiv N$, then $M \cong N$.*

Proof. By elementary amalgamation (Theorem 5.2.2) there are elementary embeddings $M \rightarrow M'$ and $N \rightarrow M'$ for some structure M' . Moving M and N by isomorphisms, we may assume $M \preceq M'$ and $N \preceq M'$. Let n be the size of M . By Lemma 5.5.4 below, $|M| = |M'| = |N| = n$. As M and N are subsets of M' , we must have $M = M' = N$. \square

Lemma 5.5.4. *Suppose $M \equiv N$. If $|M| = n < \infty$, then $|N| = n$. Consequently, M is finite iff N is finite.*

Proof. Suppose $|M| = n$. Let φ be the sentence $\exists^{=n}x \top$ (see Section 3.3). Then φ says that there are exactly n elements, so $M \models \varphi$. Therefore $N \models \varphi$ and $|N| = n$. \square

5.6 κ -categoricity

Definition 5.6.1. Let κ be an infinite cardinal. A theory T is κ -categorical if there is a unique model of cardinality κ .

Theorem 5.6.2 (Łoś-Vaught criterion). *Suppose T is κ -categorical for some $\kappa \geq |\mathcal{L}|$.*

1. *Any two infinite models of T are elementarily equivalent.*
2. *If all models of T are infinite, then T is complete.*

Proof. 1. Given infinite models M_1, M_2 , use Löwenheim-Skolem (Corollary 5.4.5) to get $N_i \equiv M_i$ with $|N_i| = \kappa$. By κ -categoricity, $N_1 \cong N_2$. Then $M_1 \equiv N_1 \equiv N_2 \equiv M_2$.

2. Clear. □

Lemma 5.6.3. *Let M, N be countable structures and let \mathcal{F} be a back-and-forth system between M and N . If $f_0 \in \mathcal{F}$, then there is an isomorphism $f : M \rightarrow N$ extending f_0 .*

Proof. Let a_1, a_2, \dots be an enumeration of M and b_1, b_2, \dots be an enumeration of N . Recursively build an increasing sequence of partial isomorphisms $f_i : A_i \rightarrow B_i$ in \mathcal{F} as follows:

- For $i = 0$, let $f_0 : A_0 \rightarrow B_0$ be the given partial isomorphism.
- For $i = 2k - 1$, take f_i to be an extension of f_{i-1} with $a_k \in \text{dom}(f_i)$. Such an f_i exists by the “forward” condition.
- For $i = 2k$, take f_i to be an extension of f_{i-1} with $b_k \in \text{im}(f_i)$. Such an f_i exists by the “backward” condition.

Take $f = \bigcup_{i=0}^{\infty} f_i$. Then f is an isomorphism from M to N . □

Theorem 5.6.4. *Let M, N be countable structures. If there is a non-empty back-and-forth system between M and N , then $M \cong N$.*

Recall the theory DLO from Definition 3.7.9. Theorem 3.7.10 gives a non-empty back-and-forth system between any two models of DLO.

Corollary 5.6.5 (Cantor). *DLO is \aleph_0 -categorical: any two countable models of DLO are isomorphic.*

Any model of DLO is infinite, so the Łoś-Vaught criterion shows that DLO is complete, as we saw earlier (Corollary 3.7.11).

Theorem 5.6.6. *DLO is not κ -categorical for $\kappa > \aleph_0$.*

Proof. If A, B are linear orders, let $A \times B$ denote the lexicographic product, the set $A \times B$ with the lexicographic order

$$\begin{aligned} (a, b) < (a', b') &\iff a < a' \text{ for } a \neq a' \\ (a, b) < (a, b') &\iff b < b'. \end{aligned}$$

Also, let A^* denote A with the reversed order.

Let $M = \kappa \times \mathbb{Q}$. It is straightforward to see that M and M^* are models of DLO, of size κ . We claim that $M \not\cong M^*$. Note that for any $a \in M$,

$$\begin{aligned} |\{x \in M : x < a\}| &< \kappa \\ |\{x \in M : x > a\}| &= \kappa. \end{aligned}$$

In M^* , the reverse properties hold, so $M \not\cong M^*$, and DLO is not κ -categorical. \square

Chapter 6

Ultraproducts

In the context of universal algebra, we had a number of ways to take models of an equational theory T and produce new models, as discussed in Chapter 2. Is there anything like this in the more general setting of first-order logic? If T is a first-order theory, and we have some models of T , can we combine them to produce new models?

We have already seen a few vague constructions of this form. For example, the upward and downward Löwenheim-Skolem theorems let us take a model M and produce elementary extensions and elementary substructures of a chosen cardinality. But this is not a very close analogue to products, subalgebras, and quotients in universal algebra.

In this chapter, we introduce the *ultraproduct* construction, which gives a much more precise, controlled way to build new models of T from old models. Let I be a set and let M_i be a model of T for each $i \in I$. To define the ultraproduct, we need one additional piece of data, a mysterious object called an *ultrafilter on I* , which is best understood as a boolean algebra homomorphism $\mathfrak{P}(I) \rightarrow \{\text{FALSE}, \text{TRUE}\}$. Given an ultrafilter \mathcal{U} on I , one can define the ultraproduct $\prod_{i \in I}^{\mathcal{U}} M_i$, which is a new model of T . The first-order properties of $\prod_{i \in I}^{\mathcal{U}} M_i$ are determined in a precise way by \mathcal{U} and the first-order properties of the M_i , a fact called *Łoś's theorem*. This can be used to produce models having desired properties. For example, we use it to give a new proof of the compactness theorem (see Theorem 6.2.6).

The actual construction of ultraproducts is a combination of products and quotients. In fact, when T is an equational class (in a functional language), the ultraproduct $\prod_{i \in I}^{\mathcal{U}} M_i$ is literally a quotient of the product $\prod_{i \in I} M_i$.

There is also an analogue of Birkhoff's HSP theorem.

Theorem (Keisler, Shelah). *Let \mathcal{K} be a class of structures. Then \mathcal{K} is an elementary class (axiomatized by a first-order theory) if and only if the following hold:*

1. \mathcal{K} is closed under isomorphisms.
2. \mathcal{K} is closed under ultraproducts: if I is a set, $M_i \in \mathcal{K}$ for $i \in I$, and \mathcal{U} is an ultrafilter on I , then the ultraproduct $\prod_{i \in I}^{\mathcal{U}} M_i$ is in \mathcal{K} .
3. \mathcal{K} is closed under “ultraroots”: if I is a set, \mathcal{U} is an ultrafilter on I , M is a structure, and the “ultrapower” $M^{\mathcal{U}} := \prod_{i \in I}^{\mathcal{U}} M$ is in \mathcal{K} , then M itself is in \mathcal{K} .

The proof of this theorem is sadly too technical to discuss here, but we do prove the following related result:

Theorem. *Let \mathcal{K} be a class of structures. Then \mathcal{K} is axiomatized by a universal theory if and only if \mathcal{K} is closed under isomorphisms, substructures, and ultraproducts.*

Here, a *universal sentence* is a sentence of the form $\forall \bar{x} \varphi(\bar{x})$ with φ being quantifier-free, and a *universal theory* is a set of universal sentences. For example, equational theories are essentially universal theories.

6.1 Ultrafilters

Let I be a set.

Definition 6.1.1. A *filter* on I is a set $\mathcal{F} \subseteq \mathfrak{P}(I)$ satisfying the following:

1. If $X, Y \in \mathcal{F}$, then $X \cap Y \in \mathcal{F}$.
2. If $X \subseteq Y \subseteq I$ and $X \in \mathcal{F}$, then $Y \in \mathcal{F}$.
3. $I \in \mathcal{F}$.

A filter is *proper* if $\emptyset \notin \mathcal{F}$.

Definition 6.1.2. A family of sets $\mathcal{S} \subseteq \mathfrak{P}(I)$ has the *finite intersection property* (FIP) if for any $n \geq 0$ and $X_1, \dots, X_n \in \mathcal{S}$, we have $\bigcap_{i=1}^n X_i \neq \emptyset$.

(When $n = 0$, $\bigcap_{i=1}^n X_i$ is defined to be I .)

Remark 6.1.3. If \mathcal{F} is a proper filter, then \mathcal{F} has the FIP, because $\emptyset \notin \mathcal{F}$, and \mathcal{F} is closed under intersection.

Lemma 6.1.4. *If $\mathcal{S} \subseteq \mathfrak{P}(I)$ has the FIP, then there is a proper filter $\mathcal{F} \supseteq \mathcal{S}$.*

Proof. Let \mathcal{F} be the set of $X \subseteq I$ such that there are $n \geq 0$ and $Y_1, \dots, Y_n \in \mathcal{S}$ with $X \supseteq \bigcap_{i=1}^n Y_i$. Then \mathcal{F} is a proper filter containing \mathcal{S} . \square

Definition 6.1.5. An *ultrafilter* on I is a proper filter \mathcal{U} such that for any $X \subseteq I$,

$$X \in \mathcal{U} \text{ or } I \setminus X \in \mathcal{U}.$$

Lemma 6.1.6. *If \mathcal{F} is a proper filter on I , then there is an ultrafilter $\mathcal{U} \supseteq \mathcal{F}$.*

Proof. By Zorn's lemma, there is a maximal proper filter $\mathcal{U} \supseteq \mathcal{F}$. We claim that \mathcal{U} is an ultrafilter. Otherwise, there is $X \subseteq I$ with $X \notin \mathcal{U}$ and $I \setminus X \notin \mathcal{U}$. By maximality, $\mathcal{U} \cup \{X\}$ is not contained in a filter, and does not have FIP. Therefore there are $Y_1, \dots, Y_n \in \mathcal{U}$ such that $X \cap \bigcap_{i=1}^n Y_i = \emptyset$. As \mathcal{U} is a filter, \mathcal{U} contains $Y := \bigcap_{i=1}^n Y_i$. Then $X \cap Y = \emptyset$, so $Y \subseteq I \setminus X$.

Applying the same argument to the complement of X , we get $Z \in \mathcal{U}$ with $Z \subseteq X$. Then $Y \cap Z \in \mathcal{U}$ because \mathcal{U} is a filter. However, $Y \cap Z \subseteq (I \setminus X) \cap X = \emptyset$, so $\emptyset \in \mathcal{U}$ contradicting the fact that \mathcal{U} is proper. \square

Theorem 6.1.7. *If $\mathcal{S} \subseteq \mathfrak{P}(I)$ has the FIP, then \mathcal{S} is contained in an ultrafilter \mathcal{U} on I .*

Fact 6.1.8. *Let \mathcal{U} be an ultrafilter on I . Let $\chi : \mathfrak{P}(I) \rightarrow \{0, 1\}$ be the characteristic function of \mathcal{U} :*

$$\chi(X) = \begin{cases} 1 & X \in \mathcal{U} \\ 0 & X \notin \mathcal{U}. \end{cases}$$

Then χ is a homomorphism of boolean algebras. In fact, ultrafilters on I correspond bijectively to homomorphisms $\mathfrak{P}(I) \rightarrow \{0, 1\}$.

6.2 Ultraproducts

Let I be a set and let \mathcal{U} be an ultrafilter on I . Let M_i be a *non-empty* \mathcal{L} -structure for each $i \in I$. Let $P = \prod_{i \in I} M_i$ and let $\pi_i : P \rightarrow M_i$ be the i th coordinate projection.

Let $\mathcal{L}(P)$ be \mathcal{L} with a new constant symbol added for each $a \in P$. Regard M_i as an $\mathcal{L}(P)$ -structure by interpreting $a \in P$ as $\pi_i(a) \in M_i$.

For any $\mathcal{L}(P)$ -sentence φ , let $\llbracket \varphi \rrbracket = \{i \in I : M_i \models \varphi\}$. Note the following:

$$\begin{aligned}\llbracket \varphi \wedge \psi \rrbracket &= \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket \\ \llbracket \varphi \vee \psi \rrbracket &= \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket \\ \llbracket \neg \varphi \rrbracket &= I \setminus \llbracket \varphi \rrbracket\end{aligned}$$

Lemma 6.2.1. *Let T be the set of $\mathcal{L}(P)$ -sentences φ such that $\llbracket \varphi \rrbracket \in \mathcal{U}$. Then T is finitely satisfiable and complete and has the witness property over P in the sense of Definition 4.3.6.*

Proof. For finite satisfiability, suppose $\varphi_1, \dots, \varphi_n \in T$. If $S = \llbracket \bigwedge_{i=1}^n \varphi_i \rrbracket = \bigcap_{i=1}^n \llbracket \varphi_i \rrbracket$, then $S \in \mathcal{U}$ so $S \neq \emptyset$. Taking $i \in S$, we have $M_i \models \bigwedge_{i=1}^n \varphi_i$.

For completeness, note that $\llbracket \varphi \rrbracket$ and $\llbracket \neg \varphi \rrbracket$ are complementary, so one of them is in \mathcal{U} , and therefore one of φ and $\neg \varphi$ is in T .

For the witness property, suppose $\exists x \varphi(x)$ is in T , meaning that $S := \llbracket \exists x \varphi(x) \rrbracket \in \mathcal{U}$. Define $c = (c_i : i \in I) \in P$ as follows:

1. If $i \in S$, then $M_i \models \exists x \varphi(x)$. Choose $c_i \in M_i$ so that $M_i \models \varphi(c_i)$.
2. If $i \notin S$, take any $c_i \in M_i$.

Then $i \in S \implies M_i \models \varphi(c)$, so $\llbracket \varphi(c) \rrbracket \supseteq S$. It follows that $\llbracket \varphi(c) \rrbracket \in \mathcal{U}$ and $\varphi(c) \in T$. \square

By Definition 4.3.5 and Theorem 4.3.7, we can talk about the canonical model of T —the unique model in which every element is named by an element of P .

Definition 6.2.2. The *ultraproduct* $\prod_{i \in I}^{\mathcal{U}} M_i$ is the canonical model of T .

Theorem 6.2.3. *Let $P = \prod_{i \in I} M_i$, let $M = \prod_{i \in I}^{\mathcal{U}} M_i$, and let $[c]$ be the element of M named by $c \in P$.*

1. *Every element of M has the form $[c]$ for some $c \in P$.*

2. $[c] = [d]$ if and only if $\{i \in I : \pi_i(c) = \pi_i(d)\} \in \mathcal{U}$.

3. Let R be a k -ary relation symbol. Then

$$R^M([c_1], \dots, [c_k]) \iff \{i \in I : M_i \models R(\pi_i(c_1), \dots, \pi_i(c_k))\} \in \mathcal{U}.$$

4. If f is a k -ary function symbol, then

$$f^M([c_1], \dots, [c_k]) = [f^P(c_1, \dots, c_k)],$$

where P is given the product structure so that

$$f^P(c_1, \dots, c_k) = (f^{M_i}(\pi_i(c_1), \dots, \pi_i(c_k)) : i \in I).$$

5. If $\varphi(x_1, \dots, x_n)$ is an \mathcal{L} -formula, then

$$M \models \varphi([c_1], \dots, [c_k]) \iff \{i \in I : M_i \models \varphi(\pi_i(c_1), \dots, \pi_i(c_k))\} \in \mathcal{U}.$$

6. If φ is an \mathcal{L} -sentence, then

$$M \models \varphi \iff \{i \in I : M_i \models \varphi\} \in \mathcal{U}.$$

Proof. Part (1) is true by construction, as M is the canonical model of T . Part (5) is also true by construction:

$$\begin{aligned} M \models \varphi([c_1], \dots, [c_k]) &\iff M \models \varphi(c_1, \dots, c_k) \iff \varphi(c_1, \dots, c_k) \in T \\ &\iff \llbracket \varphi(c_1, \dots, c_k) \rrbracket \in \mathcal{U} \\ &\iff \{i \in I : M_i \models \varphi(\pi_i(c_1), \dots, \pi_i(c_k))\} \in \mathcal{U}. \end{aligned}$$

Remember that a constant symbol $c \in P$ is interpreted as $[c]$ in M , and $\pi_i(c)$ in M_i .

Parts (2) and (3) follow by specializing part (5) to atomic formulas $x = y$ and $R(x_1, \dots, x_k)$. Part (6) is the $n = 0$ case of part (5). It remains to prove part (4). Fix $c_1, \dots, c_k \in P$ and let $d = f^P(c_1, \dots, c_k)$. Then $\pi_i(d) = f^{M_i}(\pi_i(c_1), \dots, \pi_i(c_k))$ for each $i \in I$. Then

$$\{i \in I : M_i \models \pi_i(d) = f(\pi_i(c_1), \dots, \pi_i(c_k))\} = I \in \mathcal{U},$$

so by part (5), we have

$$M \models [d] = f([c_1], \dots, [c_k]),$$

which means that $f^M([c_1], \dots, [c_k]) = [d] = [f^P(c_1, \dots, c_k)]$. \square

Remark 6.2.4. Parts (1)—(4) determine the structure of M up to isomorphism, and are usually taken as the *definition* of the ultraproduct. Then part (5) is called *Łoś's Theorem*.

Theorem 6.2.5. *Let I be a set, \mathcal{U} be an ultrafilter on I , M_i be an \mathcal{L} -structure for $i \in I$, and M be the ultraproduct $\prod_{i \in I}^{\mathcal{U}} M_i$.*

1. *Let T be a theory. If $M_i \models T$ for all i , then $M \models T$.*
2. *Let \mathcal{K} be an elementary class. If $M_i \in \mathcal{K}$ for all i , then $M \in \mathcal{K}$.*

Proof. (1) follows from Łoś's theorem: if $\varphi \in T$, then

$$\{i \in I : M_i \models \varphi\} = I \in \mathcal{U},$$

so $M \models \varphi$. (2) is a restatement of (1). □

If \mathcal{K} is a class of \mathcal{L} -structures and T is an \mathcal{L} -theory, say that T is *finitely satisfiable in \mathcal{K}* if for any finite subtheory $T_0 \subseteq_f T$ there is $M \in \mathcal{K}$ satisfying T_0 .

Theorem 6.2.6. *If T is finitely satisfiable in \mathcal{K} , then there is an ultraproduct M of structures in \mathcal{K} such that $M \models T$.*

Proof. Let $\{M_i\}_{i \in I}$ be a small collection of structures in \mathcal{K} containing at least one representative from every elementary equivalence class. If φ is an \mathcal{L} -sentence, let $\llbracket \varphi \rrbracket = \{i \in I : M_i \models \varphi\}$. Let $\mathcal{F} = \{\llbracket \varphi \rrbracket : \varphi \in T\}$. We claim that \mathcal{F} has the FIP. Indeed, if $\varphi_1, \dots, \varphi_n \in T$ then there is some M_i satisfying $\bigwedge_{j=1}^n \varphi_j$, and then $i \in \bigcap_{j=1}^n \llbracket \varphi_j \rrbracket$.

Because \mathcal{F} has the FIP, it is contained in an ultrafilter \mathcal{U} on I (Theorem 6.1.7). Let $M = \prod_{i \in I}^{\mathcal{U}} M_i$. Then for $\varphi \in T$ we have

$$\{i \in I : M_i \models \varphi\} = \llbracket \varphi \rrbracket \in \mathcal{F} \subseteq \mathcal{U},$$

and so $M \models \varphi$ by Łoś's theorem. □

Note that this gives another proof of the compactness theorem (Theorem 4.4.5).

Definition 6.2.7. If M is an \mathcal{L} -structure and \mathcal{U} is an ultrafilter on a set I , then the *ultrapower* $M^{\mathcal{U}}$ is the ultraproduct $\prod_{i \in I}^{\mathcal{U}} M_i$.

Theorem 6.2.8. *Let $\Delta : M \rightarrow M^{\mathcal{U}}$ be the diagonal map sending $a \in M$ to the class of the tuple $(a : i \in I) \in M^I = \prod_{i \in I} M$. Then Δ is an elementary embedding.*

Proof. Fix a formula $\varphi(x_1, \dots, x_n)$ and a tuple $\bar{a} = (a_1, \dots, a_n) \in M^n$. Let $S = \{i \in I : M \models \varphi(a_1, \dots, a_n)\}$. By Łoś's theorem,

$$M^{\mathcal{U}} \models \varphi(\Delta(a_1), \dots, \Delta(a_n)) \iff S \in \mathcal{U}.$$

But

$$S = \begin{cases} I & \text{if } M \models \varphi(\bar{a}) \\ \emptyset & \text{if } M \not\models \varphi(\bar{a}). \end{cases}$$

Therefore $S \in \mathcal{U} \iff M \models \varphi(\bar{a})$, and

$$M^{\mathcal{U}} \models \varphi(\Delta(\bar{a})) \iff M \models \varphi(\bar{a}). \quad \square$$

6.3 Characterizations of elementary classes

Lemma 6.3.1. *If M_1 and M_2 are elementarily equivalent, then there is an elementary embedding from M_2 to an ultrapower $M_1^{\mathcal{U}}$.*

Proof. In the proof of elementary amalgamation, we saw that $\text{eldiag}(M_2)$ is finitely satisfiable in expansions of M_1 to $\mathcal{L}(M_2)$ -structures (Lemma 5.2.3). By Theorem 6.2.6, there is an ultraproduct $N = \prod_{i \in I}^{\mathcal{U}} N_i$ satisfying $\text{eldiag}(M_2)$, where each N_i is an expansion of M_1 to an $\mathcal{L}(M_2)$ -structure. The fact that $N \models \text{eldiag}(M_2)$ gives an elementary embedding $M_2 \rightarrow N$. On the other hand, $N \upharpoonright \mathcal{L}$ is

$$\prod_{i \in I}^{\mathcal{U}} (N_i \upharpoonright \mathcal{L}) = \prod_{i \in I}^{\mathcal{U}} M_1 = M_1^{\mathcal{U}},$$

and we have the desired elementary embedding $M_2 \rightarrow M_1^{\mathcal{U}}$. \square

Theorem 6.3.2. *A class \mathcal{K} of \mathcal{L} -structures is elementary iff \mathcal{K} is closed under ultraproducts, isomorphisms, and elementary substructures.*

Proof. Suppose \mathcal{K} is closed under ultraproducts, isomorphisms, and elementary substructures.

Claim. \mathcal{K} is closed under elementary equivalence: if $M \equiv N$ and $M \in \mathcal{K}$ then $N \in \mathcal{K}$.

Proof. By Lemma 6.3.1, there is an ultrapower $M^{\mathcal{U}}$ and an elementary embedding $f : N \rightarrow M^{\mathcal{U}}$. Then $N \cong \text{im}(f) \preceq M^{\mathcal{U}}$, so

$$M \in \mathcal{K} \implies M^{\mathcal{U}} \in \mathcal{K} \implies \text{im}(f) \in \mathcal{K} \implies N \in \mathcal{K}. \quad \square_{\text{Claim}}$$

Let T be the set of all sentences φ such that every $N \in \mathcal{K}$ satisfies φ . Then $\mathcal{K} \subseteq \text{Mod}(T)$. We claim that $\mathcal{K} = \text{Mod}(T)$. Fix $M \in \text{Mod}(T)$; we claim that $M \in \mathcal{K}$. Break into two cases:

1. $\text{Th}(M)$ is finitely satisfiable in \mathcal{K} . By Theorem 6.2.6, there is a model $N \models \text{Th}(M)$ such that N is an ultraproduct of structures in \mathcal{K} . By assumption, $N \in \mathcal{K}$. The fact that $N \models \text{Th}(M)$ means that $N \equiv M$ (Theorem 3.4.1), and so $M \in \mathcal{K}$.
2. $\text{Th}(M)$ is not finitely satisfiable in \mathcal{K} . Then there is $\varphi \in \text{Th}(M)$ such that no $N \in \mathcal{K}$ satisfies φ . Then every $N \in \mathcal{K}$ satisfies $\neg\varphi$, so $\neg\varphi \in T$ and $M \models \neg\varphi$, contradicting the fact that $\varphi \in \text{Th}(M)$. \square

Fact 6.3.3 (Keisler-Shelah). *If $M \equiv N$, then there is a set I and an ultrafilter \mathcal{U} on I such that the ultraproducts $M^{\mathcal{U}}$ and $N^{\mathcal{U}}$ are isomorphic.*

Definition 6.3.4. A structure M is an *ultraroot* of a structure N if N is an ultrapower of M . A class \mathcal{K} is *closed under ultraroots* if $M^{\mathcal{U}} \in \mathcal{K} \implies M \in \mathcal{K}$.

If M is an ultraroot of N , then $N \succeq M$, so $N \equiv M$. Therefore, elementary classes are closed under ultraroots.

Theorem 6.3.5. *A class \mathcal{K} is elementary iff \mathcal{K} is closed under ultraproducts, ultraroots, and isomorphisms.*

Proof. Suppose \mathcal{K} is closed under ultraproducts, ultraroots, and isomorphisms. By the Keisler-Shelah theorem, \mathcal{K} is closed under elementary equivalence. Then the proof of Theorem 6.3.2 applies. \square

6.4 Universal classes

Lemma 6.4.1 (Lemma on constants). *Let $\bar{c} = (c_1, \dots, c_n)$ be a tuple of constant symbols not appearing in T . If $T \vdash \varphi(\bar{c})$, then $T \vdash \forall \bar{x} \varphi(\bar{x})$.*

Proof. Otherwise, there is $M \models T$ with $M \not\models \forall \bar{x} \varphi(\bar{x})$. Take $\bar{a} \in M$ with $M \models \neg \varphi(\bar{a})$. Interpreting \bar{c} as \bar{a} , we get a model of $T \cup \{\neg \varphi(\bar{c})\}$, contradicting $T \vdash \varphi(\bar{c})$. \square

A *universal formula* or \forall -*formula* is one of the form $\forall \bar{y} : \varphi(\bar{x}, \bar{y})$ where φ is quantifier-free. A *universal theory* is a set of universal sentences.

Lemma 6.4.2. *If T is a universal theory, and $M \models T$, then any substructure of M is a model of T .*

Proof. Let ψ be a sentence in T . We can write ψ as $\forall \bar{x} : \varphi(\bar{x})$ where φ is quantifier-free. Let N be a substructure of M . If $\bar{a} \in N$, then

$$M \models \forall \bar{x} : \varphi(\bar{x}) \implies M \models \varphi(\bar{a}) \implies N \models \varphi(\bar{a}),$$

where the second implication holds because $\varphi(\bar{x})$ is quantifier-free (see Remark 3.5.4). As this holds for any $\bar{a} \in N$, we have $N \models \forall \bar{x} : \varphi(\bar{x})$. \square

If T is a theory, then T_\forall denotes the set of universal sentences φ such that $T \vdash \varphi$.

Theorem 6.4.3. *$M \models T_\forall$ if and only if M is a substructure of a model of T .*

Proof. First suppose M is a substructure of $N \models T$. Then $N \models T_\forall$, so $M \models T_\forall$ as T_\forall is a universal theory.

Conversely, suppose $M \models T_\forall$. Break into two cases:

1. The $\mathcal{L}(M)$ -theory $\text{diag}(M) \cup T$ is consistent. Take a model N . Moving N by an isomorphism, we may assume that N is an extension of M , and $N \models T$. Then we are done.
2. $\text{diag}(M) \cup T$ is inconsistent. By compactness, there is a finite tuple $\bar{a} \in M$ and a quantifier-free formula $\varphi(\bar{a}) = \bigwedge_{i=1}^n \varphi_i(\bar{a})$ with $\varphi_i(\bar{a}) \in \text{diag}(M)$ such that $\{\varphi(\bar{a})\} \cup T$ is inconsistent. Then $T \vdash \neg \varphi(\bar{a})$, so $T \vdash \forall \bar{x} \neg \varphi(\bar{x})$ by the lemma on constants (Lemma 6.4.1). But then

$$(\forall \bar{x} \neg \varphi(\bar{x})) \in T_\forall,$$

so $M \models \forall \bar{x} \neg \varphi(\bar{x})$, contradicting the fact that $M \models \varphi(\bar{a})$. \square

Theorem 6.4.4. *Let T be a theory. Then T is axiomatizable by universal sentences if and only if $\text{Mod}(T)$ is closed under substructures.*

Proof. If T is axiomatizable by universal sentences, then $\text{Mod}(T)$ is closed under substructures by Lemma 6.4.2.

Next suppose that $\text{Mod}(T)$ is closed under substructures. We claim that T_{\forall} axiomatizes T . Certainly $M \models T \implies M \models T_{\forall}$. Conversely, suppose $M \models T_{\forall}$. By Theorem 6.4.3 there is an extension $N \supseteq M$ with $N \models T$. By the assumption on T , $M \models T$. \square

Theorem 6.4.5. *Let \mathcal{K} be a class of structures. Then the following are equivalent:*

1. \mathcal{K} is axiomatized by a universal theory.
2. \mathcal{K} is closed under isomorphism, substructures, and ultraproducts.

Proof. (1) \implies (2) is clear.

(2) \implies (1): by Theorem 6.3.2, \mathcal{K} is an elementary class. Then use Theorem 6.4.4. \square

Chapter 7

Topologies

This chapter is a digression into point-set topology, which will play an important role in the chapters ahead. In Section 7.1, we review the basic definitions of topological spaces, continuity, compactness, and Hausdorffness; this section can be skipped if you have ever studied topology. Section 7.2 is about the specialized topic of *ultralimits*, a variant of limits which completes the analogy

metric spaces : limits of sequences :: topological spaces : ultralimits.

(If you've studied point set topology before, ultralimits are similar to nets or filters, though better behaved in some ways.) We use ultralimits to give unusual proofs of some basic facts about compact Hausdorff spaces.

We apply point-set topology to model theory in Section 7.3, where we build a topology on the set S of complete theories (in a fixed language \mathcal{L}). The compactness theorem can be seen as a statement that this topological space is compact, and ultraproducts correspond to ultralimits in this space. The construction of the topological space S works in a more general abstract setting, which we describe in Section 7.4. This abstract generalization will be used later when we consider spaces of types and quantifier-free types in the next chapter.

7.1 Topological spaces

Fix a set S .

Definition 7.1.1. A family of sets $\mathcal{F} \subseteq \mathfrak{P}(S)$ is *closed under infinite unions* if whenever I is a set and $X_i \in \mathcal{F}$ for all $i \in I$, we have $\bigcup_{i \in I} X_i \in \mathcal{F}$. We say that $\mathcal{F} \subseteq \mathfrak{P}(S)$ is *closed under finite unions* if this holds for finite I .

Remark 7.1.2. The definition of “*closed under (in)finite intersections*” is similar, with *finite* print. We understand $\bigcap_{i \in I} X_i$ to mean

$$\{x \in S : \forall i \in I \ x \in X_i\}$$

Consequently, when $I = \emptyset$, the intersection $\bigcap_{i \in I} X_i$ is defined to be S , and not the set-theoretic universe, for example.

Definition 7.1.3. A *metric space* is a set S and function $d : S^2 \rightarrow \mathbb{R}$ satisfying the axioms:

1. $d(x, y) \geq 0$
2. $d(x, y) = d(y, x)$
3. $d(x, y) = 0 \iff x = y$.
4. $d(x, z) \leq d(x, y) + d(y, z)$.

Example 7.1.4. \mathbb{R}^2 is a metric space with respect to the metric

$$d((x_1, y_1), (x_2, y_2)) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}.$$

Fix a metric space (S, d) . An *open ball* is a set of the form

$$B_\epsilon(p) = \{x \in S : d(x, p) < \epsilon\}$$

for some $p \in S$ and $\epsilon > 0$. A set $X \subseteq S$ is *open* if for every $p \in X$, there is $\epsilon > 0$ such that $B_\epsilon(p) \subseteq X$.

Fact 7.1.5. Let (S, d) be a metric space.

1. Every open ball is open.
2. The collection of open sets is closed under infinite unions and finite intersections.

Definition 7.1.6. A *topology* on S is a family of sets τ closed under finite intersections and infinite unions. A *topological space* is a set S with a topology τ .

Fix a topological space (S, τ) .

Definition 7.1.7. A set $X \subseteq S$ is *open* if $X \in \tau$ and *closed* if $S \setminus X$ is open.

Fact 7.1.8. In a metric space, a set X is closed if and only if it “closed under limits” in the sense that:

$$\left(b_1, b_2, \dots \in X \text{ and } \lim_{i \rightarrow \infty} b_i = a \right) \implies a \in X.$$

Definition 7.1.9. A topological space S is *Hausdorff* if for any $a_1 \neq a_2$ in S , there are open sets $U_i \ni a_i$ with $U_1 \cap U_2 = \emptyset$.

Remark 7.1.10. Metric spaces are Hausdorff: if $a_1 \neq a_2$, then $B_\epsilon(a_1) \cap B_\epsilon(a_2) = \emptyset$ for $\epsilon = d(a_1, a_2)/3$.

Theorem 7.1.11. If S is Hausdorff and $p \in S$, then $\{p\}$ is closed.

Proof. For every $q \neq p$, take open sets U_q, V_q with $q \in U_q$, $p \in V_q$, and $U_q \cap V_q = \emptyset$. In particular, $q \in U_q$ and $p \notin U_q$ for each $q \neq p$. Then $S \setminus \{p\}$ is the open set $\bigcup_{q \neq p} U_q$. \square

Definition 7.1.12. An *open cover* is a collection \mathcal{C} of open sets with $\bigcup \mathcal{C} = S$. A *subcover* of \mathcal{C} is another open cover $\mathcal{C}' \subseteq \mathcal{C}$.

Definition 7.1.13. A topological space S is *compact* if every open cover has a finite subcover.

Fact 7.1.14. A metric space (S, d) is compact if and only if every sequence has a convergent subsequence: for any $a_1, a_2, a_3, \dots \in S$, there is a subsequence

$$a_{i_1}, a_{i_2}, a_{i_3}, \dots$$

(with $i_1 < i_2 < \dots$) such that $\lim_{n \rightarrow \infty} a_{i_n}$ exists.

Theorem 7.1.15. If a topological space S is compact and \mathcal{F} is a family of closed sets with the FIP, then $\bigcap \mathcal{F} \neq \emptyset$. Conversely, this property characterizes compactness.

Proof. Let X_i be open for $i \in I$, and let Y_i be the complementary closed sets. We claim that the following are equivalent:

1. If $\bigcup_{i \in I} X_i = S$, then there is $I_0 \subseteq_f I$ with $\bigcup_{i \in I_0} X_i = S$.

2. If $\bigcap_{i \in I} Y_i = \emptyset$, then there is $I_0 \subseteq_f I$ with $\bigcap_{i \in I_0} Y_i = \emptyset$.
3. If $\bigcap_{i \in I_0} Y_i \neq \emptyset$ for every $I_0 \subseteq_f I$, then $\bigcap_{i \in I} Y_i \neq \emptyset$.

Indeed, (1) and (2) are equivalent by de Morgan's laws, and (2) and (3) are contrapositives. Finally, observe that (1) is the definition of compactness, and (3) says that if $\{Y_i\}_{i \in I}$ has FIP then $\bigcap_i Y_i \neq \emptyset$. \square

Definition 7.1.16. Let S_1, S_2 be two topological spaces and $f : S_1 \rightarrow S_2$ be a function. Then f is *continuous* if for every open set $U \subseteq S_2$, the preimage $f^{-1}(U)$ is open in S_1 .

Fact 7.1.17. If S_1, S_2 are metric spaces, a function $f : S_1 \rightarrow S_2$ is continuous iff f “preserves limits”, in the sense that

$$\lim_{i \rightarrow \infty} b_i = a \implies \lim_{i \rightarrow \infty} f(b_i) = f(a).$$

Definition 7.1.18. A function $f : S_1 \rightarrow S_2$ is a *homeomorphism* if f is continuous, f is a bijection, and $f^{-1} : S_2 \rightarrow S_1$ is continuous. Two topological spaces S_1, S_2 are *homeomorphic* if there is a homeomorphism from S_1 to S_2 .

7.2 Ultralimits

Definition 7.2.1. If I is a set, $a_i \in S$ for $i \in I$, \mathcal{U} is an ultrafilter on I , and $b \in S$, then b is an *ultralimit* of the a_i , written

$$b = \lim_{i \rightarrow \mathcal{U}} a_i$$

if for every open set $N \ni b$,

$$\{i \in I : a_i \in N\} \in \mathcal{U}.$$

Fact 7.2.2. If S is Hausdorff then ultralimits are unique: for any $I, \mathcal{U}, \{a_i\}_{i \in I}$, there is at most one b with $b = \lim_{i \rightarrow \mathcal{U}} a_i$. In fact, this property holds if and only if S is Hausdorff.

Half-proof. Suppose S is Hausdorff, and

$$b = \lim_{i \rightarrow \mathcal{U}} a_i$$

$$c = \lim_{i \rightarrow \mathcal{U}} a_i.$$

If $b \neq c$, by Hausdorffness there are open sets $N_1 \ni b$ and $N_2 \ni c$ with $N_1 \cap N_2 = \emptyset$. By definition of ultralimit, the sets

$$\begin{aligned} \{i \in I : a_i \in N_1\} \\ \{i \in I : a_i \in N_2\} \end{aligned}$$

are in the ultrafilter \mathcal{U} . But their intersection is empty, and $\emptyset \notin \mathcal{U}$, a contradiction. \square

Fact 7.2.3. *If S is compact then ultralimits exist: for any $I, \mathcal{U}, \{a_i\}_{i \in I}$, there is at least one b with $b = \lim_{i \rightarrow \mathcal{U}} a_i$. In fact, this property holds if and only if S is compact.*

Half-proof. Suppose S is compact. Say an open set $N \subseteq S$ is *good* if $\{i \in I : a_i \in N\} \in \mathcal{U}$, and *bad* otherwise, i.e., if $\{i \in I : a_i \notin N\} \in \mathcal{U}$. A finite union of bad sets is bad, because \mathcal{U} is closed under finite intersections. The set S is good. There are two cases:

- S is covered by bad open sets. Then compactness gives a finite subcover, so S is a finite union of bad sets and S is bad, a contradiction.
- S is not covered by bad open sets. Take $p \in S$ such that p is in no bad set. Then every open set $N \ni p$ is good, which means $p = \lim_{i \rightarrow \mathcal{U}} a_i$. \square

Theorem 7.2.4. *A set $C \subseteq S$ is closed if and only if C is closed under ultralimits, in the sense that*

$$\left(a_i \in C \text{ for all } i \in I \text{ and } b = \lim_{i \rightarrow \mathcal{U}} a_i \right) \implies b \in C.$$

Proof. First suppose C is closed, and $a_i \in C$ for all $i \in I$. If $b \notin C$, then the complement $S \setminus C$ is an open set containing b , but $\{i \in I : a_i \in S \setminus C\} = \emptyset \notin \mathcal{U}$, contradicting the definition of ultralimits. Thus $b \in C$, and C is closed under ultralimits.

Conversely, suppose C is closed under ultralimits. Let U_0 be the union of open sets disjoint from C . It suffices to show that $C \cup U_0 = S$, as then C is the complement of the open set U_0 . Fix $p \notin U_0$; we claim $p \in C$. Let

$$\mathcal{F} = \{C \cap U : U \text{ is open and } U \ni p\}.$$

Then \mathcal{F} is closed under finite intersections, because a finite intersection of open sets is open. Moreover, $\emptyset \notin \mathcal{F}$, or else $C \cap U = \emptyset$ and U shows $p \in U_0$,

a contradiction. Thus \mathcal{F} has the FIP and is contained in an ultrafilter \mathcal{U} on C . Then

$$p = \lim_{x \rightarrow \mathcal{U}} x$$

because for any open set U containing p ,

$$\{x \in C : x \in U\} = C \cap U \in \mathcal{F} \subseteq \mathcal{U}.$$

By assumption on C , $p \in C$. □

Fact 7.2.5. *If S_1, S_2 are topological spaces and $f : S_1 \rightarrow S_2$ is continuous, then f preserves ultralimits, in the sense that*

$$b = \lim_{i \rightarrow \mathcal{U}} a_i \implies f(b) = \lim_{i \rightarrow \mathcal{U}} f(a_i).$$

In fact, this property holds if and only if f is continuous.

Half-proof. Suppose f is continuous and $b = \lim_{i \rightarrow \mathcal{U}} a_i$. If N is an open set containing $f(b)$, then $f^{-1}(N)$ is an open set containing b , and so

$$\{i \in I : f(a_i) \in N\} = \{i \in I : a_i \in f^{-1}(N)\} \in \mathcal{U}. \quad \square$$

Theorem 7.2.6. *Let S_1, S_2 be compact Hausdorff spaces and let $f : S_1 \rightarrow S_2$ be continuous.*

1. *If $X \subseteq S_1$ is closed, then the image $f(X)$ is closed.*
2. *If f is a bijection, then f is a homeomorphism.*

Proof. 1. By Theorem 7.2.4, it suffices to show that $f(X)$ is closed under ultralimits. Suppose $a_i \in f(X)$ and $\lim_{i \rightarrow \mathcal{U}} a_i = b$. We must show $b \in f(X)$. Write a_i as $f(\alpha_i)$ for some $\alpha_i \in X$. Let

$$\beta = \lim_{i \rightarrow \mathcal{U}} \alpha_i.$$

The ultralimit exists as S_1 is compact (Fact 7.2.3). The ultralimit β is in X because X is closed (Theorem 7.2.4). Then

$$f(\beta) = \lim_{i \rightarrow \mathcal{U}} f(\alpha_i) = \lim_{i \rightarrow \mathcal{U}} a_i$$

because the continuous function f preserves ultralimits (Fact 7.2.5). As S_2 is Hausdorff, ultralimits are unique (Fact 7.2.2), and so $f(\beta) = b$. Then $\beta \in X \implies b = f(\beta) \in f(X)$.

2. Let X be a subset of S_1 . Part (1) shows that

$$X \text{ is closed in } S_1 \implies f(X) \text{ is closed in } S_2.$$

The reverse direction holds by continuity. Thus f is a homeomorphism. \square

7.3 The space of complete theories

Fix a language \mathcal{L} . Let $S = \{\text{Th}(M) : M \text{ is an } \mathcal{L}\text{-structure}\}$. For each \mathcal{L} -sentence φ , let

$$\llbracket \varphi \rrbracket = \{T \in S : \varphi \in T\} = \{\text{Th}(M) : M \models \varphi\}.$$

Note that $M \models \varphi \wedge \psi \iff (M \models \varphi \text{ and } M \models \psi)$, and so

$$\llbracket \varphi \wedge \psi \rrbracket = \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket.$$

Similarly,

$$\begin{aligned} \llbracket \varphi \vee \psi \rrbracket &= \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket \\ \llbracket \neg \varphi \rrbracket &= S \setminus \llbracket \varphi \rrbracket \\ \llbracket \top \rrbracket &= S \\ \llbracket \perp \rrbracket &= \emptyset. \end{aligned}$$

Definition 7.3.1. A set $X \subseteq S$ is *clopen* if $X = \llbracket \varphi \rrbracket$ for some sentence φ , and *open* if $X = \bigcup_{i \in I} Y_i$ where each Y_i is clopen.

Theorem 7.3.2. *The collection of open sets is closed under infinite unions and finite intersections, so it defines a topology on S . Moreover, the topology is Hausdorff.*

Proof. Infinite unions are clear: an infinite union of infinite unions of clopen sets is an infinite union of clopen sets. For finite intersections, note that

$$\left(\bigcup_{i \in I} \llbracket \varphi_i \rrbracket \right) \cap \left(\bigcup_{j \in J} \llbracket \psi_j \rrbracket \right) = \bigcup_{(i,j) \in I \times J} (\llbracket \varphi_i \rrbracket \cap \llbracket \psi_j \rrbracket) = \bigcup_{(i,j) \in I \times J} \llbracket \varphi_i \wedge \psi_j \rrbracket.$$

For Hausdorffness, suppose $T_1, T_2 \in S$ and $T_1 \neq T_2$. Then there is a sentence φ such that, say, $\varphi \in T_1$ and $\varphi \notin T_2$. This means $T_1 \in \llbracket \varphi \rrbracket$ and $T_2 \notin \llbracket \varphi \rrbracket$. Then $T_2 \in S \setminus \llbracket \varphi \rrbracket = \llbracket \neg \varphi \rrbracket$. The two open sets $\llbracket \varphi \rrbracket$ and $\llbracket \neg \varphi \rrbracket$ separate T_1 from T_2 . \square

Lemma 7.3.3. *Let X_i be clopen for each $i \in I$.*

1. *Suppose $\{X_i : i \in I\}$ has the FIP: for any $I_0 \subseteq_f I$, we have $\bigcap_{i \in I_0} X_i \neq \emptyset$. Then $\bigcap_{i \in I} X_i \neq \emptyset$.*
2. *If $S = \bigcup_{i \in I} X_i$, then there is finite $I_0 \subseteq_f I$ such that $S = \bigcup_{i \in I_0} X_i$.*

Proof. 1. Let $X_i = \llbracket \varphi_i \rrbracket$. For any $I_0 \subseteq I$, there is M with $\text{Th}(M) \in \bigcap_{i \in I_0} \llbracket \varphi_i \rrbracket$, meaning that $M \models \{\varphi_i : i \in I_0\}$. By the compactness theorem, there is M satisfying $\{\varphi_i : i \in I\}$, and then $\text{Th}(M) \in \bigcap_{i \in I} \llbracket \varphi_i \rrbracket$.

2. Let $Y_i = S \setminus X_i$. Apply part (1) to the family of clopen sets $\{Y_i\}_{i \in I}$. By assumption, $\bigcap_{i \in I} Y_i = \emptyset$, so there must be $I_0 \subseteq_f I$ such that $\bigcap_{i \in I_0} Y_i = \emptyset$, or equivalently, $\bigcup_{i \in I_0} X_i = S$. \square

Theorem 7.3.4. *The topological space S is compact.*

Proof. Suppose $S = \bigcup_{i \in I} U_i$ for some open sets U_i . Let \mathcal{F} be the family of clopen sets X such that $X \subseteq U_i$ for some i . Every open set is the union of its clopen subsets, so $U_i \subseteq \bigcup \mathcal{F}$ for each i . Then $S = \bigcup_{i \in I} U_i \subseteq \bigcup \mathcal{F}$. Applying Lemma 7.3.3(2) to the clopen cover \mathcal{F} , there is a finite subcover $S = \bigcup_{j=1}^n X_j$ with $X_j \in \mathcal{F}$. For each j , choose a $U_{i_j} \supseteq X_j$. Then $S = \bigcup_{j=1}^n X_j \subseteq \bigcup_{j=1}^n U_{i_j}$, and $\{U_{i_1}, \dots, U_{i_n}\}$ is an open subcover of the original cover. \square

If S is compact, then ultralimits should exist (Fact 7.2.3). In fact, ultralimits correspond exactly to ultraproducts:

Theorem 7.3.5. *Let M be an ultraproduct $\prod_{i \in I}^{/\mathcal{U}} M_i$. Then $\text{Th}(M)$ is the ultralimit $\lim_{i \rightarrow \mathcal{U}} \text{Th}(M_i)$ in the topological space S .*

Proof. Let N be an open set containing $\text{Th}(M)$. Then there is a clopen set $\llbracket \varphi \rrbracket$ with $\text{Th}(M) \in \llbracket \varphi \rrbracket \subseteq N$, because N is a union of clopen sets. Then

$$\text{Th}(M) \in \llbracket \varphi \rrbracket \implies M \models \varphi \implies \{i \in I : M_i \models \varphi\} \in \mathcal{U}$$

by Łoś's theorem (Theorem 6.2.3(5)). If $M_i \models \varphi$, then $\text{Th}(M_i) \in \llbracket \varphi \rrbracket \subseteq N$. Therefore

$$\{i \in I : \text{Th}(M_i) \in N\} \in \mathcal{U}. \quad \square$$

Recall that X is closed iff $S \setminus X$ is open.

Theorem 7.3.6. *A set $X \subseteq S$ is clopen if and only if X is both closed and open.*

Proof. If X is clopen, then X is open. The complement $S \setminus X$ is clopen, hence open, and so X is also closed.

Conversely, suppose X is closed and open. Then $X = \bigcup_{i \in I} Y_i$ and $S \setminus X = \bigcup_{j \in J} Z_j$ where the Y_i and Z_j are clopen sets. Note that $S = \bigcup_{i \in I} Y_i \cup \bigcup_{j \in J} Z_j$. By Lemma 7.3.3(2), there are finite $I_0 \subseteq_f I$ and $J_0 \subseteq_f J$ such that $S = \bigcup_{i \in I_0} Y_i \cup \bigcup_{j \in J_0} Z_j$. Then X is the clopen set $\bigcup_{i \in I_0} Y_i$. \square

7.4 Stone spaces

Definition 7.4.1. In any topological space, a *clopen set* is a set that is both closed and open.

Definition 7.4.2. A *Stone space* is a compact, Hausdorff topological space in which every open set is a union of clopen sets.

Theorem 7.4.3. Let S be a set. Let \mathcal{B} be a boolean subalgebra of $\mathfrak{P}(S)$. Suppose the following two conditions hold:

1. If $a, b \in S$ are distinct, then there is $X \in \mathcal{B}$ with $a \in X$ and $b \notin X$, or $b \in X$ and $a \notin X$.
2. If $\{X_i\}_{i \in I}$ is a family of sets in \mathcal{B} with the FIP, then $\bigcap_{i \in I} X_i \neq \emptyset$.

Then there is a Stone space topology on S in which the clopen sets are exactly the elements of \mathcal{B} .

Proof. This follows by the arguments of Section 7.3. \square

Theorem 7.4.4. Let S_1, S_2 be Stone spaces and $f : S_1 \rightarrow S_2$ be a map. Suppose that for any clopen set X in S_2 , the preimage $f^{-1}(X)$ is clopen in S_1 . Then f is continuous.

Proof. If U is open in S_2 , then $U = \bigcup_{i \in I} X_i$ for some clopen sets X_i . Then $f^{-1}(U) = \bigcup_{i \in I} f^{-1}(X_i)$ which is open in S_1 . \square

Chapter 8

Types and quantifier elimination

This chapter introduces two important notions in model theory: *types* and *quantifier elimination*.

Types

Fix a model M . A *type* over M can be seen as an abstract description of a potential object, which might not exist in M , but does exist in a further elementary extension of M . A *partial 1-type* over M is a set $\Sigma(x)$ of formulas $\varphi(x)$ in one variable x , with parameters from M (i.e., $\mathcal{L}(M)$ -formulas), satisfying the following two conditions, which turn out to be equivalent:

1. There is a element b in an elementary extension $N \succeq M$, and b satisfies all the formulas in $\Sigma(x)$.
2. $\Sigma(x)$ is *finitely satisfiable* in M : for any finite subset $\Sigma_0 \subseteq_f \Sigma$, there is an element of M satisfying each formula in Σ_0 .

For example, in the ordered field $(\mathbb{R}, +, \cdot, -, 0, 1, \leq)$, the set of formulas

$$\Sigma(x) = \{0 \leq x, 1 \leq x, 2 \leq x, 3 \leq x, \dots\}$$

is a partial type: any finite subset is realized by a big enough real number. But no real number satisfies all the formulas in $\Sigma(x)$. Nevertheless, using compactness, one can find an elementary extension $N \succeq \mathbb{R}$ containing an element satisfying $\Sigma(x)$. The fact that partial types can be realized encapsulates many compactness arguments.

Quantifier elimination

A theory T has *quantifier elimination* if for any formula φ , there is a quantifier-free formula φ' that is equivalent to φ in models of T . When quantifier elimination holds, there is a simple criterion for testing whether models of T are elementarily equivalent. This can sometimes be used to prove that T is complete, and when T is incomplete it can be used to find the completions of T . We will see an example of this in the next chapter when we consider the theory of algebraically closed fields. Quantifier elimination also helps one analyze the structure of definable sets (see Corollary 8.4.4 for an example in dense linear orders).

Fortunately, many theories have quantifier elimination. How do we prove that a given theory T has quantifier elimination? One way is to show that an element's type is determined by its *quantifier-free type* (Theorem 8.3.3). This is often used in conjunction with the *back-and-forth systems* of Definition 3.7.5. We will see one example of this method in the current chapter (Section 8.4), another example in the next chapter (Section 9.4), and a third example in Chapter 11 (Section 11.6) after stating the definitive criterion for quantifier elimination (Theorem 11.5.4).

8.1 Types

Definition 8.1.1. Let M be an \mathcal{L} -structure, let A be a subset, and let \bar{b} be an n -tuple. The *type* of \bar{b} over A , written $\text{tp}(\bar{b}/A)$, is the set of $\mathcal{L}(A)$ -formulas $\varphi(x_1, \dots, x_n)$ such that $M \models \varphi(\bar{b})$. When $A = \emptyset$, we write $\text{tp}(\bar{b}/A)$ as $\text{tp}(\bar{b})$. We write $\text{tp}(-)$ as $\text{tp}^M(-)$ when we need to specify M .

Remark 8.1.2. Partial elementary maps preserve types: if f is a partial elementary map from M to N and \bar{a} is a tuple in $\text{dom}(f)$, then $\text{tp}(\bar{a}) = \text{tp}(f(\bar{a}))$. Indeed, for any formula $\varphi(\bar{x})$,

$$\varphi(\bar{x}) \in \text{tp}(\bar{a}) \iff M \models \varphi(\bar{a}) \iff N \models \varphi(f(\bar{a})) \iff \varphi(\bar{x}) \in \text{tp}(f(\bar{a})).$$

Remark 8.1.3. If \bar{b}, \bar{c} are two n -tuples in the same structure M , then the following are equivalent:

1. $\text{tp}(\bar{b}/A) = \text{tp}(\bar{c}/A)$.
2. For every A -definable set D , $\bar{b} \in D \iff \bar{c} \in D$.

Indeed, (1) and (2) are equivalent to

3. For every $\mathcal{L}(A)$ -formula $\varphi(\bar{x})$, $M \models \varphi(\bar{b}) \iff M \models \varphi(\bar{c})$.

Let M be an \mathcal{L} -structure and A be a set.

Definition 8.1.4. A *complete n -type* over A is something of the form $\text{tp}^N(\bar{b}/A)$ for some $N \succeq M$ and n -tuple $\bar{b} \in N^n$. The set of complete n -types is written $S_n(A)$.

Lemma 8.1.5. 1. If $p \in S_n(A)$ and $\varphi(\bar{x})$ is an $\mathcal{L}(A)$ -formula, then $\neg\varphi \in p \iff \varphi \notin p$.

2. If $p, q \in S_n(A)$ and $p \subseteq q$, then $p = q$.

Proof. 1. If $p = \text{tp}^N(\bar{b}/A)$ for some elementary extension $N \succeq M$ and $\bar{b} \in N^n$, then

$$\neg\varphi \in p \iff N \models \neg\varphi(\bar{b}) \iff N \not\models \varphi(\bar{b}) \iff \varphi \notin p.$$

2. Otherwise take $\varphi \in q \setminus p$. Then $\varphi \in q \implies \neg\varphi \notin q$, and $\varphi \notin p \implies \neg\varphi \in p$. Then $\neg\varphi \in p \setminus q$, contradicting $p \subseteq q$. \square

Definition 8.1.6. Let $\Sigma(x_1, \dots, x_n)$ be a set of $\mathcal{L}(A)$ -formulas in the variables x_1, \dots, x_n . Then $\bar{b} \in M^n$ *realizes* Σ if $\Sigma \subseteq \text{tp}(\bar{b}/A)$. We say that $\Sigma(\bar{x})$ is *realized* in M if some $\bar{a} \in M^n$ realizes Σ , and *omitted* in M otherwise.

Remark 8.1.7. If $p \in S_n(A)$ is a complete type, then a tuple \bar{b} realizes p if and only if $\text{tp}(\bar{b}/A) = p$, by Lemma 8.1.5(2) applied to the complete types p and $\text{tp}(\bar{b}/A)$.

Theorem 8.1.8. Let $\Sigma(\bar{x})$ be a set of $\mathcal{L}(A)$ -formulas in the variables x_1, \dots, x_n . The following are equivalent:

1. Every finite subset $\Sigma_0 \subseteq_f \Sigma$ is realized in M .
2. Σ is realized in an elementary extension of M .

Proof. Consider a third condition:

3. Every finite subset $\Sigma_0 \subseteq_f \Sigma$ is realized in an elementary extension of M .

We claim $(3) \implies (1) \implies (2) \implies (3)$. The direction $(2) \implies (3)$ is clear.

$(3) \implies (1)$: if $\Sigma_0 = \{\varphi_1, \dots, \varphi_n\}$, and Σ_0 is realized in $N \succeq M$, then

$$N \models \exists \bar{x} \bigwedge_{i=1}^n \varphi_i(\bar{x}).$$

As $M \preceq N$, the same sentence holds in M , which means that Σ_0 is realized in M .

$(1) \implies (2)$: Let $\mathcal{L}' = \mathcal{L}(A) \cup \{c_1, \dots, c_n\}$ where the c_i are new constant symbols. If $\Sigma_0 \subseteq_f \Sigma$, then the \mathcal{L}' -structure

$$\text{eldiag}(M) \cup \{\varphi(\bar{c}) : \varphi \in \Sigma_0\}$$

is satisfied by the structure M with \bar{c} interpreted as a realization of Σ_0 . By compactness,

$$\text{eldiag}(M) \cup \{\varphi(\bar{c}) : \varphi \in \Sigma\}$$

is satisfied by some \mathcal{L}' -structure N . The fact that $N \models \text{eldiag}(M)$ means that, up to isomorphism, $N \succeq M$. Then the interpretation \bar{c}^N of \bar{c} in N is an n -tuple realizing $\Sigma(\bar{x})$, and so (2) holds. \square

Definition 8.1.9. A *partial n -type* over A is a set $\Sigma(\bar{x})$ of $\mathcal{L}(A)$ -formulas in the variables x_1, \dots, x_n satisfying the equivalent conditions of Theorem 8.1.8.

Remark 8.1.10. Condition (2) of Theorem 8.1.8 says that $\Sigma(\bar{x})$ is a partial type if and only if $\Sigma(\bar{x}) \subseteq \text{tp}^N(\bar{b}/A)$ for some tuple \bar{b} in an elementary extension $N \succeq M$. Equivalently, $\Sigma(\bar{x})$ is a partial type if and only if Σ is a subset of a complete type. In particular, complete types are partial types.

Theorem 8.1.11. *Let $p(\bar{x})$ be a set of $\mathcal{L}(A)$ -formulas in \bar{x} . Then $p(\bar{x})$ is a complete type over A if and only if $p(\bar{x})$ is a maximal partial type over A .*

Proof. By Remark 8.1.10, every partial type is contained in a complete type. Therefore, any maximal partial type is a complete type.

Conversely, suppose p is a complete type, but not maximal. Take a larger partial type $\Sigma \supsetneq p$. Then $\Sigma \subseteq q$ for some complete type q , and we have $p \subsetneq \Sigma \subseteq q$, contradicting the incomparability of complete types (Lemma 8.1.5(2)). \square

Types over a theory

Let T be an \mathcal{L} -theory.

Definition 8.1.12. A *complete n -type* over T is something of the form $\text{tp}^M(\bar{a})$ for some $M \models T$ and n -tuple $\bar{a} \in M^n$. The set of complete n -types is written $S_n(T)$.

Lemma 8.1.13. *If $p, q \in S_n(T)$ and $p \subseteq q$, then $p = q$.*

Proof. Like Lemma 8.1.5. □

Theorem 8.1.14. *If $M \models T$ and $\bar{a} \in M^n$ realizes $p \in S_n(T)$, then $\text{tp}(\bar{a}) = p$.*

Proof. Like Remark 8.1.7. □

Theorem 8.1.15. *Let $\Sigma(\bar{x})$ be a set of \mathcal{L} -formulas in the variables x_1, \dots, x_n . The following are equivalent:*

1. *Every finite subset $\Sigma_0 \subseteq \Sigma$ is realized in a model of T .*
2. *Σ is realized in a model of T .*

Proof. Like Theorem 8.1.8, but more straightforward. □

Definition 8.1.16. A *partial n -type* over T is a set $\Sigma(\bar{x})$ of \mathcal{L} -formulas in the variables x_1, \dots, x_n satisfying the equivalent conditions of Theorem 8.1.15.

Remark 8.1.17. As in Remark 8.1.10, $\Sigma(\bar{x})$ is a partial type if and only if $\Sigma(\bar{x})$ is a subset of a complete type. In particular, complete types are partial types.

Theorem 8.1.18. *Let $p(\bar{x})$ be a set of \mathcal{L} -formulas in \bar{x} . Then $p(\bar{x})$ is a complete type over T if and only if $p(\bar{x})$ is a maximal partial type over T .*

Proof. Like Theorem 8.1.11 □

Lemma 8.1.19. *Let T be a complete theory, M be a model, and $\Sigma(\bar{x})$ be a set of \mathcal{L} -formulas in the variables \bar{x} .*

1. *If $\Sigma(\bar{x})$ is finite, then $\Sigma(\bar{x})$ is realized in a model of T if and only if $\Sigma(\bar{x})$ is realized in M .*

2. $\Sigma(\bar{x})$ is a partial type over T if and only if $\Sigma(\bar{x})$ is a partial type over $\emptyset \subseteq M$.
3. $\Sigma(\bar{x})$ is a complete type over T if and only if $\Sigma(\bar{x})$ is a complete type over $\emptyset \subseteq M$.

Proof. 1. If $N \models \exists \bar{x} \bigwedge_{\varphi \in \Sigma} \varphi(\bar{x})$ and $N \models T$ then $M \models \exists \bar{x} \bigwedge_{\varphi \in \Sigma} \varphi(\bar{x})$ because $M \equiv N$.

2. By part (1), $\Sigma(\bar{x})$ is finitely realized in models of T if and only if $\Sigma(\bar{x})$ is finitely realized in M .
3. The complete types over T are the maximal partial types over T , and similarly for types over $A = \emptyset$. Since the sets of partial types agree by part (2), the sets of maximal partial types agree. \square

Corollary 8.1.20. *If T is a complete theory and M is a model, then $S_n(T) = S_n^M(\emptyset)$.*

Fact 8.1.21. *If $A \subseteq M$ and $T_{A,M}$ is the complete $\mathcal{L}(A)$ -theory of M , then $S_n^M(A) = S_n(T_{A,M})$.*

8.2 The topology on type space

Let M be an \mathcal{L} -structure and A be a subset of M .

Theorem 8.2.1. *For each $\mathcal{L}(A)$ -formula $\varphi(x_1, \dots, x_n)$, let $\llbracket \varphi \rrbracket = \{p \in S_n(A) : \varphi \in p\}$. Then*

$$\begin{aligned} \llbracket \varphi \wedge \psi \rrbracket &= \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket \\ \llbracket \varphi \vee \psi \rrbracket &= \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket \\ \llbracket \neg \varphi \rrbracket &= S_n(A) \setminus \llbracket \varphi \rrbracket \\ \llbracket \top \rrbracket &= S_n(A) \quad \llbracket \perp \rrbracket = \emptyset \end{aligned}$$

Proof. This is a matter of unwinding definitions. For example, if $p \in S_n(A)$ is $\text{tp}^N(\bar{b}/A)$, then

$$\begin{aligned} p \in \llbracket \varphi \wedge \psi \rrbracket &\iff \varphi \wedge \psi \in p \iff N \models \varphi(\bar{b}) \wedge \psi(\bar{b}) \\ &\iff N \models \varphi(\bar{b}) \text{ and } N \models \psi(\bar{b}) \iff \dots \iff (p \in \llbracket \varphi \rrbracket \text{ and } p \in \llbracket \psi \rrbracket), \end{aligned}$$

showing that $\llbracket \varphi \wedge \psi \rrbracket = \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket$. \square

Theorem 8.2.2. *If φ, ψ are $\mathcal{L}(A)$ -formulas, then*

$$\begin{aligned} \llbracket \varphi \rrbracket \subseteq \llbracket \psi \rrbracket &\iff \varphi(M) \subseteq \psi(M) \\ \llbracket \varphi \rrbracket = \llbracket \psi \rrbracket &\iff \varphi(M) = \psi(M). \end{aligned}$$

Proof. We prove the first line, which directly implies the second line. Unwinding the definitions, $\llbracket \varphi \rrbracket \subseteq \llbracket \psi \rrbracket$ means that for every $N \succeq M$ and $\bar{b} \in N$, if $N \models \varphi(\bar{b})$ then $N \models \psi(\bar{b})$. Equivalently, every $N \succeq M$ satisfies

$$N \models \forall \bar{x} (\varphi(\bar{x}) \rightarrow \psi(\bar{x})).$$

Similarly, $\varphi(M) \subseteq \psi(M)$ means

$$M \models \forall \bar{x} (\varphi(\bar{x}) \rightarrow \psi(\bar{x})).$$

The two conditions are equivalent because $N \succeq M$. \square

Theorem 8.2.3. *The collection of sets $\llbracket \varphi \rrbracket$ form a basis for a topology making $S_n(A)$ into a Stone space, and the sets $\llbracket \varphi \rrbracket$ are exactly the clopen sets in this topology.*

Proof. We use Theorem 7.4.3. There are two things to check:

1. If $p, q \in S_n(A)$ are distinct, then there is a set $\llbracket \varphi \rrbracket$ distinguishing p and q . Indeed, take φ in $p \setminus q$.
2. If $\{\llbracket \varphi_i \rrbracket\}_{i \in I}$ has the FIP, then $\bigcap_i \llbracket \varphi_i \rrbracket \neq \emptyset$. Equivalently, if every finite subset of $\{\varphi_i : i \in I\}$ is contained in a complete type, then $\{\varphi_i : i \in I\}$ is contained in a complete type. This is clear by Theorem 8.1.8. \square

Using Theorem 8.2.2 we see the following:

Corollary 8.2.4. *The boolean algebra of clopen sets in $S_n(A)$ is isomorphic to the boolean algebra of A -definable subsets of M^n via the isomorphism $\llbracket \varphi \rrbracket \mapsto \varphi(M^n)$.*

There is an analogous picture for type spaces of a theory. Let T be an \mathcal{L} -theory:

Theorem 8.2.5. *For each \mathcal{L} -formula $\varphi(x_1, \dots, x_n)$, let $\llbracket \varphi \rrbracket = \{p \in S_n(T) : \varphi \in p\}$. Then*

$$\begin{aligned}\llbracket \varphi \wedge \psi \rrbracket &= \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket \\ \llbracket \varphi \vee \psi \rrbracket &= \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket \\ \llbracket \neg \varphi \rrbracket &= S_n(T) \setminus \llbracket \varphi \rrbracket \\ \llbracket \top \rrbracket &= S_n(T) \quad \llbracket \perp \rrbracket = \emptyset\end{aligned}$$

Moreover, $\llbracket \varphi \rrbracket = \llbracket \psi \rrbracket$ if and only if φ and ψ are equivalent modulo T , in the sense that

$$T \vdash \forall \bar{x} (\varphi(\bar{x}) \leftrightarrow \psi(\bar{x}))$$

or equivalently $\varphi(M) = \psi(M)$ for all $M \models T$.

Theorem 8.2.6. *The collection of sets $\llbracket \varphi \rrbracket$ is a basis for a topology making $S_n(T)$ a Stone space, and the clopen sets are exactly the sets $\llbracket \varphi \rrbracket$.*

8.3 Quantifier elimination

Definition 8.3.1. A theory T has *quantifier elimination* if for every formula $\varphi(\bar{x})$, there is a quantifier-free formula $\psi(\bar{x})$ that is equivalent to φ in models of T :

$$T \vdash \forall \bar{x} (\varphi(\bar{x}) \leftrightarrow \psi(\bar{x})).$$

Definition 8.3.2. If (a_1, \dots, a_n) is a tuple in a structure M , then the *quantifier-free type* $\text{qftp}(\bar{a})$ is the set of quantifier-free formulas $\varphi(x_1, \dots, x_n)$ with $M \models \varphi(\bar{a})$.

For any theory T , let $S_n^{\text{qfree}}(T)$ be the space of quantifier-free n -types, analogous to the space of complete n -types $S_n(T)$. Each quantifier-free formula $\varphi(x_1, \dots, x_n)$ defines a set $\llbracket \varphi \rrbracket_{\text{qfree}} \subseteq S_n^{\text{qfree}}(T)$, and these sets form a basis for a topology making $S_n^{\text{qfree}}(T)$ into a Stone space. Moreover, every clopen set in $S_n^{\text{qfree}}(T)$ has the form $\llbracket \varphi \rrbracket_{\text{qfree}}$ for some quantifier-free φ . The proofs are analogous to the case of $S_n(T)$ (Section 8.2). There is a restriction map

$$S_n(T) \rightarrow S_n^{\text{qfree}}(T)$$

sending $\text{tp}(\bar{a})$ to $\text{qftp}(\bar{a})$. Note that

$$\text{qftp}^M(\bar{a}) \in \llbracket \varphi \rrbracket_{\text{qfree}} \iff M \models \varphi(\bar{a}) \iff \text{tp}^M(\bar{a}) \in \llbracket \varphi \rrbracket$$

so the preimage of $\llbracket \varphi \rrbracket_{\text{qfree}}$ is $\llbracket \varphi \rrbracket$. Because preimages of clopen sets are clopen, the restriction map is continuous (Theorem 7.4.4).

Theorem 8.3.3. *The following are equivalent:*

1. *T has quantifier elimination.*
2. *For any $M, N \models T$ and $\bar{a} \in M^n$ and $\bar{b} \in N^n$,*

$$\text{qftp}^M(\bar{a}) = \text{qftp}^N(\bar{b}) \implies \text{tp}^M(\bar{a}) = \text{tp}^N(\bar{b}).$$

Proof. Consider a third condition:

3. *For every n , the restriction map $S_n(T) \rightarrow S_n^{\text{qfree}}(T)$ is a homeomorphism.*

We claim $(1) \implies (2) \implies (3) \implies (1)$.

$(1) \implies (2)$: Suppose that T has quantifier elimination and $\text{qftp}^M(\bar{a}) = \text{qftp}^N(\bar{b})$. For any formula $\varphi(\bar{x})$, there is an equivalent quantifier free-formula $\psi(\bar{x})$, and then

$$M \models \varphi(\bar{a}) \iff M \models \psi(\bar{a}) \iff N \models \psi(\bar{b}) \iff N \models \varphi(\bar{b}),$$

so that $\text{tp}^M(\bar{a}) = \text{tp}^N(\bar{b})$.

$(2) \implies (3)$: The restriction map is surjective, since $\text{qftp}^M(\bar{a})$ lifts to $\text{tp}^M(\bar{a})$. The restriction map is injective by (2). Then the restriction map is a continuous bijection between compact Hausdorff spaces, and therefore a homeomorphism (Theorem 7.2.6).

$(3) \implies (1)$: Let $\varphi(x_1, \dots, x_n)$ be a formula. Then $\llbracket \varphi \rrbracket \subseteq S_n(T)$ is clopen. The image under the homeomorphism $S_n(T) \rightarrow S_n^{\text{qfree}}(T)$ is a clopen set, which must be $\llbracket \psi \rrbracket_{\text{qfree}} \subseteq S_n^{\text{qfree}}(T)$ for some quantifier-free formula ψ . Then

$$M \models \varphi(\bar{a}) \iff \text{tp}(\bar{a}) \in \llbracket \varphi \rrbracket \iff \text{qftp}(\bar{a}) \in \llbracket \psi \rrbracket_{\text{qfree}} \iff M \models \psi(\bar{a}),$$

so $T \vdash \varphi \leftrightarrow \psi$. □

Theorem 8.3.4. *Suppose M, N are \mathcal{L} -structures and $\bar{a} \in M^n$ and $\bar{b} \in N^n$. Then the following are equivalent:*

1. $\text{qftp}(\bar{a}) = \text{qftp}(\bar{b})$.
2. *There is an isomorphism $f : \langle \bar{a} \rangle_M \rightarrow \langle \bar{b} \rangle_N$ with $f(\bar{a}) = \bar{b}$.*

Proof. (1) \implies (2): Let $p = \text{qftp}(\bar{a}) = \text{qftp}(\bar{b})$. Note that every element of $\langle \bar{a} \rangle_M$ has the form $t^M(\bar{a})$ for some term t . Similarly, $\langle \bar{b} \rangle_N = \{t^N(\bar{b}) : t(\bar{x}) \text{ a term}\}$. Moreover,

$$t^M(\bar{a}) = s^M(\bar{a}) \iff (t(\bar{x}) = s(\bar{x})) \in p \iff t^N(\bar{b}) = s^N(\bar{b}),$$

Therefore there is a bijection $f : \langle \bar{a} \rangle_M \rightarrow \langle \bar{b} \rangle_N$ sending $t^M(\bar{a})$ to $t^N(\bar{b})$. If R is a k -ary relation symbol and t_1, \dots, t_k are terms in \bar{x} , then

$$R^M(t_1^M(\bar{a}), \dots, t_k^M(\bar{a})) \iff R(t_1, \dots, t_k) \in p \iff R^N(t_1^N(\bar{b}), \dots, t_k^N(\bar{b})),$$

and so f preserves relation symbols. A similar argument shows f preserves function symbols.

(2) \implies (1): Let $A = \langle \bar{a} \rangle_M$ and $B = \langle \bar{b} \rangle_N$. Then

$$\text{qftp}^M(\bar{a}) = \text{qftp}^A(\bar{a}) = \text{qftp}^B(\bar{b}) = \text{qftp}^N(\bar{b})$$

by Theorem 3.4.3 applied to the embeddings $A \rightarrow M$, $B \rightarrow N$, and $A \xrightarrow{\cong} B$. \square

Theorem 8.3.5. *Suppose T has quantifier elimination. If $M, N \models T$, then*

$$M \equiv N \iff \langle \emptyset \rangle_M \cong \langle \emptyset \rangle_N.$$

Proof. The left hand side says that $\text{tp}^M() = \text{tp}^N()$. Indeed, $\text{tp}^M()$ is the set of formulas in 0 free variables (i.e., sentences) satisfied by the empty tuple (i.e., true in M), which is just $\text{Th}(M)$.

Meanwhile, the right hand side of $(*)$ is equivalent to $\text{qftp}^M() = \text{qftp}^N()$ by Theorem 8.3.4. We don't have to worry about how the isomorphism acts on the generators, because there are no generators to check.

Finally, quantifier elimination gives

$$\text{tp}^M() = \text{tp}^N() \iff \text{qftp}^M() = \text{qftp}^N(). \quad \square$$

Theorem 8.3.6. *Suppose T has quantifier elimination and $M, N \models T$.*

1. *If $f : M \rightarrow N$ is an embedding, then f is an elementary embedding.*
2. *If M is a substructure of N , then M is an elementary substructure of N .*

Proof. Embeddings preserve quantifier-free formulas (Theorem 3.4.3(3)), and quantifier elimination allows us to replace any formula with an equivalent quantifier-free formula. \square

8.4 Quantifier elimination in DLO

Theorem 8.4.1. *DLO has quantifier elimination.*

Proof. We use the criterion of Theorem 8.3.3. Suppose $M, N \models \text{DLO}$, $\bar{a} \in M^n$, $\bar{b} \in N^n$, and $\text{qftp}^M(\bar{a}) = \text{qftp}^N(\bar{b})$. We must show $\text{tp}^M(\bar{a}) \stackrel{?}{=} \text{tp}^N(\bar{b})$.

By Theorem 8.3.4, there is a partial isomorphism $f : \langle \bar{a} \rangle_M \rightarrow \langle \bar{b} \rangle_N$ with $f(\bar{a}) = \bar{b}$. As the language of orders is relational, all terms are trivial, and so $\text{dom}(f) = \langle \bar{a} \rangle_M = \{a_1, \dots, a_n\}$. Then f is a finite partial isomorphism. By Theorem 3.7.10, the class of finite partial isomorphisms is a back-and-forth system, and so f is a partial elementary map by Theorem 3.7.6. Therefore $\text{tp}^M(\bar{a}) = \text{tp}^N(f(\bar{a})) = \text{tp}^N(\bar{b})$. \square

If M, N are models of DLO, then $\langle \emptyset \rangle_M$ and $\langle \emptyset \rangle_N$ are both the empty order \emptyset , so $M \equiv N$. This gives another proof of the completeness of DLO.

Corollary 8.4.2. *\mathbb{Q} and the open interval $(0, 1) = \{x \in \mathbb{R} : 0 < x < 1\}$ are elementary substructures of \mathbb{R} .*

Definition 8.4.3. If $(M, \leq) \models \text{DLO}$ and $a < b$ are two points in M , then we define

$$\begin{aligned} (a, b) &:= \{x \in M : a < x < b\} \\ (a, +\infty) &:= \{x \in M : a < x\} \\ (-\infty, a) &:= \{x \in M : x < a\} \\ (-\infty, +\infty) &:= M \end{aligned}$$

Sets of these forms are called *open intervals*.

Corollary 8.4.4. *If $M \models \text{DLO}$ and $D \subseteq M^1$ is definable, then D is a finite union of points and open intervals.*

Proof. Let \mathcal{F} be the collection of sets $D \subseteq M$ such that D is a finite union of points and open intervals. It is an exercise to see that \mathcal{F} is closed under boolean operations (it is a boolean subalgebra of $\mathfrak{P}(M)$). We must show that $D \in \mathcal{F}$. Write D as $\varphi(M, \bar{b})$ for some \mathcal{L} -formula $\varphi(x, \bar{y})$ and tuple of parameters \bar{b} . By quantifier-elimination, we may assume φ is quantifier-free. Then φ is a boolean combination of atomic formulas. As \mathcal{F} is closed under

boolean combinations, we may assume φ is atomic. Then $\varphi(x, \bar{b})$ has one of the following forms:

$$x \leq x, \ x \leq b_i, \ b_i \leq x, \ b_i \leq b_j, \ x = x, \ x = b_i, \ b_i = x, \ b_i = b_j.$$

Each of these formulas defines a set in \mathcal{F} .

□

Chapter 9

Algebraically closed fields

We now turn to an important example—the theory ACF of *algebraically closed fields*. An algebraically closed field is a field in which every polynomial equation

$$a_n x^n + \cdots a_1 x + a_0 = 0$$

has a solution (except in the degenerate case where $a_0 \neq 0 = a_1 = a_2 = \cdots = a_n$). The field \mathbb{C} is algebraically closed—a fact known as the “fundamental theorem of algebra”. The class of algebraically closed fields is axiomatized by a theory ACF. For $p \in \{0, 2, 3, 5, 7, \dots\}$, ACF_p is the theory of algebraically closed fields of characteristic p . For example, $\mathbb{C} \models \text{ACF}_0$.

In the current chapter, we show that ACF has quantifier elimination (Theorem 9.4.4). From this, we see that two algebraically closed fields are elementarily equivalent if and only if they have the same characteristic (Corollary 9.4.5). As a consequence, each theory ACF_p is complete, including ACF_0 (Corollary 9.4.7). As \mathbb{C} is a model of ACF_0 , it follows that the structure \mathbb{C} is decidable—there is an algorithm which determines whether a given sentence is true or false in \mathbb{C} (Corollary 9.4.8).

We continue to use ACF as a running example in later chapters. In Chapter 10, we show that if K is algebraically closed, then any system of equations that has a solution in an extension of K already has a solution in K (Theorem 10.3.6), and we verify that algebraically closed fields of characteristic p actually exist for $p > 0$ (Corollary 10.3.5). In Chapter 13, we show that the set \mathbb{Q}^{alg} of *algebraic numbers* is an elementary substructure of \mathbb{C} (Example 13.5.5). In Chapter 14, we show that the theories ACF_0 and ACF_p are κ -categorical for any $\kappa > \aleph_0$ (Corollary 14.2.7), and use this to classify algebraically closed fields (Theorem 14.2.10). We will also see that there is

a good notion of “dimension” for definable sets in algebraically closed fields (Sections 14.5–14.6), a fact known to algebraic geometers.

9.1 Polynomial rings

Fact 9.1.1. *Let R be a ring and x be a symbol. There is a ring $R[x]$ extending R , generated by $R \cup \{x\}$, with the following properties:*

1. *Every element of $R[x]$ has the form $\sum_{i=0}^n a_i x^i$ for some $n \geq 0$ and $a_0, \dots, a_n \in R$.*
2. *Two elements $\sum_{i=0}^n a_i x^i$ and $\sum_{i=0}^n b_i x^i$ are equal if and only if the tuples \bar{a} and \bar{b} are equal.*
3. *The sum of two elements is given by*

$$\sum_{i=0}^n a_i x^i + \sum_{i=0}^n b_i x^i = \sum_{i=0}^n (a_i + b_i) x^i.$$

4. *The product of two elements is given by*

$$\left(\sum_{i=0}^n a_i x^i \right) \left(\sum_{j=0}^m b_j x^j \right) = \sum_{k=0}^{n+m} c_k x^k, \text{ where } c_k = \sum_{\substack{0 \leq i \leq n \\ 0 \leq j \leq m \\ i+j=k}} a_i b_j.$$

Elements of $R[x]$ are called polynomials.

If $P(x)$ is a non-zero polynomial, then we can write

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0,$$

where $n \geq 0$ and $a_n \neq 0$. Then $a_n x^n$ is called the *leading term*, a_n is called the *leading coefficient*, and n is called the *degree* of $P(x)$. We write the degree of P as $\deg P$. The degree of the zero polynomial is defined to be $-\infty$. A nonzero polynomial is *monic* if the leading coefficient is 1.

Definition 9.1.2 (Evaluating polynomials). If $P(x) \in R[x]$ has the form

$$P(x) = \sum_{i=0}^n a_i x^i$$

and if $b \in R$, then

$$P(b) := \sum_{i=0}^n a_i b^i.$$

Fact 9.1.3. *For fixed b , the map*

$$\begin{aligned} R[x] &\rightarrow R \\ P(x) &\rightarrow P(b) \end{aligned}$$

is a ring homomorphism, meaning among other things that

$$\begin{aligned} (P + Q)(b) &= P(b) + Q(b) \\ (PQ)(b) &= P(b)Q(b) \end{aligned}$$

Lemma 9.1.4. *Let K be a field, and let $A(x), B(x) \in K[x]$ be polynomials with $B(x)$ non-zero. then there is $R(x) \in K[x]$ such that*

$$\begin{aligned} A(x) &\equiv R(x) \pmod{B(x)} \\ \deg R(x) &< \deg B(x). \end{aligned}$$

Proof. If bx^m is the leading term of $B(x)$, replace $B(x)$ with $b^{-1}B(x)$. Then we can assume $B(x)$ is monic. Proceed by induction on $\deg A(x)$. If $\deg A(x) < \deg B(x)$, take $R(x) = A(x)$. Otherwise, let

$$\begin{aligned} A(x) &= a_n x^n + a_{n-1} x^{n-1} + \cdots \\ B(x) &= x^m + b_{m-1} x^{m-1} + \cdots \end{aligned}$$

where $m = \deg(B) \leq n = \deg(A)$. Let

$$\begin{aligned} A'(x) &= A(x) - a_n x^{n-m} B(x) \\ &= (a_n x^n + a_{n-1} x^{n-1} + \cdots) - (a_n x^n + a_n b_{m-1} x^{n-1} + \cdots + a_n b_0 x^{n-m}). \end{aligned}$$

Then $\deg A'(x) < n = \deg A(x)$, but $A(x) \equiv A'(x) \pmod{B(x)}$. By induction, $A'(x) \equiv R(x) \pmod{B(x)}$ for some suitable $R(x)$. \square

Let K be a field.

Theorem 9.1.5. *Suppose $P(x) \in K[x]$ and $P(a) = 0$ for some $a \in K$. Then $P(x) = (x - a)Q(x)$ for some $Q(x) \in K[x]$.*

Proof. Apply the division lemma to $P(x)$ and $(x - a)$, to get $R(x) \in K[x]$ with

$$\begin{aligned} P(x) &\equiv R(x) \pmod{x - a} \\ \deg R(x) &< \deg(x - a) = 1. \end{aligned}$$

The first line means there is $Q(x) \in K[x]$ with

$$P(x) = (x - a)Q(x) + R(x). \quad (*)$$

The second line means $R(x) = c$ for some constant c . Substituting $x = a$ into $(*)$, we see

$$\begin{aligned} P(a) &= (a - a)Q(a) + c \\ 0 &= 0 + c. \end{aligned}$$

Then $c = 0$, so $R(x) = 0$ and $P(x) = (x - a)Q(x)$. □

A *root* of $P(x) \in K[x]$ is an element $a \in K$ with $P(a) = 0$. Note that a is a root of $P(x)Q(x)$ if and only if a is a root of $P(x)$ or a is a root of $Q(x)$, by the zero law (Theorem 1.4.16).

Theorem 9.1.6. *If $P(x)$ is a non-zero polynomial, then the number of roots of $P(x)$ in K is at most $\deg P(x)$.*

Proof. Let $d = \deg P(x)$. If $P(x)$ has no roots, then the claim holds. Otherwise, take some root a . Then $P(x) = (x - a)Q(x)$ for some polynomial $Q(x)$ of degree $d - 1$. By induction on d , $Q(x)$ has at most $d - 1$ roots. As $(x - a)$ has one root, $P(x)$ has at most $(d - 1) + 1 = d$ roots. □

Theorem 9.1.7. *If K is a field, every ideal $I \subseteq K[x]$ is a principal ideal $P(x)K[x]$ for some polynomial $P(x)$.*

Proof. Like the proof of the same fact in \mathbb{Z} (Theorem 2.5.3), but using Lemma 9.1.4 instead of Lemma 2.5.2 and using $\deg P$ instead of $|n|$. □

Remark 9.1.8. In Theorem 9.1.7, scaling $P(x)$ by a non-zero constant, we may assume $P(x)$ is zero or monic. Then $P(x)$ is uniquely determined.

Definition 9.1.9. A non-constant polynomial $P(x)$ is *reducible* if $P(x)$ is a product of two non-constant polynomials, and *irreducible* otherwise.

Theorem 9.1.10. *If $P(x)$ is an irreducible polynomial in $K[x]$, then the quotient $K[x]/P(x)K[x]$ is a field.*

Proof. Like the proof that $\mathbb{Z}/p\mathbb{Z}$ is a field when p is prime (Theorem 2.5.7). □

9.2 The theory ACF

Definition 9.2.1. A field K is *algebraically closed* if every polynomial $P(x) \in K[x]$ with $\deg P > 0$ has a root. That is, for every $n > 0$ and $a_0, a_1, \dots, a_n \in K$ with $a_n \neq 0$, there is $x \in K$ with

$$a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0 = 0.$$

Note that the class of algebraically closed fields is elementary, defined by a theory consisting of the field axioms plus an axiom schema

$$\forall a_1, \dots, a_n \left(a_n \neq 0 \rightarrow \exists x \sum_{i=0}^n a_i x^i = 0 \right) \text{ for } n > 0.$$

This theory is usually called *ACF*.

Fact 9.2.2 (Fundamental theorem of algebra). *The field $\mathbb{C} = \{x + yi : x, y \in \mathbb{R}\}$ is algebraically closed.*

Theorem 9.2.3. *If K is algebraically closed, then K is infinite.*

Proof. If $K = \{a_1, \dots, a_n\}$, then $P(x) = 1 + \prod_{i=1}^n (x - a_i)$ has no root in K , so $K \not\models \text{ACF}$. \square

9.3 Algebraic and transcendental elements

Let L be a field and K be a subfield.

Definition 9.3.1. An element $a \in L$ is *algebraic* over K if $P(a) = 0$ for some non-zero polynomial $P(x) \in K[x]$. Otherwise, a is *transcendental* over K .

If $a \in L$, then $K[a]$ denotes the subring of L generated by $K \cup \{a\}$.

Theorem 9.3.2. *Fix $a \in L$. Let $I_{a/K} = \{P(x) \in K[x] : P(a) = 0\}$.*

1. $I_{a/K}$ is an ideal in $K[x]$.
2. $K[x]/I_{a/K}$ is isomorphic to $K[a]$ via the isomorphism sending $P(x)$ to $P(a)$.

3. If a is transcendental over K , then $I_{a/K}$ is the zero ideal $0 \cdot K[x] = \{0\}$.
4. If a is algebraic over K , then $I_{a/K} = P(x) \cdot K[x]$ for some irreducible monic polynomial $P(x)$.

Proof. Let $f : K[x] \rightarrow K[a]$ be the ring homomorphism $P(x) \mapsto P(a)$. Then $I_{a/K}$ is the kernel. The image $\text{im}(f)$ is a subring of $K[a]$ containing $f(K) = K$ and $f(x) = a$, so it must be all of $K[a]$. By the fundamental theorem on homomorphisms, $K[x]/I_{a/K}$ is isomorphic to the image $K[a]$ via the map $P(x) \mapsto P(a)$.

By Theorem 9.1.7 and Remark 9.1.8, the ideal $I_{a/K}$ is $P(x) \cdot K[x]$, where $P(x)$ is zero or a monic polynomial. In the first case, $I_{a/K} = \{0\}$, which means precisely that a is transcendental. In the second case, $P(x) \in I_{a/K}$ implies that $P(a) = 0$, and so a is algebraic. Suppose for the sake of contradiction that $P(x)$ is reducible as $P(x) = Q_1(x)Q_2(x)$. Note that $\deg P > \deg Q_i$ for $i = 1, 2$, so $Q_i \notin P(x) \cdot K[x] = I_{a/K}$, and therefore $Q_i(a) \neq 0$ for $i = 1, 2$. But then $0 = P(a) = Q_1(a)Q_2(a) \neq 0$, a contradiction. \square

Definition 9.3.3. If $a \in L$ is algebraic over K , then the *minimal polynomial* of a over K is the monic irreducible polynomial $P(x)$ appearing in Theorem 9.3.2(4).

Theorem 9.3.4. If $P(x) \in K[x]$ is a monic irreducible polynomial and $a \in L$ is a root of $P(x)$, then $P(x)$ is the minimal polynomial of a .

Proof. Let $P_0(x)$ be the actual minimal polynomial of a . Then

$$P(a) = 0 \implies P(x) \in I_{a/K} = P_0(x) \cdot K[x].$$

Thus $P(x)$ is a multiple of $P_0(x)$:

$$P(x) = P_0(x)Q(x).$$

As $P(x)$ is irreducible, this forces $Q(x) = 1$ and $P(x) = P_0(x)$. \square

9.4 Quantifier elimination in ACF

Lemma 9.4.1. Let M_1, M_2 be two fields. Let $f : R_1 \rightarrow R_2$ be a partial isomorphism. Then there is a larger partial isomorphism $g : K_1 \rightarrow K_2$ such that K_1 and K_2 are fields.

Proof. Let $K_i = \{a/b : a, b \in R_i, b \neq 0\}$. Then K_i is a subfield of M_i . For example, K_i is closed under addition because

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd}.$$

Let $g : K_1 \rightarrow K_2$ be defined by $g(a/b) = f(a)/f(b)$. This is well-defined because

$$\begin{aligned} \frac{a}{b} = \frac{a'}{b'} &\iff ab' = a'b \implies f(ab') = f(a'b) \\ &\iff f(a)f(b') = f(a')f(b) \iff \frac{f(a)}{f(b)} = \frac{f(a')}{f(b')}. \end{aligned}$$

It is an exercise in algebra to see that g is an isomorphism. For example, g preserves addition because

$$\begin{aligned} g\left(\frac{a}{b} + \frac{c}{d}\right) &= g\left(\frac{ad + bc}{bd}\right) = \frac{f(ad + bc)}{f(bd)} = \frac{f(a)f(d) + f(b)f(c)}{f(b)f(d)} \\ &= \frac{f(a)}{f(b)} + \frac{f(c)}{f(d)} = g\left(\frac{a}{b}\right) + g\left(\frac{c}{d}\right). \end{aligned} \quad \square$$

Lemma 9.4.2. *Let M_1, M_2 be two uncountable algebraically closed fields extending a countable subfield K . For any $a \in M_1$, there is $b \in M_2$ and an isomorphism $f : K[a] \rightarrow K[b]$ sending a to b and fixing K .*

Proof. Recall the notation

$$I_{a/K} = \{P(x) \in K[x] : P(a) = 0\}$$

from Theorem 9.3.2.

Claim. There is $b \in M_2$ with $I_{a/K} = I_{b/K}$.

Proof. First suppose a is transcendental. There are countably many non-zero polynomials in $K[x]$, and each has finitely many roots (Theorem 9.1.6). Therefore only countably many $b \in M_2$ are algebraic over K . Take $b \in M_2$ transcendental over K . Then $I_{a/K} = \{0\} = I_{b/K}$ by Theorem 9.3.2.

Next suppose a is algebraic with minimal polynomial $P(x) \in K[x]$. As M_2 is algebraically closed, there is $b \in M_2$ with $P(b) = 0$. By Theorem 9.3.4, $P(x)$ is the minimal polynomial of b , and then

$$I_{a/K} = P(x) \cdot K[x] = I_{b/K}. \quad \square_{\text{Claim}}$$

Fix $b \in M_2$ as in the claim, and let $I = I_{a/K} = I_{b/K}$. By Theorem 9.3.2, we have isomorphisms

$$\begin{aligned} K[x]/I &\rightarrow K[a] \\ K[x]/I &\rightarrow K[b] \end{aligned}$$

sending $P(x)$ to $P(a)$ and $P(b)$, respectively. The composition

$$K[a] \rightarrow K[x]/I \rightarrow K[b]$$

is the desired isomorphism. \square

Lemma 9.4.3. *Let M_1, M_2 be two uncountable algebraically closed fields. Let $f : R_1 \rightarrow R_2$ be a finitely-generated partial isomorphism. For any $a \in M_1$, there is $b \in M_2$ and an isomorphism $R_1[a] \rightarrow R_2[b]$ extending f .*

Proof. By Lemma 9.4.1, the isomorphism $f : R_1 \rightarrow R_2$ extends to an isomorphism $f' : K_1 \rightarrow K_2$ where the K_i are fields. Moving M_2 , we may assume $K_1 = K_2$ and f' is the identity map. Applying Lemma 9.4.2, there is an element $b \in M_2$ and an isomorphism $K_1[b] \rightarrow K_2[b]$ extending f' . This isomorphism restricts to an isomorphism $R_1[b] \rightarrow R_2[b]$ extending f . \square

Theorem 9.4.4. *The theory ACF has quantifier elimination.*

Proof. We use the criterion of Theorem 8.3.3. Suppose $M, N \models \text{ACF}$, $\bar{a} \in M^n$, $\bar{b} \in N^n$, and $\text{qftp}^M(\bar{a}) = \text{qftp}^N(\bar{b})$. We must show $\text{tp}^M(\bar{a}) = \text{tp}^N(\bar{b})$. Recall that M and N are infinite (Theorem 9.2.3). Replacing M, N with elementary extensions, we may assume M and N are uncountable. By Theorem 8.3.4, there is a partial isomorphism $f : \langle \bar{a} \rangle_M \rightarrow \langle \bar{b} \rangle_N$ with $f(\bar{a}) = \bar{b}$. Let \mathcal{F} be the collection of finitely generated partial isomorphisms from M to N . Then \mathcal{F} is a back-and-forth system: Lemma 9.4.3 gives the forward condition, and the backward condition holds by symmetry. By Theorem 3.7.6, the fact that $f \in \mathcal{F}$ implies that f is a partial elementary map, and so $\text{tp}^M(\bar{a}) = \text{tp}^N(\bar{b})$. \square

Corollary 9.4.5. *If $M, N \models \text{ACF}$, then $M \equiv N \iff \text{char}(M) = \text{char}(N)$.*

Proof. By quantifier elimination in ACF and Theorem 8.3.5, we have

$$M \equiv N \iff \langle \emptyset \rangle_M \cong \langle \emptyset \rangle_N.$$

By Theorem 2.7.5, the right hand side is equivalent to

$$\mathbb{Z}/n_M\mathbb{Z} \stackrel{?}{\cong} \mathbb{Z}/n_N\mathbb{Z}, \quad (*)$$

where $n_M = \text{char}(M)$ and $n_N = \text{char}(N)$. Clearly, $(*)$ holds if and only if $n_M = n_N$. \square

Definition 9.4.6. For $p \in \{0, 2, 3, 5, 7, 11, 13, \dots\}$, ACF_p is the theory of algebraically closed fields of characteristic p :

$$\begin{aligned} \text{ACF}_0 &= \text{ACF} \cup \left\{ \underbrace{1 + 1 + \dots + 1}_{p \text{ times}} \neq 0 : p = 2, 3, 5, 7, \dots \right\} \\ \text{ACF}_p &= \text{ACF} \cup \left\{ \underbrace{1 + 1 + \dots + 1}_{p \text{ times}} = 0 \right\} \text{ for } p > 0 \end{aligned}$$

For example, $\mathbb{C} \models \text{ACF}_0$. We will see later (Corollary 10.3.5) that ACF_p is consistent for each p .

Corollary 9.4.7. *Each theory ACF_p is complete.*

Corollary 9.4.8. *$\text{Th}(\mathbb{C})$ is decidable.*

Chapter 10

Existentially closed models

For a fixed theory T , a model M is *existentially closed* if “everything which could exist does exist,” in a certain sense. More precisely, any quantifier free $\mathcal{L}(M)$ -formula which is satisfied in an extension of M is already satisfied in M . For example, a field K is existentially closed if any system of equations which is satisfied in an extension of K is already satisfied in K . It turns out that the existentially closed fields are precisely the algebraically closed fields (Theorem 10.3.6), so “existentially closed” can be seen as a generalization of “algebraically closed from fields to other theories.

There are a couple important things to say about existentially closed models in general. First, under some technical assumptions on T , existentially closed models exist (Theorem 10.2.5). Second, if T has quantifier elimination, then all models of T are existentially closed.

10.1 Unions of chains of structures

Fix a language \mathcal{L} .

Definition 10.1.1. A *chain of \mathcal{L} -structures* is a family $\{M_i\}_{i \in I}$ where (I, \leq) is a linear order, M_i is an \mathcal{L} -structure for $i \in I$, and M_i is a substructure of M_j for $i \leq j$.

If $\{M_i\}_{i \in I}$ is a chain of \mathcal{L} -structures, we can make $M = \bigcup_{i \in I} M_i$ into a \mathcal{L} -structure by defining

$$\begin{aligned} f^M(a_1, \dots, a_n) &= f^{M_i}(a_1, \dots, a_n) \\ R^M(a_1, \dots, a_n) &\iff R^{M_i}(a_1, \dots, a_n) \end{aligned}$$

for any $i \in I$ large enough that $\{a_1, \dots, a_n\} \subseteq M_i$. The choice of i doesn't matter—if j is another choice, then

$$\begin{aligned} f^{M_i}(\bar{a}) &= f^{M_j}(\bar{a}) \\ R^{M_i}(\bar{a}) &\iff R^{M_j}(\bar{a}) \end{aligned}$$

because M_i is a substructure or extension of M_j .

Definition 10.1.2. A theory T is *inductive* if whenever $\{M_i\}_{i \in I}$ is a chain of models of T , the union $\bigcup_{i \in I} M_i$ is also a model of T .

Definition 10.1.3. An $\forall\exists$ -sentence is one of the form $\forall\bar{x} \exists\bar{y} \varphi(\bar{x}, \bar{y})$, where φ is quantifier-free. An $\forall\exists$ -theory is a set of $\forall\exists$ -sentences.

Theorem 10.1.4. *If T is an $\forall\exists$ -theory, then T is inductive.*

Proof. Suppose $\{M_i\}_{i \in I}$ is a chain of structures, and

$$M_i \models \forall\bar{x} \exists\bar{y} \varphi(\bar{x}, \bar{y}) \tag{*}$$

for all i , where φ is quantifier-free. Let $M = \bigcup_i M_i$. If $\bar{a} \in M^n$, then $\bar{a} \in M_i^n$ for large enough i . By (*), there is $\bar{b} \in M_i^m$ with $M_i \models \varphi(\bar{a}, \bar{b})$. Then $M \models \varphi(\bar{a}, \bar{b})$ because φ is quantifier-free. We have shown

$$M \models \forall\bar{x} \exists\bar{y} \varphi(\bar{x}, \bar{y}). \quad \square$$

Example 10.1.5. The theory of fields is an $\forall\exists$ -theory, so it is inductive.

10.2 Existentially closed models

Definition 10.2.1. Let $M \subseteq N$ be structures. Then M is *existentially closed* in N , written $M \preceq_1 N$, if for any quantifier-free $\mathcal{L}(M)$ -formula $\varphi(\bar{x})$,

$$N \models \exists\bar{x} \varphi(\bar{x}) \implies M \models \exists\bar{x} \varphi(\bar{x}).$$

Remark 10.2.2. $M \preceq N \implies M \preceq_1 N$.

Fix a theory T . “**Model**” will always mean **model of T** .

Definition 10.2.3. A model $M \models T$ is *existentially closed* if for any larger model $N \models T$ with $N \supseteq M$, we have $M \preceq_1 N$.

Theorem 10.2.4. *If T has quantifier-elimination, then every model of T is existentially closed.*

Proof. Suppose $M \subseteq N$ are models of T . Then $M \preceq N$ by quantifier elimination (Theorem 8.3.6), and so $M \preceq_1 N$. \square

Theorem 10.2.5. *If T is inductive, then any model M embeds into an existentially closed model $N \supseteq M$.*

Proof. If $M \models T$ and $\varphi(\bar{x})$ is a quantifier-free $\mathcal{L}(M)$ -formula, say that $\varphi(\bar{x})$ is *realizable beyond M* if it is realized in some larger model $N \supseteq M$. We make two observations:

1. A model M is existentially closed iff every quantifier-free $\mathcal{L}(M)$ -formula that is realizable beyond M is already realized in M .
2. If N extends M and a quantifier-free $\mathcal{L}(M)$ -formula φ is realizable beyond N , then it is realizable beyond M —the extension of N where φ is realized is an extension of M .

Claim. For any model M , there is a larger model $M^* \supseteq M$ such that every quantifier-free $\mathcal{L}(M)$ -formula that is realizable beyond M^* is realized in M^* .

Proof. Let $\{\varphi_\alpha\}_{\alpha < \kappa}$ enumerate all quantifier-free $\mathcal{L}(M)$ -formulas. Build an increasing chain of models $\{M_\alpha\}_{\alpha < \kappa}$ as follows:

1. $M_0 = M$.
2. If φ_α is realizable beyond M_α , then $M_{\alpha+1}$ is an extension of M_α with a realization of φ_α . Otherwise take $M_{\alpha+1} = M_\alpha$.
3. If α is a limit ordinal, take $M_\alpha = \bigcup_{\beta < \alpha} M_\beta$. This is a model because the theory is inductive.

Let $M^* = \bigcup_{\alpha < \kappa} M_\alpha$. Again, this is a model because T is inductive. Suppose φ is a quantifier-free $\mathcal{L}(M)$ -formula that is realizable beyond M^* . Then $\varphi = \varphi_\alpha$ for some α . By observation (2) above, φ_α is realizable beyond $M_\alpha \subseteq M^*$. Then φ_α is realized in $M_{\alpha+1}$ by construction. That is, $M_{\alpha+1} \models \varphi_\alpha(\bar{c})$ for some tuple \bar{c} in $M_{\alpha+1}$. As M^* extends $M_{\alpha+1}$ and φ_α is quantifier-free, $M^* \models \varphi_\alpha(\bar{c})$.

\square Claim

Using the claim, build an increasing chain of length ω :

$$M = M_0 \subseteq M_1 \subseteq M_2 \subseteq \cdots$$

where $M_{i+1} = M_i^*$. Let $N = \bigcup_{n=0}^{\infty} M_n$. Again, $N \models T$ because T is inductive. We claim N is existentially closed. Take a quantifier-free $\mathcal{L}(N)$ -formula φ that is realizable beyond N . Then φ is an $\mathcal{L}(M_i)$ -formula for some $i < \omega$. By observation (2), φ is realizable beyond $M_{i+1} = M_i^*$, so φ is realized in M_{i+1} . Then φ is realized in N , and N is existentially closed by observation (1). \square

10.3 Existentially closed fields

Lemma 10.3.1. *Let K, L be fields, and $\alpha : K \rightarrow L$ be a ring homomorphism. Then α is injective.*

Proof. Let $I = \ker(\alpha)$. We claim $I = \{0\}$. Otherwise take $a \in I \setminus \{0\}$. Then $1 = a^{-1}a \in I$, so $1^L = \alpha(1^K) = 0^L$, contradicting the definition of “field.”

As $\ker(\alpha) = \{0\}$, it follows that

$$\alpha(x) = \alpha(y) \iff x - y \in \ker(\alpha) \iff x - y = 0 \iff x = y.$$

This means that α is injective. \square

Lemma 10.3.2. *If K is a field and $P(x)$ is a polynomial of positive degree, then there is a larger field $L \supseteq K$ in which $P(x)$ has a root.*

Proof. Write $P(x)$ as a product $\prod_{i=1}^n Q_i(x)$ of irreducible factors. Replacing $P(x)$ with one of its irreducible factors, we may assume $P(x)$ is irreducible. Then $K[x]/P(x)K[x]$ is a field (Theorem 9.1.10). The composition

$$K \xrightarrow{\subseteq} K[x] \rightarrow K[x]/P(x)K[x]$$

is a homomorphism of fields, hence an embedding (Lemma 10.3.1). Up to isomorphism, $L := K[x]/P(x)K[x]$ is a field extending K . The fact that

$$P(x) \equiv 0 \pmod{P(x)} \text{ in } K[x]$$

means that

$$P(x) = 0 \text{ in } L$$

and so the element x is a root of P in L . \square

An *existentially closed field* is a field that is existentially closed among the class of all fields.

Theorem 10.3.3. *If K is an existentially closed field, then K is algebraically closed.*

Proof. Let $P(x) \in K[x]$ be a polynomial of positive degree. By Lemma 10.3.2, the quantifier-free $\mathcal{L}(K)$ -formula $P(x) = 0$ is realized in a field extending K . As K is existentially closed, it is realized in K , meaning that $P(x)$ has a root. \square

Corollary 10.3.4. *If K is a field, then there is an algebraically closed field L extending K .*

Proof. The class of fields is inductive, so Theorem 10.2.5 applies. \square

Corollary 10.3.5. *For every $p \in \{0, 2, 3, 5, 7, \dots\}$, there are algebraically closed fields of characteristic p .*

Proof. Take a field K of characteristic p (Theorem 2.7.7) and an algebraically closed field $M \supseteq K$ by Corollary 10.3.4. Then $\text{char}(M) = \text{char}(K) = p$ because characteristic doesn't change in field extensions (Theorem 2.7.10). \square

Theorem 10.3.6. *A field K is existentially closed if and only if it is algebraically closed.*

Proof. If K is existentially closed, then K is algebraically closed by Theorem 10.3.3. Conversely, suppose K is algebraically closed. We claim K is existentially closed. Let $L \supseteq K$ be an extension. By Corollary 10.3.4 there is an algebraically closed field $M \supseteq L \supseteq K$. Because ACF has quantifier elimination, algebraically closed fields are existentially closed *among algebraically closed fields*, and so $K \preceq_1 M$. But this implies $K \preceq_1 L$: if a quantifier-free $\mathcal{L}(K)$ -formula φ is realized in L , then it is realized in M , hence realized in K . \square

Chapter 11

Monster models

11.1 Pushing types along partial elementary maps

This chapter is about *monster models*, or more precisely, about κ -saturated and *strongly* κ -homogeneous models. The basic idea of saturated models is similar to existentially closed models:

Everything which could exist does exist.

For existentially closed models (Chapter 10), this meant

If a *quantifier-free formula* $\varphi(x)$ over M is satisfied in an *extension* of M , then it is satisfied in M .

For monster models, we want something slightly different:

If a *partial type* $\Sigma(x)$ over M is satisfied in an *elementary extension* of M , then it is satisfied in M .

As written above, this is usually impossible—the set of formulas $\{x \neq a : a \in M\}$ is satisfied in an elementary extension, but not satisfied in M . Instead, we only want to consider partial types that are “small.” This leads to the notion of κ -saturation for a fixed cardinal κ . A structure M is κ -saturated if every type over a small subset of M is realized in M , where “small” means “size less than κ .”

There is also a closely related notion called *strong* κ -homogeneity, which says that if two small tuples \bar{a}, \bar{b} in M have the same type, then there is

an isomorphism $\sigma : M \rightarrow M$ sending $\bar{a} \mapsto \bar{b}$. This means that M is highly symmetric.

Finally, a *monster model* is a model \mathbb{M} that is κ -saturated and strongly κ -homogeneous for some very big cardinal κ . Monster models exist (Section 11.4), and have a number of favorable properties. For example, every small model occurs as an elementary substructure of the monster \mathbb{M} (Theorem 11.2.4), definable sets satisfy a property like compactness in topology (Theorem 11.2.7), and we can detect the parameters needed to define a set by studying the symmetries of \mathbb{M} (Theorem 11.3.8). Because of these good properties, it is common in more advanced model theory to always work within a monster model. We will follow this approach in later chapters.

As an application of monster models, we give a relatively simple criterion for quantifier elimination in Theorem 11.5.4, and use it to study the model theory of discrete linear orders in Section 11.6.

Theorem 11.1.1. *Let M, N be \mathcal{L} -structures and $f : A \rightarrow B$ be a partial elementary map from M to N . If $\Sigma(\bar{x})$ is a set of $\mathcal{L}(A)$ -formulas, let $f_*\Sigma(\bar{x})$ be*

$$\{\varphi(\bar{x}, f(\bar{c})) : \bar{c} \in A, \varphi(\bar{x}, \bar{c}) \in \Sigma(\bar{x})\}.$$

1. *If $\Sigma(\bar{x})$ is finite, then $\Sigma(\bar{x})$ is realized in M if and only if $f_*\Sigma(\bar{x})$ is realized in N .*
2. *$\Sigma(\bar{x})$ is finitely realized in M if and only if $f_*\Sigma(\bar{x})$ is finitely realized in N .*
3. *f_* gives a bijection between partial types over A and partial types over B .*
4. *f_* gives a bijection between $S_n(A)$ and $S_n(B)$.*

Proof. 1. Write $\Sigma(\bar{x})$ as $\{\varphi_i(\bar{x}, \bar{c}_i) : 1 \leq i \leq n\}$. Then

$$M \models \exists \bar{x} \bigwedge_{i=1}^n \varphi_i(\bar{x}, \bar{c}_i) \iff N \models \exists \bar{x} \bigwedge_{i=1}^n \varphi_i(\bar{x}, f(\bar{c}_i)),$$

because f is a partial elementary map.

2. Clear from (1).

3. By (2), f_* is a map from partial types over A to partial types over B . The inverse map is $(f^{-1})_*$.
4. Clear from (3), since $S_n(A)$ and $S_n(B)$ are exactly the maximal partial types over A and B . \square

Corollary 11.1.2. *If $A \subseteq M \preceq N$, then $S_n^M(A) = S_n^N(A)$.*

Proof. The identity map $\text{id}_A : A \rightarrow A$ is a partial elementary map from M to N . \square

11.2 κ -saturated models

Let κ be an infinite cardinal.

Definition 11.2.1. A structure M is κ -saturated if for every $A \subseteq M$ with $|A| < \kappa$ and every $p \in S_1(A)$, p is realized in M .

Fix \mathcal{L} -structures N, M , where M is κ -saturated.

Lemma 11.2.2. *Let $f : A \rightarrow B$ be a partial elementary map from N to M , with $|A| < \kappa$. For any $\alpha \in N$ there is $\beta \in M$ such that $f \cup \{(\alpha, \beta)\}$ is a partial elementary map.*

Proof. Let $p = \text{tp}(\alpha/A) \in S_1(A)$. Let $f_*p = \{\varphi(x, f(\bar{a})) : \varphi(x, \bar{a}) \in p\}$. By Theorem 11.1.1, $f_*p \in S_1(B)$. As $|B| = |A| < \kappa$, there is $\beta \in M$ realizing f_*p . Then

$$\begin{aligned} N \models \varphi(\alpha, \bar{c}) &\iff \varphi(x, \bar{c}) \in p(x) \\ &\iff \varphi(x, f(\bar{c})) \in f_*p(x) \iff M \models \varphi(\beta, f(\bar{c})) \end{aligned}$$

for any \mathcal{L} -formula α and tuple \bar{c} in A , so $f \cup \{(\alpha, \beta)\}$ is a partial elementary map. \square

Lemma 11.2.3. *Let $f : A \rightarrow B$ be a partial elementary map from N to M . Suppose $A \subseteq A' \subseteq N$, $|A| < \kappa$, and $|A'| \leq \kappa$. Then f can be extended to a partial elementary map $f : A' \rightarrow B'$.*

Proof. Write A' as $\{a_\alpha : \alpha < \kappa\}$. Recursively choose partial elementary maps f_α for $\alpha < \kappa$ as follows:

- $f_0 = f$.
- $f_{\alpha+1} = f_\alpha \cup \{(a_\alpha, b)\}$ for some $b \in M$.
- $f_\beta = \bigcup_{\alpha < \beta} f_\alpha$ if β is a limit ordinal.

The successor step $\alpha + 1$ works because $\text{dom}(f_\alpha) \leq |A| + |\alpha| < \kappa$, so Lemma 11.2.2 applies.

Let $g = \bigcup_{\alpha < \kappa} f_\alpha$. Then g is a partial elementary map with domain A' . \square

Theorem 11.2.4 (κ -universality). *If M is κ -saturated and $N \equiv M$ with $|N| \leq \kappa$, then there is an elementary embedding $g : N \rightarrow M$. In particular, N is isomorphic to an elementary substructure of M .*

Proof. Note that $\emptyset : \emptyset \rightarrow \emptyset$ is a partial elementary map from N to M . Use Lemma 11.2.3 to extend to a partial elementary map g with $\text{dom}(g) = N$. Then g is an elementary embedding $N \rightarrow M$. \square

Theorem 11.2.5. *Suppose M is κ -saturated and $A \subseteq M$ with $|A| < \kappa$. For any finite $n < \omega$ and $p \in S_n(A)$, p is realized in M .*

Proof. Take $N \succeq M$ containing a realization $\bar{b} = p$. Note that $\text{id}_A : A \rightarrow A$ is a partial elementary map from N to M . By Lemma 11.2.3, we can extend it to a partial elementary map f with $\text{dom}(f) = A \cup \{a_1, \dots, a_n\}$. Then

$$\varphi(\bar{x}, \bar{b}) \in p(\bar{x}) \implies N \models \varphi(\bar{a}, \bar{b}) \iff M \models \varphi(f(\bar{a}), \bar{b})$$

for any formula φ and tuple \bar{b} in A , and so $f(\bar{a}) \in M^n$ realizes p . \square

Theorem 11.2.6. *Suppose M is κ -saturated, $A \subseteq M$ satisfies $|A| < \kappa$, and $\Sigma(\bar{x})$ is a partial type over A in at most κ variables. Then $\Sigma(\bar{x})$ is realized in M .*

Proof. Similar to Theorem 11.2.5. \square

Theorem 11.2.7 (κ -compactness). *Let M be κ -saturated.*

1. *Let $\Sigma(\bar{x})$ be a partial n -type over M . If $|\Sigma| < \kappa$, then $\Sigma(\bar{x})$ is realized in M .*
2. *Suppose $|I| < \kappa$ and X_i is a definable subset of M^n for each $i \in I$. If $\{X_i : i \in I\}$ has FIP, then $\bigcap_i X_i \neq \emptyset$.*

3. Suppose $X \subseteq M^n$ is definable, $|I| < \kappa$, $Y_i \subseteq M^n$ is definable for $i \in I$, and $X \subseteq \bigcup_{i \in I} Y_i$. Then there is a finite $I_0 \subseteq_f I$ such that $X \subseteq \bigcup_{i \in I_0} Y_i$.

Proof. 1. Let A be the set of parameters used in $\Sigma(\bar{x})$. Then $|A| < \kappa$, and $\Sigma(\bar{x})$ is a partial n -type over A . Take a complete n -type $p \in S_n(A)$ with $p \supseteq \Sigma(\bar{x})$. Then p is realized in M by Theorem 11.2.5.

2. Write X_i as $\varphi_i(M^n)$ for some $\mathcal{L}(M)$ -formula φ_i . Let $\Sigma = \{\varphi_i : i \in I\}$. The FIP means that Σ is finitely realized in M , i.e., a partial type. Apply (1) to find a point realizing Σ , i.e., a point in $\bigcap_{i \in I} X_i$.
3. If there is no finite subcover, then the family

$$\{X\} \cup \{M^n \setminus Y_i : i \in I\}$$

has the FIP. By (2), there is a point $\bar{a} \in X \cap \bigcap_i (M^n \setminus Y_i)$. Then $\bar{a} \in X$ but $\bar{a} \notin \bigcup_i Y_i$, a contradiction. \square

Corollary 11.2.8. *If M is κ -saturated and $D \subseteq M^n$ is definable, then one of two things happens:*

1. D is finite.
2. $|D| \geq \kappa$.

Proof. Otherwise, $\{\{p\} : p \in D\}$ is a small cover of D without a finite subcover, contradicting Theorem 11.2.7(3). \square

11.3 Strongly κ -homogeneous models

If M is a structure, then $\text{Aut}(M)$ denotes the set of automorphisms of M , i.e., isomorphisms from M to M . If $A \subseteq M$, then

$$\text{Aut}(M/A) := \{\sigma \in \text{Aut}(M) : \forall x \in A \sigma(x) = x\}$$

Definition 11.3.1. M is *strongly κ -homogeneous* if any partial elementary map f from M to M with $|\text{dom}(f)| < \kappa$ can be extended to an automorphism $\sigma \in \text{Aut}(M)$.

Theorem 11.3.2. *Suppose M is strongly κ -homogeneous. Suppose $\bar{a}, \bar{b} \in M^n$, $C \subseteq M$, and $|C| < \kappa$. Then the following are equivalent:*

1. $\text{tp}(\bar{a}/C) = \text{tp}(\bar{b}/C)$.
2. There is $\sigma \in \text{Aut}(M/C)$ with $\sigma(\bar{a}) = \bar{b}$.

Proof. (1) \implies (2): if $\text{tp}(\bar{a}/C) = \text{tp}(\bar{b}/C)$, then there is a partial elementary map

$$f : C \cup \{a_1, \dots, a_n\} \rightarrow C \cup \{b_1, \dots, b_n\}$$

$$f(x) = \begin{cases} x & \text{if } x \in C \\ b_i & \text{if } x = a_i \end{cases}$$

Then f extends to an automorphism $\sigma \in \text{Aut}(M)$. Note that $\sigma \supseteq f \supseteq \text{id}_C$, so $\sigma \in \text{Aut}(M/C)$, and $\sigma(\bar{a}) = f(\bar{a}) = \bar{b}$.

(2) \implies (1): isomorphisms preserve all formulas. \square

Definition 11.3.3. $\bar{a} \equiv_C \bar{b}$ means $\text{tp}(\bar{a}/C) = \text{tp}(\bar{b}/C)$.

Informal Definition 11.3.4. A *monster model* is a structure that is κ -saturated and strongly κ -homogeneous for some cardinal κ bigger than any cardinals we care about. A set X is “small” or “large” depending on whether $|X| < \kappa$ or $|X| \geq \kappa$.

Work in a monster model \mathbb{M} . Fix a small set $A \subseteq \mathbb{M}$.

Definition 11.3.5. A set $X \subseteq \mathbb{M}^n$ is *A-invariant* if $\sigma(X) = X$ for all $\sigma \in \text{Aut}(\mathbb{M}/A)$.

Theorem 11.3.6. *The following are equivalent for $X \subseteq \mathbb{M}^n$:*

1. X is *A-invariant*.
2. If $\bar{b}, \bar{c} \in \mathbb{M}^n$, then

$$\text{tp}(\bar{b}/A) = \text{tp}(\bar{c}/A) \implies (\bar{b} \in A \iff \bar{c} \in A).$$

3. There is a subset $X' \subseteq S_n(A)$ such that

$$X = \{\bar{b} \in \mathbb{M}^n : \text{tp}(\bar{b}/A) \in X'\}.$$

Proof. Note that (1) means the following: if $\bar{a} \in \mathbb{M}^n$ and $\sigma \in \text{Aut}(\mathbb{M}/C)$, then

$$\bar{a} \in X \iff \sigma(\bar{a}) \in X.$$

This is equivalent to (2) by Theorem 11.3.2. The equivalence of (2) and (3) is clear. \square

Remark 11.3.7. An A -definable set $\varphi(\mathbb{M}^n)$ corresponds to the clopen set $\llbracket \varphi \rrbracket \subseteq S_n(A)$, so A -definable sets are A -invariant.

Theorem 11.3.8. *If $D \subseteq \mathbb{M}^n$ is definable and A -invariant, then D is A -definable.*

Proof. Note that D is B -definable for some small $B \supseteq A$. Let $f : S_n(B) \rightarrow S_n(A)$ be the restriction map sending $\text{tp}(\bar{c}/B)$ to $\text{tp}(\bar{c}/A)$. This map is surjective because every complete type over A extends to a complete type over B . Additionally, it is continuous by Theorem 7.4.4, because the preimage of the clopen set $\llbracket \varphi \rrbracket \subseteq S_n(A)$ is the clopen set $\llbracket \varphi \rrbracket \subseteq S_n(B)$. As X is A -invariant and B -invariant, there are sets $X_A \subseteq S_n(A)$ and $X_B \subseteq S_n(B)$ such that

$$\begin{aligned} \bar{c} \in X &\iff \text{tp}(\bar{c}/A) \in X_A \\ \bar{c} \in X &\iff \text{tp}(\bar{c}/B) \in X_B. \end{aligned}$$

Then $X_B = f^{-1}(X_A)$. As f is surjective

$$\begin{aligned} X_A &= f(X_B) \\ S_n(A) \setminus X_A &= f(S_n(B) \setminus X_B). \end{aligned}$$

Because X is B -definable the set $X_B \subseteq S_n(B)$ is clopen, and so X_B and $S_n(B) \setminus X_B$ are closed. The image of a closed set is closed (Theorem 7.2.6), so X_A and its complement are both closed. Then X_A is a clopen set $\llbracket \psi \rrbracket$, and X is the A -definable set $\psi(\mathbb{M}^n)$. \square

11.4 Construction of monster models

A cardinal κ is *regular* if whenever $|I| < \kappa$ and $|X_i| < \kappa$ for every $i \in I$, we have $|\bigcup_{i \in I} X_i| < \kappa$. The cardinal \aleph_0 is regular, because a finite union of finite sets is finite. For any cardinal κ , the successor κ^+ is regular, because a union of at most κ -many sets of size at most κ has size at most $\kappa^2 = \kappa < \kappa^+$. Consequently, for any cardinal κ we can find a larger cardinal that is regular.

If κ is a regular cardinal and $A \subseteq \kappa$ with $|A| < \kappa$, then $\sup(A) < \kappa$. Otherwise, $\sup(A) = \kappa$, and then κ is a union $\bigcup_{\alpha \in A} \alpha$ where $|A| < \kappa$ and $|\alpha| < \kappa$ for each $\alpha \in \kappa$, contradicting regularity.

Lemma 11.4.1. *Let κ be a regular cardinal. Let $\{S_\alpha\}_{\alpha < \kappa}$ be an increasing chain of sets, indexed by κ . If $A \subseteq \bigcup_{\alpha < \kappa} S_\alpha$ and $|A| < \kappa$, then $A \subseteq S_\alpha$ for some $\alpha < \kappa$.*

Proof. Define $f : A \rightarrow \kappa$ by $f(x) = \min\{\alpha < \kappa : x \in S_\alpha\}$. Then $|f(A)| \leq |A| < \kappa$, so $\alpha := \sup f(A) < \kappa$. For any $x \in A$, we have $f(x) \in f(A)$ and so $f(x) \leq \alpha$. Then $x \in S_{f(x)} \subseteq S_\alpha$ for any $x \in A$, so $A \subseteq S_\alpha$. \square

Definition 11.4.2. A chain of structures $\{M_i\}_{i \in I}$ is *elementary* if $M_i \preceq M_j$ for $i \leq j$.

Theorem 11.4.3 (Tarski-Vaught). *Let $\{M_i\}_{i \in I}$ be an elementary chain of \mathcal{L} -structures. Let $M = \bigcup_{i \in I} M_i$. Then $M_i \preceq M$ for all $i \in I$.*

Proof sketch. For each i , note that $\text{eldiag}(M_i)$ is finitely satisfiable, complete, and has the witness property. Therefore the union $\bigcup_i \text{eldiag}(M_i)$ is finitely satisfiable, complete, and has the witness property. The canonical model is M . Then $M \succeq M_i$ because $M \models \text{eldiag}(M_i)$. \square

Lemma 11.4.4. *If M is a structure, there is $N \succeq M$ such that every type in $S_1(M)$ is realized in N .*

Proof. Let $\{p_i(x) : i \in I\}$ enumerate $S_1(M)$. Let $\bar{x} = (x_i : i \in I)$ be a tuple of variables, one for each $i \in I$. Let $\Sigma(\bar{x}) = \{p_i(x_i) : i \in I\}$. Then $\Sigma(\bar{x})$ is finitely satisfiable in M because each p_i is finitely satisfiable in M . Therefore, $\Sigma(\bar{x})$ is realized by some tuple $\bar{a} = (a_i : i \in I)$ in an elementary extension $N \succeq M$. The element a_i realizes p_i . \square

Theorem 11.4.5. *If M is a structure and κ is a cardinal, there is a κ -saturated $N \succeq M$.*

Proof. Replacing κ with κ^+ if necessary, we may assume κ is regular. Build an elementary chain $\{M_\alpha\}_{\alpha < \kappa}$ by recursion on α :

- $M_0 = M$.
- $M_{\alpha+1}$ is an elementary extension of M_α realizing every complete 1-type over M_α (using Lemma 11.4.4).

- If α is a limit ordinal, take $M_\alpha = \bigcup_{\beta < \alpha} M_\beta$, using the Tarski-Vaught theorem on chains (Theorem 11.4.3).

Let $N = \bigcup_{\alpha < \kappa} M_\alpha$. Then $N \succeq M_0 = M$ by Tarski-Vaught again. If $A \subseteq N$ and $|A| < \kappa$, then $A \subseteq M_\alpha$ for some $\alpha < \kappa$ by Lemma 11.4.1. If $p \in S_1(A)$, then p is a partial type over M_α , so it extends to a complete type $p' \in S_1(M_\alpha)$, which is then realized by some $b \in M_{\alpha+1} \subseteq N$. Therefore N is κ -saturated. \square

Lemma 11.4.6. *Let M be a structure and N be an $|M|^+$ -saturated elementary extension. Let f be a partial elementary map from M to M .*

1. *There is a partial elementary map g from N to N extending f , with $\text{dom}(g) = M$.*
2. *There is a partial elementary map h from N to N extending f , with $\text{im}(h) = M$.*

Proof. 1. Let $\kappa = |M|^+$. Then N is κ -saturated and $|M| < \kappa$. Note that f is a partial elementary map from M to N . By Lemma 11.2.3, we can extend f to a partial elementary map g from M to N with $\text{dom}(g) = M$.

2. This follows from (1) by symmetry. More precisely, apply part (1) to f^{-1} to get a partial elementary map g extending f^{-1} with $\text{dom}(g) = M$. Then set $h = g^{-1}$, so that $\text{im}(h) = \text{dom}(g) = M$. \square

Lemma 11.4.7. *For any M there is an elementary extension $N \succeq M$ with the following properties:*

1. *Every complete type over M is realized in N .*
2. *If f is a partial elementary map from M to M , then there is $\sigma \in \text{Aut}(N)$ extending f .*

Proof. Build an elementary chain $\{M_i\}_{i < \omega}$ by recursion on i :

- $M_0 = M$.
- M_{i+1} is an $|M_i|^+$ -saturated elementary extension of M_i . This is possible by Theorem 11.4.5.

Let $N = \bigcup_{i=0}^{\infty} M_i$. Then $M = M_0 \preceq N$ by the Tarski-Vaught theorem on chains (Theorem 11.4.3). Every complete type over M is already realized in M_1 , hence in N .

Let $f : A \rightarrow B$ be a partial elementary map from M to M . Recursively build an increasing chain of partial elementary maps $\{f_i\}_{i < \omega}$ with $\text{dom}(f_i), \text{im}(f_i) \subseteq M_i$ as follows:

- $f_0 = f$.
- If $n > 0$, then $f_n : M_n \rightarrow M_n$ is a partial elementary map extending $f_{n-1} : M_{n-1} \rightarrow M_{n-1}$ with

$$\begin{aligned} \text{dom}(f_n) &= M_{n-1} \text{ if } n \text{ is odd} \\ \text{im}(f_n) &= M_{n-1} \text{ if } n \text{ is even.} \end{aligned}$$

Take $\sigma = \bigcup_{n=0}^{\infty} f_n$. Then σ is a partial elementary map from N to N . The odd steps ensure $\text{dom}(\sigma) \supseteq \bigcup_n M_n = N$, and the even steps ensure $\text{im}(\sigma) \supseteq \bigcup_n M_n = N$. Thus $\text{dom}(\sigma) = \text{im}(\sigma) = N$, and $\sigma \in \text{Aut}(N)$. \square

Theorem 11.4.8. *If M is a structure and κ is a cardinal, then there is a κ -saturated, strongly κ -homogeneous elementary extension $N \succeq M$.*

Proof. Replacing κ with κ^+ if necessary, we may assume κ is regular. Build an elementary chain $\{M_\alpha\}_{\alpha < \kappa}$ by recursion on α as follows:

1. $M_0 = M$.
2. $M_{\alpha+1}$ is an elementary extension of M_α as in Lemma 11.4.7. In particular, every complete 1-type over M_α is realized in $M_{\alpha+1}$.
3. If α is a limit ordinal, then $M_\alpha = \bigcup_{\beta < \alpha} M_\beta$. This works by the Tarski-Vaught theorem on chains (Theorem 11.4.3).

Let $N = \bigcup_{\alpha < \kappa} M_\alpha$. As in the proof of Theorem 11.4.5, N is κ -saturated. We prove strong κ -homogeneity. Suppose f is a partial elementary map from N to N with $|\text{dom}(f)| = |\text{im}(f)| < \kappa$. By Lemma 11.4.1, there is $\alpha < \kappa$ with $\text{dom}(f) \cup \text{im}(f) \subseteq M_\alpha$. Then f is a partial elementary map from M_α to M_α . By Lemma 11.4.7, we can extend f to an automorphism $\sigma_{\alpha+1} \in \text{Aut}(M_{\alpha+1})$. Build an increasing chain $\{\sigma_\beta\}_{\alpha < \beta < \kappa}$ with $\sigma_\beta \in \text{Aut}(M_\beta)$ by repeatedly applying Lemma 11.4.7. Take $\sigma = \bigcup_{\alpha < \beta < \kappa} \sigma_\beta$. Then $\sigma \in \text{Aut}(N)$, and $\sigma \supseteq \sigma_{\alpha+1} \supseteq f$. Therefore N is strongly κ -homogeneous. \square

11.5 Saturation and back-and-forth

Theorem 11.5.1. *Let M and N be ω -saturated. For any $\bar{a} \in M^n$ and $\bar{b} \in N^n$ with $\text{tp}(\bar{a}) = \text{tp}(\bar{b})$, let $g_{\bar{a},\bar{b}}$ be the isomorphism from $\langle \bar{a} \rangle_M$ to $\langle \bar{b} \rangle_N$ sending \bar{a} to \bar{b} as in Theorem 8.3.4. Then the family $\mathcal{F} = \{g_{\bar{a},\bar{b}} : n \in \mathbb{N}, \bar{a} \in M, \bar{b} \in N, \text{tp}(\bar{a}) = \text{tp}(\bar{b})\}$ is a back-and-forth system.*

Proof. We verify the “forward” condition; “backward” is similar. Fix some $g_{\bar{a},\bar{b}}$. Let a' be an element of M . By Lemma 11.2.2 we can extend the partial elementary map $\bar{a} \mapsto \bar{b}$ to a partial elementary map $(\bar{a}, a') \mapsto (\bar{b}, b')$. Then $\text{tp}(\bar{a}, a') = \text{tp}(\bar{b}, b')$, and $g_{\bar{a}a',\bar{b}b'}$ is the desired partial isomorphism in \mathcal{F} extending $g_{\bar{a},\bar{b}}$ and containing a' in its domain. \square

Corollary 11.5.2. *If M, N are countable and ω -saturated, and $M \equiv N$, then $M \cong N$.*

Proof. Let \mathcal{F} be as in Theorem 11.5.1. By Theorem 5.6.4, it suffices to show that \mathcal{F} is non-empty. We only need to find some $n \in \mathbb{N}$, some $\bar{a} \in M^n$, and $\bar{b} \in N^n$ with $\text{tp}(\bar{a}) = \text{tp}(\bar{b})$. Take $n = 0$, $\bar{a} = () \in M^0$, and $\bar{b} = () \in N^0$. Then $\text{tp}(\bar{a}) = \text{Th}(M) = \text{Th}(N) = \text{tp}(\bar{b})$. \square

Corollary 11.5.3. *If M is countable and ω -saturated, then M is strongly ω -homogeneous.*

Proof. Let $f : A \rightarrow B$ be a partial elementary map from M to M , with A and B finite. Let \bar{a} be a tuple enumerating A . Then $\bar{b} := f(\bar{a})$ enumerates B , and $\text{tp}(\bar{a}) = \text{tp}(\bar{b})$. We must find $\sigma \in \text{Aut}(M)$ with $\sigma(\bar{a}) = \bar{b}$. If \mathcal{F} is the back-and-forth system between M and M given by Theorem 11.5.1, then $g_{\bar{a},\bar{b}} \in \mathcal{F}$. By Lemma 5.6.3, there is an isomorphism $\sigma : M \rightarrow M$ extending $g_{\bar{a},\bar{b}}$. Then $\sigma(\bar{a}) = \bar{b}$. \square

Theorem 11.5.4. *Let κ be an infinite cardinal. The following are equivalent:*

1. *T has quantifier elimination.*
2. *If M, N are κ -saturated structures, then the family \mathcal{F}_0 of all finitely generated partial isomorphisms from M to N is a back-and-forth system.*

Proof. (1) \implies (2): Note that M, N are ω -saturated. If $\bar{a} \in M^n$ and $\bar{b} \in N^n$ and $\text{qftp}(\bar{a}) = \text{qftp}(\bar{b})$, let $g_{\bar{a}, \bar{b}} : \langle \bar{a} \rangle_M \rightarrow \langle \bar{b} \rangle_N$ be the partial isomorphism sending \bar{a} to \bar{b} from Theorem 8.3.4. Then

$$\mathcal{F}_0 = \{g_{\bar{a}, \bar{b}} : \text{qftp}(\bar{a}) = \text{qftp}(\bar{b})\}.$$

By quantifier elimination, this is

$$\mathcal{F}_0 = \{g_{\bar{a}, \bar{b}} : \text{tp}(\bar{a}) = \text{tp}(\bar{b})\},$$

which is a back-and-forth system by Theorem 11.5.1.

(2) \implies (1): By Theorem 8.3.3, it suffices to show that if $M, N \models T$, if $\bar{a} \in M^n$ and $\bar{b} \in N^n$, then

$$\text{qftp}^M(\bar{a}) = \text{qftp}^N(\bar{b}) \implies \text{tp}^M(\bar{a}) = \text{tp}^N(\bar{b}).$$

Replacing M and N by elementary extensions, we may assume M and N are κ -saturated. By (2), the family \mathcal{F}_0 of finitely generated partial isomorphisms from M to N is a back-and-forth system. Suppose $\text{qftp}^M(\bar{a}) = \text{qftp}^N(\bar{b})$. Then \mathcal{F} contains the isomorphism $g_{\bar{a}, \bar{b}} : \langle \bar{a} \rangle_M \rightarrow \langle \bar{b} \rangle_N$. By Theorem 3.7.6, $g_{\bar{a}, \bar{b}}$ is a partial elementary map, so $\text{tp}(\bar{a}) = \text{tp}(\bar{b})$. \square

11.6 Application: discrete linear orders with endpoints

Definition 11.6.1. Let x, y be elements in a linear order (M, \leq) . Then y is a *successor* of x , and x is a *predecessor* of y , written $x \triangleleft y$, if $x < y$ but there is no $z \in M$ with $x < z < y$.

Let T be the theory of linear orders (M, \leq) such that the following hold:

1. $\min(M)$ and $\max(M)$ exist. In particular, M is non-empty.
2. Every $x \in M$ other than $\max(M)$ has a successor $y \triangleright x$.
3. Every $x \in M$ other than $\min(M)$ has a predecessor $y \triangleleft x$.

Example 11.6.2. Any finite non-empty linear order is a model of T .

Definition 11.6.3. Suppose $M \models T$ and $x, y \in M$. The *distance* between x and y , written $d(x, y)$, is the element of $\mathbb{N} \cup \{\infty\}$ defined as follows:

1. If $x = y$, then $d(x, y) = 0$.
2. If $x < y$, then $d(x, y)$ is $1 + |\{z \in M : x < z < y\}|$.
3. If $x > y$, then $d(y, x) = d(x, y)$.

Note that we define $d(x, y)$ to be the symbol “ ∞ ” when $[x, y]$ is infinite, rather than distinguishing cardinalities.

Let \mathcal{L}' be the base language of orders $\mathcal{L} = \{\leq\}$ plus two new constant symbols \min, \max and a binary relation R_n for each $n < \infty$. Expand any model $M \models T$ to an \mathcal{L}' -structure by interpreting the new symbols as follows:

1. $\min^M = \min(M)$.
2. $\max^M = \max(M)$.
3. $R_n^M(x, y) \iff d(x, y) = n$.

The resulting \mathcal{L}' -structures are the models of some \mathcal{L}' -theory T' .

Remark 11.6.4. Let M, N be models of T' . Suppose

$$\begin{aligned} \min(M) &= a_1 < a_2 < \cdots < a_n = \max(M) \\ \min(N) &= b_1 < b_2 < \cdots < b_n = \max(N) \\ d(a_i, a_{i+1}) &= d(b_i, b_{i+1}) \text{ for } 1 \leq i < n. \end{aligned}$$

Then there is a partial isomorphism

$$\begin{aligned} f : \{a_1, \dots, a_n\} &\rightarrow \{b_1, \dots, b_n\} \\ f(a_i) &= b_i. \end{aligned}$$

Moreover, all finite partial isomorphisms from M to N arise in this way.

Lemma 11.6.5. *Let M, N be \aleph_1 -saturated models of T' . Let \mathcal{F} be the collection of finite partial isomorphisms $f : A \rightarrow B$. Then \mathcal{F} is a back-and-forth system.*

Proof. By symmetry we only need to prove the forward condition. Let $f : A \rightarrow B$ be a finite partial isomorphism and α be an element of M . We must

find $f' \in \mathcal{F}$ extending f with $\alpha \in \text{dom}(f')$. We may assume $\alpha \notin \text{dom}(f)$; otherwise take $f' = f$. By Remark 11.6.4, f has the form

$$\begin{aligned} f : \{a_1, \dots, a_n\} &= \{b_1, \dots, b_n\} \\ f(a_i) &= b_i \end{aligned}$$

where

$$\begin{aligned} \min(M) &= a_1 < a_2 < \dots < a_n = \max(M) \\ \min(N) &= b_1 < b_2 < \dots < b_n = \max(N) \\ d(a_i, a_{i+1}) &= d(b_i, b_{i+1}) \text{ for } 1 \leq i < n. \end{aligned}$$

There must be some i such that $a_i < \alpha < a_{i+1}$. (The cases $\alpha < a_1 = \min(M)$ and $\alpha > a_n = \max(M)$ are impossible.) By Remark 11.6.4, we must find β such that

$$\begin{aligned} b_i &< \beta < b_{i+1} \\ d(b_i, \beta) &= d(a_i, \alpha) =: x \\ d(\beta, b_{i+1}) &= d(\alpha, a_{i+1}) =: y. \end{aligned}$$

Note that $d(b_i, b_{i+1}) = d(a_i, a_{i+1}) = x + y$. There are four cases:

1. $x, y < \infty$. Take β between b_i and b_{i+1} with $d(b_i, \beta) = x$. Then $d(\beta, b_{i+1}) = (x + y) - x = y$ as desired.
2. $x < \infty = y$. Then $d(b_i, b_{i+1}) = \infty$. Take β between b_i and b_{i+1} with $d(b_i, \beta) = x$. Then $d(\beta, b_{i+1}) = \infty$.
3. $y < \infty = x$. Similar.
4. $x = y = \infty$. As $d(b_i, b_{i+1}) = \infty$, there are

$$b_i \triangleleft c_1 \triangleleft c_2 \triangleleft \dots \leq \dots \triangleleft d_2 \triangleleft d_1 \triangleleft b_{i+1}.$$

The partial type $\{x > c_i : i < \omega\} \cup \{x < d_i : i < \omega\}$ is realized in M , by \aleph_1 -saturation. Let β be a realization. Then $b_i < \beta < b_{i+1}$, and $d(b_i, \beta) = d(\beta, b_{i+1}) = \infty$. \square

Theorem 11.6.6. *The theory T' has quantifier elimination.*

Corollary 11.6.7. *Let M be an infinite model of T . Let M_0 be the set of $x \in M$ such that $d(x, \min(M)) < \infty$ or $d(x, \max(M)) < \infty$. Then $M_0 \preceq M$.*

Proof. Working in the expanded language \mathcal{L}' , it is easy to see that M_0 is a substructure of M and $M_0 \models T'$. Then $M_0 \preceq M$ because submodels are elementary substructures in theories with quantifier elimination (Theorem 8.3.6(2)). \square

Definition 11.6.8. If M is a model of T , the *length* of M , written $\ell(M)$, is $d(\min(M), \max(M)) \in \mathbb{N} \cup \{\infty\}$.

Corollary 11.6.9. *Two models $M, N \models T$ are elementarily equivalent if and only if $\ell(M) = \ell(N)$.*

Proof. If $M \equiv N$, then certainly $\ell(M) = \ell(N)$, essentially because the relations R_n are definable, or simply because $\ell(M)$ is $|M| - 1$.

Conversely, suppose $\ell(M) = \ell(N)$. Let M' and N' be the expansions of M and N to models of T . Note that the minimal substructure $\langle \emptyset \rangle_{M'}$ contains only the two points $\min(M)$ and $\max(M)$. This substructure is determined up to isomorphism by $\ell(M)$. For example, $R_n(\min(M), \max(M))$ holds iff $\ell(M) = n$. Because $\ell(M) = \ell(N)$, we have $\langle \emptyset \rangle_{M'} \cong \langle \emptyset \rangle_{N'}$, and so $M' \equiv N'$ by quantifier elimination (because of Theorem 8.3.5). Restricting to \mathcal{L} -sentences, we see $M \equiv N$. \square

Corollary 11.6.10. *The theory of infinite models of T is complete and decidable.*

Corollary 11.6.11. *The class \mathcal{K} of models of T is the elementary class generated by finite non-empty linear orders.*

Proof. Certainly \mathcal{K} is an elementary class containing the finite non-empty linear orders. If \mathcal{K}' is a smaller such elementary class, take $M \in \mathcal{K} \setminus \mathcal{K}'$. Then M is infinite or else $M \in \mathcal{K}'$. Because \mathcal{K}' contains models of size $> n$ for each n , it must contain an infinite model N (Theorem 5.4.4). Then M, N are infinite models of T , so Corollary 11.6.10 shows $M \equiv N \in \mathcal{K}'$. Then $M \in \mathcal{K}'$, contradicting the choice of M . \square

Corollary 11.6.12. *If φ is a sentence in the language of orders, then the following are equivalent:*

- $T \vdash \varphi$.
- Every finite non-empty linear order satisfies φ .

Chapter 12

Countable categoricity

Recall that a theory is κ -categorical if it has exactly one model of cardinality κ . This chapter is about \aleph_0 -categoricity, which behaves very different from κ -categoricity for $\kappa > \aleph_0$. We have seen one example of an \aleph_0 -categorical theory already—the theory DLO of dense linear orders (Corollary 5.6.5). We showed that DLO is \aleph_0 -categorical via a back-and-forth argument. We will see in Theorem 12.3.5 that this is the *only* way that a theory can be \aleph_0 -categorical.

Moreover, \aleph_0 -categoricity of T is equivalent to a number of structural properties of T and its models. Specifically, the following are equivalent if T is a theory and M is a countable model:

1. T is \aleph_0 -categorical.
2. For each n , the type space $S_n(T)$ is finite.
3. For each n and finite set $A \subseteq_f M$, there are only finitely many A -definable subsets of M^n .
4. For each n , the action of $\text{Aut}(M)$ on M^n has only finitely many orbits.
5. Every countable model of T is ω -saturated.

This combination of facts is due to Engeler, Ryll-Nardzewski, and Svenonius, and is sometimes called the *Ryll-Nardzewski theorem*.

As an “application” of this theorem, we show that no theory of fields is \aleph_0 -categorical (Theorem 13.4.3). That is, if T is a theory whose models are infinite fields, then T has at least two countable models.

The proof of the Ryll-Nardzewski theorem involves another important fact, the *omitting types theorem*. A type $p \in S_n(T)$ is *omitted* in a model M if it is not realized in M . It turns out that whether a type can be omitted is closely related to the topology on $S_n(T)$. Say that p is *non-isolated* if p is a limit point of $S_n(T) \setminus \{p\}$, and *isolated* otherwise. If p is isolated, then p is realized in any model (Theorem 12.2.2), but if p is non-isolated, then p is omitted in some model (Theorem 12.2.7). In fact, given any countable set of non-isolated types, there is a model omitting all of them. We will apply this in the proof of the Ryll-Nardzewski theorem, but there are other applications.

12.1 Baire category theorem

Let S be a non-empty Stone space (Definition 7.4.2).

Definition 12.1.1. $X \subseteq S$ is *dense* if X intersects any non-empty clopen set $Y \subseteq S$.

Definition 12.1.2. A set $X \subseteq S$ is *comeager* if X contains a countable intersection of dense open sets.

Remark 12.1.3. Dense open sets are comeager. A countable intersection of comeager sets is comeager.

Theorem 12.1.4 (Baire category theorem). *If $X \subseteq S$ is comeager, then X is dense, hence non-empty.*

Proof. Because S is a Stone space, every open set is a union of clopen sets. Therefore every non-empty open set contains a non-empty clopen set.

Suppose $X \supseteq \bigcup_{i=1}^{\infty} U_i$ where each U_i is open and dense. Let V_0 be a non-empty clopen set. Then $V_0 \cap U_1$ is a non-empty open set. It contains a non-empty clopen set V_1 . Similarly, $V_1 \cap U_2$ contains a non-empty clopen set V_2 . Continuing, we can build a descending chain of clopen sets

$$V_0 \supseteq V_1 \supseteq V_2 \supseteq \cdots$$

with $V_i \subseteq U_i$. The family $\{V_0, V_1, V_2, \dots\}$ has the FIP, so the intersection is non-empty by compactness (Theorem 7.1.15). Take $p \in \bigcap_{i=0}^{\infty} V_i \subseteq V_0 \cap \bigcap_{i=1}^{\infty} U_i \subseteq V_0 \cap X$. Then X intersects V_0 as desired. \square

12.2 The omitting types theorem

Fix an \mathcal{L} -structure M , subset $A \subseteq M$, and $n < \omega$.

Definition 12.2.1. A complete type $p \in S_n(A)$ is *isolated* if $\{p\}$ is clopen.

Theorem 12.2.2. *If p is isolated, then p is realized in M .*

Proof. As $\{p\}$ is clopen, $\{p\} = \llbracket \varphi \rrbracket \subseteq S_n(A)$ for some $\mathcal{L}(A)$ -formula $\varphi(\bar{x}) \in p(\bar{x})$. Then $\varphi(\bar{x})$ is satisfied by $b \in M$, as p is finitely satisfiable. If $p' = \text{tp}(b/A) \in S_n(A)$, then $p' \in \llbracket \varphi \rrbracket = \{p\}$, so $p' = p$ and b realizes p . \square

Fix a complete theory T in a countable language \mathcal{L} .

Definition 12.2.3. If $p \in S_n(T)$ and $M \models T$, then M *omits* p if p isn't realized in M .

Let $S_\omega(T)$ be the space of complete ω -types $\text{tp}(a_0, a_1, a_2, \dots)$, i.e., complete types in the variables $\bar{x} = (x_0, x_1, x_2, \dots)$. Work in a monster model $\mathbb{M} \models T$.

Lemma 12.2.4. *There is a comeager set $W \subseteq S_\omega(T)$ such that if $\bar{c} \in \mathbb{M}^\omega$ and $\text{tp}(\bar{c}) \in W$, then $\{c_i : i \in \omega\} \preceq \mathbb{M}$.*

Proof.

Claim. For any formula $\varphi(\bar{x}, y)$, the following open set is dense:

$$U_\varphi := \llbracket \neg \exists y \varphi(\bar{x}, y) \rrbracket \cup \bigcup_{i=0}^{\infty} \llbracket \varphi(\bar{x}, x_i) \rrbracket.$$

Proof. Take non-empty $\llbracket \psi \rrbracket \subseteq S_\omega(T)$. Suppose for the sake of contradiction that $\llbracket \psi \rrbracket \cap U_\varphi = \emptyset$. Then $\llbracket \psi \rrbracket$ doesn't intersect the sets in the union, which means the following:

$$\begin{aligned} \psi(\bar{x}) \wedge (\neg \exists y \varphi(\bar{x}, y)) &\text{ is inconsistent} \\ \psi(\bar{x}) \wedge \varphi(\bar{x}, x_i) &\text{ is inconsistent, for each } i < \omega. \end{aligned}$$

Take $\bar{c} \in \mathbb{M}^\omega$ realizing $\psi(\bar{x})$. By the first line, $\mathbb{M} \models \exists y \varphi(\bar{c}, y)$. Thus $\mathbb{M} \models \varphi(\bar{c}, e)$ for some $e \in \mathbb{M}$. Take $i \gg 0$ so that x_i doesn't occur in $\psi(\bar{x})$ or $\varphi(\bar{x}, y)$. Changing c_i to e , we may assume $\mathbb{M} \models \varphi(\bar{c}, c_i)$, contradicting the second line. \square_{Claim}

Now let W be the comeager set $\bigcap_{\varphi} U_{\varphi}$. Suppose $\bar{c} \in \mathbb{M}^{\omega}$ and $\text{tp}(\bar{c}) \in W$. We claim that $M := \{c_i : i \in \omega\} \preceq \mathbb{M}$ by the Tarski-Vaught criterion (Theorem 5.3.1). Suppose $D \subseteq \mathbb{M}^1$ is M -definable, defined as $\varphi(\bar{c}, \mathbb{M})$ for some $\varphi(\bar{x}, y)$. If $D \neq \emptyset$, then $\mathbb{M} \models \exists y \varphi(\bar{c}, y)$, so $\text{tp}(\bar{c}) \notin \llbracket \neg \exists y \varphi(\bar{x}, y) \rrbracket$. But $\text{tp}(\bar{c}) \in U_{\varphi}$, so $\text{tp}(\bar{c}) \in \llbracket \varphi(\bar{x}, x_i) \rrbracket$ for some i , which means $\mathbb{M} \models \varphi(\bar{c}, c_i)$, or equivalently, $c_i \in D$. Thus $M = \{c_i : i \in \omega\}$ intersects every non-empty M -definable set $D \subseteq \mathbb{M}^1$. \square

Lemma 12.2.5. *For any $j_1, \dots, j_n < \omega$, let $f_{\bar{j}} : S_{\omega}(T) \rightarrow S_n(T)$ be the restriction map $\text{tp}(\bar{c}) \mapsto \text{tp}(c_{j_1}, \dots, c_{j_n})$.*

1. f is continuous.
2. If $X \subseteq S_{\omega}(T)$ is clopen, then $f(X) \subseteq S_n(T)$ is clopen.

Proof. 1. The preimage of $\llbracket \varphi(y_1, \dots, y_n) \rrbracket \subseteq S_n(T)$ is $\llbracket \varphi(x_{j_1}, \dots, x_{j_n}) \rrbracket \subseteq S_{\omega}(T)$.

2. If $X = \llbracket \varphi \rrbracket$, then

$$\begin{aligned} X &= \{\text{tp}(\bar{c}) : \bar{c} \in \mathbb{M}^{\omega}, \mathbb{M} \models \varphi(\bar{c})\} \\ f(X) &= \{\text{tp}(c_{j_1}, \dots, c_{j_n}) : \bar{c} \in \mathbb{M}^{\omega}, \mathbb{M} \models \varphi(\bar{c})\}. \end{aligned}$$

(The first line holds by saturation of \mathbb{M} .) Now the set

$$\{(c_{j_1}, \dots, c_{j_n}) : \bar{c} \in \mathbb{M}^{\omega}, \mathbb{M} \models \varphi(\bar{c})\}$$

is definable, defined by the formula

$$\psi(y_1, \dots, y_n) := \exists x_0, \dots, x_N \left(\varphi(\bar{x}) \wedge \bigwedge_{i=1}^n y_i = x_{j_i} \right),$$

where N is large enough to quantify away all x_i 's appearing in $\varphi(\bar{x})$. Then

$$f(X) = \{\text{tp}(\bar{a}) : \bar{a} \in \psi(\mathbb{M}^n)\} = \llbracket \psi \rrbracket. \quad \square$$

Lemma 12.2.6. *Let $p \in S_n(T)$ be non-isolated. There is a comeager set $V_p \subseteq S_{\omega}(T)$ such that if $\bar{c} \in \mathbb{M}^{\omega}$ and $\text{tp}(\bar{c}) \in V_p$, then p is not realized by any tuple in $C = \{c_i : i < \omega\}$.*

Proof.

Claim. For any $j_1, \dots, j_n \in \omega$, there is a dense open set $U_{\bar{j}} \subseteq S_\omega(T)$ such that if $\text{tp}(\bar{c}) \in U_{\bar{j}}$, then $(c_{j_1}, \dots, c_{j_n})$ doesn't realize p .

Proof. Let $f = f_{\bar{j}} : S_\omega(T) \rightarrow S_n(T)$ be the restriction map from Lemma 12.2.5. Points are closed (Theorem 7.1.11) so $S_n(T) \setminus \{p\}$ is open and the preimage $U := f^{-1}(S_n(T) \setminus \{p\})$ is open. For density, suppose $X \subseteq S_\omega(T)$ is clopen and non-empty, but $X \cap U = \emptyset$. Then $f(X) \subseteq \{p\}$. But $f(X)$ is clopen by Lemma 12.2.5(2), and non-empty, so $f(X) = \{p\}$ and p is isolated, a contradiction. Thus U is dense. Finally, if $\text{tp}(\bar{c}) \in U$, then $\text{tp}(c_{j_1}, \dots, c_{j_n}) \in S_n(T) \setminus \{p\}$, meaning that $(c_{j_1}, \dots, c_{j_n})$ doesn't realize p . □_{Claim}

Take $V_p = \bigcap_{\bar{j} \in \mathbb{N}^n} U_{\bar{j}}$. If $\text{tp}(\bar{c}) \in V_p$, then $(c_{j_1}, \dots, c_{j_n})$ doesn't realize p for any $\bar{j} \in \mathbb{N}^n$. □

Theorem 12.2.7 (Omitting types theorem). *Let p_1, p_2, \dots be a countable list of non-isolated complete types $p_i \in S_{n_i}(T)$. Then there is a countable model $M \models T$ omitting every p_i .*

Proof. Let W and V_p be the comeager sets from Lemmas 12.2.4 and 12.2.6. The set $Q = W \cap \bigcap_{i=1}^\infty V_{p_i}$ is comeager, hence non-empty by the Baire Category Theorem. Take $\bar{c} \in \mathbb{M}^\omega$ with $\text{tp}(\bar{c}) \in Q$, and let $M = \{c_i : i < \omega\}$. Then $M \preceq \mathbb{M}$ because $\text{tp}(\bar{c}) \in W$, and M omits p_i because $\text{tp}(\bar{c}) \in V_{p_i}$. □

12.3 Countably categorical theories

Work in a monster model $\mathbb{M} \models T$.

Lemma 12.3.1. *$S_n(A)$ is finite iff all types in $S_n(A)$ are isolated.*

Proof. Suppose $S_n(A)$ is finite. Every point is closed (Theorem 7.1.11), and a finite union of closed sets is closed, so every subset of $S_n(A)$ is closed, every subset is open, and every subset is clopen, implying every point is isolated.

Conversely, suppose every point is isolated. Then $S_n(A) = \bigcup_{p \in S_n(A)} \{p\}$ is an open cover. By compactness of $S_n(A)$, there is a finite subcover, and so $S_n(A)$ is finite. □

Lemma 12.3.2. *$S_n(A)$ is finite iff there are only finitely many A -definable sets $D \subseteq M^n$.*

Proof. The boolean algebra of A -definable sets is isomorphic to the boolean algebra of clopen sets in $S_n(A)$. If $S_n(A)$ is finite, then there are only finitely many clopen sets. Conversely, if there the number of clopen sets is $k < \infty$, then there are at most 2^k open sets, because every open set is a union of clopen sets. Then there are at most 2^k closed sets, hence 2^k points (as every point is closed by Theorem 7.1.11). \square

Lemma 12.3.3. *If $\bar{b}, \bar{c} \in \mathbb{M}^n$ and $A = \{a_1, \dots, a_m\} \subseteq \mathbb{M}$, then*

$$\bar{b} \equiv_A \bar{c} \iff \bar{a}\bar{b} \equiv_{\emptyset} \bar{a}\bar{c}.$$

Proof. Every $\mathcal{L}(A)$ -formula has the form $\varphi(\bar{a}, \bar{x})$ for some \mathcal{L} -formula $\varphi(\bar{w}, \bar{x})$. Therefore both sides say that for any \mathcal{L} -formula $\varphi(\bar{w}, \bar{x})$,

$$\mathbb{M} \models \varphi(\bar{a}, \bar{b}) \iff \mathbb{M} \models \varphi(\bar{a}, \bar{c}). \quad \square$$

Lemma 12.3.4. *If $|S_n(\emptyset)| < \infty$ for all n , then $|S_n(A)| < \infty$ for all n and all finite $A \subseteq \mathbb{M}$.*

Proof. Fix some $n < \omega$ and $A = \{a_1, \dots, a_m\} \subseteq M$. Then the map

$$\begin{aligned} S_n(A) &\rightarrow S_{m+n}(\emptyset) \\ \text{tp}(\bar{b}/A) &\mapsto \text{tp}(\bar{a}\bar{b}/\emptyset) \end{aligned}$$

is well-defined and injective by Lemma 12.3.3. Thus $|S_n(A)| \leq |S_{m+n}(\emptyset)| < \infty$. \square

Recall that $S_n(\emptyset) = S_n(T)$ (Corollary 8.1.20).

Theorem 12.3.5 (Engeler, Ryll-Nardzewski, Svenonius). *Let T be a complete theory in a countable language and let \mathbb{M} be a monster model. The following are equivalent:*

1. T has a unique countable model.
2. $S_n(T)$ is finite for all $n < \omega$.
3. $S_n(A)$ is finite for all $n < \omega$ and finite $A \subseteq \mathbb{M}$.
4. Every countable model of T is ω -saturated.

Proof. (1) \implies (2): Suppose $S_n(T)$ is infinite for some n . By Lemma 12.3.1, some $p \in S_n(T)$ is non-isolated. Then p is realized in some model M . By the downward Löwenheim-Skolem theorem (Theorem 5.4.3), we may assume M is countable. By the omitting types theorem (Theorem 12.2.7), there is a countable model N omitting p . Then $M \not\equiv N$, and (1) fails.

(2) \implies (3): Lemma 12.3.4.

(3) \implies (4): Assume (3). Let M be a countable model. If A is a finite subset of M , then $S_n(A)$ is finite by (3), so every $p \in S_n(A)$ is isolated by Lemma 12.3.1, and then every $p \in S_n(A)$ is realized by Theorem 12.2.2. This shows that M is ω -saturated.

(4) \implies (1): There is at most one countable ω -saturated model by Corollary 11.5.2. \square

Theorem 12.3.6. *Let M be a countable structure in a countable language. The following are equivalent:*

1. $\text{Th}(M)$ is countably categorical.
2. For every n , the action of $\text{Aut}(M)$ on M^n has finitely many orbits.

Proof. (1) \implies (2): by Theorem 12.3.5(4), M is ω -saturated, hence strongly ω -homogeneous by Corollary 11.5.3. Therefore $\bar{a}, \bar{b} \in M^n$ are in the same orbit of $\text{Aut}(M)$ iff $\text{tp}(\bar{a}) = \text{tp}(\bar{b})$. By Theorem 12.3.5(2), there are only finitely many complete n -types.

(2) \implies (1): if $D \subseteq M^n$ is 0-definable, then D is $\text{Aut}(M)$ -invariant so D is a union of orbits. There are only finitely many orbits, hence finitely many possibilities for D . Then the boolean algebra of 0-definable sets $D \subseteq M^n$ is finite. By Lemma 12.3.2, $S_n(\emptyset) = S_n(T)$ is finite. By Theorem 12.3.5, $\text{Th}(M)$ is countably categorical. \square

Chapter 13

Closure operators and pregeometries

This chapter is about the abstract notion of *closure operators*, which play a fundamental role in universal algebra and some parts of model theory. We have already seen several instances of closure operators. For example, the operation sending a set $S \subseteq M$ to the generated substructure $\langle S \rangle$ is a closure operation, and topologies can be regarded as a special kind of closure operation.

There are two closure operators which play a special role in model theory, the *definable closure* $\text{dcl}(-)$ and the *algebraic closure* $\text{acl}(-)$, which we discuss in Sections 13.2–13.3. The terminology “algebraic closure” comes from the example of ACF, where the model-theoretic algebraic closure $\text{acl}(-)$ agrees exactly with the field theoretic notion of algebraic closure K^{alg} (Theorem 13.5.2). We use this to prove a purely field-theoretic fact: the set \mathbb{Q}^{alg} of *algebraic numbers* is itself a field (Example 13.5.5), and is even an elementary substructure of \mathbb{C} .

The final two sections are on a special class of closure operators called *pregeometries* (or *matroids*), which are important in combinatorics and some branches of model theory. One of the motivating examples of pregeometries comes from *vectors spaces* (Definition 13.6.1), and we use pregeometries to classify vector spaces (Theorem 13.7.3). As an application, the theory of vector spaces is κ -categorical for certain κ (Corollary 13.7.6).

In the next chapter, we will tie these ideas together, and study theories where $\text{acl}(-)$ is a pregeometry. We will see that this happens in ACF, and more generally in any *strongly minimal* theory (Section 14.1).

13.1 Closure operators

Definition 13.1.1. A *closure operator* on a set S is an operation $\text{cl}(-) : \mathfrak{P}(S) \rightarrow \mathfrak{P}(S)$ with the following properties:

- Idempotent: $\text{cl}(\text{cl}(X)) = \text{cl}(X)$.
- Monotone: $X \subseteq Y \implies \text{cl}(X) \subseteq \text{cl}(Y)$.
- Increasing: $X \subseteq \text{cl}(X)$.

The set $\text{cl}(X)$ is called the *closure* of X .

Fix a set S and a closure operator $\text{cl}(-)$ on S . We say that $X \subseteq S$ is *closed* if $X = \text{cl}(X)$. By idempotence, the closed sets are exactly the sets of the form $\text{cl}(X)$.

Theorem 13.1.2. *If $X \subseteq S$, then $\text{cl}(X)$ is the smallest closed set containing X .*

Proof. The set $\text{cl}(X)$ contains X because $\text{cl}(-)$ is increasing, and $\text{cl}(X)$ is closed because $\text{cl}(-)$ is idempotent. Suppose Y is a closed set containing X . Then $Y = \text{cl}(Y) \supseteq \text{cl}(X)$, because $\text{cl}(-)$ is monotone. \square

Definition 13.1.3. A *closure system* on a set S is a family $\mathcal{C} \subseteq \mathfrak{P}(S)$ of sets, called *closed sets*, such that for any $X \subseteq S$ there is a smallest closed set containing X .

Remark 13.1.4. If \mathcal{C} is a closure system, define $\text{cl}(X)$ to be the smallest closed set containing X . It is easy to see that $\text{cl}(-)$ is a closure operator on S . This gives a map from closure systems to closure operators. Conversely, any closure operator $\text{cl}(-)$ defines a closure system $\{X \subseteq S : \text{cl}(X) = X\}$ by Theorem 13.1.2. It is easy to see that these two maps are inverses. Thus closure operators correspond bijectively with closure systems.

Theorem 13.1.5. *A family $\mathcal{C} \subseteq \mathfrak{P}(S)$ is a closure system iff \mathcal{C} is closed under infinite intersections.*

Proof. If \mathcal{C} is closed under infinite intersections, then the smallest closed set containing X exists—it is $\bigcap \{Y \in \mathcal{C} : Y \supseteq X\}$. Therefore \mathcal{C} is a closure system.

Conversely, suppose \mathcal{C} is a closure system. Let $\{X_i\}_{i \in I}$ be a family of closed sets and let $Y = \bigcap_{i \in I} X_i$. We claim that Y is closed. For each i , we have $\text{cl}(Y) \subseteq \text{cl}(X_i) = X_i$ by monotonicity. Thus $\text{cl}(Y) \subseteq \bigcap_{i \in I} X_i = Y$. Conversely $Y \subseteq \text{cl}(Y)$ because $\text{cl}(-)$ is increasing. \square

Example 13.1.6. If S is a topological space, the family of closed sets is a closure system. The closure $\text{cl}(X)$ is the smallest closed set containing X .

Example 13.1.7. Let M be a structure. Recall that $\langle A \rangle_M$ is the smallest substructure of M containing A . Therefore the class of substructures of M is a closure system, and $\langle - \rangle_M$ is a closure operation.

Definition 13.1.8. A closure operator $\text{cl}(-)$ on a set S is *finitary* if for any $a \in S$, $X \subseteq S$ with $a \in \text{cl}(X)$, there is a finite subset $X_0 \subseteq_f X$ with $a \in \text{cl}(X_0)$.

More generally, we say that an operation $F : \mathfrak{P}(S) \rightarrow \mathfrak{P}(S)$ is *finitary* if $F(X) \subseteq \bigcup \{F(X_0) : X_0 \subseteq_f X\}$.

Example 13.1.9. If M is a structure, the closure operation $\langle - \rangle_M$ is finitary: if $b \in \langle A \rangle$ then $b = t(\bar{a})$ for some term t and finite tuple \bar{a} . If $A_0 = \{a_1, \dots, a_n\}$, then $b \in \langle A_0 \rangle_M$.

Lemma 13.1.10. Suppose $\text{cl}(-) : \mathfrak{P}(S) \rightarrow \mathfrak{P}(S)$ is monotone, finitary, increasing, and satisfies the property

$$a \in \text{cl}(X) \text{ and } b \in \text{cl}(X \cup \{a\}) \implies b \in \text{cl}(X). \quad (*)$$

Then $\text{cl}(-)$ is idempotent, and therefore a finitary closure operator.

Proof. Note that $(*)$ implies

$$\text{cl}(X \cup \{a\}) = \text{cl}(X) \text{ for } a \in \text{cl}(X). \quad (\dagger)$$

Fix X . Then $\text{cl}(\text{cl}(X)) \supseteq \text{cl}(X)$ because $\text{cl}(-)$ is increasing. For the reverse inclusion, suppose $b \in \text{cl}(\text{cl}(X))$. As $\text{cl}(-)$ is finitary, $b \in \text{cl}(\{a_1, \dots, a_n\})$ for some $a_1, \dots, a_n \in \text{cl}(X)$. By n applications of (\dagger) ,

$$\text{cl}(X) = \text{cl}(X \cup \{a_1\}) = \text{cl}(X \cup \{a_1, a_2\}) = \dots = \text{cl}(X \cup \{a_1, \dots, a_n\}).$$

Then

$$b \in \text{cl}(\{a_1, \dots, a_n\}) \subseteq \text{cl}(X \cup \{a_1, \dots, a_n\}) = \text{cl}(X). \quad \square$$

13.2 Algebraic and definable closure

Definition 13.2.1. Let M be a structure and $A \subseteq M$ be a subset.

1. $b \in M$ is *definable* over A if $\{b\}$ is A -definable.
2. $b \in M$ is *algebraic* over A if $b \in D$ for some finite A -definable $D \subseteq M$.
3. The *definable closure* of A , written $\text{dcl}(A)$ or $\text{dcl}^M(A)$, is the set of $b \in M$ definable over A .
4. The *algebraic closure* of A , written $\text{acl}(A)$ or $\text{acl}^M(A)$, is the set of $b \in M$ algebraic over A .

Lemma 13.2.2. *If $b \in \text{acl}(A)$, then there is a finite A -definable set $X \ni b$ such that $\text{tp}(c/A) = \text{tp}(b/A)$ for every $c \in X$.*

Proof. Take a minimal finite A -definable set $X \ni b$. If X does not have the desired property, take $c \in x$ with $\text{tp}(c/A) \neq \text{tp}(b/A)$. Then there is an A -definable set D with $b \in D$ and $c \notin D$ (Remark 8.1.3). Then $X \cap D$ is a strictly smaller A -definable set containing b , a contradiction. \square

Lemma 13.2.3. *Fix a structure M .*

1. *If $A \subseteq B$ then $\text{acl}(A) \subseteq \text{acl}(B)$.*
2. *If $b \in \text{acl}(A)$, then $b \in \text{acl}(A_0)$ for some finite subset $A_0 \subseteq A$.*
3. *$A \subseteq \text{acl}(A)$.*
4. *If $b \in \text{acl}(A)$ and $c \in \text{acl}(A \cup \{b\})$, then $c \in \text{acl}(A)$.*

Proof. 1. Any finite A -definable set is a finite B -definable set.

2. Any finite A -definable set is A_0 -definable for some finite $A_0 \subseteq A$.

3. If $b \in A$, then the set $\{b\}$ is finite and A -definable, so $b \in \text{acl}(A)$.

4. As $b \in \text{acl}(A)$ and $c \in \text{acl}(A \cup \{b\})$, there is a finite A -definable set $X \ni b$ and a finite Ab -definable set $Y \ni c$. By Lemma 13.2.2, we may assume all elements of X have the same type over A . Write Y as $\varphi(M, b)$ for some $\mathcal{L}(A)$ -formula $\varphi(x, y)$. Let $n = |Y| = |\varphi(M, b)|$. Then $n = |\varphi(M, b')|$ for every $b' \in X$, since this is expressed by a formula in $\text{tp}(b'/A)$. Let $Z = \bigcup_{b' \in X} \varphi(M, b')$. Then Z is a finite A -definable set containing c . \square

Theorem 13.2.4. *Algebraic closure $\text{acl}(-)$ is a finitary closure operator.*

Proof. Lemma 13.2.3 plus Lemma 13.1.10. \square

A set $A \subseteq M$ is *algebraically closed* if $\text{acl}(A) = A$. The algebraic closure of A is the smallest algebraically closed set containing A .

Theorem 13.2.5. *Definable closure $\text{dcl}(-)$ is a finitary closure operator.*

Proof. Like Theorem 13.2.4 but slightly easier. \square

A set $A \subseteq M$ is *definably closed* if $\text{dcl}(A) = A$. The definable closure of A is the smallest definably closed set containing A .

13.3 Definable closure, algebraic closure, and substructures

Theorem 13.3.1. *Let M be a structure. If $A \subseteq M$ and $A = \text{dcl}(A)$, then A is a substructure of M .*

Proof. Let f be a k -ary function symbol, and let a_1, \dots, a_k be elements of A . Then $f(a_1, \dots, a_k)$ is in $\text{dcl}(A)$, as it is defined by the $\mathcal{L}(A)$ -formula $x = f(a_1, \dots, a_k)$. \square

Theorem 13.3.2. *Suppose $M \preceq N$ and φ is an $\mathcal{L}(M)$ -formula.*

1. $\varphi(M) \subseteq \varphi(N)$.
2. If $\varphi(M)$ is finite, then $\varphi(M) = \varphi(N)$.

Proof. 1. If $\bar{b} \in \varphi(M)$, then $M \models \varphi(\bar{b})$, so $N \models \varphi(\bar{b})$, and $\bar{b} \in \varphi(N)$.

2. Let $n = |\varphi(M)|$. Then $M \models \exists^{=n} \bar{x} \varphi(\bar{x})$, so the same sentence holds in N , and so $|\varphi(N)| = n$. As $\varphi(M) \subseteq \varphi(N)$, the two sets must be equal. \square

Theorem 13.3.3. *If $A \subseteq M \preceq N$, then $\text{acl}^M(A) = \text{acl}^N(A)$.*

Proof. If $b \in \text{acl}^M(A)$, then b is in a finite set of the form $\varphi(M)$ for some $\mathcal{L}(A)$ -formula $\varphi(x)$. By Theorem 13.3.2, $\varphi(N) = \varphi(M)$. Then $b \in \varphi(N) \implies b \in \text{acl}^N(A)$. Thus $\text{acl}^M(A) \subseteq \text{acl}^N(A)$.

The proof that $\text{acl}^N(A) \subseteq \text{acl}^M(A)$ is identical, exchanging M and N (even though there is no symmetry between M and N). \square

Theorem 13.3.4. *If $M \preceq N$, then M is algebraically closed as a subset of N .*

Proof. $\text{acl}^N(M) = \text{acl}^M(M) = M$. □

Theorem 13.3.5. *Let \mathbb{M} be a monster model, let A be a small subset, and let b be an element. Let S be the set of realizations of $\text{tp}(b/A)$.*

1. *If $b \in \text{acl}(A)$, then S is finite.*
2. *If $b \notin \text{acl}(A)$, then S is large.*

Proof. If $b \in \text{acl}(A)$, then there is a finite A -definable set $X \ni b$. Every realization of $\text{tp}(b/A)$ is in X , so $S \subseteq X$.

Conversely, suppose S is small. Let

$$\Sigma(x) = \text{tp}(b/A) \cup \{x \neq c : c \in S\}.$$

Then $\Sigma(x)$ is a set of formulas over the small set $A \cup S$ with no realizations in \mathbb{M} . By saturation, $\Sigma(x)$ isn't finitely realizable. Therefore there is $\varphi(x) \in \text{tp}(b/A)$ and $c_1, \dots, c_n \in S$ such that

$$\varphi(x) \cup \{x \neq c_i : 1 \leq i \leq n\}$$

isn't realized in \mathbb{M} . This means that $\varphi(\mathbb{M}) \subseteq \{c_1, \dots, c_n\}$. Then b is in the finite A -definable set $\varphi(\mathbb{M})$. □

Remark 13.3.6. By strong homogeneity, the set S in Theorem 13.3.5 is the set $\{\sigma(b) : \sigma \in \text{Aut}(\mathbb{M}/A)\}$.

13.4 Countably categorical fields

Theorem 13.4.1. *Let M be a field and A be a definably closed subset. Then A is a subfield.*

Proof. By Theorem 13.3.1, A is a substructure, i.e., a subring. It remains to show that A is closed under multiplicative inverses. If $a \in A$ is non-zero, then the $\mathcal{L}(A)$ -formula $xa = 1$ defines the set $\{a^{-1}\}$, so $a^{-1} \in \text{dcl}(A) = A$. □

Lemma 13.4.2. *Let T be \aleph_0 -categorical, let M be a model of T , and let A be a finite subset of M . Then $\text{dcl}(A)$ is finite.*

Proof. By Theorem 12.3.5(3), $S_1(A)$ is finite. Therefore there are only finitely many A -definable sets $D \subseteq M^1$. By definition, $b \in \text{dcl}(A)$ if and only if $\{b\}$ is definable. Thus $\text{dcl}(A)$ is finite. \square

Theorem 13.4.3. *There is no \aleph_0 -categorical theory of fields.*

Proof. Let T be an \aleph_0 -categorical theory of fields and let K be an \aleph_1 -saturated model. For any $n > 0$, the set $\{x \in K : x^n - 1 = 0\}$ is finite, by Theorem 9.1.6. Then the following partial type is finitely satisfiable in K :

$$\Sigma(x) = \{x \neq 0, x \neq 1, x^2 \neq 1, x^3 \neq 1, x^4 \neq 1, \dots\}.$$

Then $\Sigma(x)$ is realized in K . Take $b \in K$ realizing $\Sigma(x)$. Consider the group homomorphism

$$\begin{aligned} f : \mathbb{Z} &\rightarrow K^\times \\ f(n) &= b^n. \end{aligned}$$

As b realizes Σ , we have $f(b) = b^n \neq 1$ for $n \neq 0$. Therefore $\ker(f) = \{0\}$, and the image $f(\mathbb{Z})$ is isomorphic to $\mathbb{Z}/\{0\} \cong \mathbb{Z}$. In particular $f(\mathbb{Z})$ is infinite. However, $f(n) = b^n \in \text{dcl}(\{b\})$ for each n , and $\text{dcl}(\{b\})$ is finite. \square

Corollary 13.4.4. *If K is an infinite field, then there are two countable fields F_1, F_2 such that*

$$\begin{aligned} F_1 &\equiv F_2 \equiv K \\ F_1 &\not\equiv F_2. \end{aligned}$$

Proof. By Theorem 13.4.3, $\text{Th}(K)$ is not \aleph_0 -categorical. \square

13.5 Algebraic closure in ACF

If $M \models \text{ACF}$ and K is a subfield, then K^{alg} denotes the set of $a \in M$ that are algebraic over K , meaning that $P(a) = 0$ for some non-zero polynomial $P(x) \in K[x]$ (Definition 9.3.1).

Lemma 13.5.1. *Suppose $M \models \text{ACF}$ and K is a subfield. If $D \subseteq M^1$ is K -definable, then there is a finite subset $S \subseteq K^{\text{alg}}$ such that $D = S$ or $D = M \setminus S$.*

Proof. Let \mathcal{F} be the class of sets of the form S or $M \setminus S$ for finite $S \subseteq K^{\text{alg}}$. We must show $D \in \mathcal{F}$. Note that \mathcal{F} is closed under boolean combinations.

By quantifier-elimination, $D = \varphi(K)$ for some quantifier-free $\mathcal{L}(K)$ -formula φ , which is a boolean combination of atomic formulas. Because \mathcal{F} is closed under boolean combinations, we may assume that φ is atomic.

Then φ has the form $P(x) = Q(x)$ for some polynomials $P, Q \in K[x]$. If $P - Q$ is identically zero, then $\varphi(M) = M$. Otherwise, $\varphi(M)$ is the set of roots of $P - Q$, which is a subset of K^{alg} by definition of K^{alg} , and finite by Theorem 9.1.6. Either way, $\varphi(M) \in \mathcal{F}$. \square

Fix $M \models \text{ACF}$.

Theorem 13.5.2. *Let K be a subfield of M . Then $\text{acl}(K) = K^{\text{alg}}$.*

Proof. If $a \in K^{\text{alg}}$, then a is in a finite K -definable set of the form $\{x \in M : P(x) = 0\}$ for some $P(x) \in K[x]$, and so $a \in \text{acl}(K)$.

Conversely, suppose $a \in \text{acl}(K)$. Then $a \in D$ for some finite K -definable set D . By Lemma 13.5.1, $D \subseteq K^{\text{alg}}$, so $a \in K^{\text{alg}}$. \square

Theorem 13.5.3. *Let K be a subset of M . The following are equivalent:*

1. K is a subfield and $K = K^{\text{alg}}$.
2. K is a substructure and $K \models \text{ACF}$.
3. $K \preceq M$.
4. $K = \text{acl}(K)$.

Proof. We show $(1) \implies (2) \implies (3) \implies (4) \implies (1)$.

Assume (1). If $P(x) \in K[x]$ is non-constant, then there is $c \in M$ with $P(c) = 0$, because $M \models \text{ACF}$. The element c is in K^{alg} by definition, so $c \in K$. Thus $K \models \text{ACF}$.

Assume (2). Then $K \preceq M$ because ACF has quantifier elimination and both K and M are models of ACF (see Theorem 8.3.6).

Assume (3). Then $K = \text{acl}(K)$ by Theorem 13.3.4.

Assume (4). Then K is a subfield by Theorem 13.3.1 (definably closed sets are subfields) and $K = \text{acl}(K) = K^{\text{alg}}$ by Theorem 13.5.2. \square

Corollary 13.5.4. *If K is a subfield of M , then K^{alg} is also a subfield, and $K^{\text{alg}} \preceq M$.*

Proof. $\text{acl}(K^{\text{alg}}) = \text{acl}(\text{acl}(K)) = \text{acl}(K) = K^{\text{alg}}$, so K^{alg} satisfies condition (4) of Theorem 13.5.3. Then K^{alg} is a subfield and an elementary substructure by (1) and (3) of Theorem 13.5.3. \square

Example 13.5.5. The set of algebraic numbers $\mathbb{Q}^{\text{alg}} \subseteq \mathbb{C}$ is an algebraically closed field, and an elementary substructure of \mathbb{C} .

13.6 Pregeometries and vector spaces

Definition 13.6.1. Let K be a field. A K -vector space is an abelian group $(V, +)$ with a function

$$\cdot : K \times V \rightarrow V$$

satisfying the axioms

$$\begin{aligned} a \cdot (v + w) &= (a \cdot v) + (a \cdot w) \\ 1 \cdot v &= v \\ (a + b) \cdot v &= a \cdot v + b \cdot v \\ (a \cdot b) \cdot v &= a \cdot (b \cdot v). \end{aligned} \tag{\dagger}$$

For fixed K , we can regard the class of K -vector spaces as an equational class by thinking of the map $\cdot : K \times V \rightarrow V$ as a family of unary maps

$$\begin{aligned} \mu_a &: V \rightarrow V \\ \mu_a(v) &= a \cdot v, \end{aligned}$$

one for each $a \in K$. The axioms in Definition 13.6.1 become axiom schemas. For example, axiom (\dagger) becomes the axiom schema

$$\mu_{a+b}(x) = \mu_a(x) + \mu_b(x) \text{ for each } a, b \in K.$$

Subalgebras of vector spaces are called *linear subspaces*.

Remark 13.6.2. In a vector space, the following equations hold:

$$\begin{aligned} (-1) \cdot v &= -v \\ 0 \cdot v &= 0 \\ a \cdot (-v) &= -(a \cdot v) \\ a \cdot 0 &= 0. \end{aligned}$$

Fix a field K and vector space V .

Definition 13.6.3. The *span* of a set $S \subseteq V$, written $\text{span}(S)$, is the collection of elements of the form

$$a_1v_1 + \cdots + a_nv_n$$

where $n \geq 0$, $a_1, \dots, a_n \in K$, and $v_1, \dots, v_n \in S$.

Theorem 13.6.4. *The span of S is the linear subspace of V generated by S :*

$$\text{span}(S) = \langle S \rangle_V.$$

Proof. It is an exercise in algebra (using Remark 13.6.2) to see that $\text{span}(S)$ is a linear subspace. It contains S because if $v \in S$, then $v = 1 \cdot v \in \text{span}(S)$. Therefore $\text{span}(S) \supseteq \langle S \rangle_V$. On the other hand, every element of $\text{span}(S)$ has the form $t(v_1, \dots, v_n)$ for some term t and tuple \bar{v} in S , and so $\text{span}(S) \subseteq \langle S \rangle_V$. \square

Corollary 13.6.5. *Span is a finitary closure operator on V .*

Definition 13.6.6. A *pregeometry* is a pair (X, cl) where X is a set, cl is a finitary closure operator on X , and the following *exchange property* holds for $a, b \in X$ and $C \subseteq X$:

$$a \in \text{cl}(C \cup \{b\}) \setminus \text{cl}(C) \implies b \in \text{cl}(C \cup \{a\}).$$

Theorem 13.6.7. *In a vector space, $\text{span}(-)$ has the exchange property, and therefore defines a pregeometry.*

Proof. Suppose $v \in \text{span}(S \cup \{w\})$ but $v \notin \text{span}(S)$. Then

$$v = a_1v_1 + \cdots + a_nv_n + bw \tag{*}$$

for some $a_1, \dots, a_n, b \in K$ and $v_1, \dots, v_n \in S$. If $b = 0$, then $v \in \text{span}(S)$, a contradiction. Thus $b \neq 0$, and b^{-1} exists. Rearranging (*), we see

$$\begin{aligned} -bw &= a_1v_1 + \cdots + a_nv_n - v \\ w &= (-b^{-1}a_1)v_1 + \cdots + (-b^{-1}a_n)v_n + b^{-1}v \\ w &\in \text{span}(S \cup \{v\}). \end{aligned}$$

\square

Rank

Fix a pregeometry (X, cl) .

Definition 13.6.8. If $\bar{a} \in X^n$ and $B \subseteq X$, the *rank* of \bar{a} over B , written $\text{rk}(\bar{a}/B)$, is defined to be the number of $i \in \{1, \dots, n\}$ such that $a_i \notin \text{cl}(B \cup \{a_1, \dots, a_{i-1}\})$. We write $\text{rk}(\bar{a}/\emptyset)$ as $\text{rk}(\bar{a})$.

Theorem 13.6.9. $\text{rk}(\bar{a}, \bar{b}/C) = \text{rk}(\bar{a}/C) + \text{rk}(\bar{b}/C\bar{a})$.

Proof. Clear from the definition. \square

Theorem 13.6.10. $\text{rk}(\bar{a}/B) = 0 \iff \{a_1, \dots, a_n\} \subseteq \text{cl}(B)$.

Proof. If $a_i \in \text{cl}(B)$ for each i , then $a_i \in \text{cl}(B \cup \{a_1, \dots, a_{i-1}\})$ for each i , so $\text{rk}(\bar{a}/B) = 0$. Conversely, if $\text{rk}(\bar{a}/B) = 0$, then $a_i \in \text{cl}(B \cup \{a_1, \dots, a_{i-1}\})$ for each i . If S is a closed set containing B , then

$$\{a_1, \dots, a_{i-1}\} \subseteq S \implies a_i \in S$$

and so $\{a_1, \dots, a_n\} \subseteq S$ by induction. Taking $S = \text{cl}(B)$, we see $\{a_1, \dots, a_n\} \subseteq \text{cl}(B)$. \square

Lemma 13.6.11. $\text{rk}(a, b/C) = \text{rk}(b, a/C)$.

Proof. Both sides are in $\{0, 1, 2\}$. It suffices to show

$$\text{rk}(a, b/C) = 0 \iff \text{rk}(b, a/C) = 0 \quad (13.1)$$

$$\text{rk}(a, b/C) = 2 \iff \text{rk}(b, a/C) = 2 \quad (13.2)$$

By symmetry it suffices to show the \Rightarrow directions.

(13.1): by Theorem 13.6.10, both sides say $\{a, b\} \subseteq C$.

(13.2): suppose $\text{rk}(a, b/C) = 2$. Then $a \notin \text{cl}(C)$ and $b \notin \text{cl}(Ca)$. By monotonicity, $b \notin \text{cl}(C)$. If $a \in \text{cl}(Cb)$, then

$$a \in \text{cl}(Cb) \setminus \text{cl}(C) \text{ but } b \notin \text{cl}(Ca),$$

contradicting the exchange property. Thus $a \notin \text{cl}(Cb)$. With $b \notin \text{cl}(C)$, this implies $\text{rk}(b, a/C) = 2$. \square

Lemma 13.6.12. If \bar{a}, \bar{d} are tuples and b, c are elements and A is a set, then

$$\text{rk}(\bar{a}, b, c, \bar{d}/A) = \text{rk}(\bar{a}, c, b, \bar{d}/A).$$

Proof. Using Theorem 13.6.9 and Lemma 13.6.11,

$$\begin{aligned} \text{rk}(\bar{a}, b, c, \bar{d}/A) &= \text{rk}(\bar{a}/A) + \text{rk}(b, c/A, \bar{a}) + \text{rk}(\bar{d}/A, \bar{a}, b, c) \\ &= \text{rk}(\bar{a}/A) + \text{rk}(c, b/A, \bar{a}) + \text{rk}(\bar{d}/A, \bar{a}, c, b) \\ &= \text{rk}(\bar{a}, c, b, \bar{d}/A). \end{aligned} \quad \square$$

Theorem 13.6.13. *If π is a permutation of $\{1, \dots, n\}$, then*

$$\text{rk}(a_1, \dots, a_n/B) = \text{rk}(a_{\pi(1)}, \dots, a_{\pi(n)}/B).$$

Proof. Repeated applications of Lemma 13.6.12. \square

Theorem 13.6.14. *If $\bar{a} \subseteq \text{cl}(C\bar{b})$, then $\text{rk}(\bar{a}/C) \leq \text{rk}(\bar{b}/C)$.*

Proof.

$$\begin{aligned} \text{rk}(\bar{a}/C) &\leq \text{rk}(\bar{a}/C) + \text{rk}(\bar{b}/C\bar{a}) = \text{rk}(\bar{a}, \bar{b}/C) \\ &= \text{rk}(\bar{b}, \bar{a}/C) = \text{rk}(\bar{b}/C) + \text{rk}(\bar{a}/C\bar{b}) = \text{rk}(\bar{b}/C). \end{aligned} \quad \square$$

Theorem 13.6.15. *If $C \supseteq B$, then $\text{rk}(\bar{a}/C) \leq \text{rk}(\bar{a}/B)$.*

Proof. For each i , monotonicity of $\text{cl}(-)$ shows

$$a_i \notin \text{cl}(C \cup \{a_1, \dots, a_{i-1}\}) \implies a_i \notin \text{cl}(B \cup \{a_1, \dots, a_{i-1}\}). \quad \square$$

Independence and bases

Definition 13.6.16. A set $I \subseteq X$ is *independent* if $a \notin \text{cl}(I \setminus \{a\})$ for each $a \in I$.

Theorem 13.6.17. *In a vector space V , a set $I \subseteq V$ is independent if and only if the following condition holds: for any distinct $v_1, \dots, v_n \in V$ and $a_1, \dots, a_n \in K$,*

$$a_1v_1 + \dots + a_nv_n = 0 \implies v_1 = v_2 = \dots = v_n = 0. \quad (*)$$

Proof. If I is not independent, then there is some $v_1 \in \text{span}(I \setminus \{v_1\})$. Thus there are distinct $v_2, \dots, v_n \in I \setminus \{v_1\}$ with

$$v_1 = a_2v_2 + \dots + a_nv_n.$$

Setting $a_1 = -1 \in K$, we get $\sum_{i=1}^n a_i v_i = 0$, contradicting (*).

Conversely, suppose (*) fails. Permuting the v_i , we may assume $a_1 \neq 0$. Then

$$\begin{aligned} a_1 v_1 + \cdots + a_n v_n &= 0 \\ v_1 &= (-a_1^{-1} a_2) v_2 + \cdots + (-a_1^{-1} a_n) v_n \in \text{span}(v_2, \dots, v_n), \end{aligned}$$

and I is not independent. □

Now return to a general pregeometry.

Definition 13.6.18. A *basis* is a maximal independent set.

Remark 13.6.19. 1. If I is independent and $J \subseteq I$, then J is independent, by monotonicity of $\text{cl}(-)$.

2. A set $I \subseteq X$ is independent if and only if every finite subset is independent, because $\text{cl}(-)$ is finitary.

Theorem 13.6.20. *There is at least one basis.*

Proof. Zorn's lemma. □

Lemma 13.6.21. *If $a_1, \dots, a_n \in X$ are distinct, then $\text{rk}(\bar{a}) = n$ if and only if the set $I = \{a_1, \dots, a_n\}$ is independent.*

Proof. Note that $\text{rk}(\bar{a}) = n$ if and only if

$$a_i \notin \text{cl}(a_1, \dots, a_{i-1}) \text{ for all } 1 \leq i \leq n, \quad (*)$$

by definition of rank. Moreover, the condition $\text{rk}(\bar{a}) = n$ is invariant under permuting \bar{a} , by Theorem 13.6.13.

Independence clearly implies (*), as $\text{cl}(-)$ is monotone. For the converse, suppose $\text{rk}(\bar{a}) = n$ and $i \leq n$. We must show $a_i \notin \text{cl}(I \setminus \{a_i\})$. Permuting \bar{a} , we reduce to the case $i = n$. Then we must show $a_n \notin \text{cl}(a_1, \dots, a_{n-1})$, which is a case of (*). □

Lemma 13.6.22. *If I is independent and $a \notin \text{cl}(I)$, then $I \cup \{a\}$ is independent.*

Proof. By Remark 13.6.19(2), we may assume I is a finite set $\{b_1, \dots, b_n\}$. By Lemma 13.6.21 and Theorem 13.6.9

$$\text{rk}(\bar{b}, a) = \text{rk}(\bar{b}) + \text{rk}(a/\bar{b}) = n + 1,$$

and so $\{b_1, \dots, b_n, a\}$ is independent. \square

Theorem 13.6.23. *If B is a basis, then $\text{cl}(B) = X$.*

Proof. Otherwise, take $a \in X \setminus \text{cl}(B)$, and $B \cup \{a\}$ is independent, contradicting the fact that B is a maximal independent set. \square

Remark 13.6.24. Conversely, if B is independent and $\text{cl}(B) = X$, then B is a basis. Otherwise, if $I \supsetneq B$ is a larger independent set, then $I \subseteq X = \text{cl}(B)$, contradicting Lemma 13.6.25 below. Thus B is a basis if and only if B is independent and $\text{cl}(B) = X$.

Lemma 13.6.25. *If I is independent and J is a proper subset, then $I \not\subseteq \text{cl}(J)$.*

Proof. Otherwise, take $a \in I \setminus J$. Then $J \subseteq I \setminus \{a\}$, and so

$$a \in I \subseteq \text{cl}(J) \subseteq \text{cl}(I \setminus \{a\}),$$

contradicting independence. \square

Theorem 13.6.26. *If B and C are two bases, then $|B| = |C|$.*

Proof. Suppose not. Without loss of generality, $|B| < |C|$. There are two cases:

1. B and C are finite. Let \bar{b} and \bar{c} enumerate B and C . Then

$$\text{rk}(\bar{b}) = |B| < |C| = \text{rk}(\bar{c})$$

by Lemma 13.6.21. However, $C \subseteq X = \text{cl}(B)$ by Theorem 13.6.23, and so $\text{rk}(\bar{c}) \leq \text{rk}(\bar{b})$ by Theorem 13.6.14.

2. C is infinite. By Theorem 13.6.23, $B \subseteq \text{cl}(C)$ and $C \subseteq \text{cl}(B)$. As $\text{cl}(-)$ is finitary, we can find a finite subset $C_b \subseteq C$ for each $b \in B$ such that $b \in \text{cl}(C_b)$. Let $C' = \bigcup_{b \in B} C_b$. Then $B \subseteq \text{cl}(C')$, implying $C \subseteq \text{cl}(B) \subseteq \text{cl}(\text{cl}(C')) = \text{cl}(C')$. This contradicts independence of C (see Lemma 13.6.25) unless $C' = C$.

However, $|C'| < |C|$. Indeed, if B is finite then C' is finite, and so $|C'| < |C|$. If B is infinite then $|C'| \leq |B| < |C|$. Thus C' is a proper subset of C , a contradiction. \square

Definition 13.6.27. The *rank* of a pregeometry is the cardinality of any basis.

13.7 The classification of vector spaces

Fix a field K . We can regard K as a vector space by defining

$$\begin{aligned} a \cdot v &= a \cdot v \\ v + w &= v + w \end{aligned}$$

where the left-hand \cdot and $+$ are the vector space operations, and the right-hand \cdot and $+$ are the field operations.

If λ is a cardinal, let K^λ be the power vector space, i.e., the set of functions $f : \lambda \rightarrow K$ with the vector space operations defined pointwise:

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (a \cdot f)(x) &= a \cdot (f(x)). \end{aligned}$$

Say that $f \in K^\lambda$ has *finite support* if

$$\text{supp}(f) := \{x \in \lambda : f(x) \neq 0\}$$

is finite. Note that

$$\begin{aligned} \text{supp}(f + g) &\subseteq \text{supp}(f) \cup \text{supp}(g) \\ \text{supp}(af) &\subseteq \text{supp}(f) \\ \text{supp}(0) &= \emptyset. \end{aligned}$$

Therefore, $F_\lambda := \{f \in K^\lambda : \text{supp}(f) \text{ is finite}\}$ is a linear subspace of K^λ .

Lemma 13.7.1. Let V be a K -vector space. Suppose $v_i \in V$ for each $i \in \lambda$.

1. There is a homomorphism $\alpha : F_\lambda \rightarrow V$ defined by

$$\alpha(f) = \sum_{i \in \text{supp}(f)} f(i) \cdot v_i.$$

2. α is surjective if and only if $V = \text{span}\{v_i : i \in \lambda\}$.

3. α is injective if and only if $\{v_i : i \in \lambda\}$ is independent (and the v_i are distinct).
4. α is an isomorphism if and only if $\{v_i : i \in \lambda\}$ is a basis (and the v_i are distinct).
5. All homomorphisms $F_\lambda \rightarrow V$ arise as in part (1).

Proof. 1. An exercise in algebra. Essentially this works because $\alpha(f) = \sum_{i \in \lambda} f(i) \cdot v_i$, which depends linearly on f . The sum makes sense as almost all the terms in it are zero.

2. True by definition of span.
3. α is injective if and only if $\ker(\alpha) = \{0\}$. This condition means that if $f \in K^\lambda$ has finite support and

$$\sum_{i \in \lambda} f(i)v_i = 0,$$

then f must vanish. This is a rephrasing of Theorem 13.6.17.

4. α is an isomorphism if and only if it is injective and surjective. By the previous two points, this means that the set $B = \{v_i : i \in \lambda\}$ is independent and has $\text{span}(B) = V$. By Remark 13.6.24, this means that B is a basis.
5. Let $\beta : F_\lambda \rightarrow V$ be a homomorphism. Let $e_i : \lambda \rightarrow K$ be the function

$$e_i(x) = \begin{cases} 1 & \text{if } x = i \\ 0 & \text{if } x \neq i. \end{cases}$$

Then $\text{supp}(e_i) = \{i\}$, so $e_i \in F_\lambda$. Note that for any $f \in F_\lambda$,

$$f = \sum_{i \in \text{supp}(f)} f(i) \cdot e_i.$$

Therefore

$$\beta(f) = \sum_{i \in \text{supp}(f)} f(i) \cdot \beta(e_i).$$

Taking $v_i = \beta(e_i)$, we see that β has the desired form. □

Lemma 13.7.1 immediately yields the following:

Theorem 13.7.2. *Let V be a K -vector space. Then $V \cong F_\lambda$ if and only if V has a basis of cardinality λ .*

By Theorems 13.6.20 and 13.6.26, there is a unique cardinal λ such that V has a basis of cardinality λ .

Theorem 13.7.3. *Let V be a K -vector space. Then $V \cong F_\lambda$ for a unique cardinal λ , called the dimension of V .*

Lemma 13.7.4. *Let V be a K -vector space of dimension λ .*

1. *If λ is finite, then $|V| = |K|^\lambda$.*
2. *If λ or $|K|$ is infinite, then $|V| = |K| + \lambda$.*

Proof. Without loss of generality, $V = F_\lambda$.

1. $F_\lambda = K^\lambda$, whose cardinality is $|K|^\lambda$.
2. If $B \subseteq V$ is a basis of size λ , then $B \subseteq V \implies \lambda \leq |V|$. If $v \in V$ is non-zero, then there is an injection

$$\begin{aligned} K &\rightarrow V \\ a &\mapsto av, \end{aligned}$$

and so $|K| \leq |V|$. Thus $|V| \geq \max(|K|, \lambda) = |K| + \lambda$. On the other hand, V is generated by a set of size λ , and the language has size $\aleph_0 + |K|$, so that $|V| \leq \lambda + |K| + \aleph_0 = |K| + \lambda$. \square

Theorem 13.7.5. *If V is an infinite K -vector space and $|V| > |K|$, then $\dim(V) = |V|$.*

Proof. Let $\lambda = \dim(V)$. If K and λ are both finite then $|V| = |K|^\lambda < \aleph_0$, a contradiction. Therefore K or λ is infinite, and so $|V| = \max(|K|, \lambda)$. As $|V| > |K|$, this implies $|V| = \lambda$. \square

Corollary 13.7.6. *Let K be a field. The theory of infinite K -vector spaces is complete and λ -categorical for every infinite $\lambda > |K|$.*

Definition 13.7.7. A theory T in a countable language is *totally categorical* if T is λ -categorical for any λ .

Corollary 13.7.8. *If K is a finite field, then the theory of infinite K -vector spaces is totally categorical.*

Chapter 14

Strong minimality and geometric theories

A theory is *strongly minimal* if for every model M and definable set $D \subseteq M$, either D or $M \setminus D$ is finite. We have already seen one example of a strongly minimal theory, namely ACF (Lemma 13.5.1).

Although the definition of strong minimality only mentions definable sets in one variable, strong minimality has very strong consequences for definable sets $D \subseteq M^n$ in n variables. Specifically, strong minimality allows one to define the *dimension* $\dim(D)$ of any definable set. The intuition one should have is that $\dim(D)$ is something like topological dimension. For example, points have dimension 0, lines and circles have dimension 1, planes and spheres have dimension 2, and so on. In the case of algebraically closed fields, the model-theoretic dimension $\dim(-)$ corresponds to something that algebraic geometers know as “Krull dimension”.

The abstract model-theoretic dimension in strongly minimal theories satisfies many intuitive properties. For example, $\dim(D_1 \times D_2) = \dim(D_1) + \dim(D_2)$. Dimension is also invariant under definable bijections: if $f : D_1 \rightarrow D_2$ is a definable bijection, then $\dim(D_1) = \dim(D_2)$.

Another important property of strongly minimal theories is that they are κ -categorical for all $\kappa > \aleph_0$ (Corollary 14.2.7), at least when the theory is complete.¹ Even though \aleph_0 -categoricity usually fails, one can classify the countable models. For example, we classify all models of ACF (i.e., alge-

¹For example, the complete theory ACF_0 is κ -categorical, but the incomplete theory ACF is not κ -categorical by the Łoś-Vaught criterion.

braically closed fields) in Theorem 14.2.10.

The proofs of dimension theory and uncountable categoricity involve two technical properties possessed by strongly minimal theories:

1. The algebraic closure operation $\text{acl}(-)$ is a pregeometry.
2. The quantifier \exists^∞ is eliminated (see Section 14.3).

A theory is said to be *geometric* if it has these two properties. There are many geometric theories other than strongly minimal theories, including the complete theory of \mathbb{R} . As we will see, the dimension theory (but not the uncountable categoricity) generalizes from strongly minimal theories to geometric theories.

14.1 Strong minimality

Definition 14.1.1. Let S be a set. A subset $X \subseteq S$ is *cofinite* (in S) if $S \setminus X$ is finite.

Definition 14.1.2. A structure M is *minimal* if M is infinite, and every definable set $X \subseteq M$ is finite or cofinite.

Note this is only a statement about definable sets in one variable; it says nothing about definable subsets of M^n for $n > 1$.

Definition 14.1.3. A theory T is *strongly minimal* if all its models are minimal. A structure M is *strongly minimal* if its complete theory is strongly minimal.

Note that models of a strongly minimal theory are strongly minimal.

Example 14.1.4. ACF is strongly minimal by Lemma 13.5.1.

Theorem 14.1.5. Suppose M is strongly minimal and $A \subseteq M$. Then $A \preceq M$ if and only if $A = \text{acl}(A)$ and A is infinite.

Proof. First suppose $A \preceq M$. Then A is infinite because M is (Lemma 5.5.4), and $A = \text{acl}(A)$ because A is an elementary substructure (Theorem 13.3.4).

Conversely, suppose $A = \text{acl}(A)$ and A is infinite. Then $A \preceq M$ by the Tarski-Vaught criterion (Theorem 5.3.1). Indeed, suppose $D \subseteq M$ is non-empty and A -definable. By strong minimality, D is finite or cofinite. If D is finite then $D \subseteq \text{acl}(A) = A$, so $A \cap D = D \neq \emptyset$. If D is cofinite, it intersects A because A is infinite. \square

Theorem 14.1.6. *If M is strongly minimal, then $\text{acl}(-)$ satisfies the exchange property:*

$$a \in \text{acl}(Cb) \setminus \text{acl}(C) \implies b \in \text{acl}(Ca).$$

Consequently, $\text{acl}(-)$ defines a pregeometry.

Proof. Because $a \in \text{acl}(Cb)$, there is a finite Cb -definable set X containing a . Write X as $\varphi(M, b)$ for some $\mathcal{L}(C)$ -formula $\varphi(x, y)$. Let $n = |X|$. Replacing $\varphi(x, y)$ with

$$\varphi(x, y) \wedge \neg \exists^{\geq n+1} z \varphi(z, y),$$

we may assume that $|\varphi(M, b')| \leq n$ for any $b' \in M$. As $\varphi(a, b)$ holds, $b \in \varphi(a, M)$. If $\varphi(a, M)$ is finite then $b \in \text{acl}(Ca)$ as desired.

Otherwise, $\varphi(a, M)$ is cofinite, and its complement has cardinality $k < \infty$. Let D be the C -definable set of $a' \in M$ such that $|M \setminus \varphi(a', M)| = k$. Then $a \in D$, so D is infinite because $a \notin \text{acl}(C)$. Take distinct $a_1, \dots, a_{n+1} \in D$. Then $\varphi(a_i, M)$ is cofinite for each i . An intersection of cofinite sets is cofinite, hence non-empty, so there is some

$$b' \in \bigcap_{i=1}^{n+1} \varphi(a_i, M).$$

This means that $\varphi(a_i, b')$ holds for all i , and so $|\varphi(M, b')| \geq n + 1$, contradicting the choice of φ . \square

14.2 Uncountable categoricity

Let \mathbb{M} be a monster model of a *complete* strongly minimal theory in a countable language.

Theorem 14.2.1. *For any small set $A \subseteq \mathbb{M}$, there is a unique 1-type $p \in S_1(A)$ whose realizations are the elements of $\mathbb{M} \setminus \text{acl}(A)$.*

Proof. In other words, we must show $b, c \notin \text{acl}(A)$ implies $b \equiv_A c$. Otherwise there is an A -definable set D with $b \in D$ and $c \notin D$ (Remark 8.1.3). If D is finite, then $b \in \text{acl}(A)$ and if D is cofinite, then $c \in \text{acl}(A)$. Either way, we get a contradiction. \square

The pregeometry rank $\text{rk}(\bar{a}/B)$ (Definition 13.6.8) is usually written as $\dim(\bar{a}/B)$. Say that a finite tuple $\bar{a} \in \mathbb{M}^n$ is *independent* if $\dim(\bar{a}) = n$. By definition, this means $a_i \notin \text{acl}(a_1, \dots, a_{i-1})$ for each i . By Lemma 13.6.21, \bar{a} is independent if and only if the a_i are pairwise distinct and $\{a_1, \dots, a_n\}$ is independent as a set.

Lemma 14.2.2. *Let \bar{a}, \bar{b} be two finite tuples of length n in \mathbb{M} . If $\dim(\bar{a}) = n$ and $\dim(\bar{b}) = n$, then $\bar{a} \equiv_{\emptyset} \bar{b}$.*

Proof. Proceed by induction on n . The case $n = 0$ is trivial. Note that the subtuples (a_1, \dots, a_{n-1}) and (b_1, \dots, b_{n-1}) are independent. By induction, $(a_1, \dots, a_{n-1}) \equiv_{\emptyset} (b_1, \dots, b_{n-1})$. Take $\sigma \in \text{Aut}(\mathbb{M})$ moving (a_1, \dots, a_{n-1}) to (b_1, \dots, b_{n-1}) . Replacing \bar{a} with $\sigma(\bar{a})$, we may assume $a_i = b_i$ for $i < n$. By independence,

$$\begin{aligned} a_n &\notin \text{acl}(a_1, \dots, a_{n-1}) \\ b_n &\notin \text{acl}(b_1, \dots, b_{n-1}) = \text{acl}(a_1, \dots, a_{n-1}). \end{aligned}$$

By Theorem 14.2.1, a_n and b_n have the same type over (a_1, \dots, a_{n-1}) . Therefore $(a_1, \dots, a_n) \equiv_{\emptyset} (a_1, \dots, a_{n-1}, b_n) = \bar{b}$. \square

Theorem 14.2.3. *Let M be a model of T . Let $f : I_1 \rightarrow I_2$ be a bijection between two independent sets. Then f is a partial elementary map.*

Proof. If a_1, \dots, a_n are distinct elements of I_1 , then $f(a_1), \dots, f(a_n)$ are distinct elements of I_2 , and the two tuples \bar{a} and $f(\bar{a})$ are both independent n -tuples. By Lemma 14.2.2, they have the same type. \square

Definition 14.2.4. If $M \models T$, the *rank* of M , written $\text{rk}(M)$, is the rank of the pregeometry $(M, \text{acl}(-))$, i.e., the cardinality of a basis.

Remark 14.2.5. If $M \models T$, and B is a basis, then $M = \text{acl}(B)$ (Theorem 13.6.23), and so

$$|M| = |\text{acl}(B)| = |B| + \aleph_0 = \text{rk}(M) + \aleph_0.$$

Theorem 14.2.6. *Two models M_1, M_2 are isomorphic if and only if $\text{rk}(M_1) = \text{rk}(M_2)$.*

Proof. If $M_1 \cong M_2$, it is easy to see that $\text{rk}(M_1) = \text{rk}(M_2)$. Conversely, suppose the ranks are equal. Embed M_1 and M_2 into a monster model \mathbb{M} . Let B_i be a basis of M_i for $i = 1, 2$. Then $|B_1| = |B_2|$. Take a bijection $f : B_1 \rightarrow B_2$. By Theorem 14.2.3, f is a partial elementary map from \mathbb{M} to \mathbb{M} . By strong homogeneity, f extends to an automorphism $\sigma \in \text{Aut}(\mathbb{M})$. Then $\sigma(M_1) = \sigma(\text{acl}(B_1)) = \text{acl}(B_2) = M_2$. The automorphism σ restricts to an isomorphism from M_1 to M_2 . \square

Corollary 14.2.7. *T is κ -categorical for all $\kappa > \aleph_0$.*

Theorem 14.2.8. *Suppose that in models of T , algebraically closed sets are infinite. Then there is a unique model of rank κ for every cardinal κ , finite or infinite.*

Proof. Take a model M_0 of size greater than $\kappa + \aleph_0$, and let B_0 be a basis of M_0 . Then $|B_0| > \kappa$. Take a subset $I \subseteq B_0$ with $|I| = \kappa$, and let $M = \text{acl}(I)$. By assumption, M is infinite. Then $M \preceq M_0$ by Theorem 14.1.5. By Remark 13.6.24, the independent set I is a basis of M , so $\text{rk}(M) = |I| = \kappa$. \square

The classification of algebraically closed fields

Definition 14.2.9. The *transcendence degree* of an algebraically closed field K is its rank in the sense of Definition 14.2.4.

Theorem 14.2.10. *For each $p \in \{0, 2, 3, 5, 7, \dots\}$ and each cardinal κ , there is a unique algebraically closed field of characteristic p and transcendence degree κ .*

Proof. The theory ACF_p is complete and strongly minimal. Algebraically closed sets are infinite by Theorem 13.5.3(2) and Theorem 9.2.3. Therefore Theorem 14.2.8 applies, giving a unique model $M \models \text{ACF}_p$ with transcendence degree κ . \square

14.3 Uniform finiteness

The notation $\exists^\infty x P(x)$ means that there are infinitely many x such that $P(x)$ holds. In general, $\exists^\infty x$ cannot be expressed in first-order logic.

Definition 14.3.1. A structure M *eliminates* \exists^∞ if for any first-order formula $\varphi(x, \bar{y})$, there is a formula $\psi(\bar{y})$ such that for any \bar{b} ,

$$M \models \exists^\infty x \varphi(x, \bar{b}) \iff M \models \psi(\bar{b}).$$

Here, the left-hand side really means $\exists^\infty a \in M : M \models \varphi(a, \bar{b})$, or equivalently, $\varphi(M, \bar{b})$ is infinite.

Recall from Section 3.3 that the notation $\exists^{\geq n} x P(x)$ means that there are at least n values of x such that $P(x)$ holds. Unlike \exists^∞ , this can be expressed in first-order logic.

Definition 14.3.2. A structure M has *uniform finiteness* if for any formula $\varphi(x, \bar{y})$, there is a number n_φ such that for any \bar{b} in M ,

$$|\varphi(M, \bar{b})| < \infty \iff |\varphi(M, \bar{b})| < n_\varphi.$$

Theorem 14.3.3. *Let M be a structure.*

1. *If M has uniform finiteness, then M eliminates \exists^∞ .*
2. *If M has uniform finiteness and $N \equiv M$, then N has uniform finiteness.*
3. *M has uniform finiteness if and only if every $N \equiv M$ eliminates \exists^∞ .*

Proof. 1. Uniform finiteness says $(\exists^\infty x)\varphi(x, \bar{y}) \iff (\exists^{\geq n_\varphi} x)\varphi(x, \bar{y})$.

2. Given φ , if n_φ works for M then it works for N . Otherwise there is \bar{b} in N such that $n_\varphi \leq |\varphi(N, \bar{b})| < \infty$. Let $k = |\varphi(N, \bar{b})|$. Then

$$N \models \exists \bar{y} \exists^{\geq k} x \varphi(x, \bar{y}).$$

This says that there is a \bar{b} in N such that $\varphi(N, \bar{b})$ has size exactly k . As $M \equiv N$, the same holds in M , so there is a \bar{b} in M such that $\varphi(M, \bar{b})$ has size exactly $k \geq n_\varphi$, contradicting the choice of n_φ .

3. If M has uniform finiteness and $N \equiv M$, then N has uniform finiteness and therefore eliminates \exists^∞ . Conversely, suppose every $N \equiv M$ eliminates \exists^∞ . Fix an \aleph_1 -saturated elementary extension $N \succeq M$. By

part (2), it suffices to show that N has uniform finiteness. Fix $\varphi(x, \bar{y})$. Let

$$D_\infty = \{\bar{b} : |\varphi(N, \bar{b})| < \infty\}$$

$$D_k = \{\bar{b} : |\varphi(N, \bar{b})| < k\} \text{ for } k < \omega.$$

Then D_∞ is definable by elimination of \exists^∞ , and D_k is definable easily. Moreover,

$$D_0 \subseteq D_1 \subseteq \cdots \subseteq D_\infty$$

and $D_\infty = \bigcup_{i=0}^\infty D_i$. By \aleph_0 -compactness (Theorem 11.2.7(3)), $D_\infty = D_{n_\varphi}$ for some n_φ , which means

$$|\varphi(N, \bar{b})| < \infty \implies |\varphi(N, \bar{b})| < n_\varphi. \quad \square$$

Theorem 14.3.4. *If M is strongly minimal, then M has uniform finiteness.*

Proof. Without loss of generality, M is \aleph_1 -saturated. Let $\varphi(x; y_1, \dots, y_n)$ be a formula. Let D_k be the set of \bar{b} such that $\varphi(M, \bar{b})$ or $M \setminus \varphi(M, \bar{b})$ has size less than k . Then $D_0 \subseteq D_1 \subseteq \cdots \subseteq M^n$. Strong minimality means that $M^n = \bigcup_{k=0}^\infty D_k$. By \aleph_0 -compactness (Theorem 11.2.7(3)), $D_k = M^n$ for some k . This means that for any \bar{b} ,

$$|\varphi(M, \bar{b})| < k \text{ or } |M \setminus \varphi(M, \bar{b})| < k.$$

Therefore,

$$|\varphi(M, \bar{b})| < k \text{ or } |\varphi(M, \bar{b})| = \infty,$$

which is uniform finiteness. \square

14.4 Pregeometric theories

Definition 14.4.1. A theory T is *pregeometric* if $\text{acl}(-)$ satisfies exchange in models of T .

Definition 14.4.2. A complete theory T is *geometric* if it is pregeometric and has uniform finiteness.

For example, strongly minimal theories are geometric.

Let \mathbb{M} be a monster model of a pregeometric theory. Let $\dim(\bar{a}/B)$ denote the rank of \bar{a} over B with respect to $\text{acl}(-)$, i.e., the number of values $i \in \{1, \dots, n\}$ such that $a_i \notin \text{acl}(Ba_1a_2 \cdots a_{i-1})$.

Theorem 14.4.3. 1. $\dim(\bar{a}/B) = \dim(\sigma(\bar{a})/\sigma(B))$ for any $\sigma \in \text{Aut}(\mathbb{M})$.

2. If $\bar{a} \equiv_C \bar{b}$, then $\dim(\bar{a}/C) = \dim(\bar{b}/C)$.

Proof. 1. The definition of $\dim(-/-)$ is clearly automorphism-invariant.

2. Take $\sigma \in \text{Aut}(\mathbb{M}/C)$ with $\sigma(\bar{a}) = \bar{b}$ and apply part (1). \square

In particular, $\text{tp}(\bar{a}/C)$ determines $\dim(\bar{a}/C)$. If $p \in S_n(C)$, we let $\dim(p)$ denote $\dim(\bar{a}/C)$ for any \bar{a} realizing p .

Theorem 14.4.4 (Extension). *Let $B \subseteq C$ be small sets in \mathbb{M} .*

1. If $p \in S_n(B)$, then there is an extension $q \in S_n(C)$ with $q \supseteq p$.

2. If $\bar{a} \in \mathbb{M}^n$, then there is $\bar{a}' \equiv_B \bar{a}$ with $\dim(\bar{a}'/C) = \dim(\bar{a}/B)$.

Proof. Note that (1) \iff (2), by taking $p = \text{tp}(\bar{a}/B)$ and $q = \text{tp}(\bar{a}'/B)$. We prove (2) by induction on n . First suppose $n = 1$. If $a \notin \text{acl}(B)$, then the set X of realizations of $\text{tp}(a/B)$ is large by Theorem 13.3.5, so there is $a' \in X \setminus \text{acl}(C)$. Then $a' \equiv_B a$ and $\dim(a'/C) = 1 = \dim(a/B)$. On the other hand, if $a \in \text{acl}(B) \subseteq \text{acl}(C)$, then $\dim(a/C) = 0 = \dim(a/B)$, and we can take $a' = a$.

Next suppose $n > 1$. Write \bar{a} as (\bar{a}_1, \bar{a}_2) , where \bar{a}_1 and \bar{a}_2 are tuples of shorter length. By induction there is $\bar{a}'_1 \equiv_B \bar{a}_1$ with $\dim(\bar{a}'_1/C) = \dim(\bar{a}_1/B)$. Moving \bar{a} by an automorphism over B sending \bar{a}_1 to \bar{a}'_1 , we may assume $\bar{a}_1 = \bar{a}'_1$. Then

$$\dim(\bar{a}_1/C) = \dim(\bar{a}_1/B). \quad (*)$$

By induction applied to \bar{a}_2 and the inclusion $B\bar{a}_1 \subseteq C\bar{a}_1$, there is $\bar{a}'_2 \equiv_{B\bar{a}_1} \bar{a}_2$ with $\dim(\bar{a}'_2/C\bar{a}_1) = \dim(\bar{a}_2/B\bar{a}_1)$. Moving \bar{a} by an automorphism over $B\bar{a}_1$ sending \bar{a}_2 to \bar{a}'_2 , we may assume $\bar{a}'_2 = \bar{a}_2$. Then

$$\dim(\bar{a}_2/C\bar{a}_1) = \dim(\bar{a}_2/B\bar{a}_1). \quad (\dagger)$$

Adding equations $(*)$ and (\dagger) and using additivity,

$$\dim(\bar{a}_1\bar{a}_2/C) = \dim(\bar{a}_1\bar{a}_2/B). \quad \square$$

14.5 Dimension theory in the monster

Work in a monster model \mathbb{M} of a pregeometric theory.

Definition 14.5.1. If B is small and $X \subseteq \mathbb{M}^n$ is B -definable, then $\dim_B(X) = \max_{\bar{a} \in X} \dim(\bar{a}/B)$. If X is empty, we set $\dim(X) = -\infty$.

Theorem 14.5.2. If X is B -definable and C -definable, then $\dim_B(X) = \dim_C(X)$.

Proof. First suppose $B \subseteq C$. Note that $\dim(\bar{a}/B) \geq \dim(\bar{a}/C)$ for any \bar{b} (Theorem 13.6.15), so

$$\dim_B(X) = \max_{\bar{a} \in X} \dim(\bar{a}/B) \geq \max_{\bar{a} \in X} \dim(\bar{a}/C) = \dim_C(X).$$

Conversely, take $\bar{a} \in X$ with $\dim(\bar{a}/B) = \dim_B(X)$. By Theorem 14.4.4, there is $\bar{a}' \equiv_B \bar{a}$ with $\dim(\bar{a}'/C) = \dim(\bar{a}/B)$. Then $\bar{a} \in X \implies \bar{a}' \in X$ because X is B -definable (Remark 8.1.3), and so

$$\dim_C(X) \geq \dim(\bar{a}'/C) = \dim(\bar{a}/B) = \dim_B(X).$$

This completes the case where $B \subseteq C$. The general case then follows:

$$\dim_B(X) = \dim_{B \cup C}(X) = \dim_C(X). \quad \square$$

By Theorem 14.5.2, $\dim_B(X)$ doesn't depend on B , so we just write $\dim(X)$. The following two facts are a restatement of the definition:

- If X is B -definable and $\bar{a} \in X$, then $\dim(\bar{a}/B) \leq \dim(X)$.
- If X is B -definable and non-empty, then there is $\bar{a} \in X$ with $\dim(\bar{a}/B) = \dim(X)$.

We will use these repeatedly in what follows.

Theorem 14.5.3 (Basic properties of dimension). *Let X, Y be definable sets.*

1. $\dim(X) \leq 0$ if and only if $|X| < \infty$.
2. If $X \subseteq Y$, then $\dim(X) \leq \dim(Y)$.
3. If $X, Y \subseteq \mathbb{M}^n$, then $\dim(X \cup Y) = \max(\dim(X), \dim(Y))$.

4. $\dim(X \times Y) = \dim(X) + \dim(Y)$.
5. $\dim(\mathbb{M}^n) = n$, assuming \mathbb{M} is infinite.
6. If $f : X \rightarrow Y$ is a definable surjection, then $\dim(X) \geq \dim(Y)$.
7. If $f : X \rightarrow Y$ is a definable bijection, then $\dim(X) = \dim(Y)$.
8. If $f : X \rightarrow Y$ is a definable injection, then $\dim(X) \leq \dim(Y)$.

Proof. Take a small set C over which all the sets and functions are defined.

1. If X is finite, then every $\bar{a} \in X$ is algebraic over C , and so $\dim(\bar{a}/C) \leq 0$ (Theorem 13.6.10). Thus $\dim(X) \leq 0$.

Conversely, if X is infinite, then X is large (Corollary 11.2.8), so there is $\bar{a} \in X$ with $\bar{a} \notin \text{acl}(C)$. Then $\dim(X) \geq \dim(\bar{a}/C) > 0$.

2. Clear.
3. Clear.
4. If $(\bar{a}, \bar{b}) \in X \times Y$, then

$$\dim(\bar{a}\bar{b}/C) = \dim(\bar{a}/C) + \dim(\bar{b}/C\bar{a}) \leq \dim(X) + \dim(Y)$$

because \bar{a} is in the C -definable set X and \bar{b} is in the $C\bar{a}$ -definable set Y . As this holds for all $(\bar{a}, \bar{b}) \in X \times Y$, we have $\dim(X \times Y) \leq \dim(X) + \dim(Y)$.

Conversely, take $\bar{a} \in X$ with $\dim(\bar{a}/C) = \dim(X)$. The set Y is $C\bar{a}$ -definable, so there is $\bar{b} \in Y$ with $\dim(\bar{b}/C\bar{a}) = \dim(Y)$. Then

$$\dim(X \times Y) \geq \dim(\bar{a}\bar{b}/C) = \dim(\bar{a}/C) + \dim(\bar{b}/C\bar{a}) = \dim(X) + \dim(Y).$$

5. By part (4), it suffices to show $\dim(\mathbb{M}^1) = 1$. If $a \in \mathbb{M}^1$, then $\dim(a/C) \leq 1$, as a has length 1. Thus $\dim(\mathbb{M}^1) \leq 1$. On the other hand, $\dim(\mathbb{M}) \geq 1$ by part (1).
6. Take $\bar{b} \in Y$ with $\dim(\bar{b}/C) = \dim(Y)$. Since f is surjective, there is $\bar{a} \in X$ with $f(\bar{a}) = \bar{b}$. Then $\bar{b} \in \text{dcl}(C\bar{a}) \subseteq \text{acl}(C\bar{a})$, so by Theorem 13.6.14,

$$\dim(X) \geq \dim(\bar{a}/C) \geq \dim(\bar{b}/C) = \dim(Y).$$

7. Apply part (6) to f and f^{-1} .

8. By parts (2) and (7), $\dim(X) = \dim(\text{im}(f)) \leq \dim(Y)$. \square

Theorem 14.5.4 (Fiber dimension theorem). *Let $f : X \rightarrow Y$ be a definable function. For every $b \in Y$, let $X_b = f^{-1}(b) = \{x \in X : f(x) = b\}$. If $\dim(X_b) = k$ for all $b \in Y$, then $\dim(X) = k + \dim(Y)$.*

Proof. Take a small set C defining f , X , and Y . Note that \bar{a} and $(\bar{a}, f(\bar{a}))$ are interalgebraic over C for any $\bar{a} \in X$, so

$$\dim(\bar{a}/C) = \dim(\bar{a}, f(\bar{a})/C) = \dim(\bar{a}/C f(\bar{a})) + \dim(f(\bar{a})/C)$$

by Theorem 13.6.14 and additivity (Theorem 13.6.9). If $\bar{b} = f(\bar{a})$, then

$$\dim(\bar{a}/C) = \dim(\bar{a}/C\bar{b}) + \dim(\bar{b}/C) \leq \dim(X_b) + \dim(Y) = k + \dim(Y).$$

because \bar{a} is in the $C\bar{b}$ -definable set X_b , and \bar{b} is in the C -definable set Y . As this holds for any $\bar{a} \in X$, we see

$$\dim(X) \leq k + \dim(Y).$$

For the converse, take $\bar{b} \in Y$ with $\dim(\bar{b}/C) = \dim(Y)$. The set X_b is $C\bar{b}$ -definable, so there is $\bar{a} \in X_b$ with $\dim(\bar{a}/C\bar{b}) = \dim(X_b) = k$. Then $\bar{b} = f(\bar{a})$, so

$$\dim(X) \geq \dim(\bar{a}/C) = \dim(\bar{a}/C\bar{b}) + \dim(\bar{b}/C) = k + \dim(Y). \quad \square$$

Proof. Let $Z_k = f^{-1}(Y_k)$. Then $\dim(Z_k) = k + \dim(Y_k)$, and $X = \bigcup_k Z_k$. \square

Now suppose the theory is geometric (Definition 14.4.2), meaning that uniform finiteness holds.

Lemma 14.5.5. *Let $\varphi(x_1, \dots, x_n)$ be an $\mathcal{L}(\mathbb{M})$ -formula and let $\psi(x_1, \dots, x_{n-1})$ be an $\mathcal{L}(\mathbb{M})$ -formula equivalent to $\exists^\infty x_n \varphi$. Then $\varphi(\mathbb{M}^n)$ has dimension n if and only if $\psi(\mathbb{M}^{n-1})$ has dimension $n - 1$.*

Proof. Take a finite set B such that φ and ψ are $\mathcal{L}(B)$ -formulas. If $\psi(\mathbb{M}^{n-1})$ has dimension $n - 1$, take $\bar{a} \in \psi(\mathbb{M}^{n-1})$ with $\dim(\bar{a}/B) = n - 1$. Then $\varphi(\bar{a}, \mathbb{M})$ is infinite, hence large. Take $a_n \in \varphi(\bar{a}, \mathbb{M}) \setminus \text{acl}(B\bar{a})$. Then $(\bar{a}, a_n) \in \varphi(\mathbb{M}^n)$ and

$$\dim(\bar{a}, a_n/B) = \dim(\bar{a}/B) + \dim(a_n/B\bar{a}) = (n - 1) + 1 = n,$$

and $\varphi(\mathbb{M}^n)$ has dimension n .

Conversely, suppose $\varphi(\mathbb{M}^n)$ has dimension n . Take $(\bar{a}, a_n) \in \varphi(\mathbb{M}^n)$ with $\dim(\bar{a}, a_n/B) = n$. Then $a_n \notin \text{acl}(B\bar{a})$ and $\dim(\bar{a}/B) = n - 1$. As $\varphi(\bar{a}, \mathbb{M})$ is $B\bar{a}$ -definable and contains $a_n \notin \text{acl}(B\bar{a})$, it must be infinite, meaning that $\bar{a} \in \psi(\mathbb{M}^{n-1})$, and then $\dim(\psi(\mathbb{M}^{n-1})) \geq \dim(\bar{a}/B) = n - 1$. \square

Lemma 14.5.6. *Let $\varphi(x_1, \dots, x_n; y_1, \dots, y_m)$ be an \mathcal{L} -formula. The set*

$$\{\bar{b} \in \mathbb{M}^m : \dim(\varphi(\mathbb{M}^n, \bar{b})) = n\}$$

is definable.

Proof. The case $n = 0$ is easy. If $n > 0$, let $\psi(x_1, \dots, x_{n-1}; \bar{y})$ be the \mathcal{L} -formula equivalent to $\exists^\infty x_n \varphi$. Then for any $\bar{b} \in \mathbb{M}^m$,

$$\dim(\varphi(\mathbb{M}^n, \bar{b})) = n \iff \dim(\psi(\mathbb{M}^{n-1}, \bar{b})) = n - 1,$$

by Lemma 14.5.5, and the right hand side is definable by induction on n . \square

Lemma 14.5.7. *Let $D \subseteq \mathbb{M}^n$ be definable and let k be in $\{0, 1, \dots, n\}$. Then $\dim(D) \geq k$ if and only if there is a coordinate projection $\pi : \mathbb{M}^n \rightarrow \mathbb{M}^k$ such that $\pi(D) \subseteq \mathbb{M}^k$ has dimension k .*

Proof. If $\dim(\pi(D)) = k$, then $\dim(D) \geq k$ by Theorem 14.5.3(6). Conversely, suppose $\dim(D) \geq k$. Take a small set B over which D is defined, and take $\bar{a} \in D$ with $\dim(\bar{a}/B) = \dim(D) \geq k$. There are at least k values of i such that $a_i \notin \text{acl}(Ba_1, \dots, a_{i-1})$. Let $i_1 < i_2 < \dots < i_k$ be some such values. Then

$$a_{i_j} \notin \text{acl}(Ba_{i_1}a_{i_2} \cdots a_{i_{j-1}})$$

for each j , and so

$$\dim(a_{i_1}a_{i_2} \cdots a_{i_k}/B) = k.$$

Let $\pi : \mathbb{M}^n \rightarrow \mathbb{M}^k$ be the coordinate projection $\pi(x_1, \dots, x_n) = (x_{i_1}, \dots, x_{i_k})$. Then $\pi(D)$ is B -definable, $\pi(\bar{a}) \in \pi(D)$, and $\dim(\pi(\bar{a})/B) = k$. We conclude that $\dim(\pi(D)) \geq k$. \square

Theorem 14.5.8 (Definability of dimension). *Let $\varphi(x_1, \dots, x_n; y_1, \dots, y_m)$ be a formula. For each $k \leq n$, the set*

$$\{\bar{b} \in \mathbb{M}^m : \dim(\varphi(\mathbb{M}^n, \bar{b})) = k\}$$

is definable.

Proof. Let Π_k^n be the finite set of coordinate projections $\mathbb{M}^n \rightarrow \mathbb{M}^k$. For each $\pi \in \Pi_k^n$, let $\theta_\pi(z_1, \dots, z_k, \bar{y})$ be the formula

$$\exists \bar{x} \varphi(\bar{x}; \bar{y}) \wedge (\pi(\bar{x}) = \bar{z}).$$

Then $\theta_\pi(\mathbb{M}^k, \bar{b})$ is the image of $\varphi(\mathbb{M}^n, \bar{b})$ under π . By Lemma 14.5.7,

$$D_k := \{\bar{b} \in \mathbb{M}^m : \dim(\varphi(\mathbb{M}^n; \bar{b})) \geq k\} = \bigcup_{\pi \in \Pi_k^n} \{\bar{b} \in \mathbb{M}^m : \dim(\theta_\pi(\mathbb{M}^k; \bar{b})) = k\},$$

and the sets on the right-hand side are definable by Lemma 14.5.6. Thus each D_k is definable. The set we want is $D_k \setminus D_{k-1}$. \square

Theorem 14.5.9. *Let $f : X \rightarrow Y$ be a definable function. For each $b \in Y$, let $X_b = f^{-1}(b)$. For each $k \leq \dim(X)$, let $Y_k = \{b \in Y : \dim(X_b) = k\}$. Then each set Y_k is definable, and*

$$\dim(X) = \max_k (k + \dim(Y_k)).$$

Proof. Theorems 14.5.8 and 14.5.4. \square

14.6 Dimension theory in small models

If D is a definable set in a model M , and N is an elementary extension of M , then $D(N)$ denotes $\varphi(N)$, where $\varphi(\bar{x})$ is the $\mathcal{L}(M)$ -formula defining D in M . The choice of φ doesn't matter, because

$$\begin{aligned} \varphi(M) = \psi(M) &\iff M \models \forall \bar{x} (\varphi(\bar{x}) \leftrightarrow \psi(\bar{x})) \\ &\iff N \models \forall \bar{x} (\varphi(\bar{x}) \leftrightarrow \psi(\bar{x})) \iff \varphi(N) = \psi(N). \end{aligned}$$

Note that $D(M) = D$.

Lemma 14.6.1. *Let $M \preceq N$ be monster models of a pregeometric theory.*

1. *If $\bar{a} \in M^n$ and $B \subseteq M$ is small, then $\dim^M(\bar{a}/B) = \dim^N(\bar{a}/B)$.*
2. *If D is M -definable, then $\dim(D(M)) = \dim(D(N))$.*

Proof. 1. By Theorem 13.3.3, $\text{acl}(Ba_1, \dots, a_{i-1})$ is the same whether calculated in M or N .

2. Take a small set $B \subseteq M$ defining D . Take $\bar{a} \in D(M)$ with $\dim(\bar{a}/B) = \dim(D(M))$. Then $\bar{a} \in D(N)$ and $\dim(\bar{a}/B) = \dim(D(M))$, so

$$\dim(D(M)) \leq \dim(D(N)).$$

Conversely, take $\bar{a} \in D(N)$ with $\dim(\bar{a}/B) = \dim(D(N))$. Then $\text{tp}(\bar{a}/B)$ is realized in M by some \bar{c} . The formula defining $D(N)$ is in $\text{tp}(\bar{a}/B)$, so $\bar{c} \in D(N) \cap M^n = D(M)$. Also, $\dim(\bar{c}/B) = \dim(\bar{a}/B)$ by automorphism invariance. The fact that $\bar{c} \in D(M)$ then shows

$$\dim(D(M)) \geq \dim(\bar{c}/B) = \dim(\bar{a}/B) = \dim(D(N)). \quad \square$$

Definition 14.6.2. Let M be a small model of a pregeometric theory. If $D \subseteq M^n$ is definable, then $\dim(D)$ is defined to be $\dim(D(\mathbb{M}))$ where \mathbb{M} is a monster model extending M .

Theorem 14.6.3. *The choice of \mathbb{M} doesn't matter in Definition 14.6.2.*

Proof. Let $\mathbb{M}_1, \mathbb{M}_2$ be two monster models extending M . Then \mathbb{M}_1 and \mathbb{M}_2 are elementarily equivalent as $\mathcal{L}(M)$ -structures. By elementary amalgamation there is a third monster model \mathbb{M}_3 extending \mathbb{M}_1 and \mathbb{M}_2 , up to isomorphism. Then

$$\dim(D(\mathbb{M}_1)) = \dim(D(\mathbb{M}_3)) = \dim(D(\mathbb{M}_2))$$

by two applications of Lemma 14.6.1. \square

Corollary 14.6.4. *Dimension is invariant in elementary extensions: if $M \preceq N$ and D is a definable set in M , then $\dim(D(M)) = \dim(D(N))$.*

Proof. Take a monster model \mathbb{M} extending N (and M). Then $\dim(D(M)) = \dim(D(\mathbb{M})) = \dim(D(N))$ by two applications of Definition 14.6.2. \square

Theorem 14.6.5. *Let M be a model of a pregeometric theory T . Define $\dim(-)$ as in Definition 14.6.2.*

1. *The basic properties of dimension (Theorem 14.5.3) hold in M .*
2. *If T is geometric, then the fiber dimension theorem (Theorem 14.5.4) and definability of dimension (Theorem 14.5.8) hold in M .*

Proof. 1. For example, we show $\dim(D_1 \times D_2) = \dim(D_1) + \dim(D_2)$. If $\varphi(\bar{x})$ defines D_1 and $\psi(\bar{y})$ defines D_2 , then the formula $\theta(\bar{x}, \bar{y}) := \varphi(\bar{x}) \wedge \psi(\bar{y})$ defines $D_1 \times D_2$. Therefore

$$(D_1 \times D_2)(\mathbb{M}) = \theta(\mathbb{M}) = D_1(\mathbb{M}) \times D_2(\mathbb{M}).$$

Thus

$$\begin{aligned} \dim(D_1 \times D_2) &= \dim((D_1 \times D_2)(\mathbb{M})) \\ &= \dim(D_1(\mathbb{M}) \times D_2(\mathbb{M})) = \dim(D_1(\mathbb{M})) + \dim(D_2(\mathbb{M})) \\ &= \dim(D_1) + \dim(D_2). \end{aligned}$$

2. Suppose we have the configuration $f : X \rightarrow Y$ of Theorem 14.5.4, where $X_b = f^{-1}(b)$ has dimension k for every $b \in Y$. In order to reduce from M to the known case of \mathbb{M} , we need to show that the property

$$\dim(X_b) = k \text{ for every } b \in Y$$

transfers from M to \mathbb{M} . Let $f_{\mathbb{M}} : X(\mathbb{M}) \rightarrow Y(\mathbb{M})$ be the function defined by the same formula as f , and for $b \in Y(\mathbb{M})$ let $X_b(\mathbb{M})$ denote the fiber of $f_{\mathbb{M}}^{-1}(b)$. Note that when $b \in Y = Y(M)$, the set $X_b(\mathbb{M})$ really is the extension of X_b . By the definability of dimension (Theorem 14.5.8), the set

$$Z = \{b \in Y(\mathbb{M}) : \dim(X_b(\mathbb{M})) \neq k\}$$

is definable. It is $\text{Aut}(\mathbb{M}/M)$ -invariant, hence M -definable (Theorem 11.3.8). If $b \in Y(M)$, then $\dim(X_b(\mathbb{M})) = \dim(X_b) = k$, so $b \notin Z$. Therefore there are no M -points in Z . By the Tarski-Vaught criterion, $Z = \emptyset$. Thus $Z = \emptyset$, and $\dim(X_b(\mathbb{M})) = k$ for every $b \in Y(\mathbb{M})$. Then we can apply Theorem 14.5.4 over \mathbb{M} to get what we want.

For Theorem 14.5.8, fix a formula $\varphi(\bar{x}, \bar{y})$. The set

$$Z_k = \{\bar{b} \in \mathbb{M} : \dim(\varphi(\mathbb{M}; \bar{b})) = k\}$$

is definable by Theorem 14.5.8 applied to \mathbb{M} , and clearly $\text{Aut}(\mathbb{M}/\emptyset)$ -invariant. By Theorem 11.3.8, it is 0-definable, defined by some formula $\psi_k(\bar{y})$. Then for any \bar{b} in M ,

$$\begin{aligned} \dim(\varphi(M, \bar{b})) = k &\iff \dim(\varphi(\mathbb{M}, \bar{b})) = k \\ &\iff \mathbb{M} \models \psi_k(\bar{b}) \iff M \models \psi_k(\bar{b}). \end{aligned}$$

The first equivalence holds by Definition 14.6.2. The second holds by choice of ψ_k . The third holds as $M \preceq \mathbb{M}$.

Therefore $\{\bar{b} \in M^m : \dim(\varphi(M^n; \bar{b})) = k\}$ is the definable set $\psi_k(M^m)$. \square

Appendix A

Mathematical background

This appendix reviews the mathematical background assumed throughout the book.

A.1 Logical notation

We write $a := b$ or $b =: a$ to mean that a is defined to be b , i.e., “let $a = b$.”

If φ, ψ are statements, then $\varphi \implies \psi$ means that φ logically implies ψ . In other words,

$$\varphi \implies \psi$$

means

If φ then ψ .

The notation $\varphi \iff \psi$ means that $\varphi \implies \psi$ and $\psi \implies \varphi$. We say that φ holds “if and only if” ψ holds. “If and only if” is often abbreviated to iff. The notation $\forall x (\dots)$ means “for every x , ...” Similarly, the notation $\exists x (\dots)$ means “there is an x such that ...” If we are only interested in values of x coming from a set A , we write $\forall x \in A$ or $\exists x \in A$. In other words, these things

$$\forall x \in A (\dots)$$

$$\exists x \in A (\dots)$$

mean the following, respectively:

For every x in A , ...

There is an x in A such that

A.2 Sets

If A is a set and x is a value, then $x \in A$ means that x is an element of A , and $x \notin A$ means that x is not an element of A . If A and B are sets, then $A \subseteq B$ means that A is a subset of B , i.e., every element of A is an element of B . It is an axiom of set theory that $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$. The notation $A \subsetneq B$ means that A is a proper subset of B : $A \subseteq B$ and $A \neq B$. We avoid the ambiguous notation $A \subset B$. Remember to distinguish $A \subsetneq B$ (A is a proper subset of B) from $A \not\subseteq B$ (A is not a subset of B). We write $A \subseteq_f B$ to mean that A is a finite subset of B . (This relation comes up frequently in model theory.)

We write the empty set as \emptyset ; this is the unique set with no elements. If A is a set, then $\mathfrak{P}(A)$ denotes the powerset of A , the set of all subsets of A :

$$\mathfrak{P}(A) = \{X : X \subseteq A\}.$$

Here, the set-builder notation $\{x : \dots\}$ means “the set of x such that \dots .” Similarly, $\{x \in A : \dots\}$ means “the set of x in A such that \dots ” and $\{f(x) : x \in A, \dots\}$ means “the set of values $f(x)$ where x is in A and \dots ”.

If A and B are sets, then $A \cap B$ and $A \cup B$ denote the intersection and union of A and B :

$$\begin{aligned} A \cap B &= \{x : x \in A \text{ and } x \in B\} \\ A \cup B &= \{x : x \in A \text{ or } x \in B\}. \end{aligned}$$

If A_1, \dots, A_n are sets, then we write their intersection and union as follows:

$$\begin{aligned} \bigcap_{i=1}^n A_i &= A_1 \cap A_2 \cap A_3 \cap \dots \cap A_n \\ \bigcup_{i=1}^n A_i &= A_1 \cup A_2 \cup A_3 \cup \dots \cup A_n. \end{aligned}$$

Additionally, we write the “set difference” of A and B as $A \setminus B$:

$$A \setminus B = \{x : x \in A \text{ and } x \notin B\}.$$

This should be distinguished from the “symmetric difference” $(A \setminus B) \cup (B \setminus A)$, which we will not use.

Two sets A and B are *disjoint* if $A \cap B = \emptyset$. A list of sets A_1, \dots, A_n is *pairwise disjoint* if any two of them are disjoint: $A_i \cap A_j = \emptyset$ for any $i \neq j$. A list of values x_1, \dots, x_n is *pairwise distinct* if $x_i \neq x_j$ for $i \neq j$.

A.3 $\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$

We use the following blackboard bold symbols. \mathbb{N} denotes the natural numbers $\{0, 1, 2, \dots\}$. Note that we include 0 in \mathbb{N} . We sometimes denote \mathbb{N} as ω . Notation like $n < \omega$ and $n \in \omega$ means that n is a natural number. The reason for this notation comes from set theory, where ω is another name for \mathbb{N} and ω is the first ordinal number after the natural numbers. \mathbb{Z} denotes the integers $\{\dots, -2, -1, 0, 1, 2, \dots\}$. \mathbb{Q} denotes the set of rational numbers, values of the form x/y where $x, y \in \mathbb{Z}$. \mathbb{R} denotes the usual real numbers. \mathbb{C} denotes the complex numbers, values of the form $x + iy$ where $x, y \in \mathbb{R}$ and $i = \sqrt{-1}$. To summarize,

$$\mathbb{N} \subsetneq \mathbb{Z} \subsetneq \mathbb{Q} \subsetneq \mathbb{R} \subsetneq \mathbb{C}.$$

A.4 Tuples, products, and relations

An n -tuple is a list of n values $(x_1, x_2, x_3, \dots, x_n)$. 2-tuples are called *pairs* or *ordered pairs*, and 3-tuples are called *triples*. Two n -tuples

$$\begin{aligned} &(x_1, \dots, x_n) \\ &(y_1, \dots, y_n) \end{aligned}$$

are equal if and only if $x_i = y_i$ for all i between 1 and n (inclusive). For example, $(x_1, x_2) = (y_1, y_2)$ if and only if $x_1 = y_1$ and $x_2 = y_2$.

If A and B are sets, the *direct product* or *cartesian product*, written $A \times B$, is the set of ordered pairs (x, y) where $x \in A$ and $y \in B$:

$$A \times B = \{(x, y) : x \in A \text{ and } y \in B\}.$$

For example,

$$\begin{aligned} \{1, 2, 3\} \times \{\text{up}, \text{down}\} = \{ &(1, \text{up}), (1, \text{down}), (2, \text{up}), \\ &(2, \text{down}), (3, \text{up}), (3, \text{down})\}. \end{aligned}$$

A *relation* between A and B is a subset $R \subseteq A \times B$. If $x \in A$ and $y \in B$, then $x R y$ means $(x, y) \in R$.

A *relation on a set* A is a relation between A and A , i.e., a subset $R \subseteq A^2$. A relation R on A is said to be...

- *reflexive* if $x R x$ for any $x \in A$.
- *symmetric* if $x R y \implies y R x$
- *transitive* if $x R y$ and $y R z$ imply $x R z$.
- *antisymmetric* if $x R y$ and $y R x$ imply $x = y$.

For example, the relation $=$ is reflexive, symmetric, transitive, and antisymmetric. The relation $<$ is transitive, and antisymmetric, but not reflexive or symmetric. The relation \leq is reflexive, transitive, and antisymmetric, but not symmetric.

An *equivalence relation* on A is a relation that is reflexive, symmetric, and transitive. Fix an equivalence relation \sim . If $x \in A$, the *equivalence class* of x is the set of $y \in A$ equivalent to x :

$$[x] = \{y \in A : x \sim y\}.$$

The *quotient set* is the set of equivalence classes:

$$A/\sim = \{[x] : x \in A\}.$$

Using the definition of equivalence relation, one can show that $[x] = [y] \iff x \sim y$. Moreover, the collection of equivalence classes is a *partition* of A —a collection \mathcal{C} of non-empty subsets of A such that each $x \in A$ belongs to exactly one set in \mathcal{C} . In fact, there is a one-to-one correspondence between equivalence relations on A and partitions of A .

A *partial order* on A is a relation \leq that is reflexive, transitive, and antisymmetric. The prototypical examples of partial orders are:

- The relation \leq on \mathbb{R} .
- The relation \subseteq on the power set $\mathfrak{P}(X)$.

If \leq is a partial order, then $x < y$ means $x \leq y$ and $x \neq y$. The relation $<$ is always transitive and irreflexive ($x \not< x$ for any x). A *strict partial order* is a relation $<$ that is transitive and irreflexive. In fact, there is a one-to-one correspondence between partial orders on A and strict partial orders on A .

A *poset* or *partially ordered set* is a pair (A, \leq) where \leq is a partial order on A . In a poset (A, \leq) , two elements x, y are *comparable* if $x \leq y$ or $y \leq x$; otherwise they are *incomparable*. A *linear order* or *total order* is a partial order in which any two elements are comparable. The relation \leq on \mathbb{R} is a linear order, but \subseteq is usually not a linear order on $\mathfrak{P}(X)$.

A.5 Functions

A *function* or *map* from A to B is a relation $f \subseteq A \times B$ such that for every $x \in A$, there is exactly one $y \in B$ such that $x f y$. We write this unique value of y as $f(x)$, so that

$$f(x) = y \iff (x f y) \iff (x, y) \in f.$$

The notation $x \mapsto y$ means that $f(x) = y$ for whatever function f we are talking about. Note the bar at the left end of \mapsto . The notation $f : A \rightarrow B$ means that A and B are sets and f is a function from A to B . The sets A and B are called the *domain* and *codomain* of f . The domain of f is written $\text{dom}(f)$. The *range* or *image* of f is

$$\text{im}(f) = \{f(x) : x \in A\} = \{y \in B : \text{there is } x \in A \text{ such that } f(x) = y\}.$$

If X is a subset of A , then the *image* or *direct image* of X is the set

$$f(X) = \{f(x) : x \in X\} = \{y \in B : \text{there is } x \in X \text{ such that } f(x) = y\}.$$

If Y is a subset of B , then the *preimage* or *inverse image* of Y is the set

$$f^{-1}(Y) = \{x \in A : f(x) \in Y\}.$$

If f is a function from A to B , then f is a...

1. *surjection* if for every $y \in B$, there is at least one $x \in A$ with $f(x) = y$.
2. *injection* if for every $y \in B$, there is at most one $x \in A$ with $f(x) = y$.
3. *bijection* if for every $y \in B$, there is exactly one $x \in A$ with $f(x) = y$.

We say that f is *surjective*, *injective*, or *bijective* if f is a surjection, injection, or bijection. Note the following:

- f is a surjection if and only if $\text{im}(f) = B$.
- f is an injection if and only if

$$f(x) = f(y) \implies x = y$$

if and only if

$$x \neq y \implies f(x) \neq f(y).$$

- f is a bijection if and only if f is injective and surjective.

If $f : A \rightarrow B$ is a bijection, then there is an *inverse function* $f^{-1} : B \rightarrow A$ defined by

$$f^{-1}(y) = x \iff f(x) = y.$$

If $f : A \rightarrow B$ and $g : B \rightarrow C$ are functions, then the *composition* is the function $h : A \rightarrow C$ defined by

$$h(x) = f(g(x)).$$

The composition is usually written $f \circ g$. Note that $f \circ g$ is the function where g is applied first and f is applied second.

If A is a set, then id_A or id denotes the *identity function* on A , the function

$$\text{id}_A : A \rightarrow A$$

$$\text{id}_A(x) = x.$$

Note that if $f : A \rightarrow B$ is a function then

$$f \circ \text{id}_A = f$$

$$\text{id}_B \circ f = f.$$

If $f : A \rightarrow B$ is a bijection and $f^{-1} : B \rightarrow A$ is the inverse, then

$$f^{-1} \circ f = \text{id}_A$$

$$f \circ f^{-1} = \text{id}_B.$$

If A, B, C are sets, then a function $f : A \times B \rightarrow C$ is a function from the cartesian product $A \times B$ to C . If $x \in A$ and $y \in B$, we write $f((x, y))$ as $f(x, y)$, and we think of f as a function which takes one input x from A and one input y from B , and then outputs a value $f(x, y)$ in C .

If A is a set, then A^2, A^3, \dots denote the cartesian product of A with itself n times:

$$A^2 = A \times A$$

$$A^3 = A \times A \times A$$

$$\dots$$

$$A^n = \underbrace{A \times A \times \dots \times A}_{n \text{ times}}.$$

So A^n is the set of n -tuples (x_1, \dots, x_n) such that x_1, x_2, \dots, x_n are all in A . A function $f : A^n \rightarrow B$ can be thought of as a function which takes n inputs x_1, \dots, x_n from A and outputs a value in B . We think of operations like $+$ (addition) and \cdot (multiplication) as functions $\mathbb{R}^2 \rightarrow \mathbb{R}$, taking two inputs from \mathbb{R} and outputting one value in \mathbb{R} . But we usually write $x + y$ rather than $+(x, y)$.

A.6 Cardinalities

If X is a set, then $|X|$ denotes the *size* or *cardinality* of X , the number of elements of X . For example, $|\emptyset| = 0$. When X is finite, $|X|$ is an element of \mathbb{N} . Otherwise, $|X|$ is one of Cantor's *infinite cardinal numbers*, of which the first few are $\aleph_0, \aleph_1, \aleph_2, \dots$. It is worth knowing that

$$\begin{aligned} |\mathbb{N}| &= |\mathbb{Z}| = |\mathbb{Q}| = \aleph_0 \\ |\mathbb{R}| &= |\mathbb{C}| = 2^{\aleph_0} > \aleph_0. \end{aligned}$$

The axioms of set theory neither prove nor disprove the *continuum hypothesis*, the statement that $2^{\aleph_0} = \aleph_1$. If X and Y are sets, then

$$\begin{aligned} |X \times Y| &= |X| \cdot |Y| \\ |X \cup Y| &\leq |X| + |Y| \\ |X \cup Y| &= |X| + |Y| \text{ if } X \cap Y = \emptyset \\ |\mathfrak{P}(X)| &= 2^{|X|} > |X| \\ X \subseteq Y &\implies |X| \leq |Y|. \end{aligned}$$

Moreover, $|X| = |Y|$ if and only if there is a bijection $f : X \rightarrow Y$, and $|X| \leq |Y|$ if and only if there is an injection $f : X \rightarrow Y$. When X and Y are non-empty, $|X| \geq |Y|$ if and only if there is a surjection $X \rightarrow Y$.

If X is a proper subset of Y , then $|X| \leq |Y|$, but it can happen that $|X| = |Y|$ when X and Y are infinite. For example, $\mathbb{N} \subsetneq \mathbb{Z} \subsetneq \mathbb{Q}$, but $|\mathbb{N}| = |\mathbb{Z}| = |\mathbb{Q}| = \aleph_0$.

It is worth knowing that if κ, λ are infinite cardinals, then $\kappa \cdot \lambda = \kappa + \lambda = \max(\kappa, \lambda)$.

The notation $|X| < \infty$ means that the set X is finite, and $|X| = \infty$ means that X is infinite. This is an abuse of notation—there is no cardinal

number “ ∞ .” In fact,

$$\begin{aligned}|X| < \infty &\iff |X| < \aleph_0 \\ |X| = \infty &\iff |X| \geq \aleph_0.\end{aligned}$$

In set theory, \aleph_0 , \mathbb{N} , and ω are all the same thing, so we sometimes write \aleph_0 as ω .

Appendix B

Set theory

We will use the following slightly more advanced definitions and facts from set theory.

B.1 Ordinals

In set theory, one defines a class Ord of *von Neumann ordinal numbers*, also called *ordinal numbers* or *ordinals*. Rather than defining Ord , we treat it as a black box. Here are its important properties:

Fact B.1.1. 1. *There is a linear order \leq on the class of ordinals.*

2. *\leq is a well-ordering: if S is a non-empty set or class of ordinals, then $\min(S)$ exists.*

3. *For each $x \in Ord$,*

$$x = \{y \in Ord : y < x\}.$$

In particular, the right hand side is a set, not a proper class.

4. *Consequently, if $x, y \in Ord$, then $x < y \iff x \in y$.*

5. *Also, if $x, y \in Ord$, then $x \leq y \iff x \subseteq y$.*

6. *If S is a set of ordinals (not a proper class), then the supremum $\sup(S)$ exists.*

The first few ordinals are

$$0, 1, 2, 3, \dots, \omega, \omega + 1, \omega + 2, \dots, \omega 2, \omega 2 + 1, \dots$$

where

$$0 = \{\} = \emptyset$$

$$1 = \{0\}$$

$$2 = \{0, 1\}$$

$$3 = \{0, 1, 2\}$$

$$\dots$$

$$\omega = \{0, 1, 2, 3, \dots\} = \mathbb{N}$$

$$\omega + 1 = \{0, 1, 2, 3, \dots, \omega\}$$

$$\omega + 2 = \{0, 1, 2, 3, \dots, \omega, \omega + 1\}$$

$$\dots$$

$$\omega \cdot 2 = \{0, 1, 2, 3, \dots, \omega, \omega + 1, \omega + 2, \dots\}$$

$$\omega \cdot 2 + 1 = \{0, 1, 2, 3, \dots, \omega, \omega + 1, \omega + 2, \dots, \omega \cdot 2\}$$

$$\dots$$

If α is an ordinal, there is a smallest ordinal greater than α , which is called $\alpha + 1$. Ordinals of the form $\alpha + 1$ are called *successor ordinals*. Non-zero ordinals that are not successors are called *limit ordinals*. The first few limit ordinals are ω and $\omega \cdot 2$.

B.2 Induction and recursion

Let α be an ordinal and S be a subset of α , i.e., a set of ordinals less than α . Suppose that the following condition holds for any $x \in \alpha$:

$$\text{If every } y < x \text{ is in } S, \text{ then } x \in S. \quad (*_x)$$

Then $S = \alpha$, a fact called *induction*. To prove this, suppose $S \subsetneq \alpha$. Then $\alpha \setminus S$ is a non-empty set of ordinals, so it has a minimum $x = \min(\alpha \setminus S)$. If $y < x$, then $y \notin \alpha \setminus S$ because x was the minimum, and so $y \in S$. Then $(*_x)$ says that $x \in S$, contradicting the fact that we took x in the complement of S .

Closely related to induction is the fact that we can recursively define functions on α :

Fact B.2.1. *Let α be an ordinal and V be a set. Let \mathcal{P} be the set of pairs (x, f) , where $x \in \alpha$ and f is a function from x to V . Let $G : \mathcal{P} \rightarrow V$ be a function. Then there is a unique function $f : \alpha \rightarrow V$ such that for every $x \in \alpha$, $f(x) = G(f|_x)$, where $f|_x$ is the restriction of f to the set $x = \{y \in \alpha : y < x\}$.*

The notation is confusing, but the point is that the value $f(x)$ is determined by the values $f(y)$ for $y < x$. The uniqueness of f is an easy consequence of induction, but the existence of f takes more work. See a book on set theory for a proof.

B.3 Cardinals

If A, B are sets, formally define $|A| = |B|$ to mean that there is a bijection from A to B . This is an equivalence relation on the class of sets.

Fact B.3.1. *If S is any set, then there is an ordinal x with $|S| = |x|$.*

The well-ordering property lets us choose a system of representatives for the equivalence relation $|A| = |B|$.

Definition B.3.2. $|S|$ is the smallest ordinal x such that $|x| = |S|$.

Then $|A| = |B|$ agrees with the previous notation, but $|A|$ now has a concrete meaning.

Definition B.3.3. A (von Neumann) *cardinal number* or *cardinal* is an ordinal of the form $|A|$ for some set A .

Equivalently, an ordinal x is a cardinal if $|y| < |x|$ for every $y < x$ (i.e., every $y \in x$).

The class of cardinals is well-ordered: for any non-empty set of cardinals there is a smallest member. The first few cardinals are

$$0, 1, 2, 3, \dots, \aleph_0, \aleph_1, \aleph_2, \dots, \aleph_\omega, \aleph_{\omega+1}, \dots$$

The cardinal \aleph_0 is the same as the ordinal ω . The cardinal \aleph_1 equals an ordinal called ω_1 , the first uncountable ordinal. Between ω and ω_1 there are MANY countable ordinals, of which the first few are

$$\omega, \omega + 1, \dots, \omega \cdot 2, \dots, \omega \cdot 3, \dots, \omega^2, \dots, \omega^\omega, \dots, \omega^{\omega^\omega}, \dots, \omega^{\omega^{\omega^\omega}}, \dots$$

B.4 Zorn's lemma

Let (P, \leq) be a poset, i.e., a pair P and a partial order \leq on P . A *chain* is a set $I \subseteq P$ that is linearly ordered by \leq , in the sense that for any $x, y \in I$, either $x \leq y$ or $y \leq x$. If $I \subseteq P$ is a chain, an *upper bound* is an element $b \in P$ such that $b \geq I$, meaning that $b \geq x$ for all $x \in I$.

For example, if $(P, \leq) = (\mathfrak{P}(\mathbb{N}), \subseteq)$, then the family of sets

$$\begin{aligned} &\{0\}, \\ &\{0, 2\}, \\ &\{0, 2, 4\}, \\ &\{0, 2, 4, 6\}, \\ &\dots \end{aligned}$$

is a chain, and $\{0, 2, 4, 6, 8, 10, \dots\}$ is an upper bound.

An element $x \in P$ is *maximal* if there is no $y \in P$ with $y > x$. For example, in $(\mathfrak{P}(\mathbb{N}), \subseteq)$, the element \mathbb{N} is the unique maximal element. In general, there can be more than one maximal element, or none.

Fact B.4.1 (Zorn's lemma). *Let (P, \leq) be a poset such that every chain $I \subseteq P$ has an upper bound. Then P has a maximal element.*

If you're curious, here is a sketch of the proof:

Proof sketch. Suppose for the sake of contradiction that P has no maximal element. For any $x \in P$, there is $x' \in P$ with $x' > x$; otherwise x is maximal. Choose a function $x \mapsto x'$ with this property.

Take an ordinal α with $|\alpha| > |P|$.¹ Recursively define $f : \alpha \rightarrow P$ as follows. At step α , let $I_\alpha = \{f(x) : x < \alpha\}$. The set I_α will be a chain in P by induction. Take an upper bound $b \geq I_\alpha$ and set $f(\alpha) = b'$. Then $f(\alpha) > I_\alpha$, so $f(\alpha) > f(x)$ for all $x < \alpha$.

This gives a map $f : \alpha \rightarrow P$ which is strictly increasing:

$$x < y \implies f(x) < f(y).$$

Then f is an injection, and $|\alpha| \leq |P|$, contradicting the choice of α . □

For more details of the proof, see any book on set theory.

Most applications of Zorn's lemma take the following form:

¹More precisely, we want $|\alpha| \not\leq |P|$. Finding α is the hard part of the proof.

Corollary B.4.2. *Let S be a set and let $\mathcal{F} \subseteq \mathfrak{P}(S)$ be a collection of subsets of S . Suppose that whenever \mathcal{I} is a chain in (\mathcal{F}, \subseteq) , the union $\bigcup \mathcal{I}$ is in \mathcal{F} . Then \mathcal{F} has a maximal element.*

Proof. Apply Zorn's lemma to the poset (\mathcal{F}, \subseteq) . □

Example B.4.3. A *graph* is a pair (S, R) where S is a set and $R \subseteq S^2$ is a relation that is symmetric and irreflexive:

$$\begin{aligned} x R y &\implies y R x \\ x &\not R x. \end{aligned}$$

(If you know graph theory, think of S as the set of vertices, and $x R y$ as the relation saying that there is an edge from x to y . We are only considering simple, undirected graphs.)

Fix a graph (S, R) . Say that $X \subseteq S$ is *independent* if for any $x, y \in X$, we have $x \not R y$. Let $\mathcal{F} \subseteq \mathfrak{P}(S)$ be the collection of independent sets. Then there is a maximal independent set by Zorn's lemma, specifically Corollary B.4.2. To apply Corollary B.4.2, we must show that if $\mathcal{I} \subseteq \mathcal{F}$ is a chain, then $X_\infty := \bigcup \mathcal{I}$ is independent. Suppose for the sake of contradiction that $x_1, x_2 \in X_\infty$ and $x_1 R x_2$. As $x_1 \in X_\infty = \bigcup \mathcal{I}$, there is some $X_1 \in \mathcal{I}$ such that $x_1 \in X_1$. Similarly, there is $x_2 \in X_2 \in \mathcal{I}$. As \mathcal{I} is a chain, either $X_1 \subseteq X_2$ or $X_2 \subseteq X_1$. In the first case, $x_1, x_2 \in X_2$, and X_2 fails to be independent, contradicting the fact that $X_2 \in \mathcal{I} \subseteq \mathcal{F}$. Similarly, if $X_2 \subseteq X_1$ then $x_1, x_2 \in X_1$, and X_1 fails to be independent. Thus X_∞ is independent and Zorn's lemma applies.

In this case, maximality means that we have an independent set X such that $X \cup \{a\}$ is *not* independent for any $a \in S \setminus X$. Therefore, for any $a \in S \setminus X$ there is $b \in X$ with $a R b$. Thus X is an independent set and every element outside of X is adjacent (connected by an edge) to an element of X .