

Universal theories, heirs, and ultrapowers

Advanced Model Theory

April 6, 2022

Remember those weird theorems saying that some type embeds into an ultrapower of another type? (March 3, Proposition 2, and March 31, Proposition 7.) Let's get a better conceptual understanding of why they're true. Along the way, we'll learn some useful model theory.

1 Universal theories

Definition 1. A *universal formula* is a formula of the form $\forall \bar{y} \varphi(\bar{x}, \bar{y})$, where φ is quantifier-free. A universal sentence is a sentence that is universal, or equivalently, a sentence of the form $\forall \bar{x} \varphi(\bar{x})$ with φ quantifier-free.

Proposition 2. Let M be a substructure of N .

1. If $\varphi(\bar{x})$ is a quantifier-free formula and $\bar{a} \in M$, then

$$M \models \varphi(\bar{a}) \iff N \models \varphi(\bar{a}).$$

2. If $\varphi(\bar{x})$ is a universal formula and $\bar{a} \in M$, then

$$N \models \varphi(\bar{a}) \implies M \models \varphi(\bar{a})$$

Proof. “Obvious,” but here are the details for completeness:

1. If φ is atomic, this holds by definition of “substructure.” Otherwise, proceed by induction on the complexity of φ .
2. Write $\varphi(\bar{x})$ as $\forall \bar{y} \psi(\bar{x}, \bar{y})$ for quantifier-free $\psi(\bar{x}, \bar{y})$. If $N \models \varphi(\bar{a})$, then

$$\begin{aligned} N &\models \forall \bar{y} \psi(\bar{a}, \bar{y}) \\ \forall \bar{b} \in N &(N \models \psi(\bar{a}, \bar{b})) \\ \forall \bar{b} \in M &(N \models \psi(\bar{a}, \bar{b})) \quad \text{as } M \subseteq N \\ \forall \bar{b} \in M &(M \models \psi(\bar{a}, \bar{b})) \quad \text{by part (1)} \\ M &\models \forall \bar{y} \psi(\bar{a}, \bar{y}) \\ M &\models \varphi(\bar{a}). \end{aligned}$$

□

Definition 3. A *universal theory* is a set of universal sentences.

Proposition 4. Let T be a universal theory.

1. If $N \models T$ and M is a substructure of N , then $M \models T$.
2. If $f : M \rightarrow N$ is an embedding and $N \models T$, then $M \models T$.

Proof.

1. By Proposition 2(2), the axioms of T transfer from N to M .
2. If M' is the image of f , then $M \cong M'$ and M' is a substructure of N , so $N \models T \implies M' \models T \implies M \models T$. \square

Example. A *semigroup* is a structure (G, \cdot) satisfying the universal sentence

$$\forall x, y, z (x \cdot (y \cdot z) = (x \cdot y) \cdot z).$$

Proposition 4(1) implies that any substructure of a semigroup is a semigroup (which is obvious).

Definition 5. If T is a theory, then T_\forall is the set of universal sentences φ such that $T \vdash \varphi$.

Lemma 6. Suppose $N \models T$. If M is a substructure of N , then $M \models T_\forall$. More generally, if there is an embedding $M \rightarrow N$, then $M \models T_\forall$.

Proof. If $N \models T$ then $N \models T_\forall$, because the axioms of T_\forall are logical consequences of the axioms of T . Then $M \models T_\forall$ by Proposition 4. \square

Using compactness, we can prove a converse of Lemma 6.

Lemma 7. Suppose $M \models T_\forall$. Then there is a model $N \models T$ and an embedding $M \rightarrow N$. We may even take N to be an extension of M .

Proof. Consider the language $L(M)$ where we add a constant symbol for each element of M . Let D be the set of quantifier-free $L(M)$ -sentences true in M . (D is usually called the *diagram* of M .)

Claim. $D \cup T$ is consistent.

Proof. Otherwise, by compactness there are $\varphi_1, \dots, \varphi_n \in D$ such that $\{\varphi_1, \dots, \varphi_n\} \cup T$ is inconsistent. Let $\varphi = \bigwedge_{i=1}^n \varphi_i$. Then $\varphi \in D$ and $\{\varphi\} \cup T$ is inconsistent. Note φ is a

quantifier-free $L(M)$ -sentence. Write φ as $\theta(\bar{c})$ for some quantifier-free L -formula θ and some tuple $\bar{c} \in M$.¹ As $M \models \theta(\bar{c})$ and $M \models T_\forall$,

$$\begin{aligned} M &\not\models \forall \bar{x} \neg \theta(\bar{x}) \\ (\forall \bar{x} \neg \theta(\bar{x})) &\notin T_\forall \\ T &\not\models \forall \bar{x} \neg \theta(\bar{x}). \end{aligned}$$

Therefore there is a model $N \models T$ with

$$\begin{aligned} N &\models \neg \forall \bar{x} \neg \theta(\bar{x}) \\ N &\models \exists \bar{x} \theta(\bar{x}) \\ N &\models \theta(\bar{e}) \end{aligned}$$

for some $\bar{e} \in N$. Expand the L -structure N to an $L(M)$ -structure by interpreting $c_i \in M$ as e_i and interpreting the other elements of M randomly.² Then $N \models \theta(\bar{c})$. Consequently, $N \models \{\varphi\} \cup T$, contradicting the fact that $\{\varphi\} \cup T$ is inconsistent. \square_{Claim}

By the Claim and compactness, there is some $L(M)$ -structure N with $N \models D \cup T$. Then N is a model of T . Let $f : M \rightarrow N$ be the map sending $c \in M$ to its interpretation in N (which is an $L(M)$ -structure). The fact that $N \models D$ means that f is an embedding. For example:

- If $c, d \in M$ and $c \neq d$, then $M \models c \neq d$, so $(c \neq d) \in D$, so $N \models c \neq d$, which really means $f(c) \neq f(d)$.
- If R is a binary relation and $M \models R(c, d)$, then $R(c, d) \in D$, so $N \models R(c, d)$, which really means $N \models R(f(c), f(d))$.

So we have embedded M into a model of T .

For the final clause of Lemma 7, once we have an embedding $f : M \rightarrow N$ with $N \models T$, we can find an isomorphism $g : N \xrightarrow{\cong} N'$ such that $g(f(x)) = x$ for $x \in M$. Then $N' \models T$ and the embedding $M \rightarrow N'$ is an inclusion, meaning that M is a substructure of N' . \square

For future reference, we note the following Fact:

Fact 8. *Let M, N be L -structures. Let T be the complete L -theory of N . Suppose $M \models T_\forall$. (In other words, M satisfies all the universal sentences that N satisfies.)*

¹For technical reasons, we need to make two conditions hold. (1) The coordinates of \bar{c} should be pairwise distinct. (2) If $M \neq \emptyset$, then \bar{c} should have length at least one. These two conditions can be ensured as follows. Let c_1, \dots, c_n enumerate *without repeats* the elements of M mentioned in φ . But if M is non-empty and φ mentions no elements of M , take $n = 1$ and take $c_1 \in M$ arbitrary. Every element of M mentioned in φ is one of the c_i , so we can write φ as $\theta(\bar{c})$ for some θ .

²There are two technical details to check. (1) If $c_i = c_j$, we need e_i to equal e_j . This works because we made $c_i \neq c_j$ for $i \neq j$. (2) If there are elements of M outside of the c_i , we need N to be non-empty. This works because when $M \neq \emptyset$, we arranged for the length of $\bar{c}, \bar{x}, \bar{e}$ to be positive, and then $\bar{e} \in N \implies N \neq \emptyset$.

1. There is $N' \equiv N$ and an embedding $M \rightarrow N'$.

2. There is an ultrapower $N^{\mathcal{U}}$ and an embedding $M \rightarrow N^{\mathcal{U}}$.

(1) is a direct consequence of Lemma 7. (2) should sound plausible, because Lemma 7 was proved using compactness, and compactness was proved using ultraproducts.

Combining Lemmas 6 and 7, we get the following useful fact:

Theorem 9. *Let T be a theory. Then $M \models T_{\forall}$ iff M is a substructure of a model of T .*

Fact 10. *Here are two examples:*

1. DLO_{\forall} is the theory of linear orders. A structure (M, \leq) is a linear order iff (M, \leq) embeds into a model of DLO.
2. ACF_{\forall} is the theory of integral domains. A structure $(R, +, \cdot, -, 0, 1)$ is an integral domain iff $(R, +, \cdot, -, 0, 1)$ embeds into a model of ACF.

If you're curious, here is a proof of (1). If $(M, \leq) \models \text{DLO}_{\forall}$, then (M, \leq) is a substructure of a dense linear order, and so (M, \leq) is a linear order. Conversely, suppose (M, \leq) is a linear order. We claim $(M, \leq) \models \text{DLO}_{\forall}$. Otherwise, there is a universal sentence $\forall \bar{x} \varphi(\bar{x})$ such that

$$\begin{aligned} \text{DLO} &\vdash \forall \bar{x} \varphi(\bar{x}) \\ M &\not\models \forall \bar{x} \varphi(\bar{x}). \end{aligned}$$

Take $\bar{a} \in M$ with $M \models \neg \varphi(\bar{a})$. Let $A = \{a_1, \dots, a_n\}$, viewed as a substructure of M . By Proposition 2,

$$M \models \neg \varphi(\bar{a}) \implies A \models \neg \varphi(\bar{a}).$$

But then

$$A \models \neg \varphi(\bar{a}) \implies A \not\models \forall \bar{x} \varphi(\bar{x}) \implies A \not\models \text{DLO}_{\forall}.$$

By Lemma 6, A does not embed into any model of DLO. But finite linear orders embed into $(\mathbb{Q}, \leq) \models \text{DLO}$, a contradiction.

Here is another useful variant of Theorem 9:

Theorem 11. *Let T be a theory. The following are equivalent:*

1. T can be axiomatized by universal sentences.
2. If $N \models T$ and M is a substructure of N , then $M \models T$.

Proof. (1) \implies (2): Proposition 4.

(2) \implies (1): Assume (2). We claim that T and T_{\forall} are logically equivalent. Certainly $M \models T \implies M \models T_{\forall}$. Conversely, suppose $M \models T_{\forall}$. By Lemma 7, there is an extension $N \supseteq M$ with $N \models T$. By (2), $M \models T$. So T is axiomatized by the universal theory T_{\forall} . \square

Remark 12. An $\forall\exists$ -sentence is a sentence of the form $\forall\bar{x} \exists\bar{y} \varphi(\bar{x}, \bar{y})$ where φ is quantifier-free. The following fact is similar to Theorem 11 and worth knowing. (But the proof is a little harder.)

Fact 13. *If T is a theory, the following are equivalent:*

1. T can be axiomatized by $\forall\exists$ -sentences.
2. If $M_1 \subseteq M_2 \subseteq M_3 \subseteq \dots$ is an increasing chain of models of T , then $\bigcup_{i=1}^{\infty} M_i$ is a model of T .

2 The fundamental order and heirs

Recall the structures (M, dp) from the first lecture on February 24: if $M \models T$ and $p \in S_n(M)$, then (M, dp) is the expansion of M by new symbols $d\varphi$ for each L -formula $\varphi(\bar{x}, \bar{y})$, where

$$d\varphi = \{\bar{b} \in M : \varphi(\bar{x}, \bar{b}) \in p(\bar{x})\}.$$

You can check that the following equivalences hold in (M, dp) for formulas $\varphi(\bar{x}, \bar{y}), \psi(\bar{x}, \bar{y})$:

$$(d\varphi) \wedge (d\psi) \iff d(\varphi \wedge \psi) \tag{1}$$

$$(d\varphi) \vee (d\psi) \iff d(\varphi \vee \psi) \tag{2}$$

$$\neg(d\varphi) \iff d(\neg\varphi) \tag{3}$$

$$\varphi \iff d\varphi \quad (\text{when } \bar{x} \text{ has length } 0.) \tag{4}$$

Let L_d be the language of structures like $(M, dp), (N, dq)$. Using (1)–(4), one can see the following:

Fact 14. *Any quantifier-free L_d -formula $\psi(\bar{y})$ is equivalent to $d\varphi(\bar{y})$ for some L -formula $\varphi(\bar{x}, \bar{y})$.*

Also, Equation (4) means we secretly added a relation symbol for every L -formula $\varphi(\bar{y})$ on M , and so

Fact 15. *Any L -formula is equivalent to a quantifier-free L_d -formula.*

Combining these facts and the previous section, we can get a clearer proof of something mentioned in class:

Proposition 16. *If $p \in S_n(M)$ and $q \in S_n(N)$ and $[q] \leq [p]$, then there is an ultrapower $N^{\mathcal{U}}$ and an elementary embedding $f : M \rightarrow N^{\mathcal{U}}$ making $q^{\mathcal{U}}$ an extension of p .*

Proof.

Claim. (M, dp) satisfies all the universal L_d -sentences that (N, dq) satisfies.

Proof. Suppose $(N, dq) \models \forall \bar{y} \varphi(\bar{y})$ for some quantifier-free L_d -formula $\varphi(\bar{y})$. By Fact 14, $\varphi(\bar{y})$ is equivalent to $d\theta$ for some L -formula $\theta(\bar{x}, \bar{y})$. Then

$$\begin{aligned} (N, dq) &\models \forall \bar{y} d\theta(\bar{y}) \\ \forall \bar{b} \in N : \theta(\bar{x}, \bar{b}) &\in q(\bar{x}) \\ \forall \bar{b} \in N : (\neg\theta(\bar{x}, \bar{b})) &\notin q(\bar{x}). \end{aligned}$$

So q omits the formula $\neg\theta(\bar{x}, \bar{y})$. As $[q] \leq [p]$, we see that p omits the formula $\neg\theta(\bar{x}, \bar{y})$. Then

$$\begin{aligned} \forall \bar{b} \in M : (\neg\theta(\bar{x}, \bar{b})) &\notin p(\bar{x}) \\ \forall \bar{b} \in M : \theta(\bar{x}, \bar{b}) &\in p(\bar{x}) \\ (M, dp) &\models \forall \bar{y} d\theta(\bar{y}) \\ (M, dp) &\models \forall \bar{y} \varphi(\bar{y}). \end{aligned}$$

□_{Claim}

By Fact 8(2), there is an ultrafilter \mathcal{U} and embedding $(M, dp) \rightarrow (N, dq)^\mathcal{U} = (N^\mathcal{U}, dq^\mathcal{U})$. The embedding $M \rightarrow N^\mathcal{U}$ preserves all quantifier-free L_d -formulas, so it preserves *all* L -formulas by Fact 15. That is, $M \rightarrow N^\mathcal{U}$ is an elementary embedding. The fact that $(M, dp) \rightarrow (N^\mathcal{U}, dq^\mathcal{U})$ is an embedding implies that $q^\mathcal{U}$ extends p . □

What about heirs?

Proposition 17. *Suppose $N \succeq M$ and $q \in S_n(N)$ is an heir of $p \in S_n(M)$. Then there is an ultrapower $M^\mathcal{U}$ and an elementary embedding $N \rightarrow M^\mathcal{U}$ over M , making $p^\mathcal{U}$ extend q .*

Here, “over M ” means that $N \rightarrow M^\mathcal{U}$ fixes the elements of M . If we’re willing to move $M^\mathcal{U}$ by an isomorphism, this lets us arrange

$$\begin{aligned} M &\preceq N \preceq M^\mathcal{U} \\ p &\subseteq q \subseteq p^\mathcal{U}. \end{aligned}$$

Proof. Note that the following are equivalent:

1. q is an heir of p .
2. If $\varphi(\bar{x}, \bar{y})$ is an $L(M)$ -formula and $\varphi(\bar{x}, \bar{b}) \in q(\bar{x})$ for some $\bar{x} \in N$, then there is $\bar{b}' \in M$ with $\varphi(\bar{x}, \bar{b}') \in p(\bar{x})$.
3. If we expand to the language $L(M)$, then $[p] \leq [q]$ in the fundamental order.

Expand to $L(M)$. Then $[p] \leq [q]$. By Proposition 16, there is an ultrapower $M^\mathcal{U}$ and an elementary embedding $N \rightarrow M^\mathcal{U}$ making $p^\mathcal{U}$ extend q . This is an elementary embedding of $L(M)$ -structures, so it fixes the elements of M . □