

Introductory model theory with elements of  
universal algebra  
(version 3)

Will Johnson

Autumn 2022



# Preface

This is a provisional draft of the course notes for Introductory Model Theory at Fudan University in Autumn 2022. The goal is to present the basic topics of model theory without assuming much background in abstract algebra. Please send any comments or corrections to [willjohnson@fudan.edu.cn](mailto:willjohnson@fudan.edu.cn).

**The current draft is missing any examples, motivations, and explanations after Chapter 1.** I hope to remedy this soon.

## The approach of these notes

Many textbooks in model theory assume a large amount of mathematical background knowledge, especially in abstract algebra. For example, Poizat’s *Course in Model Theory*—which is better than most textbooks in this regard—assumes you know the definition of fields, rings, homomorphisms, and ideals.

There is a good reason for this approach: model theory as a subject is mostly about the applications of mathematical logic to other branches of mathematics, especially abstract algebra. However, this works poorly for our class, which is not in the mathematics department and not intended solely for mathematics majors.

In contrast, these notes will try to minimize the required mathematical background. Model theory is a branch of mathematics, so it would be impossible to explain the subject without *any* mathematical background. I assume you know what sets, functions, and ordered pairs are. (Much of this material is reviewed in Appendix A.) But I will not assume any background in abstract algebra. In theory, mathematical logic is a prerequisite for this course, so these notes will assume a little bit of background from mathematical logic, mostly set theory.

One approach would be to avoid the concepts from abstract algebra

(fields, rings, etc.) entirely, instead focusing on combinatorial examples like the random graph. Model theory *can* be presented in this way, but one loses much of the motivation, examples, intuition, and applications of the subject.

Instead, we will build up elements of abstract algebra from scratch, defining concepts like fields, rings, homomorphisms, and ideals and proving their basic properties. This will allow us to work through the model theory of algebraically closed fields, which has served as an important example in the history of the subject.

Our approach towards abstract algebra will emphasize ideas from *universal algebra*. Universal algebra is a subject which finds the parallels between ring theory, group theory, lattice theory, and other topics in algebra. For example, “ideals” in ring theory and “normal subgroups” in group theory are both instances of the more fundamental concept of “congruences” in universal algebra.

There are two reasons to use universal algebra in these notes. On a basic level, it allows us to efficiently explain concepts in ring theory and group theory simultaneously. More importantly, universal algebra is tightly connected to model theory. One could say that universal algebra is a special case of model theory, or model theory is a generalization of universal algebra. For example, Chang and Keisler describe model theory as “universal algebra plus logic” in their textbook. Consequently, one of the goals of these notes will be to emphasize the parallels between universal algebra and model theory.

## Notation and conventions

See Appendix A for a review of the notation and conventions used in this book.

# Introduction

The story of model theory is like a novel that unexpectedly changes genres halfway through. What began as a branch of mathematical logic later became “universal algebra plus logic” in Chang and Keisler’s formulation, and then “algebraic geometry minus fields” in Hodges’ formulation. This makes the subject a little hard to precisely define.

At the end of the day, model theory is a network of closely connected definitions and tools, all related to the notion of “model.” This introduction is a survey of these definitions and tools. The goal of model theory is not so much to understand “models”, but instead to build on this network and apply it to other branches of mathematics.

## 0.1 Models

Mathematical knowledge is obtained through deductive reasoning—through *proofs*. Proofs must begin somewhere, with a set of *axioms*—statements we assume without proof. When reasoning about numbers, one might take the following set of statements as axioms:

1. If  $x$  and  $y$  are numbers, then  $x + y$  and  $x \cdot y$  are numbers.
2. If  $x$  and  $y$  are numbers, then  $x + y = y + x$  and  $x \cdot y = y \cdot x$ .
3. If  $x$ ,  $y$ , and  $z$  are numbers, then  $x + (y + z) = (x + y) + z$  and  $x \cdot (y \cdot z) = (x \cdot y) \cdot z$ .
4. If  $x$ ,  $y$ , and  $z$  are numbers, then  $x \cdot (y + z) = (x \cdot y) + (x \cdot z)$ .
5. There is a number  $0$  such that  $x + 0 = x$  for any number  $x$ .
6. There is a number  $1$  such that  $x \cdot 1 = x$  for any number  $x$ .

7. For any number  $x$ , there is a number called  $-x$  such that  $x + (-x) = 0$ .
8. For any number  $x$  other than 0, there is a number called  $x^{-1}$  such that  $x \cdot (x^{-1}) = 1$ .

We can use these axioms to deduce other facts about numbers. Here is an example:

**Theorem 0.1.1.** *For any number  $x$ ,  $x \cdot 0 = 0$ .*

*Proof.* If  $y = x \cdot 0$ , then

$$\begin{aligned}
 y &\stackrel{(5)}{=} y + 0 \stackrel{(7)}{=} y + (y + (-y)) \stackrel{(3)}{=} (y + y) + (-y) \\
 &= ((x \cdot 0) + (x \cdot 0)) + (-y) \stackrel{(4)}{=} x \cdot (0 + 0) + (-y) \\
 &\stackrel{(5)}{=} x \cdot 0 + (-y) = y + (-y) \stackrel{(7)}{=} 0. \quad \square
 \end{aligned}$$

With more work (Corollary 1.4.14, Theorem 1.4.16), one can prove other algebraic facts, such as

$$\begin{aligned}
 x \cdot y = 0 &\iff (x = 0 \text{ or } y = 0) \\
 (-x) \cdot (-y) &= x \cdot y.
 \end{aligned}$$

How about the following?

$$x + x = y + y \stackrel{?}{\implies} x = y. \quad (*)$$

This is certainly true for real numbers, but can we prove  $(*)$  from the axioms? Surprisingly, **we cannot**. To see this, suppose we that 0 and 1 are the only “numbers”, and that “+” and “ $\cdot$ ” are defined as follows:

+	0	1
0	0	1
1	1	0

$\cdot$	0	1
0	0	0
1	0	1

For example, “ $1 + 1 = 0$ ” under this interpretation. By checking all the cases, one can verify that all of Axioms (1)–(8) hold. Anything which is provable from the axioms, such as Theorem 0.1.1, must be true in this interpretation. However,  $(*)$  fails:

$$1 + 1 = 0 + 0 \text{ but } 1 \neq 0.$$

Therefore  $(*)$  cannot be proven from Axioms (1)–(8).

Our strange way of interpreting “number”, “0”, and “1” is an example of a **model**—in this case a model of Axioms (1)–(8).

**Remark 0.1.2.** Axioms (1)–(8) are called the *field axioms*, and their models are called *fields*. Fields are one of the central objects of study in the branch of mathematics called *abstract algebra*.

## Some history

The toy example above is parallel to an important saga in the history of mathematics. In ancient times, the Hellenistic mathematician Euclid wrote a book called the *Elements* in which he developed geometry from a set of five axioms. Four of these axioms were simple, intuitive statements, like “any two points are on a line” or “any two right angles are congruent.” In contrast, the fifth axiom was a complicated geometric statement about parallel lines. This was the so-called *parallel postulate*.

Many people were unhappy with the parallel postulate, because it seemed less obvious and more arbitrary than the other four axioms. It felt like something that should be a theorem or a lemma, not an axiom. Euclid himself avoided using the parallel postulate until absolutely necessary. *If* one could find a proof of the parallel postulate from the other four axioms, the problem would go away: the parallel postulate could be demoted to a theorem and geometry could be developed on the firm foundation of the four intuitive axioms.

In the 2000 years after Euclid, there were many unsuccessful attempts to find such a proof. Eventually the matter was settled in 1868 when the mathematician Beltrami constructed a model of Euclid’s first four axioms in which the parallel postulate is false. This model is called *hyperbolic geometry*, and is the precursor to the sort of “curved” geometries used in Einstein’s theory of general relativity.

## 0.2 Complete theories

A *theory* is a set of axioms. If  $T$  is a theory, we write  $M \models T$  to indicate that  $M$  is a model of  $T$ , and  $M \models \varphi$  to indicate that  $M$  satisfies a sentence  $\varphi$ . For example, if  $T$  is the field axioms (1)–(8) of the previous section and  $\varphi$  is Statement (\*), then the standard real numbers  $\mathbb{R}$  satisfy both  $T$  and  $\varphi$

$$\mathbb{R} \models T \text{ and } \mathbb{R} \models \varphi,$$

but we constructed a different model  $M$  such that

$$M \models T \text{ and } M \not\models \varphi.$$

A theory  $T$  is *complete* if for every sentence  $\varphi$ , either  $T$  proves  $\varphi$  or  $T$  disproves  $\varphi$ . If we can completely axiomatize a structure, then we can determine all its logical properties:

**Fact 0.2.1.** *Let  $T$  be a complete theory.*

1. *If  $M \models T$ , then for any sentence  $\varphi$ ,*

$$M \models \varphi \iff (\varphi \text{ is provable from } T)$$

2. *If  $T$  is finite or computable, then there is an algorithm which takes a sentence  $\varphi$  as input and outputs whether or not  $\varphi$  is provable from  $T$ .*

The problem we ran into in the previous section is that the field axioms are not complete—they neither prove nor disprove the statement (\*). Can we fix the problem and write down a complete axiomatization for numbers?

To begin with, let's add the following axioms:

$$\begin{aligned} 1 + 1 &\neq 0 \\ 1 + 1 + 1 &\neq 0 \\ 1 + 1 + 1 + 1 &\neq 0 \\ &\dots \end{aligned}$$

The first of these axioms gets rid of the bad model  $M$  of the previous section, which is a good start. Models of the resulting theory are called *fields of characteristic 0*.

Sadly, the resulting theory is still not complete. For example, consider the statement

$$\text{For every number } x, \text{ there is a number } y \text{ with } y^2 = x.$$

This is true if “number” means “complex number”, and false if “number” means “real number.” In other words, the two models  $\mathbb{C}$  and  $\mathbb{R}$  show incompleteness.

We should probably decide whether we are trying to axiomatize the complex numbers  $\mathbb{C}$  or the real numbers  $\mathbb{R}$ . It turns out to be easier to axiomatize  $\mathbb{C}$ .



**Definition 0.2.2.** A field is *algebraically closed* if it satisfies the following axiom: for any numbers  $a_0, a_1, \dots, a_{n-1}$ , there is a number  $x$  such that

$$x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0 = 0.$$

In other words, every polynomial equation has a solution.

The field of complex numbers  $\mathbb{C}$  is algebraically closed, a fact known as the *fundamental theorem of algebra*. In contrast, the field of real numbers  $\mathbb{R}$  is *not* algebraically closed, as  $x^2 + 1 = 0$  has no solutions.

If we combine the axioms defining fields of characteristic zero and the axioms defining algebraically closed fields, we get a theory called  $\text{ACF}_0$ , whose models are algebraically closed fields of characteristic 0. The complex numbers are a model. In Section 9.4 we will prove the following:

**Fact 0.2.3.** *The theory  $\text{ACF}_0$  is complete.*

Consequently:

1. A sentence  $\varphi$  is true in  $\mathbb{C}$  if and only if it is provable from the  $\text{ACF}_0$  axioms.
2. There is an algorithm which takes  $\varphi$  as input and determines whether  $\varphi$  is true in  $\mathbb{C}$ .

**Remark 0.2.4.** There is also a theory called  $\text{RCF}$  which completely axiomatizes  $\mathbb{R}$ . Models of  $\text{RCF}$  are called *real closed fields*.

## 0.3 Categoricity

Model theory provides a number of methods to prove that a theory  $T$  is complete by analyzing the structure of the models of  $T$ . One such method is *categoricity*.

**Definition 0.3.1.** Let  $\kappa$  be an infinite cardinal number like  $\aleph_0$  or  $\aleph_1$ . A theory  $T$  is  $\kappa$ -categorical if  $T$  has exactly one model of size  $\kappa$ .

**Fact 0.3.2** (Łoś-Vaught criterion). *If a theory  $T$  is  $\kappa$ -categorical and all models of  $T$  are infinite, then  $T$  is complete.*

Conveniently, it is a theorem in abstract algebra that  $\text{ACF}_0$  is  $\kappa$ -categorical for all uncountable  $\kappa$ :

**Fact 0.3.3** (Steinitz). *If  $\kappa$  is an uncountable cardinal, then there is exactly one algebraically closed field of characteristic 0 and size  $\kappa$ .*

Algebraically closed fields are infinite (see Theorem 9.2.3), and so  $\text{ACF}_0$  is complete by the Łoś-Vaught criterion.

**Remark 0.3.4.** This technique is less flexible than one might hope for. In fact, algebraically closed fields are the *only* fields which can be completely axiomatized by  $\kappa$ -categorical theories. We will see a little bit of the proof in Section 13.4, but the bulk of the proof requires stability theory and is beyond the scope of this course.

## 0.4 Definable sets

If  $M$  is a model, a subset  $D \subseteq M$  is *definable* if  $D = \{x \in M : \varphi(x)\}$  where  $\varphi(x)$  is a logical statement about  $x$ . For example, the set

$$\{x \in \mathbb{R} : x \text{ has a square root}\} = \{x \in \mathbb{R} : x \geq 0\}$$

is definable in the field  $\mathbb{R}$ . More generally, a set  $D \subseteq M^n$  in  $n$  variables is definable if

$$D = \{(x_1, \dots, x_n) \in M^n : \varphi(x_1, \dots, x_n)\}$$

where  $\varphi(x_1, \dots, x_n)$  is a statement about the variables  $x_1, \dots, x_n$ . For example, the unit circle is definable in the field  $\mathbb{R}$

$$\{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}. \quad (\dagger)$$

In some models, there are very many definable sets:

**Fact 0.4.1** (Robinson). *In the field  $\mathbb{Q}$ , every computable set  $S \subseteq \mathbb{Q}^n$  is definable.*

For example, the set of Fibonacci numbers  $\{1, 2, 3, 5, 8, 13, \dots\}$  is definable. We will see a variant of Fact 0.4.1 in Section 3.9.

In other models, there are very few definable sets:

**Definition 0.4.2.** In a field  $K$ , a set is *constructible* if it is a finite union of sets of the form

$$\begin{aligned} \{(x_1, \dots, x_n) \in K^n : P_1(x_1, \dots, x_n) = P_2(x_1, \dots, x_n) = \dots \\ = P_m(x_1, \dots, x_n) = 0 \neq Q(x_1, \dots, x_n)\} \end{aligned}$$

where  $P_1, \dots, P_m, Q$  are polynomials.

For example, the unit circle  $(\dagger)$  is constructible.

**Fact 0.4.3.** *In algebraically closed fields such as  $\mathbb{C}$ , the definable sets are exactly the constructible sets.*

Constructible sets are a slight generalization of what algebraic geometers call *varieties*, and Fact 0.4.3 is equivalent to a fact called Chevalley's Theorem in algebraic geometry. We will prove Fact 0.4.3 in Section 9.4 as a consequence of *quantifier elimination*.

One consequence of Fact 0.4.3 is that definable sets in one variable are very simple:

**Corollary 0.4.4.** *If  $M$  is an algebraically closed field and  $D \subseteq M$  is definable, then  $D$  is finite or the complement of a finite set.*

One says that a theory  $T$  is *strongly minimal* if the models of  $T$  have this property. It turns out that strong minimality is closely connected to categoricity.

**Fact 0.4.5** (= Corollary 14.2.7). *If  $T$  is strongly minimal and complete, then  $T$  is  $\kappa$ -categorical for all  $\kappa > \aleph_0$ .*

**Remark 0.4.6.** In more advanced model theory, one can show that if  $T$  is  $\kappa$ -categorical for at least one  $\kappa > \aleph_0$ , then  $T$  is tightly connected to a strongly minimal theory in a certain sense. As a consequence,  $T$  is  $\kappa$ -categorical for all  $\kappa > \aleph_0$ , a fact known as *Morley's Theorem*.

Strong minimality also has an interesting consequence for definable sets in more than one variable:

**Fact 0.4.7.** *Let  $M$  be a model of a strongly minimal theory. To each definable set  $D \subseteq M^n$ , we can associate a natural number called the dimension of  $D$ , written  $\dim(D)$ , with the following properties, among others:*

1.  $\dim(X) > 0$  if and only if  $D$  is infinite.
2.  $\dim(M^n) = n$ .
3.  $\dim(X \cup Y) = \max(\dim(X), \dim(Y))$ .
4.  $\dim(X \times Y) = \dim(X) + \dim(Y)$ .
5. If  $f : X \rightarrow Y$  is a definable bijection, then  $\dim(X) = \dim(Y)$ .

The intuition is that  $\dim(D)$  measures the number of “degrees of freedom” of a value in  $D$ . For example, the unit circle  $x^2 + y^2 = 1$  has one degree of freedom, because the value  $x$  almost determines the pair  $(x, y)$ . In contrast, the plane  $\mathbb{C}^2$  has two degrees of freedom, because  $x$  and  $y$  can vary freely. We will verify Fact 0.4.7 in Sections 14.5–14.6.

In the case of algebraically closed fields,  $\dim(D)$  is something that algebraic geometers call *Krull dimension*. More generally, model theorists like to take concepts from algebraic geometry and extend them from algebraically closed fields to other strongly minimal theories. This is perhaps why Hodges characterizes model theory as “algebraic geometry minus fields” in his textbook.

## 0.5 An outline of this book

Chapters 1 and 2 are about *universal algebra*, which is the special case of model theory where axioms must be *equations* like the distributive law or associative law

$$\begin{aligned}x \cdot (y + z) &= (x \cdot y) + (x \cdot z) \\x + (y + z) &= (x + y) + z\end{aligned}$$

rather than more complicated sentences like

For every  $x$ , there is a  $y$  such that  $y \cdot y = x$  or  $y \cdot y = -x$ .

An *equational theory* is a set of equations. Many important classes of structures from abstract algebra, such as groups and rings, are defined by equational theories, as we will see in Chapter 1. Moreover, certain algebraic constructions like products and quotients make sense in any equational theory, as we will see in Chapter 2. In fact, these constructions precisely characterize which classes of structures are defined by equational theories, a fact known as *Birkhoff’s HSP theorem* (Theorem 2.9.2). Along the way, we will develop basic ring theory, using it as a running example of universal algebra. This ring theory will be used later in Chapter 9 when we axiomatize the complex numbers.

We begin the study of model theory proper in Chapter 3, where we define the basic notions of languages, structures, formulas, satisfaction, theories, models, elementary classes, elementary equivalence, elementary maps,

and definable sets. Chapter 4 is about the *compactness theorem*, the most important fundamental tool of model theory. We apply the compactness theorem in Chapter 5 on categoricity, where we prove the Łoś-Vaught criterion (Theorem 5.6.2) mentioned above.

Chapter 6 is about *ultraproducts*, a mysterious construction which plays the same role in general model theory that products, quotients, etc. play in universal algebra. We use ultraproducts to give another proof of the compactness theorem (Theorem 6.2.6), and then discuss some analogues of Birkhoff’s HSP theorem using ultraproducts in Sections 6.3–6.4.

In Chapter 7, we review basic pointset topology, and apply it to build the topological space  $S$  of complete theories. The compactness theorem can be understood as the statement that  $S$  is compact (Theorem 7.3.4), and ultraproducts correspond to a certain kind of limit in this topological space (Theorem 7.3.5). The construction of  $S$  serves as a prototype of several constructions appearing later in the book.

Chapter 8 is about types and quantifier elimination, two topics which happen to be interconnected. A *type* is a description of a potential object which may not exist in the current model, but exists in some other model. Most applications of the compactness theorem boil down to *realizing* types—finding the element described by a type. Meanwhile, *quantifier elimination* is an important technical tool that allows us to model-theoretically analyze good theories and structures. For example, quantifier elimination gives control over definable sets, and can be used to detect whether a theory is complete.

In Chapter 9, we begin studying the model theory of algebraically closed fields, leading to the complete axiomatization of the complex numbers in Corollary 9.4.7. We continue to develop the model theory of algebraically closed fields in later chapters, using it as a running example.

The next two chapters are about models where “anything which can happen does happen”, in two slightly different senses. Chapter 10 is about *existentially closed models*. Existentially closed models are a useful way of constructing new models of a given theory—for example, we will use them to show that algebraically closed fields *exist*. Existentially closed models can also be used to construct new theories and prove quantifier elimination.

Meanwhile, Chapter 11 is about *monster models*—models where every “small” type is realized. Monster models exist (Theorem 11.4.8), and possess a number of nice properties with respect to definable sets, types, and automorphisms (Sections 11.2–11.3). In advanced model theory, it is common to fix a monster model  $\mathbb{M}$  and always work inside  $\mathbb{M}$ . Monster models

also provide a nice way of thinking about quantifier elimination (see Theorem 11.5.4).

The final three chapters return to the theme of categoricity. Recall that a theory is  $\kappa$ -categorical if it has exactly one model of size  $\kappa$ . Chapter 12 is about *countable categoricity*, the case where  $\kappa = \aleph_0$ . It turns out that a structure  $M$  can be axiomatized by an  $\aleph_0$ -categorical theory if and only if  $M$  satisfies certain structural properties with respect to types, definable sets, and automorphisms (Theorems 12.3.5 and 12.3.6). Chapter 13 is a digression on abstract *closure operations* and the special family of closure operations known as *pregeometries* or *matroids*. We use pregeometries to classify vector spaces, getting more examples of  $\kappa$ -categorical theories (Corollary 13.7.6).

Finally, Chapter 14 is about *strongly minimal theories*, a special class of uncountably categorical theories including algebraically closed fields. Using pregeometries, we prove that strongly minimal theories are uncountably categorical (Corollary 14.2.7), and we develop the dimension theory described in Section 0.4 above. In fact, this dimension theory works in *any* uncountably categorical theory, but the proof is beyond the scope of this book.

# Contents

0.1	Models . . . . .	5
0.2	Complete theories . . . . .	7
0.3	Categoricity . . . . .	9
0.4	Definable sets . . . . .	10
0.5	An outline of this book . . . . .	12
<b>1</b>	<b>Algebras and equational classes</b>	<b>19</b>
1.1	Monoids and groups . . . . .	20
1.2	Rings and fields . . . . .	23
1.3	Languages, algebras, and equational classes . . . . .	24
1.4	Homomorphisms and isomorphisms . . . . .	30
<b>2</b>	<b>New algebras from old</b>	<b>37</b>
2.1	Subalgebras and generators . . . . .	37
2.2	Products . . . . .	38
2.3	Descending functions along surjections . . . . .	40
2.4	Congruences and quotients . . . . .	41
2.5	Application: finite fields . . . . .	45
2.6	The fundamental theorem on homomorphisms . . . . .	47
2.7	Application: characteristic of fields . . . . .	50
2.8	Term algebras and free algebras . . . . .	52
2.9	Birkhoff's HSP theorem . . . . .	54
<b>3</b>	<b>First-order logic</b>	<b>57</b>
3.1	Languages, structures, formulas, and satisfaction . . . . .	57
3.2	Theories and elementary classes . . . . .	60
3.3	Common abbreviations . . . . .	60
3.4	Elementary equivalence and embeddings . . . . .	61
3.5	Substructures and extensions . . . . .	63

3.6	Complete theories . . . . .	64
3.7	Completeness via back-and-forth systems . . . . .	65
3.8	Definable sets and functions . . . . .	68
3.9	Incompleteness of Peano Arithmetic . . . . .	69
<b>4</b>	<b>The compactness theorem</b>	<b>73</b>
4.1	A temporary convention . . . . .	73
4.2	Premodels . . . . .	74
4.3	The witness property . . . . .	75
4.4	Compactness via Henkin's method . . . . .	78
<b>5</b>	<b>Categoricity</b>	<b>81</b>
5.1	A useful fact around finite satisfiability . . . . .	81
5.2	Diagrams and elementary amalgamation . . . . .	81
5.3	Tarski-Vaught . . . . .	83
5.4	The Löwenheim-Skolem theorem . . . . .	84
5.5	Absolute categoricity . . . . .	86
5.6	$\kappa$ -categoricity . . . . .	87
<b>6</b>	<b>Ultraproducts</b>	<b>89</b>
6.1	Ultrafilters . . . . .	89
6.2	Ultraproducts . . . . .	90
6.3	Characterizations of elementary classes . . . . .	94
6.4	Universal classes . . . . .	95
<b>7</b>	<b>Topologies</b>	<b>97</b>
7.1	Topological spaces . . . . .	97
7.2	Ultralimits . . . . .	100
7.3	The space of complete theories . . . . .	102
7.4	Stone spaces . . . . .	104
<b>8</b>	<b>Types and quantifier elimination</b>	<b>107</b>
8.1	Types . . . . .	107
8.2	The topology on type space . . . . .	111
8.3	Quantifier elimination . . . . .	113
8.4	Quantifier elimination in DLO . . . . .	116



<b>9</b>	<b>Algebraically closed fields</b>	<b>119</b>
9.1	Polynomial rings . . . . .	119
9.2	The theory ACF . . . . .	122
9.3	Algebraic and transcendental elements . . . . .	123
9.4	Quantifier elimination in ACF . . . . .	124
<b>10</b>	<b>Existentially closed models</b>	<b>127</b>
10.1	Unions of chains of structures . . . . .	127
10.2	Existentially closed models . . . . .	128
10.3	Existentially closed fields . . . . .	130
<b>11</b>	<b>Monster models</b>	<b>133</b>
11.1	Pushing types along partial elementary maps . . . . .	133
11.2	$\kappa$ -saturated models . . . . .	134
11.3	Strongly $\kappa$ -homogeneous models . . . . .	136
11.4	Construction of monster models . . . . .	138
11.5	Saturation and back-and-forth . . . . .	142
11.6	Application: discrete linear orders with endpoints . . . . .	143
<b>12</b>	<b>Countable categoricity</b>	<b>147</b>
12.1	Baire category theorem . . . . .	147
12.2	The omitting types theorem . . . . .	148
12.3	Countably categorical theories . . . . .	150
<b>13</b>	<b>Closure operators and pregeometries</b>	<b>153</b>
13.1	Closure operators . . . . .	153
13.2	Algebraic and definable closure . . . . .	155
13.3	Definable closure, algebraic closure, and substructures . . . . .	156
13.4	Countably categorical fields . . . . .	158
13.5	Algebraic closure in ACF . . . . .	158
13.6	Pregeometries and vector spaces . . . . .	160
13.7	The classification of vector spaces . . . . .	166
<b>14</b>	<b>Strongly minimal and geometric theories</b>	<b>169</b>
14.1	Strong minimality . . . . .	169
14.2	Uncountable categoricity . . . . .	170
14.3	Uniform finiteness . . . . .	172
14.4	Pregeometric theories . . . . .	174

14.5	Dimension theory in the monster . . . . .	176
14.6	Dimension theory in small models . . . . .	180
<b>A</b>	<b>Notation and conventions</b>	<b>185</b>

# Chapter 1

## Algebras and equational classes

In this chapter, we introduce several important classes of structures from abstract algebra, including *monoids*, *groups*, *rings*, and *fields*. The definitions of these concepts look very similar to each other on an abstract level. Each definition has the form

A FOO is a set together with some operations, satisfying a certain list of equations,

where FOO is “monoid”, “group”, “ring”, and so on.

This suggests the abstract notion of an *equational class*—a class of structures defined by a set of equations. The class of monoids is an equational class, and the same holds for groups and rings. We precisely define equational classes in Section 1.3. Equational classes allow us to uniformly treat certain concepts in algebra. For example, in Section 1.4, we define the important algebraic concepts of *homomorphisms* and *isomorphisms* in a uniform way across all equational classes.

The study of equational classes is called *universal algebra*, and is the subject of this chapter and Chapter 2. Model theory proper, which begins in Chapter 3, can be seen as a generalization of universal algebra. We will use universal algebra as both a source of inspiration for model theory, and as a technical tool to develop the ring theory we will need later when we axiomatize the complex numbers in Chapter 9.

## 1.1 Monoids and groups

Let  $A$  be a set. For  $k \geq 0$ , a  $k$ -ary operation or  $k$ -ary function on  $A$  is a function  $f : A^k \rightarrow A$ , that is, a function which takes  $k$  inputs from  $A$  and outputs one value in  $A$ . *Binary*, *unary*, and *nullary* mean 2-ary, 1-ary, and 0-ary, respectively.

For example, the function

$$\max(x, y) = \begin{cases} x & \text{if } x \geq y \\ y & \text{if } x \leq y \end{cases}$$

is a function  $\mathbb{R}^2 \rightarrow \mathbb{R}$ , so it is a binary (2-ary) operation on  $\mathbb{R}$ . Other binary operations on  $\mathbb{R}$  include  $+$ ,  $-$ , and  $\times$ . Most binary operations are written in infix notation, like  $2 + 3$ , rather than prefix notation, like  $+(2, 3)$ .

Note that  $A^0$  has one element  $()$ . Therefore, we can identify nullary functions  $f : A^0 \rightarrow A$  with elements of  $A$  via the correspondence

$$f \mapsto f().$$

For example, we identify the element  $8 \in \mathbb{R}$  with the function  $f : \mathbb{R}^0 \rightarrow \mathbb{R}$  sending the one element  $()$  in  $\mathbb{R}^0$  to the element  $8 \in \mathbb{R}^1$ .

A *magma* is a pair  $(A, \star)$  where  $A$  is a set and  $\star$  is a binary operation on  $A$ . For example,  $(\mathbb{R}, +)$  and  $(\mathbb{R}, -)$  are magmas. A magma  $(A, \star)$  is a semigroup if  $\star$  is associative, meaning that:

$$x \star (y \star z) = (x \star y) \star z$$

for all  $x, y, z \in A$ . For example,  $(\mathbb{R}, +)$  and  $(\mathbb{R}, \max)$  are semigroups, because

$$\begin{aligned} x + (y + z) &= (x + y) + z \\ \max(x, \max(y, z)) &= \max(x, y, z) = \max(\max(x, y), z). \end{aligned}$$

On the other hand,  $(\mathbb{R}, -)$  is not a semigroup, because

$$5 - (3 - 2) = 4 \neq (5 - 3) - 2 = 0.$$

Let  $(A, \star)$  be a semigroup. An element  $e \in A$  is an *identity element* if

$$x \star e = x = e \star x$$

holds for all  $x \in A$ . For example, 0 is an identity element in the semigroup  $(\mathbb{R}, +)$  because  $x + 0 = 0 + x = x$ . Similarly, 1 is an identity element in the semigroup  $(\mathbb{R}, \cdot)$ , because  $x \cdot 1 = 1 \cdot x = x$ . In contrast,  $(\mathbb{R}, \max)$  has no identity element.

**Theorem 1.1.1.** *If  $(A, \star)$  is a semigroup, there is at most one identity element.*

*Proof.* Suppose  $e_1, e_2$  are identity elements. Then  $e_1 \star e_2 = e_1$  (because  $e_2$  is an identity element), and  $e_1 \star e_2 = e_2$  (because  $e_1$  is an identity element). Therefore  $e_1 = e_2$ .  $\square$

Consequently, we can talk about “the” identity element, when it exists.

**Definition 1.1.2.** A *monoid* is a triple  $(A, \star, e)$  where  $(A, \star)$  is a semigroup and  $e$  is an identity element.

For example,  $(\mathbb{R}, +, 0)$  and  $(\mathbb{R}, \cdot, 1)$  are monoids.

**Remark 1.1.3.** The traditional definition of “monoid” is “a semigroup in which an identity element exists.” With the traditional definition,  $(\mathbb{R}, +)$  is a monoid, rather than  $(\mathbb{R}, +, 0)$ . However, there is a clear correspondence between the traditional definition and Definition 1.1.2, thanks to Theorem 1.1.1. Definition 1.1.2 works slightly better for the purposes of universal algebra.

Let  $(A, \star, e)$  be a monoid. If  $x \in A$ , then an *inverse* of  $x$  is an element  $x' \in A$  such that  $x \star x' = x' \star x = e$ .

**Example 1.1.4.** In the monoid  $(\mathbb{R}, \cdot, 1)$ , the element  $2/3$  is an inverse of  $3/2$ , while the element  $0$  has no inverse.

**Theorem 1.1.5.** *Let  $(A, \star, e)$  be a monoid. If  $x \in A$ , then  $x$  has at most one inverse.*

*Proof.* Suppose  $y, z$  are both inverses of  $x$ . Then

$$y = y \star e = y \star (x \star z) = (y \star x) \star z = e \star z = z. \quad \square$$

Consequently, we can talk about “the” inverse of  $x$ , when it exists.

**Definition 1.1.6.** A *group* is a 4-tuple  $(A, \star, e, (-)')$  where  $(A, \star, e)$  is a monoid and  $(-)'$  is a unary function  $A \rightarrow A$  such that  $x'$  is an inverse of  $x$  for every  $x \in A$ .

Equivalently, a group is a 4-tuple  $(A, \star, e, (-)')$  where  $A$  is a set,  $\star$  is a binary operation on  $A$ ,  $e \in A$  is a nullary operation (an element), and  $(-)'$  is a unary operation such that the following equations hold for all  $x, y, z \in A$ .

$$\begin{aligned} x \star (y \star z) &= (x \star y) \star z \\ x \star e &= e \star x = x \\ x \star x' &= x' \star x = e. \end{aligned}$$

**Example 1.1.7.**  $(\mathbb{R}, +, 0, -)$  is a group, where  $-$  is the unary negation function  $-x$ .

Again, the traditional definition of “group” is slightly different—a group is a monoid  $(A, \star)$  in which every element has an inverse. With this definition,  $(\mathbb{R}, +)$  is a group rather than  $(\mathbb{R}, +, 0, -)$ . Again, there is a clear correspondence between the traditional definition and Definition 1.1.6, and Definition 1.1.6 is better for universal algebra.

A semigroup, monoid, or group is *commutative* if the equation

$$x \star y = y \star x$$

holds for any  $x, y$ . Commutative groups are usually called *abelian groups*. So far, all our examples have been commutative. Here is an important example of a non-abelian group.

**Example 1.1.8.** Let  $S$  be a set. Let  $\text{Perm}(S)$  be the set of bijections  $f : S \rightarrow S$ . Then  $\text{Perm}(S)$  is a group  $(\text{Perm}(S), \circ, \text{id}, (-)^{-1})$ , where

1.  $f \circ g$  is the function composition  $(f \circ g)(x) := f(g(x))$ .
2.  $\text{id} : S \rightarrow S$  is the identity function  $\text{id}(x) = x$ .
3.  $f^{-1}$  is the inverse function of  $f$ .

$\text{Perm}(S)$  is usually non-abelian, since function composition is non-abelian in general. For example, if  $f, g \in \text{Perm}(\mathbb{R})$  are

$$\begin{aligned} f(x) &= x + 1 \\ g(x) &= 2x \end{aligned}$$

then  $(f \circ g)(x) = 2x + 1$  and  $(g \circ f)(x) = 2(x + 1) = 2x + 2$ , so  $f \circ g \neq g \circ f$ .

When  $S$  is  $\{1, 2, \dots, n\}$ , the group  $\text{Perm}(S)$  is called the  *$n$ th symmetric group*.

**Example 1.1.9.** If you know linear algebra, the group of  $n \times n$  invertible matrices is a group with respect to matrix multiplication. This group is called the  *$n$ th general linear group*, usually written  $GL(n)$  or  $GL_n(\mathbb{R})$ .

**Remark 1.1.10.** Groups are usually written using multiplicative or additive notation.

	Multiplicative notation	Additive notation
$x \star y$	$x \cdot y$	$x + y$
$e$	$1$	$0$
$x'$	$x^{-1}$	$-x$

When using multiplicative notation, one uses the usual abbreviations for multiplication, writing  $xy^{-1}z$  instead of  $x \cdot (y^{-1}) \cdot z$ . Here are the group axioms in multiplicative notation:

$$\begin{aligned} x(yz) &= (xy)z \\ x \cdot 1 &= 1 \cdot x = x \\ xx^{-1} &= x^{-1}x = 1 \end{aligned}$$

Additive notation is traditionally reserved for abelian groups. Here are the abelian group axioms in additive notation:

$$\begin{aligned} x + (y + z) &= (x + y) + z \\ x + 0 &= 0 + x = x \\ x + (-x) &= (-x) + x = 0 \\ x + y &= y + x. \end{aligned}$$

## 1.2 Rings and fields

**Definition 1.2.1.** A *ring* is a 6-tuple  $(A, +, \cdot, -, 0, 1)$  such that

1.  $(A, +, 0, -)$  is an abelian group.
2.  $(A, \cdot, 1)$  is a commutative monoid.
3. The distributive law holds for  $x, y, z \in A$ :

$$x \cdot (y + z) = (x \cdot y) + (x \cdot z).$$

For example,  $(\mathbb{R}, +, \cdot, -, 0, 1)$  is a ring, where  $+, \cdot, -, 0, 1$  have their usual meanings. Similarly,  $\mathbb{Z}, \mathbb{Q}$ , and  $\mathbb{C}$  are rings with respect to the usual operations.

**Definition 1.2.2.** A *field* is a ring  $(K, +, \cdot, -, 0, 1)$  such that  $0 \neq 1$  and any  $x \neq 0$  has an inverse  $x^{-1}$  in the monoid  $(K, \cdot, 1)$ .

For example, the rings  $\mathbb{R}$ ,  $\mathbb{Q}$ , and  $\mathbb{C}$  are fields. On the other hand,  $\mathbb{Z}$  is not, because, for example,  $2 \in \mathbb{Z}$  does not have a multiplicative inverse.

**Remark 1.2.3.** 1. Again, the traditional definition of “ring” is slightly different, so that  $(\mathbb{R}, +, \cdot)$  is a ring rather than  $(\mathbb{R}, +, \cdot, -, 0, 1)$ , but the two definitions are equivalent and Definition 1.2.1 is better for universal algebra.

2. What we are calling “rings” should really be called “commutative rings.” We will follow the conventions of commutative algebra and assume all rings are commutative. In noncommutative rings,  $(A, +)$  is an abelian group but  $(A, \cdot)$  is a monoid, not necessarily commutative. One needs *two* distributive laws

$$\begin{aligned}x \cdot (y + z) &= (x \cdot y) + (x \cdot z) \\(x + y) \cdot z &= (x \cdot z) + (y \cdot z),\end{aligned}$$

as these are no longer equivalent without commutativity. A typical example of a noncommutative ring is the ring  $M_n(\mathbb{R})$  of  $n \times n$  matrices of real numbers, with  $+$  and  $\cdot$  given by matrix addition and matrix multiplication.

3. In some fields of mathematics, rings are not assumed to have a multiplicative identity element  $1 \in A$ , so that  $(A, \cdot)$  is a semigroup rather than a monoid. In this case, what we are calling “rings” should be called “commutative unital rings.”

When working with rings, we use the usual notational abbreviations from algebra. For example,  $x^2y - yz$  means  $((x \cdot x) \cdot y) + (-(y \cdot z))$ .

### 1.3 Languages, algebras, and equational classes

Each of the concepts *magma*, *semigroup*, *monoid*, *group*, *abelian group*, and *ring* is defined by a list of operations and a set of equations. More generally, an *equational class* is a class of “algebras” defined by a set of “equations”. In this section, we make these notions precise.

Before we can define equations and algebras, we need to deal with the fact that there are different kinds of algebras with different signatures. It doesn’t make sense to talk about a semigroup  $(G, \cdot)$  satisfying the distributive law



$x \cdot (y + z) = (x \cdot y) + (x \cdot z)$ , because semigroups don't have an addition operation  $x + y$ . The concept of *languages* allows us to distinguish between these different types of algebras.

**Definition 1.3.1.** A (*functional*) *language*  $\mathcal{L}$  is a set of *function symbols* and a map assigning to each function symbol  $f$  an integer  $n_f \in \mathbb{N}$  called the *arity* of  $f$ . An  *$n$ -ary function symbol* is a function symbol of arity  $n$ . Nullary function symbols are usually called *constant symbols*.

In Chapters 1–2, all “languages” will be functional languages. Later, we will see a more general notion of “language” (Definition 3.1.1). Languages are also called *signatures*, which is probably a better name. But at least in model theory, the term “language” is more common than “signature,” for historical reasons.

**Example 1.3.2.** The *language of abelian groups* has one binary function symbol  $+$ , one constant symbol  $0$ , and one unary function symbol  $-$ .

**Definition 1.3.3.** Given a language  $\mathcal{L}$ , an  $\mathcal{L}$ -*algebra*  $\mathcal{A}$  is a set  $A$  and a map assigning to each  $n$ -ary function symbol  $f$  in  $\mathcal{L}$  a corresponding  $n$ -ary function  $f^{\mathcal{A}} : A^n \rightarrow A$ . The set  $A$  is called the *underlying set* of  $\mathcal{A}$ , and  $f^{\mathcal{A}}$  is called the *interpretation of  $f$  in  $\mathcal{A}$* .

Usually we don't distinguish between an algebra  $\mathcal{A}$  and its underlying set  $A$ , writing both as  $A$ .

**Example 1.3.4.** If  $\mathcal{L}$  is the language of abelian groups, then an  $\mathcal{L}$ -algebra is essentially a 4-tuple  $(A, +^A, 0^A, -^A)$  where  $A$  is a set,  $+^A$  is a binary operation on  $A$ ,  $0^A \in A$ , and  $-^A$  is a unary operation on  $A$ .

Fix some infinite set  $\mathcal{V} = \{x, y, z, \dots\}$  of “variable symbols.”

**Definition 1.3.5.** An  $\mathcal{L}$ -*term* is a string generated by the following rules:

- If  $x$  is a variable symbol, then  $x$  is a term.
- If  $f$  is an  $n$ -ary function symbol in  $\mathcal{L}$ , and  $t_1, \dots, t_n$  are  $\mathcal{L}$ -terms, then  $f(t_1, \dots, t_n)$  is an  $\mathcal{L}$ -term.

**Example 1.3.6.** These are terms in the language of abelian groups:

$$x + (-y), \quad 0 + (x + 0), \quad z, \quad -(-(0 + x)), \quad 0.$$

When we say “let  $t(x_1, \dots, x_n)$  be a term,” we mean that  $t(x_1, \dots, x_n)$  is a term and the variables occurring in  $t(x_1, \dots, x_n)$  are contained in  $\{x_1, \dots, x_n\}$ . If  $s_1, \dots, s_n$  are terms and  $t(x_1, \dots, x_n)$  is a term, then  $t(s_1, \dots, s_n)$  denotes the result of replacing  $x_i$  with  $s_i$  in  $t(x_1, \dots, x_n)$ .

A *closed term* is a term with no variables. If  $t$  is a closed term and  $A$  is an  $\mathcal{L}$ -algebra, we define the *interpretation of  $t$  in  $A$* , written  $t^A$ , recursively as follows:

$$f(t_1, \dots, t_k)^A = f^A(t_1^A, \dots, t_k^A).$$

**Example 1.3.7.** If  $\mathcal{L}$  is the language of abelian groups, then the interpretation of the closed  $\mathcal{L}$ -term  $(0 + (-0)) + 0$  in an  $\mathcal{L}$ -algebra  $(A, +^A, 0^A, -^A)$  is the value  $(0^A +^A (-^A 0^A)) +^A 0^A$ .

The language  $\mathcal{L}(A)$  consists of  $\mathcal{L}$  with each element of  $A$  added as a new constant symbol. We regard  $A$  as an  $\mathcal{L}(A)$ -algebra by interpreting each new constant symbol as the corresponding element of  $A$ , so that  $c^A = c$  for  $c \in A$ . If  $t(x_1, \dots, x_n)$  is an  $\mathcal{L}$ -term and  $a_1, \dots, a_n \in A$ , then  $t(a_1, \dots, a_n)$  is a closed  $\mathcal{L}(A)$ -term. The *interpretation of  $t$  in  $A$* , written  $t^A$ , is the function  $t^A : A^n \rightarrow A$  defined by

$$t^A(a_1, \dots, a_n) = (t(a_1, \dots, a_n))^A.$$

**Example 1.3.8.** If  $\mathcal{L}$  is the language of abelian groups and  $t(x, y) = (-x) + (0 + y)$ , then the interpretation of  $t$  in an  $\mathcal{L}$ -algebra  $(A, +^A, 0^A, -^A)$  is the function

$$t^A(a, b) = ((-a) + (0 + b))^A = (-^A a) +^A (0^A +^A b).$$

**Definition 1.3.9.** An  $\mathcal{L}$ -*equation* is a formal expression of the form

$$t(x_1, \dots, x_n) = s(x_1, \dots, x_n)$$

for two  $\mathcal{L}$ -terms  $t(\bar{x})$  and  $s(\bar{x})$ . An  $\mathcal{L}$ -algebra  $A$  *satisfies* an equation  $(t(\bar{x}) = s(\bar{x}))$  if for any  $\bar{a} \in A^n$ ,

$$t^A(\bar{a}) = s^A(\bar{a}).$$

The notation  $A \models \varphi$  means that  $A$  satisfies  $\varphi$ .

The group  $(\mathbb{R}, +)$  satisfies the equation  $x \cdot y = y \cdot x$ , but the non-abelian group  $(\text{Perm}(S), \circ)$  does not:

$$\begin{aligned} (\mathbb{R}, +) &\models x \cdot y = y \cdot x \\ (\text{Perm}(S), \circ) &\not\models x \cdot y = y \cdot x. \end{aligned}$$

**Definition 1.3.10.** An *equational  $\mathcal{L}$ -theory* is a set  $\Sigma$  of  $\mathcal{L}$ -equations. Elements of  $\Sigma$  are called *axioms* of  $\Sigma$ . If  $\Sigma$  is an equational  $\mathcal{L}$ -theory, and  $A$  is an  $\mathcal{L}$ -algebra, then  $A$  is a *model* of  $\Sigma$ , written  $A \models \Sigma$ , if  $A \models \varphi$  for every  $\varphi \in \Sigma$ . The class of models of  $\Sigma$  is written  $\text{Mod}(\Sigma)$ . An *equational class* is a class of the form  $\text{Mod}(\Sigma)$  for some  $\Sigma$ .

**Example 1.3.11.** Let  $\mathcal{L}$  be the language of abelian groups. The *theory of abelian groups* consists of the equations

$$\begin{aligned} x + (y + z) &= (x + y) + z \\ x + 0 &= x \\ x + (-x) &= 0 \\ x + y &= y + x. \end{aligned}$$

Models are abelian groups.

**Example 1.3.12.** The classes of rings, groups, monoids, semigroups, and magmas are equational classes.

**Warning 1.3.13.** The class of fields is *not* an equational class. We will see a couple proofs of this in the next chapter (Examples 2.1.3 and 2.2.4). Nevertheless, we can still learn useful facts about fields by applying the tools of universal algebra to rings.

**Example 1.3.14.** An *idempotent monoid* is a monoid  $(A, \star, e)$  in which the equation  $x \star x = x$  holds. Idempotent monoids form an equational class.

**Example 1.3.15.** A *boolean algebra* is a 6-tuple  $(B, \wedge, \vee, \neg, 0, 1)$  where  $(B, \vee, 0)$  is an idempotent commutative monoid,  $(B, \wedge, 1)$  is an idempotent

commutative monoid, and  $\neg : B \rightarrow B$  is a unary operation such that the following equations hold:

$$\begin{aligned} x \wedge \neg x &= 0 \\ x \vee \neg x &= 1 \\ x \wedge (y \vee z) &= (x \wedge y) \vee (x \wedge z) \\ x \vee (y \wedge z) &= (x \vee y) \wedge (x \vee z) \end{aligned}$$

Boolean algebras form an equational class. If  $S$  is a set, then the powerset  $\mathfrak{P}(S)$  is a boolean algebra with

$$\begin{aligned} x \wedge y &= x \cap y \\ x \vee y &= x \cup y \\ 0 &= \emptyset \\ 1 &= S \\ \neg x &= S \setminus x \end{aligned}$$

Another important boolean algebra is the set of truth values  $\{\text{FALSE}, \text{TRUE}\}$ , with  $\wedge, \vee, \neg$  interpreted as the basic logical operations:

$$\begin{aligned} x \wedge y &= x \text{ AND } y \\ x \vee y &= x \text{ OR } y \\ \neg x &= \text{NOT } x \\ 0 &= \text{FALSE} \\ 1 &= \text{TRUE}. \end{aligned}$$

**Warning 1.3.16.** In universal algebra, equational classes are usually called *varieties*. We will avoid this terminology because “variety” tends to mean something completely different in model theory. Confusingly, “equational theory” also has a technical meaning in model theory, though it is much less common.

Note that languages are allowed to have infinitely many symbols, and theories are allowed to have infinitely many axioms. Here is an important example:

**Definition 1.3.17.** Let  $(G, \cdot)$  be a group and  $S$  be a set. An *action* of  $G$  on  $S$  is a map

$$\star : G \times S \rightarrow S$$

satisfying the following axioms for  $g, h \in G$  and  $x \in S$ :

$$\begin{aligned} (g \cdot h) \star x &= g \star (h \star x) \\ 1 \star x &= x \end{aligned}$$

Group actions are usually written using  $\cdot$  rather than  $\star$ .

**Example 1.3.18.** Suppose  $S$  is a set and  $\text{Perm}(S)$  is the group of permutations on  $S$  (Example 1.1.8). There is a natural action of  $\text{Perm}(S)$  on  $S$  given by  $f \cdot x = f(x)$ . The axioms of group actions hold as follows:

$$\begin{aligned} (f \cdot g) \cdot x &= (f \circ g)(x) = f(g(x)) = f \cdot (g \cdot x) \\ 1 \cdot x &= \text{id}(x) = x. \end{aligned}$$

**Example 1.3.19.** If you know linear algebra, another example is the group of invertible  $n \times n$  matrices, which acts by matrix multiplication on the space of  $n \times 1$  column vectors.

Fix a specific group  $G$ .

**Definition 1.3.20.** A  $G$ -set is a pair  $(S, \cdot)$  where  $S$  and  $\cdot$  is an action of  $G$  on  $S$ .

This doesn't immediately fit into the framework of universal algebra, because the binary operator  $\cdot$  involves two different sets, one of which is fixed. However, the following equivalent definition works:

**Definition 1.3.21.** A  $G$ -set is a set  $S$  and a unary operator  $\mu_g$  for each  $g \in G$ , satisfying the following identities:

$$\begin{aligned} \mu_{g \cdot h}(x) &= \mu_g(\mu_h(x)) \text{ for each } g, h \in G \\ \mu_1(x) &= x \end{aligned}$$

The translation between Definitions 1.3.20 and 1.3.21 is

$$\mu_g(x) = g \cdot x.$$

Definition 1.3.21 fits into the framework of universal algebra. Note that the language of  $G$ -sets has one symbol for each element of  $G$ . When  $G$  is an infinite group, we need infinitely many symbols. Likewise, the theory of  $G$ -sets needs infinitely many equations (when  $G$  is infinite).

**Remark 1.3.22.** If you know what vector spaces are, the same thing happens: for any fixed field  $K$ , the class of  $K$ -vector spaces is an equational class. We will review vector spaces later, in Section 13.6.

## 1.4 Homomorphisms and isomorphisms

Fix a language  $\mathcal{L}$ .

**Definition 1.4.1.** Let  $A, B$  be  $\mathcal{L}$ -algebras. A *homomorphism* from  $A$  to  $B$  is a function  $\alpha : A \rightarrow B$  such that for any  $k$ -ary function symbol  $f \in \mathcal{L}$ , and any  $a_1, \dots, a_k \in A$ ,

$$\alpha(f^A(a_1, \dots, a_k)) = f^B(\alpha(a_1), \alpha(a_2), \dots, \alpha(a_k)). \quad (*)$$

An *isomorphism* is a bijective homomorphism. Two  $\mathcal{L}$ -algebras  $A, B$  are *isomorphic*, written  $A \cong B$ , if there is an isomorphism  $\alpha : A \rightarrow B$ .

In what follows, we extend functions to tuples componentwise, so that  $\alpha(a_1, \dots, a_k) := (\alpha(a_1), \dots, \alpha(a_k))$ . Then  $(*)$  can be rewritten as  $\alpha(f^A(\bar{a})) = f^B(\alpha(\bar{a}))$ .

**Remark 1.4.2.** The idea of  $(*)$  is that  $\alpha$  “preserves” the function symbol  $f$ . It may help to rewrite  $(*)$  as the following logically equivalent condition:

$$f^A(a_1, \dots, a_k) = b \implies f^B(\alpha(a_1), \dots, \alpha(a_k)) = \alpha(b).$$

**Example 1.4.3.** Let  $\mathcal{L} = \{+, 0, -\}$  be the language of abelian groups. Let  $h : \mathbb{Z} \rightarrow \mathbb{Z}$  be  $h(x) = 2x$ . Then  $h$  is a homomorphism from  $\mathbb{Z}$  to  $\mathbb{Z}$ , because

$$\begin{aligned} 2(x + y) &= (2x) + (2y) \\ 2(0) &= 0 \\ 2(-x) &= -(2x). \end{aligned}$$

However,  $h$  is not an isomorphism, because  $h$  is not surjective.

Here is another example which is slightly more subtle.

**Example 1.4.4.** Let  $\mathcal{L} = \{\cdot, 1\}$  be the language of monoids. The exponential map  $\exp(x) = e^x$  is a homomorphism from the monoid  $(\mathbb{R}, +, 0)$  to  $(\mathbb{R}, \cdot, 1)$ , because

$$\begin{aligned}\exp(x + y) &= \exp(x) \exp(y) \\ \exp(0) &= 1\end{aligned}$$

If this looks confusing, remember that the interpretation of “ $\cdot$ ” in the monoid  $(\mathbb{R}, +, 0)$  is  $+$ , and the interpretation of “ $1$ ” is  $0$ .

The exponential map is not an isomorphism, as it is not surjective. On the other hand, if  $\mathbb{R}_{>0}$  denotes the set of positive real numbers, then  $\exp$  is an isomorphism from the monoid  $(\mathbb{R}, +, 0)$  to the monoid  $(\mathbb{R}_{>0}, \cdot, 1)$ .

**Example 1.4.5.** An action of  $G$  on a set  $S$  (Definition 1.3.17) can be thought of as a map assigning to each element  $g \in G$  a permutation  $\mu_g \in \text{Perm}(S)$  satisfying the conditions

$$\begin{aligned}\mu_{g \cdot h} &= \mu_g \circ \mu_h \\ \mu_1 &= \text{id}_S\end{aligned}$$

(See Definition 1.3.21.) These conditions precisely say that  $g \mapsto \mu_g$  is a monoid homomorphism  $(G, \cdot, 1) \rightarrow (\text{Perm}(S), \circ, \text{id}_S)$ . We will see later that such a homomorphism must actually be a group homomorphism (Theorem 1.4.13). Therefore, an action of  $G$  on  $S$  is equivalent to a group homomorphism from  $G$  to  $\text{Perm}(S)$ .

The next theorem expresses some basic facts about homomorphisms and isomorphisms: we can compose homomorphisms and take inverses of isomorphisms.

**Theorem 1.4.6.** *Let  $\mathcal{L}$  be a language and  $A, B, C$  be  $\mathcal{L}$ -algebras.*

1. *Let  $\alpha : A \rightarrow B$  and  $\beta : B \rightarrow C$  be homomorphisms. Then  $\beta \circ \alpha : A \rightarrow C$  is a homomorphism.*
2. *The identity map  $\text{id}_A : A \rightarrow A$  is an isomorphism.*
3. *If  $\alpha : A \rightarrow B$  is an isomorphism, then  $\alpha^{-1} : B \rightarrow A$  is an isomorphism.*

*Proof.* 1. If  $f \in \mathcal{L}$  is a  $k$ -ary function symbol, then

$$\beta(\alpha(f^A(\bar{a}))) = \beta(f^B(\alpha(\bar{a}))) = f^C(\beta(\alpha(\bar{a})))$$

because  $\beta$  and  $\alpha$  are homomorphisms.

2. If  $f \in \mathcal{L}$  is a  $k$ -ary relation symbol, then

$$\text{id}(f(\bar{a})) = f(\bar{a}) = f(\text{id}(\bar{a})).$$

3. Suppose  $f \in \mathcal{L}$  is a  $k$ -ary function symbol and  $b_1, \dots, b_k \in B$ . Let  $a_i = \alpha^{-1}(b_i) \in A$ . Then  $b_i = \alpha(a_i)$ . As  $\alpha$  is a homomorphism,

$$\alpha(f^A(\bar{a})) = f^B(\alpha(\bar{a})) = f^B(\bar{b}).$$

Applying  $\alpha^{-1}$  to both sides, we see that

$$\alpha^{-1}(f^B(\bar{b})) = f^A(\bar{a}) = f^A(\alpha^{-1}(\bar{b})). \quad \square$$

**Corollary 1.4.7.** *The relation of isomorphism is an equivalence relation on  $\mathcal{L}$ -algebras:*

1.  $A \cong A$  for any  $\mathcal{L}$ -algebra  $A$ .
2. If  $A \cong B$ , then  $B \cong A$ .
3. If  $A \cong B$  and  $B \cong C$ , then  $A \cong C$ .

*Proof.* 1.  $\text{id}_A : A \rightarrow A$  is an isomorphism.

2. If  $\alpha : A \rightarrow B$  is an isomorphism, then  $\alpha^{-1} : B \rightarrow A$  is an isomorphism.

3. If  $\alpha : A \rightarrow B$  and  $\beta : B \rightarrow C$  are isomorphisms, then  $\beta \circ \alpha : A \rightarrow C$  are isomorphisms.  $\square$

**Example 1.4.8.** Here is an example of an isomorphism. Let  $S$  be  $\{\{x\} : x \in \mathbb{R}\}$ , the set of one element subsets of  $\mathbb{R}$ . Define operations on  $S$  as follows:

$$\begin{aligned} \{x\} + \{y\} &:= \{x + y\} \\ \{x\} \cdot \{y\} &:= \{x \cdot y\} \\ -\{x\} &= \{-x\} \\ 1 &:= \{1\} \\ 0 &:= \{0\}. \end{aligned}$$



For example,  $\{2\} + \{3\} \cdot \{5\} = \{17\}$ . This makes  $(S, +, \cdot, -, 0, 1)$  into a ring. Then there is an isomorphism from the ring  $\mathbb{R}$  to the ring  $S$  given by

$$\begin{aligned} f : \mathbb{R} &\rightarrow S \\ f(x) &= \{x\}. \end{aligned}$$

The intuition you should have is that the ring  $S$  is a poorly disguised copy of  $\mathbb{R}$ . On some level,  $S$  is “the same thing” as  $\mathbb{R}$ .

This is the intuition we have for isomorphisms in general—if two algebras  $A$  and  $B$  are isomorphic, then we think of them as being two “copies” of the same algebra. In algebra and model theory, the objects of study are not so much algebras as isomorphism classes of algebras.

This mindset has the following corollary: if two algebras  $A$  and  $B$  are isomorphic, then they should have identical properties. For example,  $A$  and  $B$  will have the same cardinality because the isomorphism  $f : A \rightarrow B$  is a bijection, showing that  $|A| = |B|$ .

Of course, not all properties respect isomorphisms. In our example above, the property “contains 7” is true for  $\mathbb{R}$  but false for  $S$ . What we should conclude from this is that the property “contains 7” is a stupid property, or an *evil property* as they would say on the website nLab. The “meaningful” or “good” properties of algebras are the ones that respect isomorphisms. In the remainder of this section, we show that the properties defined by equations are good properties. In other words, we will show that isomorphic algebras satisfy the same equations (Theorem 1.4.11).

The next theorem shows that homomorphisms preserve not only the basic function symbols, but also all terms. This is intuitively reasonable, since terms are expressions built up from function symbols.

**Theorem 1.4.9.** *Let  $\alpha : A \rightarrow B$  be a homomorphism of  $\mathcal{L}$ -algebras. Let  $t(x_1, \dots, x_n)$  be an  $\mathcal{L}$ -term. Then for any  $a_1, \dots, a_n \in A$ , we have*

$$\alpha(t^A(\bar{a})) = t^B(\alpha(\bar{a})). \quad (\dagger)$$

*Proof.* Proceed by induction on the complexity of  $t$ .

- If  $t(\bar{x}) = x_i$ , then both sides of  $(\dagger)$  are  $\alpha(a_i)$ .
- Suppose  $t(\bar{x}) = f(s_1(\bar{x}), \dots, s_k(\bar{x}))$  for some  $k$ -ary function symbol  $f$  and some simpler  $\mathcal{L}$ -terms  $s_1, \dots, s_k$ . By definition of  $t^A$ ,

$$\alpha(t^A(\bar{a})) = \alpha(f^A(s_1^A(\bar{a}), \dots, s_k^A(\bar{a}))).$$

As  $\alpha$  is a homomorphism,

$$\alpha(f^A(s_1^A(\bar{a}), \dots, s_k^A(\bar{a}))) = f^B(\alpha(s_1^A(\bar{a})), \dots, \alpha(s_k^A(\bar{a}))).$$

By induction,

$$\alpha(s_i^A(\bar{a})) = s_i^B(\alpha(\bar{a})).$$

for each  $i$ . Therefore,

$$f^B(\alpha(s_1^A(\bar{a})), \dots, \alpha(s_k^A(\bar{a}))) = f^B(s_1^B(\alpha(\bar{a})), \dots, s_k^B(\alpha(\bar{a}))).$$

By definition, the right hand side is  $t^B(\alpha(\bar{a}))$ .  $\square$

**Lemma 1.4.10.** *Let  $\alpha : A \rightarrow B$  be a surjective homomorphism of  $\mathcal{L}$ -algebras, and let  $\varphi$  be an  $\mathcal{L}$ -equation. Then  $A \models \varphi \implies B \models \varphi$ .*

*Proof.* Suppose  $\varphi$  is  $t(x_1, \dots, x_n) = s(x_1, \dots, x_n)$ . Suppose  $b_1, \dots, b_n \in B$ . By surjectivity, we can write  $b_i$  as  $\alpha(a_i)$  for some  $a_i \in A$ . Then

$$\begin{aligned} t^B(\bar{b}) &= t^B(\alpha(\bar{a})) = \alpha(t^A(\bar{a})) \\ s^B(\bar{b}) &= s^B(\alpha(\bar{a})) = \alpha(s^A(\bar{a})) \end{aligned}$$

by Theorem 1.4.9, as  $\alpha$  is a homomorphism. Because  $A \models \varphi$ , the right hand sides are equal. Therefore the left hand sides are equal, meaning

$$t^B(\bar{b}) = s^B(\bar{b}).$$
 $\square$

This allows us to complete the proof that isomorphic algebras satisfy the same equations.

**Theorem 1.4.11.** *Suppose  $A \cong B$ .*

1.  $A \models s = t \iff B \models s = t$ .
2.  $A \models \Sigma \iff B \models \Sigma$
3.  $A \in \text{Mod}(\Sigma) \iff B \in \text{Mod}(\Sigma)$ .
4. If  $\mathcal{K}$  is an equational class, then  $A \in \mathcal{K} \iff B \in \mathcal{K}$ .

## Group homomorphisms

In this section, we study homomorphisms of groups, learning something useful about rings and fields along the way.

**Lemma 1.4.12** (Cancellation). *Let  $G$  be a group. If  $ax = ay$ , then  $x = y$ . Similarly, if  $xa = ya$ , then  $x = y$ .*

*Proof.* If  $ax = ay$ , then

$$x = 1x = (a^{-1}a)x = a^{-1}(ax) = a^{-1}(ay) = (a^{-1}a)y = 1y = y.$$

The other case is similar. □

**Theorem 1.4.13.** *Let  $G, H$  be groups. If  $f : G \rightarrow H$  is a semigroup homomorphism, meaning that  $f$  preserves multiplication*

$$f(xy) = f(x)f(y),$$

*then  $f$  also preserves 1 and  $(-)^{-1}$ :*

$$\begin{aligned} f(1) &= 1 \\ f(x^{-1}) &= f(x)^{-1}, \end{aligned}$$

*and so  $f$  is a group homomorphism.*

*Proof.* First note that

$$f(1) \cdot f(1) = f(1 \cdot 1) = f(1) = f(1) \cdot 1.$$

Canceling  $f(1)$  from both sides, we see  $f(1) = 1$ . Next,

$$f(x)f(x^{-1}) = f(xx^{-1}) = f(1) = 1 = f(x)f(x)^{-1}.$$

Canceling  $f(x)$  from both sides, we see  $f(x^{-1}) = f(x)^{-1}$ . □

**Corollary 1.4.14.** *Let  $R$  be a ring. Then the following equations hold, for  $a, x \in R$ :*

$$\begin{aligned} 0a &= 0 \\ (-x)a &= -(xa) \\ (-1)a &= -a \end{aligned}$$

*Proof.* Let  $\mu_a(x) = ax$ . Then  $\mu_a$  is a semigroup homomorphism  $(R, +) \rightarrow (R, +)$ , by the distributive law:

$$a(x + y) = ax + ay.$$

Thus  $\mu_a$  preserves zero and negation:

$$\begin{aligned}\mu_a(0) &= 0 \\ \mu_a(-x) &= -\mu_a(x),\end{aligned}$$

which are the first two equations. The third equation holds by taking  $x = 1$  in the second equation.  $\square$

**Lemma 1.4.15.** *The following holds in any field  $K$ :*

$$xy = xz \implies y = z \text{ when } x \neq 0.$$

*Proof.* Multiply both sides of  $xy = xz$  by  $x^{-1}$ , as in Lemma 1.4.12.  $\square$

**Theorem 1.4.16.** *If  $R$  is a field and  $x, y \in R$ , then*

$$xy = 0 \iff (x = 0 \text{ or } y = 0)$$

*Proof.* If  $x = 0$  or  $y = 0$ , then  $xy = 0$  by the zero law (Corollary 1.4.14). Conversely, suppose  $xy = 0$ , but  $x \neq 0$  and  $y \neq 0$ . Then

$$xy = 0 = x0$$

by the zero law. Cancelling  $x$  from both sides,  $y = 0$ , a contradiction.  $\square$

# Chapter 2

## New algebras from old

### 2.1 Subalgebras and generators

**Definition 2.1.1.** Let  $A$  be an  $\mathcal{L}$ -algebra. A *subalgebra* is a subset  $B \subseteq A$  such that for any  $k$ -ary relation symbol  $f \in \mathcal{L}$ ,

$$b_1, \dots, b_k \in B \implies f^A(b_1, \dots, b_k) \in B.$$

If  $B$  is a subalgebra of an  $\mathcal{L}$ -algebra  $A$ , then we can make  $B$  into an  $\mathcal{L}$ -algebra by defining  $f^B$  to be the restriction of  $f^A$  to  $B$ , for each function symbol  $f$ :

$$f^B(b_1, \dots, b_k) := f^A(b_1, \dots, b_k) \in B.$$

In this way, we regard subalgebras as algebras, not just sets.

**Theorem 2.1.2.** Suppose  $B$  is a subalgebra of  $A$ .

1. The inclusion  $B \rightarrow A$  is a homomorphism.
2. If  $A \models s = t$ , then  $B \models s = t$ .
3. If  $\mathcal{K} = \text{Mod}(\Sigma)$  is an equational class, then  $A \in \mathcal{K} \implies B \in \mathcal{K}$ .

*Proof.* (1) holds by choice of the structure on  $B$ . For (2), note that

$$\begin{aligned} t^B(\bar{b}) &= t^A(\bar{b}) \\ s^B(\bar{b}) &= s^A(\bar{b}) \end{aligned}$$

for  $\bar{b}$  in  $B$  by Theorem 1.4.9 and (1). If  $t^A = s^A$ , then  $t^B = s^B$ . Finally, (3) follows directly from (2).  $\square$

**Example 2.1.3.** The ring  $\mathbb{R}$  is a field, but the subring  $\mathbb{Z}$  is not. Therefore, fields are not an equational class.

**Definition 2.1.4.** If  $S$  is a subset of an algebra  $A$ , then  $\langle S \rangle$  or  $\langle S \rangle_A$  denotes the smallest subalgebra of  $A$  containing  $S$ , which is

$$\{t^A(\bar{b}) : t(x_1, \dots, x_n) \text{ is an } \mathcal{L}\text{-term and } \bar{b} \in S^n\}$$

The subalgebra  $\langle S \rangle$  is called the subalgebra *generated* by  $S$ . We say that  $A$  is *finitely generated* if  $A = \langle S \rangle$  for some finite  $S \subseteq A$ .

We often omit brackets  $\{, \}$  inside  $\langle, \rangle$ , using abbreviations like

$$\begin{aligned} \langle a_1, \dots, a_n \rangle &= \langle \{a_1, \dots, a_n\} \rangle \\ \langle A, b \rangle &= \langle A \cup \{b\} \rangle. \end{aligned}$$

## 2.2 Products

### Binary products

**Definition 2.2.1.** Let  $A, B$  be two  $\mathcal{L}$ -algebras. The *product algebra*  $A \times B$  is the  $\mathcal{L}$ -algebra with underlying set  $A \times B$  and

$$f^{A \times B}((a_1, b_1), \dots, (a_k, b_k)) := (f^A(a_1, \dots, a_k), f^B(b_1, \dots, b_k)).$$

for each  $k$ -ary function symbol  $f \in \mathcal{L}$ .

**Remark 2.2.2.** Let  $A_1, A_2$  be two  $\mathcal{L}$ -algebras. For  $i = 1, 2$ , let  $\pi_i : A_1 \times A_2 \rightarrow A_i$  be the projection map  $\pi_i(x_1, x_2) = x_i$ . Then each  $\pi_i$  is a homomorphism. For example,

$$\begin{aligned} \pi_1(f^{A_1 \times A_2}((a_1, b_1), \dots, (a_k, b_k))) &= f^{A_1}(a_1, \dots, a_k) \\ &= f^{A_1}(\pi_1(a_1, b_1), \dots, \pi_1(a_k, b_k)). \end{aligned}$$

**Theorem 2.2.3.** Let  $A_1, A_2$  be  $\mathcal{L}$ -algebras.

1. If  $\varphi$  is an equation and  $A_i \models \varphi$  for  $i = 1, 2$ , then  $A_1 \times A_2 \models \varphi$ .
2. If  $\mathcal{K}$  is an equational class and  $A_1, A_2 \in \mathcal{K}$ , then  $A_1 \times A_2 \in \mathcal{K}$ .

*Proof.* Part (2) follows formally from (1). For (1), suppose  $A_i \models t = s$  for  $i = 1, 2$ . Let  $P = A_1 \times A_2$ . We claim  $P \models t = s$ . Otherwise there are  $a_1, \dots, a_k \in P$  such that  $t^P(\bar{a}) \neq s^P(\bar{a})$ . Then there is  $i \in \{1, 2\}$  such that

$$\pi_i(t^P(\bar{a})) \neq \pi_i(s^P(\bar{a})).$$

As  $\pi_i$  is a homomorphism, we can rewrite the two sides as follows:

$$t^{A_i}(\pi_i(\bar{a})) \neq s^{A_i}(\pi_i(\bar{a})).$$

Then  $A_i \not\models t = s$ , a contradiction.  $\square$

**Example 2.2.4.** The ring  $\mathbb{R}$  is a field, but  $\mathbb{R} \times \mathbb{R}$  is not, because the zero law (Theorem 1.4.16) fails:

$$\begin{aligned} (0, 1) &\neq (0, 0) \\ (1, 0) &\neq (0, 0) \\ (0, 1) \cdot (1, 0) &= (0, 0). \end{aligned}$$

Therefore fields are not an equational class.

## Infinite products

Let  $I$  be a set.

**Definition 2.2.5.** An  $I$ -tuple is a function with domain  $I$ . If  $a$  is an  $I$ -tuple, we write  $a(i)$  as  $\pi_i(a)$ . The notation  $(a_i : i \in I)$  means the function  $i \mapsto a_i$ .

When  $I = \{1, 2, \dots, n\}$ , we identify  $n$ -tuples and  $I$ -tuples, so that

$$(a_1, a_2, \dots, a_n) = (a_i : i \in \{1, \dots, n\}).$$

Likewise, we think of  $\omega$ -tuples as tuples of length  $\omega$ , i.e., sequences, so that

$$(a_1, a_2, a_3, \dots) = (a_i : i \in \omega).$$

**Definition 2.2.6.** Let  $\{A_i\}_{i \in I}$  be a family of sets. The *direct product*  $\prod_{i \in I} A_i$  is the set of  $I$ -tuples  $(a_i : i \in I)$  such that  $a_i \in A_i$  for each  $i \in I$ .

**Remark 2.2.7.** Suppose  $A_i$  doesn't depend on  $i$ , so that  $A_i = A$  for some fixed set  $A$  as  $i$  varies. Then  $\prod_{i \in I} A$  is the set of functions from  $I$  to  $A$ . This set is also written  $A^I$ , and is called a *power* of  $A$ .

**Definition 2.2.8.** Let  $I$  be a set and let  $A_i$  be an  $\mathcal{L}$ -algebra for each  $i \in I$ . We make  $\prod_{i \in I} A_i$  into an  $\mathcal{L}$ -algebra by interpreting each  $k$ -ary function symbol  $f$  as

$$f(a_1, \dots, a_k) = (f^{A_i}(\pi_i(a_1), \dots, \pi_i(a_k)) : i \in I).$$

**Remark 2.2.9.** The structure on  $\prod_{i \in I} A_i$  is chosen to make  $\pi_i$  a homomorphism:

$$\pi_i(f(a_1, \dots, a_k)) = f^{A_i}(\pi_i(a_1), \dots, \pi_i(a_k)).$$

**Theorem 2.2.10.** Let  $A_i$  be an  $\mathcal{L}$ -algebra for each  $i \in I$ .

1. If  $\varphi$  is an equation and  $A_i \models \varphi$  for each  $i \in I$ , then  $\prod_{i \in I} A_i \models \varphi$ .
2. If  $\mathcal{K}$  is an equational class and  $A_i \in \mathcal{K}$  for all  $i \in I$ , then  $\prod_{i \in I} A_i \in \mathcal{K}$ .

*Proof.* The proof of Theorem 2.2.3 works here.  $\square$

**Example 2.2.11** (Powers). If  $I$  is a set and  $A$  is an  $\mathcal{L}$ -algebra, the power  $A^I := \prod_{i \in I} A$  is the set of functions  $I \rightarrow A$ , with all the operations defined pointwise.

## 2.3 Descending functions along surjections

**Theorem 2.3.1.** Let  $\pi : A \rightarrow A'$  be a surjection. Let  $f : A \rightarrow B$  be a function such that

$$\pi(x) = \pi(y) \implies f(x) = f(y). \quad (*)$$

Then there is a unique function  $f' : A' \rightarrow B$  such that  $f'(\pi(x)) = f(x)$ , i.e., the following diagram commutes:

$$\begin{array}{ccc} A & & \\ \pi \downarrow & \searrow f & \\ A' & \xrightarrow{f'} & B. \end{array}$$



*Proof.* For  $z \in A'$ , let  $S_z = \{f(x) : x \in A, \pi(x) = z\}$ . There is at least one  $x \in A$  with  $\pi(x) = z$  because  $\pi$  is surjective, so  $S_z$  is non-empty. If  $x_1, x_2 \in A$  satisfy  $\pi(x_i) = z$  for  $i = 1, 2$ , then  $f(x_1) = f(x_2)$  by (\*). Therefore  $S_z$  has a unique element. Let  $f'(z)$  be the unique element of  $S_z$ .

If  $x \in A$  and  $z = \pi(x)$ , then  $f(x) \in S_z = \{f'(z)\}$ , so  $f(x) = f'(z) = f'(\pi(x))$ . This proves existence of  $f'$ . For uniqueness, suppose  $f'' : A' \rightarrow B$  is another function satisfying  $f''(\pi(x)) = f(x)$ . For any  $z \in A'$ , there is  $x \in A$  with  $\pi(x) = z$ , and then

$$f''(z) = f''(\pi(x)) = f(x) = f'(\pi(x)) = f'(z),$$

so  $f'' = f'$ . □

**Example 2.3.2.** Let  $\mathcal{K}$  be the class of  $\mathcal{L}$ -algebras. If  $X, Y \in \mathcal{K}$  are isomorphic, then  $|X| = |Y|$ . Therefore, there is a map from the class  $\mathcal{K}/\cong$  of isomorphism classes to the class Card of cardinals sending the isomorphism class of  $X$  to the cardinality  $|X|$ . Ignoring the difference between sets and classes, this is an instance of Theorem 2.3.1:

$$\begin{array}{ccc} \mathcal{K} & & \\ \downarrow & \searrow & \\ \mathcal{K}/\cong & \dashrightarrow & \text{Card.} \end{array}$$

The surjectivity of  $\mathcal{K} \rightarrow \mathcal{K}/\cong$  holds because every isomorphism class is the isomorphism class of some algebra. The condition (\*) in Theorem 2.3.1 holds because

$$[X]_{\cong} = [Y]_{\cong} \iff X \cong Y \implies |X| = |Y|.$$

The conclusion of Theorem 2.3.1 says that the map  $f' : \mathcal{K}/\cong \rightarrow \text{Card}$  satisfies

$$f'([X]_{\cong}) = |X|,$$

so that  $f'$  sends the isomorphism class of  $X$  to the cardinality of  $X$ .

## 2.4 Congruences and quotients

**Definition 2.4.1.** Let  $A$  be an  $\mathcal{L}$ -algebra. A *congruence* on  $A$  is an equivalence relation  $E$  on  $A$  that is also a subalgebra of  $A \times A$ .

Unwinding the definition, an equivalence relation  $\sim$  on  $A$  is a congruence iff

$$(a_1 \sim b_1 \text{ and } a_2 \sim b_2 \text{ and } \dots \text{ and } a_k \sim b_k) \implies f(a_1, \dots, a_k) \sim f(b_1, \dots, b_k)$$

for any  $k$ -ary function symbol  $f \in \mathcal{L}$ .

**Theorem 2.4.2.** *Let  $\sim$  be an equivalence relation on an  $\mathcal{L}$ -algebra  $A$ . Then  $\sim$  is a congruence iff the following holds: for any  $k$ -ary function symbol  $f \in \mathcal{L}$  and any  $1 \leq i \leq k$ ,*

$$a_i \sim a'_i \implies f(a_1, \dots, a_k) \sim f(a_1, \dots, a_{i-1}, a'_i, a_{i+1}, \dots, a_k). \quad (*)$$

*Proof.* First suppose  $\sim$  is a congruence, and  $a_i \sim a'_i$ . For  $j \neq i$  define  $a'_j := a_j$ . Then  $a_j \sim a'_j$  because  $\sim$  is reflexive. As  $a_j \sim a'_j$  for all  $j \leq k$ , we see that

$$f(a_1, \dots, a_k) \sim f(a'_1, \dots, a'_k) = f(a_1, \dots, a_{i-1}, a'_i, a_{i+1}, \dots, a_k).$$

Conversely, suppose  $(*)$  holds. Suppose  $a_i \sim b_i$  for  $i = 1, \dots, k$ . Then  $(*)$  gives

$$\begin{aligned} & f(a_1, a_2, a_3, \dots, a_k) \\ & \sim f(b_1, a_2, a_3, \dots, a_k) \\ & \sim f(b_1, b_2, a_3, \dots, a_k) \\ & \sim \dots \\ & \sim f(b_1, b_2, \dots, b_k). \end{aligned} \quad \square$$

**Definition 2.4.3.** Let  $R$  be a ring. An *ideal* is a subset  $I \subseteq R$  such that

1.  $0 \in I$ .
2.  $x, y \in I \implies x + y \in I$ .
3.  $(x \in R \text{ and } y \in I) \implies xy \in I$ .

**Remark 2.4.4.** If  $R$  is a ring and  $a \in R$ , then the set  $aR := \{ax : x \in R\}$  is an ideal. Such ideals are called *principal ideals*.

**Theorem 2.4.5.** *Let  $R$  be a ring.*

1. If  $I$  is an ideal, define  $x \equiv_I y$  to mean  $x - y \in I$ . Then  $\equiv_I$  is a congruence.
2. This gives a bijection between congruences on  $R$  and ideals on  $R$ .

*Proof.* First, note that if  $\sim$  is a congruence, then

$$x \sim y \iff x - y \sim 0$$

for any  $x, y \in R$ . Indeed,

$$\begin{aligned} x \sim y &\implies x - y \sim y - y = 0 \\ x - y \sim 0 &\implies x = (x - y) + y \sim 0 + y = y. \end{aligned}$$

Therefore  $\sim$  must have the form

$$x \sim y \iff x - y \in I$$

for some set  $I \subseteq R$ , namely  $I = \{z \in R : z \sim 0\}$ . It remains to characterize which sets  $I$  yield congruences.

1. Reflexivity says that  $x - x \in I$  for any  $x$ . This holds iff  $0 \in I$ .
2. Symmetry says that  $x - y \in I \iff y - x \in I$ . This holds iff  $I$  is closed under negation.
3. Transitivity says that if  $x - y \in I$  and  $y - z \in I$ , then  $x - z \in I$ . This holds iff  $I$  is closed under addition.
4. Compatibility with  $+$  says that if  $x - y \in I$ , then  $(x + a) - (y + a) \in I$ . This condition holds for any  $I$ .
5. Compatibility with  $\cdot$  says that if  $x - y \in I$ , then  $(ax) - (ay) \in I$ . This condition holds iff  $I$  is closed under multiplication by  $R$ .

In summary,  $I$  yields a congruence if and only if the following four properties hold:

- $0 \in I$
- $x \in I \implies -x \in I$
- $x, y \in I \implies x + y \in I$

- $a \in R, x \in I \implies ax \in I$ .

The second condition is an instance of the fourth (take  $a = -1$ ), so it can be removed. The remaining three conditions are the definition of “ideal.”  $\square$

**Remark 2.4.6.** The relation  $x \equiv_I y$  is called “congruence modulo  $I$ ”, and is usually written like  $x \equiv y \pmod{I}$ . When  $I$  is a principal ideal  $aR$ , it is usually written  $x \equiv y \pmod{a}$ .

If  $X$  is a set and  $E$  is an equivalence relation on  $X$ , let  $X/E$  denote the quotient, the set  $\{[a]_E : a \in X\}$ , where  $[a]_E$  is the  $E$ -equivalence class  $[a]_E = \{b \in X : a E b\}$ . Recall that  $[a]_E = [b]_E \iff a E b$ . We omit the subscript when  $E$  is clear from context.

**Theorem 2.4.7** (Quotients). *Let  $A$  be an  $\mathcal{L}$ -algebra and  $E$  be a congruence on  $A$ . Then there is a unique  $\mathcal{L}$ -algebra with underlying set  $A/E$  such that  $A \rightarrow A/E$  is a homomorphism, meaning that*

$$f^{A/E}([a_1], \dots, [a_k]) = [f^A(a_1, \dots, a_k)] \quad (\dagger)$$

for any  $k$ -ary function symbol in  $\mathcal{L}$ .

*Proof.* By Theorem 2.3.1,  $f^{A/E} : (A/E)^k \rightarrow A/E$  is uniquely determined by  $(\dagger)$ , as long as

$$([a_1], \dots, [a_k]) = ([b_1], \dots, [b_k]) \implies [f^A(\bar{a})] = [f^A(\bar{b})],$$

or equivalently,

$$(a_i E b_i \text{ for } i = 1, \dots, k) \implies (f^A(\bar{a}) E f^A(\bar{b})).$$

This holds because  $E$  is a congruence.  $\square$

**Definition 2.4.8.** If  $A$  is an  $\mathcal{L}$ -algebra and  $E$  is a congruence, then  $A/E$  is called the *quotient algebra* of  $A$  by  $E$ .

**Theorem 2.4.9.** *Let  $A$  be an  $\mathcal{L}$ -algebra and  $E$  be a congruence on  $A$ .*

1. *If  $A$  satisfies an equation  $\varphi$ , then  $A/E$  satisfies  $\varphi$ .*
2. *If  $\mathcal{K} = \text{Mod}(\Sigma)$  is an equational class, then  $A \in \mathcal{K} \implies A/E \in \mathcal{K}$ .*

*Proof.* By construction, there is a surjective homomorphism  $A \rightarrow A/E$ . Then  $A \models \varphi \implies A/E \models \varphi$  (Lemma 1.4.10).  $\square$

If  $E$  is an equivalence relation on  $A$ , a *set of representatives* for  $E$  is a set  $S \subseteq A$  containing exactly one element from each  $E$ -equivalence class.

**Theorem 2.4.10.** *Let  $A$  be an  $\mathcal{L}$ -algebra and  $E$  be a congruence. Let  $S$  be a set of representatives for  $E$ . Let  $\rho : A \rightarrow S$  be the map sending  $x \in A$  to the unique  $y \in S \cap [x]_E$ . For each  $k$ -ary function symbol  $f \in \mathcal{L}$ , define*

$$f^S(x_1, \dots, x_k) = \rho(f^A(x_1, \dots, x_k)).$$

*This makes  $S$  into an  $\mathcal{L}$ -algebra isomorphic to  $A/E$ .*

*Proof.* The construction certainly makes  $S$  into an  $\mathcal{L}$ -algebra. Define  $\alpha : S \rightarrow A/E$  by  $\alpha(x) = [x]$ . We claim that  $\alpha$  is an isomorphism.

1.  $\alpha$  is a homomorphism: if  $f$  is a  $k$ -ary function symbol, then

$$\alpha(f^S(x_1, \dots, x_k)) = [\rho(f^A(x_1, \dots, x_k))]$$

But  $\rho(y) E y$  for any  $y \in A$ , so  $[\rho(y)] = [y]$ . Therefore

$$[\rho(f^A(x_1, \dots, x_k))] = [f^A(x_1, \dots, x_k)].$$

By construction of the quotient,

$$[f^A(x_1, \dots, x_k)] = f^{A/E}([x_1], \dots, [x_k]) = f^{A/E}(\alpha(x_1), \dots, \alpha(x_k)).$$

Putting everything together,

$$\alpha(f^S(x_1, \dots, x_k)) = f^{A/E}(\alpha(x_1), \dots, \alpha(x_k)).$$

2.  $\alpha$  is a bijection: clear by choice of  $S$ .  $\square$

## 2.5 Application: finite fields

Work in the ring of integers  $\mathbb{Z}$ .

**Lemma 2.5.1.** *If  $n > 0$ , then  $\{0, 1, \dots, n-1\}$  is a system of representatives for  $\equiv_n$ . That is, for every  $x \in \mathbb{Z}$  there is a unique  $y \in \{0, \dots, n-1\}$  with  $x \equiv y \pmod{n}$ .*

*Proof.* An exercise, by induction on  $x$ . □

**Lemma 2.5.2.** *If  $0 \neq n \in \mathbb{Z}$ , and  $x \in \mathbb{Z}$ , there is  $y \in \mathbb{Z}$  with*

$$\begin{aligned} x &\equiv y \pmod{n} \\ |y| &< |n|. \end{aligned}$$

*Proof.* Since  $n\mathbb{Z} = (-n)\mathbb{Z}$ , we may assume  $n > 0$  by replacing  $n$  with  $-n$  if necessary. Then there is some  $y \in \{0, 1, 2, \dots, n-1\}$  with  $x \equiv y \pmod{n}$  by Lemma 2.5.1. □

**Theorem 2.5.3.** *Every ideal  $I \subseteq \mathbb{Z}$  is a principal ideal  $I = n\mathbb{Z}$  for some  $n \geq 0$ .*

*Proof.* Note that  $\{0\} \subseteq I$ . If  $I = \{0\}$ , take  $n = 0$ . Otherwise, take  $n \in I \setminus \{0\}$  minimizing  $|n|$ . Replacing  $n$  with  $-n$  if necessary, we may assume  $n \geq 0$ . Then  $n\mathbb{Z} \subseteq I$  because  $I$  is an ideal. We claim  $n\mathbb{Z} = I$ . Otherwise, take  $a \in I \setminus n\mathbb{Z}$ . By Lemma 2.5.2 there is  $b \in \mathbb{Z}$  with

$$\begin{aligned} b &\equiv a \pmod{n} \\ |b| &< |n|. \end{aligned}$$

Then  $b - a \in n\mathbb{Z} \subseteq I$ , and  $a \in I$ , so  $b \in I$ . If  $b \neq 0$  then  $b$  contradicts the choice of  $n$ . If  $b = 0$ , then  $a \equiv b = 0 \pmod{n}$ , so  $a \in n\mathbb{Z}$ , contradicting the choice of  $a$ . □

**Theorem 2.5.4.** *If  $n > 0$ , then  $\mathbb{Z}/n\mathbb{Z}$  is isomorphic to  $R = (\{0, 1, \dots, n-1\}, +^R, \cdot^R)$ , where  $x +^R y$  is the unique  $z \in R$  with  $x + y \equiv z \pmod{n}$ , and  $x \cdot^R y$  is the unique  $w \in R$  with  $xy \equiv w \pmod{n}$ . This follows by Theorem 2.4.10 applied to the system of representatives in Lemma 2.5.1. In particular,  $\mathbb{Z}/n\mathbb{Z}$  is finite, of size  $n$ .*

**Lemma 2.5.5.** *Let  $I, J$  be ideals in a ring  $R$ . Let  $I + J := \{x + y : x \in I, y \in J\}$ . Then  $I + J$  is an ideal containing  $I$  and  $J$ .*

*Proof.* To show that  $I + J$  is an ideal, there are three things to check:

1.  $0 \in I + J$ : take  $x = 0 \in I$  and  $y = 0 \in J$ .
2.  $I + J$  is closed under addition: if  $x, x' \in I$  and  $y, y' \in J$ , then  $(x + y) + (x' + y') = (x + x') + (y + y') \in I + J$ .

3.  $I + J$  is closed under multiplication by  $R$ : if  $x \in I, y \in J$ , and  $a \in R$ , then  $a(x + y) = (ax) + (ay) \in I + J$ .

Finally, if  $x \in I$  then  $x + 0 \in I + J$  because  $0 \in J$ . This shows  $I \subseteq I + J$ , and  $J \subseteq I + J$  follows similarly.  $\square$

An ideal  $I \subseteq R$  is *proper* if  $1 \notin I$ . A *maximal ideal* is a maximal proper ideal.

**Theorem 2.5.6.** *If  $I$  is a maximal ideal, then  $R/I$  is a field.*

*Proof.* Let  $[a] \in R/I$  denote the image of  $a \in R$ . First we show that  $1 \neq 0$  in  $R/I$ . The fact that  $1 \notin I$  means that  $1 \not\equiv 0 \pmod{I}$ , so  $[1] \neq [0]$ . Next we show that any  $[a] \neq 0$  has a multiplicative inverse. By Lemma 2.5.5,  $aR + I$  is an ideal containing  $aR$  and  $I$ . The fact that  $[a] \neq 0$  means that  $a = a - 0 \notin I$ . Therefore  $aR + I \supsetneq I$ , as  $a \in aR \subseteq aR + I$ . By maximality of  $I$ ,  $aR + I$  is improper, so  $1 \in aR + I$ . Therefore there are  $x \in R$  and  $y \in I$  with  $1 = ax + y$ . Then  $ax - 1 = y \in I$ , so that  $ax \equiv 1 \pmod{I}$  and  $[a][x] = [ax] = [1]$ . Then  $[x]$  is the multiplicative inverse of  $[a]$ .  $\square$

**Theorem 2.5.7.** *If  $p$  is prime, then  $\mathbb{Z}/p\mathbb{Z}$  is a field.*

*Proof.* It suffices to show that the ideal  $p\mathbb{Z}$  is maximal. If not, take a larger proper ideal  $I \supsetneq p\mathbb{Z}$ . By Theorem 2.5.3,  $I = n\mathbb{Z}$  for some  $n \in \mathbb{Z}$ . Then  $p \in p\mathbb{Z} \subseteq n\mathbb{Z}$ , so  $p$  is a multiple of  $n$ . In other words,  $p = nm$  for some  $m \in \mathbb{Z}$ . As  $p$  is prime, one of  $n$  or  $m$  is  $\pm 1$ .

- If  $n = \pm 1$ , then  $n\mathbb{Z}$  is improper, a contradiction.
- If  $m = \pm 1$ , then  $n = \pm p$ , and  $n\mathbb{Z} = p\mathbb{Z}$ , a contradiction.  $\square$

## 2.6 The fundamental theorem on homomorphisms

**Theorem 2.6.1 (Images).** *Let  $\alpha : A \rightarrow B$  be a homomorphism of  $\mathcal{L}$ -algebras. Then the image  $\text{im}(\alpha) = \alpha(A) = \{\alpha(x) : x \in A\}$  is a subalgebra of  $B$ .*

*Proof.* Suppose  $f \in \mathcal{L}$  is a  $k$ -ary function symbol, and  $b_1, \dots, b_k \in \text{im}(\alpha)$ . Each  $b_i$  can be written as  $\alpha(a_i)$  for some  $a_i \in A$ . Then

$$f(b_1, \dots, b_k) = f(\alpha(a_1), \dots, \alpha(a_k)) = \alpha(f(a_1, \dots, a_k)) \in \text{im}(\alpha). \quad \square$$

**Definition 2.6.2.** Let  $\alpha : A \rightarrow B$  be a homomorphism of  $\mathcal{L}$ -algebras. The *kernel* of  $\alpha$  is the equivalence relation

$$a \sim b \iff \alpha(a) = \alpha(b).$$

We write the kernel as  $\ker(\alpha)$ .

**Theorem 2.6.3.** If  $\alpha : A \rightarrow B$  is a homomorphism of  $\mathcal{L}$ -algebras, then the kernel is a congruence on  $A$ .

*Proof.* Let  $E = \ker(\alpha)$ . Let  $f \in \mathcal{L}$  be a  $k$ -ary function symbol. If  $a_i E b_i$  for  $i = 1, \dots, k$ , then  $\alpha(a_i) = \alpha(b_i)$  for each  $i$ . As  $\alpha$  is a homomorphism,

$$\alpha(f(\bar{a})) = f(\alpha(\bar{a})) = f(\alpha(\bar{b})) = \alpha(f(\bar{b})).$$

Therefore  $f(\bar{a}) E f(\bar{b})$ . □

**Remark 2.6.4.** In ring theory, the kernel of a homomorphism  $f : R \rightarrow S$  is the *ideal*  $\{x \in R : f(x) = 0\}$ . This is the ideal corresponding to the kernel in the sense of Definition 2.6.2.

**Lemma 2.6.5.** Let  $A, B, C$  be algebras. Let  $\alpha : A \rightarrow B$  be a surjective homomorphism and  $\beta : B \rightarrow C$  be a function such that  $\beta \circ \alpha : A \rightarrow C$  is a homomorphism. Then  $\beta$  is a homomorphism.

*Proof.* Let  $f$  be a  $k$ -ary function symbol. If  $\bar{b} \in B^k$ , then  $\bar{b} = \alpha(\bar{a})$  for some  $\bar{a} \in A^k$ . Then

$$\beta(f(\bar{b})) = \beta(f(\alpha(\bar{a}))) = \beta(\alpha(f(\bar{a}))) = f(\beta(\alpha(\bar{a}))) = f(\beta(\bar{b}))$$

because  $\alpha$  and  $\beta \circ \alpha$  are homomorphisms. Therefore  $\beta$  is a homomorphism. □

**Theorem 2.6.6** (Universal property of quotients). Let  $A$  be an algebra and  $E$  be a congruence on  $A$ . If  $\alpha : A \rightarrow B$  is a homomorphism and  $E \subseteq \ker(\alpha)$ , then there is a unique homomorphism  $\beta : A/E \rightarrow B$  such that  $\alpha(x) = \beta([x])$ , or equivalently, the following diagram commutes:

$$\begin{array}{ccc} A & & \\ \downarrow & \searrow \alpha & \\ A/E & \xrightarrow{\beta} & B \end{array}$$



*Proof.* The condition  $E \subseteq \ker(\alpha)$  means that

$$[x]_E = [y]_E \implies \alpha(x) = \alpha(y).$$

By Theorem 2.3.1, there is a unique function  $\beta : A/E \rightarrow B$  such that  $\alpha(x) = \beta([x])$ . We need  $\beta$  to be a homomorphism. The composition  $A \rightarrow A/E \rightarrow B$  is the homomorphism  $\alpha$ , and  $A \rightarrow A/E$  is a surjective homomorphism, so  $\beta : A/E \rightarrow B$  is a homomorphism by Lemma 2.6.5.  $\square$

**Theorem 2.6.7** (Fundamental theorem on homomorphisms). *Let  $\alpha : A \rightarrow B$  be a homomorphism of  $\mathcal{L}$ -algebras. Let  $E$  be the kernel (a congruence on  $A$ ) and let  $\text{im}(\alpha)$  be the image (a subalgebra of  $B$ ). There is an isomorphism  $\beta : A/E \rightarrow \text{im}(\alpha)$ , and  $\alpha$  is the composition of the following three homomorphisms:*

$$A \rightarrow A/E \xrightarrow{\beta} \text{im}(\alpha) \xhookrightarrow{\subseteq} B$$

*Proof.* We can regard  $\alpha : A \rightarrow B$  as a surjective homomorphism  $\alpha' : A \rightarrow \text{im}(\alpha)$ . Then  $\ker(\alpha') = \ker(\alpha) \supseteq E$ , so Theorem 2.6.6 gives a homomorphism  $\beta : A/E \rightarrow \text{im}(\alpha)$  making the diagram commute:

$$\begin{array}{ccc} A & \xrightarrow{\alpha} & B \\ & \searrow \alpha' & \nearrow \\ & A/E \xrightarrow{\beta} \text{im}(\alpha) & \end{array}$$

Then  $\beta$  is an isomorphism:

- $\beta$  is surjective because any element of  $\text{im}(\alpha)$  has the form  $\alpha(a) = \beta([a])$  for some  $a \in A$ .
- $\beta$  is injective because for  $a, b \in A$ ,

$$\beta([a]) = \beta([b]) \iff \alpha(a) = \alpha(b) \iff a E b \iff [a] = [b]. \quad \square$$

In the case of surjective homomorphisms, Theorem 2.6.7 says the following:

**Corollary 2.6.8.** *If  $\alpha : A \rightarrow B$  is a surjective homomorphism, then there is an isomorphism  $A/\ker(\alpha) \rightarrow B$ .*

## 2.7 Application: characteristic of fields

If  $R$  is a ring and  $n \in \mathbb{Z}$ , let  $n^R$  be the interpretation of  $n$  in  $R$ , that is,

$$n^R = \begin{cases} \underbrace{(1 + \cdots + 1)}_{n \text{ times}}^R & \text{if } n > 0 \\ 0^R & \text{if } n = 0 \\ \underbrace{(-(1 + \cdots + 1))}_{n \text{ times}}^R & \text{if } n < 0 \end{cases}$$

For example,  $n^{\mathbb{Z}} = n$ .

**Lemma 2.7.1.** *The map  $\alpha : n \mapsto n^R$  is the unique homomorphism from  $\mathbb{Z}$  to  $R$ .*

*Proof.* The fact that  $\alpha$  is a homomorphism is an exercise in induction and the ring axioms. If  $\beta : \mathbb{Z} \rightarrow R$  is another homomorphism, then  $\beta(n) = \alpha(n)$  by Theorem 1.4.9. For example,

$$\beta(3) = \beta(1 + 1 + 1) = (1 + 1 + 1)^R = \alpha(3). \quad \square$$

The subring of  $R$  generated by the empty set, written  $\langle \emptyset \rangle_R$ , can be described abstractly as the smallest subring of  $R$ , or explicitly as the set of things of the form  $t^R$  for  $t$  a closed term.

**Lemma 2.7.2.** *The image  $\text{im}(\alpha)$  equals  $\langle \emptyset \rangle_R$ .*

*Proof.* The image  $\text{im}(\alpha)$  is a subring, so  $\text{im}(\alpha) \supseteq \langle \emptyset \rangle_R$ . On the other hand,  $\alpha(n) = n^R$  is clearly  $t^R$  for some closed term  $t$ , so  $\alpha(n) \in \langle \emptyset \rangle_R$ . Thus  $\text{im}(\alpha) \subseteq \langle \emptyset \rangle_R$ .  $\square$

We summarize the situation below:

**Theorem 2.7.3.** *If  $R$  is a ring, then there is a unique homomorphism  $\alpha : \mathbb{Z} \rightarrow R$  given by  $\alpha(n) = n^R$ , and the image  $\text{im}(\alpha)$  is the minimal subring  $\langle \emptyset \rangle_R$ .*

**Definition 2.7.4.** The *characteristic* of  $R$ , written  $\text{char}(R)$ , is the unique  $n \in \mathbb{N}$  such that the kernel of  $\mathbb{Z} \rightarrow R$  is the principal ideal  $n\mathbb{Z}$ .

**Theorem 2.7.5.** *If  $R$  has characteristic  $n$ , then the minimal subring  $\langle \emptyset \rangle_R$  is isomorphic to  $\mathbb{Z}/n\mathbb{Z}$ .*

*Proof.* By Theorem 2.7.3,  $\langle \emptyset \rangle_R$  is the image of the unique homomorphism  $\alpha : \mathbb{Z} \rightarrow R$ . By the fundamental theorem on homomorphisms,  $\langle \emptyset \rangle_R = \text{im}(\alpha) \cong \mathbb{Z} / \ker(\alpha) = \mathbb{Z} / n\mathbb{Z}$ , where  $n = \text{char}(R)$ .  $\square$

**Theorem 2.7.6.** *If  $K$  is a field, then  $\text{char}(K) \in \{0, 2, 3, 5, 7, 11, \dots\}$ .*

*Proof.* If  $n = \text{char}(K)$ , and  $\alpha : \mathbb{Z} \rightarrow R$  is the unique homomorphism, then  $\ker(\alpha) = n\mathbb{Z}$ . We must rule out the following cases:

- $n = 1$ . Then  $1 \in 1\mathbb{Z} = \ker(\alpha)$ , so  $1^K = \alpha(1) = 0^K$ , and  $K$  is not a field.
- $n$  is a composite number  $ab$ , for some integers  $a, b > 1$ . Then  $a, b \notin n\mathbb{Z}$  and  $n \in n\mathbb{Z}$ , so  $a, b \notin \ker(\alpha)$  but  $ab = n \in \ker(\alpha)$ . This means that

$$\begin{aligned}\alpha(a) &\neq 0 \\ \alpha(b) &\neq 0 \\ \alpha(a)\alpha(b) &= \alpha(ab) = 0,\end{aligned}$$

contradicting the zero law (Theorem 1.4.16).  $\square$

**Theorem 2.7.7.** *If  $p \in \{0, 2, 3, 5, 7, \dots\}$ , then there is a field of characteristic  $p$ .*

*Proof.* The field  $\mathbb{R}$  has characteristic 0. If  $p > 0$ , the kernel of  $\mathbb{Z} \rightarrow \mathbb{Z}/p\mathbb{Z}$  is  $p\mathbb{Z}$ , so the field  $\mathbb{Z}/p\mathbb{Z}$  has characteristic  $p$ .  $\square$

**Lemma 2.7.8.** *Let  $R$  be a ring. Then  $R/0R \cong R$ .*

*Proof.* By the fundamental theorem on homomorphisms (Corollary 2.6.8) applied to the surjective homomorphism  $\text{id}_R : R \rightarrow R$ , we have  $R \cong R/I$ , where  $I = \ker(\text{id}_R) = \{x \in R : \text{id}(x) = 0\} = \{0\} = 0R$ .  $\square$

**Theorem 2.7.9.** *If  $K$  is a field, then the minimal subring  $\langle \emptyset \rangle_K$  is isomorphic to  $\mathbb{Z}$  if  $\text{char}(K) = 0$ , and  $\mathbb{Z}/p\mathbb{Z}$  if  $\text{char}(K) = p > 0$ .*

**Theorem 2.7.10.** *Let  $L \supseteq K$  be an extension of fields. Then  $\text{char}(L) = \text{char}(K)$ .*

*Proof.* Let  $\alpha : \mathbb{Z} \rightarrow K$  be the unique homomorphism from  $\mathbb{Z}$  to  $K$ . Then  $\alpha$  is also the unique homomorphism from  $\mathbb{Z}$  to  $L$ . The kernel of  $\alpha$  is the same whether we regard  $\alpha$  as a homomorphism to  $K$  or to  $L$ .  $\square$

## 2.8 Term algebras and free algebras

Let  $\bar{x}$  be a tuple of variables, possibly infinite. Let  $T(\bar{x})$  be the set of terms in the variables  $\bar{x}$ . We make  $T(\bar{x})$  into an algebra by setting

$$f^{T(\bar{x})}(t_1, \dots, t_k) = f(t_1, \dots, t_k).$$

**Theorem 2.8.1.** *If  $A$  is an algebra and  $\bar{a}$  is a tuple in  $A$  of the same length as  $\bar{x}$ , then the map*

$$\begin{aligned} \eta : T(\bar{x}) &\rightarrow A \\ \eta(t(\bar{x})) &= t^A(\bar{a}) \end{aligned}$$

*is a homomorphism  $T(\bar{x}) \rightarrow A$ .*

*Proof.* Suppose  $f$  is a  $k$ -ary function symbol, and  $t_1(\bar{x}), \dots, t_k(\bar{x}) \in T(\bar{x})$ . Then

$$\begin{aligned} \eta(f^{T(\bar{x})}(t_1(\bar{x}), \dots, t_k(\bar{x}))) &= \eta(f(t_1(\bar{x}), \dots, t_k(\bar{x}))) \\ &= f^A(t_1^A(\bar{a}), \dots, t_k^A(\bar{a})) = f^A(\eta(t_1(\bar{x})), \dots, \eta(t_k(\bar{x}))). \end{aligned} \quad \square$$

Fix an equational class  $\mathcal{K}$ . Let  $\text{Eq}(\mathcal{K})$  denote the set of equations  $\varphi$  such that every  $A \in \mathcal{K}$  satisfies  $\varphi$ .

**Definition 2.8.2.** If  $t, s$  are two terms, then  $t \equiv_{\mathcal{K}} s$  if the equation  $t = s$  holds for all  $A \in \mathcal{K}$ . In other words,  $t \equiv_{\mathcal{K}} s$  means that  $(t = s) \in \text{Eq}(\mathcal{K})$ .

**Theorem 2.8.3.** *If  $\bar{x}$  is a tuple of variables, then  $\equiv_{\mathcal{K}}$  is a congruence on the term algebra  $T(\bar{x})$ .*

*Proof.* It is easy to see that  $\equiv_{\mathcal{K}}$  is an equivalence relation. Suppose  $t_1, \dots, t_k$  and  $s_1, \dots, s_k$  are terms in  $T(\bar{x})$  with  $t_i \equiv_{\mathcal{K}} s_i$  for each  $i$ , and  $f$  is a  $k$ -ary function symbol. Then for any  $A \in \mathcal{K}$  and tuple  $\bar{a}$  in  $A$ ,

$$t_i^A(\bar{a}) = s_i^A(\bar{a}) \text{ for } i = 1, \dots, k,$$

and so

$$f(t_1(\bar{a}), \dots, t_k(\bar{a}))^A = f(s_1(\bar{a}), \dots, s_k(\bar{a}))^A.$$

It follows that

$$f(t_1(\bar{x}), \dots, t_k(\bar{x})) \equiv_{\mathcal{K}} f(s_1(\bar{x}), \dots, s_k(\bar{x})). \quad \square$$

**Definition 2.8.4.** The  $(\mathcal{K})$ -free algebra on the variables  $\bar{x}$ , written  $F_{\mathcal{K}}(\bar{x})$  or  $F(\bar{x})$ , is the quotient  $T(\bar{x})/\equiv_{\mathcal{K}}$ .

**Theorem 2.8.5.**  $F_{\mathcal{K}}(\bar{x}) \in \mathcal{K}$ .

*Proof.* Let  $t(y_1, \dots, y_k) = s(y_1, \dots, y_k)$  be any of the axioms defining  $\mathcal{K}$ . For any terms  $u_1, \dots, u_k \in T(\bar{x})$ , any  $A \in \mathcal{K}$ , and any  $\bar{a}$  in  $A$ , we have

$$t(u_1(\bar{a}), \dots, u_k(\bar{a})) = s(u_1(\bar{a}), \dots, u_k(\bar{a}))$$

because  $A \models t = s$ . Therefore

$$\begin{aligned} t(u_1(\bar{x}), \dots, u_k(\bar{x})) &\equiv_{\mathcal{K}} s(u_1(\bar{x}), \dots, u_k(\bar{x})) \\ [t(u_1(\bar{x}), \dots, u_k(\bar{x}))] &= [s(u_1(\bar{x}), \dots, u_k(\bar{x}))] \text{ in } F(\bar{x}) \\ t([u_1(\bar{x})], \dots, [u_k(\bar{x})]) &= s([u_1(\bar{x})], \dots, [u_k(\bar{x})]) \text{ in } F(\bar{x}) \end{aligned}$$

Thus  $F(\bar{x}) \models t = s$ . □

**Theorem 2.8.6** (Universal mapping property). *For any  $A \in \mathcal{K}$  and tuple  $\bar{a}$  in  $A$  (of the same length as  $\bar{x}$ ), there is a homomorphism  $\alpha : F(\bar{x}) \rightarrow A$  sending  $[t(\bar{x})]$  to  $t(\bar{a})$ .*

*Proof.* Let  $\beta : T(\bar{x}) \rightarrow A$  be the evaluation map  $t(\bar{x}) \mapsto t(\bar{a})$ . Note that

$$t \equiv_{\mathcal{K}} s \implies t(\bar{a}) = s(\bar{a}) \iff \beta(t) = \beta(s).$$

Therefore  $(\equiv_{\mathcal{K}}) \subseteq \ker(\beta)$ . By the universal property of quotients (Theorem 2.6.6), there is a homomorphism  $\alpha : F(\bar{x}) \rightarrow A$  making the diagram commute

$$\begin{array}{ccc} T(\bar{x}) & & \\ \downarrow & \searrow \beta & \\ F(\bar{x}) & \xrightarrow{\alpha} & A. \end{array}$$

Thus  $\alpha([t(\bar{x})]) = \beta(t(\bar{x})) = t(\bar{a})$ . □

**Corollary 2.8.7.** *If  $A \in \mathcal{K}$ , then there is a surjective homomorphism  $\alpha : F(\bar{x}) \rightarrow A$  for some  $\bar{x}$ .*

*Proof.* Let  $\bar{a} = (a_i : i \in I)$  be a tuple (probably infinite) enumerating all of  $A$ . Let  $\bar{x} = (x_i : i \in I)$  be a tuple of variables of the same length. Let  $\alpha : F(\bar{x}) \rightarrow A$  be the homomorphism sending  $[t(\bar{x})]$  to  $t^A(\bar{a})$ . Letting  $t(\bar{x}) = x_i$ , we see that  $\alpha([x_i]) = a_i$ , so  $\alpha$  is surjective. □

## 2.9 Birkhoff's HSP theorem

**Lemma 2.9.1.** *Let  $\alpha_i : A \rightarrow B_i$  be a homomorphism of  $\mathcal{L}$ -algebras for  $i \in I$ . Let  $\alpha : A \rightarrow \prod_{i \in I} B_i$  be the map  $\alpha(a) = (\alpha_i(a) : i \in I)$ . Then  $\alpha$  is a homomorphism.*

*Proof.* Let  $\pi_j$  be the  $j$ th coordinate projection from  $\prod_{i \in I} B_i$  to  $B_j$ . If  $f$  is a  $k$ -ary function symbol and  $a_1, \dots, a_k \in A$ , we must show

$$\alpha(f(\bar{a})) \stackrel{?}{=} f(\alpha(\bar{a})).$$

Both sides are in  $\prod_{i \in I} B_i$ . If the two sides disagree, then there is  $i \in I$  such that

$$\pi_i(\alpha(f(\bar{a}))) \neq \pi_i(f(\alpha(\bar{a}))).$$

As  $\pi_i$  is a homomorphism (Remark 2.2.9), we can change the right hand side:

$$\pi_i(\alpha(f(\bar{a}))) \neq f(\pi_i(\alpha(\bar{a}))).$$

Now  $\pi_i \circ \alpha = \alpha_i$  by definition of  $\alpha$ , so

$$\alpha_i(f(\bar{a})) \neq f(\alpha_i(\bar{a})).$$

This contradicts the fact that  $\alpha_i$  is a homomorphism. □

**Theorem 2.9.2** (Birkhoff's HSP theorem). *Let  $\mathcal{K}$  be a class of algebras. Then  $\mathcal{K}$  is an equational class if and only if  $\mathcal{K}$  is closed under isomorphisms, subalgebras, products, and quotients.*

*Proof.* Equational classes are closed under isomorphisms (Theorem 1.4.11), subalgebras (Theorem 2.1.2), products (Theorem 2.2.10), and quotients (Theorem 2.4.9).

Conversely, suppose  $\mathcal{K}$  is closed under isomorphisms, subalgebras, products, and quotients. Let  $\Sigma$  be  $\text{Eq}(\mathcal{K})$ , the set of equations holding on  $\mathcal{K}$ . If  $A \in \mathcal{K}$ , then  $A \models \Sigma$ , and so  $\mathcal{K} \subseteq \text{Mod}(\Sigma) =: \bar{\mathcal{K}}$ .

*Claim.* If  $\bar{x}$  is a tuple of variables, then the  $\bar{\mathcal{K}}$ -free algebra  $F_{\bar{\mathcal{K}}}(\bar{x})$  is in  $\mathcal{K}$ .

*Proof.* Let  $\{(a_i, b_i)\}_{i \in I}$  enumerate all the pairs of distinct elements of  $F = F_{\bar{\mathcal{K}}}(\bar{x})$ . For each  $i$ , we can write  $a_i = [t_i(\bar{x})]$  and  $b_i = [s_i(\bar{x})]$  for terms  $t_i, s_i \in T(\bar{x})$ . The fact that  $a_i \neq b_i$  means that  $t_i(\bar{x}) \not\equiv_{\bar{\mathcal{K}}} s_i(\bar{x})$ . Thus  $(t_i = s_i) \notin \Sigma = \text{Eq}(\mathcal{K})$ , so there is  $A_i \in \mathcal{K}$  with  $A_i \not\models t_i = s_i$ . Then there

is  $\bar{a}_i \in A_i$  with  $t_i^A(\bar{a}_i) \neq s_i^A(\bar{a}_i)$ . As  $A_i \in \mathcal{K} \subseteq \bar{\mathcal{K}}$ , there is a homomorphism  $\beta_i : F(\bar{x}) \rightarrow A_i$  sending  $[t(\bar{x})]$  to  $t^{A_i}(\bar{a}_i)$  (see Theorem 2.8.6). Then

$$\beta_i(a_i) = \beta_i([t_i(\bar{x})]) = t_i^{A_i}(\bar{a}_i) \neq s_i^{A_i}(\bar{a}_i) = \beta_i([s_i(\bar{x})]) = \beta_i(b_i).$$

Thus, for every  $i \in I$ ,

$$\beta_i(a_i) \neq \beta_i(b_i). \quad (*)$$

By Lemma 2.9.1, there is a homomorphism  $\beta : F(\bar{x}) \rightarrow \prod_{i \in I} A_i$  with  $\beta(x) = (\beta_i(x) : i \in I)$ . Note that  $\prod_{i \in I} A_i \in \mathcal{K}$  because  $\mathcal{K}$  is closed under products. We claim that  $\beta$  is injective. Indeed, if  $a, b \in F(\bar{x})$  and  $a \neq b$ , then  $(a, b) = (a_i, b_i)$  for some  $i \in I$ . Then  $\beta_i(a) = \beta_i(a_i) \neq \beta_i(b_i) = \beta_i(b)$ , so  $\beta(a)$  and  $\beta(b)$  differ at the  $i$ th coordinate.

Then  $\beta$  is an isomorphism from  $F(\bar{x})$  to  $\text{im}(\beta)$ . As  $\text{im}(\beta)$  is a subalgebra of  $\prod_{i \in I} A_i \in \mathcal{K}$ , we have  $\text{im}(\beta) \in \mathcal{K}$  and then  $F(\bar{x}) \in \mathcal{K}$ .  $\square_{\text{Claim}}$

Now if  $A \in \bar{\mathcal{K}}$  then there is a surjective homomorphism  $F(\bar{x}) \rightarrow A$  for some  $\bar{\mathcal{K}}$ -free algebra  $F(\bar{x})$  (Corollary 2.8.7). By the fundamental theorem on homomorphisms (Corollary 2.6.8),  $A$  is isomorphic to a quotient of  $F(\bar{x})$ . By the claim,  $F(\bar{x}) \in \mathcal{K}$ , and thus  $A \in \mathcal{K}$ .

This shows that  $\bar{\mathcal{K}} \subseteq \mathcal{K}$ . As  $\mathcal{K} \subseteq \bar{\mathcal{K}}$ , the class  $\mathcal{K}$  equals the equational class  $\bar{\mathcal{K}}$ .  $\square$





# Chapter 3

## First-order logic

### 3.1 Languages, structures, formulas, and satisfaction

**Definition 3.1.1.** A language  $\mathcal{L}$  consists of a set of *function symbols*, a set of *relation symbols*, and a function assigning to each function or relation symbol  $X$  an integer  $n_X \in \mathbb{N}$  called the *arity* of  $X$ . A symbol  $X$  is said to be *k-ary* if  $n_X = k$ . Nullary function symbols are called *constant symbols*.

**Example 3.1.2.** The *language of posets* contains one binary relation symbol  $\leq$ . The *language of monoids* contains one binary function symbol  $\cdot$  and one constant symbol  $1$ .

Let  $M$  be a set. For  $n \geq 0$ , an *n-ary relation* on  $M$  is a subset  $R \subseteq M^n$ . If  $a_1, \dots, a_n \in M$ , then “ $R(a_1, \dots, a_n)$ ” means “ $(a_1, \dots, a_n) \in R$ ”. We think of  $R$  as an *n-ary function*  $M^n \rightarrow \{\text{FALSE}, \text{TRUE}\}$ .

**Definition 3.1.3.** Let  $\mathcal{L}$  be a language. An  $\mathcal{L}$ -*structure*  $\mathcal{M}$  consists of the following:

1. A set  $M$ , called the *underlying set* of  $\mathcal{M}$ .
2. A map assigning an *n-ary operation*  $f^{\mathcal{M}} : M^n \rightarrow M$  to each *n-ary function symbol*  $f$ .
3. A map assigning an *n-ary relation*  $R^{\mathcal{M}} \subseteq M^n$  to each *n-ary relation symbol*  $R$ .

Usually we don't distinguish between a structure  $\mathcal{M}$  and its underlying set  $M$ , writing both as  $M$ .

**Example 3.1.4.** If  $\mathcal{L}$  is the language of posets, then an  $\mathcal{L}$ -structure is essentially a pair  $(M, \leq^M)$  where  $M$  is a set and  $\leq^M$  is a binary relation on  $M$ .

**Definition 3.1.5.** Let  $\mathcal{L}^+$  be a language and  $\mathcal{L}^-$  be a sublanguage. For any  $\mathcal{L}^+$ -structure  $M$ , we can form an  $\mathcal{L}^-$ -structure from  $M$  by forgetting about the symbols in  $\mathcal{L}^+ \setminus \mathcal{L}^-$ . The resulting structure  $M \upharpoonright \mathcal{L}^-$  is called a *reduct* of  $M$ . Conversely,  $M$  is an *expansion* of  $M \upharpoonright \mathcal{L}^-$ .

Fix a language  $\mathcal{L}$  and a class  $\mathcal{V} = \{x, y, z, \dots\}$  of *variable symbols* disjoint from the symbols in  $\mathcal{L}$ .

**Definition 3.1.6.** An  $\mathcal{L}$ -term is a string generated by the following rules:

- If  $x$  is a variable symbol, then  $x$  is a term.
- If  $f$  is an  $n$ -ary function symbol in  $\mathcal{L}$ , and  $t_1, \dots, t_n$  are  $\mathcal{L}$ -terms, then  $f(t_1, \dots, t_n)$  is an  $\mathcal{L}$ -term.

**Definition 3.1.7.** An  $\mathcal{L}$ -formula is a string generated by the following rules:

1. If  $R$  is a  $k$ -ary relation symbol in  $\mathcal{L}$  and  $t_1, \dots, t_k$  are  $\mathcal{L}$ -terms, then  $R(t_1, \dots, t_k)$  is an  $\mathcal{L}$ -formula.
2. If  $t$  and  $s$  are  $\mathcal{L}$ -terms, then  $(t = s)$  is an  $\mathcal{L}$ -formula.
3. If  $\varphi$  and  $\psi$  are  $\mathcal{L}$ -formulas, then so are  $\varphi \wedge \psi$ ,  $\varphi \vee \psi$ , and  $\neg\varphi$ .
4.  $\top$  and  $\perp$  are  $\mathcal{L}$ -formulas.
5. If  $\varphi$  is an  $\mathcal{L}$ -formula and  $x$  is a variable symbol, then the following are  $\mathcal{L}$ -formulas:

$$\exists x (\varphi) \in \mathcal{F}$$

$$\forall x (\varphi) \in \mathcal{F}.$$

The set of *atomic  $\mathcal{L}$ -formulas* is the subset generated by (1)–(2), while the set of *quantifier-free  $\mathcal{L}$ -formulas* is generated by (1)–(4). If  $x$  is a variable symbol and  $\alpha$  is a term or formula, an occurrence of  $x$  in  $\alpha$  is *bound* if it

occurs inside a quantifier  $\exists x (\dots)$  or  $\forall x (\dots)$ , and *free* otherwise. The *free variables* of  $\alpha$  are the variables with free occurrences in  $\alpha$ . A *closed term* is a term with no free variables (i.e., no variables). A *sentence* is a formula with no free variables.

We write “let  $\varphi(x_1, \dots, x_n)$  be a formula” to mean that  $\varphi(x_1, \dots, x_n)$  is a formula whose free variables are contained in  $\{x_1, \dots, x_n\}$ . We use the same convention for terms. If  $\varphi(x_1, \dots, x_n)$  is a formula and  $t_1, \dots, t_n$  are terms, then  $\varphi(t_1, \dots, t_n)$  denotes the result of replacing the free occurrences of  $x_i$  with  $t_i$  in  $\varphi$ .

Let  $M$  be an  $\mathcal{L}$ -structure and  $A$  be a subset of  $M$ . The language  $\mathcal{L}(A)$  is obtained by adding each element of  $A$  as a new constant symbol. We regard  $M$  as an  $\mathcal{L}(A)$ -structure by interpreting each new constant symbol as the corresponding element of  $A$ :

$$a^M = a.$$

If  $t$  is a closed  $\mathcal{L}(M)$ -term, define the interpretation  $t^M$  recursively as follows:

$$f(t_1, \dots, t_k)^M = f^M(t_1^M, \dots, t_k^M),$$

If  $t(x_1, \dots, x_n)$  is an  $\mathcal{L}$ -term or  $\mathcal{L}(M)$ -term, then the interpretation  $t^M$  is the function  $M^n \rightarrow M$  defined by

$$t^M(a_1, \dots, a_n) = (t(a_1, \dots, a_n))^M.$$

If  $\varphi$  is an  $\mathcal{L}(M)$ -sentence, we define  $M \models \varphi$  recursively:

$$\begin{aligned} M \models t = s &\iff t^M = s^M \\ M \models R(t_1, \dots, t_k) &\iff (t_1^M, \dots, t_k^M) \in R^M \\ M \models \varphi \vee \psi &\iff (M \models \varphi \text{ or } M \models \psi) \\ M \models \varphi \wedge \psi &\iff (M \models \varphi \text{ and } M \models \psi) \\ M \models \neg \varphi &\iff M \not\models \varphi \\ M \models \top &\text{ is always true} \\ M \models \perp &\text{ is always false} \\ M \models \exists x \varphi(x) &\iff \exists a \in M : M \models \varphi(a) \\ M \models \forall x \varphi(x) &\iff \forall a \in M : M \models \varphi(a). \end{aligned}$$

If  $M \models \varphi$ , we say that  $M$  *satisfies*  $\varphi$ , or  $\varphi$  is *true* in  $M$ .

## 3.2 Theories and elementary classes

Fix a language  $\mathcal{L}$ . An  $\mathcal{L}$ -theory is a set  $T$  of  $\mathcal{L}$ -sentences. Elements of  $T$  are called *axioms*. An  $\mathcal{L}$ -structure  $M$  *satisfies*  $T$ , or is a *model* of  $T$ , written  $M \models T$ , if

$$M \models \varphi \text{ for every } \varphi \in T.$$

The set of models of  $T$  is written  $\text{Mod}(T)$ . An *elementary class* is a class of the form  $\text{Mod}(T)$ .

**Example 3.2.1.** Let  $\mathcal{L}$  be the language of rings. The *theory of rings*  $T_{\text{rings}}$  has the following axioms:

$$\begin{aligned} & \forall x \forall y (x + y = y + x) \\ & \forall x \forall y \forall z (x + (y + z) = (x + y) + z) \\ & \dots \end{aligned}$$

Models of  $T_{\text{rings}}$  are exactly rings. The *theory of fields*  $T_{\text{fields}}$  is  $T_{\text{rings}}$  plus the following two axioms:

$$\begin{aligned} & \neg(0 = 1) \\ & \forall x (x = 0 \vee \exists y (x \cdot y = 1)). \end{aligned}$$

Models of  $T_{\text{fields}}$  are fields.

## 3.3 Common abbreviations

The following abbreviations are standard:

Abbreviation	Meaning
$\varphi \rightarrow \psi$	$\neg\varphi \vee \psi$
$\varphi \leftarrow \psi$	$\psi \rightarrow \varphi$
$\varphi \leftrightarrow \psi$	$(\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi)$
$t \neq s$	$\neg t = s$
$\bigwedge_{i=1}^n \varphi_i$	$\varphi_1 \wedge \dots \wedge \varphi_n$ [or $\top$ if $n = 0$ ]
$\bigvee_{i=1}^n \varphi_i$	$\varphi_1 \vee \dots \vee \varphi_n$ [or $\perp$ if $n = 0$ ]
$\exists^{\geq n} \varphi(x)$	$\exists y_1, \dots, y_n \left( \bigwedge_{i=1}^n \varphi(y_i) \wedge \bigwedge_{1 \leq i < j \leq n} (y_i \neq y_j) \right)$
$\exists^{=n} \varphi(x)$	$\exists^{\geq n} x \varphi(x) \wedge \neg \exists^{\geq n+1} x \varphi(x)$
$\exists! x \varphi(x)$	$\exists^{=1} x \varphi(x).$

Additionally, if  $\mathcal{L}$  contains a symbol  $\leq$ , then

$$\begin{aligned} t \geq s &\text{ means } s \leq t \\ t < s &\text{ means } t \leq s \wedge t \neq s \\ t > s &\text{ means } s < t. \end{aligned}$$

**Example 3.3.1.** Let  $\mathcal{L} = \{\leq\}$  be the language of posets. The *theory of posets*, whose models are posets, has the following axioms:

$$\begin{aligned} &\forall x(x \leq x) \\ &\forall x, y, z(x \leq y \wedge y \leq z \rightarrow x \leq z) \\ &\forall x, y(x \leq y \wedge y \leq x \rightarrow x = y). \end{aligned}$$

The *theory of linear orders* adds one more axiom:

$$\forall x, y(x \leq y \vee y \leq x).$$

### 3.4 Elementary equivalence and embeddings

The *complete theory* of an  $\mathcal{L}$ -structure  $M$  is the set

$$\text{Th}(M) := \{\varphi : \varphi \text{ is an } \mathcal{L}\text{-sentence and } M \models \varphi\}.$$

Two  $\mathcal{L}$ -structures  $M$  and  $N$  are *elementarily equivalent*, written  $M \equiv N$ , if  $\text{Th}(M) = \text{Th}(N)$ , meaning that  $M \models \varphi \iff N \models \varphi$  for every  $\mathcal{L}$ -sentence  $\varphi$ .

**Theorem 3.4.1.** *A structure  $N$  is a model of  $\text{Th}(M)$  if and only if  $N \equiv M$ .*

*Proof.* Note that  $N \models \text{Th}(M)$  iff  $\text{Th}(N) \supseteq \text{Th}(M)$ . (More generally,  $N \models T$  iff  $\text{Th}(N) \supseteq T$ .) So we must rule out the case that  $\text{Th}(N) \supsetneq \text{Th}(M)$ . Suppose  $\varphi \in \text{Th}(N) \setminus \text{Th}(M)$ . Then  $N \models \varphi$  and  $M \not\models \varphi$ , implying that  $N \not\models \neg\varphi$  and  $M \models \neg\varphi$ . Then  $(\neg\varphi) \in \text{Th}(M) \setminus \text{Th}(N)$ , contradicting the fact that  $\text{Th}(N) \supseteq \text{Th}(M)$ .  $\square$

**Definition 3.4.2.** Let  $M, N$  be  $\mathcal{L}$ -structures. A function  $\alpha : M \rightarrow N$  is an *embedding* if  $\alpha$  is injective, and  $\alpha$  strictly preserves the function and relation symbols:

$$\begin{aligned} \alpha(f^M(b_1, \dots, b_n)) &= f^N(\alpha(b_1), \dots, \alpha(b_n)) \\ R^M(b_1, \dots, b_n) &\iff R^N(\alpha(b_1), \dots, \alpha(b_n)). \end{aligned}$$

An *isomorphism* is a bijective embedding. Two structures  $M$  and  $N$  are *isomorphic*, written  $M \cong N$ , if there is an isomorphism from  $M$  to  $N$ .

The analogue of Theorem 1.4.6 holds, and so  $\cong$  is an equivalence relation.

**Theorem 3.4.3.** *Let  $f : M \rightarrow N$  be an embedding and  $a_1, \dots, a_n$  be in  $M$ .*

1. *If  $t(x_1, \dots, x_n)$  is a term, then*

$$f(t^M(a_1, \dots, a_n)) = t^N(f(a_1), \dots, f(a_n)).$$

2. *If  $\varphi(x_1, \dots, x_n)$  is a quantifier-free formula, then*

$$M \models \varphi(a_1, \dots, a_n) \iff N \models \varphi(f(a_1), \dots, f(a_n)).$$

3. *If  $f$  is an isomorphism and  $\varphi(x_1, \dots, x_n)$  is any formula, then*

$$M \models \varphi(a_1, \dots, a_n) \iff N \models \varphi(f(a_1), \dots, f(a_n)).$$

*Proof.* By induction on the complexity of  $t$  or  $\varphi$ , like in Theorem 1.4.9. (In fact, part (1) is an instance of Theorem 1.4.9.)  $\square$

**Definition 3.4.4.** A function  $\alpha : M \rightarrow N$  is an *elementary embedding* if  $\alpha$  preserves all  $\mathcal{L}$ -formulas  $\varphi(x_1, \dots, x_n)$ , in the sense that

$$M \models \varphi(b_1, \dots, b_n) \iff N \models \varphi(\alpha(b_1), \dots, \alpha(b_n)). \quad (*)$$

**Remark 3.4.5.** 1. Any elementary embedding is an embedding: apply (\*) to the following formulas:

$$\begin{aligned} x_1 &= x_2 \\ R(x_1, \dots, x_k) \\ f(x_1, \dots, x_k) &= x_{k+1} \end{aligned}$$

2. Theorem 3.4.3(3) says that isomorphisms are elementary embeddings.

3. If  $\alpha : M \rightarrow N$  is an elementary embedding, then  $M \equiv N$ . To see this, take  $n = 0$  in (\*), so that  $\varphi$  is a sentence. Then (\*) says that  $M \models \varphi \iff N \models \varphi$ .

**Corollary 3.4.6.** *If  $M \cong N$ , then  $M \equiv N$ .*

### 3.5 Substructures and extensions

Let  $M$  be an  $\mathcal{L}$ -structure.

**Definition 3.5.1.** A *substructure* of  $M$  is a subset  $A \subseteq M$  such that for any  $k$ -ary function symbol  $f$  in  $\mathcal{L}$ ,

$$a_1, \dots, a_k \in A \implies f^M(a_1, \dots, a_k) \in A.$$

If  $A$  is a substructure of  $M$ , we regard  $A$  as an  $\mathcal{L}$ -structure by defining  $f^A$  and  $R^A$  to be the restrictions of  $f^M$  and  $R^M$  to  $A$ . In this way, substructures are structures, not just sets.

If  $M$  is an  $\mathcal{L}$ -structure, an *extension* of  $M$  is an  $\mathcal{L}$ -structure  $N$  such that  $M$  is a substructure of  $N$ .

**Definition 3.5.2.** If  $A$  is a subset of  $M$ , then  $\langle A \rangle_M$  denotes the smallest substructure of  $M$  containing  $A$ , which is

$$\{t^M(\bar{b}) : t(x_1, \dots, x_n) \text{ is an } \mathcal{L}\text{-term and } \bar{b} \in A^n\}$$

The substructure  $\langle A \rangle_M$  is called the substructure *generated* by  $A$ . We say that  $M$  is *finitely generated* if  $M = \langle A \rangle$  for some finite  $A \subseteq M$ .

**Remark 3.5.3.** If the language  $\mathcal{L}$  has no function symbols or constant symbols, then every subset of  $M$  is a substructure, and so  $\langle A \rangle_M = A$ .

**Remark 3.5.4.** If  $A$  is a substructure of  $M$ , then the inclusion  $A \hookrightarrow M$  is an embedding. Therefore

$$A \models \varphi(\bar{a}) \iff M \models \varphi(\bar{a})$$

for *quantifier-free* formulas  $\varphi$  and tuples  $\bar{a}$  in  $A$ .

**Definition 3.5.5.** Let  $M$  and  $N$  be  $\mathcal{L}$ -structures. We say that  $M$  is an *elementary substructure* of  $N$ , written  $M \preceq N$ , or that  $N$  is an *elementary extension* of  $M$ , written  $N \succeq M$ , if  $M$  is a substructure of  $N$  and the inclusion  $M \rightarrow N$  is an elementary embedding, meaning that

$$M \models \varphi(a_1, \dots, a_n) \iff N \models \varphi(a_1, \dots, a_n) \quad (*)$$

for  $\mathcal{L}$ -formulas  $\varphi$  and  $\bar{a} \in M^n$ .

Equation (\*) says that  $M$  and  $N$  satisfy the same  $\mathcal{L}(M)$ -sentences.

**Theorem 3.5.6.** *Let  $f$  be an embedding from  $M$  to  $N$ .*

1. *The image  $\text{im}(f)$  is a substructure of  $N$ , and  $M \rightarrow \text{im}(f)$  is an isomorphism.*
2. *If  $f$  is an elementary embedding, then  $\text{im}(f) \preceq N$ .*

*Proof.* 1. Straightforward. The fact that  $\text{im}(f)$  is a substructure is like the analogous fact for homomorphisms (Theorem 2.6.1).

2. If  $M' = \text{im}(f)$ , then the inclusion  $M' \hookrightarrow N$  is the composition of two elementary maps:

$$M' \xrightarrow{\cong} M \xrightarrow{f} N. \quad \square$$

### 3.6 Complete theories

We write  $T \vdash \varphi$  if  $\varphi$  is provable from  $T$ , and  $T \models \varphi$  if every model of  $T$  satisfies  $\varphi$ .

**Fact 3.6.1** (Soundness and completeness theorem). *If  $T$  is a theory and  $\varphi$  is a sentence, then*

$$T \vdash \varphi \iff T \models \varphi.$$

A theory  $T$  is *inconsistent* if the following equivalent conditions hold:

1.  $T \vdash \perp$ .
2.  $T \vdash \varphi$  for all  $\varphi$ .
3.  $T \vdash \varphi$  and  $T \vdash \neg\varphi$  for some  $\varphi$ .

Otherwise,  $T$  is *consistent*. Note that  $T \models \perp$  if and only if  $T$  has no models, because no structure  $M$  satisfies  $\perp$ . Therefore the completeness theorem implies the following:

**Theorem 3.6.2.**  *$T$  is consistent if and only if  $T$  has a model.*

A theory  $T$  is *complete* if  $T$  is consistent and for any sentence  $\varphi$ ,

$$T \vdash \varphi \text{ or } T \vdash \neg\varphi.$$



**Theorem 3.6.3.** *If  $T$  is complete and  $M \models T$ , then for any sentence  $\varphi$ ,*

$$M \models \varphi \iff T \vdash \varphi.$$

*Proof.* As  $M \models T$  and  $T$  is complete,

$$\begin{aligned} T \vdash \varphi &\implies M \models \varphi \\ T \not\vdash \varphi &\implies T \vdash \neg\varphi \implies M \models \neg\varphi \implies M \not\models \varphi. \end{aligned} \quad \square$$

**Theorem 3.6.4.** *Let  $T$  be a theory. Then  $T$  is complete if and only if any two models  $M_1, M_2 \models T$  are elementarily equivalent.*

*Proof.* First suppose  $T$  is complete. If  $M_1, M_2$  are two models, then  $\text{Th}(M_1) = \{\varphi : T \vdash \varphi\} = \text{Th}(M_2)$  by Theorem 3.6.3, and so  $M_1 \equiv M_2$ .

Conversely suppose  $T$  is incomplete, with  $T \not\vdash \varphi$  and  $T \not\vdash \neg\varphi$ . By the completeness theorem, there are models  $M_1, M_2 \models T$  with  $M_1 \models \neg\varphi$  and  $M_2 \models \varphi$ . Then  $M_1 \not\equiv M_2$ .  $\square$

**Fact 3.6.5.** *Let  $\mathcal{L}$  be a countable language. If  $T$  is finite or more generally if  $T$  is computably enumerable, then the set  $\overline{T} = \{\varphi : T \vdash \varphi\}$  is computably enumerable.*

**Theorem 3.6.6.** *If  $\mathcal{L}$  is countable,  $T$  is complete and computably enumerable, and  $M$  is a model of  $T$ , then  $\text{Th}(M)$  is computable—there is an algorithm which takes an  $\mathcal{L}$ -sentence  $\varphi$  and determines whether  $M \models \varphi$ .*

*Proof.* By Theorem 3.6.3,  $\text{Th}(M) = \overline{T}$ , which is computably enumerable by Fact 3.6.5. However,

$$\varphi \notin \text{Th}(M) \iff \neg\varphi \in \text{Th}(M),$$

and so  $\text{Th}(M)$  is also co-c.e. As  $\text{Th}(M)$  is both c.e. and co-c.e., it is computable.  $\square$

### 3.7 Completeness via back-and-forth systems

Let  $M, N$  be two  $\mathcal{L}$ -structures.

**Definition 3.7.1.** A *partial elementary map* from  $M$  to  $N$  is a bijection  $f : A \rightarrow B$  where  $A$  and  $B$  are subsets of  $M$  and  $N$ , respectively, and  $f$  preserves all  $\mathcal{L}$ -formulas, in the sense that

$$M \models \varphi(a_1, \dots, a_n) \iff N \models \varphi(f(a_1), \dots, f(a_n))$$

for  $n \geq 0$ ,  $\varphi(x_1, \dots, x_n)$  an  $\mathcal{L}$ -formula, and  $a_1, \dots, a_n \in A$ .

**Remark 3.7.2.** Taking  $n = 0$ , we see that if  $f$  is a partial elementary map from  $M$  to  $N$ , then  $M \equiv N$ .

Conversely,  $\emptyset : \emptyset \rightarrow \emptyset$  is a partial elementary map from  $M$  to  $N$  if and only if  $M \equiv N$ .

**Remark 3.7.3.** An elementary embedding from  $M$  to  $N$  is the same thing as a partial elementary map  $f$  from  $M$  to  $N$  with  $\text{dom}(f) = M$ .

**Definition 3.7.4.** A *partial isomorphism* from  $M$  to  $N$  is an isomorphism  $f : A \rightarrow B$  where  $A$  is a substructure of  $M$  and  $B$  is a substructure of  $N$ .

By Theorem 3.4.3, partial isomorphisms preserve quantifier-free formulas: if  $f : A \rightarrow B$  is a partial isomorphism and  $\bar{a} \in A^n$  and  $\varphi(\bar{x})$  is quantifier-free, then

$$M \models \varphi(\bar{a}) \iff A \models \varphi(\bar{a}) \iff B \models \varphi(f(\bar{a})) \iff N \models \varphi(f(\bar{a})).$$

**Definition 3.7.5.** A *back-and-forth system* between  $M$  and  $N$  is a family  $\mathcal{F}$  of partial isomorphisms satisfying the following conditions:

1. Forward: if  $f : A \rightarrow B$  is in  $\mathcal{F}$ , and  $a' \in M$ , then there is  $f' : A' \rightarrow B'$  in  $\mathcal{F}$  such that  $f'$  extends  $f$  and  $a' \in A'$ .
2. Backward: if  $f : A \rightarrow B$  is in  $\mathcal{F}$ , and  $b' \in N$ , then there is  $f' : A' \rightarrow B'$  in  $\mathcal{F}$  such that  $f'$  extends  $f$  and  $b' \in B'$ .

**Theorem 3.7.6.** If  $\mathcal{F}$  is a back-and-forth system, then every  $f \in \mathcal{F}$  is a partial elementary map from  $M$  to  $N$ .

*Proof.* We show by induction on the complexity of  $\varphi(x_1, \dots, x_n)$  that if  $f : A \rightarrow B$  is in  $\mathcal{F}$  and  $a_1, \dots, a_n \in A$ , then

$$M \models \varphi(a_1, \dots, a_n) \iff N \models \varphi(f(a_1), \dots, f(a_n)). \quad (\dagger)$$

The case of atomic formulas holds because  $f$  is a partial isomorphism. The case of boolean operations is straightforward. It remains to consider the case where  $\varphi(\bar{x})$  is  $\exists y : \psi(\bar{x}, y)$ . By symmetry, we only need to prove the  $(\Rightarrow)$  direction of  $(\dagger)$ . Suppose  $M \models \exists y : \psi(\bar{a}, y)$ . Take some  $a' \in M$  such that  $M \models \psi(\bar{a}, a')$ . By the “forward” condition, there is  $g \in \mathcal{F}$  extending  $f$  with  $a' \in \text{dom}(g)$ . By induction,  $M \models \psi(\bar{a}, a') \implies N \models \psi(g(\bar{a}), g(a'))$ , and then  $N \models \varphi(g(\bar{a}))$ . But  $g(\bar{a}) = f(\bar{a})$ , so  $N \models \varphi(f(\bar{a}))$  as desired.  $\square$

**Corollary 3.7.7.** *Let  $M, N$  be two structures. If there is a non-empty back-and-forth system between  $M$  and  $N$ , then  $M \equiv N$ .*

**Definition 3.7.8.** A linear order  $(M, \leq)$  is *dense* if for any  $x < y$  there is  $z$  with  $x < z < y$ .

**Definition 3.7.9.** DLO is the theory of non-empty dense linear orders without endpoints. That is,  $(M, \leq) \models \text{DLO}$  if  $(M, \leq)$  is a non-empty dense linear order without a greatest or least element.

**Theorem 3.7.10.** *Let  $M$  and  $N$  be models of DLO and let  $\mathcal{F}$  be the class of finite partial isomorphisms  $f : A \rightarrow B$  between  $M$  and  $N$ . Then  $\mathcal{F}$  is a back-and-forth system.*

*Proof.* We prove the forward condition; the backward condition follows by symmetry. Fix a finite partial isomorphism  $f : A \rightarrow B$  and an element  $a' \in M$ . We must extend  $f$  to a larger finite partial isomorphism  $f'$  containing  $a'$ . Enumerate the elements of  $A$ :

$$\begin{aligned} A &= \{a_1, \dots, a_n\} \\ a_1 &< a_2 < \dots < a_n \end{aligned}$$

Let  $b_i = f(a_i)$ . Then

$$\begin{aligned} B &= \{b_1, \dots, b_n\} \\ b_1 &< b_2 < \dots < b_n. \end{aligned}$$

There are five cases.

1.  $n = 0$ , so that  $A, B$  are empty. Take any  $b' \in N$ . ( $N$  is non-empty.) Then  $f' : \{a'\} \rightarrow \{b'\}$  extends  $f$  and is defined at  $a'$ .
2.  $a' = a_i \in A$ . Then we can take  $f' = f$ .

3.  $a' < a_1$ . Take  $b' \in N$  with  $b' < b_1$ . ( $N$  has no minimum.) Then there is a partial isomorphism

$$f' : \{a', a_1, \dots, a_n\} \rightarrow \{b', b_1, \dots, b_n\}$$

extending  $f$  and sending  $a'$  to  $b'$ .

4.  $a' > a_n$ . This case is similar to the previous case.
5.  $a_i < a' < a_{i+1}$  for some  $i$ . Take  $b' \in N$  with  $b_i < b' < b_{i+1}$ . ( $N$  is densely ordered.) Then there is a partial isomorphism

$$f' : \{a_1, \dots, a_i, a', a_{i+1}, \dots, a_n\} \rightarrow \{b_1, \dots, b_i, b', b_{i+1}, \dots, b_n\}$$

extending  $f$  and sending  $a'$  to  $b'$ . □

**Corollary 3.7.11.** 1. *The theory DLO is complete—any two models  $M, N$  are elementarily equivalent.*

2. *If  $M \models \text{DLO}$ , then  $\text{Th}(M)$  is decidable.*

For example, the complete theory of  $(\mathbb{Q}, \leq)$  is decidable.

### 3.8 Definable sets and functions

Fix a language  $\mathcal{L}$  and an  $\mathcal{L}$ -structure  $M$ . If  $\varphi(x_1, \dots, x_n)$  is an  $\mathcal{L}$ -formula, then  $\varphi(M^n)$  or  $\varphi(M)$  denotes the set *defined by*  $\varphi$ :

$$\varphi(M^n) := \{\bar{a} \in M^n : M \models \varphi(\bar{a})\}.$$

More generally, if  $\varphi(x_1, \dots, x_n; y_1, \dots, y_m)$  is an  $\mathcal{L}$ -formula and  $\bar{b} \in M^m$ , then  $\varphi(M^n, \bar{b})$  or  $\varphi(M, \bar{b})$  denotes the set defined by  $\varphi(\bar{x}, \bar{b})$ :

$$\varphi(M^n, \bar{b}) := \{\bar{a} \in M^n : M \models \varphi(\bar{a}, \bar{b})\}.$$

Sets of this form are called *definable sets*. A set  $D \subseteq M^n$  is *A-definable* or *definable over A* if  $D = \varphi(M^n, \bar{b})$  with  $\bar{b} \in A^m$ . That is, an  $A$ -definable set is a set defined by an  $\mathcal{L}(A)$ -formula. Definable sets are the same thing as  $M$ -definable sets. A *0-definable set* is a  $\emptyset$ -definable set. If  $D, E$  are definable sets, a function  $f : D \rightarrow E$  is *definable* if the graph  $\Gamma(f) = \{(x, y) \in D \times E : f(x) = y\}$  is a definable set.  $A$ -definable functions are defined similarly.

**Remark 3.8.1.** If  $t(x_1, \dots, x_n)$  is an  $\mathcal{L}(M)$ -term, then the function  $t^M : M^n \rightarrow M$  is definable, defined by the formula  $t(\bar{x}) = y$ .

**Theorem 3.8.2.** If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are definable, then  $g \circ f : X \rightarrow Z$  is definable.

*Proof.* Let  $\varphi(\bar{x}, \bar{y})$  be an  $\mathcal{L}(M)$ -formula defining  $f$ , meaning that

$$f(\bar{a}) = \bar{b} \iff M \models \varphi(\bar{a}, \bar{b}).$$

Similarly, let  $\psi(\bar{y}, \bar{z})$  define  $g$ . Let  $\theta(\bar{x}, \bar{z})$  be

$$\exists \bar{y} (\varphi(\bar{x}, \bar{y}) \wedge \psi(\bar{y}, \bar{z})).$$

Then  $\theta$  defines  $g \circ f$ :

$$\begin{aligned} M \models \theta(\bar{a}, \bar{c}) &\iff \exists \bar{b} (M \models \varphi(\bar{a}, \bar{b}) \text{ and } M \models \psi(\bar{b}, \bar{c})) \\ &\iff \exists \bar{b} (f(\bar{a}) = \bar{b} \text{ and } g(\bar{b}) = \bar{c}) \\ &\iff g(f(\bar{a})) = \bar{c}. \end{aligned} \quad \square$$

### 3.9 Incompleteness of Peano Arithmetic

Consider  $\mathbb{N}$  as a structure  $(\mathbb{N}, +, \cdot, 0, 1)$ .

**Remark 3.9.1.** The relation  $x \leq y$  is definable, defined by

$$\exists z : z + x = y.$$

**Remark 3.9.2.** The function  $|x - y|$  is definable.

*Proof.*  $|x - y| = z$  if and only if  $x = y + z \vee y = x + z$ .  $\square$

**Definition 3.9.3.** If  $n > 0$  and  $a \in \mathbb{N}$ , then  $a \bmod n$  denotes the unique  $x \in \{0, 1, \dots, n-1\}$  such that  $x \equiv a \pmod{n}$ .

**Lemma 3.9.4.** The function  $x \bmod y$  is definable.

*Proof.*  $(x \bmod y) = z$  if and only if

$$z + 1 \leq y \wedge (\exists k : k \cdot y + z = x). \quad \square$$

**Definition 3.9.5.** For  $a, b, x \in \mathbb{N}$ ,  $\gamma(a, b, x) = (a \bmod ((x + 1)b + 1))$ .

**Lemma 3.9.6.** For any finite sequence  $c_0, c_1, \dots, c_n \in \mathbb{N}$ , there are  $a, b \in \mathbb{N}$  such that  $c_i = \gamma(a, b, i)$  for  $0 \leq i \leq n$ .

*Proof.* Take  $m > \max(c_0, \dots, c_n, n)$ , and let  $b = m! = 1 \cdot 2 \cdot 3 \cdot 4 \cdots m$ .

*Claim.* The numbers  $\{b + 1, 2b + 1, 3b + 1, \dots, (n + 1)b + 1\}$  are pairwise coprime.

*Proof.* Suppose a prime number  $p$  divides both  $ib + 1$  and  $jb + 1$  for some  $1 \leq i < j \leq n + 1$ . Then  $p$  divides the difference  $(j - i)b$ , and one of two things happens:

- $p$  divides  $b$ .
- $p$  divides  $j - i$ . Then  $p \leq j - i \leq n \leq m$ , so  $p$  divides  $m! = b$ .

Either way,  $p$  divides  $b$ , and so

$$0 \equiv ib + 1 \equiv i0 + 1 = 1 \pmod{p},$$

a contradiction. □<sub>Claim</sub>

By the Chinese remainder theorem, there is an  $a$  such that

$$a \equiv c_i \pmod{(i + 1)b + 1}$$

for each  $0 \leq i \leq n$ . But  $c_i \leq m \leq b < (i + 1)b + 1$ , and so

$$c_i = (a \bmod ((i + 1)b + 1)) = \gamma(a, b, i). \quad \square$$

**Theorem 3.9.7.** Let  $f(n)$  be the  $n$ th Fibonacci number  $(1, 1, 2, 3, 5, 8, 13, \dots)$

$$\begin{aligned} f(0) &= f(1) = 1 \\ f(n + 2) &= f(n) + f(n + 1) \end{aligned}$$

Then the function  $f : \mathbb{N} \rightarrow \mathbb{N}$  is definable.

*Proof.* For fixed  $n$  and  $k$ , the following are equivalent:

1.  $f(n) = k$ .

2. There are  $c_0, \dots, c_n$  such that the following hold:

$$\begin{aligned} c_0 &= 1 \\ c_1 &= 1 \\ c_i &= c_{i-1} + c_{i-2} \text{ for all } 2 \leq i \leq n. \end{aligned}$$

3. There are  $a$  and  $b$  such that the following hold:

$$\begin{aligned} \gamma(a, b, 0) &= 1 \\ \gamma(a, b, 1) &= 1 \\ \gamma(a, b, i) &= \gamma(a, b, i-1) + \gamma(a, b, i-2) \text{ for all } 2 \leq i \leq n. \end{aligned}$$

Indeed, (2) and (3) are equivalent by Lemma 3.9.6. Condition (3) is easily expressed by a formula.  $\square$

**Theorem 3.9.8.** *If  $f : \mathbb{N}^k \rightarrow \mathbb{N}$  is computable, then  $f$  is definable.*

*Proof sketch.* The only hard thing to check is primitive recursion, which is handled by the method of Theorem 3.9.7.  $\square$

**Corollary 3.9.9.** *If  $S \subseteq \mathbb{N}^k$  is computable, then  $S$  is definable.*

**Theorem 3.9.10.**  *$\text{Th}(\mathbb{N})$  is incomputable.*

*Proof.* Fix a reasonable enumeration of Turing machines. Let  $S \subseteq \mathbb{N}^2$  be the set of pairs  $(n, k)$  such that the  $n$ th Turing machine halts within  $k$  steps. By Corollary 3.9.9,  $S$  is definable, defined by an  $\mathcal{L}(\mathbb{N})$ -formula  $\varphi(x, y, n_1, n_2, \dots)$ . Replacing  $n_i$  with  $\underbrace{1 + 1 + \dots + 1}_{n_i \text{ times}}$ , we see that  $S$  is  $\emptyset$ -definable, defined by an  $\mathcal{L}$ -formula  $\varphi(x, y)$ . Then for any  $n \in \mathbb{N}$ ,

$$\mathbb{N} \models \exists y \varphi(\underbrace{1 + 1 + \dots + 1}_{n \text{ times}}, y) \iff (\text{the } n\text{th Turing machine halts}).$$

Therefore the halting problem reduces to  $\text{Th}(\mathbb{N})$ .  $\square$

Let  $\mathcal{L}$  be the language  $\{+, \cdot, 0, 1\}$ . *Peano Arithmetic* (PA) consists of the following axioms:

$$\begin{aligned} & \forall x : x + 1 \neq 0 \\ & \forall x : (x = 0 \vee (\exists y : x = y + 1)) \\ & \forall x, y : (x + 1 = y + 1 \rightarrow x = y) \\ & \forall x : x + 0 = x \\ & \forall x, y : x + (y + 1) = (x + y) + 1 \\ & \forall x : x \cdot 0 = 0 \\ & \forall x, y : x \cdot (y + 1) = (x \cdot y) + x \end{aligned}$$

as well as the infinite set of axioms  $\{Ind_\varphi : \varphi(x, y_1, \dots, y_n) \text{ an } \mathcal{L}\text{-formula}\}$ , where  $Ind_\varphi$  is the induction axiom

$$\forall \bar{y} (\varphi(0, \bar{y}) \wedge (\forall x \varphi(x, \bar{y}) \rightarrow \varphi(x + 1, \bar{y})) \rightarrow \forall x \varphi(x, \bar{y})).$$

For example,  $(\mathbb{N}, +, \cdot, 0, 1) \models \text{PA}$ .

**Remark 3.9.11.** The induction schema says that if  $D \subseteq M^1$  is a definable set such that  $0 \in D$  and  $x \in D \implies x + 1 \in D$ , then  $D$  contains every  $x \in M$ .

PA is computably axiomatized, so Theorems 3.6.6 and 3.9.10 yield the following:

**Corollary 3.9.12.** *Peano arithmetic is incomplete.*

Then Theorem 3.6.4 gives the following:

**Corollary 3.9.13.** *There is a model  $M \models \text{PA}$  with  $M \not\cong \mathbb{N}$ .*



# Chapter 4

## The compactness theorem

### 4.1 A temporary convention

When we defined formulas (Definition 3.1.7), we chose the following logical operators to be fundamental:

$$\wedge, \vee, \top, \perp, \exists, \forall, \neg.$$

In contrast, notions like  $\leftrightarrow$  and  $\exists!$  are defined in terms the fundamental operators (see Section 3.3). For example,  $\varphi \rightarrow \psi$  is an abbreviation for  $\neg\varphi \vee \psi$ .

In this chapter, we temporarily switch conventions, regarding

$$\wedge, \top, \exists, \neg,$$

as fundamental logical operators, and  $\vee, \forall, \perp$  as derived notions like  $\leftrightarrow$  or  $\exists!$ . Specifically:

$$\begin{aligned}\varphi \vee \psi &:= \neg(\neg\varphi \wedge \neg\psi) \\ \perp &:= \neg\top \\ \exists x \varphi(x) &:= \neg\forall x \neg\varphi(x).\end{aligned}$$

This convention simplifies the proofs of the compactness and completeness theorems, without impacting the conclusions of these theorems.

## 4.2 Premodels

A  $\mathcal{L}$ -prestructure is a pair  $(M, \approx)$  where  $M$  is an  $\mathcal{L}$ -structure and  $\approx$  is an equivalence relation on  $M$  such that the following hold:

1. If  $f$  is a  $k$ -ary function symbol and  $a_i \approx b_i$  for  $i = 1, \dots, k$ , then

$$f(a_1, \dots, a_k) \approx f(b_1, \dots, b_k).$$

2. If  $R$  is a  $k$ -ary relation symbol and  $a_i \approx b_i$  for  $i = 1, \dots, k$ , then

$$R(a_1, \dots, a_k) \iff R(b_1, \dots, b_k).$$

The first condition says that  $\approx$  is a congruence with respect to the function symbols.

**Definition 4.2.1.** If  $(M, \approx)$  is an  $\mathcal{L}$ -prestructure, then  $M/\approx$  is the following  $\mathcal{L}$ -structure:

1. The underlying set is  $M/\approx = \{[a] : a \in M\}$ , where  $[a] = \{b \in M : b \approx a\}$ .
2. If  $f$  is a  $k$ -ary function symbol, then

$$f([a_1], \dots, [a_k]) = [f(a_1, \dots, a_k)]$$

3. If  $R$  is a  $k$ -ary relation symbol, then

$$R([a_1], \dots, [a_n]) \iff R(a_1, \dots, a_n).$$

This is well-defined by definition of prestructure (using Theorem 2.3.1).

If  $M$  is an  $\mathcal{L}$ -prestructure, if  $\varphi(x_1, \dots, x_n)$  is an  $\mathcal{L}$ -sentence, and  $a_1, \dots, a_n \in M$ , then we define  $M \models \varphi(a_1, \dots, a_n)$  exactly as for ordinary structures, except that

$$M \models t(\bar{a}) = s(\bar{a}) \iff t^M(\bar{a}) \approx s^M(\bar{a}).$$

In other words, we treat the symbol “=” as a non-logical symbol whose interpretation is  $\approx$ .

**Definition 4.2.2.** A *premodel* of  $T$  is an  $\mathcal{L}$ -prestructure satisfying  $T$ .

**Theorem 4.2.3.** *If  $(M, \approx)$  is an  $\mathcal{L}$ -prestructure, then*

$$M \models \varphi(a_1, \dots, a_n) \iff (M/\approx) \models \varphi([a_1], \dots, [a_n]).$$

*Proof.* By induction on the complexity of  $\varphi$ . For example, if  $\varphi(x_1, \dots, x_n)$  is  $\exists y : \psi(\bar{x}, y)$ , then

$$\begin{aligned} M \models \varphi(a_1, \dots, a_n) &\iff \exists b \in M : M \models \psi(\bar{a}, b) \\ &\iff \exists b \in M : (M/\approx) \models \psi([a_1], \dots, [a_n], [b]) \\ &\iff \exists c \in (M/\approx) : (M/\approx) \models \psi([a_1], \dots, [a_n], c) \\ &\iff (M/\approx) \models \varphi([a_1], \dots, [a_n]), \end{aligned}$$

as every element of  $M/\approx$  has the form  $[b]$  for some  $b \in M/\approx$ .  $\square$

**Corollary 4.2.4.** *If  $(M, \approx)$  is a premodel of  $T$ , then  $M/\approx$  is a model of  $T$ .*

## 4.3 The witness property

**Definition 4.3.1.** Let  $T$  be an  $\mathcal{L}$ -theory.

- $T$  is *satisfiable* if  $T$  has a model.
- $T$  is *finitely satisfiable* if every  $T_0 \subseteq_f T$  is satisfiable.
- $T$  is *complete* if for every  $\mathcal{L}$ -sentence  $\varphi$ ,  $\varphi \in T$  or  $\neg\varphi \in T$ .
- $T$  has the *witness property* if whenever  $(\exists x \varphi(x)) \in T$ , there is a closed term  $t$  such that  $\varphi(t) \in T$ .

Recall from §3.6 that the notation  $T \models \varphi$  means that every model of  $T$  satisfies  $\varphi$ .

**Lemma 4.3.2.** *Suppose an  $\mathcal{L}$ -theory  $T$  is finitely satisfiable and complete,  $T_0 \subseteq_f T$ , and  $T_0 \models \varphi$ . Then  $\varphi \in T$ .*

*Proof.* Otherwise,  $\neg\varphi \in T$ . Then  $T_0 \cup \{\neg\varphi\}$  shows  $T$  is not finitely satisfiable.  $\square$

**Lemma 4.3.3.** *Suppose an  $\mathcal{L}$ -theory  $T$  is finitely satisfiable and complete and has the witness property. Let  $\varphi, \psi$  be sentences and  $\theta(x)$  be a formula.*

1.  $\neg\varphi \in T \iff \varphi \notin T$ .
2.  $\varphi \wedge \psi \in T \iff (\varphi \in T \text{ and } \psi \in T)$
3.  $\top \in T$ .
4.  $(\exists x \theta(x)) \in T$  if and only if there is a closed term  $t$  such that  $\theta(t) \in T$ .

*Proof.* 1. Either  $\varphi \in T$  or  $\neg\varphi \in T$  by completeness. If both  $\varphi$  and  $\neg\varphi$  are in  $T$ , then  $T$  is not finitely satisfiable.

2. Both directions hold by Lemma 4.3.2, because

$$\begin{aligned} \{\varphi, \psi\} &\models \varphi \wedge \psi \\ \varphi \wedge \psi &\models \varphi \text{ and } \varphi \wedge \psi \models \psi. \end{aligned}$$

3. Lemma 4.3.2 again.

4. The  $\Leftarrow$  direction holds by Lemma 4.3.2, and the  $\Rightarrow$  direction holds because  $T$  has the witness property.  $\square$

**Theorem 4.3.4.** *Let  $T$  be finitely satisfiable and complete, with the witness property. Then  $T$  has a model.*

*Proof.* By Corollary 4.2.4, it suffices to build a premodel  $(M, \approx) \models T$ . The underlying set of  $M$  will be the set of closed  $\mathcal{L}$ -terms. If  $f$  is a  $k$ -ary function symbol and  $t_1, \dots, t_k \in M$ , let  $f^M(t_1, \dots, t_k) = f(t_1, \dots, t_k)$ . If  $R$  is a  $k$ -ary function symbol and  $t_1, \dots, t_k \in M$ , let  $R^M(t_1, \dots, t_k)$  hold iff  $R(t_1, \dots, t_k) \in T$ . Finally, let  $t \approx s$  hold iff  $(t = s) \in T$ .

*Claim.* The relation  $\approx$  is transitive.

*Proof.* Suppose  $\{s = t, t = u\} \subseteq T$ . Note that  $\{s = t, t = u\} \models \{s = u\}$ . (Any structure which satisfies  $s = t$  and  $t = u$  must also satisfy  $s = u$ .) By Lemma 4.3.2,  $(s = u) \in T$ .  $\square_{\text{Claim}}$

Similar arguments show that  $\approx$  is symmetric, reflexive, and respects the function and relation symbols. Thus  $M$  is an  $\mathcal{L}$ -prestructure.

*Claim.* If  $\varphi$  is an  $\mathcal{L}$ -sentence, then  $M \models \varphi \iff \varphi \in T$ .

*Proof.* Proceed by induction on  $\varphi$ :

1. If  $\varphi$  is an atomic formula, then  $M \models \varphi \iff \varphi \in T$  by choice of the  $\mathcal{L}$ -structure:
  - (a)  $M \models R(t_1, \dots, t_k) \iff R(t_1, \dots, t_k) \in T$  by choice of  $R^M$ .
  - (b)  $M \models t = s \iff (t = s) \in T$  by choice of  $\approx$ .
  - (c)  $M \models f(t_1, \dots, t_k) = s \iff (f(t_1, \dots, t_k) = s) \in T$  by choice of  $f^M$ .
2. The logical operators  $\neg, \top, \wedge, \exists$  work correctly by Lemma 4.3.3 and induction. For example,

$$\begin{aligned}
 M \models \exists x \varphi(x) &\iff \exists t \in M : M \models \varphi(t) \\
 &\iff \exists t \in M : \varphi(t) \in T \quad \text{by induction} \\
 &\iff (\exists x \varphi(x)) \in T \quad \text{by Lemma 4.3.3(4).} \quad \square_{\text{Claim}}
 \end{aligned}$$

By the Claim,  $M$  is a premodel of  $T$ . By Corollary 4.2.4, the quotient  $M/\approx$  is a model of  $T$ .  $\square$

**Definition 4.3.5.** Suppose  $T$  is finitely satisfiable and complete and has the witness property. The *canonical model* of  $T$  is the model constructed in the proof of Theorem 4.3.4.

The canonical model of  $T$  is characterized up to isomorphism by the fact that every element  $a \in M$  is named by a closed term  $t$ , in the sense that  $a = t^M$ .

**Definition 4.3.6.** Let  $\mathcal{C}$  be a collection of closed  $\mathcal{L}$ -terms. An  $\mathcal{L}$ -theory *has the witness property over  $\mathcal{C}$*  if whenever  $(\exists x \varphi(x)) \in T$ , there is a term  $t \in \mathcal{C}$  such that  $\varphi(t) \in T$ .

**Theorem 4.3.7.** Let  $T$  be a finitely satisfiable, complete theory with the witness property over  $\mathcal{C}$ , and let  $M$  be the canonical model of  $T$ . Then every element of  $M$  is named by a term in  $\mathcal{C}$ .

*Proof.* Fix  $a \in M$ . Then  $a = t^M$  for some term  $t$ . The  $\mathcal{L}$ -sentence

$$\exists x x = t$$

holds in any  $\mathcal{L}$ -structure, so it must be in  $T$  by Lemma 4.3.2. By the witness property over  $\mathcal{C}$ , there is  $t' \in \mathcal{C}$  such that  $(t' = t) \in T$ . Then  $M \models t' = t$ , so  $a = t^M = (t')^M$ , and  $a$  is named by a term in  $\mathcal{C}$ .  $\square$

## 4.4 Compactness via Henkin's method

**Theorem 4.4.1.** *Let  $I$  be a linear order and  $\{T_i\}_{i \in I}$  be a chain of finitely satisfiable theories, meaning that  $i < i' \implies T_i \subseteq T_{i'}$ . Then the union  $T = \bigcup_{i \in I} T_i$  is finitely satisfiable.*

*Proof.* Suppose  $T_0 \subseteq_f T$ . Let  $T_0 = \{\varphi_1, \dots, \varphi_n\}$ . For each  $j \leq n$ , we have  $\varphi_j \in T = \bigcup_i T_i$ , so there is some  $i_j \in I$  with  $\varphi_j \in T_{i_j}$ . Let  $\ell = \max(i_1, \dots, i_n) \in I$ . Then for each  $j \leq n$ , we have  $\varphi_j \in T_{i_j} \subseteq T_\ell$ . Thus  $T_0 \subseteq_f T_\ell$ , and  $T_0$  is satisfiable because  $T_\ell$  is finitely satisfiable.  $\square$

**Lemma 4.4.2.** *If  $T$  is a finitely satisfiable  $\mathcal{L}$ -theory, then there is a complete, finitely satisfiable  $\mathcal{L}$ -theory  $T' \supseteq T$ .*

*Proof.* By Zorn's lemma and Theorem 4.4.1 there is a  $T' \supseteq T$  which is maximal among finitely satisfiable  $\mathcal{L}$ -theories. We claim  $T'$  is complete. Otherwise there is a sentence  $\varphi$  with  $\varphi \notin T'$  and  $\neg\varphi \notin T'$ . By maximality,  $T' \cup \{\varphi\}$  is not finitely satisfiable, so there is  $T_1 \subseteq_f T$  with  $T_1 \models \neg\varphi$ . Similarly,  $\neg\varphi \notin T'$  implies that there is  $T_2 \subseteq_f T$  with  $T_2 \models \varphi$ . Then  $T_1 \cup T_2$  is not satisfiable, a contradiction.  $\square$

**Lemma 4.4.3.** *Let  $T$  be a finitely satisfiable  $\mathcal{L}$ -theory containing a sentence  $\exists x \varphi(x)$ . Let  $\mathcal{L}' = \mathcal{L} \cup \{c\}$  where  $c$  is a new constant symbol. Then  $T \cup \{\varphi(c)\}$  is a finitely satisfiable  $\mathcal{L}'$ -theory.*

*Proof.* Otherwise,  $T_0 \cup \{\varphi(c)\}$  is unsatisfiable for some  $T_0 \subseteq_f T$ . Take  $M \models T_0 \cup \{\exists x \varphi(x)\}$  and take  $b \in M$  such that  $M \models \varphi(b)$ . Expand the  $\mathcal{L}$ -structure  $M$  to an  $\mathcal{L}'$ -structure by interpreting  $c$  as  $b$ . Then  $M \models T_0 \cup \{\varphi(c)\}$ , contradicting the choice of  $T_0$ .  $\square$

**Lemma 4.4.4.** *Let  $T$  be a finitely satisfiable  $\mathcal{L}$ -theory. There is a language  $\mathcal{L}' \supseteq \mathcal{L}$  and an  $\mathcal{L}'$ -theory  $T' \supseteq T$  that is finitely satisfiable and complete, and has the witness property.*

*Proof.* Build increasing chains

$$\begin{aligned} \mathcal{L}_0 &\subseteq \mathcal{L}_1 \subseteq \dots \\ T_0 &\subseteq T_1 \subseteq \dots \end{aligned}$$

where  $T_i$  is a finitely satisfiable  $\mathcal{L}_i$ -theory as follows:

1.  $\mathcal{L}_0 = \mathcal{L}$  and  $T_0 = T$ .
2. If  $n > 0$  and  $n$  is odd, then  $\mathcal{L}_n = \mathcal{L}_{n-1}$  and  $T_n$  is a completion of  $T_{n-1}$  from Lemma 4.4.2.
3. If  $n > 0$  and  $n$  is even, let  $\{\varphi_i(x) : i \in I\}$  enumerate the  $\mathcal{L}_{n-1}$ -formulas such that  $(\exists x \varphi(x)) \in T_{n-1}$ . Let  $\mathcal{L}_n = \mathcal{L}_{n-1} \cup \{c_i : i \in I\}$  where the  $c_i$  are new constant symbols. Let  $T_n = T_{n-1} \cup \{\varphi_i(c_i) : i \in I\}$ . Then  $T_n$  is finitely satisfiable by Lemma 4.4.3.

Finally, take  $\mathcal{L}' = \bigcup_n \mathcal{L}_n$  and  $T' = \bigcup_n T_n$ . Then  $T'$  is finitely satisfiable because each  $T_i$  is,  $T'$  is complete because of the odd-numbered steps, and  $T'$  has the witness property because of the even-numbered steps.  $\square$

**Theorem 4.4.5** (Compactness). *If  $T$  is finitely satisfiable, then  $T$  has a model.*

*Proof.* By Lemma 4.4.4, there is a language  $\mathcal{L}' \supseteq \mathcal{L}$  and an  $\mathcal{L}'$ -theory  $T' \supseteq T$  such that  $T'$  is finitely satisfiable and complete and has the witness property. By Theorem 4.3.4 there is an  $\mathcal{L}'$ -structure  $M$  satisfying  $T'$ , and therefore  $T$ .  $\square$





# Chapter 5

## Categoricity

### 5.1 A useful fact around finite satisfiability

Let  $T$  be a set of sentences. Suppose  $T$  is closed under finite conjunctions, in the sense that

$$\varphi_1, \dots, \varphi_n \in T \implies \bigwedge_{i=1}^n \varphi_i \in T.$$

Then  $T$  is finitely satisfiable if and only if every  $\varphi \in T$  is satisfiable, because any finite subset  $\{\varphi_1, \dots, \varphi_n\} \subseteq T$  is equivalent to a single formula  $\bigwedge_{i=1}^n \varphi_i$  in  $T$ .

More generally, suppose  $T_1, T_2$  are two sets of sentences, and  $T_1$  is closed under finite conjunctions. Then the following are equivalent:

- $T_1 \cup T_2$  is finitely satisfiable.
- For every  $\varphi \in T_1$  and finite subset  $T'_2 \subseteq_f T_2$ , the theory  $\{\varphi\} \cup \{T'_2\}$  is satisfiable.

We will use this fact implicitly in what follows.

### 5.2 Diagrams and elementary amalgamation

Let  $M$  be an  $\mathcal{L}$ -structure and  $A$  be a subset of  $M$ . The language  $\mathcal{L}(A)$  is obtained by adding each element of  $A$  as a new constant symbol. We regard

$M$  as an  $\mathcal{L}(A)$ -structure by interpreting each new constant symbol as the corresponding element of  $A$ :

$$a^M = a.$$

**Definition 5.2.1.** The *elementary diagram* of  $M$ , written  $\text{eldiag}(M)$ , is the set of  $\mathcal{L}(M)$ -sentences  $\varphi$  such that  $M \models \varphi$ . The *diagram* of  $M$  is the set of atomic and negated atomic formulas in  $\text{eldiag}(M)$ .

A model of  $\text{diag}(M)$  consists of an  $\mathcal{L}$ -structure  $N$  and a map

$$\begin{aligned} M &\rightarrow N \\ a &\mapsto a^N \end{aligned}$$

such that for any atomic  $\mathcal{L}$ -formula  $\varphi(x_1, \dots, x_n)$  and any  $\bar{a} \in M^n$ ,

$$\begin{aligned} M \models \varphi(a_1, \dots, a_n) &\implies N \models \varphi(a_1^N, \dots, a_n^N) \\ M \models \neg\varphi(a_1, \dots, a_n) &\implies N \models \neg\varphi(a_1^N, \dots, a_n^N). \end{aligned}$$

This means that  $a \mapsto a^N$  is an embedding.

Therefore, a model of  $\text{diag}(M)$  is the same thing as an  $\mathcal{L}$ -structure  $N$  with an embedding  $f : M \rightarrow N$ . In particular, if  $N$  is an extension of  $M$ , then  $N$  is a model of  $\text{diag}(M)$  in a natural way, interpreting  $a \in M$  as  $a \in N$ . Every model of  $\text{diag}(M)$  is isomorphic as an  $\mathcal{L}(M)$ -structure to an extension of  $M$ .

Similarly, a model of  $\text{eldiag}(M)$  is the same thing as an  $\mathcal{L}$ -structure  $N$  with an elementary embedding  $f : M \rightarrow N$ . Any elementary extension of  $M$  is a model of  $\text{eldiag}(M)$ , and every model of  $\text{eldiag}(M)$  is isomorphic as an  $\mathcal{L}(M)$ -structure to an elementary extension of  $M$ .

**Theorem 5.2.2** (Elementary amalgamation). *If  $M_1$  and  $M_2$  are elementarily equivalent, then there is a structure  $N$  and elementary embeddings  $M_1 \rightarrow N$  and  $M_2 \rightarrow N$ .*

We first prove a lemma. Recall the notion of *expansion* from Definition 3.1.5.

**Lemma 5.2.3.** *If  $M_1 \equiv M_2$  and  $\varphi(\bar{c}) \in \text{eldiag}(M_2)$ , then there is an  $\mathcal{L}(M_2)$ -structure  $N$  expanding the  $\mathcal{L}$ -structure  $M_1$  such that  $N \models \varphi$ .*

*Proof.* The fact that  $M_2 \models \varphi(\bar{c})$  implies that  $M_2 \models \exists \bar{x} \varphi(\bar{x})$  and then  $M_1 \models \exists \bar{x} \varphi(\bar{x})$ . Take  $\bar{a}$  in  $M_1$  such that  $M_1 \models \varphi(\bar{a})$ . Expand  $M_1$  to an  $\mathcal{L}(M_2)$ -structure  $N$  by interpreting the symbols  $c_1, \dots, c_n$  as  $a_1, \dots, a_n$ , respectively. Then  $N \models \varphi(\bar{c})$  because  $\bar{c}^N = \bar{a}$  and  $M_1 \models \varphi(\bar{a})$ .  $\square$

Now we prove Theorem 5.2.2.

*Proof.* It suffices to show that  $\text{eldiag}(M_1) \cup \text{eldiag}(M_2)$  is consistent. Otherwise, by compactness, there is a tuple  $\bar{c}$  in  $M_2$  and formula  $\varphi(\bar{c}) \in \text{eldiag}(M_2)$  such that  $\text{eldiag}(M_1) \cup \{\varphi(\bar{c})\}$  is inconsistent. But  $M_1 \models \text{eldiag}(M_1)$ , and Lemma 5.2.3 gives an expansion of  $M_1$  satisfying  $\{\varphi(\bar{c})\}$ , and so  $\text{eldiag}(M_1) \cup \{\varphi(\bar{c})\}$  is consistent.  $\square$

## 5.3 Tarski-Vaught

**Theorem 5.3.1.** *Let  $M$  be a structure and  $A$  be a subset. Then  $A \preceq M$  iff the following holds: for every non-empty  $A$ -definable  $D \subseteq M$ , we have  $D \cap A \neq \emptyset$ .*

This condition is called the *Tarski-Vaught criterion*.

*Proof.* First suppose  $A \preceq M$ . Suppose  $D$  is  $A$ -definable and non-empty. Write  $D$  as  $\varphi(M, \bar{a})$  for some  $\mathcal{L}(A)$ -formula  $\varphi(x, \bar{a})$ . Then

$$M \models \exists x : \varphi(x, \bar{a}) \implies A \models \exists x : \varphi(x, \bar{a})$$

so there is  $b \in A$  with  $A \models \varphi(b, \bar{a})$ , which implies  $M \models \varphi(b, \bar{a})$  as  $A \preceq M$ . Then  $b \in A$  and  $b \in D = \varphi(M, \bar{a})$ .

Conversely, suppose the Tarski-Vaught criterion holds. First we show that  $A$  is a substructure. Suppose  $f$  is a  $k$ -ary function symbol and  $a_1, \dots, a_k \in A$ . The set  $\{f(\bar{a})\}$  is non-empty and  $A$ -definable, so  $A$  intersects it, meaning that  $f(\bar{a}) \in A$ .

Next we show that for any  $\mathcal{L}$ -formula  $\varphi(\bar{x})$  and any  $\bar{a} \in A$ ,

$$M \models \varphi(\bar{a}) \iff A \models \varphi(\bar{a}). \quad (*)$$

We may assume  $\varphi$  makes no use of  $\wedge, \forall, \top, \perp$ . Proceed by induction on the complexity of  $\varphi$ . If  $\varphi$  is atomic or more generally quantifier-free, then  $(*)$  holds because  $A$  is a substructure.

If  $\varphi$  is  $\psi \vee \theta$ , then

$$\begin{aligned} M \models \varphi(\bar{a}) &\iff M \models \psi(\bar{a}) \text{ or } M \models \theta(\bar{a}) \\ &\iff A \models \psi(\bar{a}) \text{ or } A \models \theta(\bar{a}) \\ &\iff A \models \varphi(\bar{a}) \end{aligned}$$

by induction. Finally, if  $\varphi(\bar{x})$  is  $\exists y : \psi(\bar{x}, y)$ , then

$$\begin{aligned} M \models \varphi(\bar{a}) &\iff \exists b \in M : M \models \psi(\bar{a}, b) \\ &\stackrel{\text{TV}}{\iff} \exists b \in A : M \models \psi(\bar{a}, b) \\ &\stackrel{\text{ind}}{\iff} \exists b \in A : A \models \psi(\bar{a}, b) \\ &\iff A \models \varphi(\bar{a}). \end{aligned}$$

The second line uses the Tarski-Vaught criterion, and the third line uses induction.  $\square$

## 5.4 The Löwenheim-Skolem theorem

**Definition 5.4.1.** If  $\mathcal{L}$  is a language, the *size* of  $\mathcal{L}$ , written  $|\mathcal{L}|$ , is  $\aleph_0$  plus the number of symbols in  $\mathcal{L}$ .

If  $\bar{x}$  is a finite tuple of variables, then  $|\mathcal{L}|$  is the number of  $\mathcal{L}$ -formulas  $\varphi(\bar{x})$ .

**Remark 5.4.2.** If  $M$  is a  $\mathcal{L}$ -structure and  $A \subseteq M$ , then the number of  $A$ -definable sets in  $M$  is at most  $|A| + |\mathcal{L}|$ .

**Theorem 5.4.3** (Downward Löwenheim-Skolem theorem). *Let  $M$  be an  $\mathcal{L}$ -structure.*

1. *If  $A \subseteq M$ , there is an elementary substructure  $N \preceq M$  with  $N \supseteq A$  and  $|N| \leq |A| + |\mathcal{L}|$ .*
2. *For any  $\kappa$  with  $|\mathcal{L}| \leq \kappa \leq |M|$ , there is  $N \preceq M$  with  $|N| = \kappa$ .*

*Proof.* 1. Let  $F : \mathfrak{P}(M) \setminus \{\emptyset\} \rightarrow M$  be a function such that  $F(X) \in X$ . Recursively define  $A_0 \subseteq A_1 \subseteq \dots$  by letting  $A_0 = A$  and

$$A_{n+1} = A_n \cup \{F(X) : X \subseteq M \text{ is } A_n\text{-definable and non-empty}\}.$$

Let  $N = \bigcup_{n=0}^{\infty} A_n$ . Then  $A = A_0 \subseteq N$ . By induction on  $n$ , each  $A_n$  has size at most  $|A| + |\mathcal{L}|$ , and therefore  $|N| \leq |A| + |\mathcal{L}|$ . If  $X$  is  $N$ -definable and non-empty, then  $X$  is  $A_n$ -definable for some  $n$ , so  $F(X) \in A_{n+1} \subseteq N$ . Thus  $N$  intersects every  $N$ -definable set, and  $N \preceq M$  by Tarski-Vaught.

2. Take a subset  $A \subseteq M$  with  $|A| = \kappa$ , and take  $N \preceq M$  as in part (1). Then  $|N| \leq |A| + |\mathcal{L}| = \kappa + |\mathcal{L}| = \kappa$ . On the other hand,  $|N| \geq \kappa$  because  $N \supseteq A$ .  $\square$

**Theorem 5.4.4** (Löwenheim-Skolem theorem). *Let  $T$  be an  $\mathcal{L}$ -theory. Suppose  $T$  has an infinite model, or more generally that for every  $n < \omega$ ,  $T$  has a model of size at least  $n$ . Then for any  $\kappa \geq |\mathcal{L}|$ ,  $T$  has a model of size  $\kappa$ .*

*Proof.* Let  $\mathcal{L}' = \mathcal{L} \cup \{c_\alpha : \alpha < \kappa\}$ , where the  $c_\alpha$  are new constant symbols. Let  $\Sigma = \{(c_\alpha \neq c_\beta) : \alpha < \beta < \kappa\}$ .

*Claim.*  $T \cup \Sigma$  is finitely satisfiable.

*Proof.* Suppose  $\Sigma_0 \subseteq_f \Sigma$ . Let  $S$  be the finite set of  $\alpha \in \kappa$  such that  $c_\alpha$  appears in  $\Sigma_0$ . Take a model  $M \models T$  with  $|M| > |S|$ . Expand  $M$  to an  $\mathcal{L}'$ -structure by interpreting the  $c_\alpha$  for  $\alpha \in S$  as distinct elements of  $M$ , and interpreting  $c_\alpha$  for  $\alpha \in \kappa \setminus S$  arbitrarily. Then  $M \models T \cup \Sigma_0$ .  $\square_{\text{Claim}}$

By compactness,  $T \cup \Sigma$  has a model  $M$ . Then the  $c_\alpha^M$  are pairwise distinct, so  $|M| \geq \kappa$ . By downward Löwenheim-Skolem (Theorem 5.4.3), there is an elementary substructure  $N \preceq M$  with  $|N| = \kappa$ . Then  $N \equiv M$  and  $M \models T$ , so  $N \models T$ .  $\square$

**Corollary 5.4.5** (Löwenheim-Skolem theorem, second version). *Let  $M$  be an infinite  $\mathcal{L}$ -structure. For any  $\kappa \geq |\mathcal{L}|$ , there is  $N \equiv M$  with  $|N| = \kappa$ .*

*Proof.* Apply Theorem 5.4.4 to the  $\mathcal{L}$ -theory  $\text{Th}(M)$ . Models of  $\text{Th}(M)$  are elementarily equivalent to  $M$  (Theorem 3.4.1).  $\square$

**Example 5.4.6.** Earlier we saw that Peano Arithmetic has models which are not elementarily equivalent to  $\mathbb{N}$  (Corollary 3.9.13). Using the Löwenheim-Skolem theorem, we can also produce models that are elementarily equivalent to  $\mathbb{N}$  but not isomorphic to it. For example, Corollary 5.4.5 gives  $M \equiv \mathbb{N}$  with  $|M| > \aleph_0$ . Then  $M \models \text{PA}$  but  $M \not\cong \mathbb{N}$ .

**Theorem 5.4.7** (Upward Löwenheim-Skolem theorem). *If  $M$  is an infinite  $\mathcal{L}$ -structure and  $\kappa \geq |M| + |\mathcal{L}|$ , then there is an elementary extension  $N \succeq M$  with  $|N| = \kappa$ .*

*Proof.* As  $\kappa \geq |M| + |\mathcal{L}| = |\mathcal{L}(M)|$ , we can apply the Löwenheim-Skolem theorem (Theorem 5.4.4) to the  $\mathcal{L}(M)$ -theory  $\text{eldiag}(M)$  to get a model  $N \models \text{eldiag}(M)$  with  $|N| = \kappa$ . Moving  $N$  by an isomorphism, we may assume  $N \succeq M$ .  $\square$

## 5.5 Absolute categoricity

**Definition 5.5.1.** A theory  $T$  is *absolutely categorical* if there is a unique model of  $T$ , up to isomorphism.

**Theorem 5.5.2.** *If  $T$  is absolutely categorical, then the unique model  $M$  is finite.*

*Proof.* Take  $\kappa > |M| + |\mathcal{L}|$ . If  $M$  is infinite, then there is a model  $N \models T$  with  $|N| = \kappa$  by Theorem 5.4.4. Then  $M$  and  $N$  are two non-isomorphic models, contradicting categoricity.  $\square$

Conversely, if  $M$  is finite, then  $\text{Th}(M)$  is absolutely categorical. Recall that models of  $\text{Th}(M)$  are the same thing as structures elementarily equivalent to  $M$  (Theorem 3.4.1).

**Theorem 5.5.3.** *If  $M$  is finite and  $M \equiv N$ , then  $M \cong N$ .*

*Proof.* By elementary amalgamation (Theorem 5.2.2) there are elementary embeddings  $M \rightarrow M'$  and  $N \rightarrow M'$  for some structure  $M'$ . Moving  $M$  and  $N$  by isomorphisms, we may assume  $M \preceq M'$  and  $N \preceq M'$ . Let  $n$  be the size of  $M$ . By Lemma 5.5.4 below,  $|M| = |M'| = |N| = n$ . As  $M$  and  $N$  are subsets of  $M'$ , we must have  $M = M' = N$ .  $\square$

**Lemma 5.5.4.** *Suppose  $M \equiv N$ . If  $|M| = n < \infty$ , then  $|N| = n$ . Consequently,  $M$  is finite iff  $N$  is finite.*

*Proof.* Suppose  $|M| = n$ . Let  $\varphi$  be the sentence  $\exists^{=n}x \top$  (see Section 3.3). Then  $\varphi$  says that there are exactly  $n$  elements, so  $M \models \varphi$ . Therefore  $N \models \varphi$  and  $|N| = n$ .  $\square$

## 5.6 $\kappa$ -categoricity

**Definition 5.6.1.** Let  $\kappa$  be an infinite cardinal. A theory  $T$  is  $\kappa$ -categorical if there is a unique model of cardinality  $\kappa$ .

**Theorem 5.6.2** (Łoś-Vaught criterion). *Suppose  $T$  is  $\kappa$ -categorical for some  $\kappa \geq |\mathcal{L}|$ .*

1. *Any two infinite models of  $T$  are elementarily equivalent.*
2. *If all models of  $T$  are infinite, then  $T$  is complete.*

*Proof.* 1. Given infinite models  $M_1, M_2$ , use Löwenheim-Skolem (Corollary 5.4.5) to get  $N_i \equiv M_i$  with  $|N_i| = \kappa$ . By  $\kappa$ -categoricity,  $N_1 \cong N_2$ . Then  $M_1 \equiv N_1 \equiv N_2 \equiv M_2$ .

2. Clear. □

**Lemma 5.6.3.** *Let  $M, N$  be countable structures and let  $\mathcal{F}$  be a back-and-forth system between  $M$  and  $N$ . If  $f_0 \in \mathcal{F}$ , then there is an isomorphism  $f : M \rightarrow N$  extending  $f_0$ .*

*Proof.* Let  $a_1, a_2, \dots$  be an enumeration of  $M$  and  $b_1, b_2, \dots$  be an enumeration of  $N$ . Recursively build an increasing sequence of partial isomorphisms  $f_i : A_i \rightarrow B_i$  in  $\mathcal{F}$  as follows:

- For  $i = 0$ , let  $f_0 : A_0 \rightarrow B_0$  be the given partial isomorphism.
- For  $i = 2k - 1$ , take  $f_i$  to be an extension of  $f_{i-1}$  with  $a_k \in \text{dom}(f_i)$ . Such an  $f_i$  exists by the “forward” condition.
- For  $i = 2k$ , take  $f_i$  to be an extension of  $f_{i-1}$  with  $b_k \in \text{im}(f_i)$ . Such an  $f_i$  exists by the “backward” condition.

Take  $f = \bigcup_{i=0}^{\infty} f_i$ . Then  $f$  is an isomorphism from  $M$  to  $N$ . □

**Theorem 5.6.4.** *Let  $M, N$  be countable structures. If there is a non-empty back-and-forth system between  $M$  and  $N$ , then  $M \cong N$ .*

Recall the theory DLO from Definition 3.7.9. Theorem 3.7.10 gives a non-empty back-and-forth system between any two models of DLO.

**Corollary 5.6.5** (Cantor). *DLO is  $\aleph_0$ -categorical: any two countable models of DLO are isomorphic.*

Any model of DLO is infinite, so the Łoś-Vaught criterion shows that DLO is complete, as we saw earlier (Corollary 3.7.11).

**Theorem 5.6.6.** *DLO is not  $\kappa$ -categorical for  $\kappa > \aleph_0$ .*

*Proof.* If  $A, B$  are linear orders, let  $A \times B$  denote the lexicographic product, the set  $A \times B$  with the lexicographic order

$$\begin{aligned} (a, b) < (a', b') &\iff a < a' \text{ for } a \neq a' \\ (a, b) < (a, b') &\iff b < b'. \end{aligned}$$

Also, let  $A^*$  denote  $A$  with the reversed order.

Let  $M = \kappa \times \mathbb{Q}$ . It is straightforward to see that  $M$  and  $M^*$  are models of DLO, of size  $\kappa$ . We claim that  $M \not\cong M^*$ . Note that for any  $a \in M$ ,

$$\begin{aligned} |\{x \in M : x < a\}| &< \kappa \\ |\{x \in M : x > a\}| &= \kappa. \end{aligned}$$

In  $M^*$ , the reverse properties hold, so  $M \not\cong M^*$ , and DLO is not  $\kappa$ -categorical.  $\square$



# Chapter 6

## Ultraproducts

### 6.1 Ultrafilters

Let  $I$  be a set.

**Definition 6.1.1.** A *filter* on  $I$  is a set  $\mathcal{F} \subseteq \mathfrak{P}(I)$  satisfying the following:

1. If  $X, Y \in \mathcal{F}$ , then  $X \cap Y \in \mathcal{F}$ .
2. If  $X \subseteq Y \subseteq I$  and  $X \in \mathcal{F}$ , then  $Y \in \mathcal{F}$ .
3.  $I \in \mathcal{F}$ .

A filter is *proper* if  $\emptyset \notin \mathcal{F}$ .

**Definition 6.1.2.** A family of sets  $\mathcal{S} \subseteq \mathfrak{P}(I)$  has the *finite intersection property* (FIP) if for any  $n \geq 0$  and  $X_1, \dots, X_n \in \mathcal{S}$ , we have  $\bigcap_{i=1}^n X_i \neq \emptyset$ .

(When  $n = 0$ ,  $\bigcap_{i=1}^n X_i$  is defined to be  $I$ .)

**Remark 6.1.3.** If  $\mathcal{F}$  is a proper filter, then  $\mathcal{F}$  has the FIP, because  $\emptyset \notin \mathcal{F}$ , and  $\mathcal{F}$  is closed under intersection.

**Lemma 6.1.4.** If  $\mathcal{S} \subseteq \mathfrak{P}(I)$  has the FIP, then there is a proper filter  $\mathcal{F} \supseteq \mathcal{S}$ .

*Proof.* Let  $\mathcal{F}$  be the set of  $X \subseteq I$  such that there are  $n \geq 0$  and  $Y_1, \dots, Y_n \in \mathcal{S}$  with  $X \supseteq \bigcap_{i=1}^n Y_i$ . Then  $\mathcal{F}$  is a proper filter containing  $\mathcal{S}$ .  $\square$

**Definition 6.1.5.** An *ultrafilter* on  $I$  is a proper filter  $\mathcal{U}$  such that for any  $X \subseteq I$ ,

$$X \in \mathcal{U} \text{ or } I \setminus X \in \mathcal{U}.$$

**Lemma 6.1.6.** *If  $\mathcal{F}$  is a proper filter on  $I$ , then there is an ultrafilter  $\mathcal{U} \supseteq \mathcal{F}$ .*

*Proof.* By Zorn's lemma, there is a maximal proper filter  $\mathcal{U} \supseteq \mathcal{F}$ . We claim that  $\mathcal{U}$  is an ultrafilter. Otherwise, there is  $X \subseteq I$  with  $X \notin \mathcal{U}$  and  $I \setminus X \notin \mathcal{U}$ . By maximality,  $\mathcal{U} \cup \{X\}$  is not contained in a filter, and does not have FIP. Therefore there are  $Y_1, \dots, Y_n \in \mathcal{U}$  such that  $X \cap \bigcap_{i=1}^n Y_i = \emptyset$ . As  $\mathcal{U}$  is a filter,  $\mathcal{U}$  contains  $Y := \bigcap_{i=1}^n Y_i$ . Then  $X \cap Y = \emptyset$ , so  $Y \subseteq I \setminus X$ .

Applying the same argument to the complement of  $X$ , we get  $Z \in \mathcal{U}$  with  $Z \subseteq X$ . Then  $Y \cap Z \in \mathcal{U}$  because  $\mathcal{U}$  is a filter. However,  $Y \cap Z \subseteq (I \setminus X) \cap X = \emptyset$ , so  $\emptyset \in \mathcal{U}$  contradicting the fact that  $\mathcal{U}$  is proper.  $\square$

**Theorem 6.1.7.** *If  $\mathcal{S} \subseteq \mathfrak{P}(I)$  has the FIP, then  $\mathcal{S}$  is contained in an ultrafilter  $\mathcal{U}$  on  $I$ .*

**Fact 6.1.8.** *Let  $\mathcal{U}$  be an ultrafilter on  $I$ . Let  $\chi : \mathfrak{P}(I) \rightarrow \mathcal{U}$  be the characteristic function of  $\mathcal{U}$ :*

$$\chi(X) = \begin{cases} 1 & X \in \mathcal{U} \\ 0 & X \notin \mathcal{U}. \end{cases}$$

*Then  $\chi$  is a homomorphism of boolean algebras. In fact, ultrafilters on  $I$  correspond bijectively to homomorphisms  $\mathfrak{P}(I) \rightarrow \{0, 1\}$ .*

## 6.2 Ultraproducts

Let  $I$  be a set and let  $\mathcal{U}$  be an ultrafilter on  $I$ . Let  $M_i$  be a *non-empty*  $\mathcal{L}$ -structure for each  $i \in I$ . Let  $P = \prod_{i \in I} M_i$  and let  $\pi_i : P \rightarrow M_i$  be the  $i$ th coordinate projection.

Let  $\mathcal{L}(P)$  be  $\mathcal{L}$  with a new constant symbol added for each  $a \in P$ . Regard  $M_i$  as an  $\mathcal{L}(P)$ -structure by interpreting  $a \in P$  as  $\pi_i(a) \in M_i$ .

For any  $\mathcal{L}(P)$ -sentence  $\varphi$ , let  $\llbracket \varphi \rrbracket = \{i \in I : M_i \models \varphi\}$ . Note the following:

$$\begin{aligned} \llbracket \varphi \wedge \psi \rrbracket &= \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket \\ \llbracket \varphi \vee \psi \rrbracket &= \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket \\ \llbracket \neg \varphi \rrbracket &= I \setminus \llbracket \varphi \rrbracket \end{aligned}$$

**Lemma 6.2.1.** *Let  $T$  be the set of  $\mathcal{L}(P)$ -sentences  $\varphi$  such that  $\llbracket \varphi \rrbracket \in \mathcal{U}$ . Then  $T$  is finitely satisfiable and complete and has the witness property over  $P$  in the sense of Definition 4.3.6.*

*Proof.* For finite satisfiability, suppose  $\varphi_1, \dots, \varphi_n \in T$ . If  $S = \llbracket \bigwedge_{i=1}^n \varphi_i \rrbracket = \bigcap_{i=1}^n \llbracket \varphi_i \rrbracket$ , then  $S \in \mathcal{U}$  so  $S \neq \emptyset$ . Taking  $i \in S$ , we have  $M_i \models \bigwedge_{i=1}^n \varphi_i$ .

For completeness, note that  $\llbracket \varphi \rrbracket$  and  $\llbracket \neg \varphi \rrbracket$  are complementary, so one of them is in  $\mathcal{U}$ , and therefore one of  $\varphi$  and  $\neg \varphi$  is in  $T$ .

For the witness property, suppose  $\exists x \varphi(x)$  is in  $T$ , meaning that  $S := \llbracket \exists x \varphi(x) \rrbracket \in \mathcal{U}$ . Define  $c = (c_i : i \in I) \in P$  as follows:

1. If  $i \in S$ , then  $M_i \models \exists x \varphi(x)$ . Choose  $c_i \in M_i$  so that  $M_i \models \varphi(c_i)$ .
2. If  $i \notin S$ , take any  $c_i \in M_i$ .

Then  $i \in S \implies M_i \models \varphi(c)$ , so  $\llbracket \varphi(c) \rrbracket \supseteq S$ . It follows that  $\llbracket \varphi(c) \rrbracket \in \mathcal{U}$  and  $\varphi(c) \in T$ .  $\square$

By Definition 4.3.5 and Theorem 4.3.7, we can talk about the canonical model of  $T$ —the unique model in which every element is named by an element of  $P$ .

**Definition 6.2.2.** The *ultraproduct*  $\prod_{i \in I}^{\mathcal{U}} M_i$  is the canonical model of  $T$ .

**Theorem 6.2.3.** Let  $P = \prod_{i \in I} M_i$ , let  $M = \prod_{i \in I}^{\mathcal{U}} M_i$ , and let  $[c]$  be the element of  $M$  named by  $c \in P$ .

1. Every element of  $M$  has the form  $[c]$  for some  $c \in P$ .
2.  $[c] = [d]$  if and only if  $\{i \in I : \pi_i(c) = \pi_i(d)\} \in \mathcal{U}$ .
3. Let  $R$  be a  $k$ -ary relation symbol. Then

$$R^M([c_1], \dots, [c_k]) \iff \{i \in I : M_i \models R(\pi_i(c_1), \dots, \pi_i(c_k))\} \in \mathcal{U}.$$

4. If  $f$  is a  $k$ -ary function symbol, then

$$f^M([c_1], \dots, [c_k]) = [f^P(c_1, \dots, c_k)],$$

where  $P$  is given the product structure so that

$$f^P(c_1, \dots, c_k) = (f^{M_i}(\pi_i(c_1), \dots, \pi_i(c_k)) : i \in I).$$

5. If  $\varphi(x_1, \dots, x_n)$  is an  $\mathcal{L}$ -formula, then

$$M \models \varphi([c_1], \dots, [c_k]) \iff \{i \in I : M_i \models \varphi(\pi_i(c_1), \dots, \pi_i(c_k))\} \in \mathcal{U}.$$

6. If  $\varphi$  is an  $\mathcal{L}$ -sentence, then

$$M \models \varphi \iff \{i \in I : M_i \models \varphi\} \in \mathcal{U}.$$

*Proof.* Part (1) is true by construction, as  $M$  is the canonical model of  $T$ . Part (5) is also true by construction:

$$\begin{aligned} M \models \varphi([c_1], \dots, [c_k]) &\iff M \models \varphi(c_1, \dots, c_k) \iff \varphi(c_1, \dots, c_k) \in T \\ &\iff \llbracket \varphi(c_1, \dots, c_k) \rrbracket \in \mathcal{U} \\ &\iff \{i \in I : M_i \models \varphi(\pi_i(c_1), \dots, \pi_i(c_k))\} \in \mathcal{U}. \end{aligned}$$

Remember that a constant symbol  $c \in P$  is interpreted as  $[c]$  in  $M$ , and  $\pi_i(c)$  in  $M_i$ .

Parts (2) and (3) follow by specializing part (5) to atomic formulas  $x = y$  and  $R(x_1, \dots, x_k)$ . Part (6) is the  $n = 0$  case of part (5). It remains to prove part (4). Fix  $c_1, \dots, c_k \in P$  and let  $d = f^P(c_1, \dots, c_k)$ . Then  $\pi_i(d) = f^{M_i}(\pi_i(c_1), \dots, \pi_i(c_k))$  for each  $i \in I$ . Then

$$\{i \in I : M_i \models \pi_i(d) = f(\pi_i(c_1), \dots, \pi_i(c_k))\} = I \in \mathcal{U},$$

so by part (5), we have

$$M \models [d] = f([c_1], \dots, [c_k]),$$

which means that  $f^M([c_1], \dots, [c_k]) = [d] = [f^P(c_1, \dots, c_k)]$ .  $\square$

**Remark 6.2.4.** Parts (1)–(4) determine the structure of  $M$  up to isomorphism, and are usually taken as the *definition* of the ultraproduct. Then part (5) is called *Łoś's Theorem*.

**Theorem 6.2.5.** Let  $I$  be a set,  $\mathcal{U}$  be an ultrafilter on  $I$ ,  $M_i$  be an  $\mathcal{L}$ -structure for  $i \in I$ , and  $M$  be the ultraproduct  $\prod_{i \in I}^{\mathcal{U}} M_i$ .

1. Let  $T$  be a theory. If  $M_i \models T$  for all  $i$ , then  $M \models T$ .

2. Let  $\mathcal{K}$  be an elementary class. If  $M_i \in \mathcal{K}$  for all  $i$ , then  $M \in \mathcal{K}$ .

*Proof.* (1) follows from Łoś's theorem: if  $\varphi \in T$ , then

$$\{i \in I : M_i \models \varphi\} = I \in \mathcal{U},$$

so  $M \models \varphi$ . (2) is a restatement of (1).  $\square$

If  $\mathcal{K}$  is a class of  $\mathcal{L}$ -structures and  $T$  is an  $\mathcal{L}$ -theory, say that  $T$  is *finitely satisfiable in  $\mathcal{K}$*  if for any finite subtheory  $T_0 \subseteq_f T$  there is  $M \in \mathcal{K}$  satisfying  $T_0$ .

**Theorem 6.2.6.** *If  $T$  is finitely satisfiable in  $\mathcal{K}$ , then there is an ultraproduct  $M$  of structures in  $\mathcal{K}$  such that  $M \models T$ .*

*Proof.* Let  $\{M_i\}_{i \in I}$  be a small collection of structures in  $\mathcal{K}$  containing at least one representative from every elementary equivalence class. If  $\varphi$  is an  $\mathcal{L}$ -sentence, let  $\llbracket \varphi \rrbracket = \{i \in I : M_i \models \varphi\}$ . Let  $\mathcal{F} = \{\llbracket \varphi \rrbracket : \varphi \in T\}$ . We claim that  $\mathcal{F}$  has the FIP. Indeed, if  $\varphi_1, \dots, \varphi_n \in T$  then there is some  $M_i$  satisfying  $\bigwedge_{j=1}^n \varphi_j$ , and then  $i \in \bigcap_{j=1}^n \llbracket \varphi_j \rrbracket$ .

Because  $\mathcal{F}$  has the FIP, it is contained in an ultrafilter  $\mathcal{U}$  on  $I$  (Theorem 6.1.7). Let  $M = \prod_{i \in I}^{\mathcal{U}} M_i$ . Then for  $\varphi \in T$  we have

$$\{i \in I : M_i \models \varphi\} = \llbracket \varphi \rrbracket \in \mathcal{F} \subseteq \mathcal{U},$$

and so  $M \models \varphi$  by Łoś's theorem.  $\square$

Note that this gives another proof of the compactness theorem (Theorem 4.4.5).

**Definition 6.2.7.** If  $M$  is an  $\mathcal{L}$ -structure and  $\mathcal{U}$  is an ultrafilter on a set  $I$ , then the *ultrapower*  $M^{\mathcal{U}}$  is the ultraproduct  $\prod_{i \in I}^{\mathcal{U}} M_i$ .

**Theorem 6.2.8.** *Let  $\Delta : M \rightarrow M^{\mathcal{U}}$  be the diagonal map sending  $a \in M$  to the class of the tuple  $(a : i \in I) \in M^I = \prod_{i \in I} M$ . Then  $\Delta$  is an elementary embedding.*

*Proof.* Fix a formula  $\varphi(x_1, \dots, x_n)$  and a tuple  $\bar{a} = (a_1, \dots, a_n) \in M^n$ . Let  $S = \{i \in I : M \models \varphi(a_1, \dots, a_n)\}$ . By Łoś's theorem,

$$M^{\mathcal{U}} \models \varphi(\Delta(a_1), \dots, \Delta(a_n)) \iff S \in \mathcal{U}.$$

But

$$S = \begin{cases} I & \text{if } M \models \varphi(\bar{a}) \\ \emptyset & \text{if } M \not\models \varphi(\bar{a}). \end{cases}$$

Therefore  $S \in \mathcal{U} \iff M \models \varphi(\bar{a})$ , and

$$M^{\mathcal{U}} \models \varphi(\Delta(\bar{a})) \iff M \models \varphi(\bar{a}). \quad \square$$

### 6.3 Characterizations of elementary classes

**Lemma 6.3.1.** *If  $M_1$  and  $M_2$  are elementarily equivalent, then there is an elementary embedding from  $M_2$  to an ultrapower  $M_1^{\mathcal{U}}$ .*

*Proof.* In the proof of elementary amalgamation, we saw that  $\text{eldiag}(M_2)$  is finitely satisfiable in expansions of  $M_1$  to  $\mathcal{L}(M_2)$ -structures (Lemma 5.2.3). By Theorem 6.2.6, there is an ultraproduct  $N = \prod_{i \in I}^{\mathcal{U}} N_i$  satisfying  $\text{eldiag}(M_2)$ , where each  $N_i$  is an expansion of  $M_1$  to an  $\mathcal{L}(M_2)$ -structure. The fact that  $N \models \text{eldiag}(M_2)$  gives an elementary embedding  $M_2 \rightarrow N$ . On the other hand,  $N \upharpoonright \mathcal{L}$  is

$$\prod_{i \in I}^{\mathcal{U}} (N_i \upharpoonright \mathcal{L}) = \prod_{i \in I}^{\mathcal{U}} M_1 = M_1^{\mathcal{U}},$$

and we have the desired elementary embedding  $M_2 \rightarrow M_1^{\mathcal{U}}$ .  $\square$

**Theorem 6.3.2.** *A class  $\mathcal{K}$  of  $\mathcal{L}$ -structures is elementary iff  $\mathcal{K}$  is closed under ultraproducts, isomorphisms, and elementary substructures.*

*Proof.* Suppose  $\mathcal{K}$  is closed under ultraproducts, isomorphisms, and elementary substructures.

*Claim.*  $\mathcal{K}$  is closed under elementary equivalence: if  $M \equiv N$  and  $M \in \mathcal{K}$  then  $N \in \mathcal{K}$ .

*Proof.* By Lemma 6.3.1, there is an ultrapower  $M^{\mathcal{U}}$  and an elementary embedding  $f : N \rightarrow M^{\mathcal{U}}$ . Then  $N \cong \text{im}(f) \preceq M^{\mathcal{U}}$ , so

$$M \in \mathcal{K} \implies M^{\mathcal{U}} \in \mathcal{K} \implies \text{im}(f) \in \mathcal{K} \implies N \in \mathcal{K}. \quad \square_{\text{Claim}}$$

Let  $T$  be the set of all sentences  $\varphi$  such that every  $N \in \mathcal{K}$  satisfies  $\varphi$ . Then  $\mathcal{K} \subseteq \text{Mod}(T)$ . We claim that  $\mathcal{K} = \text{Mod}(T)$ . Fix  $M \in \text{Mod}(T)$ ; we claim that  $M \in \mathcal{K}$ . Break into two cases:

1.  $\text{Th}(M)$  is finitely satisfiable in  $\mathcal{K}$ . By Theorem 6.2.6, there is a model  $N \models \text{Th}(M)$  such that  $N$  is an ultraproduct of structures in  $\mathcal{K}$ . By assumption,  $N \in \mathcal{K}$ . The fact that  $N \models \text{Th}(M)$  means that  $N \equiv M$  (Theorem 3.4.1), and so  $M \in \mathcal{K}$ .
2.  $\text{Th}(M)$  is not finitely satisfiable in  $\mathcal{K}$ . Then there is  $\varphi \in \text{Th}(M)$  such that no  $N \in \mathcal{K}$  satisfies  $\varphi$ . Then every  $N \in \mathcal{K}$  satisfies  $\neg\varphi$ , so  $\neg\varphi \in T$  and  $M \models \neg\varphi$ , contradicting the fact that  $\varphi \in \text{Th}(M)$ .  $\square$

**Fact 6.3.3** (Keisler-Shelah). *If  $M \equiv N$ , then there is a set  $I$  and an ultrafilter  $\mathcal{U}$  on  $I$  such that the ultraproducts  $M^{\mathcal{U}}$  and  $N^{\mathcal{U}}$  are isomorphic.*

**Definition 6.3.4.** A structure  $M$  is an *ultraroot* of a structure  $N$  if  $N$  is an ultrapower of  $M$ . A class  $\mathcal{K}$  is *closed under ultraroots* if  $M^{\mathcal{U}} \in \mathcal{K} \implies M \in \mathcal{K}$ .

If  $M$  is an ultraroot of  $N$ , then  $N \succeq M$ , so  $N \equiv M$ . Therefore, elementary classes are closed under ultraroots.

**Theorem 6.3.5.** *A class  $\mathcal{K}$  is elementary iff  $\mathcal{K}$  is closed under ultraproducts, ultraroots, and isomorphisms.*

*Proof.* Suppose  $\mathcal{K}$  is closed under ultraproducts, ultraroots, and isomorphisms. By the Keisler-Shelah theorem,  $\mathcal{K}$  is closed under elementary equivalence. Then the proof of Theorem 6.3.2 applies.  $\square$

## 6.4 Universal classes

**Lemma 6.4.1** (Lemma on constants). *Let  $\bar{c} = (c_1, \dots, c_n)$  be a tuple of constant symbols not appearing in  $T$ . If  $T \vdash \varphi(\bar{c})$ , then  $T \vdash \forall \bar{x} \varphi(\bar{x})$ .*

*Proof.* Otherwise, there is  $M \models T$  with  $M \not\models \forall \bar{x} \varphi(\bar{x})$ . Take  $\bar{a} \in M$  with  $M \models \neg \varphi(\bar{a})$ . Interpreting  $\bar{c}$  as  $\bar{a}$ , we get a model of  $T \cup \{\neg \varphi(\bar{c})\}$ , contradicting  $T \vdash \varphi(\bar{c})$ .  $\square$

A *universal formula* or  $\forall$ -*formula* is one of the form  $\forall \bar{y} : \varphi(\bar{x}, \bar{y})$  where  $\varphi$  is quantifier-free. A *universal theory* is a set of universal sentences.

**Lemma 6.4.2.** *If  $T$  is a universal theory, and  $M \models T$ , then any substructure of  $M$  is a model of  $T$ .*

*Proof.* Let  $\psi$  be a sentence in  $T$ . We can write  $\psi$  as  $\forall \bar{x} : \varphi(\bar{x})$  where  $\varphi$  is quantifier-free. Let  $N$  be a substructure of  $M$ . If  $\bar{a} \in N$ , then

$$M \models \forall \bar{x} : \varphi(\bar{x}) \implies M \models \varphi(\bar{a}) \implies N \models \varphi(\bar{a}),$$

where the second implication holds because  $\varphi(\bar{x})$  is quantifier-free (see Remark 3.5.4). As this holds for any  $\bar{a} \in N$ , we have  $N \models \forall \bar{x} : \varphi(\bar{x})$ .  $\square$

If  $T$  is a theory, then  $T_{\forall}$  denotes the set of universal sentences  $\varphi$  such that  $T \vdash \varphi$ .

**Theorem 6.4.3.**  $M \models T_\forall$  if and only if  $M$  is a substructure of a model of  $T$ .

*Proof.* First suppose  $M$  is a substructure of  $N \models T$ . Then  $N \models T_\forall$ , so  $M \models T_\forall$  as  $T_\forall$  is a universal theory.

Conversely, suppose  $M \models T_\forall$ . Break into two cases:

1. The  $\mathcal{L}(M)$ -theory  $\text{diag}(M) \cup T$  is consistent. Take a model  $N$ . Moving  $N$  by an isomorphism, we may assume that  $N$  is an extension of  $M$ , and  $N \models T$ . Then we are done.
2.  $\text{diag}(M) \cup T$  is inconsistent. By compactness, there is a finite tuple  $\bar{a} \in M$  and a quantifier-free formula  $\varphi(\bar{a}) = \bigwedge_{i=1}^n \varphi_i(\bar{a})$  with  $\varphi_i(\bar{a}) \in \text{diag}(M)$  such that  $\{\varphi(\bar{a})\} \cup T$  is inconsistent. Then  $T \vdash \neg\varphi(\bar{a})$ , so  $T \vdash \forall \bar{x} \neg\varphi(\bar{x})$  by the lemma on constants (Lemma 6.4.1). But then

$$(\forall \bar{x} \neg\varphi(\bar{x})) \in T_\forall,$$

so  $M \models \forall \bar{x} \neg\varphi(\bar{x})$ , contradicting the fact that  $M \models \varphi(\bar{a})$ .  $\square$

**Theorem 6.4.4.** Let  $T$  be a theory. Then  $T$  is axiomatizable by universal sentences if and only if  $\text{Mod}(T)$  is closed under substructures.

*Proof.* If  $T$  is axiomatizable by universal sentences, then  $\text{Mod}(T)$  is closed under substructures by Lemma 6.4.2.

Next suppose that  $\text{Mod}(T)$  is closed under substructures. We claim that  $T_\forall$  axiomatizes  $T$ . Certainly  $M \models T \implies M \models T_\forall$ . Conversely, suppose  $M \models T_\forall$ . By Theorem 6.4.3 there is an extension  $N \supseteq M$  with  $N \models T$ . By the assumption on  $T$ ,  $M \models T$ .  $\square$

**Theorem 6.4.5.** Let  $\mathcal{K}$  be a class of structures. Then the following are equivalent:

1.  $\mathcal{K}$  is axiomatized by a universal theory.
2.  $\mathcal{K}$  is closed under isomorphism, substructures, and ultraproducts.

*Proof.* (1)  $\implies$  (2) is clear.

(2)  $\implies$  (1): by Theorem 6.3.2,  $\mathcal{K}$  is an elementary class. Then use Theorem 6.4.4.  $\square$



# Chapter 7

## Topologies

### 7.1 Topological spaces

Fix a set  $S$ .

**Definition 7.1.1.** A family of sets  $\mathcal{F} \subseteq \mathfrak{P}(S)$  is *closed under infinite unions* if whenever  $I$  is a set and  $X_i \in \mathcal{F}$  for all  $i \in I$ , we have  $\bigcup_{i \in I} X_i \in \mathcal{F}$ . We say that  $\mathcal{F} \subseteq \mathfrak{P}(S)$  is *closed under finite unions* if this holds for finite  $I$ .

**Remark 7.1.2.** The definition of “*closed under (in)finite intersections*” is similar, with fine print. We understand  $\bigcap_{i \in I} X_i$  to mean

$$\{x \in S : \forall i \in I \ x \in X_i\}$$

Consequently, when  $I = \emptyset$ , the intersection  $\bigcap_{i \in I} X_i$  is defined to be  $S$ , and not the set-theoretic universe, for example.

**Definition 7.1.3.** A *metric space* is a set  $S$  and function  $d : S^2 \rightarrow \mathbb{R}$  satisfying the axioms:

1.  $d(x, y) \geq 0$
2.  $d(x, y) = d(y, x)$
3.  $d(x, y) = 0 \iff x = y$ .
4.  $d(x, z) \leq d(x, y) + d(y, z)$ .

**Example 7.1.4.**  $\mathbb{R}^2$  is a metric space with respect to the metric

$$d((x_1, y_1), (x_2, y_2)) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}.$$

Fix a metric space  $(S, d)$ . An *open ball* is a set of the form

$$B_\epsilon(p) = \{x \in S : d(x, p) < \epsilon\}$$

for some  $p \in S$  and  $\epsilon > 0$ . A set  $X \subseteq S$  is *open* if for every  $p \in X$ , there is  $\epsilon > 0$  such that  $B_\epsilon(p) \subseteq X$ .

**Fact 7.1.5.** Let  $(S, d)$  be a metric space.

1. Every open ball is open.
2. The collection of open sets is closed under infinite unions and finite intersections.

**Definition 7.1.6.** A *topology* on  $S$  is a family of sets  $\tau$  closed under finite intersections and infinite unions. A *topological space* is a set  $S$  with a topology  $\tau$ .

Fix a topological space  $(S, \tau)$ .

**Definition 7.1.7.** A set  $X \subseteq S$  is *open* if  $X \in \tau$  and *closed* if  $S \setminus X$  is open.

**Fact 7.1.8.** In a metric space, a set  $X$  is closed if and only if it “closed under limits” in the sense that:

$$\left(b_1, b_2, \dots \in X \text{ and } \lim_{i \rightarrow \infty} b_i = a\right) \implies a \in X.$$

**Definition 7.1.9.** A topological space  $S$  is *Hausdorff* if for any  $a_1 \neq a_2$  in  $S$ , there are open sets  $U_i \ni a_i$  with  $U_1 \cap U_2 = \emptyset$ .

**Remark 7.1.10.** Metric spaces are Hausdorff: if  $a_1 \neq a_2$ , then  $B_\epsilon(a_1) \cap B_\epsilon(a_2) = \emptyset$  for  $\epsilon = d(a_1, a_2)/3$ .

**Theorem 7.1.11.** If  $S$  is Hausdorff and  $p \in S$ , then  $\{p\}$  is closed.

*Proof.* For every  $q \neq p$ , take open sets  $U_q, V_q$  with  $q \in U_q$ ,  $p \in V_q$ , and  $U_q \cap V_q = \emptyset$ . In particular,  $q \in U_q$  and  $p \notin U_q$  for each  $q \neq p$ . Then  $S \setminus \{p\}$  is the open set  $\bigcup_{q \neq p} U_q$ .  $\square$

**Definition 7.1.12.** An *open cover* is a collection  $\mathcal{C}$  of open sets with  $\bigcup \mathcal{C} = S$ . A *subcover* of  $\mathcal{C}$  is another open cover  $\mathcal{C}' \subseteq \mathcal{C}$ .

**Definition 7.1.13.** A topological space  $S$  is *compact* if every open cover has a finite subcover.

**Fact 7.1.14.** A metric space  $(S, d)$  is compact if and only if every sequence has a convergent subsequence: for any  $a_1, a_2, a_3, \dots \in S$ , there is a subsequence

$$a_{i_1}, a_{i_2}, a_{i_3}, \dots$$

(with  $i_1 < i_2 < \dots$ ) such that  $\lim_{n \rightarrow \infty} a_{i_n}$  exists.

**Theorem 7.1.15.** If a topological space  $S$  is compact and  $\mathcal{F}$  is a family of closed sets with the FIP, then  $\bigcap \mathcal{F} \neq \emptyset$ . Conversely, this property characterizes compactness.

*Proof.* Let  $X_i$  be open for  $i \in I$ , and let  $Y_i$  be the complementary closed sets. We claim that the following are equivalent:

1. If  $\bigcup_{i \in I} X_i = S$ , then there is  $I_0 \subseteq_f I$  with  $\bigcup_{i \in I_0} X_i = S$ .
2. If  $\bigcap_{i \in I} Y_i = \emptyset$ , then there is  $I_0 \subseteq_f I$  with  $\bigcap_{i \in I_0} Y_i = \emptyset$ .
3. If  $\bigcap_{i \in I_0} Y_i \neq \emptyset$  for every  $I_0 \subseteq_f I$ , then  $\bigcap_{i \in I} Y_i \neq \emptyset$ .

Indeed, (1) and (2) are equivalent by de Morgan's laws, and (2) and (3) are contrapositives. Finally, observe that (1) is the definition of compactness, and (3) says that if  $\{Y_i\}_{i \in I}$  has FIP then  $\bigcap_i Y_i \neq \emptyset$ .  $\square$

**Definition 7.1.16.** Let  $S_1, S_2$  be two topological spaces and  $f : S_1 \rightarrow S_2$  be a function. Then  $f$  is *continuous* if for every open set  $U \subseteq S_2$ , the preimage  $f^{-1}(U)$  is open in  $S_1$ .

**Fact 7.1.17.** If  $S_1, S_2$  are metric spaces, a function  $f : S_1 \rightarrow S_2$  is continuous iff  $f$  “preserves limits”, in the sense that

$$\lim_{i \rightarrow \infty} b_i = a \implies \lim_{i \rightarrow \infty} f(b_i) = f(a).$$

**Definition 7.1.18.** A function  $f : S_1 \rightarrow S_2$  is a *homeomorphism* if  $f$  is continuous,  $f$  is a bijection, and  $f^{-1} : S_2 \rightarrow S_1$  is continuous. Two topological spaces  $S_1, S_2$  are *homeomorphic* if there is a homeomorphism from  $S_1$  to  $S_2$ .

## 7.2 Ultralimits

**Definition 7.2.1.** If  $I$  is a set,  $a_i \in S$  for  $i \in I$ ,  $\mathcal{U}$  is an ultrafilter on  $I$ , and  $b \in S$ , then  $b$  is an *ultralimit* of the  $a_i$ , written

$$b = \lim_{i \rightarrow \mathcal{U}} a_i$$

if for every open set  $N \ni b$ ,

$$\{i \in I : a_i \in N\} \in \mathcal{U}.$$

**Fact 7.2.2.** *If  $S$  is Hausdorff then ultralimits are unique: for any  $I, \mathcal{U}, \{a_i\}_{i \in I}$ , there is at most one  $b$  with  $b = \lim_{i \rightarrow \mathcal{U}} a_i$ . In fact, this property holds if and only if  $S$  is Hausdorff.*

*Half-proof.* Suppose  $S$  is Hausdorff, and

$$\begin{aligned} b &= \lim_{i \rightarrow \mathcal{U}} a_i \\ c &= \lim_{i \rightarrow \mathcal{U}} a_i. \end{aligned}$$

If  $b \neq c$ , by Hausdorffness there are open sets  $N_1 \ni b$  and  $N_2 \ni c$  with  $N_1 \cap N_2 = \emptyset$ . By definition of ultralimit, the sets

$$\begin{aligned} \{i \in I : a_i \in N_1\} \\ \{i \in I : a_i \in N_2\} \end{aligned}$$

are in the ultrafilter  $\mathcal{U}$ . But their intersection is empty, and  $\emptyset \notin \mathcal{U}$ , a contradiction.  $\square$

**Fact 7.2.3.** *If  $S$  is compact then ultralimits exist: for any  $I, \mathcal{U}, \{a_i\}_{i \in I}$ , there is at least one  $b$  with  $b = \lim_{i \rightarrow \mathcal{U}} a_i$ . In fact, this property holds if and only if  $S$  is compact.*

*Half-proof.* Suppose  $S$  is compact. Say an open set  $N \subseteq S$  is *good* if  $\{i \in I : a_i \in N\} \in \mathcal{U}$ , and *bad* otherwise, i.e., if  $\{i \in I : a_i \notin N\} \in \mathcal{U}$ . A finite union of bad sets is bad, because  $\mathcal{U}$  is closed under finite intersections. The set  $S$  is good. There are two cases:

- $S$  is covered by bad open sets. Then compactness gives a finite subcover, so  $S$  is a finite union of bad sets and  $S$  is bad, a contradiction.

- $S$  is not covered by bad open sets. Take  $p \in S$  such that  $p$  is in no bad set. Then every open set  $N \ni p$  is good, which means  $p = \lim_{i \rightarrow \mathcal{U}} a_i$ .  $\square$

**Theorem 7.2.4.** *A set  $C \subseteq S$  is closed if and only if  $C$  is closed under ultralimits, in the sense that*

$$\left( a_i \in C \text{ for all } i \in I \text{ and } b = \lim_{i \rightarrow \mathcal{U}} a_i \right) \implies b \in C.$$

*Proof.* First suppose  $C$  is closed, and  $a_i \in C$  for all  $i \in I$ . If  $b \notin C$ , then the complement  $S \setminus C$  is an open set containing  $b$ , but  $\{i \in I : a_i \in S \setminus C\} = \emptyset \notin \mathcal{U}$ , contradicting the definition of ultralimits. Thus  $b \in C$ , and  $C$  is closed under ultralimits.

Conversely, suppose  $C$  is closed under ultralimits. Let  $U_0$  be the union of open sets disjoint from  $C$ . It suffices to show that  $C \cup U_0 = S$ , as then  $C$  is the complement of the open set  $U_0$ . Fix  $p \notin U_0$ ; we claim  $p \in C$ . Let

$$\mathcal{F} = \{C \cap U : U \text{ is open and } U \ni p\}.$$

Then  $\mathcal{F}$  is closed under finite intersections, because a finite intersection of open sets is open. Moreover,  $\emptyset \notin \mathcal{F}$ , or else  $C \cap U = \emptyset$  and  $U$  shows  $p \in U_0$ , a contradiction. Thus  $\mathcal{F}$  has the FIP and is contained in an ultrafilter  $\mathcal{U}$  on  $C$ . Then

$$p = \lim_{x \rightarrow \mathcal{U}} x$$

because for any open set  $U$  containing  $p$ ,

$$\{x \in C : x \in U\} = C \cap U \in \mathcal{F} \subseteq \mathcal{U}.$$

By assumption on  $C$ ,  $p \in C$ .  $\square$

**Fact 7.2.5.** *If  $S_1, S_2$  are topological spaces and  $f : S_1 \rightarrow S_2$  is continuous, then  $f$  preserves ultralimits, in the sense that*

$$b = \lim_{i \rightarrow \mathcal{U}} a_i \implies f(b) = \lim_{i \rightarrow \mathcal{U}} f(a_i).$$

*In fact, this property holds if and only if  $f$  is continuous.*

*Half-proof.* Suppose  $f$  is continuous and  $b = \lim_{i \rightarrow \mathcal{U}} a_i$ . If  $N$  is an open set containing  $f(b)$ , then  $f^{-1}(N)$  is an open set containing  $b$ , and so

$$\{i \in I : f(a_i) \in N\} = \{i \in I : a_i \in f^{-1}(N)\} \in \mathcal{U}. \quad \square$$

**Theorem 7.2.6.** *Let  $S_1, S_2$  be compact Hausdorff spaces and let  $f : S_1 \rightarrow S_2$  be continuous.*

1. *If  $X \subseteq S_1$  is closed, then the image  $f(X)$  is closed.*
2. *If  $f$  is a bijection, then  $f$  is a homeomorphism.*

*Proof.* 1. By Theorem 7.2.4, it suffices to show that  $f(X)$  is closed under ultralimits. Suppose  $a_i \in f(X)$  and  $\lim_{i \rightarrow \mathcal{U}} a_i = b$ . We must show  $b \in f(X)$ . Write  $a_i$  as  $f(\alpha_i)$  for some  $\alpha_i \in X$ . Let

$$\beta = \lim_{i \rightarrow \mathcal{U}} \alpha_i.$$

The ultralimit exists as  $S_1$  is compact (Fact 7.2.3). The ultralimit  $\beta$  is in  $X$  because  $X$  is closed (Theorem 7.2.4). Then

$$f(\beta) = \lim_{i \rightarrow \mathcal{U}} f(\alpha_i) = \lim_{i \rightarrow \mathcal{U}} a_i$$

because the continuous function  $f$  preserves ultralimits (Fact 7.2.5). As  $S_2$  is Hausdorff, ultralimits are unique (Fact 7.2.2), and so  $f(\beta) = b$ . Then  $\beta \in X \implies b = f(\beta) \in f(X)$ .

2. Let  $X$  be a subset of  $S_1$ . Part (1) shows that

$$X \text{ is closed in } S_1 \implies f(X) \text{ is closed in } S_2.$$

The reverse direction holds by continuity. Thus  $f$  is a homeomorphism.  $\square$

### 7.3 The space of complete theories

Fix a language  $\mathcal{L}$ . Let  $S = \{\text{Th}(M) : M \text{ is an } \mathcal{L}\text{-structure}\}$ . For each  $\mathcal{L}$ -sentence  $\varphi$ , let

$$\llbracket \varphi \rrbracket = \{T \in S : \varphi \in T\} = \{\text{Th}(M) : M \models \varphi\}.$$

Note that  $M \models \varphi \wedge \psi \iff (M \models \varphi \text{ and } M \models \psi)$ , and so

$$\llbracket \varphi \wedge \psi \rrbracket = \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket.$$

Similarly,

$$\begin{aligned}\llbracket \varphi \vee \psi \rrbracket &= \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket \\ \llbracket \neg \varphi \rrbracket &= S \setminus \llbracket \varphi \rrbracket \\ \llbracket \top \rrbracket &= S \\ \llbracket \perp \rrbracket &= \emptyset.\end{aligned}$$

**Definition 7.3.1.** A set  $X \subseteq S$  is *clopen* if  $X = \llbracket \varphi \rrbracket$  for some sentence  $\varphi$ , and *open* if  $X = \bigcup_{i \in I} Y_i$  where each  $Y_i$  is clopen.

**Theorem 7.3.2.** *The collection of open sets is closed under infinite unions and finite intersections, so it defines a topology on  $S$ . Moreover, the topology is Hausdorff.*

*Proof.* Infinite unions are clear: an infinite union of infinite unions of clopen sets is an infinite union of clopen sets. For finite intersections, note that

$$\left( \bigcup_{i \in I} \llbracket \varphi_i \rrbracket \right) \cap \left( \bigcup_{j \in J} \llbracket \psi_j \rrbracket \right) = \bigcup_{(i,j) \in I \times J} (\llbracket \varphi_i \rrbracket \cap \llbracket \psi_j \rrbracket) = \bigcup_{(i,j) \in I \times J} \llbracket \varphi_i \wedge \psi_j \rrbracket.$$

For Hausdorffness, suppose  $T_1, T_2 \in S$  and  $T_1 \neq T_2$ . Then there is a sentence  $\varphi$  such that, say,  $\varphi \in T_1$  and  $\varphi \notin T_2$ . This means  $T_1 \in \llbracket \varphi \rrbracket$  and  $T_2 \notin \llbracket \varphi \rrbracket$ . Then  $T_2 \in S \setminus \llbracket \varphi \rrbracket = \llbracket \neg \varphi \rrbracket$ . The two open sets  $\llbracket \varphi \rrbracket$  and  $\llbracket \neg \varphi \rrbracket$  separate  $T_1$  from  $T_2$ .  $\square$

**Lemma 7.3.3.** *Let  $X_i$  be clopen for each  $i \in I$ .*

1. *Suppose  $\{X_i : i \in I\}$  has the FIP: for any  $I_0 \subseteq_f I$ , we have  $\bigcap_{i \in I_0} X_i \neq \emptyset$ . Then  $\bigcap_{i \in I} X_i \neq \emptyset$ .*
2. *If  $S = \bigcup_{i \in I} X_i$ , then there is finite  $I_0 \subseteq_f I$  such that  $S = \bigcup_{i \in I_0} X_i$ .*

*Proof.* 1. Let  $X_i = \llbracket \varphi_i \rrbracket$ . For any  $I_0 \subseteq I$ , there is  $M$  with  $\text{Th}(M) \in \bigcap_{i \in I_0} \llbracket \varphi_i \rrbracket$ , meaning that  $M \models \{\varphi_i : i \in I_0\}$ . By the compactness theorem, there is  $M$  satisfying  $\{\varphi_i : i \in I\}$ , and then  $\text{Th}(M) \in \bigcap_{i \in I} \llbracket \varphi_i \rrbracket$ .

2. Let  $Y_i = S \setminus X_i$ . Apply part (1) to the family of clopen sets  $\{Y_i\}_{i \in I}$ . By assumption,  $\bigcap_{i \in I} Y_i = \emptyset$ , so there must be  $I_0 \subseteq_f I$  such that  $\bigcap_{i \in I_0} Y_i = \emptyset$ , or equivalently,  $\bigcup_{i \in I_0} X_i = S$ .  $\square$

**Theorem 7.3.4.** *The topological space  $S$  is compact.*

*Proof.* Suppose  $S = \bigcup_{i \in I} U_i$  for some open sets  $U_i$ . Let  $\mathcal{F}$  be the family of clopen sets  $X$  such that  $X \subseteq U_i$  for some  $i$ . Every open set is the union of its clopen subsets, so  $U_i \subseteq \bigcup \mathcal{F}$  for each  $i$ . Then  $S = \bigcup_{i \in I} U_i \subseteq \bigcup \mathcal{F}$ . Applying Lemma 7.3.3(2) to the clopen cover  $\mathcal{F}$ , there is a finite subcover  $S = \bigcup_{j=1}^n X_j$  with  $X_j \in \mathcal{F}$ . For each  $j$ , choose a  $U_{i_j} \supseteq X_j$ . Then  $S = \bigcup_{j=1}^n X_j \subseteq \bigcup_{j=1}^n U_{i_j}$ , and  $\{U_{i_1}, \dots, U_{i_n}\}$  is an open subcover of the original cover.  $\square$

If  $S$  is compact, then ultralimits should exist (Fact 7.2.3). In fact, ultralimits correspond exactly to ultraproducts:

**Theorem 7.3.5.** *Let  $M$  be an ultraproduct  $\prod_{i \in I}^{/\mathcal{U}} M_i$ . Then  $\text{Th}(M)$  is the ultralimit  $\lim_{i \rightarrow \mathcal{U}} \text{Th}(M_i)$  in the topological space  $S$ .*

*Proof.* Let  $N$  be an open set containing  $\text{Th}(M)$ . Then there is a clopen set  $\llbracket \varphi \rrbracket$  with  $\text{Th}(M) \in \llbracket \varphi \rrbracket \subseteq N_0$ , because  $N_0$  is a union of clopen sets. Then

$$\text{Th}(M) \in \llbracket \varphi \rrbracket \implies M \models \varphi \implies \{i \in I : M_i \models \varphi\} \in \mathcal{U}$$

by Łoś's theorem (Theorem 6.2.3(5)). If  $M_i \models \varphi$ , then  $\text{Th}(M_i) \in \llbracket \varphi \rrbracket \subseteq N$ . Therefore

$$\{i \in I : \text{Th}(M_i) \in N\} \in \mathcal{U}.$$

$\square$

Recall that  $X$  is closed iff  $S \setminus X$  is open.

**Theorem 7.3.6.** *A set  $X \subseteq S$  is clopen if and only if  $X$  is both closed and open.*

*Proof.* If  $X$  is clopen, then  $X$  is open. The complement  $S \setminus X$  is clopen, hence open, and so  $X$  is also closed.

Conversely, suppose  $X$  is closed and open. Then  $X = \bigcup_{i \in I} Y_i$  and  $S \setminus X = \bigcup_{j \in J} Z_j$  where the  $Y_i$  and  $Z_j$  are clopen sets. Note that  $S = \bigcup_{i \in I} Y_i \cup \bigcup_{j \in J} Z_j$ . By Lemma 7.3.3(2), there are finite  $I_0 \subseteq_f I$  and  $J_0 \subseteq_f J$  such that  $S = \bigcup_{i \in I_0} Y_i \cup \bigcup_{j \in J_0} Z_j$ . Then  $X$  is the clopen set  $\bigcup_{i \in I_0} Y_i$ .  $\square$

## 7.4 Stone spaces

**Definition 7.4.1.** In any topological space, a *clopen set* is a set that is both closed and open.

**Definition 7.4.2.** A *Stone space* is a compact, Hausdorff topological space in which every open set is a union of clopen sets.



**Theorem 7.4.3.** *Let  $S$  be a set. Let  $\mathcal{B}$  be a boolean subalgebra of  $\mathfrak{P}(S)$ . Suppose the following two conditions hold:*

1. *If  $a, b \in S$  are distinct, then there is  $X \in \mathcal{B}$  with  $a \in X$  and  $b \notin X$ , or  $b \in X$  and  $a \notin X$ .*
2. *If  $\{X_i\}_{i \in I}$  is a family of sets in  $\mathcal{B}$  with the FIP, then  $\bigcap_{i \in I} X_i \neq \emptyset$ .*

*Then there is a Stone space topology on  $S$  in which the clopen sets are exactly the elements of  $\mathcal{B}$ .*

*Proof.* This follows by the arguments of Section 7.3. □

**Theorem 7.4.4.** *Let  $S_1, S_2$  be Stone spaces and  $f : S_1 \rightarrow S_2$  be a map. Suppose that for any clopen set  $X$  in  $S_2$ , the preimage  $f^{-1}(X)$  is clopen in  $S_1$ . Then  $f$  is continuous.*

*Proof.* If  $U$  is open in  $S_2$ , then  $U = \bigcup_{i \in I} X_i$  for some clopen sets  $X_i$ . Then  $f^{-1}(U) = \bigcup_{i \in I} f^{-1}(X_i)$  which is open in  $S_1$ . □



# Chapter 8

## Types and quantifier elimination

### 8.1 Types

**Definition 8.1.1.** Let  $M$  be an  $\mathcal{L}$ -structure, let  $A$  be a subset, and let  $\bar{b}$  be an  $n$ -tuple. The *type* of  $\bar{b}$  over  $A$ , written  $\text{tp}(\bar{b}/A)$ , is the set of  $\mathcal{L}(A)$ -formulas  $\varphi(x_1, \dots, x_n)$  such that  $M \models \varphi(\bar{b})$ . When  $A = \emptyset$ , we write  $\text{tp}(\bar{b}/A)$  as  $\text{tp}(\bar{b})$ . We write  $\text{tp}(-)$  as  $\text{tp}^M(-)$  when we need to specify  $M$ .

**Remark 8.1.2.** Partial elementary maps preserve types: if  $f$  is a partial elementary map from  $M$  to  $N$  and  $\bar{a}$  is a tuple in  $\text{dom}(f)$ , then  $\text{tp}(\bar{a}) = \text{tp}(f(\bar{a}))$ . Indeed, for any formula  $\varphi(\bar{x})$ ,

$$\varphi(\bar{x}) \in \text{tp}(\bar{a}) \iff M \models \varphi(\bar{a}) \iff N \models \varphi(f(\bar{a})) \iff \varphi(\bar{x}) \in \text{tp}(f(\bar{a})).$$

**Remark 8.1.3.** If  $\bar{b}, \bar{c}$  are two  $n$ -tuples in the same structure  $M$ , then the following are equivalent:

1.  $\text{tp}(\bar{b}/A) = \text{tp}(\bar{c}/A)$ .
2. For every  $A$ -definable set  $D$ ,  $\bar{b} \in D \iff \bar{c} \in D$ .

Indeed, (1) and (2) are equivalent to

3. For every  $\mathcal{L}(A)$ -formula  $\varphi(\bar{x})$ ,  $M \models \varphi(\bar{b}) \iff M \models \varphi(\bar{c})$ .

Let  $M$  be an  $\mathcal{L}$ -structure and  $A$  be a set.

**Definition 8.1.4.** A *complete  $n$ -type* over  $A$  is something of the form  $\text{tp}^N(\bar{b}/A)$  for some  $N \succeq M$  and  $n$ -tuple  $\bar{b} \in N^n$ . The set of complete  $n$ -types is written  $S_n(A)$ .

**Lemma 8.1.5.** 1. If  $p \in S_n(A)$  and  $\varphi(\bar{x})$  is an  $\mathcal{L}(A)$ -formula, then  $\neg\varphi \in p \iff \varphi \notin p$ .

2. If  $p, q \in S_n(A)$  and  $p \subseteq q$ , then  $p = q$ .

*Proof.* 1. If  $p = \text{tp}^N(\bar{b}/A)$  for some elementary extension  $N \succeq M$  and  $\bar{b} \in N^n$ , then

$$\neg\varphi \in p \iff N \models \neg\varphi(\bar{b}) \iff N \not\models \varphi(\bar{b}) \iff \varphi \notin p.$$

2. Otherwise take  $\varphi \in q \setminus p$ . Then  $\varphi \in q \implies \neg\varphi \notin q$ , and  $\varphi \notin p \implies \neg\varphi \in p$ . Then  $\neg\varphi \in p \setminus q$ , contradicting  $p \subseteq q$ .  $\square$

**Definition 8.1.6.** Let  $\Sigma(x_1, \dots, x_n)$  be a set of  $\mathcal{L}(A)$ -formulas in the variables  $x_1, \dots, x_n$ . Then  $\bar{b} \in M^n$  *realizes*  $\Sigma$  if  $\Sigma \subseteq \text{tp}(\bar{b}/A)$ . We say that  $\Sigma(\bar{x})$  is *realized* in  $M$  if some  $\bar{a} \in M^n$  realizes  $\Sigma$ , and *omitted* in  $M$  otherwise.

**Remark 8.1.7.** If  $p \in S_n(A)$  is a complete type, then a tuple  $\bar{b}$  realizes  $p$  if and only if  $\text{tp}(\bar{b}/A) = p$ , by Lemma 8.1.5(2) applied to the complete types  $p$  and  $\text{tp}(\bar{b}/A)$ .

**Theorem 8.1.8.** Let  $\Sigma(\bar{x})$  be a set of  $\mathcal{L}(A)$ -formulas in the variables  $x_1, \dots, x_n$ . The following are equivalent:

1. Every finite subset  $\Sigma_0 \subseteq_f \Sigma$  is realized in  $M$ .
2.  $\Sigma$  is realized in an elementary extension of  $M$ .

*Proof.* Consider a third condition:

3. Every finite subset  $\Sigma_0 \subseteq_f \Sigma$  is realized in an elementary extension of  $M$ .

We claim  $(3) \implies (1) \implies (2) \implies (3)$ . The direction  $(2) \implies (3)$  is clear.

$(3) \implies (1)$ : if  $\Sigma_0 = \{\varphi_1, \dots, \varphi_n\}$ , and  $\Sigma_0$  is realized in  $N \succeq M$ , then

$$N \models \exists \bar{x} \bigwedge_{i=1}^n \varphi_i(\bar{x}).$$

As  $M \preceq N$ , the same sentence holds in  $M$ , which means that  $\Sigma_0$  is realized in  $M$ .

(1)  $\implies$  (2): Let  $\mathcal{L}' = \mathcal{L}(A) \cup \{c_1, \dots, c_n\}$  where the  $c_i$  are new constant symbols. If  $\Sigma_0 \subseteq_f \Sigma$ , then the  $\mathcal{L}'$ -structure

$$\text{eldiag}(M) \cup \{\varphi(\bar{c}) : \varphi \in \Sigma_0\}$$

is satisfied by the structure  $M$  with  $\bar{c}$  interpreted as a realization of  $\Sigma_0$ . By compactness,

$$\text{eldiag}(M) \cup \{\varphi(\bar{c}) : \varphi \in \Sigma\}$$

is satisfied by some  $\mathcal{L}'$ -structure  $N$ . The fact that  $N \models \text{eldiag}(M)$  means that, up to isomorphism,  $N \succeq M$ . Then the interpretation  $\bar{c}^N$  of  $\bar{c}$  in  $N$  is an  $n$ -tuple realizing  $\Sigma(\bar{x})$ , and so (2) holds.  $\square$

**Definition 8.1.9.** A *partial  $n$ -type* over  $A$  is a set  $\Sigma(\bar{x})$  of  $\mathcal{L}(A)$ -formulas in the variables  $x_1, \dots, x_n$  satisfying the equivalent conditions of Theorem 8.1.8.

**Remark 8.1.10.** Condition (2) of Theorem 8.1.8 says that  $\Sigma(\bar{x})$  is a partial type if and only if  $\Sigma(\bar{x}) \subseteq \text{tp}^N(\bar{b}/A)$  for some tuple  $\bar{b}$  in an elementary extension  $N \succeq M$ . Equivalently,  $\Sigma(\bar{x})$  is a partial type if and only if  $\Sigma$  is a subset of a complete type. In particular, complete types are partial types.

**Theorem 8.1.11.** Let  $p(\bar{x})$  be a set of  $\mathcal{L}(A)$ -formulas in  $\bar{x}$ . Then  $p(\bar{x})$  is a complete type over  $A$  if and only if  $p(\bar{x})$  is a maximal partial type over  $A$ .

*Proof.* By Remark 8.1.10, every partial type is contained in a complete type. Therefore, any maximal partial type is a complete type.

Conversely, suppose  $p$  is a complete type, but not maximal. Take a larger partial type  $\Sigma \supsetneq p$ . Then  $\Sigma \subseteq q$  for some complete type  $q$ , and we have  $p \subsetneq \Sigma \subseteq q$ , contradicting the incomparability of complete types (Lemma 8.1.5(2)).  $\square$

## Types over a theory

Let  $T$  be an  $\mathcal{L}$ -theory.

**Definition 8.1.12.** A *complete  $n$ -type* over  $T$  is something of the form  $\text{tp}^M(\bar{a})$  for some  $M \models T$  and  $n$ -tuple  $\bar{a} \in M^n$ . The set of complete  $n$ -types is written  $S_n(T)$ .

**Lemma 8.1.13.** *If  $p, q \in S_n(T)$  and  $p \subseteq q$ , then  $p = q$ .*

*Proof.* Like Lemma 8.1.5. □

**Theorem 8.1.14.** *If  $M \models T$  and  $\bar{a} \in M^n$  realizes  $p \in S_n(T)$ , then  $\text{tp}(\bar{a}) = p$ .*

*Proof.* Like Remark 8.1.7. □

**Theorem 8.1.15.** *Let  $\Sigma(\bar{x})$  be a set of  $\mathcal{L}$ -formulas in the variables  $x_1, \dots, x_n$ . The following are equivalent:*

1. *Every finite subset  $\Sigma_0 \subseteq \Sigma$  is realized in a model of  $T$ .*
2.  *$\Sigma$  is realized in a model of  $T$ .*

*Proof.* Like Theorem 8.1.8, but more straightforward. □

**Definition 8.1.16.** *A partial  $n$ -type over  $T$  is a set  $\Sigma(\bar{x})$  of  $\mathcal{L}$ -formulas in the variables  $x_1, \dots, x_n$  satisfying the equivalent conditions of Theorem 8.1.15.*

**Remark 8.1.17.** As in Remark 8.1.10,  $\Sigma(\bar{x})$  is a partial type if and only if  $\Sigma(\bar{x})$  is a subset of a complete type. In particular, complete types are partial types.

**Theorem 8.1.18.** *Let  $p(\bar{x})$  be a set of  $\mathcal{L}$ -formulas in  $\bar{x}$ . Then  $p(\bar{x})$  is a complete type over  $T$  if and only if  $p(\bar{x})$  is a maximal partial type over  $T$ .*

*Proof.* Like Theorem 8.1.11 □

**Lemma 8.1.19.** *Let  $T$  be a complete theory,  $M$  be a model, and  $\Sigma(\bar{x})$  be a set of  $\mathcal{L}$ -formulas in the variables  $\bar{x}$ .*

1. *If  $\Sigma(\bar{x})$  is finite, then  $\Sigma(\bar{x})$  is realized in a model of  $T$  if and only if  $\Sigma(\bar{x})$  is realized in  $M$ .*
2.  *$\Sigma(\bar{x})$  is a partial type over  $T$  if and only if  $\Sigma(\bar{x})$  is a partial type over  $\emptyset \subseteq M$ .*
3.  *$\Sigma(\bar{x})$  is a complete type over  $T$  if and only if  $\Sigma(\bar{x})$  is a complete type over  $\emptyset \subseteq M$ .*

*Proof.* 1. If  $N \models \exists \bar{x} \bigwedge_{\varphi \in \Sigma} \varphi(\bar{x})$  and  $N \models T$  then  $M \models \exists \bar{x} \bigwedge_{\varphi \in \Sigma} \varphi(\bar{x})$  because  $M \equiv N$ .

2. By part (1),  $\Sigma(\bar{x})$  is finitely realized in models of  $T$  if and only if  $\Sigma(\bar{x})$  is finitely realized in  $M$ .
3. The complete types over  $T$  are the maximal partial types over  $T$ , and similarly for types over  $A = \emptyset$ . Since the sets of partial types agree by part (2), the sets of maximal partial types agree.  $\square$

**Corollary 8.1.20.** *If  $T$  is a complete theory and  $M$  is a model, then  $S_n(T) = S_n^M(\emptyset)$ .*

**Fact 8.1.21.** *If  $A \subseteq M$  and  $T_{A,M}$  is the complete  $\mathcal{L}(A)$ -theory of  $M$ , then  $S_n^M(A) = S_n(T_{A,M})$ .*

## 8.2 The topology on type space

Let  $M$  be an  $\mathcal{L}$ -structure and  $A$  be a subset of  $M$ .

**Theorem 8.2.1.** *For each  $\mathcal{L}(A)$ -formula  $\varphi(x_1, \dots, x_n)$ , let  $\llbracket \varphi \rrbracket = \{p \in S_n(A) : \varphi \in p\}$ . Then*

$$\begin{aligned} \llbracket \varphi \wedge \psi \rrbracket &= \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket \\ \llbracket \varphi \vee \psi \rrbracket &= \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket \\ \llbracket \neg \varphi \rrbracket &= S_n(A) \setminus \llbracket \varphi \rrbracket \\ \llbracket \top \rrbracket &= S_n(A) \quad \llbracket \perp \rrbracket = \emptyset \end{aligned}$$

*Proof.* This is a matter of unwinding definitions. For example, if  $p \in S_n(A)$  is  $\text{tp}^N(\bar{b}/A)$ , then

$$\begin{aligned} p \in \llbracket \varphi \wedge \psi \rrbracket &\iff \varphi \wedge \psi \in p \iff N \models \varphi(\bar{b}) \wedge \psi(\bar{b}) \\ &\iff N \models \varphi(\bar{b}) \text{ and } N \models \psi(\bar{b}) \iff \dots \iff (p \in \llbracket \varphi \rrbracket \text{ and } p \in \llbracket \psi \rrbracket), \end{aligned}$$

showing that  $\llbracket \varphi \wedge \psi \rrbracket = \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket$ .  $\square$

**Theorem 8.2.2.** *If  $\varphi, \psi$  are  $\mathcal{L}(A)$ -formulas, then*

$$\begin{aligned} \llbracket \varphi \rrbracket \subseteq \llbracket \psi \rrbracket &\iff \varphi(M) \subseteq \psi(M) \\ \llbracket \varphi \rrbracket = \llbracket \psi \rrbracket &\iff \varphi(M) = \psi(M). \end{aligned}$$

*Proof.* We prove the first line, which directly implies the second line. Unwinding the definitions,  $\llbracket \varphi \rrbracket \subseteq \llbracket \psi \rrbracket$  means that for every  $N \succeq M$  and  $\bar{b} \in N$ , if  $N \models \varphi(\bar{b})$  then  $N \models \psi(\bar{b})$ . Equivalently, every  $N \succeq M$  satisfies

$$N \models \forall \bar{x} (\varphi(\bar{x}) \rightarrow \psi(\bar{x})).$$

Similarly,  $\varphi(M) \subseteq \psi(M)$  means

$$M \models \forall \bar{x} (\varphi(\bar{x}) \rightarrow \psi(\bar{x})).$$

The two conditions are equivalent because  $N \succeq M$ . □

**Theorem 8.2.3.** *The collection of sets  $\llbracket \varphi \rrbracket$  form a basis for a topology making  $S_n(A)$  into a Stone space, and the sets  $\llbracket \varphi \rrbracket$  are exactly the clopen sets in this topology.*

*Proof.* We use Theorem 7.4.3. There are two things to check:

1. If  $p, q \in S_n(A)$  are distinct, then there is a set  $\llbracket \varphi \rrbracket$  distinguishing  $p$  and  $q$ . Indeed, take  $\varphi$  in  $p \setminus q$ .
2. If  $\{\llbracket \varphi_i \rrbracket\}_{i \in I}$  has the FIP, then  $\bigcap_i \llbracket \varphi_i \rrbracket \neq \emptyset$ . Equivalently, if every finite subset of  $\{\varphi_i : i \in I\}$  is contained in a complete type, then  $\{\varphi_i : i \in I\}$  is contained in a complete type. This is clear by Theorem 8.1.8. □

Using Theorem 8.2.2 we see the following:

**Corollary 8.2.4.** *The boolean algebra of clopen sets in  $S_n(A)$  is isomorphic to the boolean algebra of  $A$ -definable subsets of  $M^n$  via the isomorphism  $\llbracket \varphi \rrbracket \mapsto \varphi(M^n)$ .*

There is an analogous picture for type spaces of a theory. Let  $T$  be an  $\mathcal{L}$ -theory:

**Theorem 8.2.5.** *For each  $\mathcal{L}$ -formula  $\varphi(x_1, \dots, x_n)$ , let  $\llbracket \varphi \rrbracket = \{p \in S_n(T) : \varphi \in p\}$ . Then*

$$\begin{aligned} \llbracket \varphi \wedge \psi \rrbracket &= \llbracket \varphi \rrbracket \cap \llbracket \psi \rrbracket \\ \llbracket \varphi \vee \psi \rrbracket &= \llbracket \varphi \rrbracket \cup \llbracket \psi \rrbracket \\ \llbracket \neg \varphi \rrbracket &= S_n(T) \setminus \llbracket \varphi \rrbracket \\ \llbracket \top \rrbracket &= S_n(T) \quad \llbracket \perp \rrbracket = \emptyset \end{aligned}$$



Moreover,  $\llbracket \varphi \rrbracket = \llbracket \psi \rrbracket$  if and only if  $\varphi$  and  $\psi$  are equivalent modulo  $T$ , in the sense that

$$T \vdash \forall \bar{x} (\varphi(\bar{x}) \leftrightarrow \psi(\bar{x}))$$

or equivalently  $\varphi(M) = \psi(M)$  for all  $M \models T$ .

**Theorem 8.2.6.** *The collection of sets  $\llbracket \varphi \rrbracket$  is a basis for a topology making  $S_n(T)$  a Stone space, and the clopen sets are exactly the sets  $\llbracket \varphi \rrbracket$ .*

### 8.3 Quantifier elimination

**Definition 8.3.1.** A theory  $T$  has *quantifier elimination* if for every formula  $\varphi(\bar{x})$ , there is a quantifier-free formula  $\psi(\bar{x})$  that is equivalent to  $\varphi$  in models of  $T$ :

$$T \vdash \forall \bar{x} (\varphi(\bar{x}) \leftrightarrow \psi(\bar{x})).$$

**Definition 8.3.2.** If  $(a_1, \dots, a_n)$  is a tuple in a structure  $M$ , then the *quantifier-free type*  $\text{qftp}(\bar{a})$  is the set of quantifier-free formulas  $\varphi(x_1, \dots, x_n)$  with  $M \models \varphi(\bar{a})$ .

For any theory  $T$ , let  $S_n^{\text{qfree}}(T)$  be the space of quantifier-free  $n$ -types, analogous to the space of complete  $n$ -types  $S_n(T)$ . Each quantifier-free formula  $\varphi(x_1, \dots, x_n)$  defines a set  $\llbracket \varphi \rrbracket_{\text{qfree}} \subseteq S_n^{\text{qfree}}(T)$ , and these sets form a basis for a topology making  $S_n^{\text{qfree}}(T)$  into a Stone space. Moreover, every clopen set in  $S_n^{\text{qfree}}(T)$  has the form  $\llbracket \varphi \rrbracket_{\text{qfree}}$  for some quantifier-free  $\varphi$ . The proofs are analogous to the case of  $S_n(T)$  (Section 8.2). There is a restriction map

$$S_n(T) \rightarrow S_n^{\text{qfree}}(T)$$

sending  $\text{tp}(\bar{a})$  to  $\text{qftp}(\bar{a})$ . Note that

$$\text{qftp}^M(\bar{a}) \in \llbracket \varphi \rrbracket_{\text{qfree}} \iff M \models \varphi(\bar{a}) \iff \text{tp}^M(\bar{a}) \in \llbracket \varphi \rrbracket$$

so the preimage of  $\llbracket \varphi \rrbracket_{\text{qfree}}$  is  $\llbracket \varphi \rrbracket$ . Because preimages of clopen sets are clopen, the restriction map is continuous (Theorem 7.4.4).

**Theorem 8.3.3.** *The following are equivalent:*

1.  $T$  has quantifier elimination.

2. For any  $M, N \models T$  and  $\bar{a} \in M^n$  and  $\bar{b} \in N^n$ ,

$$\text{qftp}^M(\bar{a}) = \text{qftp}^N(\bar{b}) \implies \text{tp}^M(\bar{a}) = \text{tp}^N(\bar{b}).$$

*Proof.* Consider a third condition:

3. For every  $n$ , the restriction map  $S_n(T) \rightarrow S_n^{\text{qfree}}(T)$  is a homeomorphism.

We claim  $(1) \implies (2) \implies (3) \implies (1)$ .

$(1) \implies (2)$ : Suppose that  $T$  has quantifier elimination and  $\text{qftp}^M(\bar{a}) = \text{qftp}^N(\bar{b})$ . For any formula  $\varphi(\bar{x})$ , there is an equivalent quantifier free-formula  $\psi(\bar{x})$ , and then

$$M \models \varphi(\bar{a}) \iff M \models \psi(\bar{a}) \iff N \models \psi(\bar{b}) \iff N \models \varphi(\bar{b}),$$

so that  $\text{tp}^M(\bar{a}) = \text{tp}^N(\bar{b})$ .

$(2) \implies (3)$ : The restriction map is surjective, since  $\text{qftp}^M(\bar{a})$  lifts to  $\text{tp}^M(\bar{a})$ . The restriction map is injective by (2). Then the restriction map is a continuous bijection between compact Hausdorff spaces, and therefore a homeomorphism (Theorem 7.2.6).

$(3) \implies (1)$ : Let  $\varphi(x_1, \dots, x_n)$  be a formula. Then  $\llbracket \varphi \rrbracket \subseteq S_n(T)$  is clopen. The image under the homeomorphism  $S_n(T) \rightarrow S_n^{\text{qfree}}(T)$  is a clopen set, which must be  $\llbracket \psi \rrbracket_{\text{qfree}} \subseteq S_n^{\text{qfree}}(T)$  for some quantifier-free formula  $\psi$ . Then

$$M \models \varphi(\bar{a}) \iff \text{tp}(\bar{a}) \in \llbracket \varphi \rrbracket \iff \text{qftp}(\bar{a}) \in \llbracket \psi \rrbracket_{\text{qfree}} \iff M \models \psi(\bar{a}),$$

so  $T \vdash \varphi \leftrightarrow \psi$ . □

**Theorem 8.3.4.** Suppose  $M, N$  are  $\mathcal{L}$ -structures and  $\bar{a} \in M^n$  and  $\bar{b} \in N^n$ . Then the following are equivalent:

1.  $\text{qftp}(\bar{a}) = \text{qftp}(\bar{b})$ .

2. There is an isomorphism  $f : \langle \bar{a} \rangle_M \rightarrow \langle \bar{b} \rangle_N$  with  $f(\bar{a}) = \bar{b}$ .

*Proof.*  $(1) \implies (2)$ : Let  $p = \text{qftp}(\bar{a}) = \text{qftp}(\bar{b})$ . Note that every element of  $\langle \bar{a} \rangle_M$  has the form  $t^M(\bar{a})$  for some term  $t$ . Similarly,  $\langle \bar{b} \rangle_N = \{t^N(\bar{b}) : t(\bar{x}) \text{ a term}\}$ . Moreover,

$$t^M(\bar{a}) = s^M(\bar{a}) \iff (t(\bar{x}) = s(\bar{x})) \in p \iff t^N(\bar{b}) = s^N(\bar{b}),$$

Therefore there is a bijection  $f : \langle \bar{a} \rangle_M \rightarrow \langle \bar{b} \rangle_N$  sending  $t^M(\bar{a})$  to  $t^N(\bar{b})$ . If  $R$  is a  $k$ -ary relation symbol and  $t_1, \dots, t_k$  are terms in  $\bar{x}$ , then

$$R^M(t_1^M(\bar{a}), \dots, t_k^M(\bar{a})) \iff R(t_1, \dots, t_k) \in p \iff R^N(t_1^N(\bar{a}), \dots, t_k^N(\bar{a})),$$

and so  $f$  preserves relation symbols. A similar argument shows  $f$  preserves function symbols.

(2)  $\implies$  (1): Let  $A = \langle \bar{a} \rangle_M$  and  $B = \langle \bar{b} \rangle_N$ . Then

$$\text{qftp}^M(\bar{a}) = \text{qftp}^A(\bar{a}) = \text{qftp}^B(\bar{b}) = \text{qftp}^N(\bar{b})$$

by Theorem 3.4.3 applied to the embeddings  $A \rightarrow M$ ,  $B \rightarrow N$ , and  $A \xrightarrow{\cong} B$ .  $\square$

**Theorem 8.3.5.** *Suppose  $T$  has quantifier elimination. If  $M, N \models T$ , then*

$$M \equiv N \iff \langle \emptyset \rangle_M \cong \langle \emptyset \rangle_N.$$

*Proof.* The left hand side says that  $\text{tp}^M() = \text{tp}^N()$ . Indeed,  $\text{tp}^M()$  is the set of formulas in 0 free variables (i.e., sentences) satisfied by the empty tuple (i.e., true in  $M$ ), which is just  $\text{Th}(M)$ .

Meanwhile, the right hand side of  $(*)$  is equivalent to  $\text{qftp}^M() = \text{qftp}^N()$  by Theorem 8.3.4. We don't have to worry about how the isomorphism acts on the generators, because there are no generators to check.

Finally, quantifier elimination gives

$$\text{tp}^M() = \text{tp}^N() \iff \text{qftp}^M() = \text{qftp}^N(). \quad \square$$

**Theorem 8.3.6.** *Suppose  $T$  has quantifier elimination and  $M, N \models T$ .*

1. *If  $f : M \rightarrow N$  is an embedding, then  $f$  is an elementary embedding.*
2. *If  $M$  is a substructure of  $N$ , then  $M$  is an elementary substructure of  $N$ .*

*Proof.* Embeddings preserve quantifier-free formulas (Theorem 3.4.3(3)), and quantifier elimination allows us to replace any formula with an equivalent quantifier-free formula.  $\square$

## 8.4 Quantifier elimination in DLO

**Theorem 8.4.1.** *DLO has quantifier elimination.*

*Proof.* We use the criterion of Theorem 8.3.3. Suppose  $M, N \models \text{DLO}$ ,  $\bar{a} \in M^n$ ,  $\bar{b} \in N^n$ , and  $\text{qftp}^M(\bar{a}) = \text{qftp}^N(\bar{b})$ . We must show  $\text{tp}^M(\bar{a}) \stackrel{?}{=} \text{tp}^N(\bar{b})$ .

By Theorem 8.3.4, there is a partial isomorphism  $f : \langle \bar{a} \rangle_M \rightarrow \langle \bar{b} \rangle_N$  with  $f(\bar{a}) = \bar{b}$ . As the language of orders is relational, all terms are trivial, and so  $\text{dom}(f) = \langle \bar{a} \rangle_M = \{a_1, \dots, a_n\}$ . Then  $f$  is a finite partial isomorphism. By Theorem 3.7.10, the class of finite partial isomorphisms is a back-and-forth system, and so  $f$  is a partial elementary map by Theorem 3.7.6. Therefore  $\text{tp}^M(\bar{a}) = \text{tp}^N(f(\bar{a})) = \text{tp}^N(\bar{b})$ .  $\square$

If  $M, N$  are models of DLO, then  $\langle \emptyset \rangle_M$  and  $\langle \emptyset \rangle_N$  are both the empty order  $\emptyset$ , so  $M \equiv N$ . This gives another proof of the completeness of DLO.

**Corollary 8.4.2.**  *$\mathbb{Q}$  and the open interval  $(0, 1) = \{x \in \mathbb{R} : 0 < x < 1\}$  are elementary substructures of  $\mathbb{R}$ .*

**Definition 8.4.3.** If  $(M, \leq) \models \text{DLO}$  and  $a < b$  are two points in  $M$ , then we define

$$\begin{aligned} (a, b) &:= \{x \in M : a < x < b\} \\ (a, +\infty) &:= \{x \in M : a < x\} \\ (-\infty, a) &:= \{x \in M : x < a\} \\ (-\infty, +\infty) &:= M \end{aligned}$$

Sets of these forms are called *open intervals*.

**Corollary 8.4.4.** *If  $M \models \text{DLO}$  and  $D \subseteq M^1$  is definable, then  $D$  is a finite union of points and open intervals.*

*Proof.* Let  $\mathcal{F}$  be the collection of sets  $D \subseteq M$  such that  $D$  is a finite union of points and open intervals. It is an exercise to see that  $\mathcal{F}$  is closed under boolean operations (it is a boolean subalgebra of  $\mathfrak{P}(M)$ ). We must show that  $D \in \mathcal{F}$ . Write  $D$  as  $\varphi(M, \bar{b})$  for some  $\mathcal{L}$ -formula  $\varphi(x, \bar{y})$  and tuple of parameters  $\bar{b}$ . By quantifier-elimination, we may assume  $\varphi$  is quantifier-free. Then  $\varphi$  is a boolean combination of atomic formulas. As  $\mathcal{F}$  is closed under

boolean combinations, we may assume  $\varphi$  is atomic. Then  $\varphi(x, \bar{b})$  has one of the following forms:

$$x \leq x, \ x \leq b_i, \ b_i \leq x, \ b_i \leq b_j, \ x = x, \ x = b_i, \ b_i = x, \ b_i = b_j.$$

Each of these formulas defines a set in  $\mathcal{F}$ .

□



# Chapter 9

## Algebraically closed fields

### 9.1 Polynomial rings

**Fact 9.1.1.** *Let  $R$  be a ring and  $x$  be a symbol. There is a ring  $R[x]$  extending  $R$ , generated by  $R \cup \{x\}$ , with the following properties:*

1. *Every element of  $R[x]$  has the form  $\sum_{i=0}^n a_i x^i$  for some  $n \geq 0$  and  $a_0, \dots, a_n \in R$ .*
2. *Two elements  $\sum_{i=0}^n a_i x^i$  and  $\sum_{i=0}^n b_i x^i$  are equal if and only if the tuples  $\bar{a}$  and  $\bar{b}$  are equal.*
3. *The sum of two elements is given by*

$$\sum_{i=0}^n a_i x^i + \sum_{i=0}^n b_i x^i = \sum_{i=0}^n (a_i + b_i) x^i.$$

4. *The product of two elements is given by*

$$\left( \sum_{i=0}^n a_i x^i \right) \left( \sum_{j=0}^m b_j x^j \right) = \sum_{k=0}^{n+m} c_k x^k, \text{ where } c_k = \sum_{\substack{0 \leq i \leq n \\ 0 \leq j \leq m \\ i+j=k}} a_i b_j.$$

*Elements of  $R[x]$  are called polynomials.*

If  $P(x)$  is a non-zero polynomial, then we can write

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

where  $n \geq 0$  and  $a_n \neq 0$ . Then  $a_n x^n$  is called the *leading term*,  $a_n$  is called the *leading coefficient*, and  $n$  is called the *degree* of  $P(x)$ . We write the degree of  $P$  as  $\deg P$ . The degree of the zero polynomial is defined to be  $-\infty$ . A nonzero polynomial is *monic* if the leading coefficient is 1.

**Definition 9.1.2** (Evaluating polynomials). If  $P(x) \in R[x]$  has the form

$$P(x) = \sum_{i=0}^n a_i x^i$$

and if  $b \in R$ , then

$$P(b) := \sum_{i=0}^n a_i b^i.$$

**Fact 9.1.3.** For fixed  $b$ , the map

$$\begin{aligned} R[x] &\rightarrow R \\ P(x) &\rightarrow P(b) \end{aligned}$$

is a ring homomorphism, meaning among other things that

$$\begin{aligned} (P + Q)(b) &= P(b) + Q(b) \\ (PQ)(b) &= P(b)Q(b) \end{aligned}$$

**Lemma 9.1.4.** Let  $K$  be a field, and let  $A(x), B(x) \in K[x]$  be polynomials with  $B(x)$  non-zero. then there is  $R(x) \in K[x]$  such that

$$\begin{aligned} A(x) &\equiv R(x) \pmod{B(x)} \\ \deg R(x) &< \deg B(x). \end{aligned}$$

*Proof.* If  $bx^m$  is the leading term of  $B(x)$ , replace  $B(x)$  with  $b^{-1}B(x)$ . Then we can assume  $B(x)$  is monic. Proceed by induction on  $\deg A(x)$ . If  $\deg A(x) < \deg B(x)$ , take  $R(x) = A(x)$ . Otherwise, let

$$\begin{aligned} A(x) &= a_n x^n + a_{n-1} x^{n-1} + \cdots \\ B(x) &= x^m + b_{m-1} x^{m-1} + \cdots \end{aligned}$$



where  $m = \deg(B) \leq n = \deg(A)$ . Let

$$\begin{aligned} A'(x) &= A(x) - a_n x^{n-m} B(x) \\ &= (a_n x^n + a_{n-1} x^{n-1} + \cdots) - (a_n x^n + a_n b_{m-1} x^{n-1} + \cdots + a_n b_0 x^{n-m}). \end{aligned}$$

Then  $\deg A'(x) < n = \deg A(x)$ , but  $A(x) \equiv A'(x) \pmod{B(x)}$ . By induction,  $A'(x) \equiv R(x) \pmod{B(x)}$  for some suitable  $R(x)$ .  $\square$

Let  $K$  be a field.

**Theorem 9.1.5.** *Suppose  $P(x) \in K[x]$  and  $P(a) = 0$  for some  $a \in K$ . Then  $P(x) = (x - a)Q(x)$  for some  $Q(x) \in K[x]$ .*

*Proof.* Apply the division lemma to  $P(x)$  and  $(x - a)$ , to get  $R(x) \in K[x]$  with

$$\begin{aligned} P(x) &\equiv R(x) \pmod{x - a} \\ \deg R(x) &< \deg(x - a) = 1. \end{aligned}$$

The first line means there is  $Q(x) \in K[x]$  with

$$P(x) = (x - a)Q(x) + R(x). \quad (*)$$

The second line means  $R(x) = c$  for some constant  $c$ . Substituting  $x = a$  into  $(*)$ , we see

$$\begin{aligned} P(a) &= (a - a)Q(a) + c \\ 0 &= 0 + c. \end{aligned}$$

Then  $c = 0$ , so  $R(x) = 0$  and  $P(x) = (x - a)Q(x)$ .  $\square$

A *root* of  $P(x) \in K[x]$  is an element  $a \in K$  with  $P(a) = 0$ . Note that  $a$  is a root of  $P(x)Q(x)$  if and only if  $a$  is a root of  $P(x)$  or  $a$  is a root of  $Q(x)$ , by the zero law (Theorem 1.4.16).

**Theorem 9.1.6.** *If  $P(x)$  is a non-zero polynomial, then the number of roots of  $P(x)$  in  $K$  is at most  $\deg P(x)$ .*

*Proof.* Let  $d = \deg P(x)$ . If  $P(x)$  has no roots, then the claim holds. Otherwise, take some root  $a$ . Then  $P(x) = (x - a)Q(x)$  for some polynomial  $Q(x)$  of degree  $d - 1$ . By induction on  $d$ ,  $Q(x)$  has at most  $d - 1$  roots. As  $(x - a)$  has one root,  $P(x)$  has at most  $(d - 1) + 1 = d$  roots.  $\square$

**Theorem 9.1.7.** *If  $K$  is a field, every ideal  $I \subseteq K[x]$  is a principal ideal  $P(x)K[x]$  for some polynomial  $P(x)$ .*

*Proof.* Like the proof of the same fact in  $\mathbb{Z}$  (Theorem 2.5.3), but using Lemma 9.1.4 instead of Lemma 2.5.2 and using  $\deg P$  instead of  $|n|$ .  $\square$

**Remark 9.1.8.** In Theorem 9.1.7, scaling  $P(x)$  by a non-zero constant, we may assume  $P(x)$  is zero or monic. Then  $P(x)$  is uniquely determined.

**Definition 9.1.9.** A non-constant polynomial  $P(x)$  is *reducible* if  $P(x)$  is a product of two non-constant polynomials, and *irreducible* otherwise.

**Theorem 9.1.10.** *If  $P(x)$  is an irreducible polynomial in  $K[x]$ , then the quotient  $K[x]/P(x)K[x]$  is a field.*

*Proof.* Like the proof that  $\mathbb{Z}/p\mathbb{Z}$  is a field when  $p$  is prime (Theorem 2.5.7).  $\square$

## 9.2 The theory ACF

**Definition 9.2.1.** A field  $K$  is *algebraically closed* if every polynomial  $P(x) \in K[x]$  with  $\deg P > 0$  has a root. That is, for every  $n > 0$  and  $a_0, a_1, \dots, a_n \in K$  with  $a_n \neq 0$ , there is  $x \in K$  with

$$a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0 = 0.$$

Note that the class of algebraically closed fields is elementary, defined by a theory consisting of the field axioms plus an axiom schema

$$\forall a_1, \dots, a_n \left( a_n \neq 0 \rightarrow \exists x \sum_{i=0}^n a_i x^i = 0 \right) \text{ for } n > 0.$$

This theory is usually called *ACF*.

**Fact 9.2.2** (Fundamental theorem of algebra). *The field  $\mathbb{C} = \{x + yi : x, y \in \mathbb{R}\}$  is algebraically closed.*

**Theorem 9.2.3.** *If  $K$  is algebraically closed, then  $K$  is infinite.*

*Proof.* If  $K = \{a_1, \dots, a_n\}$ , then  $P(x) = 1 + \prod_{i=1}^n (x - a_i)$  has no root in  $K$ , so  $K \not\models \text{ACF}$ .  $\square$

## 9.3 Algebraic and transcendental elements

Let  $L$  be a field and  $K$  be a subfield.

**Definition 9.3.1.** An element  $a \in L$  is *algebraic* over  $K$  if  $P(a) = 0$  for some non-zero polynomial  $P(x) \in K[x]$ . Otherwise,  $a$  is *transcendental* over  $K$ .

If  $a \in L$ , then  $K[a]$  denotes the subring of  $L$  generated by  $K \cup \{a\}$ .

**Theorem 9.3.2.** Fix  $a \in L$ . Let  $I_{a/K} = \{P(x) \in K[x] : P(a) = 0\}$ .

1.  $I_{a/K}$  is an ideal in  $K[x]$ .
2.  $K[x]/I_{a/K}$  is isomorphic to  $K[a]$  via the isomorphism sending  $P(x)$  to  $P(a)$ .
3. If  $a$  is transcendental over  $K$ , then  $I_{a/K}$  is the zero ideal  $0 \cdot K[x] = \{0\}$ .
4. If  $a$  is algebraic over  $K$ , then  $I_{a/K} = P(x) \cdot K[x]$  for some irreducible monic polynomial  $P(x)$ .

*Proof.* Let  $f : K[x] \rightarrow K[a]$  be the ring homomorphism  $P(x) \mapsto P(a)$ . Then  $I_{a/K}$  is the kernel. The image  $\text{im}(f)$  is a subring of  $K[a]$  containing  $f(K) = K$  and  $f(x) = a$ , so it must be all of  $K[a]$ . By the fundamental theorem on homomorphisms,  $K[x]/I_{a/K}$  is isomorphic to the image  $K[a]$  via the map  $P(x) \mapsto P(a)$ .

By Theorem 9.1.7 and Remark 9.1.8, the ideal  $I_{a/K}$  is  $P(x) \cdot K[x]$ , where  $P(x)$  is zero or a monic polynomial. In the first case,  $I_{a/K} = \{0\}$ , which means precisely that  $a$  is transcendental. In the second case,  $P(x) \in I_{a/K}$  implies that  $P(a) = 0$ , and so  $a$  is algebraic. Suppose for the sake of contradiction that  $P(x)$  is reducible as  $P(x) = Q_1(x)Q_2(x)$ . Note that  $\deg P > \deg Q_i$  for  $i = 1, 2$ , so  $Q_i \notin P(x) \cdot K[x] = I_{a/K}$ , and therefore  $Q_i(a) \neq 0$  for  $i = 1, 2$ . But then  $0 = P(a) = Q_1(a)Q_2(a) \neq 0$ , a contradiction.  $\square$

**Definition 9.3.3.** If  $a \in L$  is algebraic over  $K$ , then the *minimal polynomial* of  $a$  over  $K$  is the monic irreducible polynomial  $P(x)$  appearing in Theorem 9.3.2(4).

**Theorem 9.3.4.** If  $P(x) \in K[x]$  is a monic irreducible polynomial and  $a \in L$  is a root of  $P(x)$ , then  $P(x)$  is the minimal polynomial of  $a$ .

*Proof.* Let  $P_0(x)$  be the actual minimal polynomial of  $a$ . Then

$$P(a) = 0 \implies P(x) \in I_{a/K} = P_0(x) \cdot K[x].$$

Thus  $P(x)$  is a multiple of  $P_0(x)$ :

$$P(x) = P_0(x)Q(x).$$

As  $P(x)$  is irreducible, this forces  $Q(x) = 1$  and  $P(x) = P_0(x)$ .  $\square$

## 9.4 Quantifier elimination in ACF

**Lemma 9.4.1.** *Let  $M_1, M_2$  be two fields. Let  $f : R_1 \rightarrow R_2$  be a partial isomorphism. Then there is a larger partial isomorphism  $g : K_1 \rightarrow K_2$  such that  $K_1$  and  $K_2$  are fields.*

*Proof.* Let  $K_i = \{a/b : a, b \in R_i, b \neq 0\}$ . Then  $K_i$  is a subfield of  $M_i$ . For example,  $K_i$  is closed under addition because

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd}.$$

Let  $g : K_1 \rightarrow K_2$  be defined by  $g(a/b) = f(a)/f(b)$ . This is well-defined because

$$\begin{aligned} \frac{a}{b} = \frac{a'}{b'} &\iff ab' = a'b \implies f(ab') = f(a'b) \\ &\iff f(a)f(b') = f(a')f(b) \iff \frac{f(a)}{f(b)} = \frac{f(a')}{f(b')}. \end{aligned}$$

It is an exercise in algebra to see that  $g$  is an isomorphism. For example,  $g$  preserves addition because

$$\begin{aligned} g\left(\frac{a}{b} + \frac{c}{d}\right) &= g\left(\frac{ad + bc}{bd}\right) = \frac{f(ad + bc)}{f(bd)} = \frac{f(a)f(d) + f(b)f(c)}{f(b)f(d)} \\ &= \frac{f(a)}{f(b)} + \frac{f(c)}{f(d)} = g\left(\frac{a}{b}\right) + g\left(\frac{c}{d}\right). \end{aligned} \quad \square$$

**Lemma 9.4.2.** *Let  $M_1, M_2$  be two uncountable algebraically closed fields extending a countable subfield  $K$ . For any  $a \in M_1$ , there is  $b \in M_2$  and an isomorphism  $f : K[a] \rightarrow K[b]$  sending  $a$  to  $b$  and fixing  $K$ .*

*Proof.* Recall the notation

$$I_{a/K} = \{P(x) \in K[x] : P(a) = 0\}$$

from Theorem 9.3.2.

*Claim.* There is  $b \in M_2$  with  $I_{a/K} = I_{b/K}$ .

*Proof.* First suppose  $a$  is transcendental. There are countably many non-zero polynomials in  $K[x]$ , and each has finitely many roots (Theorem 9.1.6). Therefore only countably many  $b \in M_2$  are algebraic over  $K$ . Take  $b \in M_2$  transcendental over  $K$ . Then  $I_{a/K} = \{0\} = I_{b/K}$  by Theorem 9.3.2.

Next suppose  $a$  is algebraic with minimal polynomial  $P(x) \in K[x]$ . As  $M_2$  is algebraically closed, there is  $b \in M_2$  with  $P(b) = 0$ . By Theorem 9.3.4,  $P(x)$  is the minimal polynomial of  $b$ , and then

$$I_{a/K} = P(x) \cdot K[x] = I_{b/K}. \quad \square_{\text{Claim}}$$

Fix  $b \in M_2$  as in the claim, and let  $I = I_{a/K} = I_{b/K}$ . By Theorem 9.3.2, we have isomorphisms

$$\begin{aligned} K[x]/I &\rightarrow K[a] \\ K[x]/I &\rightarrow K[b] \end{aligned}$$

sending  $P(x)$  to  $P(a)$  and  $P(b)$ , respectively. The composition

$$K[a] \rightarrow K[x]/I \rightarrow K[b]$$

is the desired isomorphism.  $\square$

**Lemma 9.4.3.** *Let  $M_1, M_2$  be two uncountable algebraically closed fields. Let  $f : R_1 \rightarrow R_2$  be a finitely-generated partial isomorphism. For any  $a \in M_1$ , there is  $b \in M_2$  and an isomorphism  $R_1[a] \rightarrow R_2[b]$  extending  $f$ .*

*Proof.* By Lemma 9.4.1, the isomorphism  $f : R_1 \rightarrow R_2$  extends to an isomorphism  $f' : K_1 \rightarrow K_2$  where the  $K_i$  are fields. Moving  $M_2$ , we may assume  $K_1 = K_2$  and  $f'$  is the identity map. Applying Lemma 9.4.2, there is an element  $b \in M_2$  and an isomorphism  $K_1[b] \rightarrow K_2[b]$  extending  $f'$ . This isomorphism restricts to an isomorphism  $R_1[b] \rightarrow R_2[b]$  extending  $f$ .  $\square$

**Theorem 9.4.4.** *The theory ACF has quantifier elimination.*

*Proof.* We use the criterion of Theorem 8.3.3. Suppose  $M, N \models \text{ACF}$ ,  $\bar{a} \in M^n$ ,  $\bar{b} \in N^n$ , and  $\text{qftp}^M(\bar{a}) = \text{qftp}^N(\bar{b})$ . We must show  $\text{tp}^M(\bar{a}) = \text{tp}^N(\bar{b})$ . Recall that  $M$  and  $N$  are infinite (Theorem 9.2.3). Replacing  $M, N$  with elementary extensions, we may assume  $M$  and  $N$  are uncountable. By Theorem 8.3.4, there is a partial isomorphism  $f : \langle \bar{a} \rangle_M \rightarrow \langle \bar{b} \rangle_N$  with  $f(\bar{a}) = \bar{b}$ . Let  $\mathcal{F}$  be the collection of finitely generated partial isomorphisms from  $M$  to  $N$ . Then  $\mathcal{F}$  is a back-and-forth system: Lemma 9.4.3 gives the forward condition, and the backward condition holds by symmetry. By Theorem 3.7.6, the fact that  $f \in \mathcal{F}$  implies that  $f$  is a partial elementary map, and so  $\text{tp}^M(\bar{a}) = \text{tp}^N(\bar{b})$ .  $\square$

**Corollary 9.4.5.** *If  $M, N \models \text{ACF}$ , then  $M \equiv N \iff \text{char}(M) = \text{char}(N)$ .*

*Proof.* By quantifier elimination in ACF and Theorem 8.3.5, we have

$$M \equiv N \iff \langle \emptyset \rangle_M \cong \langle \emptyset \rangle_N.$$

By Theorem 2.7.5, the right hand side is equivalent to

$$\mathbb{Z}/n_M\mathbb{Z} \stackrel{?}{\cong} \mathbb{Z}/n_N\mathbb{Z}, \tag{*}$$

where  $n_M = \text{char}(M)$  and  $n_N = \text{char}(N)$ . Clearly,  $(*)$  holds if and only if  $n_M = n_N$ .  $\square$

**Definition 9.4.6.** For  $p \in \{0, 2, 3, 5, 7, 11, 13, \dots\}$ ,  $\text{ACF}_p$  is the theory of algebraically closed fields of characteristic  $p$ :

$$\begin{aligned} \text{ACF}_0 &= \text{ACF} \cup \left\{ \underbrace{1 + 1 + \dots + 1}_{p \text{ times}} \neq 0 : p = 2, 3, 5, 7, \dots \right\} \\ \text{ACF}_p &= \text{ACF} \cup \left\{ \underbrace{1 + 1 + \dots + 1}_{p \text{ times}} = 0 \right\} \text{ for } p > 0 \end{aligned}$$

For example,  $\mathbb{C} \models \text{ACF}_0$ . We will see later (Corollary 10.3.5) that  $\text{ACF}_p$  is consistent for each  $p$ .

**Corollary 9.4.7.** *Each theory  $\text{ACF}_p$  is complete.*

**Corollary 9.4.8.**  *$\text{Th}(\mathbb{C})$  is decidable.*

# Chapter 10

## Existentially closed models

### 10.1 Unions of chains of structures

Fix a language  $\mathcal{L}$ .

**Definition 10.1.1.** A *chain of  $\mathcal{L}$ -structures* is a family  $\{M_i\}_{i \in I}$  where  $(I, \leq)$  is a linear order,  $M_i$  is an  $\mathcal{L}$ -structure for  $i \in I$ , and  $M_i$  is a substructure of  $M_j$  for  $i \leq j$ .

If  $\{M_i\}_{i \in I}$  is a chain of  $\mathcal{L}$ -structures, we can make  $M = \bigcup_{i \in I} M_i$  into a  $\mathcal{L}$ -structure by defining

$$\begin{aligned} f^M(a_1, \dots, a_n) &= f^{M_i}(a_1, \dots, a_n) \\ R^M(a_1, \dots, a_n) &\iff R^{M_i}(a_1, \dots, a_n) \end{aligned}$$

for any  $i \in I$  large enough that  $\{a_1, \dots, a_n\} \subseteq M_i$ . The choice of  $i$  doesn't matter—if  $j$  is another choice, then

$$\begin{aligned} f^{M_i}(\bar{a}) &= f^{M_j}(\bar{a}) \\ R^{M_i}(\bar{a}) &\iff R^{M_j}(\bar{a}) \end{aligned}$$

because  $M_i$  is a substructure or extension of  $M_j$ .

**Definition 10.1.2.** A theory  $T$  is *inductive* if whenever  $\{M_i\}_{i \in I}$  is a chain of models of  $T$ , the union  $\bigcup_{i \in I} M_i$  is also a model of  $T$ .

**Definition 10.1.3.** An  $\forall\exists$ -*sentence* is one of the form  $\forall \bar{x} \exists \bar{y} \varphi(\bar{x}, \bar{y})$ , where  $\varphi$  is quantifier-free. An  $\forall\exists$ -*theory* is a set of  $\forall\exists$ -sentences.

**Theorem 10.1.4.** *If  $T$  is an  $\forall\exists$ -theory, then  $T$  is inductive.*

*Proof.* Suppose  $\{M_i\}_{i \in I}$  is a chain of structures, and

$$M_i \models \forall \bar{x} \exists \bar{y} \varphi(\bar{x}, \bar{y}) \quad (*)$$

for all  $i$ , where  $\varphi$  is quantifier-free. Let  $M = \bigcup_i M_i$ . If  $\bar{a} \in M^n$ , then  $\bar{a} \in M_i^n$  for large enough  $i$ . By  $(*)$ , there is  $\bar{b} \in M_i^m$  with  $M_i \models \varphi(\bar{a}, \bar{b})$ . Then  $M \models \varphi(\bar{a}, \bar{b})$  because  $\varphi$  is quantifier-free. We have shown

$$M \models \forall \bar{x} \exists \bar{y} \varphi(\bar{x}, \bar{y}). \quad \square$$

**Example 10.1.5.** The theory of fields is an  $\forall\exists$ -theory, so it is inductive.

## 10.2 Existentially closed models

**Definition 10.2.1.** Let  $M \subseteq N$  be structures. Then  $M$  is *existentially closed* in  $N$ , written  $M \preceq_1 N$ , if for any quantifier-free  $\mathcal{L}(M)$ -formula  $\varphi(\bar{x})$ ,

$$N \models \exists \bar{x} \varphi(\bar{x}) \implies M \models \exists \bar{x} \varphi(\bar{x}).$$

**Remark 10.2.2.**  $M \preceq N \implies M \preceq_1 N$ .

Fix a theory  $T$ . “**Model**” will always mean **model of  $T$** .

**Definition 10.2.3.** A model  $M \models T$  is *existentially closed* if for any larger model  $N \models T$  with  $N \supseteq M$ , we have  $M \preceq_1 N$ .

**Theorem 10.2.4.** *If  $T$  has quantifier-elimination, then every model of  $T$  is existentially closed.*

*Proof.* Suppose  $M \subseteq N$  are models of  $T$ . Then  $M \preceq N$  by quantifier elimination (Theorem 8.3.6), and so  $M \preceq_1 N$ .  $\square$

**Theorem 10.2.5.** *If  $T$  is inductive, then any model  $M$  embeds into an existentially closed model  $N \supseteq M$ .*

*Proof.* If  $M \models T$  and  $\varphi(\bar{x})$  is a quantifier-free  $\mathcal{L}(M)$ -formula, say that  $\varphi(\bar{x})$  is *realizable beyond  $M$*  if it is realized in some larger model  $N \supseteq M$ . We make two observations:



1. A model  $M$  is existentially closed iff every quantifier-free  $\mathcal{L}(M)$ -formula that is realizable beyond  $M$  is already realized in  $M$ .
2. If  $N$  extends  $M$  and a quantifier-free  $\mathcal{L}(M)$ -formula  $\varphi$  is realizable beyond  $N$ , then it is realizable beyond  $M$ —the extension of  $N$  where  $\varphi$  is realized is an extension of  $M$ .

*Claim.* For any model  $M$ , there is a larger model  $M^* \supseteq M$  such that every quantifier-free  $\mathcal{L}(M)$ -formula that is realizable beyond  $M^*$  is realized in  $M^*$ .

*Proof.* Let  $\{\varphi_\alpha\}_{\alpha < \kappa}$  enumerate all quantifier-free  $\mathcal{L}(M)$ -formulas. Build an increasing chain of models  $\{M_\alpha\}_{\alpha < \kappa}$  as follows:

1.  $M_0 = M$ .
2. If  $\varphi_\alpha$  is realizable beyond  $M_\alpha$ , then  $M_{\alpha+1}$  is an extension of  $M_\alpha$  with a realization of  $\varphi_\alpha$ . Otherwise take  $M_{\alpha+1} = M_\alpha$ .
3. If  $\alpha$  is a limit ordinal, take  $M_\alpha = \bigcup_{\beta < \alpha} M_\beta$ . This is a model because the theory is inductive.

Let  $M^* = \bigcup_{\alpha < \kappa} M_\alpha$ . Again, this is a model because  $T$  is inductive. Suppose  $\varphi$  is a quantifier-free  $\mathcal{L}(M)$ -formula that is realizable beyond  $M^*$ . Then  $\varphi = \varphi_\alpha$  for some  $\alpha$ . By observation (2) above,  $\varphi_\alpha$  is realizable beyond  $M_\alpha \subseteq M^*$ . Then  $\varphi_\alpha$  is realized in  $M_{\alpha+1}$  by construction. That is,  $M_{\alpha+1} \models \varphi_\alpha(\bar{c})$  for some tuple  $\bar{c}$  in  $M_{\alpha+1}$ . As  $M^*$  extends  $M_{\alpha+1}$  and  $\varphi_\alpha$  is quantifier-free,  $M^* \models \varphi_\alpha(\bar{c})$ .

□<sub>Claim</sub>

Using the claim, build an increasing chain of length  $\omega$ :

$$M = M_0 \subseteq M_1 \subseteq M_2 \subseteq \cdots$$

where  $M_{i+1} = M_i^*$ . Let  $N = \bigcup_{n=0}^{\infty} M_n$ . Again,  $N \models T$  because  $T$  is inductive. We claim  $N$  is existentially closed. Take a quantifier-free  $\mathcal{L}(N)$ -formula  $\varphi$  that is realizable beyond  $N$ . Then  $\varphi$  is an  $\mathcal{L}(M_i)$ -formula for some  $i < \omega$ . By observation (2),  $\varphi$  is realizable beyond  $M_{i+1} = M_i^*$ , so  $\varphi$  is realized in  $M_{i+1}$ . Then  $\varphi$  is realized in  $N$ , and  $N$  is existentially closed by observation (1). □

### 10.3 Existentially closed fields

**Lemma 10.3.1.** *Let  $K, L$  be fields, and  $\alpha : K \rightarrow L$  be a ring homomorphism. Then  $\alpha$  is injective.*

*Proof.* Let  $I = \ker(\alpha)$ . We claim  $I = \{0\}$ . Otherwise take  $a \in I \setminus \{0\}$ . Then  $1 = a^{-1}a \in I$ , so  $1^L = \alpha(1^K) = 0^L$ , contradicting the definition of “field.”

As  $\ker(\alpha) = \{0\}$ , it follows that

$$\alpha(x) = \alpha(y) \iff x - y \in \ker(\alpha) \iff x - y = 0 \iff x = y.$$

This means that  $\alpha$  is injective. □

**Lemma 10.3.2.** *If  $K$  is a field and  $P(x)$  is a polynomial of positive degree, then there is a larger field  $L \supseteq K$  in which  $P(x)$  has a root.*

*Proof.* Write  $P(x)$  as a product  $\prod_{i=1}^n Q_i(x)$  of irreducible factors. Replacing  $P(x)$  with one of its irreducible factors, we may assume  $P(x)$  is irreducible. Then  $K[x]/P(x)K[x]$  is a field (Theorem 9.1.10). The composition

$$K \xrightarrow{\subseteq} K[x] \rightarrow K[x]/P(x)K[x]$$

is a homomorphism of fields, hence an embedding (Lemma 10.3.1). Up to isomorphism,  $L := K[x]/P(x)K[x]$  is a field extending  $K$ . The fact that

$$P(x) \equiv 0 \pmod{P(x)} \text{ in } K[x]$$

means that

$$P(x) = 0 \text{ in } L$$

and so the element  $x$  is a root of  $P$  in  $L$ . □

An *existentially closed field* is a field that is existentially closed among the class of all fields.

**Theorem 10.3.3.** *If  $K$  is an existentially closed field, then  $K$  is algebraically closed.*

*Proof.* Let  $P(x) \in K[x]$  be a polynomial of positive degree. By Lemma 10.3.2, the quantifier-free  $\mathcal{L}(K)$ -formula  $P(x) = 0$  is realized in a field extending  $K$ . As  $K$  is existentially closed, it is realized in  $K$ , meaning that  $P(x)$  has a root. □

**Corollary 10.3.4.** *If  $K$  is a field, then there is an algebraically closed field  $L$  extending  $K$ .*

*Proof.* The class of fields is inductive, so Theorem 10.2.5 applies.  $\square$

**Corollary 10.3.5.** *For every  $p \in \{0, 2, 3, 5, 7, \dots\}$ , there are algebraically closed fields of characteristic  $p$ .*

*Proof.* Take a field  $K$  of characteristic  $p$  (Theorem 2.7.7) and an algebraically closed field  $M \supseteq K$  by Corollary 10.3.4. Then  $\text{char}(M) = \text{char}(K) = p$  because characteristic doesn't change in field extensions (Theorem 2.7.10).  $\square$

**Theorem 10.3.6.** *A field  $K$  is existentially closed if and only if it is algebraically closed.*

*Proof.* If  $K$  is existentially closed, then  $K$  is algebraically closed by Theorem 10.3.3. Conversely, suppose  $K$  is algebraically closed. We claim  $K$  is existentially closed. Let  $L \supseteq K$  be an extension. By Corollary 10.3.4 there is an algebraically closed field  $M \supseteq L \supseteq K$ . Because ACF has quantifier elimination, algebraically closed fields are existentially closed *among algebraically closed fields*, and so  $K \preceq_1 M$ . But this implies  $K \preceq_1 L$ : if a quantifier-free  $\mathcal{L}(K)$ -formula  $\varphi$  is realized in  $L$ , then it is realized in  $M$ , hence realized in  $K$ .  $\square$



# Chapter 11

## Monster models

### 11.1 Pushing types along partial elementary maps

**Theorem 11.1.1.** *Let  $M, N$  be  $\mathcal{L}$ -structures and  $f : A \rightarrow B$  be a partial elementary map from  $M$  to  $N$ . If  $\Sigma(\bar{x})$  is a set of  $\mathcal{L}(A)$ -formulas, let  $f_*\Sigma(\bar{x})$  be*

$$\{\varphi(\bar{x}, f(\bar{c})) : \bar{c} \in A, \varphi(\bar{x}, \bar{c}) \in \Sigma(\bar{x})\}.$$

1. *If  $\Sigma(\bar{x})$  is finite, then  $\Sigma(\bar{x})$  is realized in  $M$  if and only if  $f_*\Sigma(\bar{x})$  is realized in  $N$ .*
2.  *$\Sigma(\bar{x})$  is finitely realized in  $M$  if and only if  $f_*\Sigma(\bar{x})$  is finitely realized in  $N$ .*
3.  *$f_*$  gives a bijection between partial types over  $A$  and partial types over  $B$ .*
4.  *$f_*$  gives a bijection between  $S_n(A)$  and  $S_n(B)$ .*

*Proof.* 1. Write  $\Sigma(\bar{x})$  as  $\{\varphi_i(\bar{x}, \bar{c}_i) : 1 \leq i \leq n\}$ . Then

$$M \models \exists \bar{x} \bigwedge_{i=1}^n \varphi_i(\bar{x}, \bar{c}_i) \iff N \models \exists \bar{x} \bigwedge_{i=1}^n \varphi_i(\bar{x}, f(\bar{c}_i)),$$

because  $f$  is a partial elementary map.

2. Clear from (1).

3. By (2),  $f_*$  is a map from partial types over  $A$  to partial types over  $B$ . The inverse map is  $(f^{-1})_*$ .
4. Clear from (3), since  $S_n(A)$  and  $S_n(B)$  are exactly the maximal partial types over  $A$  and  $B$ .  $\square$

**Corollary 11.1.2.** *If  $A \subseteq M \preceq N$ , then  $S_n^M(A) = S_n^N(A)$ .*

*Proof.* The identity map  $\text{id}_A : A \rightarrow A$  is a partial elementary map from  $M$  to  $N$ .  $\square$

## 11.2 $\kappa$ -saturated models

Let  $\kappa$  be an infinite cardinal.

**Definition 11.2.1.** A structure  $M$  is  $\kappa$ -saturated if for every  $A \subseteq M$  with  $|A| < \kappa$  and every  $p \in S_1(A)$ ,  $p$  is realized in  $M$ .

Fix  $\mathcal{L}$ -structures  $N, M$ , where  $M$  is  $\kappa$ -saturated.

**Lemma 11.2.2.** *Let  $f : A \rightarrow B$  be a partial elementary map from  $N$  to  $M$ , with  $|A| < \kappa$ . For any  $\alpha \in N$  there is  $\beta \in M$  such that  $f \cup \{(\alpha, \beta)\}$  is a partial elementary map.*

*Proof.* Let  $p = \text{tp}(\alpha/A) \in S_1(A)$ . Let  $f_*p = \{\varphi(x, f(\bar{a})) : \varphi(x, \bar{a}) \in p\}$ . By Theorem 11.1.1,  $f_*p \in S_1(B)$ . As  $|B| = |A| < \kappa$ , there is  $\beta \in M$  realizing  $f_*p$ . Then

$$\begin{aligned} N \models \varphi(\alpha, \bar{c}) &\iff \varphi(x, \bar{c}) \in p(x) \\ &\iff \varphi(x, f(\bar{c})) \in f_*p(x) \iff M \models \varphi(\beta, f(\bar{c})) \end{aligned}$$

for any  $\mathcal{L}$ -formula  $\alpha$  and tuple  $\bar{c}$  in  $A$ , so  $f \cup \{(\alpha, \beta)\}$  is a partial elementary map.  $\square$

**Lemma 11.2.3.** *Let  $f : A \rightarrow B$  be a partial elementary map from  $N$  to  $M$ . Suppose  $A \subseteq A' \subseteq N$ ,  $|A| < \kappa$ , and  $|A'| \leq \kappa$ . Then  $f$  can be extended to a partial elementary map  $f : A' \rightarrow B'$ .*

*Proof.* Write  $A'$  as  $\{a_\alpha : \alpha < \kappa\}$ . Recursively choose partial elementary maps  $f_\alpha$  for  $\alpha < \kappa$  as follows:

- $f_0 = f$ .
- $f_{\alpha+1} = f_\alpha \cup \{(a_\alpha, b)\}$  for some  $b \in M$ .
- $f_\beta = \bigcup_{\alpha < \beta} f_\alpha$  if  $\beta$  is a limit ordinal.

The successor step  $\alpha + 1$  works because  $\text{dom}(f_\alpha) \leq |A| + |\alpha| < \kappa$ , so Lemma 11.2.2 applies.

Let  $g = \bigcup_{\alpha < \kappa} f_\alpha$ . Then  $g$  is a partial elementary map with domain  $A'$ .  $\square$

**Theorem 11.2.4** ( $\kappa$ -universality). *If  $M$  is  $\kappa$ -saturated and  $N \equiv M$  with  $|N| \leq \kappa$ , then there is an elementary embedding  $g : N \rightarrow M$ . In particular,  $N$  is isomorphic to an elementary substructure of  $M$ .*

*Proof.* Note that  $\emptyset : \emptyset \rightarrow \emptyset$  is a partial elementary map from  $N$  to  $M$ . Use Lemma 11.2.3 to extend to a partial elementary map  $g$  with  $\text{dom}(g) = N$ . Then  $g$  is an elementary embedding  $N \rightarrow M$ .  $\square$

**Theorem 11.2.5.** *Suppose  $M$  is  $\kappa$ -saturated and  $A \subseteq M$  with  $|A| < \kappa$ . For any finite  $n < \omega$  and  $p \in S_n(A)$ ,  $p$  is realized in  $M$ .*

*Proof.* Take  $N \succeq M$  containing a realization  $\bar{b} = p$ . Note that  $\text{id}_A : A \rightarrow A$  is a partial elementary map from  $N$  to  $M$ . By Lemma 11.2.3, we can extend it to a partial elementary map  $f$  with  $\text{dom}(f) = A \cup \{a_1, \dots, a_n\}$ . Then

$$\varphi(\bar{x}, \bar{b}) \in p(\bar{x}) \implies N \models \varphi(\bar{a}, \bar{b}) \iff M \models \varphi(f(\bar{a}), \bar{b})$$

for any formula  $\varphi$  and tuple  $\bar{b}$  in  $A$ , and so  $f(\bar{a}) \in M^n$  realizes  $p$ .  $\square$

**Theorem 11.2.6.** *Suppose  $M$  is  $\kappa$ -saturated,  $A \subseteq M$  satisfies  $|A| < \kappa$ , and  $\Sigma(\bar{x})$  is a partial type over  $A$  in at most  $\kappa$  variables. Then  $\Sigma(\bar{x})$  is realized in  $M$ .*

*Proof.* Similar to Theorem 11.2.5.  $\square$

**Theorem 11.2.7** ( $\kappa$ -compactness). *Let  $M$  be  $\kappa$ -saturated.*

1. *Let  $\Sigma(\bar{x})$  be a partial  $n$ -type over  $M$ . If  $|\Sigma| < \kappa$ , then  $\Sigma(\bar{x})$  is realized in  $M$ .*
2. *Suppose  $|I| < \kappa$  and  $X_i$  is a definable subset of  $M^n$  for each  $i \in I$ . If  $\{X_i : i \in I\}$  has FIP, then  $\bigcap_i X_i \neq \emptyset$ .*

3. Suppose  $X \subseteq M^n$  is definable,  $|I| < \kappa$ ,  $Y_i \subseteq M^n$  is definable for  $i \in I$ , and  $X \subseteq \bigcup_{i \in I} Y_i$ . Then there is a finite  $I_0 \subseteq_f I$  such that  $X \subseteq \bigcup_{i \in I_0} Y_i$ .

*Proof.* 1. Let  $A$  be the set of parameters used in  $\Sigma(\bar{x})$ . Then  $|A| < \kappa$ , and  $\Sigma(\bar{x})$  is a partial  $n$ -type over  $A$ . Take a complete  $n$ -type  $p \in S_n(A)$  with  $p \supseteq \Sigma(\bar{x})$ . Then  $p$  is realized in  $M$  by Theorem 11.2.5.

2. Write  $X_i$  as  $\varphi_i(M^n)$  for some  $\mathcal{L}(M)$ -formula  $\varphi_i$ . Let  $\Sigma = \{\varphi_i : i \in I\}$ . The FIP means that  $\Sigma$  is finitely realized in  $M$ , i.e., a partial type. Apply (1) to find a point realizing  $\Sigma$ , i.e., a point in  $\bigcap_{i \in I} X_i$ .

3. If there is no finite subcover, then the family

$$\{X\} \cup \{M^n \setminus Y_i : i \in I\}$$

has the FIP. By (2), there is a point  $\bar{a} \in X \cap \bigcap_i (M^n \setminus Y_i)$ . Then  $\bar{a} \in X$  but  $\bar{a} \notin \bigcup_i Y_i$ , a contradiction.  $\square$

**Corollary 11.2.8.** *If  $M$  is  $\kappa$ -saturated and  $D \subseteq M^n$  is definable, then one of two things happens:*

1.  $D$  is finite.
2.  $|D| \geq \kappa$ .

*Proof.* Otherwise,  $\{\{p\} : p \in D\}$  is a small cover of  $D$  without a finite subcover, contradicting Theorem 11.2.7(3).  $\square$

### 11.3 Strongly $\kappa$ -homogeneous models

If  $M$  is a structure, then  $\text{Aut}(M)$  denotes the set of automorphisms of  $M$ , i.e., isomorphisms from  $M$  to  $M$ . If  $A \subseteq M$ , then

$$\text{Aut}(M/A) := \{\sigma \in \text{Aut}(M) : \forall x \in A \sigma(x) = x\}$$

**Definition 11.3.1.**  $M$  is *strongly  $\kappa$ -homogeneous* if any partial elementary map  $f$  from  $M$  to  $M$  with  $|\text{dom}(f)| < \kappa$  can be extended to an automorphism  $\sigma \in \text{Aut}(M)$ .

**Theorem 11.3.2.** *Suppose  $M$  is strongly  $\kappa$ -homogeneous. Suppose  $\bar{a}, \bar{b} \in M^n$ ,  $C \subseteq M$ , and  $|C| < \kappa$ . Then the following are equivalent:*



1.  $\text{tp}(\bar{a}/C) = \text{tp}(\bar{b}/C)$ .
2. There is  $\sigma \in \text{Aut}(M/C)$  with  $\sigma(\bar{a}) = \bar{b}$ .

*Proof.* (1)  $\implies$  (2): if  $\text{tp}(\bar{a}/C) = \text{tp}(\bar{b}/C)$ , then there is a partial elementary map

$$f : C \cup \{a_1, \dots, a_n\} \rightarrow C \cup \{b_1, \dots, b_n\}$$

$$f(x) = \begin{cases} x & \text{if } x \in C \\ b_i & \text{if } x = a_i \end{cases}$$

Then  $f$  extends to an automorphism  $\sigma \in \text{Aut}(M)$ . Note that  $\sigma \supseteq f \supseteq \text{id}_C$ , so  $\sigma \in \text{Aut}(M/C)$ , and  $\sigma(\bar{a}) = f(\bar{a}) = \bar{b}$ .

(2)  $\implies$  (1): isomorphisms preserve all formulas.  $\square$

**Definition 11.3.3.**  $\bar{a} \equiv_C \bar{b}$  means  $\text{tp}(\bar{a}/C) = \text{tp}(\bar{b}/C)$ .

**Informal Definition 11.3.4.** A *monster model* is a structure that is  $\kappa$ -saturated and strongly  $\kappa$ -homogeneous for some cardinal  $\kappa$  bigger than any cardinals we care about. A set  $X$  is “small” or “large” depending on whether  $|X| < \kappa$  or  $|X| \geq \kappa$ .

Work in a monster model  $\mathbb{M}$ . Fix a small set  $A \subseteq \mathbb{M}$ .

**Definition 11.3.5.** A set  $X \subseteq \mathbb{M}^n$  is *A-invariant* if  $\sigma(X) = X$  for all  $\sigma \in \text{Aut}(\mathbb{M}/A)$ .

**Theorem 11.3.6.** *The following are equivalent for  $X \subseteq \mathbb{M}^n$ :*

1.  $X$  is *A-invariant*.
2. If  $\bar{b}, \bar{c} \in \mathbb{M}^n$ , then

$$\text{tp}(\bar{b}/A) = \text{tp}(\bar{c}/A) \implies (\bar{b} \in A \iff \bar{c} \in A).$$

3. There is a subset  $X' \subseteq S_n(A)$  such that

$$X = \{\bar{b} \in \mathbb{M}^n : \text{tp}(\bar{b}/A) \in X'\}.$$

*Proof.* Note that (1) means the following: if  $\bar{a} \in \mathbb{M}^n$  and  $\sigma \in \text{Aut}(\mathbb{M}/C)$ , then

$$\bar{a} \in X \iff \sigma(\bar{a}) \in X.$$

This is equivalent to (2) by Theorem 11.3.2. The equivalence of (2) and (3) is clear.  $\square$

**Remark 11.3.7.** An  $A$ -definable set  $\varphi(\mathbb{M}^n)$  corresponds to the clopen set  $\llbracket \varphi \rrbracket \subseteq S_n(A)$ , so  $A$ -definable sets are  $A$ -invariant.

**Theorem 11.3.8.** *If  $D \subseteq \mathbb{M}^n$  is definable and  $A$ -invariant, then  $D$  is  $A$ -definable.*

*Proof.* Note that  $D$  is  $B$ -definable for some small  $B \supseteq A$ . Let  $f : S_n(B) \rightarrow S_n(A)$  be the restriction map sending  $\text{tp}(\bar{c}/B)$  to  $\text{tp}(\bar{c}/A)$ . This map is surjective because every complete type over  $A$  extends to a complete type over  $B$ . Additionally, it is continuous by Theorem 7.4.4, because the preimage of the clopen set  $\llbracket \varphi \rrbracket \subseteq S_n(A)$  is the clopen set  $\llbracket \varphi \rrbracket \subseteq S_n(B)$ . As  $X$  is  $A$ -invariant and  $B$ -invariant, there are sets  $X_A \subseteq S_n(A)$  and  $X_B \subseteq S_n(B)$  such that

$$\begin{aligned} \bar{c} \in X &\iff \text{tp}(\bar{c}/A) \in X_A \\ \bar{c} \in X &\iff \text{tp}(\bar{c}/B) \in X_B. \end{aligned}$$

Then  $X_B = f^{-1}(X_A)$ . As  $f$  is surjective

$$\begin{aligned} X_A &= f(X_B) \\ S_n(A) \setminus X_A &= f(S_n(B) \setminus X_B). \end{aligned}$$

Because  $X$  is  $B$ -definable the set  $X_B \subseteq S_n(B)$  is clopen, and so  $X_B$  and  $S_n(B) \setminus X_B$  are closed. The image of a closed set is closed (Theorem 7.2.6), so  $X_A$  and its complement are both closed. Then  $X_A$  is a clopen set  $\llbracket \psi \rrbracket$ , and  $X$  is the  $A$ -definable set  $\psi(\mathbb{M}^n)$ .  $\square$

## 11.4 Construction of monster models

A cardinal  $\kappa$  is *regular* if whenever  $|I| < \kappa$  and  $|X_i| < \kappa$  for every  $i \in I$ , we have  $|\bigcup_{i \in I} X_i| < \kappa$ . The cardinal  $\aleph_0$  is regular, because a finite union of finite sets is finite. For any cardinal  $\kappa$ , the successor  $\kappa^+$  is regular, because a union of at most  $\kappa$ -many sets of size at most  $\kappa$  has size at most  $\kappa^2 = \kappa < \kappa^+$ . Consequently, for any cardinal  $\kappa$  we can find a larger cardinal that is regular.

If  $\kappa$  is a regular cardinal and  $A \subseteq \kappa$  with  $|A| < \kappa$ , then  $\sup(A) < \kappa$ . Otherwise,  $\sup(A) = \kappa$ , and then  $\kappa$  is a union  $\bigcup_{\alpha \in A} \alpha$  where  $|A| < \kappa$  and  $|\alpha| < \kappa$  for each  $\alpha \in \kappa$ , contradicting regularity.

**Lemma 11.4.1.** *Let  $\kappa$  be a regular cardinal. Let  $\{S_\alpha\}_{\alpha < \kappa}$  be an increasing chain of sets, indexed by  $\kappa$ . If  $A \subseteq \bigcup_{\alpha < \kappa} S_\alpha$  and  $|A| < \kappa$ , then  $A \subseteq S_\alpha$  for some  $\alpha < \kappa$ .*

*Proof.* Define  $f : A \rightarrow \kappa$  by  $f(x) = \min\{\alpha < \kappa : x \in S_\alpha\}$ . Then  $|f(A)| \leq |A| < \kappa$ , so  $\alpha := \sup f(A) < \kappa$ . For any  $x \in A$ , we have  $f(x) \in f(A)$  and so  $f(x) \leq \alpha$ . Then  $x \in S_{f(x)} \subseteq S_\alpha$  for any  $x \in A$ , so  $A \subseteq S_\alpha$ .  $\square$

**Definition 11.4.2.** A chain of structures  $\{M_i\}_{i \in I}$  is *elementary* if  $M_i \preceq M_j$  for  $i \leq j$ .

**Theorem 11.4.3** (Tarski-Vaught). *Let  $\{M_i\}_{i \in I}$  be an elementary chain of  $\mathcal{L}$ -structures. Let  $M = \bigcup_{i \in I} M_i$ . Then  $M_i \preceq M$  for all  $i \in I$ .*

*Proof sketch.* For each  $i$ , note that  $\text{eldiag}(M_i)$  is finitely satisfiable, complete, and has the witness property. Therefore the union  $\bigcup_i \text{eldiag}(M_i)$  is finitely satisfiable, complete, and has the witness property. The canonical model is  $M$ . Then  $M \succeq M_i$  because  $M \models \text{eldiag}(M_i)$ .  $\square$

**Lemma 11.4.4.** *If  $M$  is a structure, there is  $N \succeq M$  such that every type in  $S_1(M)$  is realized in  $N$ .*

*Proof.* Let  $\{p_i(x) : i \in I\}$  enumerate  $S_1(M)$ . Let  $\bar{x} = (x_i : i \in I)$  be a tuple of variables, one for each  $i \in I$ . Let  $\Sigma(\bar{x}) = \{p_i(x_i) : i \in I\}$ . Then  $\Sigma(\bar{x})$  is finitely satisfiable in  $M$  because each  $p_i$  is finitely satisfiable in  $M$ . Therefore,  $\Sigma(\bar{x})$  is realized by some tuple  $\bar{a} = (a_i : i \in I)$  in an elementary extension  $N \succeq M$ . The element  $a_i$  realizes  $p_i$ .  $\square$

**Theorem 11.4.5.** *If  $M$  is a structure and  $\kappa$  is a cardinal, there is a  $\kappa$ -saturated  $N \succeq M$ .*

*Proof.* Replacing  $\kappa$  with  $\kappa^+$  if necessary, we may assume  $\kappa$  is regular. Build an elementary chain  $\{M_\alpha\}_{\alpha < \kappa}$  by recursion on  $\alpha$ :

- $M_0 = M$ .
- $M_{\alpha+1}$  is an elementary extension of  $M_\alpha$  realizing every complete 1-type over  $M_\alpha$  (using Lemma 11.4.4).

- If  $\alpha$  is a limit ordinal, take  $M_\alpha = \bigcup_{\beta < \alpha} M_\beta$ , using the Tarski-Vaught theorem on chains (Theorem 11.4.3).

Let  $N = \bigcup_{\alpha < \kappa} M_\alpha$ . Then  $N \succeq M_0 = M$  by Tarski-Vaught again. If  $A \subseteq N$  and  $|A| < \kappa$ , then  $A \subseteq M_\alpha$  for some  $\alpha < \kappa$  by Lemma 11.4.1. If  $p \in S_1(A)$ , then  $p$  is a partial type over  $M_\alpha$ , so it extends to a complete type  $p' \in S_1(M_\alpha)$ , which is then realized by some  $b \in M_{\alpha+1} \subseteq N$ . Therefore  $N$  is  $\kappa$ -saturated.  $\square$

**Lemma 11.4.6.** *Let  $M$  be a structure and  $N$  be an  $|M|^+$ -saturated elementary extension. Let  $f$  be a partial elementary map from  $M$  to  $M$ .*

1. *There is a partial elementary map  $g$  from  $N$  to  $N$  extending  $f$ , with  $\text{dom}(g) = M$ .*
2. *There is a partial elementary map  $h$  from  $N$  to  $N$  extending  $f$ , with  $\text{im}(h) = M$ .*

*Proof.* 1. Let  $\kappa = |M|^+$ . Then  $N$  is  $\kappa$ -saturated and  $|M| < \kappa$ . Note that  $f$  is a partial elementary map from  $M$  to  $N$ . By Lemma 11.2.3, we can extend  $f$  to a partial elementary map  $g$  from  $M$  to  $N$  with  $\text{dom}(g) = M$ .

2. This follows from (1) by symmetry. More precisely, apply part (1) to  $f^{-1}$  to get a partial elementary map  $g$  extending  $f^{-1}$  with  $\text{dom}(g) = M$ . Then set  $h = g^{-1}$ , so that  $\text{im}(h) = \text{dom}(g) = M$ .  $\square$

**Lemma 11.4.7.** *For any  $M$  there is an elementary extension  $N \succeq M$  with the following properties:*

1. *Every complete type over  $M$  is realized in  $N$ .*
2. *If  $f$  is a partial elementary map from  $M$  to  $M$ , then there is  $\sigma \in \text{Aut}(N)$  extending  $f$ .*

*Proof.* Build an elementary chain  $\{M_i\}_{i < \omega}$  by recursion on  $i$ :

- $M_0 = M$ .
- $M_{i+1}$  is an  $|M_i|^+$ -saturated elementary extension of  $M_i$ . This is possible by Theorem 11.4.5.

Let  $N = \bigcup_{i=0}^{\infty} M_i$ . Then  $M = M_0 \preceq N$  by the Tarski-Vaught theorem on chains (Theorem 11.4.3). Every complete type over  $M$  is already realized in  $M_1$ , hence in  $N$ .

Let  $f : A \rightarrow B$  be a partial elementary map from  $M$  to  $M$ . Recursively build an increasing chain of partial elementary maps  $\{f_i\}_{i < \omega}$  with  $\text{dom}(f_i), \text{im}(f_i) \subseteq M_i$  as follows:

- $f_0 = f$ .
- If  $n > 0$ , then  $f_n : M_n \dashrightarrow M_n$  is a partial elementary map extending  $f_{n-1} : M_{n-1} \dashrightarrow M_{n-1}$  with

$$\begin{aligned} \text{dom}(f_n) &= M_{n-1} \text{ if } n \text{ is odd} \\ \text{im}(f_n) &= M_{n-1} \text{ if } n \text{ is even.} \end{aligned}$$

Take  $\sigma = \bigcup_{n=0}^{\infty} f_n$ . Then  $\sigma$  is a partial elementary map from  $N$  to  $N$ . The odd steps ensure  $\text{dom}(\sigma) \supseteq \bigcup_n M_n = N$ , and the even steps ensure  $\text{im}(\sigma) \supseteq \bigcup_n M_n = N$ . Thus  $\text{dom}(\sigma) = \text{im}(\sigma) = N$ , and  $\sigma \in \text{Aut}(N)$ .  $\square$

**Theorem 11.4.8.** *If  $M$  is a structure and  $\kappa$  is a cardinal, then there is a  $\kappa$ -saturated, strongly  $\kappa$ -homogeneous elementary extension  $N \succeq M$ .*

*Proof.* Replacing  $\kappa$  with  $\kappa^+$  if necessary, we may assume  $\kappa$  is regular. Build an elementary chain  $\{M_\alpha\}_{\alpha < \kappa}$  by recursion on  $\alpha$  as follows:

1.  $M_0 = M$ .
2.  $M_{\alpha+1}$  is an elementary extension of  $M_\alpha$  as in Lemma 11.4.7. In particular, every complete 1-type over  $M_\alpha$  is realized in  $M_{\alpha+1}$ .
3. If  $\alpha$  is a limit ordinal, then  $M_\alpha = \bigcup_{\beta < \alpha} M_\beta$ . This works by the Tarski-Vaught theorem on chains (Theorem 11.4.3).

Let  $N = \bigcup_{\alpha < \kappa} M_\alpha$ . As in the proof of Theorem 11.4.5,  $N$  is  $\kappa$ -saturated. We prove strong  $\kappa$ -homogeneity. Suppose  $f$  is a partial elementary map from  $N$  to  $N$  with  $|\text{dom}(f)| = |\text{im}(f)| < \kappa$ . By Lemma 11.4.1, there is  $\alpha < \kappa$  with  $\text{dom}(f) \cup \text{im}(f) \subseteq M_\alpha$ . Then  $f$  is a partial elementary map from  $M_\alpha$  to  $M_\alpha$ . By Lemma 11.4.7, we can extend  $f$  to an automorphism  $\sigma_{\alpha+1} \in \text{Aut}(M_{\alpha+1})$ . Build an increasing chain  $\{\sigma_\beta\}_{\alpha < \beta < \kappa}$  with  $\sigma_\beta \in \text{Aut}(M_\beta)$  by repeatedly applying Lemma 11.4.7. Take  $\sigma = \bigcup_{\alpha < \beta < \kappa} \sigma_\beta$ . Then  $\sigma \in \text{Aut}(N)$ , and  $\sigma \supseteq \sigma_{\alpha+1} \supseteq f$ . Therefore  $N$  is strongly  $\kappa$ -homogeneous.  $\square$

## 11.5 Saturation and back-and-forth

**Theorem 11.5.1.** *Let  $M$  and  $N$  be  $\omega$ -saturated. For any  $\bar{a} \in M^n$  and  $\bar{b} \in N^n$  with  $\text{tp}(\bar{a}) = \text{tp}(\bar{b})$ , let  $g_{\bar{a},\bar{b}}$  be the isomorphism from  $\langle \bar{a} \rangle_M$  to  $\langle \bar{b} \rangle_N$  sending  $\bar{a}$  to  $\bar{b}$  as in Theorem 8.3.4. Then the family  $\mathcal{F} = \{g_{\bar{a},\bar{b}} : n \in \mathbb{N}, \bar{a} \in M, \bar{b} \in N, \text{tp}(\bar{a}) = \text{tp}(\bar{b})\}$  is a back-and-forth system.*

*Proof.* We verify the “forward” condition; “backward” is similar. Fix some  $g_{\bar{a},\bar{b}}$ . Let  $a'$  be an element of  $M$ . By Lemma 11.2.2 we can extend the partial elementary map  $\bar{a} \mapsto \bar{b}$  to a partial elementary map  $(\bar{a}, a') \mapsto (\bar{b}, b')$ . Then  $\text{tp}(\bar{a}, a') = \text{tp}(\bar{b}, b')$ , and  $g_{\bar{a}a',\bar{b}b'}$  is the desired partial isomorphism in  $\mathcal{F}$  extending  $g_{\bar{a},\bar{b}}$  and containing  $a'$  in its domain.  $\square$

**Corollary 11.5.2.** *If  $M, N$  are countable and  $\omega$ -saturated, and  $M \equiv N$ , then  $M \cong N$ .*

*Proof.* Let  $\mathcal{F}$  be as in Theorem 11.5.1. By Theorem 5.6.4, it suffices to show that  $\mathcal{F}$  is non-empty. We only need to find some  $n \in \mathbb{N}$ , some  $\bar{a} \in M^n$ , and  $\bar{b} \in N^n$  with  $\text{tp}(\bar{a}) = \text{tp}(\bar{b})$ . Take  $n = 0$ ,  $\bar{a} = () \in M^0$ , and  $\bar{b} = () \in N^0$ . Then  $\text{tp}(\bar{a}) = \text{Th}(M) = \text{Th}(N) = \text{tp}(\bar{b})$ .  $\square$

**Corollary 11.5.3.** *If  $M$  is countable and  $\omega$ -saturated, then  $M$  is strongly  $\omega$ -homogeneous.*

*Proof.* Let  $f : A \rightarrow B$  be a partial elementary map from  $M$  to  $M$ , with  $A$  and  $B$  finite. Let  $\bar{a}$  be a tuple enumerating  $A$ . Then  $\bar{b} := f(\bar{a})$  enumerates  $B$ , and  $\text{tp}(\bar{a}) = \text{tp}(\bar{b})$ . We must find  $\sigma \in \text{Aut}(M)$  with  $\sigma(\bar{a}) = \bar{b}$ . If  $\mathcal{F}$  is the back-and-forth system between  $M$  and  $M$  given by Theorem 11.5.1, then  $g_{\bar{a},\bar{b}} \in \mathcal{F}$ . By Lemma 5.6.3, there is an isomorphism  $\sigma : M \rightarrow M$  extending  $g_{\bar{a},\bar{b}}$ . Then  $\sigma(\bar{a}) = \bar{b}$ .  $\square$

**Theorem 11.5.4.** *Let  $\kappa$  be an infinite cardinal. The following are equivalent:*

1.  *$T$  has quantifier elimination.*
2. *If  $M, N$  are  $\kappa$ -saturated structures, then the family  $\mathcal{F}_0$  of all finitely generated partial isomorphisms from  $M$  to  $N$  is a back-and-forth system.*

*Proof.* (1)  $\implies$  (2): Note that  $M, N$  are  $\omega$ -saturated. If  $\bar{a} \in M^n$  and  $\bar{b} \in N^n$  and  $\text{qftp}(\bar{a}) = \text{qftp}(\bar{b})$ , let  $g_{\bar{a}, \bar{b}} : \langle \bar{a} \rangle_M \rightarrow \langle \bar{b} \rangle_N$  be the partial isomorphism sending  $\bar{a}$  to  $\bar{b}$  from Theorem 8.3.4. Then

$$\mathcal{F}_0 = \{g_{\bar{a}, \bar{b}} : \text{qftp}(\bar{a}) = \text{qftp}(\bar{b})\}.$$

By quantifier elimination, this is

$$\mathcal{F}_0 = \{g_{\bar{a}, \bar{b}} : \text{tp}(\bar{a}) = \text{tp}(\bar{b})\},$$

which is a back-and-forth system by Theorem 11.5.1.

(2)  $\implies$  (1): By Theorem 8.3.3, it suffices to show that if  $M, N \models T$ , if  $\bar{a} \in M^n$  and  $\bar{b} \in N^n$ , then

$$\text{qftp}^M(\bar{a}) = \text{qftp}^N(\bar{b}) \implies \text{tp}^M(\bar{a}) = \text{tp}^N(\bar{b}).$$

Replacing  $M$  and  $N$  by elementary extensions, we may assume  $M$  and  $N$  are  $\kappa$ -saturated. By (2), the family  $\mathcal{F}_0$  of finitely generated partial isomorphisms from  $M$  to  $N$  is a back-and-forth system. Suppose  $\text{qftp}^M(\bar{a}) = \text{qftp}^N(\bar{b})$ . Then  $\mathcal{F}$  contains the isomorphism  $g_{\bar{a}, \bar{b}} : \langle \bar{a} \rangle_M \rightarrow \langle \bar{b} \rangle_N$ . By Theorem 3.7.6,  $g_{\bar{a}, \bar{b}}$  is a partial elementary map, so  $\text{tp}(\bar{a}) = \text{tp}(\bar{b})$ .  $\square$

## 11.6 Application: discrete linear orders with endpoints

**Definition 11.6.1.** Let  $x, y$  be elements in a linear order  $(M, \leq)$ . Then  $y$  is a *successor* of  $x$ , and  $x$  is a *predecessor* of  $y$ , written  $x \triangleleft y$ , if  $x < y$  but there is no  $z \in M$  with  $x < z < y$ .

Let  $T$  be the theory of linear orders  $(M, \leq)$  such that the following hold:

1.  $\min(M)$  and  $\max(M)$  exist. In particular,  $M$  is non-empty.
2. Every  $x \in M$  other than  $\max(M)$  has a successor  $y \triangleright x$ .
3. Every  $x \in M$  other than  $\min(M)$  has a predecessor  $y \triangleleft x$ .

**Example 11.6.2.** Any finite non-empty linear order is a model of  $T$ .

**Definition 11.6.3.** Suppose  $M \models T$  and  $x, y \in M$ . The *distance* between  $x$  and  $y$ , written  $d(x, y)$ , is the element of  $\mathbb{N} \cup \{\infty\}$  defined as follows:

1. If  $x = y$ , then  $d(x, y) = 0$ .
2. If  $x < y$ , then  $d(x, y)$  is  $1 + |\{z \in M : x < z < y\}|$ .
3. If  $x > y$ , then  $d(y, x) = d(x, y)$ .

Note that we define  $d(x, y)$  to be the symbol “ $\infty$ ” when  $[x, y]$  is infinite, rather than distinguishing cardinalities.

Let  $\mathcal{L}'$  be the base language of orders  $\mathcal{L} = \{\leq\}$  plus two new constant symbols  $\min, \max$  and a binary relation  $R_n$  for each  $n < \infty$ . Expand any model  $M \models T$  to an  $\mathcal{L}'$ -structure by interpreting the new symbols as follows:

1.  $\min^M = \min(M)$ .
2.  $\max^M = \max(M)$ .
3.  $R_n^M(x, y) \iff d(x, y) = n$ .

The resulting  $\mathcal{L}'$ -structures are the models of some  $\mathcal{L}'$ -theory  $T'$ .

**Remark 11.6.4.** Let  $M, N$  be models of  $T'$ . Suppose

$$\begin{aligned} \min(M) &= a_1 < a_2 < \cdots < a_n = \max(M) \\ \min(N) &= b_1 < b_2 < \cdots < b_n = \max(N) \\ d(a_i, a_{i+1}) &= d(b_i, b_{i+1}) \text{ for } 1 \leq i < n. \end{aligned}$$

Then there is a partial isomorphism

$$\begin{aligned} f : \{a_1, \dots, a_n\} &\rightarrow \{b_1, \dots, b_n\} \\ f(a_i) &= b_i. \end{aligned}$$

Moreover, all finite partial isomorphisms from  $M$  to  $N$  arise in this way.

**Lemma 11.6.5.** *Let  $M, N$  be  $\aleph_1$ -saturated models of  $T'$ . Let  $\mathcal{F}$  be the collection of finite partial isomorphisms  $f : A \rightarrow B$ . Then  $\mathcal{F}$  is a back-and-forth system.*

*Proof.* By symmetry we only need to prove the forward condition. Let  $f : A \rightarrow B$  be a finite partial isomorphism and  $\alpha$  be an element of  $M$ . We must



find  $f' \in \mathcal{F}$  extending  $f$  with  $\alpha \in \text{dom}(f')$ . We may assume  $\alpha \notin \text{dom}(f)$ ; otherwise take  $f' = f$ . By Remark 11.6.4,  $f$  has the form

$$\begin{aligned} f : \{a_1, \dots, a_n\} &= \{b_1, \dots, b_n\} \\ f(a_i) &= b_i \end{aligned}$$

where

$$\begin{aligned} \min(M) &= a_1 < a_2 < \dots < a_n = \max(M) \\ \min(N) &= b_1 < b_2 < \dots < b_n = \max(N) \\ d(a_i, a_{i+1}) &= d(b_i, b_{i+1}) \text{ for } 1 \leq i < n. \end{aligned}$$

There must be some  $i$  such that  $a_i < \alpha < a_{i+1}$ . (The cases  $\alpha < a_1 = \min(M)$  and  $\alpha > a_n = \max(M)$  are impossible.) By Remark 11.6.4, we must find  $\beta$  such that

$$\begin{aligned} b_i &< \beta < b_{i+1} \\ d(b_i, \beta) &= d(a_i, \alpha) =: x \\ d(\beta, b_{i+1}) &= d(\alpha, a_{i+1}) =: y. \end{aligned}$$

Note that  $d(b_i, b_{i+1}) = d(a_i, a_{i+1}) = x + y$ . There are four cases:

1.  $x, y < \infty$ . Take  $\beta$  between  $b_i$  and  $b_{i+1}$  with  $d(b_i, \beta) = x$ . Then  $d(\beta, b_{i+1}) = (x + y) - x = y$  as desired.
2.  $x < \infty = y$ . Then  $d(b_i, b_{i+1}) = \infty$ . Take  $\beta$  between  $b_i$  and  $b_{i+1}$  with  $d(b_i, \beta) = x$ . Then  $d(\beta, b_{i+1}) = \infty$ .
3.  $y < \infty = x$ . Similar.
4.  $x = y = \infty$ . As  $d(b_i, b_{i+1}) = \infty$ , there are

$$b_i \triangleleft c_1 \triangleleft c_2 \triangleleft \dots \leq \dots \triangleleft d_2 \triangleleft d_1 \triangleleft b_{i+1}.$$

The partial type  $\{x > c_i : i < \omega\} \cup \{x < d_i : i < \omega\}$  is realized in  $M$ , by  $\aleph_1$ -saturation. Let  $\beta$  be a realization. Then  $b_i < \beta < b_{i+1}$ , and  $d(b_i, \beta) = d(\beta, b_{i+1}) = \infty$ .  $\square$

**Theorem 11.6.6.** *The theory  $T'$  has quantifier elimination.*

**Corollary 11.6.7.** *Let  $M$  be an infinite model of  $T$ . Let  $M_0$  be the set of  $x \in M$  such that  $d(x, \min(M)) < \infty$  or  $d(x, \max(M)) < \infty$ . Then  $M_0 \preceq M$ .*

*Proof.* Working in the expanded language  $\mathcal{L}'$ , it is easy to see that  $M_0$  is a substructure of  $M$  and  $M_0 \models T'$ . Then  $M_0 \preceq M$  because submodels are elementary substructures in theories with quantifier elimination (Theorem 8.3.6(2)).  $\square$

**Definition 11.6.8.** If  $M$  is a model of  $T$ , the *length* of  $M$ , written  $\ell(M)$ , is  $d(\min(M), \max(M)) \in \mathbb{N} \cup \{\infty\}$ .

**Corollary 11.6.9.** *Two models  $M, N \models T$  are elementarily equivalent if and only if  $\ell(M) = \ell(N)$ .*

*Proof.* If  $M \equiv N$ , then certainly  $\ell(M) = \ell(N)$ , essentially because the relations  $R_n$  are definable, or simply because  $\ell(M)$  is  $|M| - 1$ .

Conversely, suppose  $\ell(M) = \ell(N)$ . Let  $M'$  and  $N'$  be the expansions of  $M$  and  $N$  to models of  $T$ . Note that the minimal substructure  $\langle \emptyset \rangle_{M'}$  contains only the two points  $\min(M)$  and  $\max(M)$ . This substructure is determined up to isomorphism by  $\ell(M)$ . For example,  $R_n(\min(M), \max(M))$  holds iff  $\ell(M) = n$ . Because  $\ell(M) = \ell(N)$ , we have  $\langle \emptyset \rangle_{M'} \cong \langle \emptyset \rangle_{N'}$ , and so  $M' \equiv N'$  by quantifier elimination (because of Theorem 8.3.5). Restricting to  $\mathcal{L}$ -sentences, we see  $M \equiv N$ .  $\square$

**Corollary 11.6.10.** *The theory of infinite models of  $T$  is complete and decidable.*

**Corollary 11.6.11.** *The class  $\mathcal{K}$  of models of  $T$  is the elementary class generated by finite non-empty linear orders.*

*Proof.* Certainly  $\mathcal{K}$  is an elementary class containing the finite non-empty linear orders. If  $\mathcal{K}'$  is a smaller such elementary class, take  $M \in \mathcal{K} \setminus \mathcal{K}'$ . Then  $M$  is infinite or else  $M \in \mathcal{K}'$ . Because  $\mathcal{K}'$  contains models of size  $> n$  for each  $n$ , it must contain an infinite model  $N$  (Theorem 5.4.4). Then  $M, N$  are infinite models of  $T$ , so Corollary 11.6.10 shows  $M \equiv N \in \mathcal{K}'$ . Then  $M \in \mathcal{K}'$ , contradicting the choice of  $M$ .  $\square$

**Corollary 11.6.12.** *If  $\varphi$  is a sentence in the language of orders, then the following are equivalent:*

- $T \vdash \varphi$ .
- Every finite non-empty linear order satisfies  $\varphi$ .

# Chapter 12

## Countable categoricity

### 12.1 Baire category theorem

Let  $S$  be a non-empty Stone space (Definition 7.4.2).

**Definition 12.1.1.**  $X \subseteq S$  is *dense* if  $X$  intersects any non-empty clopen set  $Y \subseteq S$ .

**Definition 12.1.2.** A set  $X \subseteq S$  is *comeager* if  $X$  contains a countable intersection of dense open sets.

**Remark 12.1.3.** Dense open sets are comeager. A countable intersection of comeager sets is comeager.

**Theorem 12.1.4** (Baire category theorem). *If  $X \subseteq S$  is comeager, then  $X$  is dense, hence non-empty.*

*Proof.* Because  $S$  is a Stone space, every open set is a union of clopen sets. Therefore every non-empty open set contains a non-empty clopen set.

Suppose  $X \supseteq \bigcup_{i=1}^{\infty} U_i$  where each  $U_i$  is open and dense. Let  $V_0$  be a non-empty clopen set. Then  $V_0 \cap U_1$  is a non-empty open set. It contains a non-empty clopen set  $V_1$ . Similarly,  $V_1 \cap U_2$  contains a non-empty clopen set  $V_2$ . Continuing, we can build a descending chain of clopen sets

$$V_0 \supseteq V_1 \supseteq V_2 \supseteq \cdots$$

with  $V_i \subseteq U_i$ . The family  $\{V_0, V_1, V_2, \dots\}$  has the FIP, so the intersection is non-empty by compactness (Theorem 7.1.15). Take  $p \in \bigcap_{i=0}^{\infty} V_i \subseteq V_0 \cap \bigcap_{i=1}^{\infty} U_i \subseteq V_0 \cap X$ . Then  $X$  intersects  $V_0$  as desired.  $\square$

## 12.2 The omitting types theorem

Fix an  $\mathcal{L}$ -structure  $M$ , subset  $A \subseteq M$ , and  $n < \omega$ .

**Definition 12.2.1.** A complete type  $p \in S_n(A)$  is *isolated* if  $\{p\}$  is clopen.

**Theorem 12.2.2.** *If  $p$  is isolated, then  $p$  is realized in  $M$ .*

*Proof.* As  $\{p\}$  is clopen,  $\{p\} = \llbracket \varphi \rrbracket \subseteq S_n(A)$  for some  $\mathcal{L}(A)$ -formula  $\varphi(\bar{x}) \in p(\bar{x})$ . Then  $\varphi(\bar{x})$  is satisfied by  $b \in M$ , as  $p$  is finitely satisfiable. If  $p' = \text{tp}(b/A) \in S_n(A)$ , then  $p' \in \llbracket \varphi \rrbracket = \{p\}$ , so  $p' = p$  and  $b$  realizes  $p$ .  $\square$

Fix a complete theory  $T$  in a countable language  $\mathcal{L}$ .

**Definition 12.2.3.** If  $p \in S_n(T)$  and  $M \models T$ , then  $M$  *omits*  $p$  if  $p$  isn't realized in  $M$ .

Let  $S_\omega(T)$  be the space of complete  $\omega$ -types  $\text{tp}(a_0, a_1, a_2, \dots)$ , i.e., complete types in the variables  $\bar{x} = (x_0, x_1, x_2, \dots)$ . Work in a monster model  $\mathbb{M} \models T$ .

**Lemma 12.2.4.** *There is a comeager set  $W \subseteq S_\omega(T)$  such that if  $\bar{c} \in \mathbb{M}^\omega$  and  $\text{tp}(\bar{c}) \in W$ , then  $\{c_i : i \in \omega\} \preceq \mathbb{M}$ .*

*Proof.*

*Claim.* For any formula  $\varphi(\bar{x}, y)$ , the following open set is dense:

$$U_\varphi := \llbracket \neg \exists y \varphi(\bar{x}, y) \rrbracket \cup \bigcup_{i=0}^{\infty} \llbracket \varphi(\bar{x}, x_i) \rrbracket.$$

*Proof.* Take non-empty  $\llbracket \psi \rrbracket \subseteq S_\omega(T)$ . Suppose for the sake of contradiction that  $\llbracket \psi \rrbracket \cap U_\varphi = \emptyset$ . Then  $\llbracket \psi \rrbracket$  doesn't intersect the sets in the union, which means the following:

$$\begin{aligned} \psi(\bar{x}) \wedge (\neg \exists y \varphi(\bar{x}, y)) &\text{ is inconsistent} \\ \psi(\bar{x}) \wedge \varphi(\bar{x}, x_i) &\text{ is inconsistent, for each } i < \omega. \end{aligned}$$

Take  $\bar{c} \in \mathbb{M}^\omega$  realizing  $\psi(\bar{x})$ . By the first line,  $\mathbb{M} \models \exists y \varphi(\bar{c}, y)$ . Thus  $\mathbb{M} \models \varphi(\bar{c}, e)$  for some  $e \in \mathbb{M}$ . Take  $i \gg 0$  so that  $x_i$  doesn't occur in  $\psi(\bar{x})$  or  $\varphi(\bar{x}, y)$ . Changing  $c_i$  to  $e$ , we may assume  $\mathbb{M} \models \varphi(\bar{c}, c_i)$ , contradicting the second line.  $\square$  Claim

Now let  $W$  be the comeager set  $\bigcap_{\varphi} U_{\varphi}$ . Suppose  $\bar{c} \in \mathbb{M}^{\omega}$  and  $\text{tp}(\bar{c}) \in W$ . We claim that  $M := \{c_i : i \in \omega\} \preceq \mathbb{M}$  by the Tarski-Vaught criterion (Theorem 5.3.1). Suppose  $D \subseteq \mathbb{M}^1$  is  $M$ -definable, defined as  $\varphi(\bar{c}, \mathbb{M})$  for some  $\varphi(\bar{x}, y)$ . If  $D \neq \emptyset$ , then  $\mathbb{M} \models \exists y \varphi(\bar{c}, y)$ , so  $\text{tp}(\bar{c}) \notin \llbracket \neg \exists y \varphi(\bar{x}, y) \rrbracket$ . But  $\text{tp}(\bar{c}) \in U_{\varphi}$ , so  $\text{tp}(\bar{c}) \in \llbracket \varphi(\bar{x}, x_i) \rrbracket$  for some  $i$ , which means  $\mathbb{M} \models \varphi(\bar{c}, c_i)$ , or equivalently,  $c_i \in D$ . Thus  $M = \{c_i : i \in \omega\}$  intersects every non-empty  $M$ -definable set  $D \subseteq \mathbb{M}^1$ .  $\square$

**Lemma 12.2.5.** *For any  $j_1, \dots, j_n < \omega$ , let  $f_{\bar{j}} : S_{\omega}(T) \rightarrow S_n(T)$  be the restriction map  $\text{tp}(\bar{c}) \mapsto \text{tp}(c_{j_1}, \dots, c_{j_n})$ .*

1.  $f$  is continuous.
2. If  $X \subseteq S_{\omega}(T)$  is clopen, then  $f(X) \subseteq S_n(T)$  is clopen.

*Proof.* 1. The preimage of  $\llbracket \varphi(y_1, \dots, y_n) \rrbracket \subseteq S_n(T)$  is  $\llbracket \varphi(x_{j_1}, \dots, x_{j_n}) \rrbracket \subseteq S_{\omega}(T)$ .

2. If  $X = \llbracket \varphi \rrbracket$ , then

$$\begin{aligned} X &= \{\text{tp}(\bar{c}) : \bar{c} \in \mathbb{M}^{\omega}, \mathbb{M} \models \varphi(\bar{c})\} \\ f(X) &= \{\text{tp}(c_{j_1}, \dots, c_{j_n}) : \bar{c} \in \mathbb{M}^{\omega}, \mathbb{M} \models \varphi(\bar{c})\}. \end{aligned}$$

(The first line holds by saturation of  $\mathbb{M}$ .) Now the set

$$\{(c_{j_1}, \dots, c_{j_n}) : \bar{c} \in \mathbb{M}^{\omega}, \mathbb{M} \models \varphi(\bar{c})\}$$

is definable, defined by the formula

$$\psi(y_1, \dots, y_n) := \exists x_0, \dots, x_N \left( \varphi(\bar{x}) \wedge \bigwedge_{i=1}^n y_i = x_{j_i} \right),$$

where  $N$  is large enough to quantify away all  $x_i$ 's appearing in  $\varphi(\bar{x})$ . Then

$$f(X) = \{\text{tp}(\bar{a}) : \bar{a} \in \psi(\mathbb{M}^n)\} = \llbracket \psi \rrbracket. \quad \square$$

**Lemma 12.2.6.** *Let  $p \in S_n(T)$  be non-isolated. There is a comeager set  $V_p \subseteq S_{\omega}(T)$  such that if  $\bar{c} \in \mathbb{M}^{\omega}$  and  $\text{tp}(\bar{c}) \in V_p$ , then  $p$  is not realized by any tuple in  $C = \{c_i : i < \omega\}$ .*

*Proof.*

*Claim.* For any  $j_1, \dots, j_n \in \omega$ , there is a dense open set  $U_{\bar{j}} \subseteq S_\omega(T)$  such that if  $\text{tp}(\bar{c}) \in U_{\bar{j}}$ , then  $(c_{j_1}, \dots, c_{j_n})$  doesn't realize  $p$ .

*Proof.* Let  $f = f_{\bar{j}} : S_\omega(T) \rightarrow S_n(T)$  be the restriction map from Lemma 12.2.5. Points are closed (Theorem 7.1.11) so  $S_n(T) \setminus \{p\}$  is open and the preimage  $U := f^{-1}(S_n(T) \setminus \{p\})$  is open. For density, suppose  $X \subseteq S_\omega(T)$  is clopen and non-empty, but  $X \cap U = \emptyset$ . Then  $f(X) \subseteq \{p\}$ . But  $f(X)$  is clopen by Lemma 12.2.5(2), and non-empty, so  $f(X) = \{p\}$  and  $p$  is isolated, a contradiction. Thus  $U$  is dense. Finally, if  $\text{tp}(\bar{c}) \in U$ , then  $\text{tp}(c_{j_1}, \dots, c_{j_n}) \in S_n(T) \setminus \{p\}$ , meaning that  $(c_{j_1}, \dots, c_{j_n})$  doesn't realize  $p$ . □<sub>Claim</sub>

Take  $V_p = \bigcap_{\bar{j} \in \mathbb{N}^\omega} U_{\bar{j}}$ . If  $\text{tp}(\bar{c}) \in V_p$ , then  $(c_{j_1}, \dots, c_{j_n})$  doesn't realize  $p$  for any  $\bar{j} \in \mathbb{N}^\omega$ . □

**Theorem 12.2.7** (Omitting types theorem). *Let  $p_1, p_2, \dots$  be a countable list of non-isolated complete types  $p_i \in S_{n_i}(T)$ . Then there is a countable model  $M \models T$  omitting every  $p_i$ .*

*Proof.* Let  $W$  and  $V_p$  be the comeager sets from Lemmas 12.2.4 and 12.2.6. The set  $Q = W \cap \bigcap_{i=1}^\infty V_{p_i}$  is comeager, hence non-empty by the Baire Category Theorem. Take  $\bar{c} \in \mathbb{M}^\omega$  with  $\text{tp}(\bar{c}) \in Q$ , and let  $M = \{c_i : i < \omega\}$ . Then  $M \preceq \mathbb{M}$  because  $\text{tp}(\bar{c}) \in W$ , and  $M$  omits  $p_i$  because  $\text{tp}(\bar{c}) \in V_{p_i}$ . □

## 12.3 Countably categorical theories

Work in a monster model  $\mathbb{M} \models T$ .

**Lemma 12.3.1.**  *$S_n(A)$  is finite iff all types in  $S_n(A)$  are isolated.*

*Proof.* Suppose  $S_n(A)$  is finite. Every point is closed (Theorem 7.1.11), and a finite union of closed sets is closed, so every subset of  $S_n(A)$  is closed, every subset is open, and every subset is clopen, implying every point is isolated.

Conversely, suppose every point is isolated. Then  $S_n(A) = \bigcup_{p \in S_n(A)} \{p\}$  is an open cover. By compactness of  $S_n(A)$ , there is a finite subcover, and so  $S_n(A)$  is finite. □

**Lemma 12.3.2.**  *$S_n(A)$  is finite iff there are only finitely many  $A$ -definable sets  $D \subseteq M^n$ .*

*Proof.* The boolean algebra of  $A$ -definable sets is isomorphic to the boolean algebra of clopen sets in  $S_n(A)$ . If  $S_n(A)$  is finite, then there are only finitely many clopen sets. Conversely, if there the number of clopen sets is  $k < \infty$ , then there are at most  $2^k$  open sets, because every open set is a union of clopen sets. Then there are at most  $2^k$  closed sets, hence  $2^k$  points (as every point is closed by Theorem 7.1.11).  $\square$

**Lemma 12.3.3.** *If  $\bar{b}, \bar{c} \in \mathbb{M}^n$  and  $A = \{a_1, \dots, a_m\} \subseteq \mathbb{M}$ , then*

$$\bar{b} \equiv_A \bar{c} \iff \bar{a}\bar{b} \equiv_{\emptyset} \bar{a}\bar{c}.$$

*Proof.* Every  $\mathcal{L}(A)$ -formula has the form  $\varphi(\bar{a}, \bar{x})$  for some  $\mathcal{L}$ -formula  $\varphi(\bar{w}, \bar{x})$ . Therefore both sides say that for any  $\mathcal{L}$ -formula  $\varphi(\bar{w}, \bar{x})$ ,

$$\mathbb{M} \models \varphi(\bar{a}, \bar{b}) \iff \mathbb{M} \models \varphi(\bar{a}, \bar{c}). \quad \square$$

**Lemma 12.3.4.** *If  $|S_n(\emptyset)| < \infty$  for all  $n$ , then  $|S_n(A)| < \infty$  for all  $n$  and all finite  $A \subseteq \mathbb{M}$ .*

*Proof.* Fix some  $n < \omega$  and  $A = \{a_1, \dots, a_m\} \subseteq M$ . Then the map

$$\begin{aligned} S_n(A) &\rightarrow S_{m+n}(\emptyset) \\ \text{tp}(\bar{b}/A) &\mapsto \text{tp}(\bar{a}\bar{b}/\emptyset) \end{aligned}$$

is well-defined and injective by Lemma 12.3.3. Thus  $|S_n(A)| \leq |S_{m+n}(\emptyset)| < \infty$ .  $\square$

Recall that  $S_n(\emptyset) = S_n(T)$  (Corollary 8.1.20).

**Theorem 12.3.5** (Engeler, Ryll-Nardzewski, Svenonius). *Let  $T$  be a complete theory in a countable language and let  $\mathbb{M}$  be a monster model. The following are equivalent:*

1.  $T$  has a unique countable model.
2.  $S_n(T)$  is finite for all  $n < \omega$ .
3.  $S_n(A)$  is finite for all  $n < \omega$  and finite  $A \subseteq \mathbb{M}$ .
4. Every countable model of  $T$  is  $\omega$ -saturated.

*Proof.* (1)  $\implies$  (2): Suppose  $S_n(T)$  is infinite for some  $n$ . By Lemma 12.3.1, some  $p \in S_n(T)$  is non-isolated. Then  $p$  is realized in some model  $M$ . By the downward Löwenheim-Skolem theorem (Theorem 5.4.3), we may assume  $M$  is countable. By the omitting types theorem (Theorem 12.2.7), there is a countable model  $N$  omitting  $p$ . Then  $M \not\cong N$ , and (1) fails.

(2)  $\implies$  (3): Lemma 12.3.4.

(3)  $\implies$  (4): Assume (3). Let  $M$  be a countable model. If  $A$  is a finite subset of  $M$ , then  $S_n(A)$  is finite by (3), so every  $p \in S_n(A)$  is isolated by Lemma 12.3.1, and then every  $p \in S_n(A)$  is realized by Theorem 12.2.2. This shows that  $M$  is  $\omega$ -saturated.

(4)  $\implies$  (1): There is at most one countable  $\omega$ -saturated model by Corollary 11.5.2.  $\square$

**Theorem 12.3.6.** *Let  $M$  be a countable structure in a countable language. The following are equivalent:*

1.  $\text{Th}(M)$  is countably categorical.
2. For every  $n$ , the action of  $\text{Aut}(M)$  on  $M^n$  has finitely many orbits.

*Proof.* (1)  $\implies$  (2): by Theorem 12.3.5(4),  $M$  is  $\omega$ -saturated, hence strongly  $\omega$ -homogeneous by Corollary 11.5.3. Therefore  $\bar{a}, \bar{b} \in M^n$  are in the same orbit of  $\text{Aut}(M)$  iff  $\text{tp}(\bar{a}) = \text{tp}(\bar{b})$ . By Theorem 12.3.5(2), there are only finitely many complete  $n$ -types.

(2)  $\implies$  (1): if  $D \subseteq M^n$  is 0-definable, then  $D$  is  $\text{Aut}(M)$ -invariant so  $D$  is a union of orbits. There are only finitely many orbits, hence finitely many possibilities for  $D$ . Then the boolean algebra of 0-definable sets  $D \subseteq M^n$  is finite. By Lemma 12.3.2,  $S_n(\emptyset) = S_n(T)$  is finite. By Theorem 12.3.5,  $\text{Th}(M)$  is countably categorical.  $\square$



# Chapter 13

## Closure operators and pregeometries

### 13.1 Closure operators

**Definition 13.1.1.** A *closure operator* on a set  $S$  is an operation  $\text{cl}(-) : \mathfrak{P}(S) \rightarrow \mathfrak{P}(S)$  with the following properties:

- Idempotent:  $\text{cl}(\text{cl}(X)) = \text{cl}(X)$ .
- Monotone:  $X \subseteq Y \implies \text{cl}(X) \subseteq \text{cl}(Y)$ .
- Increasing:  $X \subseteq \text{cl}(X)$ .

The set  $\text{cl}(X)$  is called the *closure* of  $X$ .

Fix a set  $S$  and a closure operator  $\text{cl}(-)$  on  $S$ . We say that  $X \subseteq S$  is *closed* if  $X = \text{cl}(X)$ . By idempotence, the closed sets are exactly the sets of the form  $\text{cl}(X)$ .

**Theorem 13.1.2.** If  $X \subseteq S$ , then  $\text{cl}(X)$  is the smallest closed set containing  $X$ .

*Proof.* The set  $\text{cl}(X)$  contains  $X$  because  $\text{cl}(-)$  is increasing, and  $\text{cl}(X)$  is closed because  $\text{cl}(-)$  is idempotent. Suppose  $Y$  is a closed set containing  $X$ . Then  $Y = \text{cl}(Y) \supseteq \text{cl}(X)$ , because  $\text{cl}(-)$  is monotone.  $\square$

**Definition 13.1.3.** A *closure system* on a set  $S$  is a family  $\mathcal{C} \subseteq \mathfrak{P}(S)$  of sets, called *closed sets*, such that for any  $X \subseteq S$  there is a smallest closed set containing  $X$ .

**Remark 13.1.4.** If  $\mathcal{C}$  is a closure system, define  $\text{cl}(X)$  to be the smallest closed set containing  $X$ . It is easy to see that  $\text{cl}(-)$  is a closure operator on  $S$ . This gives a map from closure systems to closure operators. Conversely, any closure operator  $\text{cl}(-)$  defines a closure system  $\{X \subseteq S : \text{cl}(X) = X\}$  by Theorem 13.1.2. It is easy to see that these two maps are inverses. Thus closure operators correspond bijectively with closure systems.

**Theorem 13.1.5.** *A family  $\mathcal{C} \subseteq \mathfrak{P}(S)$  is a closure system iff  $\mathcal{C}$  is closed under infinite intersections.*

*Proof.* If  $\mathcal{C}$  is closed under infinite intersections, then the smallest closed set containing  $X$  exists—it is  $\bigcap \{Y \in \mathcal{C} : Y \supseteq X\}$ . Therefore  $\mathcal{C}$  is a closure system.

Conversely, suppose  $\mathcal{C}$  is a closure system. Let  $\{X_i\}_{i \in I}$  be a family of closed sets and let  $Y = \bigcap_{i \in I} X_i$ . We claim that  $Y$  is closed. For each  $i$ , we have  $\text{cl}(Y) \subseteq \text{cl}(X_i) = X_i$  by monotonicity. Thus  $\text{cl}(Y) \subseteq \bigcap_{i \in I} X_i = Y$ . Conversely  $Y \subseteq \text{cl}(Y)$  because  $\text{cl}(-)$  is increasing.  $\square$

**Example 13.1.6.** If  $S$  is a topological space, the family of closed sets is a closure system. The closure  $\text{cl}(X)$  is the smallest closed set containing  $X$ .

**Example 13.1.7.** Let  $M$  be a structure. Recall that  $\langle A \rangle_M$  is the smallest substructure of  $M$  containing  $A$ . Therefore the class of substructures of  $M$  is a closure system, and  $\langle - \rangle_M$  is a closure operation.

**Definition 13.1.8.** A closure operator  $\text{cl}(-)$  on a set  $S$  is *finitary* if for any  $a \in S$ ,  $X \subseteq S$  with  $a \in \text{cl}(X)$ , there is a finite subset  $X_0 \subseteq_f X$  with  $a \in \text{cl}(X_0)$ .

More generally, we say that an operation  $F : \mathfrak{P}(S) \rightarrow \mathfrak{P}(S)$  is *finitary* if  $F(X) \subseteq \bigcup \{F(X_0) : X_0 \subseteq_f X\}$ .

**Example 13.1.9.** If  $M$  is a structure, the closure operation  $\langle - \rangle_M$  is finitary: if  $b \in \langle A \rangle$  then  $b = t(\bar{a})$  for some term  $t$  and finite tuple  $\bar{a}$ . If  $A_0 = \{a_1, \dots, a_n\}$ , then  $b \in \langle A_0 \rangle_M$ .

**Lemma 13.1.10.** *Suppose  $\text{cl}(-) : \mathfrak{P}(S) \rightarrow \mathfrak{P}(S)$  is monotone, finitary, increasing, and satisfies the property*

$$a \in \text{cl}(X) \text{ and } b \in \text{cl}(X \cup \{a\}) \implies b \in \text{cl}(X). \quad (*)$$

*Then  $\text{cl}(-)$  is idempotent, and therefore a finitary closure operator.*

*Proof.* Note that  $(*)$  implies

$$\text{cl}(X \cup \{a\}) = \text{cl}(X) \text{ for } a \in \text{cl}(X). \quad (\dagger)$$

Fix  $X$ . Then  $\text{cl}(\text{cl}(X)) \supseteq \text{cl}(X)$  because  $\text{cl}(-)$  is increasing. For the reverse inclusion, suppose  $b \in \text{cl}(\text{cl}(X))$ . As  $\text{cl}(-)$  is finitary,  $b \in \text{cl}(\{a_1, \dots, a_n\})$  for some  $a_1, \dots, a_n \in \text{cl}(X)$ . By  $n$  applications of  $(\dagger)$ ,

$$\text{cl}(X) = \text{cl}(X \cup \{a_1\}) = \text{cl}(X \cup \{a_1, a_2\}) = \dots = \text{cl}(X \cup \{a_1, \dots, a_n\}).$$

Then

$$b \in \text{cl}(\{a_1, \dots, a_n\}) \subseteq \text{cl}(X \cup \{a_1, \dots, a_n\}) = \text{cl}(X). \quad \square$$

## 13.2 Algebraic and definable closure

**Definition 13.2.1.** Let  $M$  be a structure and  $A \subseteq M$  be a subset.

1.  $b \in M$  is *definable* over  $A$  if  $\{b\}$  is  $A$ -definable.
2.  $b \in M$  is *algebraic* over  $A$  if  $b \in D$  for some finite  $A$ -definable  $D \subseteq M$ .
3. The *definable closure* of  $A$ , written  $\text{dcl}(A)$  or  $\text{dcl}^M(A)$ , is the set of  $b \in M$  definable over  $A$ .
4. The *algebraic closure* of  $A$ , written  $\text{acl}(A)$  or  $\text{acl}^M(A)$ , is the set of  $b \in M$  algebraic over  $A$ .

**Lemma 13.2.2.** *If  $b \in \text{acl}(A)$ , then there is a finite  $A$ -definable set  $X \ni b$  such that  $\text{tp}(c/A) = \text{tp}(b/A)$  for every  $c \in X$ .*

*Proof.* Take a minimal finite  $A$ -definable set  $X \ni b$ . If  $X$  does not have the desired property, take  $c \in x$  with  $\text{tp}(c/A) \neq \text{tp}(b/A)$ . Then there is an  $A$ -definable set  $D$  with  $b \in D$  and  $c \notin D$  (Remark 8.1.3). Then  $X \cap D$  is a strictly smaller  $A$ -definable set containing  $b$ , a contradiction.  $\square$

**Lemma 13.2.3.** *Fix a structure  $M$ .*

1. *If  $A \subseteq B$  then  $\text{acl}(A) \subseteq \text{acl}(B)$ .*
2. *If  $b \in \text{acl}(A)$ , then  $b \in \text{acl}(A_0)$  for some finite subset  $A_0 \subseteq A$ .*
3.  *$A \subseteq \text{acl}(A)$ .*

4. If  $b \in \text{acl}(A)$  and  $c \in \text{acl}(A \cup \{b\})$ , then  $c \in \text{acl}(A)$ .

*Proof.* 1. Any finite  $A$ -definable set is a finite  $B$ -definable set.

2. Any finite  $A$ -definable set is  $A_0$ -definable for some finite  $A_0 \subseteq A$ .

3. If  $b \in A$ , then the set  $\{b\}$  is finite and  $A$ -definable, so  $b \in \text{acl}(A)$ .

4. As  $b \in \text{acl}(A)$  and  $c \in \text{acl}(A \cup \{b\})$ , there is a finite  $A$ -definable set  $X \ni b$  and a finite  $Ab$ -definable set  $Y \ni c$ . By Lemma 13.2.2, we may assume all elements of  $X$  have the same type over  $A$ . Write  $Y$  as  $\varphi(M, b)$  for some  $\mathcal{L}(A)$ -formula  $\varphi(x, y)$ . Let  $n = |Y| = |\varphi(M, b)|$ . Then  $n = |\varphi(M, b')|$  for every  $b' \in X$ , since this is expressed by a formula in  $\text{tp}(b'/A)$ . Let  $Z = \bigcup_{b' \in X} \varphi(M, b')$ . Then  $Z$  is a finite  $A$ -definable set containing  $c$ .  $\square$

**Theorem 13.2.4.** *Algebraic closure  $\text{acl}(-)$  is a finitary closure operator.*

*Proof.* Lemma 13.2.3 plus Lemma 13.1.10.  $\square$

A set  $A \subseteq M$  is *algebraically closed* if  $\text{acl}(A) = A$ . The algebraic closure of  $A$  is the smallest algebraically closed set containing  $A$ .

**Theorem 13.2.5.** *Definable closure  $\text{dcl}(-)$  is a finitary closure operator.*

*Proof.* Like Theorem 13.2.4 but slightly easier.  $\square$

A set  $A \subseteq M$  is *definably closed* if  $\text{dcl}(A) = A$ . The definable closure of  $A$  is the smallest definably closed set containing  $A$ .

### 13.3 Definable closure, algebraic closure, and substructures

**Theorem 13.3.1.** *Let  $M$  be a structure. If  $A \subseteq M$  and  $A = \text{dcl}(A)$ , then  $A$  is a substructure of  $M$ .*

*Proof.* Let  $f$  be a  $k$ -ary function symbol, and let  $a_1, \dots, a_k$  be elements of  $A$ . Then  $f(a_1, \dots, a_k)$  is in  $\text{dcl}(A)$ , as it is defined by the  $\mathcal{L}(A)$ -formula  $x = f(a_1, \dots, a_k)$ .  $\square$

**Theorem 13.3.2.** *Suppose  $M \preceq N$  and  $\varphi$  is an  $\mathcal{L}(M)$ -formula.*

1.  $\varphi(M) \subseteq \varphi(N)$ .
2. If  $\varphi(M)$  is finite, then  $\varphi(M) = \varphi(N)$ .

*Proof.* 1. If  $\bar{b} \in \varphi(M)$ , then  $M \models \varphi(\bar{b})$ , so  $N \models \varphi(\bar{b})$ , and  $\bar{b} \in \varphi(N)$ .

2. Let  $n = |\varphi(M)|$ . Then  $M \models \exists^{=n} \bar{x} \varphi(\bar{x})$ , so the same sentence holds in  $N$ , and so  $|\varphi(N)| = n$ . As  $\varphi(M) \subseteq \varphi(N)$ , the two sets must be equal.  $\square$

**Theorem 13.3.3.** *If  $A \subseteq M \preceq N$ , then  $\text{acl}^M(A) = \text{acl}^N(A)$ .*

*Proof.* If  $b \in \text{acl}^M(A)$ , then  $b$  is in a finite set of the form  $\varphi(M)$  for some  $\mathcal{L}(A)$ -formula  $\varphi(x)$ . By Theorem 13.3.2,  $\varphi(N) = \varphi(M)$ . Then  $b \in \varphi(N) \implies b \in \text{acl}^N(A)$ . Thus  $\text{acl}^M(A) \subseteq \text{acl}^N(A)$ .

The proof that  $\text{acl}^N(A) \subseteq \text{acl}^M(A)$  is identical, exchanging  $M$  and  $N$  (even though there is no symmetry between  $M$  and  $N$ ).  $\square$

**Theorem 13.3.4.** *If  $M \preceq N$ , then  $M$  is algebraically closed as a subset of  $N$ .*

*Proof.*  $\text{acl}^N(M) = \text{acl}^M(M) = M$ .  $\square$

**Theorem 13.3.5.** *Let  $\mathbb{M}$  be a monster model, let  $A$  be a small subset, and let  $b$  be an element. Let  $S$  be the set of realizations of  $\text{tp}(b/A)$ .*

1. *If  $b \in \text{acl}(A)$ , then  $S$  is finite.*
2. *If  $b \notin \text{acl}(A)$ , then  $S$  is large.*

*Proof.* If  $b \in \text{acl}(A)$ , then there is a finite  $A$ -definable set  $X \ni b$ . Every realization of  $\text{tp}(b/A)$  is in  $X$ , so  $S \subseteq X$ .

Conversely, suppose  $S$  is small. Let

$$\Sigma(x) = \text{tp}(b/A) \cup \{x \neq c : c \in S\}.$$

Then  $\Sigma(x)$  is a set of formulas over the small set  $A \cup S$  with no realizations in  $\mathbb{M}$ . By saturation,  $\Sigma(x)$  isn't finitely realizable. Therefore there is  $\varphi(x) \in \text{tp}(b/A)$  and  $c_1, \dots, c_n \in S$  such that

$$\varphi(x) \cup \{x \neq c_i : 1 \leq i \leq n\}$$

isn't realized in  $\mathbb{M}$ . This means that  $\varphi(\mathbb{M}) \subseteq \{c_1, \dots, c_n\}$ . Then  $b$  is in the finite  $A$ -definable set  $\varphi(\mathbb{M})$ .  $\square$

**Remark 13.3.6.** By strong homogeneity, the set  $S$  in Theorem 13.3.5 is the set  $\{\sigma(b) : \sigma \in \text{Aut}(\mathbb{M}/A)\}$ .

## 13.4 Countably categorical fields

**Theorem 13.4.1.** *Let  $M$  be a field and  $A$  be a definably closed subset. Then  $A$  is a subfield.*

*Proof.* By Theorem 13.3.1,  $A$  is a substructure, i.e., a subring. It remains to show that  $A$  is closed under multiplicative inverses. If  $a \in A$  is non-zero, then the  $\mathcal{L}(A)$ -formula  $xa = 1$  defines the set  $\{a^{-1}\}$ , so  $a^{-1} \in \text{dcl}(A) = A$ .  $\square$

**Lemma 13.4.2.** *Let  $T$  be  $\aleph_0$ -categorical, let  $M$  be a model of  $T$ , and let  $A$  be a finite subset of  $M$ . Then  $\text{dcl}(A)$  is finite.*

*Proof.* By Theorem 12.3.5(3),  $S_1(A)$  is finite. Therefore there are only finitely many  $A$ -definable sets  $D \subseteq M^1$ . By definition,  $b \in \text{dcl}(A)$  if and only if  $\{b\}$  is definable. Thus  $\text{dcl}(A)$  is finite.  $\square$

**Theorem 13.4.3.** *There is no  $\aleph_0$ -categorical theory of fields.*

*Proof.* Let  $T$  be an  $\aleph_0$ -categorical theory of fields and let  $K$  be an  $\aleph_1$ -saturated model. For any  $n > 0$ , the set  $\{x \in K : x^n - 1 = 0\}$  is finite, by Theorem 9.1.6. Then the following partial type is finitely satisfiable in  $K$ :

$$\Sigma(x) = \{x \neq 0, x \neq 1, x^2 \neq 1, x^3 \neq 1, x^4 \neq 1, \dots\}.$$

Then  $\Sigma(x)$  is realized in  $K$ . Take  $b \in K$  realizing  $\Sigma(x)$ . Consider the group homomorphism

$$\begin{aligned} f : \mathbb{Z} &\rightarrow K^\times \\ f(n) &= b^n. \end{aligned}$$

As  $b$  realizes  $\Sigma$ , we have  $f(b) = b^n \neq 1$  for  $n \neq 0$ . Therefore  $\ker(f) = \{0\}$ , and the image  $f(\mathbb{Z})$  is isomorphic to  $\mathbb{Z}/\{0\} \cong \mathbb{Z}$ . In particular  $f(\mathbb{Z})$  is infinite. However,  $f(n) = b^n \in \text{dcl}(\{b\})$  for each  $n$ , and  $\text{dcl}(\{b\})$  is finite.  $\square$

## 13.5 Algebraic closure in ACF

If  $M \models \text{ACF}$  and  $K$  is a subfield, then  $K^{\text{alg}}$  denotes the set of  $a \in M$  that are algebraic over  $K$ , meaning that  $P(a) = 0$  for some non-zero polynomial  $P(x) \in K[x]$  (Definition 9.3.1).

**Lemma 13.5.1.** *Suppose  $M \models \text{ACF}$  and  $K$  is a subfield. If  $D \subseteq M^1$  is  $K$ -definable, then there is a finite subset  $S \subseteq K^{\text{alg}}$  such that  $D = S$  or  $D = M \setminus S$ .*

*Proof.* Let  $\mathcal{F}$  be the class of sets of the form  $S$  or  $M \setminus S$  for finite  $S \subseteq K^{\text{alg}}$ . We must show  $D \in \mathcal{F}$ . Note that  $\mathcal{F}$  is closed under boolean combinations.

By quantifier-elimination,  $D = \varphi(K)$  for some quantifier-free  $\mathcal{L}(K)$ -formula  $\varphi$ , which is a boolean combination of atomic formulas. Because  $\mathcal{F}$  is closed under boolean combinations, we may assume that  $\varphi$  is atomic.

Then  $\varphi$  has the form  $P(x) = Q(x)$  for some polynomials  $P, Q \in K[x]$ . If  $P - Q$  is identically zero, then  $\varphi(M) = M$ . Otherwise,  $\varphi(M)$  is the set of roots of  $P - Q$ , which is a subset of  $K^{\text{alg}}$  by definition of  $K^{\text{alg}}$ , and finite by Theorem 9.1.6. Either way,  $\varphi(M) \in \mathcal{F}$ .  $\square$

Fix  $M \models \text{ACF}$ .

**Theorem 13.5.2.** *Let  $K$  be a subfield of  $M$ . Then  $\text{acl}(K) = K^{\text{alg}}$ .*

*Proof.* If  $a \in K^{\text{alg}}$ , then  $a$  is in a finite  $K$ -definable set of the form  $\{x \in M : P(x) = 0\}$  for some  $P(x) \in K[x]$ , and so  $a \in \text{acl}(K)$ .

Conversely, suppose  $a \in \text{acl}(K)$ . Then  $a \in D$  for some finite  $K$ -definable set  $D$ . By Lemma 13.5.1,  $D \subseteq K^{\text{alg}}$ , so  $a \in K^{\text{alg}}$ .  $\square$

**Theorem 13.5.3.** *Let  $K$  be a subset of  $M$ . The following are equivalent:*

1.  $K$  is a subfield and  $K = K^{\text{alg}}$ .
2.  $K$  is a substructure and  $K \models \text{ACF}$ .
3.  $K \preceq M$ .
4.  $K = \text{acl}(K)$ .

*Proof.* We show  $(1) \implies (2) \implies (3) \implies (4) \implies (1)$ .

Assume (1). If  $P(x) \in K[x]$  is non-constant, then there is  $c \in M$  with  $P(c) = 0$ , because  $M \models \text{ACF}$ . The element  $c$  is in  $K^{\text{alg}}$  by definition, so  $c \in K$ . Thus  $K \models \text{ACF}$ .

Assume (2). Then  $K \preceq M$  because ACF has quantifier elimination and both  $K$  and  $M$  are models of ACF (see Theorem 8.3.6).

Assume (3). Then  $K = \text{acl}(K)$  by Theorem 13.3.4.

Assume (4). Then  $K$  is a subfield by Theorem 13.3.1 (definably closed sets are subfields) and  $K = \text{acl}(K) = K^{\text{alg}}$  by Theorem 13.5.2.  $\square$

**Corollary 13.5.4.** *If  $K$  is a subfield of  $M$ , then  $K^{\text{alg}}$  is also a subfield, and  $K^{\text{alg}} \preceq M$ .*

*Proof.*  $\text{acl}(K^{\text{alg}}) = \text{acl}(\text{acl}(K)) = \text{acl}(K) = K^{\text{alg}}$ , so  $K^{\text{alg}}$  satisfies condition (4) of Theorem 13.5.3. Then  $K^{\text{alg}}$  is a subfield and an elementary substructure by (1) and (3) of Theorem 13.5.3.  $\square$

**Example 13.5.5.** The set of algebraic numbers  $\mathbb{Q}^{\text{alg}} \subseteq \mathbb{C}$  is an algebraically closed field, and an elementary substructure of  $\mathbb{C}$ .

## 13.6 Pregeometries and vector spaces

**Definition 13.6.1.** Let  $K$  be a field. A  $K$ -vector space is an abelian group  $(V, +)$  with a function

$$\cdot : K \times V \rightarrow V$$

satisfying the axioms

$$\begin{aligned} a \cdot (v + w) &= (a \cdot v) + (a \cdot w) \\ 1 \cdot v &= v \\ (a + b) \cdot v &= a \cdot v + b \cdot v \\ (a \cdot b) \cdot v &= a \cdot (b \cdot v). \end{aligned} \tag{†}$$

For fixed  $K$ , we can regard the class of  $K$ -vector spaces as an equational class by thinking of the map  $\cdot : K \times V \rightarrow V$  as a family of unary maps

$$\begin{aligned} \mu_a : V &\rightarrow V \\ \mu_a(v) &= a \cdot v, \end{aligned}$$

one for each  $a \in K$ . The axioms in Definition 13.6.1 become axiom schemas. For example, axiom (†) becomes the axiom schema

$$\mu_{a+b}(x) = \mu_a(x) + \mu_b(x) \text{ for each } a, b \in K.$$

Subalgebras of vector spaces are called *linear subspaces*.

**Remark 13.6.2.** In a vector space, the following equations hold:

$$\begin{aligned} (-1) \cdot v &= -v \\ 0 \cdot v &= 0 \\ a \cdot (-v) &= -(a \cdot v) \\ a \cdot 0 &= 0. \end{aligned}$$



Fix a field  $K$  and vector space  $V$ .

**Definition 13.6.3.** The *span* of a set  $S \subseteq V$ , written  $\text{span}(S)$ , is the collection of elements of the form

$$a_1v_1 + \cdots + a_nv_n$$

where  $n \geq 0$ ,  $a_1, \dots, a_n \in K$ , and  $v_1, \dots, v_n \in S$ .

**Theorem 13.6.4.** *The span of  $S$  is the linear subspace of  $V$  generated by  $S$ :*

$$\text{span}(S) = \langle S \rangle_V.$$

*Proof.* It is an exercise in algebra (using Remark 13.6.2) to see that  $\text{span}(S)$  is a linear subspace. It contains  $S$  because if  $v \in S$ , then  $v = 1 \cdot v \in \text{span}(S)$ . Therefore  $\text{span}(S) \supseteq \langle S \rangle_V$ . On the other hand, every element of  $\text{span}(S)$  has the form  $t(v_1, \dots, v_n)$  for some term  $t$  and tuple  $\bar{v}$  in  $S$ , and so  $\text{span}(S) \subseteq \langle S \rangle_V$ .  $\square$

**Corollary 13.6.5.** *Span is a finitary closure operator on  $V$ .*

**Definition 13.6.6.** A *pregeometry* is a pair  $(X, \text{cl})$  where  $X$  is a set,  $\text{cl}$  is a finitary closure operator on  $X$ , and the following *exchange property* holds for  $a, b \in X$  and  $C \subseteq X$ :

$$a \in \text{cl}(C \cup \{b\}) \setminus \text{cl}(C) \implies b \in \text{cl}(C \cup \{a\}).$$

**Theorem 13.6.7.** *In a vector space,  $\text{span}(-)$  has the exchange property, and therefore defines a pregeometry.*

*Proof.* Suppose  $v \in \text{span}(S \cup \{w\})$  but  $v \notin \text{span}(S)$ . Then

$$v = a_1v_1 + \cdots + a_nv_n + bw \tag{*}$$

for some  $a_1, \dots, a_n, b \in K$  and  $v_1, \dots, v_n \in S$ . If  $b = 0$ , then  $v \in \text{span}(S)$ , a contradiction. Thus  $b \neq 0$ , and  $b^{-1}$  exists. Rearranging (\*), we see

$$\begin{aligned} -bw &= a_1v_1 + \cdots + a_nv_n - v \\ w &= (-b^{-1}a_1)v_1 + \cdots + (-b^{-1}a_n)v_n + b^{-1}v \\ w &\in \text{span}(S \cup \{v\}). \end{aligned} \tag{*} \quad \square$$

## Rank

Fix a pregeometry  $(X, \text{cl})$ .

**Definition 13.6.8.** If  $\bar{a} \in X^n$  and  $B \subseteq X$ , the *rank* of  $\bar{a}$  over  $B$ , written  $\text{rk}(\bar{a}/B)$ , is defined to be the number of  $i \in \{1, \dots, n\}$  such that  $a_i \notin \text{cl}(B \cup \{a_1, \dots, a_{i-1}\})$ . We write  $\text{rk}(\bar{a}/\emptyset)$  as  $\text{rk}(\bar{a})$ .

**Theorem 13.6.9.**  $\text{rk}(\bar{a}, \bar{b}/C) = \text{rk}(\bar{a}/C) + \text{rk}(\bar{b}/C\bar{a})$ .

*Proof.* Clear from the definition.  $\square$

**Theorem 13.6.10.**  $\text{rk}(\bar{a}/B) = 0 \iff \{a_1, \dots, a_n\} \subseteq \text{cl}(B)$ .

*Proof.* If  $a_i \in \text{cl}(B)$  for each  $i$ , then  $a_i \in \text{cl}(B \cup \{a_1, \dots, a_{i-1}\})$  for each  $i$ , so  $\text{rk}(\bar{a}/B) = 0$ . Conversely, if  $\text{rk}(\bar{a}/B) = 0$ , then  $a_i \in \text{cl}(B \cup \{a_1, \dots, a_{i-1}\})$  for each  $i$ . If  $S$  is a closed set containing  $B$ , then

$$\{a_1, \dots, a_{i-1}\} \subseteq S \implies a_i \in S$$

and so  $\{a_1, \dots, a_n\} \subseteq S$  by induction. Taking  $S = \text{cl}(B)$ , we see  $\{a_1, \dots, a_n\} \subseteq \text{cl}(B)$ .  $\square$

**Lemma 13.6.11.**  $\text{rk}(a, b/C) = \text{rk}(b, a/C)$ .

*Proof.* Both sides are in  $\{0, 1, 2\}$ . It suffices to show

$$\text{rk}(a, b/C) = 0 \iff \text{rk}(b, a/C) = 0 \tag{13.1}$$

$$\text{rk}(a, b/C) = 2 \iff \text{rk}(b, a/C) = 2 \tag{13.2}$$

By symmetry it suffices to show the  $\Rightarrow$  directions.

(13.1): by Theorem 13.6.10, both sides say  $\{a, b\} \subseteq C$ .

(13.2): suppose  $\text{rk}(a, b/C) = 2$ . Then  $a \notin \text{cl}(C)$  and  $b \notin \text{cl}(Ca)$ . By monotonicity,  $b \notin \text{cl}(C)$ . If  $a \in \text{cl}(Cb)$ , then

$$a \in \text{cl}(Cb) \setminus \text{cl}(C) \text{ but } b \notin \text{cl}(Ca),$$

contradicting the exchange property. Thus  $a \notin \text{cl}(Cb)$ . With  $b \notin \text{cl}(C)$ , this implies  $\text{rk}(b, a/C) = 2$ .  $\square$

**Lemma 13.6.12.** If  $\bar{a}, \bar{d}$  are tuples and  $b, c$  are elements and  $A$  is a set, then

$$\text{rk}(\bar{a}, b, c, \bar{d}/A) = \text{rk}(\bar{a}, c, b, \bar{d}/A).$$

*Proof.* Using Theorem 13.6.9 and Lemma 13.6.11,

$$\begin{aligned} \text{rk}(\bar{a}, b, c, \bar{d}/A) &= \text{rk}(\bar{a}/A) + \text{rk}(b, c/A, \bar{a}) + \text{rk}(\bar{d}/A, \bar{a}, b, c) \\ &= \text{rk}(\bar{a}/A) + \text{rk}(c, b/A, \bar{a}) + \text{rk}(\bar{d}/A, \bar{a}, c, b) \\ &= \text{rk}(\bar{a}, c, b, \bar{d}/A). \end{aligned} \quad \square$$

**Theorem 13.6.13.** *If  $\pi$  is a permutation of  $\{1, \dots, n\}$ , then*

$$\text{rk}(a_1, \dots, a_n/B) = \text{rk}(a_{\pi(1)}, \dots, a_{\pi(n)}/B).$$

*Proof.* Repeated applications of Lemma 13.6.12.  $\square$

**Theorem 13.6.14.** *If  $\bar{a} \subseteq \text{cl}(C\bar{b})$ , then  $\text{rk}(\bar{a}/C) \leq \text{rk}(\bar{b}/C)$ .*

*Proof.*

$$\begin{aligned} \text{rk}(\bar{a}/C) &\leq \text{rk}(\bar{a}/C) + \text{rk}(\bar{b}/C\bar{a}) = \text{rk}(\bar{a}, \bar{b}/C) \\ &= \text{rk}(\bar{b}, \bar{a}/C) = \text{rk}(\bar{b}/C) + \text{rk}(\bar{a}/C\bar{b}) = \text{rk}(\bar{b}/C). \end{aligned} \quad \square$$

**Theorem 13.6.15.** *If  $C \supseteq B$ , then  $\text{rk}(\bar{a}/C) \leq \text{rk}(\bar{a}/B)$ .*

*Proof.* For each  $i$ , monotonicity of  $\text{cl}(-)$  shows

$$a_i \notin \text{cl}(C \cup \{a_1, \dots, a_{i-1}\}) \implies a_i \notin \text{cl}(B \cup \{a_1, \dots, a_{i-1}\}). \quad \square$$

## Independence and bases

**Definition 13.6.16.** A set  $I \subseteq X$  is *independent* if  $a \notin \text{cl}(I \setminus \{a\})$  for each  $a \in I$ .

**Theorem 13.6.17.** *In a vector space  $V$ , a set  $I \subseteq V$  is independent if and only if the following condition holds: for any distinct  $v_1, \dots, v_n \in V$  and  $a_1, \dots, a_n \in K$ ,*

$$a_1 v_1 + \dots + a_n v_n = 0 \implies v_1 = v_2 = \dots = v_n = 0. \quad (*)$$

*Proof.* If  $I$  is not independent, then there is some  $v_1 \in \text{span}(I \setminus \{v_1\})$ . Thus there are distinct  $v_2, \dots, v_n \in I \setminus \{v_1\}$  with

$$v_1 = a_2 v_2 + \dots + a_n v_n.$$

Setting  $a_1 = -1 \in K$ , we get  $\sum_{i=1}^n a_i v_i = 0$ , contradicting  $(*)$ .

Conversely, suppose  $(*)$  fails. Permuting the  $v_i$ , we may assume  $a_1 \neq 0$ . Then

$$\begin{aligned} a_1 v_1 + \cdots + a_n v_n &= 0 \\ v_1 &= (-a_1^{-1} a_2) v_2 + \cdots + (-a_1^{-1} a_n) v_n \in \text{span}(v_2, \dots, v_n), \end{aligned}$$

and  $I$  is not independent. □

Now return to a general pregeometry.

**Definition 13.6.18.** A *basis* is a maximal independent set.

**Remark 13.6.19.** 1. If  $I$  is independent and  $J \subseteq I$ , then  $J$  is independent, by monotonicity of  $\text{cl}(-)$ .

2. A set  $I \subseteq X$  is independent if and only if every finite subset is independent, because  $\text{cl}(-)$  is finitary.

**Theorem 13.6.20.** *There is at least one basis.*

*Proof.* Zorn's lemma. □

**Lemma 13.6.21.** *If  $a_1, \dots, a_n \in X$  are distinct, then  $\text{rk}(\bar{a}) = n$  if and only if the set  $I = \{a_1, \dots, a_n\}$  is independent.*

*Proof.* Note that  $\text{rk}(\bar{a}) = n$  if and only if

$$a_i \notin \text{cl}(a_1, \dots, a_{i-1}) \text{ for all } 1 \leq i \leq n, \quad (*)$$

by definition of rank. Moreover, the condition  $\text{rk}(\bar{a}) = n$  is invariant under permuting  $\bar{a}$ , by Theorem 13.6.13.

Independence clearly implies  $(*)$ , as  $\text{cl}(-)$  is monotone. For the converse, suppose  $\text{rk}(\bar{a}) = n$  and  $i \leq n$ . We must show  $a_i \notin \text{cl}(I \setminus \{a_i\})$ . Permuting  $\bar{a}$ , we reduce to the case  $i = n$ . Then we must show  $a_n \notin \text{cl}(a_1, \dots, a_{n-1})$ , which is a case of  $(*)$ . □

**Lemma 13.6.22.** *If  $I$  is independent and  $a \notin \text{cl}(I)$ , then  $I \cup \{a\}$  is independent.*

*Proof.* By Remark 13.6.19(2), we may assume  $I$  is a finite set  $\{b_1, \dots, b_n\}$ . By Lemma 13.6.21 and Theorem 13.6.9

$$\text{rk}(\bar{b}, a) = \text{rk}(\bar{b}) + \text{rk}(a/\bar{b}) = n + 1,$$

and so  $\{b_1, \dots, b_n, a\}$  is independent.  $\square$

**Theorem 13.6.23.** *If  $B$  is a basis, then  $\text{cl}(B) = X$ .*

*Proof.* Otherwise, take  $a \in X \setminus \text{cl}(B)$ , and  $B \cup \{a\}$  is independent, contradicting the fact that  $B$  is a maximal independent set.  $\square$

**Remark 13.6.24.** Conversely, if  $B$  is independent and  $\text{cl}(B) = X$ , then  $B$  is a basis. Otherwise, if  $I \supsetneq B$  is a larger independent set, then  $I \subseteq X = \text{cl}(B)$ , contradicting Lemma 13.6.25 below. Thus  $B$  is a basis if and only if  $B$  is independent and  $\text{cl}(B) = X$ .

**Lemma 13.6.25.** *If  $I$  is independent and  $J$  is a proper subset, then  $I \not\subseteq \text{cl}(J)$ .*

*Proof.* Otherwise, take  $a \in I \setminus J$ . Then  $J \subseteq I \setminus \{a\}$ , and so

$$a \in I \subseteq \text{cl}(J) \subseteq \text{cl}(I \setminus \{a\}),$$

contradicting independence.  $\square$

**Theorem 13.6.26.** *If  $B$  and  $C$  are two bases, then  $|B| = |C|$ .*

*Proof.* Suppose not. Without loss of generality,  $|B| < |C|$ . There are two cases:

1.  $B$  and  $C$  are finite. Let  $\bar{b}$  and  $\bar{c}$  enumerate  $B$  and  $C$ . Then

$$\text{rk}(\bar{b}) = |B| < |C| = \text{rk}(\bar{c})$$

by Lemma 13.6.21. However,  $C \subseteq X = \text{cl}(B)$  by Theorem 13.6.23, and so  $\text{rk}(\bar{c}) \leq \text{rk}(\bar{b})$  by Theorem 13.6.14.

2.  $C$  is infinite. By Theorem 13.6.23,  $B \subseteq \text{cl}(C)$  and  $C \subseteq \text{cl}(B)$ . As  $\text{cl}(-)$  is finitary, we can find a finite subset  $C_b \subseteq C$  for each  $b \in B$  such that  $b \in \text{cl}(C_b)$ . Let  $C' = \bigcup_{b \in B} C_b$ . Then  $B \subseteq \text{cl}(C')$ , implying  $C \subseteq \text{cl}(B) \subseteq \text{cl}(\text{cl}(C')) = \text{cl}(C')$ . This contradicts independence of  $C$  (see Lemma 13.6.25) unless  $C' = C$ .

However,  $|C'| < |C|$ . Indeed, if  $B$  is finite then  $C'$  is finite, and so  $|C'| < |C|$ . If  $B$  is infinite then  $|C'| \leq |B| < |C|$ . Thus  $C'$  is a proper subset of  $C$ , a contradiction.  $\square$

**Definition 13.6.27.** The *rank* of a pregeometry is the cardinality of any basis.

## 13.7 The classification of vector spaces

Fix a field  $K$ . We can regard  $K$  as a vector space by defining

$$\begin{aligned} a \cdot v &= a \cdot v \\ v + w &= v + w \end{aligned}$$

where the left-hand  $\cdot$  and  $+$  are the vector space operations, and the right-hand  $\cdot$  and  $+$  are the field operations.

If  $\lambda$  is a cardinal, let  $K^\lambda$  be the power vector space, i.e., the set of functions  $f : \lambda \rightarrow K$  with the vector space operations defined pointwise:

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (a \cdot f)(x) &= a \cdot (f(x)). \end{aligned}$$

Say that  $f \in K^\lambda$  has *finite support* if

$$\text{supp}(f) := \{x \in \lambda : f(x) \neq 0\}$$

is finite. Note that

$$\begin{aligned} \text{supp}(f + g) &\subseteq \text{supp}(f) \cup \text{supp}(g) \\ \text{supp}(af) &\subseteq \text{supp}(f) \\ \text{supp}(0) &= \emptyset. \end{aligned}$$

Therefore,  $F_\lambda := \{f \in K^\lambda : \text{supp}(f) \text{ is finite}\}$  is a linear subspace of  $K^\lambda$ .

**Lemma 13.7.1.** Let  $V$  be a  $K$ -vector space. Suppose  $v_i \in V$  for each  $i \in \lambda$ .

1. There is a homomorphism  $\alpha : F_\lambda \rightarrow V$  defined by

$$\alpha(f) = \sum_{i \in \text{supp}(f)} f(i) \cdot v_i.$$

2.  $\alpha$  is surjective if and only if  $V = \text{span}\{v_i : i \in \lambda\}$ .

3.  $\alpha$  is injective if and only if  $\{v_i : i \in \lambda\}$  is independent (and the  $v_i$  are distinct).
4.  $\alpha$  is an isomorphism if and only if  $\{v_i : i \in \lambda\}$  is a basis (and the  $v_i$  are distinct).
5. All homomorphisms  $F_\lambda \rightarrow V$  arise as in part (1).

*Proof.* 1. An exercise in algebra. Essentially this works because  $\alpha(f) = \sum_{i \in \lambda} f(i) \cdot v_i$ , which depends linearly on  $f$ . The sum makes sense as almost all the terms in it are zero.

2. True by definition of span.

3.  $\alpha$  is injective if and only if  $\ker(\alpha) = \{0\}$ . This condition means that if  $f \in K^\lambda$  has finite support and

$$\sum_{i \in \lambda} f(i)v_i = 0,$$

then  $f$  must vanish. This is a rephrasing of Theorem 13.6.17.

4.  $\alpha$  is an isomorphism if and only if it is injective and surjective. By the previous two points, this means that the set  $B = \{v_i : i \in \lambda\}$  is independent and has  $\text{span}(B) = V$ . By Remark 13.6.24, this means that  $B$  is a basis.

5. Let  $\beta : F_\lambda \rightarrow V$  be a homomorphism. Let  $e_i : \lambda \rightarrow K$  be the function

$$e_i(x) = \begin{cases} 1 & \text{if } x = i \\ 0 & \text{if } x \neq i. \end{cases}$$

Then  $\text{supp}(e_i) = \{i\}$ , so  $e_i \in F_\lambda$ . Note that for any  $f \in F_\lambda$ ,

$$f = \sum_{i \in \text{supp}(f)} f(i) \cdot e_i.$$

Therefore

$$\beta(f) = \sum_{i \in \text{supp}(f)} f(i) \cdot \beta(e_i).$$

Taking  $v_i = \beta(e_i)$ , we see that  $\beta$  has the desired form.  $\square$

Lemma 13.7.1 immediately yields the following:

**Theorem 13.7.2.** *Let  $V$  be a  $K$ -vector space. Then  $V \cong F_\lambda$  if and only if  $V$  has a basis of cardinality  $\lambda$ .*

By Theorems 13.6.20 and 13.6.26, there is a unique cardinal  $\lambda$  such that  $V$  has a basis of cardinality  $\lambda$ .

**Theorem 13.7.3.** *Let  $V$  be a  $K$ -vector space. Then  $V \cong F_\lambda$  for a unique cardinal  $\lambda$ , called the dimension of  $V$ .*

**Lemma 13.7.4.** *Let  $V$  be a  $K$ -vector space of dimension  $\lambda$ .*

1. *If  $\lambda$  is finite, then  $|V| = |K|^\lambda$ .*
2. *If  $\lambda$  or  $|K|$  is infinite, then  $|V| = |K| + \lambda$ .*

*Proof.* Without loss of generality,  $V = F_\lambda$ .

1.  $F_\lambda = K^\lambda$ , whose cardinality is  $|K|^\lambda$ .
2. If  $B \subseteq V$  is a basis of size  $\lambda$ , then  $B \subseteq V \implies \lambda \leq |V|$ . If  $v \in V$  is non-zero, then there is an injection

$$\begin{aligned} K &\rightarrow V \\ a &\mapsto av, \end{aligned}$$

and so  $|K| \leq |V|$ . Thus  $|V| \geq \max(|K|, \lambda) = |K| + \lambda$ . On the other hand,  $V$  is generated by a set of size  $\lambda$ , and the language has size  $\aleph_0 + |K|$ , so that  $|V| \leq \lambda + |K| + \aleph_0 = |K| + \lambda$ .  $\square$

**Theorem 13.7.5.** *If  $V$  is an infinite  $K$ -vector space and  $|V| > |K|$ , then  $\dim(V) = |V|$ .*

*Proof.* Let  $\lambda = \dim(V)$ . If  $K$  and  $\lambda$  are both finite then  $|V| = |K|^\lambda < \aleph_0$ , a contradiction. Therefore  $K$  or  $\lambda$  is infinite, and so  $|V| = \max(|K|, \lambda)$ . As  $|V| > |K|$ , this implies  $|V| = \lambda$ .  $\square$

**Corollary 13.7.6.** *Let  $K$  be a field. The theory of infinite  $K$ -vector spaces is complete and  $\lambda$ -categorical for every infinite  $\lambda > |K|$ .*

**Definition 13.7.7.** A theory  $T$  in a countable language is *totally categorical* if  $T$  is  $\lambda$ -categorical for any  $\lambda$ .

**Corollary 13.7.8.** *If  $K$  is a finite field, then the theory of infinite  $K$ -vector spaces is totally categorical.*



# Chapter 14

## Strongly minimal and geometric theories

### 14.1 Strong minimality

**Definition 14.1.1.** Let  $S$  be a set. A subset  $X \subseteq S$  is *cofinite* (in  $S$ ) if  $S \setminus X$  is finite.

**Definition 14.1.2.** A structure  $M$  is *minimal* if  $M$  is infinite, and every definable set  $X \subseteq M$  is finite or cofinite.

Note this is only a statement about definable sets in one variable; it says nothing about definable subsets of  $M^n$  for  $n > 1$ .

**Definition 14.1.3.** A theory  $T$  is *strongly minimal* if all its models are minimal. A structure  $M$  is *strongly minimal* if its complete theory is strongly minimal.

Note that models of a strongly minimal theory are strongly minimal.

**Example 14.1.4.** ACF is strongly minimal by Lemma 13.5.1.

**Theorem 14.1.5.** Suppose  $M$  is strongly minimal and  $A \subseteq M$ . Then  $A \preceq M$  if and only if  $A = \text{acl}(A)$  and  $A$  is infinite.

*Proof.* First suppose  $A \preceq M$ . Then  $A$  is infinite because  $M$  is (Lemma 5.5.4), and  $A = \text{acl}(A)$  because  $A$  is an elementary substructure (Theorem 13.3.4).

Conversely, suppose  $A = \text{acl}(A)$  and  $A$  is infinite. Then  $A \preceq M$  by the Tarski-Vaught criterion (Theorem 5.3.1). Indeed, suppose  $D \subseteq M$  is non-empty and  $A$ -definable. By strong minimality,  $D$  is finite or cofinite. If  $D$  is finite then  $D \subseteq \text{acl}(A) = A$ , so  $A \cap D = D \neq \emptyset$ . If  $D$  is cofinite, it intersects  $A$  because  $A$  is infinite.  $\square$

**Theorem 14.1.6.** *If  $M$  is strongly minimal, then  $\text{acl}(-)$  satisfies the exchange property:*

$$a \in \text{acl}(Cb) \setminus \text{acl}(C) \implies b \in \text{acl}(Ca).$$

Consequently,  $\text{acl}(-)$  defines a pregeometry.

*Proof.* Because  $a \in \text{acl}(Cb)$ , there is a finite  $Cb$ -definable set  $X$  containing  $a$ . Write  $X$  as  $\varphi(M, b)$  for some  $\mathcal{L}(C)$ -formula  $\varphi(x, y)$ . Let  $n = |X|$ . Replacing  $\varphi(x, y)$  with

$$\varphi(x, y) \wedge \neg \exists^{\geq n+1} z \varphi(z, y),$$

we may assume that  $|\varphi(M, b')| \leq n$  for any  $b' \in M$ . As  $\varphi(a, b)$  holds,  $b \in \varphi(a, M)$ . If  $\varphi(a, M)$  is finite then  $b \in \text{acl}(Ca)$  as desired.

Otherwise,  $\varphi(a, M)$  is cofinite, and its complement has cardinality  $k < \infty$ . Let  $D$  be the  $C$ -definable set of  $a' \in M$  such that  $|M \setminus \varphi(a', M)| = k$ . Then  $a \in D$ , so  $D$  is infinite because  $a \notin \text{acl}(C)$ . Take distinct  $a_1, \dots, a_{n+1} \in D$ . Then  $\varphi(a_i, M)$  is cofinite for each  $i$ . An intersection of cofinite sets is cofinite, hence non-empty, so there is some

$$b' \in \bigcap_{i=1}^{n+1} \varphi(a_i, M).$$

This means that  $\varphi(a_i, b')$  holds for all  $i$ , and so  $|\varphi(M, b')| \geq n+1$ , contradicting the choice of  $\varphi$ .  $\square$

## 14.2 Uncountable categoricity

Let  $\mathbb{M}$  be a monster model of a *complete* strongly minimal theory in a countable language.

**Theorem 14.2.1.** *For any small set  $A \subseteq \mathbb{M}$ , there is a unique 1-type  $p \in S_1(A)$  whose realizations are the elements of  $\mathbb{M} \setminus \text{acl}(A)$ .*

*Proof.* In other words, we must show  $b, c \notin \text{acl}(A)$  implies  $b \equiv_A c$ . Otherwise there is an  $A$ -definable set  $D$  with  $b \in D$  and  $c \notin D$  (Remark 8.1.3). If  $D$  is finite, then  $b \in \text{acl}(A)$  and if  $D$  is cofinite, then  $c \in \text{acl}(A)$ . Either way, we get a contradiction.  $\square$

The pregeometry rank  $\text{rk}(\bar{a}/B)$  (Definition 13.6.8) is usually written as  $\dim(\bar{a}/B)$ . Say that a finite tuple  $\bar{a} \in \mathbb{M}^n$  is *independent* if  $\dim(\bar{a}) = n$ . By definition, this means  $a_i \notin \text{acl}(a_1, \dots, a_{i-1})$  for each  $i$ . By Lemma 13.6.21,  $\bar{a}$  is independent if and only if the  $a_i$  are pairwise distinct and  $\{a_1, \dots, a_n\}$  is independent as a set.

**Lemma 14.2.2.** *Let  $\bar{a}, \bar{b}$  be two finite tuples of length  $n$  in  $\mathbb{M}$ . If  $\dim(\bar{a}) = n$  and  $\dim(\bar{b}) = n$ , then  $\bar{a} \equiv_{\emptyset} \bar{b}$ .*

*Proof.* Proceed by induction on  $n$ . The case  $n = 0$  is trivial. Note that the subtuples  $(a_1, \dots, a_{n-1})$  and  $(b_1, \dots, b_{n-1})$  are independent. By induction,  $(a_1, \dots, a_{n-1}) \equiv_{\emptyset} (b_1, \dots, b_{n-1})$ . Take  $\sigma \in \text{Aut}(\mathbb{M})$  moving  $(a_1, \dots, a_{n-1})$  to  $(b_1, \dots, b_{n-1})$ . Replacing  $\bar{a}$  with  $\sigma(\bar{a})$ , we may assume  $a_i = b_i$  for  $i < n$ . By independence,

$$\begin{aligned} a_n &\notin \text{acl}(a_1, \dots, a_{n-1}) \\ b_n &\notin \text{acl}(b_1, \dots, b_{n-1}) = \text{acl}(a_1, \dots, a_{n-1}). \end{aligned}$$

By Theorem 14.2.1,  $a_n$  and  $b_n$  have the same type over  $(a_1, \dots, a_{n-1})$ . Therefore  $(a_1, \dots, a_n) \equiv_{\emptyset} (a_1, \dots, a_{n-1}, b_n) = \bar{b}$ .  $\square$

**Theorem 14.2.3.** *Let  $M$  be a model of  $T$ . Let  $f : I_1 \rightarrow I_2$  be a bijection between two independent sets. Then  $f$  is a partial elementary map.*

*Proof.* If  $a_1, \dots, a_n$  are distinct elements of  $I_1$ , then  $f(a_1), \dots, f(a_n)$  are distinct elements of  $I_2$ , and the two tuples  $\bar{a}$  and  $f(\bar{a})$  are both independent  $n$ -tuples. By Lemma 14.2.2, they have the same type.  $\square$

**Definition 14.2.4.** If  $M \models T$ , the *rank* of  $M$ , written  $\text{rk}(M)$ , is the rank of the pregeometry  $(M, \text{acl}(-))$ , i.e., the cardinality of a basis.

**Remark 14.2.5.** If  $M \models T$ , and  $B$  is a basis, then  $M = \text{acl}(B)$  (Theorem 13.6.23), and so

$$|M| = |\text{acl}(B)| = |B| + \aleph_0 = \text{rk}(M) + \aleph_0.$$

**Theorem 14.2.6.** *Two models  $M_1, M_2$  are isomorphic if and only if  $\text{rk}(M_1) = \text{rk}(M_2)$ .*

*Proof.* If  $M_1 \cong M_2$ , it is easy to see that  $\text{rk}(M_1) = \text{rk}(M_2)$ . Conversely, suppose the ranks are equal. Embed  $M_1$  and  $M_2$  into a monster model  $\mathbb{M}$ . Let  $B_i$  be a basis of  $M_i$  for  $i = 1, 2$ . Then  $|B_1| = |B_2|$ . Take a bijection  $f : B_1 \rightarrow B_2$ . By Theorem 14.2.3,  $f$  is a partial elementary map from  $\mathbb{M}$  to  $\mathbb{M}$ . By strong homogeneity,  $f$  extends to an automorphism  $\sigma \in \text{Aut}(\mathbb{M})$ . Then  $\sigma(M_1) = \sigma(\text{acl}(B_1)) = \text{acl}(B_2) = M_2$ . The automorphism  $\sigma$  restricts to an isomorphism from  $M_1$  to  $M_2$ .  $\square$

**Corollary 14.2.7.**  *$T$  is  $\kappa$ -categorical for all  $\kappa > \aleph_0$ .*

**Theorem 14.2.8.** *Suppose that in models of  $T$ , algebraically closed sets are infinite. Then there is a unique model of rank  $\kappa$  for every cardinal  $\kappa$ , finite or infinite.*

*Proof.* Take a model  $M_0$  of size greater than  $\kappa + \aleph_0$ , and let  $B_0$  be a basis of  $M_0$ . Then  $|B_0| > \kappa$ . Take a subset  $I \subseteq B_0$  with  $|I| = \kappa$ , and let  $M = \text{acl}(I)$ . By assumption,  $M$  is infinite. Then  $M \preceq M_0$  by Theorem 14.1.5. By Remark 13.6.24, the independent set  $I$  is a basis of  $M$ , so  $\text{rk}(M) = |I| = \kappa$ .  $\square$

## The classification of algebraically closed fields

**Definition 14.2.9.** The *transcendence degree* of an algebraically closed field  $K$  is its rank in the sense of Definition 14.2.4.

**Theorem 14.2.10.** *For each  $p \in \{0, 2, 3, 5, 7, \dots\}$  and each cardinal  $\kappa$ , there is a unique algebraically closed field of characteristic  $p$  and transcendence degree  $\kappa$ .*

*Proof.* The theory  $\text{ACF}_p$  is complete and strongly minimal. Algebraically closed sets are infinite by Theorem 13.5.3(2) and Theorem 9.2.3. Therefore Theorem 14.2.8 applies, giving a unique model  $M \models \text{ACF}_p$  with transcendence degree  $\kappa$ .  $\square$

## 14.3 Uniform finiteness

The notation  $\exists^\infty x P(x)$  means that there are infinitely many  $x$  such that  $P(x)$  holds. In general,  $\exists^\infty x$  cannot be expressed in first-order logic.

**Definition 14.3.1.** A structure  $M$  *eliminates*  $\exists^\infty$  if for any first-order formula  $\varphi(x, \bar{y})$ , there is a formula  $\psi(\bar{y})$  such that for any  $\bar{b}$ ,

$$M \models \exists^\infty x \varphi(x, \bar{b}) \iff M \models \psi(\bar{b}).$$

Here, the left-hand side really means  $\exists^\infty a \in M : M \models \varphi(a, \bar{b})$ , or equivalently,  $\varphi(M, \bar{b})$  is infinite.

Recall from Section 3.3 that the notation  $\exists^{\geq n} x P(x)$  means that there are at least  $n$  values of  $x$  such that  $P(x)$  holds. Unlike  $\exists^\infty$ , this can be expressed in first-order logic.

**Definition 14.3.2.** A structure  $M$  has *uniform finiteness* if for any formula  $\varphi(x, \bar{y})$ , there is a number  $n_\varphi$  such that for any  $\bar{b}$  in  $M$ ,

$$|\varphi(M, \bar{b})| < \infty \iff |\varphi(M, \bar{b})| < n_\varphi.$$

**Theorem 14.3.3.** *Let  $M$  be a structure.*

1. *If  $M$  has uniform finiteness, then  $M$  eliminates  $\exists^\infty$ .*
2. *If  $M$  has uniform finiteness and  $N \equiv M$ , then  $N$  has uniform finiteness.*
3.  *$M$  has uniform finiteness if and only if every  $N \equiv M$  eliminates  $\exists^\infty$ .*

*Proof.* 1. Uniform finiteness says  $(\exists^\infty x) \varphi(x, \bar{y}) \iff (\exists^{\geq n_\varphi} x) \varphi(x, \bar{y})$ .

2. Given  $\varphi$ , if  $n_\varphi$  works for  $M$  then it works for  $N$ . Otherwise there is  $\bar{b}$  in  $N$  such that  $n_\varphi \leq |\varphi(N, \bar{b})| < \infty$ . Let  $k = |\varphi(N, \bar{b})|$ . Then

$$N \models \exists \bar{y} \exists^=k x \varphi(x, \bar{y}).$$

This says that there is a  $\bar{b}$  in  $N$  such that  $\varphi(N, \bar{b})$  has size exactly  $k$ . As  $M \equiv N$ , the same holds in  $M$ , so there is a  $\bar{b}$  in  $M$  such that  $\varphi(M, \bar{b})$  has size exactly  $k \geq n_\varphi$ , contradicting the choice of  $n_\varphi$ .

3. If  $M$  has uniform finiteness and  $N \equiv M$ , then  $N$  has uniform finiteness and therefore eliminates  $\exists^\infty$ . Conversely, suppose every  $N \equiv M$  eliminates  $\exists^\infty$ . Fix an  $\aleph_1$ -saturated elementary extension  $N \succeq M$ . By

part (2), it suffices to show that  $N$  has uniform finiteness. Fix  $\varphi(x, \bar{y})$ . Let

$$D_\infty = \{\bar{b} : |\varphi(N, \bar{b})| < \infty\}$$

$$D_k = \{\bar{b} : |\varphi(N, \bar{b})| < k\} \text{ for } k < \omega.$$

Then  $D_\infty$  is definable by elimination of  $\exists^\infty$ , and  $D_k$  is definable easily. Moreover,

$$D_0 \subseteq D_1 \subseteq \cdots \subseteq D_\infty$$

and  $D_\infty = \bigcup_{i=0}^\infty D_i$ . By  $\aleph_0$ -compactness (Theorem 11.2.7(3)),  $D_\infty = D_{n_\varphi}$  for some  $n_\varphi$ , which means

$$|\varphi(N, \bar{b})| < \infty \implies |\varphi(N, \bar{b})| < n_\varphi. \quad \square$$

**Theorem 14.3.4.** *If  $M$  is strongly minimal, then  $M$  has uniform finiteness.*

*Proof.* Without loss of generality,  $M$  is  $\aleph_1$ -saturated. Let  $\varphi(x; y_1, \dots, y_n)$  be a formula. Let  $D_k$  be the set of  $\bar{b}$  such that  $\varphi(M, \bar{b})$  or  $M \setminus \varphi(M, \bar{b})$  has size less than  $k$ . Then  $D_0 \subseteq D_1 \subseteq \cdots \subseteq M^n$ . Strong minimality means that  $M^n = \bigcup_{k=0}^\infty D_k$ . By  $\aleph_0$ -compactness (Theorem 11.2.7(3)),  $D_k = M^n$  for some  $k$ . This means that for any  $\bar{b}$ ,

$$|\varphi(M, \bar{b})| < k \text{ or } |M \setminus \varphi(M, \bar{b})| < k.$$

Therefore,

$$|\varphi(M, \bar{b})| < k \text{ or } |\varphi(M, \bar{b})| = \infty,$$

which is uniform finiteness.  $\square$

## 14.4 Pregeometric theories

**Definition 14.4.1.** A theory  $T$  is *pregeometric* if  $\text{acl}(-)$  satisfies exchange in models of  $T$ .

**Definition 14.4.2.** A complete theory  $T$  is *geometric* if it is pregeometric and has uniform finiteness.

For example, strongly minimal theories are geometric.

Let  $\mathbb{M}$  be a monster model of a pregeometric theory. Let  $\dim(\bar{a}/B)$  denote the rank of  $\bar{a}$  over  $B$  with respect to  $\text{acl}(-)$ , i.e., the number of values  $i \in \{1, \dots, n\}$  such that  $a_i \notin \text{acl}(Ba_1a_2 \cdots a_{i-1})$ .

**Theorem 14.4.3.** 1.  $\dim(\bar{a}/B) = \dim(\sigma(\bar{a})/\sigma(B))$  for any  $\sigma \in \text{Aut}(\mathbb{M})$ .

2. If  $\bar{a} \equiv_C \bar{b}$ , then  $\dim(\bar{a}/C) = \dim(\bar{b}/C)$ .

*Proof.* 1. The definition of  $\dim(-/-)$  is clearly automorphism-invariant.

2. Take  $\sigma \in \text{Aut}(\mathbb{M}/C)$  with  $\sigma(\bar{a}) = \bar{b}$  and apply part (1).  $\square$

In particular,  $\text{tp}(\bar{a}/C)$  determines  $\dim(\bar{a}/C)$ . If  $p \in S_n(C)$ , we let  $\dim(p)$  denote  $\dim(\bar{a}/C)$  for any  $\bar{a}$  realizing  $p$ .

**Theorem 14.4.4** (Extension). *Let  $B \subseteq C$  be small sets in  $\mathbb{M}$ .*

1. If  $p \in S_n(B)$ , then there is an extension  $q \in S_n(C)$  with  $q \supseteq p$ .

2. If  $\bar{a} \in \mathbb{M}^n$ , then there is  $\bar{a}' \equiv_B \bar{a}$  with  $\dim(\bar{a}'/C) = \dim(\bar{a}/B)$ .

*Proof.* Note that (1)  $\iff$  (2), by taking  $p = \text{tp}(\bar{a}/B)$  and  $q = \text{tp}(\bar{a}'/B)$ . We prove (2) by induction on  $n$ . First suppose  $n = 1$ . If  $a \notin \text{acl}(B)$ , then the set  $X$  of realizations of  $\text{tp}(a/B)$  is large by Theorem 13.3.5, so there is  $a' \in X \setminus \text{acl}(C)$ . Then  $a' \equiv_B a$  and  $\dim(a'/C) = 1 = \dim(a/B)$ . On the other hand, if  $a \in \text{acl}(B) \subseteq \text{acl}(C)$ , then  $\dim(a/C) = 0 = \dim(a/B)$ , and we can take  $a' = a$ .

Next suppose  $n > 1$ . Write  $\bar{a}$  as  $(\bar{a}_1, \bar{a}_2)$ , where  $\bar{a}_1$  and  $\bar{a}_2$  are tuples of shorter length. By induction there is  $\bar{a}'_1 \equiv_B \bar{a}_1$  with  $\dim(\bar{a}'_1/C) = \dim(\bar{a}_1/B)$ . Moving  $\bar{a}$  by an automorphism over  $B$  sending  $\bar{a}_1$  to  $\bar{a}'_1$ , we may assume  $\bar{a}_1 = \bar{a}'_1$ . Then

$$\dim(\bar{a}_1/C) = \dim(\bar{a}_1/B). \quad (*)$$

By induction applied to  $\bar{a}_2$  and the inclusion  $B\bar{a}_1 \subseteq C\bar{a}_1$ , there is  $\bar{a}'_2 \equiv_{B\bar{a}_1} \bar{a}_2$  with  $\dim(\bar{a}'_2/C\bar{a}_1) = \dim(\bar{a}_2/B\bar{a}_1)$ . Moving  $\bar{a}$  by an automorphism over  $B\bar{a}_1$  sending  $\bar{a}_2$  to  $\bar{a}'_2$ , we may assume  $\bar{a}'_2 = \bar{a}_2$ . Then

$$\dim(\bar{a}_2/C\bar{a}_1) = \dim(\bar{a}_2/B\bar{a}_1). \quad (\dagger)$$

Adding equations  $(*)$  and  $(\dagger)$  and using additivity,

$$\dim(\bar{a}_1\bar{a}_2/C) = \dim(\bar{a}_1\bar{a}_2/B). \quad \square$$

## 14.5 Dimension theory in the monster

Work in a monster model  $\mathbb{M}$  of a pregeometric theory.

**Definition 14.5.1.** If  $B$  is small and  $X \subseteq \mathbb{M}^n$  is  $B$ -definable, then  $\dim_B(X) = \max_{\bar{a} \in X} \dim(\bar{a}/B)$ . If  $X$  is empty, we set  $\dim(X) = -\infty$ .

**Theorem 14.5.2.** If  $X$  is  $B$ -definable and  $C$ -definable, then  $\dim_B(X) = \dim_C(X)$ .

*Proof.* First suppose  $B \subseteq C$ . Note that  $\dim(\bar{a}/B) \geq \dim(\bar{a}/C)$  for any  $\bar{b}$  (Theorem 13.6.15), so

$$\dim_B(X) = \max_{\bar{a} \in X} \dim(\bar{a}/B) \geq \max_{\bar{a} \in X} \dim(\bar{a}/C) = \dim_C(X).$$

Conversely, take  $\bar{a} \in X$  with  $\dim(\bar{a}/B) = \dim_B(X)$ . By Theorem 14.4.4, there is  $\bar{a}' \equiv_B \bar{a}$  with  $\dim(\bar{a}'/C) = \dim(\bar{a}/B)$ . Then  $\bar{a} \in X \implies \bar{a}' \in X$  because  $X$  is  $B$ -definable (Remark 8.1.3), and so

$$\dim_C(X) \geq \dim(\bar{a}'/C) = \dim(\bar{a}/B) = \dim_B(X).$$

This completes the case where  $B \subseteq C$ . The general case then follows:

$$\dim_B(X) = \dim_{B \cup C}(X) = \dim_C(X). \quad \square$$

By Theorem 14.5.2,  $\dim_B(X)$  doesn't depend on  $B$ , so we just write  $\dim(X)$ . The following two facts are a restatement of the definition:

- If  $X$  is  $B$ -definable and  $\bar{a} \in X$ , then  $\dim(\bar{a}/B) \leq \dim(X)$ .
- If  $X$  is  $B$ -definable and non-empty, then there is  $\bar{a} \in X$  with  $\dim(\bar{a}/B) = \dim(X)$ .

We will use these repeatedly in what follows.

**Theorem 14.5.3** (Basic properties of dimension). *Let  $X, Y$  be definable sets.*

1.  $\dim(X) \leq 0$  if and only if  $|X| < \infty$ .
2. If  $X \subseteq Y$ , then  $\dim(X) \leq \dim(Y)$ .
3. If  $X, Y \subseteq \mathbb{M}^n$ , then  $\dim(X \cup Y) = \max(\dim(X), \dim(Y))$ .



4.  $\dim(X \times Y) = \dim(X) + \dim(Y)$ .
5.  $\dim(\mathbb{M}^n) = n$ , assuming  $\mathbb{M}$  is infinite.
6. If  $f : X \rightarrow Y$  is a definable surjection, then  $\dim(X) \geq \dim(Y)$ .
7. If  $f : X \rightarrow Y$  is a definable bijection, then  $\dim(X) = \dim(Y)$ .
8. If  $f : X \rightarrow Y$  is a definable injection, then  $\dim(X) \leq \dim(Y)$ .

*Proof.* Take a small set  $C$  over which all the sets and functions are defined.

1. If  $X$  is finite, then every  $\bar{a} \in X$  is algebraic over  $C$ , and so  $\dim(\bar{a}/C) \leq 0$  (Theorem 13.6.10). Thus  $\dim(X) \leq 0$ .

Conversely, if  $X$  is infinite, then  $X$  is large (Corollary 11.2.8), so there is  $\bar{a} \in X$  with  $\bar{a} \notin \text{acl}(C)$ . Then  $\dim(X) \geq \dim(\bar{a}/C) > 0$ .

2. Clear.

3. Clear.

4. If  $(\bar{a}, \bar{b}) \in X \times Y$ , then

$$\dim(\bar{a}\bar{b}/C) = \dim(\bar{a}/C) + \dim(\bar{b}/C\bar{a}) \leq \dim(X) + \dim(Y)$$

because  $\bar{a}$  is in the  $C$ -definable set  $X$  and  $\bar{b}$  is in the  $C\bar{a}$ -definable set  $Y$ . As this holds for all  $(\bar{a}, \bar{b}) \in X \times Y$ , we have  $\dim(X \times Y) \leq \dim(X) + \dim(Y)$ .

Conversely, take  $\bar{a} \in X$  with  $\dim(\bar{a}/C) = \dim(X)$ . The set  $Y$  is  $C\bar{a}$ -definable, so there is  $\bar{b} \in Y$  with  $\dim(\bar{b}/C\bar{a}) = \dim(Y)$ . Then

$$\dim(X \times Y) \geq \dim(\bar{a}\bar{b}/C) = \dim(\bar{a}/C) + \dim(\bar{b}/C\bar{a}) = \dim(X) + \dim(Y).$$

5. By part (4), it suffices to show  $\dim(\mathbb{M}^1) = 1$ . If  $a \in \mathbb{M}^1$ , then  $\dim(a/C) \leq 1$ , as  $a$  has length 1. Thus  $\dim(\mathbb{M}^1) \leq 1$ . On the other hand,  $\dim(\mathbb{M}) \geq 1$  by part (1).
6. Take  $\bar{b} \in Y$  with  $\dim(\bar{b}/C) = \dim(Y)$ . Since  $f$  is surjective, there is  $\bar{a} \in X$  with  $f(\bar{a}) = \bar{b}$ . Then  $\bar{b} \in \text{dcl}(C\bar{a}) \subseteq \text{acl}(C\bar{a})$ , so by Theorem 13.6.14,

$$\dim(X) \geq \dim(\bar{a}/C) \geq \dim(\bar{b}/C) = \dim(Y).$$

7. Apply part (6) to  $f$  and  $f^{-1}$ .

8. By parts (2) and (7),  $\dim(X) = \dim(\text{im}(f)) \leq \dim(Y)$ .  $\square$

**Theorem 14.5.4** (Fiber dimension theorem). *Let  $f : X \rightarrow Y$  be a definable function. For every  $b \in Y$ , let  $X_b = f^{-1}(b) = \{x \in X : f(x) = b\}$ . If  $\dim(X_b) = k$  for all  $b \in Y$ , then  $\dim(X) = k + \dim(Y)$ .*

*Proof.* Take a small set  $C$  defining  $f$ ,  $X$ , and  $Y$ . Note that  $\bar{a}$  and  $(\bar{a}, f(\bar{a}))$  are interalgebraic over  $C$  for any  $\bar{a} \in X$ , so

$$\dim(\bar{a}/C) = \dim(\bar{a}, f(\bar{a})/C) = \dim(\bar{a}/Cf(\bar{a})) + \dim(f(\bar{a})/C)$$

by Theorem 13.6.14 and additivity (Theorem 13.6.9). If  $\bar{b} = f(\bar{a})$ , then

$$\dim(\bar{a}/C) = \dim(\bar{a}/C\bar{b}) + \dim(\bar{b}/C) \leq \dim(X_b) + \dim(Y) = k + \dim(Y).$$

because  $\bar{a}$  is in the  $C\bar{b}$ -definable set  $X_b$ , and  $\bar{b}$  is in the  $C$ -definable set  $Y$ . As this holds for any  $\bar{a} \in X$ , we see

$$\dim(X) \leq k + \dim(Y).$$

For the converse, take  $\bar{b} \in Y$  with  $\dim(\bar{b}/C) = \dim(Y)$ . The set  $X_{\bar{b}}$  is  $C\bar{b}$ -definable, so there is  $\bar{a} \in X_{\bar{b}}$  with  $\dim(\bar{a}/C\bar{b}) = \dim(X_b) = k$ . Then  $\bar{b} = f(\bar{a})$ , so

$$\dim(X) \geq \dim(\bar{a}/C) = \dim(\bar{a}/C\bar{b}) + \dim(\bar{b}/C) = k + \dim(Y). \quad \square$$

*Proof.* Let  $Z_k = f^{-1}(Y_k)$ . Then  $\dim(Z_k) = k + \dim(Y_k)$ , and  $X = \bigcup_k Z_k$ .  $\square$

**Now suppose the theory is geometric (Definition 14.4.2),** meaning that uniform finiteness holds.

**Lemma 14.5.5.** *Let  $\varphi(x_1, \dots, x_n)$  be an  $\mathcal{L}(\mathbb{M})$ -formula and let  $\psi(x_1, \dots, x_{n-1})$  be an  $\mathcal{L}(\mathbb{M})$ -formula equivalent to  $\exists^\infty x_n \varphi$ . Then  $\varphi(\mathbb{M}^n)$  has dimension  $n$  if and only if  $\psi(\mathbb{M}^{n-1})$  has dimension  $n - 1$ .*

*Proof.* Take a finite set  $B$  such that  $\varphi$  and  $\psi$  are  $\mathcal{L}(B)$ -formulas. If  $\psi(\mathbb{M}^{n-1})$  has dimension  $n - 1$ , take  $\bar{a} \in \psi(\mathbb{M}^{n-1})$  with  $\dim(\bar{a}/B) = n - 1$ . Then  $\varphi(\bar{a}, \mathbb{M})$  is infinite, hence large. Take  $a_n \in \varphi(\bar{a}, \mathbb{M}) \setminus \text{acl}(B\bar{a})$ . Then  $(\bar{a}, a_n) \in \varphi(\mathbb{M}^n)$  and

$$\dim(\bar{a}, a_n/B) = \dim(\bar{a}/B) + \dim(a_n/B\bar{a}) = (n - 1) + 1 = n,$$

and  $\varphi(\mathbb{M}^n)$  has dimension  $n$ .

Conversely, suppose  $\varphi(\mathbb{M}^n)$  has dimension  $n$ . Take  $(\bar{a}, a_n) \in \varphi(\mathbb{M}^n)$  with  $\dim(\bar{a}, a_n/B) = n$ . Then  $a_n \notin \text{acl}(B\bar{a})$  and  $\dim(\bar{a}/B) = n - 1$ . As  $\varphi(\bar{a}, \mathbb{M})$  is  $B\bar{a}$ -definable and contains  $a_n \notin \text{acl}(B\bar{a})$ , it must be infinite, meaning that  $\bar{a} \in \psi(\mathbb{M}^{n-1})$ , and then  $\dim(\psi(\mathbb{M}^{n-1})) \geq \dim(\bar{a}/B) = n - 1$ .  $\square$

**Lemma 14.5.6.** *Let  $\varphi(x_1, \dots, x_n; y_1, \dots, y_m)$  be an  $\mathcal{L}$ -formula. The set*

$$\{\bar{b} \in \mathbb{M}^m : \dim(\varphi(\mathbb{M}^n, \bar{b})) = n\}$$

*is definable.*

*Proof.* The case  $n = 0$  is easy. If  $n > 0$ , let  $\psi(x_1, \dots, x_{n-1}; \bar{y})$  be the  $\mathcal{L}$ -formula equivalent to  $\exists^\infty x_n \varphi$ . Then for any  $\bar{b} \in \mathbb{M}^m$ ,

$$\dim(\varphi(\mathbb{M}^n, \bar{b})) = n \iff \dim(\psi(\mathbb{M}^{n-1}, \bar{b})) = n - 1,$$

by Lemma 14.5.5, and the right hand side is definable by induction on  $n$ .  $\square$

**Lemma 14.5.7.** *Let  $D \subseteq \mathbb{M}^n$  be definable and let  $k$  be in  $\{0, 1, \dots, n\}$ . Then  $\dim(D) \geq k$  if and only if there is a coordinate projection  $\pi : \mathbb{M}^n \rightarrow \mathbb{M}^k$  such that  $\pi(D) \subseteq \mathbb{M}^k$  has dimension  $k$ .*

*Proof.* If  $\dim(\pi(D)) = k$ , then  $\dim(D) \geq k$  by Theorem 14.5.3(6). Conversely, suppose  $\dim(D) \geq k$ . Take a small set  $B$  over which  $D$  is defined, and take  $\bar{a} \in D$  with  $\dim(\bar{a}/B) = \dim(D) \geq k$ . There are at least  $k$  values of  $i$  such that  $a_i \notin \text{acl}(Ba_1, \dots, a_{i-1})$ . Let  $i_1 < i_2 < \dots < i_k$  be some such values. Then

$$a_{i_j} \notin \text{acl}(Ba_{i_1} a_{i_2} \dots a_{i_{j-1}})$$

for each  $j$ , and so

$$\dim(a_{i_1} a_{i_2} \dots a_{i_k}/B) = k.$$

Let  $\pi : \mathbb{M}^n \rightarrow \mathbb{M}^k$  be the coordinate projection  $\pi(x_1, \dots, x_n) = (x_{i_1}, \dots, x_{i_k})$ . Then  $\pi(D)$  is  $B$ -definable,  $\pi(\bar{a}) \in \pi(D)$ , and  $\dim(\pi(\bar{a})/B) = k$ . We conclude that  $\dim(\pi(D)) \geq k$ .  $\square$

**Theorem 14.5.8** (Definability of dimension). *Let  $\varphi(x_1, \dots, x_n; y_1, \dots, y_m)$  be a formula. For each  $k \leq n$ , the set*

$$\{\bar{b} \in \mathbb{M}^m : \dim(\varphi(\mathbb{M}^n; \bar{b})) = k\}$$

*is definable.*

*Proof.* Let  $\Pi_k^n$  be the finite set of coordinate projections  $\mathbb{M}^n \rightarrow \mathbb{M}^k$ . For each  $\pi \in \Pi_k^n$ , let  $\theta_\pi(z_1, \dots, z_k, \bar{y})$  be the formula

$$\exists \bar{x} \varphi(\bar{x}; \bar{y}) \wedge (\pi(\bar{x}) = \bar{z}).$$

Then  $\theta_\pi(\mathbb{M}^k, \bar{b})$  is the image of  $\varphi(\mathbb{M}^n, \bar{b})$  under  $\pi$ . By Lemma 14.5.7,

$$D_k := \{\bar{b} \in \mathbb{M}^m : \dim(\varphi(\mathbb{M}^n; \bar{b})) \geq k\} = \bigcup_{\pi \in \Pi_k^n} \{\bar{b} \in \mathbb{M}^m : \dim(\theta_\pi(\mathbb{M}^k; \bar{b})) = k\},$$

and the sets on the right-hand side are definable by Lemma 14.5.6. Thus each  $D_k$  is definable. The set we want is  $D_k \setminus D_{k-1}$ .  $\square$

**Theorem 14.5.9.** *Let  $f : X \rightarrow Y$  be a definable function. For each  $b \in Y$ , let  $X_b = f^{-1}(b)$ . For each  $k \leq \dim(X)$ , let  $Y_k = \{b \in Y : \dim(X_b) = k\}$ . Then each set  $Y_k$  is definable, and*

$$\dim(X) = \max_k (k + \dim(Y_k)).$$

*Proof.* Theorems 14.5.8 and 14.5.4.  $\square$

## 14.6 Dimension theory in small models

If  $D$  is a definable set in a model  $M$ , and  $N$  is an elementary extension of  $M$ , then  $D(N)$  denotes  $\varphi(N)$ , where  $\varphi(\bar{x})$  is the  $\mathcal{L}(M)$ -formula defining  $D$  in  $M$ . The choice of  $\varphi$  doesn't matter, because

$$\begin{aligned} \varphi(M) = \psi(M) &\iff M \models \forall \bar{x} (\varphi(\bar{x}) \leftrightarrow \psi(\bar{x})) \\ \iff N \models \forall \bar{x} (\varphi(\bar{x}) \leftrightarrow \psi(\bar{x})) &\iff \varphi(N) = \psi(N). \end{aligned}$$

Note that  $D(M) = D$ .

**Lemma 14.6.1.** *Let  $M \preceq N$  be monster models of a pregeometric theory.*

1. *If  $\bar{a} \in M^n$  and  $B \subseteq M$  is small, then  $\dim^M(\bar{a}/B) = \dim^N(\bar{a}/B)$ .*
2. *If  $D$  is  $M$ -definable, then  $\dim(D(M)) = \dim(D(N))$ .*

*Proof.* 1. By Theorem 13.3.3,  $\text{acl}(Ba_1, \dots, a_{i-1})$  is the same whether calculated in  $M$  or  $N$ .

2. Take a small set  $B \subseteq M$  defining  $D$ . Take  $\bar{a} \in D(M)$  with  $\dim(\bar{a}/B) = \dim(D(M))$ . Then  $\bar{a} \in D(N)$  and  $\dim(\bar{a}/B) = \dim(D(M))$ , so

$$\dim(D(M)) \leq \dim(D(N)).$$

Conversely, take  $\bar{a} \in D(N)$  with  $\dim(\bar{a}/B) = \dim(D(N))$ . Then  $\text{tp}(\bar{a}/B)$  is realized in  $M$  by some  $\bar{c}$ . The formula defining  $D(N)$  is in  $\text{tp}(\bar{a}/B)$ , so  $\bar{c} \in D(N) \cap M^n = D(M)$ . Also,  $\dim(\bar{c}/B) = \dim(\bar{a}/B)$  by automorphism invariance. The fact that  $\bar{c} \in D(M)$  then shows

$$\dim(D(M)) \geq \dim(\bar{c}/B) = \dim(\bar{a}/B) = \dim(D(N)). \quad \square$$

**Definition 14.6.2.** Let  $M$  be a small model of a pregeometric theory. If  $D \subseteq M^n$  is definable, then  $\dim(D)$  is defined to be  $\dim(D(\mathbb{M}))$  where  $\mathbb{M}$  is a monster model extending  $M$ .

**Theorem 14.6.3.** *The choice of  $\mathbb{M}$  doesn't matter in Definition 14.6.2.*

*Proof.* Let  $\mathbb{M}_1, \mathbb{M}_2$  be two monster models extending  $M$ . Then  $\mathbb{M}_1$  and  $\mathbb{M}_2$  are elementarily equivalent as  $\mathcal{L}(M)$ -structures. By elementary amalgamation there is a third monster model  $\mathbb{M}_3$  extending  $\mathbb{M}_1$  and  $\mathbb{M}_2$ , up to isomorphism. Then

$$\dim(D(\mathbb{M}_1)) = \dim(D(\mathbb{M}_3)) = \dim(D(\mathbb{M}_2))$$

by two applications of Lemma 14.6.1.  $\square$

**Corollary 14.6.4.** *Dimension is invariant in elementary extensions: if  $M \preceq N$  and  $D$  is a definable set in  $M$ , then  $\dim(D(M)) = \dim(D(N))$ .*

*Proof.* Take a monster model  $\mathbb{M}$  extending  $N$  (and  $M$ ). Then  $\dim(D(M)) = \dim(D(\mathbb{M})) = \dim(D(N))$  by two applications of Definition 14.6.2.  $\square$

**Theorem 14.6.5.** *Let  $M$  be a model of a pregeometric theory  $T$ . Define  $\dim(-)$  as in Definition 14.6.2.*

1. *The basic properties of dimension (Theorem 14.5.3) hold in  $M$ .*
2. *If  $T$  is geometric, then the fiber dimension theorem (Theorem 14.5.4) and definability of dimension (Theorem 14.5.8) hold in  $M$ .*

*Proof.* 1. For example, we show  $\dim(D_1 \times D_2) = \dim(D_1) + \dim(D_2)$ . If  $\varphi(\bar{x})$  defines  $D_1$  and  $\psi(\bar{y})$  defines  $D_2$ , then the formula  $\theta(\bar{x}, \bar{y}) := \varphi(\bar{x}) \wedge \psi(\bar{y})$  defines  $D_1 \times D_2$ . Therefore

$$(D_1 \times D_2)(\mathbb{M}) = \theta(\mathbb{M}) = D_1(\mathbb{M}) \times D_2(\mathbb{M}).$$

Thus

$$\begin{aligned} \dim(D_1 \times D_2) &= \dim((D_1 \times D_2)(\mathbb{M})) \\ &= \dim(D_1(\mathbb{M}) \times D_2(\mathbb{M})) = \dim(D_1(\mathbb{M})) + \dim(D_2(\mathbb{M})) \\ &= \dim(D_1) + \dim(D_2). \end{aligned}$$

2. Suppose we have the configuration  $f : X \rightarrow Y$  of Theorem 14.5.4, where  $X_b = f^{-1}(b)$  has dimension  $k$  for every  $b \in Y$ . In order to reduce from  $M$  to the known case of  $\mathbb{M}$ , we need to show that the property

$$\dim(X_b) = k \text{ for every } b \in Y$$

transfers from  $M$  to  $\mathbb{M}$ . Let  $f_{\mathbb{M}} : X(\mathbb{M}) \rightarrow Y(\mathbb{M})$  be the function defined by the same formula as  $f$ , and for  $b \in Y(\mathbb{M})$  let  $X_b(\mathbb{M})$  denote the fiber of  $f_{\mathbb{M}}^{-1}(b)$ . Note that when  $b \in Y = Y(M)$ , the set  $X_b(\mathbb{M})$  really is the extension of  $X_b$ . By the definability of dimension (Theorem 14.5.8), the set

$$Z = \{b \in Y(\mathbb{M}) : \dim(X_b(\mathbb{M})) \neq k\}$$

is definable. It is  $\text{Aut}(\mathbb{M}/M)$ -invariant, hence  $M$ -definable (Theorem 11.3.8). If  $b \in Y(M)$ , then  $\dim(X_b(\mathbb{M})) = \dim(X_b) = k$ , so  $b \notin Z$ . Therefore there are no  $M$ -points in  $Z$ . By the Tarski-Vaught criterion,  $Z = \emptyset$ . Thus  $Z = \emptyset$ , and  $\dim(X_b(\mathbb{M})) = k$  for every  $b \in Y(\mathbb{M})$ . Then we can apply Theorem 14.5.4 over  $\mathbb{M}$  to get what we want.

For Theorem 14.5.8, fix a formula  $\varphi(\bar{x}, \bar{y})$ . The set

$$Z_k = \{\bar{b} \in \mathbb{M} : \dim(\varphi(\mathbb{M}; \bar{b})) = k\}$$

is definable by Theorem 14.5.8 applied to  $\mathbb{M}$ , and clearly  $\text{Aut}(\mathbb{M}/\emptyset)$ -invariant. By Theorem 11.3.8, it is 0-definable, defined by some formula  $\psi_k(\bar{y})$ . Then for any  $\bar{b}$  in  $M$ ,

$$\begin{aligned} \dim(\varphi(M, \bar{b})) = k &\iff \dim(\varphi(\mathbb{M}, \bar{b})) = k \\ &\iff \mathbb{M} \models \psi_k(\bar{b}) \iff M \models \psi_k(\bar{b}). \end{aligned}$$

The first equivalence holds by Definition 14.6.2. The second holds by choice of  $\psi_k$ . The third holds as  $M \preceq \mathbb{M}$ .

Therefore  $\{\bar{b} \in M^m : \dim(\varphi(M^n; \bar{b})) = k\}$  is the definable set  $\psi_k(M^m)$ .

□





# Appendix A

## Notation and conventions

This appendix reviews the mathematical notation used in this book, which is mostly standard. In the process, we will also review a little bit of the mathematical background assumed throughout the book.

If  $\varphi, \psi$  are statements, then  $\varphi \implies \psi$  means that  $\varphi$  logically implies  $\psi$ . In other words,

$$\varphi \implies \psi$$

means

If  $\varphi$  then  $\psi$ .

The notation  $\varphi \iff \psi$  means that  $\varphi \implies \psi$  and  $\psi \implies \varphi$ . We say that  $\varphi$  holds “if and only if”  $\psi$  holds. “If and only if” is often abbreviated to iff. The notation  $\forall x (\dots)$  means “for every  $x$ ,  $\dots$ ” Similarly, the notation  $\exists x (\dots)$  means “there is an  $x$  such that  $\dots$ ” If we are only interested in values of  $x$  coming from a set  $A$ , we write  $\forall x \in A$  or  $\exists x \in A$ . In other words, these things

$$\forall x \in A (\dots)$$

$$\exists x \in A (\dots)$$

mean the following, respectively:

For every  $x$  in  $A$ ,  $\dots$

There is an  $x$  in  $A$  such that  $\dots$

If  $A$  is a set and  $x$  is a value, then  $x \in A$  means that  $x$  is an element of  $A$ , and  $x \notin A$  means that  $x$  is not an element of  $A$ . If  $A$  and  $B$  are sets,

then  $A \subseteq B$  means that  $A$  is a subset of  $B$ , i.e., every element of  $A$  is an element of  $B$ . It is an axiom of set theory that  $A = B$  if and only if  $A \subseteq B$  and  $B \subseteq A$ . The notation  $A \subsetneq B$  means that  $A$  is a proper subset of  $B$ :  $A \subseteq B$  and  $A \neq B$ . We avoid the ambiguous notation  $A \subset B$ . Remember to distinguish  $A \subsetneq B$  ( $A$  is a proper subset of  $B$ ) from  $A \not\subseteq B$  ( $A$  is not a subset of  $B$ ). We write  $A \subseteq_f B$  to mean that  $A$  is a finite subset of  $B$ . (This relation comes up frequently in model theory.)

We write the empty set as  $\emptyset$ ; this is the unique set with no elements. If  $A$  is a set, then  $\mathfrak{P}(A)$  denotes the powerset of  $A$ , the set of all subsets of  $A$ :

$$\mathfrak{P}(A) = \{X : X \subseteq A\}.$$

Here, the set-builder notation  $\{x : \dots\}$  means “the set of  $x$  such that  $\dots$ ”. Similarly,  $\{x \in A : \dots\}$  means “the set of  $x$  in  $A$  such that  $\dots$ ” and  $\{f(x) : x \in A, \dots\}$  means “the set of values  $f(x)$  where  $x$  is in  $A$  and  $\dots$ ”.

If  $A$  and  $B$  are sets, then  $A \cap B$  and  $A \cup B$  denote the intersection and union of  $A$  and  $B$ :

$$\begin{aligned} A \cap B &= \{x : x \in A \text{ and } x \in B\} \\ A \cup B &= \{x : x \in A \text{ or } x \in B\}. \end{aligned}$$

If  $A_1, \dots, A_n$  are sets, then we write their intersection and union as follows:

$$\begin{aligned} \bigcap_{i=1}^n A_i &= A_1 \cap A_2 \cap A_3 \cap \dots \cap A_n \\ \bigcup_{i=1}^n A_i &= A_1 \cup A_2 \cup A_3 \cup \dots \cup A_n. \end{aligned}$$

Additionally, we write the “set difference” of  $A$  and  $B$  as  $A \setminus B$ :

$$A \setminus B = \{x : x \in A \text{ and } x \notin B\}.$$

This should be distinguished from the “symmetric difference”  $(A \setminus B) \cup (B \setminus A)$ , which we will not use.

Two sets  $A$  and  $B$  are *disjoint* if  $A \cap B = \emptyset$ . A list of sets  $A_1, \dots, A_n$  is *pairwise disjoint* if any two of them are disjoint:  $A_i \cap A_j = \emptyset$  for any  $i \neq j$ . A list of values  $x_1, \dots, x_n$  is *pairwise distinct* if  $x_i \neq x_j$  for  $i \neq j$ .

We use the following blackboard bold symbols.  $\mathbb{N}$  denotes the natural numbers  $\{0, 1, 2, \dots\}$ . Note that we include 0 in  $\mathbb{N}$ . We sometimes denote

$\mathbb{N}$  as  $\omega$ . Notation like  $n < \omega$  and  $n \in \omega$  means that  $n$  is a natural number. The reason for this notation comes from set theory, where  $\omega$  is another name for  $\mathbb{N}$  and the first ordinal number after the natural numbers.  $\mathbb{Z}$  denotes the integers  $\{\dots, -2, -1, 0, 1, 2, \dots\}$ .  $\mathbb{Q}$  denotes the set of rational numbers, values of the form  $x/y$  where  $x, y \in \mathbb{Z}$ .  $\mathbb{R}$  denotes the usual real numbers.  $\mathbb{C}$  denotes the complex numbers, values of the form  $x + iy$  where  $x, y \in \mathbb{R}$  and  $i = \sqrt{-1}$ . To summarize,

$$\mathbb{N} \subsetneq \mathbb{Z} \subsetneq \mathbb{Q} \subsetneq \mathbb{R} \subsetneq \mathbb{C}.$$

An *n-tuple* is a list of  $n$  values  $(x_1, x_2, x_3, \dots, x_n)$ . 2-tuples are called *pairs* or *ordered pairs*, and 3-tuples are called *triples*. Two  $n$ -tuples

$$\begin{aligned} &(x_1, \dots, x_n) \\ &(y_1, \dots, y_n) \end{aligned}$$

are equal if and only if  $x_i = y_i$  for all  $i$  between 1 and  $n$  (inclusive). For example,  $(x_1, x_2) = (y_1, y_2)$  if and only if  $x_1 = y_1$  and  $x_2 = y_2$ .

If  $A$  and  $B$  are sets, the *direct product* or *cartesian product*, written  $A \times B$ , is the set of ordered pairs  $(x, y)$  where  $x \in A$  and  $y \in B$ :

$$A \times B = \{(x, y) : x \in A \text{ and } y \in B\}.$$

For example,

$$\begin{aligned} \{1, 2, 3\} \times \{\text{up}, \text{down}\} = &\{(1, \text{up}), (1, \text{down}), (2, \text{up}), \\ &(2, \text{down}), (3, \text{up}), (3, \text{down})\}. \end{aligned}$$

A *relation* between  $A$  and  $B$  is a subset  $R \subseteq A \times B$ . If  $x \in A$  and  $y \in B$ , then  $x R y$  means  $(x, y) \in R$ .

TODO: partial orders and linear orders.

A *function* or *map* from  $A$  to  $B$  is a relation  $f \subseteq A \times B$  such that for every  $x \in A$ , there is exactly one  $y \in B$  such that  $x f y$ . We write this unique value of  $y$  as  $f(x)$ , so that

$$f(x) = y \iff (x f y) \iff (x, y) \in f.$$

The notation  $x \mapsto y$  means that  $f(x) = y$  for whatever function  $f$  we are talking about. Note the bar at the left end of  $\mapsto$ . The notation  $f : A \rightarrow B$  means that  $A$  and  $B$  are sets and  $f$  is a function from  $A$  to  $B$ . The sets  $A$

and  $B$  are called the *domain* and *codomain* of  $f$ . The domain of  $f$  is written  $\text{dom}(f)$ . The *range* or *image* of  $f$  is

$$\text{im}(f) = \{f(x) : x \in A\} = \{y \in B : \text{there is } x \in A \text{ such that } f(x) = y\}.$$

If  $X$  is a subset of  $A$ , then the *image* or *direct image* of  $X$  is the set

$$f(X) = \{f(x) : x \in X\} = \{y \in B : \text{there is } x \in X \text{ such that } f(x) = y\}.$$

If  $Y$  is a subset of  $B$ , then the *preimage* or *inverse image* of  $Y$  is the set

$$f^{-1}(Y) = \{x \in A : f(x) \in Y\}.$$

If  $f$  is a function from  $A$  to  $B$ , then  $f$  is a . . .

1. *surjection* if for every  $y \in B$ , there is at least one  $x \in A$  with  $f(x) = y$ .
2. *injection* if for every  $y \in B$ , there is at most one  $x \in A$  with  $f(x) = y$ .
3. *bijection* if for every  $y \in B$ , there is exactly one  $x \in A$  with  $f(x) = y$ .

We say that  $f$  is *surjective*, *injective*, or *bijective* if  $f$  is a surjection, injection, or bijection. Note the following:

- $f$  is a surjection if and only if  $\text{im}(f) = B$ .
- $f$  is an injection if and only if

$$f(x) = f(y) \implies x = y$$

if and only if

$$x \neq y \implies f(x) \neq f(y).$$

- $f$  is a bijection if and only if  $f$  is injective and surjective.

If  $f : A \rightarrow B$  is a bijection, then there is an *inverse function*  $f^{-1} : B \rightarrow A$  defined by

$$f^{-1}(y) = x \iff f(x) = y.$$

If  $f : A \rightarrow B$  and  $g : B \rightarrow C$  are functions, then the *composition* is the function  $h : A \rightarrow C$  defined by

$$h(x) = f(g(x)).$$

The composition is usually written  $f \circ g$ . Note that  $f \circ g$  is the function where  $g$  is applied first and  $f$  is applied second.

If  $A$  is a set, then  $\text{id}_A$  or  $\text{id}$  denotes the *identity function* on  $A$ , the function

$$\begin{aligned}\text{id}_A : A &\rightarrow A \\ \text{id}_A(x) &= x.\end{aligned}$$

Note that if  $f : A \rightarrow B$  is a function then

$$\begin{aligned}f \circ \text{id}_A &= f \\ \text{id}_B \circ f &= f.\end{aligned}$$

If  $f : A \rightarrow B$  is a bijection and  $f^{-1} : B \rightarrow A$  is the inverse, then

$$\begin{aligned}f^{-1} \circ f &= \text{id}_A \\ f \circ f^{-1} &= \text{id}_B.\end{aligned}$$

If  $A, B, C$  are sets, then a function  $f : A \times B \rightarrow C$  is a function from the cartesian product  $A \times B$  to  $C$ . If  $x \in A$  and  $y \in B$ , we write  $f((x, y))$  as  $f(x, y)$ , and we think of  $f$  as a function which takes one input  $x$  from  $A$  and one input  $y$  from  $B$ , and then outputs a value  $f(x, y)$  in  $C$ .

If  $A$  is a set, then  $A^2, A^3, \dots$  denote the cartesian product of  $A$  with itself  $n$  times:

$$\begin{aligned}A^2 &= A \times A \\ A^3 &= A \times A \times A \\ &\dots \\ A^n &= \underbrace{A \times A \times \dots \times A}_{n \text{ times}}.\end{aligned}$$

So  $A^n$  is the set of  $n$ -tuples  $(x_1, \dots, x_n)$  such that  $x_1, x_2, \dots, x_n$  are all in  $A$ . A function  $f : A^n \rightarrow B$  can be thought of as a function which takes  $n$  inputs  $x_1, \dots, x_n$  from  $A$  and outputs a value in  $B$ . We think of operations like  $+$  (addition) and  $\cdot$  (multiplication) as functions  $\mathbb{R}^2 \rightarrow \mathbb{R}$ , taking two inputs from  $\mathbb{R}$  and outputting one value in  $\mathbb{R}$ . But we usually write  $x + y$  rather than  $+(x, y)$ .

If  $X$  is a set, then  $|X|$  denotes the *size* or *cardinality* of  $X$ , the number of elements of  $X$ . For example,  $|\emptyset| = 0$ . When  $X$  is finite,  $|X|$  is an element

of  $\mathbb{N}$ . Otherwise,  $|X|$  is one of Cantor's *infinite cardinal numbers*, of which the first few are  $\aleph_0, \aleph_1, \aleph_2, \dots$ . It is worth knowing that

$$\begin{aligned} |\mathbb{N}| &= |\mathbb{Z}| = |\mathbb{Q}| = \aleph_0 \\ |\mathbb{R}| &= |\mathbb{C}| = 2^{\aleph_0} > \aleph_0. \end{aligned}$$

The axioms of set theory neither prove nor disprove the *continuum hypothesis*, the statement that  $2^{\aleph_0} = \aleph_1$ . If  $X$  and  $Y$  are sets, then

$$\begin{aligned} |X \times Y| &= |X| \cdot |Y| \\ |X \cup Y| &\leq |X| + |Y| \\ |X \cup Y| &= |X| + |Y| \text{ if } X \cap Y = \emptyset \\ |\mathfrak{P}(X)| &= 2^{|X|} > |X| \\ X \subseteq Y &\implies |X| \leq |Y|. \end{aligned}$$

Moreover,  $|X| = |Y|$  if and only if there is a bijection  $f : X \rightarrow Y$ , and  $|X| \leq |Y|$  if and only if there is an injection  $f : X \rightarrow Y$ . When  $X$  and  $Y$  are non-empty,  $|X| \geq |Y|$  if and only if there is a surjection  $X \rightarrow Y$ .

If  $X$  is a proper subset of  $Y$ , then  $|X| < |Y|$ , but it can happen that  $|X| = |Y|$  when  $X$  and  $Y$  are infinite. For example,  $\mathbb{N} \subsetneq \mathbb{Z} \subsetneq \mathbb{Q}$ , but  $|\mathbb{N}| = |\mathbb{Z}| = |\mathbb{Q}| = \aleph_0$ .

It is worth knowing that if  $\kappa, \lambda$  are infinite cardinals, then  $\kappa \cdot \lambda = \kappa + \lambda = \max(\kappa, \lambda)$ .

The notation  $|X| < \infty$  means that the set  $X$  is finite, and  $|X| = \infty$  means that  $X$  is infinite. This is an abuse of notation—there is no cardinal number “ $\infty$ .” In fact,

$$\begin{aligned} |X| < \infty &\iff |X| < \aleph_0 \\ |X| = \infty &\iff |X| \geq \aleph_0. \end{aligned}$$

In set theory,  $\aleph_0$ ,  $\mathbb{N}$ , and  $\omega$  are all the same thing, so we sometimes write  $\aleph_0$  as  $\omega$ .