

Orbifold Stiefel-Whitney classes on right-angled Coxeter complexes

Lisu Wu

School of Mathematical Sciences, Fudan University


Toric Topology 2021 in Osaka (Online)

March 25, 2021, Osaka City University, Osaka

1. Right-angled Coxeter complexes and its homology groups
2. Orbifold Stiefel-Whitney classes on RACC

- An n -orbifold is a singular space locally modelled on the quotient of an open set of \mathbb{R}^n by a finite group action.

- An n -orbifold is a singular space locally modelled on the quotient of an open set of \mathbb{R}^n by a finite group action.



♣ A right-angled Coxeter n -orbifold (RACO) is an orbifold locally modelled on $\mathbb{R}^n/(\mathbb{Z}_2)^k$, where $(\mathbb{Z}_2)^k$ -action is generated by standard reflections (across the coordinate hyperplanes).

♣ A right-angled Coxeter n -orbifold (RACO) is an orbifold locally modelled on $\mathbb{R}^n/(\mathbb{Z}_2)^k$, where $(\mathbb{Z}_2)^k$ -action is generated by standard reflections (across the coordinate hyperplanes).

♣ Right-angled Coxeter cell: $e^n/(\mathbb{Z}_2)^k$.

If $k > 0$, $e^n/(\mathbb{Z}_2)^k$ is singular cell, otherwise, regular cell.

♣ A right-angled Coxeter n -orbifold (RACO) is an orbifold locally modelled on $\mathbb{R}^n/(\mathbb{Z}_2)^k$, where $(\mathbb{Z}_2)^k$ -action is generated by standard reflections (across the coordinate hyperplanes).

♣ Right-angled Coxeter cell: $e^n/(\mathbb{Z}_2)^k$.

If $k > 0$, $e^n/(\mathbb{Z}_2)^k$ is singular cell, otherwise, regular cell.

♣ Right-angled Coxeter complexes(RACC): Every attaching map

$$\phi : \partial \overline{e^n}/(\mathbb{Z}_2)^k \rightarrow X^{n-1}$$

preserves the local groups.

♣ A right-angled Coxeter n -orbifold (RACO) is an orbifold locally modelled on $\mathbb{R}^n/(\mathbb{Z}_2)^k$, where $(\mathbb{Z}_2)^k$ -action is generated by standard reflections (across the coordinate hyperplanes).

♣ Right-angled Coxeter cell: $e^n/(\mathbb{Z}_2)^k$.

If $k > 0$, $e^n/(\mathbb{Z}_2)^k$ is singular cell, otherwise, regular cell.

♣ Right-angled Coxeter complexes(RACC): Every attaching map


$$\phi : \partial \overline{e^n}/(\mathbb{Z}_2)^k \rightarrow X^{n-1}$$

preserves the local groups.


An example




$$\mathbb{e}^2/\mathbb{Z}_2$$




A




B



E



C




D


An example




e^2/\mathbb{Z}_2




A




B



E








C








D

(*) E is not a RACC.

Blow-up of RACC



Blow-up of RACC



$$(*) \quad \tilde{X}^2 = Y/[p, q] \sim [p, q'] \simeq T^2.$$

Boundary maps of RACC are defined by boundary maps of its blow-up:

$$d(e^n/W) = \sum n_\beta \left(\frac{|W|}{|W_\beta|} \bmod 2 \right) e_\beta^{n-1}/W_\beta$$

where W_β is a subgroup of W .

Boundary maps of RACC are defined by boundary maps of its blow-up:

$$d(e^n/W) = \sum n_\beta \left(\frac{|W|}{|W_\beta|} \bmod 2 \right) e_\beta^{n-1}/W_\beta$$

where W_β is a subgroup of W .

Theorem (Lü-Wu-Yu)

Let X be an n -dimensional right-angled Coxeter complex, and T be the face set of X . Then

$$H_i^{\text{orb}}(X) = \bigoplus_{f \in T} H_{i-l(f)}(X_f) \quad (1)$$

where $l(f)$ is the local codimension of f .

The cup product of RACC is defined by its blow-up,

$$H_{orb}^*(X) := H^*(\tilde{X}).$$

The cup product of RACC is defined by its blow-up,

$$H_{orb}^*(X) := H^*(\tilde{X}).$$

Example (Simple polytope)

Let P be a simple polytope equipped with a right-angled Coxeter orbifold structure. Then the standard cubical decomposition of P is a RACC. Then

$$H_i^{orb}(P) = \mathbb{Z}^{f_{n-i}} \quad (2)$$

where f_{n-i} is the number of $(n-i)$ -faces of P .



$$H_{orb}^*(P; \mathbb{Z}_2) = \mathbb{Z}_2[v_1, \dots, v_m] / I + J$$

where I is the Stanley-Reisner ideal of P , $J = (v_i^2, \forall i)$.

E, X be two orbifolds with orbifold structures $\{U^*, \psi^*, G^*\}$ and $\{U, \psi, G\}$, an orbifold vector bundle $\pi : E \rightarrow X$ satisfies:



$$\begin{array}{ccc}
 E & \begin{array}{c} E_{U/G} \xleftarrow{\psi_{U^*}} U^* \cong U \times \mathbb{R}^m \\ \downarrow \pi_H \qquad \downarrow \tilde{\pi} \\ U/G \xleftarrow{\psi_U} U \end{array} & \begin{array}{c} \curvearrowright G^* \\ \\ \curvearrowright G \end{array}
 \end{array}$$

- Compatibility conditions.


 \bar{E}

 \tilde{E}

$$\bar{E} = D^1 \times \mathbb{R} / (x, y) \sim (-x, y)$$



$$\tilde{E} = D^1 \times \mathbb{R} / (x, y) \sim (-x, -y)$$


 \bar{E}

 \tilde{E}

$$\bar{E} = D^1 \times \mathbb{R} / (x, y) \sim (-x, y)$$

$$\tilde{E} = D^1 \times \mathbb{R} / (x, y) \sim (-x, -y)$$

$$H^*(D^1/\mathbb{Z}_2; \mathbb{Z}_2) = \mathbb{Z}_2[s]/(s^2)$$




 \bar{E}

 \tilde{E}

$$\bar{E} = D^1 \times \mathbb{R} / (x, y) \sim (-x, y)$$

$$\tilde{E} = D^1 \times \mathbb{R} / (x, y) \sim (-x, -y)$$

$$H^*(D^1/\mathbb{Z}_2; \mathbb{Z}_2) = \mathbb{Z}_2[s] / (s^2)$$

$$w(\bar{E}) := 1 \quad w(\tilde{E}) := 1 + s$$


 \bar{E}

 \tilde{E}

$$\bar{E} = D^1 \times \mathbb{R} / (x, y) \sim (-x, y)$$

$$\tilde{E} = D^1 \times \mathbb{R} / (x, y) \sim (-x, -y)$$

$$H^*(D^1/\mathbb{Z}_2; \mathbb{Z}_2) = \mathbb{Z}_2[s] / (s^2)$$

$$w(\bar{E}) := 1 \quad w(\tilde{E}) := 1 + s$$

(Orbifold vector bundles over a RACC maybe not a RACC.)

In general, for $\pi : E \rightarrow D^n/(\mathbb{Z}_2)^k$, there is a representation

$$\rho : (\mathbb{Z}_2)^k \longrightarrow GL_m(\mathbb{R}).$$

The image of ρ on the generator set of $(\mathbb{Z}_2)^k$ gives a $m \times k$ matrix with elements ± 1 .

$$C = \begin{pmatrix} x_1^1 & x_1^2 & \cdots & x_1^k \\ x_2^1 & x_2^2 & \cdots & x_2^k \\ \vdots & \vdots & \ddots & \vdots \\ x_m^1 & x_m^2 & \cdots & x_m^k \end{pmatrix}_{m \times k}$$

In general, for $\pi : E \rightarrow D^n/(\mathbb{Z}_2)^k$, there is a representation

$$\rho : (\mathbb{Z}_2)^k \longrightarrow GL_m(\mathbb{R}).$$

The image of ρ on the generator set of $(\mathbb{Z}_2)^k$ gives a $m \times k$ matrix with elements ± 1 .

$$C = \begin{pmatrix} x_1^1 & x_1^2 & \cdots & x_1^k \\ x_2^1 & x_2^2 & \cdots & x_2^k \\ \vdots & \vdots & \ddots & \vdots \\ x_m^1 & x_m^2 & \cdots & x_m^k \end{pmatrix}_{m \times k}$$

$$H^*(D^n/(\mathbb{Z}_2)^k; \mathbb{Z}_2) \cong \mathbb{Z}_2[s_1, \dots, s_k]/(s_i^2, \forall i).$$

Orbifold SW classes on $D^n/(\mathbb{Z}_2)^k$

In general, for $\pi : E \rightarrow D^n/(\mathbb{Z}_2)^k$, there is a representation

$$\rho : (\mathbb{Z}_2)^k \longrightarrow GL_m(\mathbb{R}).$$

The image of ρ on the generator set of $(\mathbb{Z}_2)^k$ gives a $m \times k$ matrix with elements ± 1 .


$$C = \begin{pmatrix} x_1^1 & x_1^2 & \cdots & x_1^k \\ x_2^1 & x_2^2 & \cdots & x_2^k \\ \vdots & \vdots & \ddots & \vdots \\ x_m^1 & x_m^2 & \cdots & x_m^k \end{pmatrix}_{m \times k}$$

$$H^*(D^n/(\mathbb{Z}_2)^k; \mathbb{Z}_2) \cong \mathbb{Z}_2[s_1, \dots, s_k]/(s_i^2, \forall i).$$

The total orbifold Stiefel-Whitney class of $\pi : E \rightarrow D^n/(\mathbb{Z}_2)^k$ is defined:

$$w(E) = \prod_{i=1}^m \left(1 + \sum_{j=1}^k \frac{1 - x_i^j}{2} s_j \right) \in H^*(D^n/(\mathbb{Z}_2)^k; \mathbb{Z}_2). \quad (3)$$

Orbifold SW classes on RACC– An example



$$H^i(X; \mathbb{Z}_2) = \begin{cases} \mathbb{Z}_2, & i = 0 \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2, & i = 1 \\ \mathbb{Z}_2, & i = 2 \\ 0, & \text{otherwise.} \end{cases}$$

Figure: $X = S^1 \times [-1, 1]/\mathbb{Z}_2$

All regular cells in a RACC X give a subcomplex of X , denoted by X_{reg} , then

$$X/X_{reg} = \bigvee_H H.$$

All regular cells in a RACC X give a subcomplex of X , denoted by X_{reg} , then

$$X/X_{reg} = \bigvee_H H.$$

For each H , we have a

$$\mathcal{R}_H = \mathbb{Z}_2[s_1, \dots, s_n]/(I_H + J_H) \quad (4)$$

where I_H is the Stanley-Reisner ideal determined by H , $J_H = (s_i^2, \forall i)$.

All regular cells in a RACC X give a subcomplex of X , denoted by X_{reg} , then

$$X/X_{reg} = \bigvee_H H.$$

For each H , we have a

$$\mathcal{R}_H = \mathbb{Z}_2[s_1, \dots, s_n]/(I_H + J_H) \quad (4)$$


where I_H is the Stanley-Reisner ideal determined by H , $J_H = (s_i^2, \forall i)$.

$$\mathcal{R}_H < H^*(H; \mathbb{Z}_2) < H^*(X; \mathbb{Z}_2)$$

$$\pi : E \longrightarrow X \Rightarrow \pi_H : E_H \longrightarrow H \Rightarrow \text{matrix } C_H \Rightarrow \tilde{\pi} : E_{C_H} \longrightarrow D^n/(\mathbb{Z}_2)^n.$$


Orbifold SW classes on RACC

$$\pi : E \longrightarrow X \Rightarrow \pi_H : E_H \longrightarrow H \Rightarrow \text{matrix } C_H \Rightarrow \tilde{\pi} : E_{C_H} \longrightarrow D^n/(\mathbb{Z}_2)^n.$$



Orbifold SW classes on RACC

$$\pi : E \longrightarrow X \Rightarrow \pi_H : E_H \longrightarrow H \Rightarrow \text{matrix } C_H \Rightarrow \tilde{\pi} : E_{C_H} \longrightarrow D^n/(\mathbb{Z}_2)^\eta.$$



$$\begin{array}{ccc} E_H & \xrightarrow{\quad} & E_{C_H} \\ \pi_H \downarrow & & \downarrow \tilde{\pi} \\ H & \xrightarrow{j} & D^n/(\mathbb{Z}_2)^\eta \end{array}$$

In general, let $j : \mathcal{N}(H) \rightarrow \mathcal{N}(D^\eta/(\mathbb{Z}_2)^\eta) = \Delta^{\eta-1}$ be a simplicial map.

$$j^* : \mathbb{Z}_2[s_1, \dots, s_\eta]/(s_i^2, \forall i) \longrightarrow \mathcal{R}_H \subset H^*(X; \mathbb{Z}_2).$$

In general, let $j : \mathcal{N}(H) \rightarrow \mathcal{N}(D^\eta/(\mathbb{Z}_2)^\eta) = \Delta^{\eta-1}$ be a simplicial map.

$$j^* : \mathbb{Z}_2[s_1, \dots, s_\eta]/(s_i^2, \forall i) \longrightarrow \mathcal{R}_H \subset H^*(X; \mathbb{Z}_2).$$

Total orbifold SW class of $\pi_H : E_H \longrightarrow X_H$ is defined:

$$w(E_H) = j^*(w(E_{C_H})) \in \mathcal{R}_H \subset H^*(X; \mathbb{Z}_2) \quad (5)$$

In general, let $j : \mathcal{N}(H) \rightarrow \mathcal{N}(D^\eta/(\mathbb{Z}_2)^\eta) = \Delta^{\eta-1}$ be a simplicial map.

$$j^* : \mathbb{Z}_2[s_1, \dots, s_\eta]/(s_i^2, \forall i) \longrightarrow \mathcal{R}_H < H^*(X; \mathbb{Z}_2).$$

Total orbifold SW class of $\pi_H : E_H \longrightarrow X_H$ is defined:


$$w(E_H) = j^*(w(E_{C_H})) \in \mathcal{R}_H < H^*(X; \mathbb{Z}_2) \quad (5)$$

Total orbifold SW class of $\pi : E \longrightarrow X$ is defined:

$$w(E) = w(E(X_{reg})) \cdot \prod_H w(E_H). \quad (6)$$

where $E(X_{reg}) = \pi|_{X_{reg}}$.

Example



$$H^i(X; \mathbb{Z}_2) = \begin{cases} \mathbb{Z}_2, & i = 0, 2 \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2, & i = 1 \\ 0, & \text{otherwise.} \end{cases}$$

$$w(TX) = w(TS^1)(1 + s) = 1 + s.$$

Figure: $X = S^1 \times [-1, 1]/\mathbb{Z}_2$

Thank You

Wu, Li-Su

Email: wulisuwulisu@qq.com

School of Mathematical Sciences, Fudan University