

Fundamental groups of small covers

Wu, Lisu

School of Mathematical Sciences, Fudan University

The 5th Korea Toric Topology Winter Workshop

Gyeongju, Korea, January 21-23, 2019

1. Introduction
2. Presentations of Fundamental Groups
3. Main Results and Applications

- An n -dimensional small cover is a closed n -manifold M with a locally standard \mathbb{Z}_2^n -action whose orbit space is a simple polytope P .

$$\pi : M \longrightarrow P$$

- An n -dimensional small cover is a closed n -manifold M with a locally standard \mathbb{Z}_2^n -action whose orbit space is a simple polytope P .

$$\pi : M \longrightarrow P$$

- The \mathbb{Z}_2^n -action on M determines a \mathbb{Z}_2^n -valued characteristic function λ on the set of facets of P

$$\lambda : \{F_1, F_2, \dots, F_m\} \longrightarrow \mathbb{Z}_2^n$$

- An n -dimensional small cover is a closed n -manifold M with a locally standard \mathbb{Z}_2^n -action whose orbit space is a simple polytope P .

$$\pi : M \longrightarrow P$$

- The \mathbb{Z}_2^n -action on M determines a \mathbb{Z}_2^n -valued characteristic function λ on the set of facets of P

$$\lambda : \{F_1, F_2, \dots, F_m\} \longrightarrow \mathbb{Z}_2^n$$

such that

$$\forall f = F_1 \cap F_2 \cap \dots \cap F_k,$$

$$G_f \triangleq \langle \lambda(F_1), \lambda(F_2), \dots, \lambda(F_k) \rangle \cong \mathbb{Z}_2^k.$$

- An n -dimensional small cover is a closed n -manifold M with a locally standard \mathbb{Z}_2^n -action whose orbit space is a simple polytope P .

$$\pi : M \longrightarrow P$$

- The \mathbb{Z}_2^n -action on M determines a \mathbb{Z}_2^n -valued characteristic function λ on the set of facets of P

$$\lambda : \{F_1, F_2, \dots, F_m\} \longrightarrow \mathbb{Z}_2^n$$

such that

$$\forall f = F_1 \cap F_2 \cap \dots \cap F_k,$$

$$G_f \triangleq \langle \lambda(F_1), \lambda(F_2), \dots, \lambda(F_k) \rangle \cong \mathbb{Z}_2^k.$$

$$\text{Rk: } \mathcal{F}(P) \triangleq \{F_1, F_2, \dots, F_m\}.$$

- Small cover

$$M = P \times \mathbb{Z}_2^n / \sim$$

where $(p, g) \sim (q, h)$ iff $p = q$, $g^{-1}h \in G_{f(p)}$, and $f(p)$ is the unique face of P that contains p in its relative interior, $G_{f(p)} = \{1\}$ if $p \in P^\circ$.

- Small cover

$$M = P \times \mathbb{Z}_2^n / \sim$$

where $(p, g) \sim (q, h)$ iff $p = q$, $g^{-1}h \in G_{f(p)}$, and $f(p)$ is the unique face of P that contains p in its relative interior, $G_{f(p)} = \{1\}$ if $p \in P^\circ$.

- Define right-angled Coxeter group of P

$$W = \langle s_F, \forall F \in \mathcal{F}(P) \mid s_F^2 = 1; (s_F s_{F'})^2 = 1, F \cap F' \neq \emptyset \rangle$$

- The Borel construction of \mathbb{Z}_2^n on M

$$BP = M \times_{\mathbb{Z}_2^n} E\mathbb{Z}_2^n$$

where $E\mathbb{Z}_2^n = (S^\infty)^n$. And $\pi_1(BP) \cong W$.

- The Borel construction of \mathbb{Z}_2^n on M

$$BP = M \times_{\mathbb{Z}_2^n} E\mathbb{Z}_2^n$$



where $E\mathbb{Z}_2^n = (S^\infty)^n$. And $\pi_1(BP) \cong W$.

- $M \rightarrow BP \rightarrow B\mathbb{Z}_2^n$ induces a right-split exact sequence


$$1 \longrightarrow \pi_1(M) \longrightarrow W \begin{array}{c} \xrightarrow{\phi} \\ \xleftarrow{\gamma} \end{array} \mathbb{Z}_2^n \longrightarrow 1$$

where $\phi(s_F) = \lambda(F)$, $\forall F \in \mathcal{F}(P)$.

Cell decomposition



Generators and relations




Cell-①

Relation-1: $x_{F,g}x_{F,g\lambda(F)} = 1$

Relation-2: $x_{F,g}x_{F',g\lambda(F')} = x_{F',g}x_{F,g\lambda(F')}$

Generators and relations



Relation-2: $x_{F,g}x_{F',g\lambda}(F) = x_{F',g}x_{F,g\lambda}(F')$


Relation-3: $x_{F,g} = 1, p_0 \subset F$

Presentation of $\pi_1(M, p_0)$

$$\begin{aligned}\pi_1(M, p_0) = \langle x_{F,g}, \forall F, g \mid & x_{F,g} x_{F,g\lambda(F)} = 1; \\ & x_{F,g} x_{F',g\lambda(F)} = x_{F',g} x_{F,g\lambda(F')}, F \cap F' \neq \emptyset; \\ & x_{F,g} = 1, p_0 \in F \rangle\end{aligned}$$

Relation between $\pi_1(M)$ and W


$$\tilde{M} = Q \times \pi_1(M) / \sim = P \times W / \sim$$



Rk: $\lambda(F_1, F_2, F, F') = (e_1, e_2, e_1 e_2, e_2).$

Relation between $\pi_1(M)$ and W

$$\tilde{M} = Q \times \pi_1(M) / \sim = P \times W / \sim$$




$$x_{F,1}(Q, 1) \mapsto (Q, x_{F,1})$$

Rk: $\lambda(F_1, F_2, F, F') = (e_1, e_2, e_1 e_2, e_2).$

Relation between $\pi_1(M)$ and W

$$\tilde{M} = Q \times \pi_1(M) / \sim = P \times W / \sim$$




$$x_{F,1}(Q, 1) \mapsto (Q, x_{F,1})$$

$$s_F(P, 1) \mapsto (P, s_F)$$

Rk: $\lambda(F_1, F_2, F, F') = (e_1, e_2, e_1 e_2, e_2).$

Relation between $\pi_1(M)$ and W

$$\tilde{M} = Q \times \pi_1(M) / \sim = P \times W / \sim$$



$$x_{F,1}(Q, 1) \mapsto (Q, x_{F,1})$$

$$s_F(P, 1) \mapsto (P, s_F)$$

$$x_{F,1}(P, 1) \mapsto (P, s_{F_2} s_{F_1} s_F)$$

$$\begin{array}{ccc} & & \uparrow \\ & \downarrow & \\ \gamma(\lambda(F)) \cdot s_F(P, 1) & \mapsto & s_{F_2} s_{F_1} s_F(P, 1) \end{array}$$

Rk: $\lambda(F_1, F_2, F, F') = (e_1, e_2, e_1 e_2, e_2).$

$$\begin{aligned}\alpha : \pi_1(M, p_0) &\longrightarrow W \\ x_{F,g} &\longmapsto \gamma(g\lambda(F)) \cdot \gamma(\lambda(F))s_F \cdot (\gamma(g\lambda(F)))^{-1} \\ &= \gamma(g)s_F\gamma(g\lambda(F))\end{aligned}$$

$$\alpha : \pi_1(M, p_0) \longrightarrow W$$

$$\begin{aligned} x_{F,g} &\longmapsto \gamma(g\lambda(F)) \cdot \gamma(\lambda(F))s_F \cdot (\gamma(g\lambda(F)))^{-1} \\ &= \gamma(g)s_F\gamma(g\lambda(F)) \end{aligned}$$


$$1 \longrightarrow \pi_1(M) \xrightarrow{\alpha} W \xrightarrow{\phi} \mathbb{Z}_2^n \longrightarrow 1 \quad (1)$$

$$\begin{array}{ccc} M & \xrightarrow{\quad \quad \quad} & \pi_1(M) \\ \pi \downarrow & & \downarrow \alpha \\ P & \xrightarrow{\quad \quad \quad} & W \end{array}$$

$$\begin{array}{ccc} M & \xrightarrow{\quad \quad \quad} & \pi_1(M) \\ \pi \downarrow & & \downarrow \alpha \\ P & \xrightarrow{\quad \quad \quad} & W \end{array}$$

$$\begin{array}{ccc} M & \xrightarrow{\quad\quad\quad} & \pi_1(M) \\ \pi \downarrow & & \downarrow \alpha \\ P & \xrightarrow{\quad\quad\quad} & W \end{array}$$

$$\begin{array}{ccc} M & \xrightarrow{\quad \quad \quad} & \pi_1(M) \\ \pi \downarrow & & \downarrow \alpha \\ P & \xrightarrow{\quad \quad \quad} & W \end{array}$$



For any proper face f of P ,

- Define $\mathcal{F}(f^\perp) \triangleq \{F \in \mathcal{F}(P) \mid \dim(f \cap F) = \dim(f) - 1\}$. So $\mathcal{F}(f^\perp)$ consists of those facets of P that intersect f transversely.

For any proper face f of P ,

- Define $\mathcal{F}(f^\perp) \triangleq \{F \in \mathcal{F}(P) \mid \dim(f \cap F) = \dim(f) - 1\}$. So $\mathcal{F}(f^\perp)$ consists of those facets of P that intersect f transversely.
- A submanifold Σ in M is called π_1 -injective if the inclusion $\Sigma \hookrightarrow M$ induces a monomorphism in the fundamental group.

Theorem (Wu-Yu, 2017)

Let M be a small cover over a simple polytope P and f be a proper face of P . The following two statements are equivalent.

- > The facial submanifold M_f is π_1 -injective.*
- > For any $F, F' \in \mathcal{F}(f^\perp)$, we have $f \cap F \cap F' \neq \emptyset$ whenever $F \cap F' \neq \emptyset$.*

Theorem (Wu-Yu, 2017)

Let M be a small cover over a simple polytope P and f be a proper face of P . The following two statements are equivalent.

- > The facial submanifold M_f is π_1 -injective.*
- > For any $F, F' \in \mathcal{F}(f^\perp)$, we have $f \cap F \cap F' \neq \emptyset$ whenever $F \cap F' \neq \emptyset$.*

Rk: The π_1 -injectivity of a facial submanifold of small cover only depends on the local face structure of f in P .

Theorem (Wu-Yu, 2017)

Let M be a small cover over a simple polytope P and f be a proper face of P . The following two statements are equivalent.

- > The facial submanifold M_f is π_1 -injective.*
- > For any $F, F' \in \mathcal{F}(f^\perp)$, we have $f \cap F \cap F' \neq \emptyset$ whenever $F \cap F' \neq \emptyset$.*

Rk: The π_1 -injectivity of a facial submanifold of small cover only depends on the local face structure of f in P .

Rk: We can determine the kernel of $i_* : \pi_1(M_f) \longrightarrow \pi_1(M)$.

A simple polytope P is called flag if a collection of facets of P has common intersection whenever they pairwise intersect.

A simple polytope P is called flag if a collection of facets of P has common intersection whenever they pairwise intersect.

Proposition (Wu-Yu, 2017)

Let M be a small cover over P . Then P is flag if and only if every facial submanifold of M is π_1 -injective.

A simple polytope P is called flag if a collection of facets of P has common intersection whenever they pairwise intersect.

Proposition (Wu-Yu, 2017)

Let M be a small cover over P . Then P is flag if and only if every facial submanifold of M is π_1 -injective.

Proposition (Wu-Yu, 2017)

For any small cover M over a 3-dimensional simple polytope P , there always exists a facet F of P so that the facial submanifold M_F is π_1 -injective.

For a 3-dimensional simple polytope P ,

- A k -circuit in P is a simple loop on the boundary of P which intersects transversely with the interior of exactly k distinct edges,

For a 3-dimensional simple polytope P ,

- A k -circuit in P is a simple loop on the boundary of P which intersects transversely with the interior of exactly k distinct edges, and a k -circuit is called prismatic if the endpoints of those edges are distinct.

For a 3-dimensional simple polytope P ,

- A k -circuit in P is a simple loop on the boundary of P which intersects transversely with the interior of exactly k distinct edges, and a k -circuit is called prismatic if the endpoints of those edges are distinct.
- A k -belt in P is a set of k distinct faces F_1, \dots, F_k of P such that $F_i \cap F_{i+1} \neq \emptyset$ for $1 \leq i \leq k-1$, $F_k \cap F_1 \neq \emptyset$, and any three faces in the belt have no common intersection.

For a 3-dimensional simple polytope P ,

- A k -circuit in P is a simple loop on the boundary of P which intersects transversely with the interior of exactly k distinct edges, and a k -circuit is called prismatic if the endpoints of those edges are distinct.
- A k -belt in P is a set of k distinct faces F_1, \dots, F_k of P such that $F_i \cap F_{i+1} \neq \emptyset$ for $1 \leq i \leq k-1$, $F_k \cap F_1 \neq \emptyset$, and any three faces in the belt have no common intersection.
- Each k -belt can determine a prismatic k -circuit.

For a 3-dimensional simple polytope P ,

- A k -circuit in P is a simple loop on the boundary of P which intersects transversely with the interior of exactly k distinct edges, and a k -circuit is called prismatic if the endpoints of those edges are distinct.
- A k -belt in P is a set of k distinct faces F_1, \dots, F_k of P such that $F_i \cap F_{i+1} \neq \emptyset$ for $1 \leq i \leq k-1$, $F_k \cap F_1 \neq \emptyset$, and any three faces in the belt have no common intersection.
- Each k -belt can determine a prismatic k -circuit. **A prismatic 3-circuit determines a 3-belt;**

For a 3-dimensional simple polytope P ,

- A k -circuit in P is a simple loop on the boundary of P which intersects transversely with the interior of exactly k distinct edges, and a k -circuit is called prismatic if the endpoints of those edges are distinct.
- A k -belt in P is a set of k distinct faces F_1, \dots, F_k of P such that $F_i \cap F_{i+1} \neq \emptyset$ for $1 \leq i \leq k-1$, $F_k \cap F_1 \neq \emptyset$, and any three faces in the belt have no common intersection.
- Each k -belt can determine a prismatic k -circuit. A prismatic 3-circuit determines a 3-belt; and if there is no prismatic 3-circuit, then a prismatic 4-circuit determines a 4-belt.

Let M be a 3-small cover over $P(\neq \Delta^3)$, the following facts was referred in DJ's paper(corrected):

- If there exist prismatic 3-circuits in P , then M can be decomposed into prime pieces glued along S^2 or $\mathbb{R}P^2$.

Let M be a 3-small cover over $P(\neq \Delta^3)$, the following facts was referred in DJ's paper(corrected):

- If there exist prismatic 3-circuits in P , then M can be decomposed into prime pieces glued along S^2 or $\mathbb{R}P^2$.
- If there is no prismatic 3-circuit but prismatic 4-circuits in P , then M can be decomposed into atoroidal and Seifert fibered pieces glued along tori or Klein bottles.

Let M be a 3-small cover over $P(\neq \Delta^3)$, the following facts was referred in DJ's paper(corrected):

- If there exist prismatic 3-circuits in P , then M can be decomposed into prime pieces glued along S^2 or $\mathbb{R}P^2$.
- If there is no prismatic 3-circuit but prismatic 4-circuits in P , then M can be decomposed into atoroidal and Seifert fibered pieces glued along tori or Klein bottles.
- If there is no prismatic 3 or 4-circuit in P , then M is hyperbolic.

Let M be a connected 3-manifold.

- M is called prime if $M = M_1 \# M_2$ implies $M_1 = S^3$ or $M_2 = S^3$.

Let M be a connected 3-manifold.

- M is called prime if $M = M_1 \# M_2$ implies $M_1 = S^3$ or $M_2 = S^3$.
- M is called irreducible if every embedded 2-sphere bounds a 3-ball. An prime 3-manifold is irreducible except S^2 -bundle over S^1 .

Let M be a connected 3-manifold.

- M is called prime if $M = M_1 \# M_2$ implies $M_1 = S^3$ or $M_2 = S^3$.
- M is called irreducible if every embedded 2-sphere bounds a 3-ball. An prime 3-manifold is irreducible except S^2 -bundle over S^1 .

Theorem (Kneser, Milnor, Prime Decomposition Theorem)

Each compact 3-manifold M can factor as a connected sum of prime manifolds. This decomposition is unique under the assumption that M is orientable.

Proposition

Let M be a 3-dimensional small cover over a simple polytope P , then M is irreducible if and only if there is no prismatic 3-circuit in P .

Proposition

Let M be a 3-dimensional small cover over a simple polytope P , then M is irreducible if and only if there is no prismatic 3-circuit in P .

In particular, the prime decomposition of an oriented 3-small cover is equivalent to cutting surgery along prismatic 3-circuits in P .

- A compact 3-manifold M is called atoroidal if it contains no essential torus.

- A compact 3-manifold M is called atoroidal if it contains no essential torus.
- A manifold M is called hyperbolic if it admits a complete Riemannian metric of constant sectional curvature -1 .

- A compact 3-manifold M is called atoroidal if it contains no essential torus.
- A manifold M is called hyperbolic if it admits a complete Riemannian metric of constant sectional curvature -1 .

Theorem (Jaco-Shalen-Johannson, Torus Decomposition Theorem)

For an oriented, irreducible, closed 3-manifold, there exists a (possibly empty) collection of disjointly embedded incompressible tori T_1, \dots, T_m such that each component of M cut along $T_1 \cup \dots \cup T_m$ is atoroidal or Seifert fibered, and a minimal such collection T_1, \dots, T_m is unique up to isotopy.

Theorem (Perelman, Geometrization Theorem)

Let M be a irreducible colsed 3-manifold. There exists a (possibly empty) collection of disjointly embedded incompressible surface S_1, \dots, S_m which are either tori or Klein bottles, such that each component of M cut along $S_1 \cup \dots \cup S_m$ is geometric. Any such collection of tori with a minimal number of components is unique up to isotopy.

Theorem (Perelman, Geometrization Theorem)

Let M be a irreducible colsed 3-manifold. There exists a (possibly empty) collection of disjointly embedded incompressible surface S_1, \dots, S_m which are either tori or Klein bottles, such that each component of M cut along $S_1 \cup \dots \cup S_m$ is geometric. Any such collection of tori with a minimal number of components is unique up to isotopy.

Proposition

Let M be a 3-small cover over a simple polytope P , then M is atoroidal if and only if there is no 4-belt in P .

In particular, the JSJ decomposition or geometric decomposition of a irreducible 3-small cover is equivalent to cutting surgery along prismatic 4-circuits in P .

Theorem (Thurston, Hyperbolization Theorem)

Each irreducible, atoroidal, closed 3-manifold with infinite fundamental group is hyperbolic.

Theorem (Thurston, Hyperbolization Theorem)

Each irreducible, atoroidal, closed 3-manifold with infinite fundamental group is hyperbolic.

Proposition

Let M be a 3-small cover over a simple polytope $P(\neq \Delta^3)$, then M is hyperbolic if and only if there is no prismatic 3 or 4-circuit in P .

Theorem (Thurston, Hyperbolization Theorem)

Each irreducible, atoroidal, closed 3-manifold with infinite fundamental group is hyperbolic.


Proposition

Let M be a 3-small cover over a simple polytope $P(\neq \Delta^3)$, then M is hyperbolic if and only if there is no prismatic 3 or 4-circuit in P .

Proposition (Wu-Yu, 2017)

A small cover M over a simple 3-polytope P can admit a Riemannian metric with nonnegative scalar curvature if and only if P is combinatorially equivalent to the cube $[0, 1]^3$ or a polytope obtained from Δ^3 by a sequence of vertex cuts.

A classification for 3-small cover



End of Talk

The 5th Korea Toric Topology Winter Workshop

Gyeongju, Korea. 10:30 - 11:10 January 22, 2019

Some references

-  Wu and Yu, Fundamental groups of small covers revisited. (2018).
-  Wu, Atoroidal manifolds in small covers. (2018).
-  Agol's *talk-1* & *talk-2* (2012, 2014).
-  Aschenbrenner-Friedl-Wilton, 3-manifold groups, (2013).
-  Buchstaber and Panov, Torus actions and their applications in topology and combinatorics. (2002).
-  Davis and Januszkiewicz, Convex polytopes, coxeter orbifolds and torus actions, (1991).
-  Chen, A homotopy theory of orbispaces. (2001).
-  Kapovich, Hyperbolic manifolds and discrete groups. (2010).



Email: wulisuwulisu@qq.com

Homepage: <http://algebraic.top/>