$$\bar{x} = \frac{\sum_{i=1}^{n} x_{i}}{n}$$

$$s^{2} = \frac{\sum_{i=1}^{n} (x_{i} - \bar{x})^{2}}{n-1}$$

$$s = \sqrt{s^{2}}$$

$$\bar{x} \pm z_{\frac{q}{2}}(\frac{s}{\sqrt{n}})$$

$$\bar{x} \pm t_{\frac{q}{2}}(\frac{s}{\sqrt{n}})...df = n-1$$

$$z = \frac{\bar{x} - x_{0}}{\frac{s}{\sqrt{n}}}$$

$$t = \frac{\bar{x} - x_{0}}{\frac{s}{\sqrt{n}}}...df = n-1$$

$$\hat{p} \pm z_{\frac{q}{2}}\sqrt{\frac{\hat{p}\hat{q}}{n}}$$

$$z = \frac{\hat{p} - p_{0}}{\sqrt{\frac{p_{0}q_{0}}{n}}}$$

$$(\bar{x}_{1} - \bar{x}_{2}) \pm t_{\frac{q}{2}}\sqrt{\frac{(n_{1} - 1)s_{1}^{2} + (n_{2} - 1)s_{2}^{2}}{n_{1} + n_{2} - 2}} * (\frac{1}{n_{1}} + \frac{1}{n_{2}})$$

$$t = \frac{(\bar{x}_{1} - \bar{x}_{2})}{\sqrt{\frac{(n_{1} - 1)s_{1}^{2} + (n_{2} - 1)s_{2}^{2}}{n_{1} + n_{2} - 2}} * (\frac{1}{n_{1}} + \frac{1}{n_{2}})$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$t = \frac{(\bar{x}_1 - \bar{x}_2)}{\sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}} * (\frac{1}{n_1} + \frac{1}{n_2})}$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A/B) = \frac{P(A \cap B)}{P(B)}$$

$${}_{N}C_{r} = \frac{N!}{r!(N-r)!}$$

 $f(x) = \binom{n}{n} p^x q^{n-x}$


$$(\bar{x}_1 - \bar{x}_2) \pm z_{\alpha/2} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

$$z = \frac{(\bar{x}_1 - \bar{x}_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

$$(\hat{p}_1 - \hat{p}_2) \pm z_{\alpha/2} \sqrt{\frac{\hat{p}_1 \hat{q}_1}{n_1} + \frac{\hat{p}_2 \hat{q}_2}{n_2}}$$

$$z = \frac{(\hat{p}_1 - \hat{p}_2)}{\sqrt{\hat{p}\hat{q}(\frac{1}{n_1} + \frac{1}{n_2})}} \dots \hat{p} = \frac{x_1 + x_2}{n_1 + n_2}$$

TABLE II Binomial Probabilities



Tabulated values are $\sum_{x=0}^{k} p(x)$. (Computations are rounded at the third decimal place.)

a. n = 5

a. # — J													
k p	.01	.05	.10	.20	.30	.40	.50	.60	.70	.80	.90	.95	.99
0	.951	.774	.590	.328	.168	.078	.031	.010	.002	.000	.000	.000	.000
1	.999	.977	.919	.737	.528	.337	.188	.087	.031	.007	.000	.000	.000
2	1.000	.999	.991	.942	.837	.683	.500	.317	.163	.058	.009	.001	.000
3	1.000	1.000	1.000	.993	.969	.913	.812	.663	.472	.263	.081	.023	.001
4	1.000	1.000	1.000	1.000	.998	.990	.969	.922	.832	.672	.410	.226	.049

b, n = 6

<i>k p</i>	.01	.05	.10	.20	.30	.40	.50	.60	.70	.80	.90	.95	.99
0	.941	.735	.531	.262	.118	.047	.016	.004	.001	.000	.000	.000	.000
1	.999	.967	.886	.655	.420	.233	.109	.041	.011	.002	.000	.000	.000
2	1.000	.998	.984	.901	.744	.544	.344	.179	.070	.017	.001	.000	.000
3	1.000	1.000	.999	.983	.930	.821	.656	.456	.256	.099	.016	.002	.000
4	1.000	1.000	1.000	.998	.989	. 95 9	.891	.767	.580	.345	.114	.033	.001
5	1.000	1.000	1.000	1.000	.999	.996	.984	.953	.882	.738	.469	.265	.059

c. n = 7

P													
k	.01	.05	.10	.20	.30	.40	.50	.60	.70	.80	.90	.95	.99
0	.932	.698	.478	.210	.082	.028	.008	.002	.000	.000	.000	.000	.000
1	.998	.956	.850	.577	.329	.159	.063	.019	.004	.000	.000	.000	.000
2	1.000	.996	.974	.852	.647	.420	.227	.096	.029	.005	.000	.000	.000
3	1.000	1.000	.997	.967	.874	.710	.500	.290	.126	.033	.003	.000	.000
4	1.000	1.000	1.000	.995	.971	.904	.773	.580	.353	.148	.026	.004	.000
5	1.000	1.000	1.000	1.000	.996	.981	.937	.841	.671	.423	.150	.044	.002
6	1.000	1.000	1.000	1.000	1.000	.998	.992	.972	.918	.790	.522	.302	.068

TABLE II (continued)


P													
	.01	.05	.10	.20	,30	.40	.50	.60	.70	.80	.90	.95	.99
0	.923	.663	.430	.168	.058	.017	.004	.001	.000	.000	.000	.000	.000
1	.997	.943	.813	.503	.255	.106	.035	.009	.001	.000	.000	.000	.000
2	1.000	.994	.962	.797	.552	.315	.145	.050	.011	.001	.000	.000	.000
3	1.000	1.000	.995	.944	.806	.594	.363	.174	.058	.010	.000	.000	.000
4	1.000	1.000	1.000	.990	.942	.826	.637	.406	.194	.056	.005	.000	.000
5	1.000	1.000	1.000	.999	.989	.950	.855	.685	.448	.203	.038	.006	.000
6	1.000	1.000	1.000	1.000	.999	.991	.965	.894	.745	.497	.187	.057	.003
7	1.000	1.000	1.000	1.000	1.000	.999	.996	.983	.942	.832	.570	.337	.077

$\mathbf{e.}\ n=9$													
k P	.01	.05	.10	.20	.30	.40	.50	.60	.70	.80	.90	.95	.99
0	.914	.630	.387	.134	.040	.010	.002	.000	.000	.000	.000	.000	.000
1	.997	.929	.775	.436	.196	.071	.020	.004	.000	.000	.000	.000	.000
2	1.000	.992	.947	.738	.463	.232	.090	.025	.004	.000	.000	.000	.000
3	1.000	.999	.992	.914	.730	.483	.254	.099	.025	.003	.000	.000	.000
4	1.000	1.000	.999	.980	.901	.733	.500	.267	.099	.020	.001	.000	.000
5	1.000	1.000	1.000	.997	.975	.901	.746	.517	.270	.086	.008	.001	.000
6	1.000	1.000	1.000	1.000	.996	.975	.910	.768	.537	.262	.053	.008	.000
7	1.000	1.000	1.000	1.000	1.000	.996	.980	.929	.804	.564	.225	.071	.003
8	1.000	1.000	1.000	1.000	1.000	1.000	.998	.990	.960	.866	.613	.370	.086

f. n = 10													
k p	.01	.05	.10	.20	.30	.40	.50	.60	.70	.80	.90	.95	.99
0	.904	.599	.349	.107	.028	.006	.001	.000	.000	.000	.000	.000	.000
1	.996	.914	.376	.376	.149	.046	.011	.002	.000	.000	.000	.000	.000
2	1.000	.988	.930	.678	.383	.167	.055	.012	.002	.000	.000	.000	.000
3	1.000	.999	.987	.879	.650	.382	.172	.055	.011	.001	.000	.000	.000
4	1.000	1.000	.998	.967	.850	.633	.377	.166	.047	.006	.000	.000	.000
5	1.000	1.000	1.000	.999	.953	.834	.623	.367	.150	.033	.002	.000	.000
6	1.000	1.000	1.000	.999	.989	.945	.828	.618	.350	.121	.013	.001	.000
7	1.000	1.000	1.000	1.000	.998	.988	.945	.833	.617	.322	.070	.012	.000
8	1.000	1.000	1.000	1.000	1.000	.998	.989	.954	.851	.624	.264	.086	.004
9	1.000	1.000	1.000	1.000	1.000	1.000	.999	.994	.972	.893	.651	.401	.096

(continues on next page)


TABLE III Normal Curve Areas



	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
.0	.0000	.0040	.0080	0120	0160	0100	0220	0070	0210	0250
	.0398	.0438	.0080	.0120 .0517	.0160 .0557	.0199	.0239	.0279	.0319	.0359
.1 .2	.0793	.0438	.0478	.0910	.0337	.0596 .0987	.0636 .1026	.0675	.0714	.0753
.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1004	.1103	.1141
.4	.1554	.1591	.1628	1664	.1700	.1736	.1772	.1808	.1844	.1517
. 4 .5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	
	1	1	l .	1	1		1	i	1	.2224
.6	.2257	.2291	.2324	.2357	.2389	.2422	.2454	.2486	.2517	.2549
.7	.2580	.2611	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
.8	.2881	.2910	.2939	.2967	.2995	.3023	.3051	.3078	.3106	.3133
.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4265	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4625	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4960	.4961	.4962	.4963	.4964
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981
2.9	.4981	.4982	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986
3.0	.4987	.4987	.4987	.4988	.4988	.4989	.4989	.4989	.4990	.4990

Source: Abridged from Table I of A. Hald, Statistical Tables and Formulas (New York: Wiley), 1952. Reproduced by permission of A. Hald.

TABLE IV Critical Values of t



ν	t _{.100}	t,050	t _{.025}	t _{.010}	t.005	t _{.001}	t.0005
1	3.078	6.314	12.706	31.821	63.657	318.31	636.62
2	1.886	2.920	4.303	6.965	9.925	22.326	31.598
3	1.638	2.353	3.182	4.541	5.841	10.213	12.924
4	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	1.476	2.015	2.571	3.365	4.032	5.893	6.869
6	1.440	1.943	2.447	3.143	3.707	5.208	5.959
7	1.415	1.895	2.365	2.998	3.499	4.785	5.408
8	1.397	1.860	2.306	2.896	3.355	4.501	5.041
9	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	1.372	1.812	2.228	2.764	3.169	4.144	4.587
11	1.363	1.796	2.201	2.718	3.106	4.025	4.437
12	1.356	1.782	2.179	2.681	3.055	3.930	4.318
13	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	1.341	1.753	2.131	2,602	2.947	3.733	4.073
16	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	1.333	1.740	2.110	2.567	2.898	3.646	3.965
18	1.330	1.734	2.101	2,552	2.878	3.610	3.922
19	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	1.325	1.725	2.086	2.528	2.845	3.552	3.850
21	1.323	1.721	2.080	2.518	2.831	3.527	3.819
22	1.321	1,717	2.074	2.508	2.819	3.505	3.792
. 23	1.319	1.714	2.069	2.500	2.807	3.485	3.767
24	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	1.314	1.703	2.052	2.473	2.771	3.421	3.690
28	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	1.310	1.697	2.042	2,457	2.750	3.385	3.646
40	1.303	1.684	2.021	2.423	2,704	3.307	3.551
60	1.296	1.671	2.000	2.390	2.660	3.232	3.460
120	1.289	1.658	1.980	2.358	2.617	3.160	3.373
	1.282	1.645	1.960	2.326	2.576	3.090	3.291

Source: This table is reproduced with the kind permission of the Trustees of Biometrika from E. S. Pearson and H.O. Hartley (eds.), The Biometrika Tables for Statisticians, Vol. 1, 3d ed., Biometrika, 1966.

TABLEV Critical Values of T_L and T_U for the Wilcoxon Rank Sum Test: Independent Samples

Test statistic is the rank sum associated with the smaller sample (if equal sample sizes, either rank sum can be used).

a. $\alpha = .025$ one-tailed; $\alpha = .05$ two-tailed

n_2		3		1		5		6		7		3	9)	1	0
	T _L	T _U	T _L	Tu	T _L	T_{U}	T _L	Tu	T _L	T _U	T _L	$T_{\mathbf{U}}$	T _L	T_{U}	T _L	T _U
3	5	16	6	18	6	21	7	23	7	26	8	28	8	31	9	33
4	6	18	11	25	12	28	12	32	13	35	14	38	15	41	16	44
5	6	21	12	28	18	37	19	41	20	45	21	49	22	53	24	56
6	7	23	12	32	19	41	26	52	28	56	29	61	31	65	32	70
7	7	26	13	35	20	45	28	56	37	68	39	73	41	78	43	83
8	8	28	14	38	21	49	29	61	39	73	49	87	51	93	54	98
9	8	31	15	41	22	53	31	65	41	78	51	93	63	108	66	114
10	9	33	16	44	24	56	32	70	43	83	54	98	66	114	79	131

b. $\alpha = .05$ one-tailed; $\alpha = .10$ two-tailed

n_2		3		4		5		6	•	7	8	})	1	.0
	$T_{\rm L}$	T_{U}	TL	T_{U}	T _L	T _U	T _L	T _U	T _L	T _U	T _L	T_{U}	T _L	Tu	T _L	T _U
3	6	15	7	17	7	20	8	22	9	24	9	27	10	29	11	31
4	7	17	12	24	13	27	14	30	15	33	16	36	17	39	18	42
5	7	20	13	27	19	36	20	40	22	43	24	46	25	50	26	54
6	8	22	14	30	20	40	28	50	30	54	32	58	33	63	35	67
7	9	24	15	33	22	43	30	54	39	66	41	71	43	76	46	80
8	9	27	16	36	24	46	32	58	41	71	52	84	54	90	57	95
9	10	29	17	39	25	50	33	63	43	76	54	90	66	105	69	111
10	11	31	18	42	26	54	35	67	46	80	57	95	69	111	83	127

Source: From F. Wilcoxon and R. A. Wilcox, "Some Rapid Approximate Statistical Procedures," 1964, 20-23.