

Original: 4925 tokens

Ours: 1047 tokens




Figure 3. LncRNA-CGB knock-out (KO) mice developed more severe TB than did control mice. (a) Northern blotting analysis of lncRNA expression in splenocytes isolated from wild-type (WT) or lncRNA-CGB KO mice. (b) The experimental diagram showed that WT or lncRNA-CGB KO mice were infected with *M. tuberculosis* and sacrificed at the indicated time point. (c) Gross pathology of representative lungs isolated from WT and lncRNA-CGB KO mice. The red circle marked more severe pathological lesions. (d) H&E staining of sections of representative lungs derived from *M. tuberculosis*-infected lncRNA-CGB KO or WT mice. Yellow arrows heads marked the granuloma lesions or accumulation of inflammatory cells in the lungs. Note that larger-scale and more granuloma lesions were observed in lncRNA-CGB KO mice. (e) Macroscopic evaluation of *M. tuberculosis* bacilli burdens and distribution on the sections of lungs isolated from *M. tuberculosis*-infected WT and lncRNA-CGB KO mice. Representative fields were shown. Note that more bacilli positive for acid-fast staining was observed in lesions of *M. tuberculosis*-infected lncRNA-CGB KO mice. Lower panel: higher-resolution view of the regions as indicated by the box in the upper panel. (f) *M. tuberculosis* CFU analysis of bacillus burdens in the lungs (per gram) of *M. tuberculosis*-infected WT or lncRNA-CGB KO mice. $N = 7$ SPF mice for each group for each experiment with at least two biological repeats. Error bars indicate average values $\pm$ SEM. ** $p < .01$.




Figure 3. LncRNA-CGB knock-out (KO) mice developed more severe TB than did control mice. (a) Northern blotting analysis of lncRNA expression in splenocytes isolated from wild-type (WT) or lncRNA-CGB KO mice. (b) The experimental diagram showed that WT or lncRNA-CGB KO mice were infected with *M. tuberculosis* and sacrificed at the indicated time point. (c) Gross pathology of representative lungs isolated from WT and lncRNA-CGB KO mice. The red circle marked more severe pathological lesions. (d) H&E staining of sections of representative lungs derived from *M. tuberculosis*-infected lncRNA-CGB KO or WT mice. Yellow arrows heads marked the granuloma lesions or accumulation of inflammatory cells in the lungs. Note that larger-scale and more granuloma lesions were observed in lncRNA-CGB KO mice. (e) Macroscopic evaluation of *M. tuberculosis* bacilli burdens and distribution on the sections of lungs isolated from *M. tuberculosis*-infected WT and lncRNA-CGB KO mice. Representative fields were shown. Note that more bacilli positive for acid-fast staining was observed in lesions of *M. tuberculosis*-infected lncRNA-CGB KO mice. Lower panel: higher-resolution view of the regions as indicated by the box in the upper panel. (f) *M. tuberculosis* CFU analysis of bacillus burdens in the lungs (per gram) of *M. tuberculosis*-infected WT or lncRNA-CGB KO mice. $N = 7$ SPF mice for each group for each experiment with at least two biological repeats. Error bars indicate average values $\pm$ SEM. ** $p < .01$.