



Click "Global", "Regions", or "Projects" to browse AntWeb's catalogs.


[X](#)Species: *Acanthomyrmex concavus*

Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in mature wet forest, 1 times found in lowland rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected from 350 - 500 meters, 400 meters average


Type specimens: Holotype of *Acanthomyrmex concavus*: casent0901700Species: *Acanthomyrmex ferox*

Specimen Habitat Summary

Found most commonly in these habitats: 3 times found in tropical rainforest, 2 times found in mature wet forest, 1 times found in on rainforest floor, 2 times found in rainforest, 3 times found in lowland rainforest, 1 times found in second-growth rainforest, 1 times found in 2nd growth wet forest

Collected most commonly using these methods or in the following microhabitats: 6 times winkler, 1 times In dead branch, 1 times pitfall, litter, 1 times search

Elevations: collected from 88 - 515 meters, 260 meters average

Type specimens: Syntype of *Acanthomyrmex ferox*: casent0904703Species: *Acropyga my02*

Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in litter, rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler



Elevations: collected at 225 m

Type specimens:

Species: *Aenictus chapmani_cf*



Species: *Aenictus levior*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in 2nd growth wet forest

Collected most commonly using these methods or in the following microhabitats: 1 times search

Elevations: collected from 250 - 450 meters, 383 meters average

Type specimens:

Species: *Aenictus montivagus*





Specimen Habitat Summary

Elevations: collected at 1600 m

Type specimens: paratype of *Aenictus montivagus*: casent0903689

Species: *Anochetus agilis*




Specimen Habitat Summary

Found most commonly in these habitats: 0 times found in Degraded coastal hill forest

Type specimens: Holotype of Anochetus agilis: casent0903984

Species: Anochetus my02



Specimen Habitat Summary



Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 18 m

Type specimens:

Species: Anochetus my03



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 28 m

Type specimens:

Species: Anochetus my04



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 20 m


Type specimens:

Species: Anomalomyrma taylori





Species: *Aphaenogaster my01*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 180 m

Type specimens:


Species: *Calyptomyrmex danum*



Specimen Habitat Summary

Type specimens: Holotype of *Calyptomyrmex danum*: casent0900995

Species: *Calyptomyrmex my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 937 m

Type specimens:

Species: *Calyptomyrmex sabahensis*



Specimen Habitat Summary


Found most commonly in these habitats: 9 times found in Rainforest

Collected most commonly using these methods or in the following microhabitats: 10 times Berlesate

Elevations: collected from 228 - 229 meters, 228 meters average

Type specimens: paratype of *Calyptomyrmex sabahensis*: casent0900999

Species: *Camponotus cylindricus_nr*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Fogging, canopy

Elevations: collected at 558 m

Type specimens:

Species: *Camponotus gigas*



Specimen Habitat Summary


Found most commonly in these habitats: 4 times found in rainforest, 1 times found in rainforest edge, 1 times found in tropical rainforest, 1 times found in mature wet forest


Collected most commonly using these methods or in the following microhabitats: 2 times search, 2 times Hand

Elevations: collected from 100 - 558 meters, 240 meters average

Type specimens:

Species: *Camponotus my01*





Specimen Habitat Summary



Found most commonly in these habitats: 2 times found in litter, rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times pitfall

Elevations: collected from 130 - 173 meters, 158 meters average

Type specimens:

Species: *Camponotus saundersi*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Fogging, canopy

Elevations: collected from 200 - 558 meters, 289 meters average

Type specimens: syntype of *Camponotus saundersi*: casent0905463

Species: *Cardiocondyla my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in rainforest


Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, 1 times Winkler

Elevations: collected from 1 - 104 meters, 46 meters average

Type specimens:

Species: *Carebara affinis*





Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in lowland wet forest, 1 times found in oil palm plantation, 1 times found in sclerophyl forest, 1 times found in litter, rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler, 1 times pitfall, litter, 1 times search

Elevations: collected from 50 - 180 meters, 144 meters average

Type specimens: Holotype of *Solenopsis calida*: casent0901423; syntype of *Solenopsis laboriosa*: casent0901419, casent0901420

Species: *Carebara alperti*





Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 175 m

Type specimens:

Species: *Carebara my01*



Specimen Habitat Summary



Found most commonly in these habitats: 1 times found in litter, rainforest, 2 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, 2 times Winkler

Elevations: collected from 56 - 937 meters, 496 meters average

Type specimens:

Species: *Carebara my04*



Specimen Habitat Summary


Found most commonly in these habitats: 3 times found in rainforest, 1 times found in oil palm plantation

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, litter, 2 times Winkler, 1 times pitfall

Elevations: collected from 57 - 104 meters, 84 meters average

Type specimens:

Species: Carebara silena



Specimen Habitat Summary



Found most commonly in these habitats: 2 times found in litter, rainforest, 1 times found in oil palm plantation

Collected most commonly using these methods or in the following microhabitats: 2 times pitfall, 1 times pitfall, litter

Elevations: collected from 104 - 173 meters, 132 meters average

Type specimens: syntype of *Pheidologeton silenus*: casent0900737, casent0900738

Species: Cataulacus hispidulus



Specimen Habitat Summary



Found most commonly in these habitats: 1 times found in tropical rainforest edge, 1 times found in rainforest edge, 1 times found in wet forest

Collected most commonly using these methods or in the following microhabitats: 2 times search

Elevations: collected from 170 - 230 meters, 182 meters average

Type specimens: Holotype of *Cataulacus hispidulus*: casent0901452; syntype of *Cataulacus brookei*: casent0909236

Species: Cataulacus my01



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest


Collected most commonly using these methods or in the following microhabitats: 1 times Hand

Elevations: collected at 525 m

Type specimens:

Species: Centromyrmex my01





Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 20 m

Type specimens:

Species: *Cerapachys cerberus*



Specimen Habitat Summary

Elevations: collected at 250 m

Type specimens: Holotype of *Cerapachys cerberus*: casent0902727

Species: *Cerapachys doryloides*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times hypogaeic at oil bait, 1 times soil core

Type specimens: Holotype of *Cerapachys doryloides*: casent0902717

Species: *Cerapachys mlb-browni*



Specimen Habitat Summary

Elevations: collected at 600 m

Type specimens: Holotype of Cerapachys browni: casent0902736

Species: Cerapachys mlb-dyak




Specimen Habitat Summary

Elevations: collected at 1270 m

Type specimens: Holotype of Cerapachys dyak: casent0902729

Species: Cerapachys mlb-venator




Specimen Habitat Summary

Elevations: collected at 1880 m

Type specimens: Holotype of Cerapachys venator: casent0902724

Species: Cerapachys mlb-vermiculus



Specimen Habitat Summary

Elevations: collected at 1600 m

Type specimens: Holotype of Cerapachys vermiculus: casent0902735

Species: Cerapachys mlb-verminosus






Specimen Habitat Summary

Elevations: collected at 250 m

Type specimens: Holotype of *Cerapachys verminosus*: casent0902734

Species: *Cerapachys suscitatus_nr*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 225 m

Type specimens:

Species: *Chimaeridris burckhardti*




Specimen Habitat Summary

Elevations: collected at 1500 m

Type specimens:

Species: *Cladomyrma* (indet)



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in 2nd growth vegetation, 1 times found in wet forest

Collected most commonly using these methods or in the following microhabitats: 2 times search

Elevations: collected from 250 - 320 meters, 292 meters average

Type specimens:


Species: Cladomyrma andrei



Specimen Habitat Summary

Type specimens: Holotype of Cladomyrma cryptata: casent0903186; paratype: casent0173912, casent0173913; syntype of Cladomyrma andrei: casent0905172

Species: Cladomyrma andrei_nr




Specimen Habitat Summary

Elevations: collected at 1000 m

Type specimens:

Species: Cladomyrma crypteroniae



Specimen Habitat Summary

Type specimens: paratype: casent0173916, casent0173917, casent0173918

Species: Cladomyrma dianeae




Specimen Habitat Summary

Type specimens: paratype: casent0173914, casent0173915

Species: *Cladomyrma hewitti*



Species: *Cladomyrma maryatiae*




Specimen Habitat Summary

Elevations: collected at 600 m

Type specimens: paratype: casent0173900, casent0173901, casent0173902

Species: *Cladomyrma my03*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 558 m

Type specimens:

Species: *Crematogaster (Crematogaster) (inflata) ampullaris*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 220 m

Type specimens:

Species: *Crematogaster decamera*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in secondary and primary rainforest

Elevations: collected at 180 m

Type specimens: Lectotype of *Crematogaster decamera*: casent0908461; paralectotype of *Crematogaster decamera*: casent0908462; syntype of *Crematogaster decamera*: casent0902110

Species: *Crematogaster fraxatrix*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 175 m

Type specimens: Lectotype of *Crematogaster fraxatrix*: casent0908581

Species: *Crematogaster gavapiga_cf*





Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in rainforest

Elevations: collected from 175 - 220 meters, 197 meters average

Type specimens:

Species: *Crematogaster inflata*





Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in 2nd growth wet forest, 1 times found in on low vegetation

Collected most commonly using these methods or in the following microhabitats: 2 times search

Elevations: collected from 175 - 450 meters, 321 meters average

Type specimens: syntype of *Crematogaster inflata*: casent0901438

Species: *Crematogaster masukoi*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 220 m

Type specimens: paratype of *Crematogaster masukoi*: casent0911144

Species: *Crematogaster modiglianii*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest, 1 times found in tropical rainforest edge, 1 times found in tropical rainforest


Collected most commonly using these methods or in the following microhabitats: 2 times search

Elevations: collected from 100 - 190 meters, 167 meters average

Type specimens: syntype of *Crematogaster modiglianii*: casent0904531; syntype of *Crematogaster modiglianii clemensae*: casent0908635; syntype of *Crematogaster modiglianii sarawakana*: casent0908636; syntype of *Crematogaster modiglianii anamita*: casent0912787; syntype of *Crematogaster modiglianii anoemica*: casent0912788; syntype of *Crematogaster modiglianii surbeki*: casent0912789

Species: *Crematogaster my01*





Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 937 m

Type specimens:

Species: *Crematogaster my02*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in oil palm plantation, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Pitfall, litter, 1 times pitfall

Elevations: collected from 64 - 173 meters, 91 meters average

Type specimens:

Species: *Crematogaster ochracea_cf*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest edge

Elevations: collected at 180 m

Type specimens:

Species: *Crematogaster reticulata*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in secondary rainforest

Elevations: collected at 300 m

Type specimens: paratype of *Crematogaster reticulata*: casent0905905; syntype of *Crematogaster reticulata*: casent0912718

Species: *Crematogaster rogenhoferi_cf*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest edge

Elevations: collected at 180 m

Type specimens:

Species: *Cryptopone my01*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 180 m

Type specimens:

Species: *Cryptopone testacea*



Specimen Habitat Summary

Found most commonly in these habitats: 2 times found in Rainforest, 0 times found in Lowland rainforest, 0 times found in Partly cleared forest, 0 times found in Ridge forest, 1 times found in around camp, 0 times found in Bank Pagota R.

Collected most commonly using these methods or in the following microhabitats: 2 times Hand-Collected from 1 square meter of sifted leaf litter, 1 times 6L sifted litter, extracted in Winkler sack over 48 hours

Elevations: collected from 10 - 1400 meters, 472 meters average

Type specimens:

Species: *Dacetinops concinnus*






Specimen Habitat Summary

Found most commonly in these habitats: 3 times found in rainforest, 1 times found in lowland rainforest

Elevations: collected at 229 m

Type specimens:

Species: *Dacetinops my01*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected from 106 - 558 meters, 377 meters average

Type specimens:

Species: *Dacetinops solivagus*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest


Collected most commonly using these methods or in the following microhabitats: 1 times berlesate

Elevations: collected at 229 m

Type specimens: paratype: casent0172457

Species: *Diacamma intricatum*





Specimen Habitat Summary



Found most commonly in these habitats: 1 times found in tropical rainforest, 2 times found in 2nd growth wet forest, 1 times found in mature wet forest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 3 times search, 1 times bait, 1 times Winkler

Elevations: collected from 190 - 760 meters, 408 meters average

Type specimens: syntype of *Diacamma intricatum*: casent0900669, casent0901341

Species: *Diacamma rugosum*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in wet forest, 2 times found in rainforest, 1 times found in lowland wet forest, 1 times found in Patch of rainforest on coffee farm

Collected most commonly using these methods or in the following microhabitats: 3 times search, 2 times Winkler, 1 times Pitfall

Elevations: collected from 26 - 558 meters, 226 meters average

Type specimens: Holotype of *Diacamma rugosum japensis*: casent0900673; Holotype of *Ponera geometrica*: casent0901342; Holotype of *Ponera sculpturata*: casent0901343; Holotype of *Ponera vagans*: casent0901344; syntype of *Diacamma rugosum*: casent0913723; syntype of *Diacamma vagans papuanum*: casent0915179; syntype of *Diacamma versicolor*: casent0900672; syntype of *Diacamma bispinosum saussurei*: casent0907220; syntype of *Diacamma rugosum geminatum*: casent0903878; syntype of *Ponera tortuolosa*: casent0901345

Species: *Dilobocondyla my01*





Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected from 150 - 225 meters, 175 meters average

Type specimens:

Species: *Discothyrea my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected from 130 - 558 meters, 272 meters average

Type specimens:

Species: Dolichoderus (indet)



Specimen Habitat Summary


Found most commonly in these habitats: 10 times found in Rainforest, 5 times found in Mallee, 1 times found in near river, 0 times found in Tropical hardwood forest, 2 times found in mature wet forest, 0 times found in montane rainforest, 2 times found in dry sclerophyll, 3 times found in eucalypt woodland, 2 times found in tropical wet forest, 2 times found in montane wet forest, ...

Collected most commonly using these methods or in the following microhabitats: 1 times day sweeping along forest trail, 8 times Malaise, 4 times blacklight, 5 times search, 1 times Malaise trap, 3 times winkler, 2 times pan trap, 1 times FIT, 0 times Hollow twig, 2 times Beating, 1 times ex hanging nest, ...

Elevations: collected from 10 - 1670 meters, 360 meters average

Type specimens:

Species: Dolichoderus cuspidatus



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in Rainforest, 1 times found in litter, rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, 1 times Winkler

Elevations: collected from 106 - 1050 meters, 301 meters average

Type specimens: syntype of Dolichoderus cuspidatus: casent0901856; syntype of Dolichoderus cuspidatus surbecki: casent0911495; syntype of Dolichoderus cuspidatus tenuispina: casent0909481

Species: Dolichoderus indrapurensis




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in tropical rainforest, 1 times found in wet forest

Collected most commonly using these methods or in the following microhabitats: 2 times search

Elevations: collected from 190 - 230 meters, 203 meters average

Species: Dolichoderus kinabaluensis



Specimen Habitat Summary

Found most commonly in these habitats: 2 times found in cloud forest

Collected most commonly using these methods or in the following microhabitats: 2 times search

Elevations: collected from 1560 - 2260 meters, 2143 meters average

Type specimens: Holotype of Dolichoderus kinabaluensis: casent0902965

Species: Dolichoderus magnipastor



Specimen Habitat Summary

Elevations: collected at 600 m

Type specimens: Holotype of Dolichoderus magnipastor: casent0902962

Species: Dolichoderus maschwitzi





Specimen Habitat Summary

Elevations: collected at 600 m

Type specimens: Holotype of Dolichoderus maschwitzi: casent0902964

Species: Dolichoderus my01





Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, forest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected at 24 m

Type specimens:

Species: *Dolichoderus my02*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, forest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected at 937 m

Type specimens:

Species: *Dolichoderus my03*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected at 558 m

Type specimens:

Species: *Dolichoderus pastorulus*




Specimen Habitat Summary

Elevations: collected at 600 m

Type specimens: Holotype of Dolichoderus pastorulus: casent0902966

Species: *Dolichoderus thoracicus_group*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in wet forest, 1 times found in 2nd growth wet forest, 1 times found in mature wet forest

Collected most commonly using these methods or in the following microhabitats: 5 times search

Elevations: collected from 200 - 1670 meters, 422 meters average

Type specimens:

Species: *Dorylus laevigatus*



Specimen Habitat Summary


Found most commonly in these habitats: 3 times found in rainforest, 1 times found in oil palm plantation

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, litter, 1 times Winkler

Elevations: collected from 28 - 556 meters, 157 meters average

Type specimens: syntype of *Dorylus laevigatus*: casent0902696; syntype of *Dorylus breviceps*: casent0903709

Species: *Echinopla lineata*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in mangrove forest, 1 times found in rainforest


Collected most commonly using these methods or in the following microhabitats: 1 times search, 1 times Fogging, canopy

Elevations: collected from 5 - 937 meters, 335 meters average

Type specimens: syntype of *Echinopla sucki*: casent0910967

Species: *Ectomomyrmex astutus*





Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 34 m

Type specimens: syntype of *Pachycondyla astuta*: casent0902492

Species: *Ectomomyrmex leeuwenhoeki*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in tropical rainforest, 1 times found in 2nd growth wet forest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler, 1 times 6L sifted litter, extracted in Winkler sack over 36 hours

Elevations: collected from 210 - 330 meters, 242 meters average

Type specimens: syntype of *Pachycondyla leeuwenhoeki*: casent0907272

Species: *Emeryopone my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest


Collected most commonly using these methods or in the following microhabitats: 1 times night 9pm winkler

Elevations: collected at 180 m

Type specimens:

Species: *Epelysidris brocha*





Specimen Habitat Summary



Found most commonly in these habitats: 1 times found in tropical rainforest, 1 times found in Moss Forest

Collected most commonly using these methods or in the following microhabitats: 1 times search, 1 times search in soil

Elevations: collected from 210 - 1219 meters, 546 meters average

Type specimens:

Species: *Euprenolepis* (indet)



Specimen Habitat Summary



Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in tropical rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, 1 times search

Elevations: collected from 173 - 500 meters, 256 meters average

Type specimens:

Species: *Euprenolepis* my01



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in litter, rainforest, 2 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, 1 times Winkler

Elevations: collected from 57 - 180 meters, 145 meters average

Type specimens:

Species: *Eurhopalothrix* *cinnamea*



Species: *Eurhopalothrix coronata*




Specimen Habitat Summary

Elevations: collected at 600 m

Type specimens:

Species: *Eurhopalothrix dubia*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in tropical rainforest, 1 times found in 2nd growth wet forest, 2 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 5 times Winkler

Elevations: collected from 28 - 330 meters, 200 meters average

Type specimens:

Species: *Forelophilus my01*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected from 500 - 525 meters, 512 meters average

Type specimens:

Species: *Forelophilus overbecki*






Specimen Habitat Summary

Elevations: collected at 1750 m

Type specimens: paralectotype of *Forelophilus overbecki*: casent0907588

Species: *Gauromyrmex my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest edge

Collected most commonly using these methods or in the following microhabitats: 1 times at light

Elevations: collected at 190 m

Type specimens:

Species: *Gesomyrmex my01*





Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 525 m

Type specimens:

Species: *Gnamptogenys binghamii*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in tropical rainforest, 0 times found in Rubber plantn. edge rainforest, 1 times found in tropical broadleaf moist forest

Collected most commonly using these methods or in the following microhabitats: 2 times search, 1 times Winkler

Elevations: collected from 190 - 900 meters, 329 meters average

Type specimens: syntype of *Gnamptogenys binghamii*: casent0907195; syntype of *Stictoponera borneensis*: casent0905738

Species: *Gnamptogenys cibrata*



Specimen Habitat Summary


Found most commonly in these habitats: 4 times found in tropical rainforest, 2 times found in 2nd growth wet forest, 3 times found in mature wet forest, 1 times found in wet forest, 1 times found in peak of limestone pinnacle, scrubby veg.

Collected most commonly using these methods or in the following microhabitats: 10 times Winkler, 1 times search

Elevations: collected from 180 - 1010 meters, 416 meters average

Type specimens: syntype of *Ectatomma diehlii*: casent0907130; syntype of *Gnamptogenys cibrata*: casent0903809; syntype of *Havilandius diehlii*: casent0900587

Species: *Gnamptogenys fontana*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in mature wet forest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected from 350 - 900 meters, 625 meters average

Type specimens: Holotype of *Gnamptogenys fontana*: casent0911163; paratype of *Gnamptogenys fontana*: casent0900558

Species: *Gnamptogenys lacunosa*





Specimen Habitat Summary

Elevations: collected at 1600 m

Type specimens: paratype of *Gnamptogenys lacunosa*: casent0900591

Species: *Gnamptogenys leiolabia*





Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in Rainforest, 1 times found in mature wet forest, 1 times found in tropical rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected from 350 - 515 meters, 432 meters average

Type specimens: holotype: anic32-017684

Species: *Gnamptogenys menadensis*



Specimen Habitat Summary



Found most commonly in these habitats: 2 times found in wet forest, 1 times found in primary dipterocarp forest, 1 times found in tropical rainforest

Collected most commonly using these methods or in the following microhabitats: 4 times search

Elevations: collected from 200 - 760 meters, 320 meters average

Type specimens:

Species: *Gnamptogenys my01*



Specimen Habitat Summary



Found most commonly in these habitats: 2 times found in rainforest, 1 times found in litter, rainforest, 1 times found in oil palm plantation


Collected most commonly using these methods or in the following microhabitats: 3 times Winkler, 1 times pitfall, litter

Elevations: collected from 28 - 150 meters, 106 meters average

Type specimens:

Species: *Gnamptogenys my02*





1 mm

Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected at 28 m

Type specimens:

Species: *Gnamptogenys my03*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected at 558 m

Type specimens:

Species: *Gnamptogenys my05*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected at 34 m

Type specimens:

Species: *Gnamptogenys my06*



Specimen Habitat Summary



Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 34 m

Type specimens:

Species: *Gnamptogenys ortostoma*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in Degraded coastal hill forest on granite

Type specimens: holotype: anic32-017689

Species: *Gnamptogenys pertusa*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in Rainforest

Elevations: collected at 500 m

Type specimens: holotype: anic32-017691

Species: *Gnamptogenys scalpta*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in montane rainforest

Elevations: collected at 1585 m

Type specimens: Holotype of *Gnamptogenys scalpta*: casent0911168

Species: *Gnamptogenys sila*





Specimen Habitat Summary

Elevations: collected at 330 m

Type specimens: Holotype of *Gnamptogenys sila*: casent0900589

Species: *Gnamptogenys treta*




Specimen Habitat Summary

Elevations: collected at 500 m

Type specimens: Holotype of *Gnamptogenys treta*: casent0911170

Species: *Harpegnathos macgregori*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 558 m

Type specimens:

Species: *Leptanilla my01*




Specimen Habitat Summary

Elevations: collected at 1610 m

Type specimens:

Species: *Leptogenys my01*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in tropical rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times search, 2 times Winkler

Elevations: collected from 18 - 190 meters, 155 meters average

Type specimens:

Species: *Leptogenys my02*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in oil palm plantation, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler, 1 times pitfall, litter

Elevations: collected from 104 - 558 meters, 205 meters average

Type specimens:

Species: *Leptogenys my03*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in oil palm plantation, 1 times found in rainforest


Collected most commonly using these methods or in the following microhabitats: 3 times Winkler

Elevations: collected from 150 - 558 meters, 231 meters average

Type specimens:

Species: *Leptogenys my04*





Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 88 m

Type specimens:

Species: *Leptogenys my05*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected from 57 - 150 meters, 119 meters average

Type specimens:

Species: *Leptogenys my06*



Specimen Habitat Summary



Found most commonly in these habitats: 3 times found in mature wet forest, 2 times found in tropical rainforest, 1 times found in 2nd growth wet forest, 1 times found in rainforest


Collected most commonly using these methods or in the following microhabitats: 7 times winkler

Elevations: collected from 18 - 760 meters, 356 meters average

Type specimens:

Species: *Leptogenys my07*





Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 173 m

Type specimens:

Species: *Leptogenys my08*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in oil palm plantation, 1 times found in tropical rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Pitfall, litter, 1 times search

Elevations: collected from 170 - 173 meters, 171 meters average

Type specimens:

Species: *Liomyrmex gestroi*



Specimen Habitat Summary



Found most commonly in these habitats: 2 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times at night 9pm

Elevations: collected from 20 - 180 meters, 60 meters average

Type specimens: Holotype of *Liomyrmex froggatti* major: casent0901738; syntype of *Liomyrmex froggatti*: casent0901736, casent0901737; syntype of *Liomyrmex gestroi*: casent0905754; syntype of *Liomyrmex reneae*: casent0901739; syntype of *Liomyrmex aurianus*: casent0904615; syntype of *Myrmica caeca*: casent0901735

Species: *Lophomyrmex longicornis*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in lowland rainforest, 2 times found in tropical rainforest, 1 times found in mature wet forest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler, 1 times bait, 1 times Pitfall trap, litter

Elevations: collected from 190 - 937 meters, 403 meters average

Type specimens: Holotype of *Lophomyrmex longicornis*: casent0901267; paratype: anic32-015010, casent0178544

Species: *Lophomyrmex my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 57 m

Type specimens:

Species: *Lordomyrma* (indet)



Specimen Habitat Summary


Found most commonly in these habitats: 21 times found in montane rainforest, 16 times found in Rainforest, 6 times found in Montane Forest, 3 times found in Moss for., 3 times found in wet sclerophyll, 4 times found in Eucalypt forest, 0 times found in Logs, 0 times found in Lowland Rainforest, 3 times found in tropical rainforest, 1 times found in plot - 3, ...

Collected most commonly using these methods or in the following microhabitats: 2 times Berlesate, 5 times at honey bait, 0 times Pyrethrum, 4 times pitfall, 0 times Pyrethrum knockdown, 3 times search, 0 times Intercept, 1 times Winkler, 0 times flight intercept trap

Elevations: collected from 5 - 4500 meters, 1784 meters average

Type specimens:

Species: *Lordomyrma my01*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in litter, rainforest


Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 64 m

Type specimens:

Species: *Lordomyrma reticulata*





Specimen Habitat Summary


Found most commonly in these habitats: 5 times found in tropical rainforest, 2 times found in 2nd growth wet forest, 2 times found in clay bank in mature wet forest, 1 times found in litter, rainforest, 1 times found in logged forest

Collected most commonly using these methods or in the following microhabitats: 8 times search, 2 times Winkler, 1 times mini winkler

Elevations: collected from 64 - 360 meters, 219 meters average

Type specimens:

Species: *Loweriella boltoni*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in primary forest, 1 times found in wet forest, 1 times found in Borneo rainforest, 1 times found in lowl.rainfor., 1 times found in lowland rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times hand collection, 1 times search

Elevations: collected from 350 - 620 meters, 496 meters average

Type specimens: paratype: anic32-015034, casent0172291, casent0178492

Species: *Mayriella my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in rainforest


Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, 1 times Winkler

Elevations: collected from 56 - 937 meters, 349 meters average

Type specimens:

Species: *Meranoplus castaneus*






Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest, 1 times found in rainforest edge

Elevations: collected from 100 - 525 meters, 272 meters average

Type specimens: Holotype of *Meranoplus castaneus*: casent0901456; Holotype of *Meranoplus cordatus*: casent0901457; Lectotype of *Meranoplus castaneus hammaceros*: casent0908935

Species: *Meranoplus malaysianus*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in mature wet forest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected from 24 - 350 meters, 187 meters average

Type specimens: Holotype of *Meranoplus malaysianus*: casent0902029

Species: *Meranoplus mucronatus*



Specimen Habitat Summary



Found most commonly in these habitats: 3 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 4 times Winkler

Elevations: collected from 106 - 1219 meters, 251 meters average

Type specimens: Holotype of *Meranoplus mucronatus*: casent0902028

Species: *Myopias my01*



0.5 mm

Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in mature wet forest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected from 350 - 937 meters, 752 meters average

Type specimens:

Species: *Myopopone my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in oil palm plantation, 1 times found in rainforest, 1 times found in rotten log, rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, 1 times Hand, 1 times Winkler

Elevations: collected from 173 - 558 meters, 271 meters average

Type specimens:

Species: *Myrmecina my02*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in oil palm plantation, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler, 1 times pitfall

Elevations: collected from 57 - 173 meters, 140 meters average

Type specimens:

Species: *Myrmicaria bey-mary*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 558 m

Type specimens:

Species: *Myrmicaria carinata*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall

Elevations: collected from 173 - 460 meters, 268 meters average

Type specimens: Holotype of *Myrmicaria carinata*: casent0901427; Holotype of *Physatta dromedarius*: casent0901429

Species: *Myrmicaria melanogaster*



Specimen Habitat Summary



Found most commonly in these habitats: 2 times found in tropical rainforest, 1 times found in 2nd growth wet forest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times search, 1 times light

Elevations: collected from 20 - 450 meters, 220 meters average

Type specimens: Lectotype of *Myrmicaria melanogaster*: casent0904469

Species: *Myrmicaria pseudoflava*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 20 m

Type specimens:

Species: *Myrmoteras arcoelinae*






Specimen Habitat Summary

Elevations: collected at 1610 m

Type specimens: Holotype of *Myrmoteras arcoelinae*: casent0910206; paratype of *Myrmoteras arcoelinae*: casent0903245

Species: *Myrmoteras bakeri*



Specimen Habitat Summary

Found most commonly in these habitats: 2 times found in mature wet forest, 0 times found in Rainforest, 1 times found in tropical rainforest, 1 times found in 2nd growth wet forest, 0 times found in Degraded coastal hill forest, 0 times found in lowland Dipterocarp

Collected most commonly using these methods or in the following microhabitats: 4 times Winkler

Elevations: collected from 304 - 2140 meters, 650 meters average

Type specimens:

Species: *Myrmoteras danieli*





Specimen Habitat Summary

Elevations: collected at 1540 m

Type specimens: Holotype of *Myrmoteras danieli*: casent0910209; paratype of *Myrmoteras danieli*: casent0903247

Species: *Myrmoteras mjoebergi*




Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 180 m

Type specimens:

Species: Myrmoteras my03



Specimen Habitat Summary



Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 24 m

Type specimens:

Species: Myrarium camillae



Specimen Habitat Summary



Found most commonly in these habitats: 18 times found in Rainforest, 1 times found in K mixed beach forest, 1 times found in wet forest, 1 times found in montane forest, 1 times found in lowland rainforest, 1 times found in Narra plantation with Abaca understory, 1 times found in seasonal rainforest, 1 times found in SL rainforest, 1 times found in 2nd growth wet forest, 1 times found in degraded rainforest

Collected most commonly using these methods or in the following microhabitats: 3 times Winkler, 3 times Berlesate, 1 times W53-5, C88 54-1, C88 53-5, 1 times QM Berlesate No. 697, 1 times 3 MaxiWinklers, 1 times berlesate no. 20, 1 times ground sample, 1 times berlesate ANIC 381

Elevations: collected from 6 - 1100 meters, 479 meters average

Type specimens: paratype of *Mystrium oculatum*: casent0104982; syntype of *Mystrium camillae*: casent0903790

Species: Nylanderia my01



Specimen Habitat Summary

Found most commonly in these habitats: 2 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 3 times Winkler

Elevations: collected from 57 - 106 meters, 96 meters average

Type specimens:

Species: Odontomachus latidens





Specimen Habitat Summary

Elevations: collected at 600 m

Type specimens: syntype of *Odontomachus latidens procerus*: casent0903998; syntype of *Odontomachus latidens sumatranaus*: casent0903999

Species: *Odontomachus monticola*



Specimen Habitat Summary



Found most commonly in these habitats: 4 times found in litter, rainforest, 2 times found in 2nd growth wet forest, 1 times found in rainforest, 1 times found in Schima-Castanopsis forest, 1 times found in mature wet forest

Collected most commonly using these methods or in the following microhabitats: 4 times pitfall, 3 times Winkler, 2 times search

Elevations: collected from 104 - 1350 meters, 403 meters average

Type specimens: Lectotype of *Odontomachus monticola*: casent0905746; paralectotype of *Odontomachus monticola*: casent0904000; syntype of *Odontomachus monticola formosae*: casent0900651; syntype of *Odontomachus monticola major*: casent0900657; syntype of *Odontomachus monticola formosae*: casent0907427; syntype of *Odontomachus monticola longi*: casent0907428; syntype of *Odontomachus monticola major*: casent0907429; syntype of *Odontomachus monticola punctulatus*: casent0907430

Species: *Odontomachus rixosus*



Specimen Habitat Summary


Found most commonly in these habitats: 23 times found in Rainforest, 0 times found in 1° forest, 1 times found in Rainforest (disturbed), 1 times found in secondary forest


Collected most commonly using these methods or in the following microhabitats: 21 times Berlesate, 1 times litter sample, 0 times Berlesate No. 330, 0 times Berlesate No. 40

Elevations: collected from 228 - 725 meters, 315 meters average

Type specimens: syntype of *Odontomachus rixosus*: casent0900656; syntype of *Odontomachus rixosus conifera*: casent0907432; syntype of *Odontomachus rixosus obscurior*: casent0907431

Species: *Odontomachus simillimus*





Specimen Habitat Summary


Found most commonly in these habitats: 113 times found in primary rainforest, 86 times found in secondary forest, 79 times found in Disturbed forest, 60 times found in dry forest, 53 times found in coastal forest, 30 times found in Pandanus & Casuarina forest, 27 times found in Rainforest, 20 times found in forest, 11 times found in forest edge, 8 times found in coastal scrub, ...

Collected most commonly using these methods or in the following microhabitats: 18 times hand collection, 23 times davis-sifting; incidental aspirated, 5 times 9 MaxiWinks, mixed samples, 4 times Malaise, 3 times winkler 2, 6 times aspirating, 4 times davis-sifting; incidental aspirated; PB bait, 6 times aspirating; PB & maple syrup bait, 1 times 9 MaxiWinks.mixed samples, 3 times aspirated, 3 times aspirating; PB bait, ...

Elevations: collected from 1 - 1828 meters, 245 meters average

Type specimens: Holotype of *Ponera pallidicornis*: casent0901350; syntype of *Odontomachus simillimus*: casent0900649, casent0900650; syntype of *Odontomachus haematodes fuscipennis*: casent0907433; type of *Odontomachus haematoda breviceps*: casent0901486

Species: *Odontoponera my02*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, litter, 1 times Winkler

Elevations: collected from 57 - 130 meters, 105 meters average

Type specimens:

Species: *Odontoponera my03*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in oil palm plantation, 1 times found in litter, rainforest


Collected most commonly using these methods or in the following microhabitats: 2 times Pitfall, litter, 1 times pitfall

Elevations: collected from 56 - 173 meters, 79 meters average

Type specimens:

Species: *Pachycondyla my01*





Specimen Habitat Summary



Found most commonly in these habitats: 4 times found in litter, rainforest, 2 times found in litter, forest, 2 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 6 times Winkler, 2 times pitfall

Elevations: collected from 20 - 150 meters, 109 meters average

Type specimens:

Species: *Pachycondyla my02*



Specimen Habitat Summary



Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 937 m

Type specimens:

Species: *Pachycondyla my06*






Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 180 m

Type specimens:

Species: *Paraparatrechina my01*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in litter, rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall

Elevations: collected from 104 - 225 meters, 144 meters average

Type specimens:

Species: *Paratopula longispina*



Specimen Habitat Summary

Elevations: collected at 1006 m

Type specimens:

Species: *Paratopula macta*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in slightly disturbed seasonal tropical forest

Collected most commonly using these methods or in the following microhabitats: 1 times Light Trap

Elevations: collected from 283 - 800 meters, 547 meters average

Type specimens: Holotype of *Paratopula macta*: casent0901740; paratype: casent0178560

Species: *Pheidole* (indet)



Specimen Habitat Summary

Found most commonly in these habitats: 184 times found in rainforest, 91 times found in tropical wet forest, 184 times found in tropical rainforest, 140 times found in cloud forest, 111 times found in montane wet forest, 95 times found in wet forest, 94 times found in mature wet forest, 97 times found in Bamboo forest, 71 times found in tropical moist forest, 48 times found in lowland rainforest, ...

Collected most commonly using these methods or in the following microhabitats: 464 times MiniWinkler, 291 times Winkler, 312 times search, 50 times davis-sifting; incidental aspirated, 72 times hand collecting, 4 times Winkler sifter, 144 times MaxiWinkler, 88 times Malaise, 37 times Blacklight, 17 times aspirating; PB & maple syrup bait, 82 times pan trap, ...

Elevations: collected from 1 - 17300 meters, 687 meters average

Type specimens:

Species: *Pheidole acantha*





Specimen Habitat Summary

Type specimens: paratype of *Pheidole acantha*: casent0901618, casent0901619

Species: *Pheidole angulicollis*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in alluvial forest

Collected most commonly using these methods or in the following microhabitats: 1 times soil core

Elevations: collected at 1000 m

Type specimens: paratype of *Pheidole angulicollis*: casent0901622, casent0901623

Species: *Pheidole annexa*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in tropical rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times search

Elevations: collected at 190 m

Type specimens: paratype of *Pheidole annexus*: casent0901626, casent0901627

Species: *Pheidole cariniceps*





0.5 mm

Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in tropical rainforest, 1 times found in mixed dipterocarp forest

Collected most commonly using these methods or in the following microhabitats: 1 times search

Elevations: collected from 130 - 185 meters, 166 meters average

Type specimens: paratype of *Pheidole cariniceps*: casent0901628, casent0901629

Species: *Pheidole clypeocornis*



Specimen Habitat Summary



Found most commonly in these habitats: 3 times found in tropical rainforest, 2 times found in mature wet forest, 1 times found in lowland rainforest

Collected most commonly using these methods or in the following microhabitats: 4 times Winkler, 1 times search

Elevations: collected from 190 - 515 meters, 332 meters average

Type specimens: paratype of *Pheidole clypeocornis*: casent0901661, casent0901662

Species: *Pheidole comata*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in 2nd growth wet forest, 1 times found in tropical rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times search, 1 times bait

Elevations: collected from 180 - 1000 meters, 490 meters average

Type specimens: Holotype of *Pheidole comata*: casent0901504

Species: *Pheidole deltea*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in 2nd growth wet forest, 1 times found in tropical rainforest, 1 times found in mature wet forest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times bait, 2 times Winkler

Elevations: collected from 190 - 558 meters, 305 meters average

Type specimens: paratype of *Pheidole deltea*: casent0901640, casent0901641

Species: *Pheidole elisae*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in lowland rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 558 m

Type specimens: Lectotype of *Pheidole elisae nenia*: casent0907973; paralectotype of *Pheidole elisae nenia*: casent0907974; syntype of *Pheidole elisae*: casent0904257

Species: *Pheidole hortensis*



Specimen Habitat Summary

Found most commonly in these habitats: 2 times found in rainforest, 1 times found in tropical rainforest

Collected most commonly using these methods or in the following microhabitats: 3 times Winkler

Elevations: collected from 28 - 210 meters, 88 meters average

Type specimens: syntype of *Pheidole hortensis*: casent0904271, casent0904272, casent0907975, casent0907976

Species: *Pheidole inornata*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in mixed dipterocarp forest

Elevations: collected from 500 - 1000 meters, 750 meters average

Type specimens: paratype of *Pheidole inornata*: casent0901630, casent0901631

Species: *Pheidole kikutai*






Specimen Habitat Summary

Elevations: collected at 900 m

Type specimens: paratype of *Pheidole kikutai*: casent0901648, casent0901649

Species: *Pheidole lokitae*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in tropical rainforest, 1 times found in 2nd growth wet forest, 1 times found in swamp forest

Collected most commonly using these methods or in the following microhabitats: 4 times Winkler

Elevations: collected from 180 - 1060 meters, 573 meters average

Type specimens: syntype of *Pheidole lokitae*: casent0907766, casent0907767

Species: *Pheidole lucioccipitalis*



Specimen Habitat Summary

Found most commonly in these habitats: 4 times found in tropical rainforest, 1 times found in 2nd growth wet forest, 1 times found in swamp forest, 1 times found in mature wet forest, 1 times found in peak of limestone pinnacle, scrubby veg., 1 times found in rainforest



Collected most commonly using these methods or in the following microhabitats: 6 times Winkler, 2 times search, 1 times L

Elevations: collected from 190 - 1398 meters, 465 meters average

Type specimens: paratype of *Pheidole lucioccipitalis*: casent0901644, casent0901645

Species: *Pheidole manukana*






0.2 mm

Specimen Habitat Summary

Type specimens: paratype of *Pheidole manukana*: casent0901632, casent0901633

Species: *Pheidole modiglianii*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in tropical rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times search

Elevations: collected at 190 m

Type specimens: syntype of *Pheidole modiglianii*: casent0904264, casent0904265

Species: *Pheidole montana*




Specimen Habitat Summary

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall trap

Elevations: collected from 500 - 1500 meters, 1000 meters average

Type specimens:

Species: *Pheidole my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest


Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 558 m

Type specimens:

Species: *Pheidole parvicorpus*





Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in swamp forest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 1060 m

Type specimens: paratype of *Pheidole parvicorpus*: casent0901646, casent0901647


Species: *Pheidole planidorsum*



Specimen Habitat Summary

Type specimens: paratype of *Pheidole planidorsum*: casent0901638, casent0901639

Species: *Pheidole poringensis*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected from 88 - 600 meters, 344 meters average

Type specimens: paratype of *Pheidole poringensis*: casent0901660

Species: *Philidris my01*






Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 13 m

Type specimens:

Species: *Philidris my02*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Fogging, canopy

Elevations: collected at 937 m

Type specimens:

Species: *Philidris my03*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 104 m

Type specimens:

Species: *Plagiolepis my01*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in litter, rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler, 1 times pitfall

Elevations: collected from 104 - 130 meters, 121 meters average

Type specimens:

Species: [Platythyrea my01](#)



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times hand

Elevations: collected at 525 m

Type specimens:

Species: [Platythyrea my02](#)



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest, 1 times found in litter, rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, litter

Elevations: collected from 173 - 180 meters, 175 meters average

Type specimens:

Species: [Polyrhachis danum](#)





Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Type specimens: paratype: casent0217423; paratype of Polyrhachis danum: casent0903418

Species: [Polyrhachis maryatiae](#)





Specimen Habitat Summary


Elevations: collected at 700 m

Type specimens: paratype: casent0217434

Species: *Polyrhachis montana*



Species: *Polyrhachis my01*






Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 225 m

Type specimens:

Species: *Polyrhachis my02*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 225 m

Type specimens:

Species: *Polyrhachis my03*





Specimen Habitat Summary



Found most commonly in these habitats: 2 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, litter

Elevations: collected from 104 - 225 meters, 144 meters average

Type specimens:

Species: *Ponera my01*



Specimen Habitat Summary



Found most commonly in these habitats: 2 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler, 1 times pitfall

Elevations: collected from 57 - 104 meters, 78 meters average

Type specimens:

Species: *Ponera my02*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in litter, rainforest


Collected most commonly using these methods or in the following microhabitats: 2 times winkler

Elevations: collected at 64 m

Type specimens:

Species: *Prenolepis my01*






Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 180 m

Type specimens:

Species: *Pristomyrmex my01*



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in rainforest, 1 times found in litter, rainforest, 1 times found in oil palm plantation

Collected most commonly using these methods or in the following microhabitats: 3 times Winkler

Elevations: collected from 104 - 180 meters, 116 meters average

Type specimens:

Species: *Probolomyrmex my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected from 104 - 500 meters, 236 meters average

Type specimens:

Species: *Proceratium my02*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 57 m

Type specimens:

Species: *Pseudolasius* my01



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 28 m

Type specimens:

Species: *Recurvidris* my01



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Pitfall, litter

Elevations: collected at 173 m

Type specimens:

Species: *Rhopalomastix* my01



Specimen Habitat Summary


Found most commonly in these habitats: 2 times found in rainforest

Elevations: collected at 525 m

Type specimens:

Species: *Solenopsis* my01





Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected from 88 - 130 meters, 116 meters average

Type specimens:

Species: *Solenopsis my02*



Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in ecotone, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Winkler

Elevations: collected from 24 - 64 meters, 50 meters average

Type specimens:

Species: *Stigmatomma my01*




Specimen Habitat Summary

Elevations: collected at 1140 m

Type specimens:

Species: *Stigmatomma my04*





0.2 mm

0.5 r


Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Elevations: collected at 180 m

Type specimens:

Species: *Strumigenys my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 57 m

Type specimens:

Species: *Strumigenys my02*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, forest, 1 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 3 times Winkler

Elevations: collected from 56 - 106 meters, 87 meters average

Type specimens:

Species: *Strumigenys my03*



Specimen Habitat Summary



Found most commonly in these habitats: 1 times found in litter, rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 106 m

Type specimens:

Species: *Strumigenys my04*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in litter, rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 64 m

Type specimens:

Species: *Tapinoma (indet)*



Specimen Habitat Summary


Found most commonly in these habitats: 45 times found in tropical moist forest, 22 times found in 2° lowland rainforest, 20 times found in lowland wet forest, 20 times found in mesophyll forest, 8 times found in rainforest, 19 times found in cloud forest, 1 times found in veg seems especially short, savannah, no pandanus immediately nearby; monomorium? nest in dead branch of living tree. Camponotus nest in dead stick on ground at base of same tree; 1 or 2 Tapinoma ne, 17 times found in montane wet forest, 1 times found in dry sclerophyll, 13 times found in 2° wet forest, ...

Collected most commonly using these methods or in the following microhabitats: 123 times Beating, 54 times MiniWinkler, 30 times search, 45 times Malaise, 16 times aspirating; PB bait, 6 times aspirating, 12 times aspirating; PB & maple syrup bait, 5 times davis-sifting; incidental aspirated, 7 times aspirated, 10 times Baiting, 2 times aspirated; PB bait, ...

Elevations: collected from 2 - 3064 meters, 581 meters average

Type specimens:

Species: *Tapinoma my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in tropical rainforest, 1 times found in litter, forest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected from 100 - 465 meters, 319 meters average

Type specimens:

Species: *Technomyrmex modiglianii*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in cloud forest

Collected most commonly using these methods or in the following microhabitats: 0 times Berlesate, 1 times search

Elevations: collected at 1575 m

Type specimens: syntype of *Technomyrmex modiglianii*: casent0905065; syntype of *Technomyrmex modiglianii javanus*: casent0909805

Species: *Technomyrmex my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 558 m

Type specimens:

Species: *Technomyrmex my02*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in rainforest


Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 104 m

Type specimens:

Species: *Tetraponera my01*





Specimen Habitat Summary



Found most commonly in these habitats: 1 times found in litter, rainforest, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, 1 times Winkler

Elevations: collected from 173 - 973 meters, 439 meters average

Type specimens:

Species: *Vollenhovia my01*



Specimen Habitat Summary


Found most commonly in these habitats: 1 times found in oil palm plantation, 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 2 times Pitfall, litter

Elevations: collected from 57 - 64 meters, 61 meters average

Type specimens:

Species: *Vollenhovia my02*



Specimen Habitat Summary



Found most commonly in these habitats: 1 times found in oil palm plantation

Collected most commonly using these methods or in the following microhabitats: 1 times pitfall, litter

Elevations: collected at 173 m

Type specimens:

Species: *Vollenhovia my03*



 0.2 mm

0.5 mm

Specimen Habitat Summary

Found most commonly in these habitats: 1 times found in rainforest

Collected most commonly using these methods or in the following microhabitats: 1 times Winkler

Elevations: collected at 104 m

Type specimens:

Copyright © 2002-2014.

AntWeb content is licensed under a Creative Commons Attribution License. We encourage use of AntWeb images. In print, each image must include attribution to its photographer and "from www.AntWeb.org" in the figure caption. For websites, images must be clearly identified as coming from www.AntWeb.org, with a backward link to the respective source page. See How to Cite AntWeb.

Antweb is funded from private donations and from grants from the National Science Foundation, DEB-0344731, EF-0431330 and DEB-0842395. c:3

 AntWeb is hosted and supported by The California Academy of Sciences.