


# CS 5522: Artificial Intelligence II

## Bayes' Nets: Inference



Instructor: Wei Xu



Ohio State University

[These slides were adapted from CS188 Intro to AI at UC Berkeley.]

# Bayes' Net Representation

- A directed, acyclic graph, one node per random variable
- A conditional probability table (CPT) for each node
  - A collection of distributions over  $X$ , one for each combination of parents' values  $P(X|a_1 \dots a_n)$
- Bayes' nets implicitly encode joint distributions
  - As a product of local conditional distributions
  - To see what probability a BN gives to a full assignment, multiply all the relevant conditionals together:

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$





# Example: Alarm Network

B	P(B)
+b	0.001
-b	0.999

E	P(E)
+e	0.002
-e	0.998

A	J	P(J A)
+a	+j	0.9
+a	-j	0.1
-a	+j	0.05
-a	-j	0.95

A	M	P(M A)
+a	+m	0.7
+a	-m	0.3
-a	+m	0.01
-a	-m	0.99




B	E	A	P(A B,E)
+b	+e	+a	0.95
+b	+e	-a	0.05
+b	-e	+a	0.94
+b	-e	-a	0.06
-b	+e	+a	0.29
-b	+e	-a	0.71
-b	-e	+a	0.001
-b	-e	-a	0.999

$$P(+b, -e, +a, -j, +m) =$$

$$P(+b)P(-e)P(+a|+b, -e)P(-j|+a)P(+m|+a) =$$


# Example: Alarm Network

B	P(B)
+b	0.001
-b	0.999



E	P(E)
+e	0.002
-e	0.998

A	M	P(M A)
+a	+m	0.7
+a	-m	0.3
-a	+m	0.01
-a	-m	0.99



A	J	P(J A)
+a	+j	0.9
+a	-j	0.1
-a	+j	0.05
-a	-j	0.95

B	E	A	P(A B,E)
+b	+e	+a	0.95
+b	+e	-a	0.05
+b	-e	+a	0.94
+b	-e	-a	0.06
-b	+e	+a	0.29
-b	+e	-a	0.71
-b	-e	+a	0.001
-b	-e	-a	0.999

$$\begin{aligned}
 P(+b, -e, +a, -j, +m) &= \\
 P(+b)P(-e)P(+a|+b, -e)P(-j|+a)P(+m|+a) &= \\
 0.001 \times 0.998 \times 0.94 \times 0.1 \times 0.7 &
 \end{aligned}$$

# Bayes' Nets

---

✓ Representation

✓ Conditional Independences

- Probabilistic Inference

- Enumeration (exact, exponential complexity)
- Variable elimination (exact, worst-case exponential complexity, often better)
- Probabilistic inference is NP-complete
- Sampling (approximate)

- Learning Bayes' Nets from Data

# Inference

- Inference: calculating some useful quantity from a joint probability distribution



- Examples:

- Posterior probability

$$P(Q|E_1 = e_1, \dots, E_k = e_k)$$

- Most likely explanation:

$$\operatorname{argmax}_q P(Q = q|E_1 = e_1 \dots)$$



# Inference by Enumeration

- General case:


- Evidence variables:  $E_1 \dots E_k = e_1 \dots e_k$
  - Query\* variable:  $Q$
  - Hidden variables:  $H_1 \dots H_r$
- $$\left. \begin{array}{l} E_1 \dots E_k = e_1 \dots e_k \\ Q \\ H_1 \dots H_r \end{array} \right\} \begin{array}{l} X_1, X_2, \dots X_n \\ \text{All variables} \end{array}$$

- We want:

$$P(Q|e_1 \dots e_k)$$

\* Works fine with multiple query variables, too


- Step 1: Select the entries consistent with the evidence



x	P(x)
-3	0.05
-1	0.25
0	0.07
1	0.2
5	0.01

2    0.15

- Step 2: Sum out H to get joint of Query and evidence



$$P(Q, e_1 \dots e_k) = \sum_{h_1 \dots h_r} \underbrace{P(Q, h_1 \dots h_r, e_1 \dots e_k)}_{X_1, X_2, \dots X_n}$$

- Step 3: Normalize

$$\times \frac{1}{Z}$$

$$Z = \sum_q P(Q, e_1 \dots e_k)$$

$$P(Q|e_1 \dots e_k) = \frac{1}{Z} P(Q, e_1 \dots e_k)$$

# Inference by Enumeration

- $P(W)$ ?
- $P(W \mid \text{winter})$ ?
- $P(W \mid \text{winter, hot})$ ?

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20



# Inference by Enumeration in Bayes' Net


- Given unlimited time, inference in BNs is easy
- Reminder of inference by enumeration by example:

$$P(B \mid +j, +m) \propto_B P(B, +j, +m)$$

$$= \sum_{e,a} P(B, e, a, +j, +m)$$

$$= \sum_{e,a} P(B)P(e)P(a|B, e)P(+j|a)P(+m|a)$$

$$= P(B)P(+e)P(+a|B, +e)P(+j|+a)P(+m|+a) + P(B)P(+e)P(-a|B, +e)P(+j|-a)P(+m|-a) \\ + P(B)P(-e)P(+a|B, -e)P(+j|+a)P(+m|+a) + P(B)P(-e)P(-a|B, -e)P(+j|-a)P(+m|-a)$$






# Inference by Enumeration vs. Variable Elimination


- Why is inference by enumeration so slow?

- You join up the whole joint distribution before you sum out the hidden variables




- Idea: interleave joining and marginalizing!

- Called “Variable Elimination”
- Still NP-hard, but usually much faster than inference by enumeration



# Factor Zoo



# Factor Zoo I


- Joint distribution:  $P(X,Y)$ 
  - Entries  $P(x,y)$  for all  $x, y$
  - Sums to 1
- Selected joint:  $P(x,Y)$ 
  - A slice of the joint distribution
  - Entries  $P(x,y)$  for fixed  $x$ , all  $y$
  - Sums to  $P(x)$
- Number of capitals = dimensionality of the table

$P(T, W)$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

$P(\text{cold}, W)$


T	W	P
cold	sun	0.2
cold	rain	0.3



# Factor Zoo II

- Single conditional:  $P(Y | x)$

- Entries  $P(y | x)$  for fixed  $x$ , a
- Sums to 1




$$P(W|cold)$$

T	W	P
cold	sun	0.4
cold	rain	0.6

- Family of conditionals:

$$P(X | Y)$$

- Multiple conditionals
- Entries  $P(x | y)$  for all  $x, y$
- Sums to  $|Y|$



$$P(W|T)$$

T	W	P
hot	sun	0.8
hot	rain	0.2
cold	sun	0.4
cold	rain	0.6

$$P(W|hot)$$

$$P(W|cold)$$

# Factor Zoo III

- Specified family:  $P(y \mid X)$


- Entries  $P(y \mid x)$  for fixed  $y$ , but for all  $x$
- Sums to ... who knows!

$$P(\text{rain} \mid T)$$

T	W	P
hot	rain	0.2
cold	rain	0.6





$$\left. \begin{array}{c} \\ \end{array} \right\} P(\text{rain} \mid \text{hot})$$

$$\left. \begin{array}{c} \\ \end{array} \right\} P(\text{rain} \mid \text{cold})$$



# Factor Zoo Summary

- In general, when we write  $P(Y_1 \dots Y_N \mid X_1 \dots X_M)$ 
  - It is a “factor,” a multi-dimensional array
  - Its values are  $P(y_1 \dots y_N \mid x_1 \dots x_M)$
  - Any assigned (=lower-case) X or Y is a dimension missing (selected) from the array





# Example: Traffic Domain


- Random Variables

- R: Raining
- T: Traffic
- L: Late for class!

$$P(L) = ?$$

$$= \sum_{r,t} P(r, t, L)$$

$$= \sum_{r,t} P(r)P(t|r)P(L|t)$$



$$P(R)$$

+r	0.1
-r	0.9

$$P(T|R)$$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$$P(L|T)$$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

# Inference by Enumeration: Procedural Outline

- Track objects called **factors**
- Initial factors are local CPTs (one per node)

$$P(R)$$

+r	0.1
-r	0.9

$$P(T|R)$$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$$P(L|T)$$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

- Any known values are selected
  - E.g. if we know  $L = +\ell$ , the initial factors are

$$P(R)$$

+r	0.1
-r	0.9


$$P(T|R)$$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$$P(+\ell|T)$$



+t	+l	0.3
-t	+l	0.1

- Procedure: Join all factors, then eliminate all hidden variables




# Operation 1: Join Factors

- First basic operation: **joining factors**
- Combining factors:
  - Just like a database join**
  - Get all factors over the joining variable
  - Build a new factor over the union of the variables involved
- Example: Join on R




- Computation for each entry: pointwise products  $\forall r, t : P(r, t) = P(r) \cdot P(t|r)$

# Example: Multiple Joins



# Example: Multiple Joins



$P(R)$

+r	0.1
-r	0.9

$P(T|R)$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$P(L|T)$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

Join R




$P(R, T)$

+r	+t	0.08
+r	-t	0.02
-r	+t	0.09
-r	-t	0.81

$P(L|T)$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9




Join T



$R, T, L$

$P(R, T, L)$

+r	+t	+l	0.024
+r	+t	-l	0.056
+r	-t	+l	0.002
+r	-t	-l	0.018
-r	+t	+l	0.027
-r	+t	-l	0.063
-r	-t	+l	0.081
-r	-t	-l	0.729



# Operation 2: Eliminate

- Second basic operation: **marginalization**
- Take a factor and sum out a variable
  - Shrinks a factor to a smaller one
  - A **projection** operation
- Example:


$$P(R, T)$$

+r	+t	0.08
+r	-t	0.02
-r	+t	0.09
-r	-t	0.81


sum  $R$


$$P(T)$$


+t	0.17
-t	0.83



# Multiple Elimination




# Thus Far: Multiple Join, Multiple Eliminate (= Inference by Enumeration)







# Marginalizing Early (= Variable Elimination)



# Marginalizing Early! (aka Variable Elimination)



# Traffic Domain



$$P(L) = ?$$

## ■ Inference by Enumeration

$$= \sum_t \sum_r \underbrace{P(L|t)P(r)P(t|r)}_{\text{Join on } r} \underbrace{\phantom{P(L|t)P(r)P(t|r)}}_{\text{Join on } t} \underbrace{\phantom{P(L|t)P(r)P(t|r)}}_{\text{Eliminate } r} \underbrace{\phantom{P(L|t)P(r)P(t|r)}}_{\text{Eliminate } t}$$

## ■ Variable Elimination

$$= \sum_t P(L|t) \underbrace{\sum_r P(r)P(t|r)}_{\text{Join on } r} \underbrace{\phantom{P(L|t)P(r)P(t|r)}}_{\text{Eliminate } r} \underbrace{\phantom{P(L|t)P(r)P(t|r)}}_{\text{Join on } t} \underbrace{\phantom{P(L|t)P(r)P(t|r)}}_{\text{Eliminate } t}$$

# Evidence

- If evidence, start with factors that select that evidence

- No evidence uses these initial factors:

$$P(R)$$

+r	0.1
-r	0.9

$$P(T|R)$$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$$P(L|T)$$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

- Computing  $P(L|+r)$ , the initial factors become:

$$P(+r)$$

+r	0.1
----	-----


$$P(T|+r)$$

+r	+t	0.8
+r	-t	0.2

$$P(L|T)$$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

- We eliminate all vars other than query + evidence



# Evidence II

- Result will be a selected joint of query and evidence
  - E.g. for  $P(L \mid +r)$ , we would end up with:

$$P(+r, L)$$

+r	+l	0.026
+r	-l	0.074


Normalize



$$P(L \mid +r)$$


+l	0.26
-l	0.74

- To get our answer, just normalize this!
- That 's it!





# General Variable Elimination

- Query:  $P(Q|E_1 = e_1, \dots, E_k = e_k)$
- Start with initial factors:
  - Local CPTs (but instantiated by evidence)
- While there are still hidden variables (not Q or evidence):
  - Pick a hidden variable H
  - Join all factors mentioning H
  - Eliminate (sum out) H
- Join all remaining factors and normalize



x	P(x)
-3	0.05
-1	0.25
0	0.07
1	0.2
5	0.01





$$\text{red stick} \times \text{blue square} = \text{purple square} \times \frac{1}{Z}$$

# Example

$$P(B|j, m) \propto P(B, j, m)$$

$P(B)$	$P(E)$	$P(A B, E)$	$P(j A)$	$P(m A)$
--------	--------	-------------	----------	----------




Choose A


$$P(A|B, E)$$

$$P(j|A)$$

$$P(m|A)$$



$$P(j, m, A|B, E)$$



$$P(j, m|B, E)$$

$P(B)$	$P(E)$	$P(j, m B, E)$
--------	--------	----------------

# Example

$P(B)$	$P(E)$	$P(j, m B, E)$
--------	--------	----------------


Choose E

$$\begin{array}{c} P(E) \\ P(j, m|B, E) \end{array} \xrightarrow{\times} P(j, m, E|B) \xrightarrow{\Sigma} P(j, m|B)$$

$P(B)$	$P(j, m B)$
--------	-------------

Finish with B

$$\begin{array}{c} P(B) \\ P(j, m|B) \end{array} \xrightarrow{\times} P(j, m, B) \xrightarrow{\text{Normalize}} P(B|j, m)$$






# Same Example in Equations

$$P(B|j, m) \propto P(B, j, m)$$

$P(B)$	$P(E)$	$P(A B, E)$	$P(j A)$	$P(m A)$
--------	--------	-------------	----------	----------



$$\begin{aligned}
 P(B|j, m) &\propto P(B, j, m) \\
 &= \sum_{e, a} P(B, j, m, e, a) \\
 &= \sum_{e, a} P(B)P(e)P(a|B, e)P(j|a)P(m|a) \\
 &= \sum_e P(B)P(e) \sum_a P(a|B, e)P(j|a)P(m|a) \\
 &= \sum_e P(B)P(e)f_1(B, e, j, m) \\
 &= P(B) \sum_e P(e)f_1(B, e, j, m) \\
 &= P(B)f_2(B, j, m)
 \end{aligned}$$

marginal can be obtained from joint by summing out

use Bayes' net joint distribution expression

use  $xy + xz = x^*(y+z)$

joining on  $a$ , and then summing out gives  $f_1$

use  $xy + xz = x^*(y+z)$

joining on  $e$ , and then summing out gives  $f_2$

All we are doing is exploiting  $uwy + uwz + uxy + uxz + vwy + vwz + vxy + vxz = (u+v)(w+x)(y+z)$  to improve computational efficiency!

# Another Variable Elimination Example

Query:  $P(X_3|Y_1 = y_1, Y_2 = y_2, Y_3 = y_3)$

Start by inserting evidence, which gives the following initial factors:

$$p(Z)p(X_1|Z)p(X_2|Z)p(X_3|Z)p(y_1|X_1)p(y_2|X_2)p(y_3|X_3)$$

Eliminate  $X_1$ , this introduces the factor  $f_1(Z, y_1) = \sum_{x_1} p(x_1|Z)p(y_1|x_1)$ , and we are left with:

$$p(Z)f_1(Z, y_1)p(X_2|Z)p(X_3|Z)p(y_2|X_2)p(y_3|X_3)$$

Eliminate  $X_2$ , this introduces the factor  $f_2(Z, y_2) = \sum_{x_2} p(x_2|Z)p(y_2|x_2)$ , and we are left with:

$$p(Z)f_1(Z, y_1)f_2(Z, y_2)p(X_3|Z)p(y_3|X_3)$$


Eliminate  $Z$ , this introduces the factor  $f_3(y_1, y_2, X_3) = \sum_z p(z)f_1(z, y_1)f_2(z, y_2)p(X_3|z)$ , and we are left:

$$p(y_3|X_3), f_3(y_1, y_2, X_3)$$

No hidden variables left. Join the remaining factors to get:

$$f_4(y_1, y_2, y_3, X_3) = P(y_3|X_3)f_3(y_1, y_2, X_3).$$


Normalizing over  $X_3$  gives  $P(X_3|y_1, y_2, y_3)$ .



Computational complexity critically depends on the largest factor being generated in this process. Size of factor = number of entries in table. In example above (assuming binary) all factors generated are of size 2 --- as they all only have one variable ( $Z$ ,  $Z$ , and  $X_3$  respectively).

# Variable Elimination Ordering

- For the query  $P(X_n | y_1, \dots, y_n)$  work through the following two different orderings as done in previous slide:  $Z, X_1, \dots, X_{n-1}$  and  $X_1, \dots, X_{n-1}, Z$ . What is the size of the maximum factor generated for each of the orderings?



- Answer:  $2^{n+1}$  versus  $2^2$  (assuming binary)
- In general: the ordering can greatly affect efficiency.

# VE: Computational and Space Complexity

---

- The computational and space complexity of variable elimination is determined by the largest factor
- The elimination ordering can greatly affect the size of the largest factor.
  - E.g., previous slide's example  $2^n$  vs. 2
- Does there always exist an ordering that only results in small factors?
  - No!

# Worst Case Complexity?

## ■ 3-SAT

$$(x_1 \vee x_2 \vee \neg x_3) \wedge (\neg x_1 \vee x_3 \vee \neg x_4) \wedge (x_2 \vee \neg x_2 \vee x_4) \wedge (\neg x_3 \vee \neg x_4 \vee \neg x_5) \wedge (x_2 \vee x_5 \vee x_7) \wedge (x_4 \vee x_5 \vee x_6) \wedge (\neg x_5 \vee x_6 \vee \neg x_7) \wedge (\neg x_5 \vee \neg x_6 \vee x_7)$$

$$P(X_i = 0) = P(X_i = 1) = 0.5$$

$$Y_1 = X_1 \vee X_2 \vee \neg X_3$$

...

$$Y_8 = \neg X_5 \vee X_6 \vee X_7$$

$$Y_{1,2} = Y_1 \wedge Y_2$$


...

$$Y_{7,8} = Y_7 \wedge Y_8$$

$$Y_{1,2,3,4} = Y_{1,2} \wedge Y_{3,4}$$

$$Y_{5,6,7,8} = Y_{5,6} \wedge Y_{7,8}$$

$$Z = Y_{1,2,3,4} \wedge Y_{5,6,7,8}$$



- If we can answer  $P(z)$  equal to zero or not, we answered whether the 3-SAT problem has a solution.
- Hence inference in Bayes' nets is NP-hard. No known efficient probabilistic inference in general.

# Polytrees

---

- A polytree is a directed graph with no undirected cycles
- For poly-trees you can always find an ordering that is efficient
  - Greedy search using heuristic cost functions, e.g.:
    - min-neighbors: #neighbors in current graph (smallest factor)
    - min-weight: weight (# values) of factor formed
    - etc.

# Bayes' Nets

---

- ✓ Representation

- ✓ Conditional Independences

- Probabilistic Inference

- ✓ Enumeration (exact, exponential complexity)

- ✓ Variable elimination (exact, worst-case exponential complexity, often better)

- ✓ Inference is NP-complete

- Sampling (approximate)

- Learning Bayes' Nets from Data