

TAPENTADOL -> obesity

For clinical-collaborator outreach.

Why this hypothesis is real

Drug-target convergence on the disease scores **0.96** (noisy-OR over OT target-disease associations). This pair is fully novel relative to all known drug-indication assignments in ChEMBL. Lead target is well expressed in the disease-relevant tissue (adipose tissue). Orthologous-gene model-organism evidence is strong (phylo_score 0.71, n=7). Reactome co-membership reinforces the indirect mechanism with **4** shared specific pathways.

Why now (IP runway + literature footprint)

Loss-of-exclusivity year **2029**, with 7 generic ANDAs already on the market.

Clinical opportunity

US population: approximately **108,000,000** patients -- a mass-market opportunity. (source: CDC NHANES adult obesity prevalence)

What we propose

BioBix (Department of Mathematical Modelling, Statistics and Bioinformatics, Ghent University) brings bioinformatics support to the collaboration:

- Target validation across blood / immune / cancer multi-omics datasets
- Biomarker candidate identification from public + private cohorts
- Patient stratification analytics for trial-enrichment design
- Mechanism-of-action mining via systems-biology priors
- Quick-turn evidence syntheses for IIT and partnered programs

Proposed collaboration model:

- Investigator-initiated trial framework with the substance owner
- BioBix as bioinformatics analytics partner (target/biomarker/biostatistics)
- Joint publication strategy; data-sharing agreement gated on IP terms

Next step

A 30-minute scoping call to align on feasibility. We would come prepared with a brief mechanistic dossier, an annotated literature pull, and a pilot biomarker shortlist for the disease.

Wim Van Criekinge, Professor (PhD, h-index 73)

Department of Mathematical Modelling, Statistics and Bioinformatics, Ghent University

Wim.VanCriekinge@UGent.be