
k+3=16
k+2=15
k+1=14
Author(s) and Year
Vaccinated
Control
TB+
TB−
TB+
TB−
Log[RR] [95% CI]
k=13 Aronson, 1948 4 119 11 128 −0.89 [−2.01, 0.23]
12 Ferguson & Simes, 1949 6 300 29 274 −1.59 [−2.45, −0.72]
11 Rosenthal et al, 1960 3 228 11 209 −1.35 [−2.61, −0.08]
10 Hart & Sutherland, 1977 62 13536 248 12619 −1.44 [−1.72, −1.16]
9 Frimodt−Moller et al, 1973 33 5036 47 5761 −0.22 [−0.66, 0.23]
8 Stein & Aronson, 1953 180 1361 372 1079 −0.79 [−0.95, −0.62]
7 Vandiviere et al, 1973 8 2537 10 619 −1.62 [−2.55, −0.70]
6 TPT Madras, 1980 505 87886 499 87892 0.01 [−0.11, 0.14]
5 Coetzee & Berjak, 1968 29 7470 45 7232 −0.47 [−0.94, −0.00]
4 Rosenthal et al, 1961 17 1699 65 1600 −1.37 [−1.90, −0.84]
3 Comstock et al, 1974 186 50448 141 27197 −0.34 [−0.56, −0.12]
2 Comstock & Webster, 1969 5 2493 3 2338 0.45 [−0.98, 1.88]
1 Comstock et al, 1976 27 16886 29 17825 −0.02 [−0.54, 0.51]
0
−1 Random−Effects Model −0.71 [−1.07, −0.36]
−2
Axis line at −2.
−3 −2 −1 0 1 2
Log Risk Ratio
(use par("mgp") to adjust the margin line for the axis title, axis labels, and axis line; use par("tcl") to adjust the length of the tick marks)
−9.5 −8 −6 −4.5
ilab.xpos values
−3 alim (or range(at)) 2
Study labels to the right of xlim[1].
Annotations to the left of xlim[2].
−16
xlim
7