

Phase structure of the Quark-Meson model

Rui Wen,¹ Yong-rui Chen,¹ and Wei-jie Fu^{1,*}

¹*School of Physics, Dalian University of Technology, Dalian, 116024, P.R. China*

PACS numbers: 11.30.Rd, 11.10.Wx, 05.10.Cc, 12.38.Mh

I. INTRODUCTION

A

II. 2 FLAVOR QUARK MESON MODEL

The effective action of the 2 flavor quark-meson model reads [1]

$$\Gamma_k[\Phi] = \int_x \{ Z_{q,k} \bar{q} (\gamma_\mu \partial_\mu - \gamma_0 \mu) q + \frac{1}{2} Z_{\phi,k} (\partial_\mu \phi)^2 + h_k \bar{q} (T^0 \sigma + i \gamma_5 \vec{T} \cdot \vec{\pi}) q + V_k(\rho) - c\sigma \} \quad (1)$$

with the integral $\int_x = \int_0^{1/T} d_{x_0} \int d^3x$. T and μ denote the temperature and quark chemical potential, respectively. The meson/quark wave function $Z_{\phi,k}/Z_{q,k}$ are scale and field dependent. Moreover, in this work we ignore the longitudinal and transversal splitting of Z_k , i.e., Z_k^\parallel and Z_k^\perp , respectively, and an interesting discussion is presented in Ref. [2]. T^0 and \vec{T} are generators of $SU(N_f = 2)$ with $\text{Tr}(T^i T^j) = \frac{1}{2} \delta^{ij}$ and $T^0 = \frac{1}{\sqrt{N_f}} \mathbb{1}_{N_f \times N_f}$. Quark and meson interaction is encoded in Yukawa coupling h_k in Eq. (1). The effective potential $V_k(\rho)$ is scale and field dependent and chirally invariant with chirally invariant variable $\rho = \frac{1}{2} \phi^2$. The linear sigma term $-c\sigma$ explicitly breaks the chiral symmetry.

Then with the Wetterich equation

$$\partial_t \Gamma_k[\Phi] = \frac{1}{2} \text{Tr} G_{\phi\phi}[\Phi] \partial_t R_k^\phi - \text{Tr} G_{q\bar{q}}[\Phi] \partial_t R_k^q \quad (2)$$

with the regulators

The renormalized dimensionless meson and quark

$$\begin{aligned} \partial_t V_k(\rho) = & \frac{k^4}{4\pi^2} \{ (N_f^2 - 1) l_0^B(m_{\pi,k}, \eta_{\phi,k}; T) \\ & + l_0^B(m_{\sigma,k}, \eta_{\phi,k}; T) \\ & - 4N_c N_f l_0^F(m_{q,k}, \eta_{q,k}; T, \mu) \} \end{aligned} \quad (6)$$

here $l_0^{B/F}$ are threshold functions which are defined as

$$l_0^B(\bar{m}_{\phi,k}^2, \eta_{\phi,k}) = \frac{2}{3} \frac{1}{\sqrt{1 + \bar{m}_{\phi,k}^2}} \left(1 - \frac{\eta_{\phi,k}}{5} \right) \left(\frac{1}{2} + n_B(\bar{m}_{\phi,k}^2, T) \right) \quad (7)$$

$$l_0^F(\bar{m}_{q,k}^2, \eta_{q,k}) = \frac{1}{3} \frac{1}{\sqrt{1 + \bar{m}_{q,k}^2}} \left(1 - \frac{\eta_{q,k}}{4} \right) (1 - n_F(E - \mu) - n_F(E + \mu)) \quad (8)$$

$$\eta_{\phi,k} = \frac{1}{6\pi^2} \left\{ \frac{4}{k^2} \rho (V'_k(\rho))^2 \right\} \quad (9)$$

III. RESULTS

A. phase diagram at chiral limit

IV. SUMMARY AND DISCUSSIONS

In this work, we have studied

ACKNOWLEDGMENTS

The work was supported by the National Natural Science Foundation of China under Contracts Nos. 11775041.

-
- [1] W.-j. Fu and J. M. Pawłowski, Phys. Rev. **D92**, 116006 (2015), arXiv:1508.06504 [hep-ph].
 [2] S. Yin, R. Wen, and W.-j. Fu, (2019), arXiv:1907.10262 [hep-ph].

masses are obtained by

* wjfu@dlut.edu.cn

$$\bar{m}_\pi = \frac{V'_k(\rho)}{k^2 Z_{\phi,k}} \quad (3)$$

$$\bar{m}_\sigma = \frac{V'_k(\rho) + 2\rho V''_k(\rho)}{k^2 Z_{\phi,k}} \quad (4)$$

$$\bar{m}_q = \frac{h_k^2 \rho}{2k^2 Z_{q,k}^2} \quad (5)$$