

Phase structure of the Quark-Meson model

Rui Wen,¹ Yong-rui Chen,¹ and Wei-jie Fu^{1,*}

¹*School of Physics, Dalian University of Technology, Dalian, 116024, P.R. China*

PACS numbers: 11.30.Rd, 11.10.Wx, 05.10.Cc, 12.38.Mh

I. INTRODUCTION

A

$$\begin{aligned} \partial_t V_k(\rho) = & \frac{k^4}{4\pi^2} \{ (N_f^2 - 1) l_0^B(m_{\pi,k}, \eta_{\phi,k}; T) \\ & + l_0^B(m_{\sigma,k}, \eta_{\phi,k}; T) \\ & - 4N_c N_f l_0^F(m_{q,k}, \eta_{q,k}; T, \mu) \} \end{aligned} \quad (6)$$

II. 2 FLAVOR QUARK MESON MODEL

The effective action of the 2 flavor quark-meson model reads

$$\begin{aligned} \Gamma_k[\Phi] = & \int_x \{ Z_{q,k} \bar{q} (\gamma_\mu \partial_\mu - \gamma_0 \mu) q + \frac{1}{2} Z_{\phi,k} (\partial_\mu \phi)^2 \\ & + k_k \bar{q} (T^0 \sigma + i \gamma_5 T \cdot \pi) q + V_k(\rho) - c\sigma \} \end{aligned} \quad (1)$$

where $\rho = \frac{1}{2}\phi^2$ is chirally invariant variable

The dimensionless meson and quark masses are obtained by

$$\bar{m}_\pi = \frac{V'_k(\rho)}{k^2 Z_{\phi,k}} \quad (2)$$

$$\bar{m}_\sigma = \frac{V'_k(\rho) + 2\rho V''_k(\rho)}{k^2 Z_{\phi,k}} \quad (3)$$

$$\bar{m}_q = \frac{h_k^2 \rho}{2k^2 Z_{q,k}^2} \quad (4)$$

with the Wetterich equation

$$\partial_t \Gamma_k[\Phi] = \frac{1}{2} \text{Tr} G_{\phi\phi}[\Phi] \partial_t R_k^\phi - \text{Tr} G_{q\bar{q}}[\Phi] \partial_t R_k^q \quad (5)$$

the regulator

* wjfu@dlut.edu.cn

here $l_0^{B/F}$ are threshold functions which are defined as

$$\begin{aligned} l_0^B(\bar{m}_{\phi,k}^2, \eta_{\phi,k}) = & \frac{2}{3} \frac{1}{\sqrt{1 + \bar{m}_{\phi,k}^2}} \left(1 - \frac{\eta_{\phi,k}}{5} \right) \\ & \left(\frac{1}{2} + n_B(\bar{m}_{\phi,k}^2, T) \right) \end{aligned} \quad (7)$$

$$\begin{aligned} l_0^F(\bar{m}_{q,k}^2, \eta_{q,k}) = & \frac{1}{3} \frac{1}{\sqrt{1 + \bar{m}_{q,k}^2}} \left(1 - \frac{\eta_{q,k}}{4} \right) \\ & \left(1 - n_F(E - \mu) - n_F(E + \mu) \right) \end{aligned} \quad (8)$$

$$\eta_{\phi,k} = \frac{1}{6\pi^2} \left\{ \frac{4}{k^2} \rho (V'_k(\rho))^2 \right\} \quad (9)$$

III. RESULTS

A. phase diagram at chiral limit

IV. SUMMARY AND DISCUSSIONS

In this work, we have studied

ACKNOWLEDGMENTS

The work was supported by the National Natural Science Foundation of China under Contracts Nos. 11775041.