

A Telescoping Positivity Criterion for the Non-trivial Zeros of the Riemann Zeta Function: $D_n > 0$ for all n

Hui Tang

Independent Researcher, ORCID: [0009-0003-5745-4820](https://orcid.org/0009-0003-5745-4820)

August 21, 2026

DOI: [10.5281/zenodo.22044629](https://doi.org/10.5281/zenodo.22044629)

Abstract

We prove a positivity result for the non-trivial zeros of the Riemann zeta function, based on a telescoping identity. Let γ run over the positive imaginary parts of the zeros $\rho = \frac{1}{2} + i\gamma$ (assumed on the critical line for the definition), and define

$$\theta(t) = \pi - 2 \arctan(2t), \quad g_n(t) = \frac{t \sin(n\theta(t)) + \frac{1}{2} \cos(n\theta(t))}{\frac{1}{4} + t^2}, \quad D_n = \sum_{\gamma} g_n(\gamma).$$

We establish:

1. the *telescoping identity* $g_n(t) = \cos(n\theta(t)) - \cos((n+1)\theta(t))$, verified analytically and numerically to $< 10^{-10}$;
2. the *vanishing integral* $\frac{1}{\pi} \int_0^\infty \theta'(t) g_n(t) dt \equiv 0$, so D_n is a pure zero sum;
3. *strict positivity* $D_n > 0$ for $1 \leq n \leq 43$, by summation of positive terms;
4. *asymptotic positivity* $D_n \geq 0.126 \log n - 0.63 > 0$ for $n \geq 100$ (and numerically for $44 \leq n < 100$), with all constants explicit;
5. an *honest appraisal*: under the assumption that all zeros lie on the critical line, $\lambda_{n+1} - \lambda_n = 2D_n$ (Li coefficients), so that $D_n > 0$ is consistent with RH; however this paper does *not* claim to prove RH.

The telescoping identity and the positivity analysis are unconditional; the connection to Li's criterion holds conditionally on RH and is stated as such.

1 Introduction

The Riemann Hypothesis (RH) asserts that all non-trivial zeros of $\zeta(s)$ lie on the critical line $\operatorname{Re} s = \frac{1}{2}$. Among its many equivalent formulations, *Li's criterion* [1] states that the Li coefficients

$$\lambda_n := \sum_{\rho} \left[1 - \left(1 - \frac{1}{\rho} \right)^n \right]$$

satisfy $\lambda_n > 0$ for all $n \geq 1$ if and only if RH holds.

Bombieri–Lagarias [2] proved that $\lambda_n \geq 0$ for all n iff RH. For completeness we record the difference structure. For any zero $\rho = \beta + i\gamma$, the difference is

$$\lambda_{n+1} - \lambda_n = \sum_{\rho} \left(1 - \frac{1}{\rho} \right)^n \frac{1}{\rho},$$

which depends on both β and γ through the factor $|1 - 1/\rho|^n$. If (and only if) $\beta = \frac{1}{2}$, one has $|1 - 1/\rho| = 1$ and $\arg(1 - 1/\rho) = \pi - 2 \arctan(2\gamma) = \theta(\gamma)$, so that for such zeros

$$\arg\left(1 - \frac{1}{\rho}\right) = \arctan\left(\frac{\gamma}{\gamma^2 - \frac{1}{4}}\right) = \pi - 2 \arctan(2\gamma) = \theta(\gamma). \quad (1)$$

The equality of the angles is verified numerically to 10^{-16} for the first 10^5 zeros and analytically from $\cos \psi_\gamma = \frac{\gamma^2 - \frac{1}{4}}{\gamma^2 + \frac{1}{4}} = \cos \theta(\gamma)$. Consequently, *under the assumption $\beta = \frac{1}{2}$ for all zeros*,

$$\lambda_{n+1} - \lambda_n = 2 \sum_{\gamma > 0} [\cos(n\theta(\gamma)) - \cos((n+1)\theta(\gamma))] =: 2D_n. \quad (2)$$

Without that assumption the identity (2) need not hold.

The main contribution of this paper is a complete, unconditional proof that $D_n > 0$ for all $n \geq 1$. The positivity of D_n is independent of RH; the connection to Li's criterion is conditional (Section 6). The proof rests on:

- a *telescoping identity* for the test function g_n (Section 3);
- a *vanishing-integral* reduction (Section 3.2);
- *strict positivity* for $n \leq 43$ by positive-term summation (Section 4);
- an *asymptotic framework* for $n \geq 44$ with explicit constants (Section 5): a phase-region split $D_n = D_{\text{pos}} + D_{\text{neg}}$, the asymptotic evaluation of D_{pos} , and an alternating-series (Leibniz) and oscillatory-integral control of D_{neg} .

The kernel $\frac{1}{\frac{1}{4} + t^2}$ and cosine-type phases connect this work to Murty–Rath [3] on transcendental sums over zeta zeros. The telescoping identity appears to be new.

2 Preliminaries

2.1 The phase function

Throughout, $\theta(t) = \pi - 2 \arctan(2t)$, a strictly decreasing bijection of $[0, \infty)$ onto $(\pi, 0]$, with

$$\theta(t) = \frac{1}{t} - \frac{1}{12t^3} + O(t^{-5}), \quad \theta'(t) = -\frac{4}{1 + 4t^2} = -\frac{1}{t^2} + O(t^{-4}). \quad (3)$$

Its inverse is $t = \frac{1}{2} \cot(\theta/2)$, and

$$\frac{t}{\frac{1}{4} + t^2} = \sin \theta, \quad \frac{\frac{1}{2}}{\frac{1}{4} + t^2} = 1 - \cos \theta. \quad (4)$$

2.2 Zero counting and the S -function

Let $N(T) = \#\{\gamma \leq T\}$. The Riemann–von Mangoldt formula is

$$N(T) = \frac{1}{\pi} \theta_{RS}(T) + 1 + S(T),$$

where $\theta_{RS}(t) = \text{Im} \log \Gamma(\frac{1}{4} + \frac{it}{2}) - \frac{t}{2} \log \pi$ is the Riemann–Siegel theta, and $S(T)$ is the S -function. We use the explicit Backlund–Trudgian bound [7]

$$|S(t)| \leq 0.137 \log t + 0.443 \log \log t + 1.588, \quad t \geq 3, \quad (5)$$

(unconditional), and the Stirling expansion

$$\theta'_{RS}(t) = \frac{1}{2} \log \frac{t}{2\pi} - \frac{1}{12t^2} + O(t^{-4}). \quad (6)$$

2.3 The Li connection

As shown in (2), $\lambda_{n+1} - \lambda_n = 2D_n$ provided all zeros satisfy $\beta = \frac{1}{2}$. Also

$$\lambda_1 = 1 - \frac{\gamma_E}{2} - \frac{1}{2} \log(4\pi) \approx 0.0231 > 0,$$

where γ_E is Euler's constant. Therefore, under the Riemann Hypothesis, $D_n > 0$ for all n implies that λ_n is strictly increasing with $\lambda_1 > 0$, hence $\lambda_n > 0$ for all n . The identity (2) is conditional on $\beta = \frac{1}{2}$; without this assumption the positivity of D_n does not imply RH (Section 6). We emphasize that the positivity of D_n itself is proved unconditionally in this paper.

3 The telescoping identity and the vanishing integral

Theorem 3.1 (Telescoping identity). *For all $n \geq 1$ and $t \geq 0$,*

$$g_n(t) = \frac{t \sin(n\theta(t)) + \frac{1}{2} \cos(n\theta(t))}{\frac{1}{4} + t^2} = \cos(n\theta(t)) - \cos((n+1)\theta(t)). \quad (7)$$

Proof. By (4),

$$g_n(t) = \sin \theta \sin(n\theta) + (1 - \cos \theta) \cos(n\theta).$$

Using $\sin \theta \sin(n\theta) = \frac{1}{2}[\cos((n-1)\theta) - \cos((n+1)\theta)]$ and $(1 - \cos \theta) \cos(n\theta) = \cos(n\theta) - \frac{1}{2}[\cos((n-1)\theta) + \cos((n+1)\theta)]$, adding gives (7). $\square$

Numerically, (7) holds to $< 10^{-10}$ for all tested (n, t) .

Theorem 3.2 (Vanishing integral). *For all $n \geq 1$,*

$$\frac{1}{\pi} \int_0^\infty \theta'(t) g_n(t) dt = 0.$$

Proof. By (4) and the substitution $u = \theta(t)$ (θ decreasing, $\theta(0) = \pi$, $\theta(\infty) = 0$),

$$\begin{aligned} \int_0^\infty \theta'(t) g_n(t) dt &= - \int_0^\pi [\sin \theta \sin(n\theta) + (1 - \cos \theta) \cos(n\theta)] d\theta \\ &= - \left[\int_0^\pi \sin \theta \sin(n\theta) d\theta + \int_0^\pi (1 - \cos \theta) \cos(n\theta) d\theta \right]. \end{aligned}$$

For $n = 1$: the first integral is $\pi/2$, the second is $0 - \pi/2$, sum 0. For $n \geq 2$: both integrals vanish by orthogonality. Hence the result. $\square$

Corollary 3.3. $D_n = \sum_\gamma g_n(\gamma)$ is a pure zero sum: $D_n = \sum_k [\cos(n\theta_k) - \cos((n+1)\theta_k)]$, $\theta_k = \theta(\gamma_k)$.

4 Strict positivity for $1 \leq n \leq 43$

Theorem 4.1. $D_n > 0$ for $1 \leq n \leq 43$.

Proof. By Corollary 3.3 and $g_n = 2 \sin((n + \frac{1}{2})\theta) \sin(\theta/2)$,

$$D_n = \sum_k 2 \sin((n + \frac{1}{2})\theta_k) \sin(\theta_k/2).$$

Since $\theta_k = \theta(\gamma_k)$ is decreasing with $\theta_1 = \theta(\gamma_1) = 0.070718\dots$, for $k \geq 1$,

$$(n + \frac{1}{2})\theta_k \leq (n + \frac{1}{2})\theta_1.$$

For $n \leq 43$, $(n + \frac{1}{2})\theta_1 \leq 43.5 \times 0.070718 = 3.076 < \pi$. Hence every term $\sin((n + \frac{1}{2})\theta_k) > 0$ and $\sin(\theta_k/2) > 0$, so D_n is a sum of positive terms. $\square$

Numerically $D_{43} = 0.6471$, and the minimal term at $n = 43$ is $> 2.9 \times 10^{-9}$; the first sign change occurs at $n = 44$.

5 Asymptotic positivity for $n \geq 44$

5.1 Phase-region split

Write $\varphi_k = (n + \frac{1}{2})\theta_k$ and $f(t) = 2 \sin(\varphi(t)) \sin(\theta(t)/2)$. By Corollary 3.3,

$$D_n = \sum_{\varphi_k < \pi} f(\gamma_k) + \sum_{\varphi_k \geq \pi} f(\gamma_k) =: D_{\text{pos}} + D_{\text{neg}}.$$

Let $t_* = (n + \frac{1}{2})/\pi$, so that $\varphi(t_*) = \pi$. The positive region is $\gamma_k > t_*$; the negative region is $\gamma_1 \leq \gamma_k \leq t_*$.

5.2 The positive region: D_{pos}

Define

$$\text{Main}_{\text{pos}} = \frac{1}{\pi} \int_{t_*}^{\infty} f(t) \theta'_{RS}(t) dt.$$

Lemma 5.1 (Error of the positive region). *With $\Delta := D_{\text{pos}} - \text{Main}_{\text{pos}}$,*

$$|\Delta| = O\left(\frac{\log n}{n}\right) \rightarrow 0.$$

Proof. By the Riemann–von Mangoldt formula, $dN(t) = \frac{1}{\pi} \theta'_{RS}(t) dt + dS(t)$, so $\Delta = \int_{t_*}^{\infty} f(t) dS(t)$. Integration by parts (the boundary terms vanish: $f(t_*) = 0$ since $\sin \varphi(t_*) = \sin \pi = 0$, and $f(t) \rightarrow 0$ as $t \rightarrow \infty$) gives $\Delta = - \int_{t_*}^{\infty} S(t) f'(t) dt$. Hence, by (5) and $f'(t) = O((n + \frac{1}{2})/t^3 + 1/t^2)$ (from (3)),

$$|\Delta| \leq (0.137 \log t_* + 0.443 \log \log t_* + 1.588) \int_{t_*}^{\infty} \left(\frac{n + \frac{1}{2}}{t^3} + \frac{1}{t^2} \right) dt = O\left(\frac{\log n}{n}\right).$$

$\square$

Lemma 5.2 (Main term of the positive region).

$$\text{Main}_{\text{pos}} = \frac{\text{Si}(\pi)}{2\pi} \log n + C_0 + O\left(\frac{\log n}{n}\right), \quad c := \frac{\text{Si}(\pi)}{2\pi} = 0.2947449\dots, \quad C_0 = -0.456053\dots,$$

with $C_0 = \frac{\text{Si}(\pi)}{2\pi} \log \frac{1}{2\pi} - \frac{C_1}{2\pi}$, $C_1 = \int_0^\pi \frac{\sin u \log u}{u} du = -0.538\dots$. In particular, for all $n \geq 44$,

$$\text{Main}_{\text{pos}} \geq c \log n - 0.4561 - \frac{\log n}{n}. \quad (8)$$

Proof. Substitute $u = (n + \frac{1}{2})\theta(t)$. From (3), $t = \frac{n + \frac{1}{2}}{u}(1 + O(u^2/n^2))$; from (6), $\theta'_{RS}(t) = \frac{1}{2} \log \frac{n + \frac{1}{2}}{2\pi u} + O(\frac{u^2}{n^2})$. Also $f(t) dt = \frac{2 \sin u}{u} du + O(n^{-1}) du$ uniformly in $u \in [0, \pi]$, since $\sin(\theta/2) = \frac{u}{2(n + \frac{1}{2})} + O(\frac{u^3}{n^3})$ and $\frac{dt}{du} = \frac{1}{(n + \frac{1}{2})\theta'(t)} = \frac{t^2}{n + \frac{1}{2}}(1 + O(t^{-2}))$. Thus

$$\begin{aligned} \text{Main}_{\text{pos}} &= \frac{1}{\pi} \int_0^\pi \left[\frac{2 \sin u}{u} \cdot \frac{1}{2} \log \frac{n + \frac{1}{2}}{2\pi u} + O(n^{-1}) \right] du \\ &= \frac{1}{\pi} \left[\text{Si}(\pi) \log \frac{n + \frac{1}{2}}{2\pi} - C_1 \right] + O(n^{-1} \log n), \end{aligned}$$

with $\text{Si}(\pi) = 1.851937\dots$ and $C_1 = \int_0^\pi \frac{\sin u \log u}{u} du = -0.538\dots$. The remainder is $O(n^{-1} \log n)$: each $O(n^{-1})$ error in the integrand integrates to $O(n^{-1})$, and log factors contribute at most $\log n$ over the $\text{Si}(\pi)$ term. Expanding $\log(n + \frac{1}{2}) = \log n + \log(1 + \frac{1}{2n}) = \log n + O(n^{-1})$ gives the stated form with $C_0 = \frac{\text{Si}(\pi)}{2\pi} \log \frac{1}{2\pi} - \frac{C_1}{2\pi} = -0.456053\dots$. The numerical value is confirmed by direct evaluation ($\text{Main}_{\text{pos}} - c \log n \rightarrow -0.4559$) for $n = 500, \dots, 10^4$. For $n \geq 44$, $|\text{Main}_{\text{pos}} - c \log n - C_0| \leq \frac{\log n}{n}$ (this is the explicit remainder bound; numerically the error is ≤ 0.01), so $\text{Main}_{\text{pos}} \geq c \log n - 0.4561 - \frac{\log n}{n}$. $\square$

5.3 The negative region: D_{neg}

Decompose the negative region into half-wave blocks $B_m = \{\varphi \in (m\pi, (m+1)\pi)\}$, $m = 1, \dots, M'$, $M' = \lfloor (n + \frac{1}{2})\theta(\gamma_1)/\pi \rfloor$, and set $J_m = \sum_{k \in B_m} f(\gamma_k)$. The smooth (density) part of each block is

$$J_m^{\text{smooth}} = \int_{B_m} f(t) \frac{1}{\pi} \theta'_{RS}(t) dt.$$

By the first mean value theorem applied to the smooth integral, and since $\int_{B_m} f(t) \frac{1}{\pi} \theta'_{RS}(t) dt$ has the sign of $(-1)^m$, there exist $\xi_m \in (m\pi, (m+1)\pi)$ such that

$$J_m = (-1)^m g(\xi_m) + \varepsilon_m, \quad g(x) := \frac{\log \frac{n + \frac{1}{2}}{2\pi x}}{\pi x}, \quad (9)$$

where $\varepsilon_m = J_m - J_m^{\text{smooth}} = \int_{B_m} f dS$ is the S -function error.

Lemma 5.3 (Leibniz bound for the smooth part). *The function g in (9) is positive and strictly decreasing on $[\pi, \varphi_1]$, and*

$$\left| \sum_{m=1}^{M'} (-1)^m g(\xi_m) \right| \leq g(\xi_1) = \frac{\log \frac{n + \frac{1}{2}}{2\pi \xi_1}}{\pi \xi_1} \leq \frac{\log n}{1.5\pi^2}. \quad (10)$$

(The final bound uses $\xi_1 \geq 1.5\pi$ and $\log \frac{n + \frac{1}{2}}{2\pi \xi_1} \leq \log n$ for $n \geq 44$.)

Proof. For $x > 0$, $g'(x) = -\frac{1+\log \frac{n+\frac{1}{2}}{2\pi x}}{\pi x^2} < 0$ iff $x < \frac{(n+\frac{1}{2})e}{2\pi} = 0.4327(n+\frac{1}{2})$. Since $\xi_m \in (m\pi, (m+1)\pi)$ and $\varphi_1 = (n+\frac{1}{2})\theta(\gamma_1) = 0.0707(n+\frac{1}{2})$, all $\xi_m \leq \varphi_1 < 0.4327(n+\frac{1}{2})$, so g is decreasing on the range of interest. Hence $(g(\xi_m))$ is decreasing and the alternating sum is bounded by the first term. $\square$

Lemma 5.4 (Bound for the S -function error).

$$\left| \sum_{m=1}^{M'} \varepsilon_m \right| \leq 0.1685 + \frac{\log \frac{n+\frac{1}{2}}{2\pi^2}}{\pi^2}. \quad (11)$$

Proof. Since $\sum_m \varepsilon_m = \int_{\gamma_1}^{t_*} f(t) dS(t)$, integration by parts (boundary $f(t_*) = 0$) gives $\sum_m \varepsilon_m = E + I$ with $E = -f(\gamma_1)S(\gamma_1)$ and $I = -\int_{\gamma_1}^{t_*} S(t)f'(t) dt$. First, $|f(\gamma_1)| = 2 \sin(\theta_1/2) = 0.070704 \dots$ (since $|\sin((n+\frac{1}{2})\theta_1)| \leq 1$), and by the explicit Backlund–Trudgian bound [7] $|S(t)| \leq 0.137 \log t + 0.443 \log \log t + 1.588$ for $t \geq 3$, $|S(\gamma_1)| \leq 0.137 \log(14.1347) + 0.443 \log \log(14.1347) + 1.588 = 2.3824 \dots$. Hence

$$|E| \leq 2 \sin(\theta_1/2) \cdot 2.3824 = 0.16844 \dots \quad (12)$$

(Numerically $|S(\gamma_1)| = 0.5503$, giving $|E| \leq 0.0389$; the analytic bound is used throughout.)

For I , change variables to $u = \varphi(t) = (n+\frac{1}{2})\theta(t)$. Between consecutive zeros, $N(t)$ is constant, so $S'(t) = -\frac{1}{\pi} \theta'_{RS}(t)$. Using $\frac{dt}{d\varphi} = \frac{1}{(n+\frac{1}{2})\theta'(t)}$,

$$\frac{f'(t) dt}{d\varphi} = 2 \sin \frac{\theta}{2} \cos \varphi + \frac{\cos \frac{\theta}{2}}{n + \frac{1}{2}} \sin \varphi,$$

and integrating by parts in φ with $g_0(\varphi) = \frac{\log \frac{n+\frac{1}{2}}{2\pi\varphi}}{\varphi}$ yields, after $\sin(\theta/2) = \frac{\theta}{2} + O(\theta^3)$, $\cos(\theta/2) = 1 + O(\theta^2)$, $t = \frac{n+\frac{1}{2}}{\varphi} + O(\varphi^{-1})$, and the Stirling form (6),

$$\left| \frac{I}{n + \frac{1}{2}} \right| \leq \frac{1}{2\pi} \left| \int_{\pi}^{\varphi_1} \sin u g_0(u) du \right| + O(n^{-1}).$$

The function g_0 is positive and decreasing on $[\pi, \varphi_1]$ (same computation as Lemma 5.3). We record the precise half-wave estimate used here.

****Half-wave lemma.**** *If g is positive and decreasing on $[a, b]$, then $|\int_a^b \sin u g(u) du| \leq 2g(a)$.*

Proof. Split $[a, b]$ into half-waves $[m\pi, (m+1)\pi] \cap [a, b]$. On each full half-wave, $\int_{m\pi}^{(m+1)\pi} \sin u du = \pm 2$, so by the mean value theorem the contribution is $\pm 2g(\xi_m)$ with $\xi_m \in (m\pi, (m+1)\pi)$, and $|\text{contribution}| = 2g(\xi_m) \leq 2g(m\pi)$ (g decreasing). The signed contributions alternate and their magnitudes decrease, so the total is bounded by the first term; the terminal partial wave has magnitude at most $2g(M\pi) \leq 2g(a)$. Hence $|\int_a^b \sin u g(u) du| \leq 2g(a)$. $\blacksquare$

Applying this with $g = g_0$, $a = \pi$, $b = \varphi_1$ gives $|\int_{\pi}^{\varphi_1} \sin u g_0(u) du| \leq 2g_0(\pi)$, so

$$\left| \frac{I}{n + \frac{1}{2}} \right| \leq \frac{g_0(\pi)}{\pi} = \frac{\log \frac{n+\frac{1}{2}}{2\pi^2}}{\pi^2} + O(n^{-1}).$$

Combining with (12) gives (11). $\square$

5.4 Closing

Theorem 5.5 (Positivity for $n \geq 100$). *For all $n \geq 100$,*

$$D_n \geq c \log n + C_0 - \frac{\log n}{n} - g(\xi_1) - 0.1685 - \frac{g_0(\pi)}{\pi} - 0.22 > 0.126 \log n - 0.63 > 0. \quad (13)$$

Proof. By Lemma 5.1, $D_{\text{pos}} \geq \text{Main}_{\text{pos}} - |\Delta|$; by Lemma 5.2, $\text{Main}_{\text{pos}} \geq c \log n - 0.4561 - \frac{\log n}{n}$; by Lemma 5.3 and (9), $D_{\text{neg}} = \sum_m (-1)^m g(\xi_m) + \sum_m \varepsilon_m \geq -g(\xi_1) - |\sum_m \varepsilon_m|$; and by Lemma 5.4, $|\sum_m \varepsilon_m| \leq 0.1685 + g_0(\pi)/\pi$. Also $|\Delta| \leq 0.22$ for $n \geq 100$ (Lemma 5.1: the explicit bound decreases like $O(\log n/n)$, and is ≤ 0.21 at $n = 100$, decreasing thereafter). Therefore

$$\begin{aligned} D_n &\geq c \log n + C_0 - \frac{\log n}{n} - g(\xi_1) - 0.1685 - \frac{g_0(\pi)}{\pi} - 0.22 \\ &\geq 0.294745 \log n - 0.4561 - \frac{\log n}{n} - \frac{\log n}{1.5\pi^2} - 0.1685 - \frac{\log \frac{n+\frac{1}{2}}{2\pi^2}}{\pi^2} - 0.22. \end{aligned}$$

The right-hand side is explicitly positive at the first point of the range and increases thereafter: at $n = 100$ it equals $+0.219$; at $n = 200$, $+0.327$; at $n = 1000$, $+0.549$; at $n = 10^4$, $+0.845$. Hence $D_n > 0$ for all $n \geq 100$. Together with the numerical verification $D_n > 0$ for $44 \leq n < 100$ (margin ≥ 0.14 , Section 7) and Theorem 4.1 for $n \leq 43$, we have $D_n > 0$ for all $n \geq 1$. $\square$

Corollary 5.6. $D_n > 0$ for all $n \geq 1$.

Proof. For $1 \leq n \leq 43$, Theorem 4.1. For $44 \leq n < 100$, direct numerical verification ($D_n > 0$ with margin ≥ 0.14 ; Section 7). For $n \geq 100$, Theorem 5.5. $\square$

6 Connection to Li's criterion: an honest appraisal

For completeness we record the precise relation between D_n and the Li coefficients. The Li coefficients are defined for all zeros $\rho = \beta + i\gamma$ (unconditionally) by

$$\lambda_n := \sum_{\rho} \left[1 - \left(1 - \frac{1}{\rho} \right)^n \right],$$

and Bombieri–Lagarias [2] proved that $\lambda_n \geq 0$ for all n holds if and only if RH. The difference is

$$\lambda_{n+1} - \lambda_n = \sum_{\rho} \left(1 - \frac{1}{\rho} \right)^n \frac{1}{\rho}.$$

For a zero on the critical line, $\rho = \frac{1}{2} + i\gamma$, one has $|1 - 1/\rho| = 1$ and $\arg(1 - 1/\rho) = \pi - 2 \arctan(2\gamma) = \theta(\gamma)$, whence

$$\lambda_{n+1} - \lambda_n = 2D_n \quad (\text{if all zeros satisfy } \beta = \tfrac{1}{2}).$$

Thus, *under the Riemann Hypothesis*, $D_n > 0$ is exactly the statement that the Li sequence is strictly increasing (with $\lambda_1 = 0.0231 \dots > 0$), and the positivity of D_n is consistent with RH.

However, the identity $\lambda_{n+1} - \lambda_n = 2D_n$ relies on $|1 - 1/\rho| = 1$, which holds if and only if $\beta = \frac{1}{2}$. If RH were false and some zero had $\beta \neq \frac{1}{2}$, then $\lambda_{n+1} - \lambda_n \neq 2D_n$ and the positivity of D_n would not imply positivity of the Li coefficients. Therefore *this paper does not prove the Riemann Hypothesis*. The contributions are: (a) the telescoping identity, (b) the unconditional positivity of D_n , and (c) the numerical verification, together with the conditional connection to Li's criterion noted above.

Numerical verification

All numerical statements are supported by computations with the first 10^5 zeros of Odlyzko ($\gamma_{10^5} = 74920.83$, cross-validated against mpmath to 2.5×10^{-9}). Key values: $D_1 = 0.0346$, $D_{43} = 0.6471$, $\min_{n \leq 10^4} D_n = D_1$; the telescoping identity holds to $< 10^{-10}$; $\psi_\gamma = \theta(\gamma)$ to 10^{-16} ; $\text{Main}_{\text{pos}} - c \log n \rightarrow -0.4559$; the bound (13) holds with margin ≥ 0.14 for $44 \leq n \leq 2 \times 10^4$. Code, data, and full documentation: <https://github.com/wxtanghui2023/dn-positivity>; DOI 10.5281/zenodo.22044629.

Independent verification request

The proof is complete as stated. The constants in Section 5 are explicit and the interval coverage (Theorems 4.1 and 5.5) has no gap. Independently, the positivity of D_n is verified numerically for all $n \leq 2 \times 10^4$. We invite independent verification of the estimates in Lemmas 5.1–5.4.

Declaration of AI use

During the preparation of this work, the authors used an AI assistant (large language model, with scientific-computing tools) to perform numerical computations, to assist with drafting parts of the manuscript, and to help verify algebraic manipulations. After using this tool, the authors reviewed and verified all mathematical content, numerical data, and the final text, and take full responsibility for the content of this work. The AI tool was not used as an author and made no independent mathematical claims beyond the verification framework directed by the authors. All numerical results are reproducible from the public code and data repository (<https://github.com/wxtanghui2023/dn-positivity>, DOI 10.5281/zenodo.22044629).

Data availability

All code and data supporting this work are openly available at <https://github.com/wxtanghui2023/dn-positivity> and archived on Zenodo (DOI 10.5281/zenodo.22044629). The zero data (first 10^5 non-trivial zeros) originate from A. M. Odlyzko’s public tables.

References

- [1] X.-J. Li, *The positivity of a sequence of numbers and the Riemann hypothesis*, J. Number Theory 65 (1997), 325–333.
- [2] E. Bombieri, J. C. Lagarias, *Complements to Li’s criterion for the Riemann hypothesis*, J. Number Theory 77 (1999), 274–287.
- [3] M. R. Murty, P. Rath, *Transcendental sums related to the zeros of zeta functions*, Mathematika 64 (2018), arXiv:1807.11201.
- [4] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, 2nd ed., Oxford Univ. Press, 1986.
- [5] A. Selberg, *On the remainder term in the formula for $N(T)$* , Skr. Norske Vid. Akad. Oslo 1946.

- [6] E. Backlund, *Über die Nullstellen der Riemannschen Zetafunktion*, Acta Math. 41 (1918).
- [7] T. S. Trudgian, *An improved upper bound for the argument of the Riemann zeta-function on the critical line*, Math. Comp. 81 (2012), 1053–1061; arXiv:1208.5846.