

vv214: Eigenvalue problems. Diagonalization.

Dr.Olga Danilkina

UM-SJTU Joint Institute



July 18, 2019

This week

1. Discrete dynamical systems
2. Eigenvectors/eigenfunctions and eigenvalues
3. Algebraic and geometric multiplicities
4. Eigenbasis
5. Diagonalization

Discrete Dynamical Systems

Consider a mathematical model

$$\begin{cases} c(t+1) = a_1 c(t) + b_1 r(t), \\ r(t+1) = a_2 c(t) + b_2 r(t) \end{cases}$$

that describes the dynamic of two populations as time changes (say, a **predator-prey model**).

Discrete Dynamical Systems

Consider a mathematical model

$$\begin{cases} c(t+1) = a_1 c(t) + b_1 r(t), \\ r(t+1) = a_2 c(t) + b_2 r(t) \end{cases}$$

that describes the dynamic of two populations as time changes (say, a **predator-prey model**).

$$\Rightarrow 0 < a_1, b_1 < 1, b_2 > 1, a_2 < 0$$

Discrete Dynamical Systems

Consider a mathematical model

$$\begin{cases} c(t+1) = a_1 c(t) + b_1 r(t), \\ r(t+1) = a_2 c(t) + b_2 r(t) \end{cases}$$

that describes the dynamic of two populations as time changes (say, a **predator-prey model**).

$$\Rightarrow 0 < a_1, b_1 < 1, b_2 > 1, a_2 < 0$$

$$a_1 = 0.86, b_1 = 0.08, a_2 = -0.12, b_2 = 1.14$$

Discrete Dynamical Systems

Consider a mathematical model

$$\begin{cases} c(t+1) = a_1 c(t) + b_1 r(t), \\ r(t+1) = a_2 c(t) + b_2 r(t) \end{cases}$$

that describes the dynamic of two populations as time changes (say, a **predator-prey model**).

$$\Rightarrow 0 < a_1, b_1 < 1, b_2 > 1, a_2 < 0$$

$$a_1 = 0.86, b_1 = 0.08, a_2 = -0.12, b_2 = 1.14$$

$$\bar{x}(t) = \begin{pmatrix} c(t) \\ r(t) \end{pmatrix} \Rightarrow$$

Discrete Dynamical Systems

Consider a mathematical model

$$\begin{cases} c(t+1) = a_1 c(t) + b_1 r(t), \\ r(t+1) = a_2 c(t) + b_2 r(t) \end{cases}$$

that describes the dynamic of two populations as time changes (say, a **predator-prey model**).

$$\Rightarrow 0 < a_1, b_1 < 1, b_2 > 1, a_2 < 0$$

$$a_1 = 0.86, b_1 = 0.08, a_2 = -0.12, b_2 = 1.14$$

$$\bar{x}(t) = \begin{pmatrix} c(t) \\ r(t) \end{pmatrix} \Rightarrow \bar{x}(t+1) = A\bar{x}(t), A = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix}$$

Discrete Dynamical Systems

Consider a mathematical model

$$\begin{cases} c(t+1) = a_1 c(t) + b_1 r(t), \\ r(t+1) = a_2 c(t) + b_2 r(t) \end{cases}$$

that describes the dynamic of two populations as time changes (say, a **predator-prey model**).

$$\Rightarrow 0 < a_1, b_1 < 1, b_2 > 1, a_2 < 0$$

$$a_1 = 0.86, b_1 = 0.08, a_2 = -0.12, b_2 = 1.14$$

$$\bar{x}(t) = \begin{pmatrix} c(t) \\ r(t) \end{pmatrix} \Rightarrow \bar{x}(t+1) = A\bar{x}(t), A = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix}$$

$$\bar{x}_0(t) = \begin{pmatrix} c_0 \\ r_0 \end{pmatrix}$$

Discrete Dynamical Systems

Consider a mathematical model

$$\begin{cases} c(t+1) = a_1 c(t) + b_1 r(t), \\ r(t+1) = a_2 c(t) + b_2 r(t) \end{cases}$$

that describes the dynamic of two populations as time changes (say, a **predator-prey model**).

$$\Rightarrow 0 < a_1, b_1 < 1, b_2 > 1, a_2 < 0$$

$$a_1 = 0.86, b_1 = 0.08, a_2 = -0.12, b_2 = 1.14$$

$$\bar{x}(t) = \begin{pmatrix} c(t) \\ r(t) \end{pmatrix} \Rightarrow \bar{x}(t+1) = A\bar{x}(t), A = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix}$$

$$\bar{x}_0(t) = \begin{pmatrix} c_0 \\ r_0 \end{pmatrix} \xrightarrow{A} \bar{x}_1 = \bar{x}(1) = \begin{pmatrix} c(1) \\ r(1) \end{pmatrix}$$

Discrete Dynamical Systems

Consider a mathematical model

$$\begin{cases} c(t+1) = a_1 c(t) + b_1 r(t), \\ r(t+1) = a_2 c(t) + b_2 r(t) \end{cases}$$

that describes the dynamic of two populations as time changes (say, a **predator-prey model**).

$$\Rightarrow 0 < a_1, b_1 < 1, b_2 > 1, a_2 < 0$$

$$a_1 = 0.86, b_1 = 0.08, a_2 = -0.12, b_2 = 1.14$$

$$\bar{x}(t) = \begin{pmatrix} c(t) \\ r(t) \end{pmatrix} \Rightarrow \bar{x}(t+1) = A\bar{x}(t), A = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix}$$

$$\bar{x}_0(t) = \begin{pmatrix} c_0 \\ r_0 \end{pmatrix} \xrightarrow{A} \bar{x}_1 = \bar{x}(1) = \begin{pmatrix} c(1) \\ r(1) \end{pmatrix} \xrightarrow{A} \dots \xrightarrow{A} \bar{x}(t) \xrightarrow{A} \dots$$

Discrete Dynamical Systems

Consider a mathematical model

$$\begin{cases} c(t+1) = a_1 c(t) + b_1 r(t), \\ r(t+1) = a_2 c(t) + b_2 r(t) \end{cases}$$

that describes the dynamic of two populations as time changes (say, a **predator-prey model**).

$$\Rightarrow 0 < a_1, b_1 < 1, b_2 > 1, a_2 < 0$$

$$a_1 = 0.86, b_1 = 0.08, a_2 = -0.12, b_2 = 1.14$$

$$\bar{x}(t) = \begin{pmatrix} c(t) \\ r(t) \end{pmatrix} \Rightarrow \bar{x}(t+1) = A\bar{x}(t), A = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix}$$

$$\bar{x}_0(t) = \begin{pmatrix} c_0 \\ r_0 \end{pmatrix} \xrightarrow{A} \bar{x}_1 = \bar{x}(1) = \begin{pmatrix} c(1) \\ r(1) \end{pmatrix} \xrightarrow{A} \dots \xrightarrow{A} \bar{x}(t) \xrightarrow{A} \dots$$

$$\bar{x}_1 = A\bar{x}_0,$$

Discrete Dynamical Systems

Consider a mathematical model

$$\begin{cases} c(t+1) = a_1 c(t) + b_1 r(t), \\ r(t+1) = a_2 c(t) + b_2 r(t) \end{cases}$$

that describes the dynamic of two populations as time changes (say, a **predator-prey model**).

$$\Rightarrow 0 < a_1, b_1 < 1, b_2 > 1, a_2 < 0$$

$$a_1 = 0.86, b_1 = 0.08, a_2 = -0.12, b_2 = 1.14$$

$$\bar{x}(t) = \begin{pmatrix} c(t) \\ r(t) \end{pmatrix} \Rightarrow \bar{x}(t+1) = A\bar{x}(t), A = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix}$$

$$\bar{x}_0(t) = \begin{pmatrix} c_0 \\ r_0 \end{pmatrix} \xrightarrow{A} \bar{x}_1 = \bar{x}(1) = \begin{pmatrix} c(1) \\ r(1) \end{pmatrix} \xrightarrow{A} \dots \xrightarrow{A} \bar{x}(t) \xrightarrow{A} \dots$$

$$\bar{x}_1 = A\bar{x}_0, \bar{x}_2 = A\bar{x}_1 =$$

Discrete Dynamical Systems

Consider a mathematical model

$$\begin{cases} c(t+1) = a_1 c(t) + b_1 r(t), \\ r(t+1) = a_2 c(t) + b_2 r(t) \end{cases}$$

that describes the dynamic of two populations as time changes (say, a **predator-prey model**).

$$\Rightarrow 0 < a_1, b_1 < 1, b_2 > 1, a_2 < 0$$

$$a_1 = 0.86, b_1 = 0.08, a_2 = -0.12, b_2 = 1.14$$

$$\bar{x}(t) = \begin{pmatrix} c(t) \\ r(t) \end{pmatrix} \Rightarrow \bar{x}(t+1) = A\bar{x}(t), A = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix}$$

$$\bar{x}_0(t) = \begin{pmatrix} c_0 \\ r_0 \end{pmatrix} \xrightarrow{A} \bar{x}_1 = \bar{x}(1) = \begin{pmatrix} c(1) \\ r(1) \end{pmatrix} \xrightarrow{A} \dots \xrightarrow{A} \bar{x}(t) \xrightarrow{A} \dots$$

$$\bar{x}_1 = A\bar{x}_0, \bar{x}_2 = A\bar{x}_1 = A^2\bar{x}_0,$$

Discrete Dynamical Systems

Consider a mathematical model

$$\begin{cases} c(t+1) = a_1 c(t) + b_1 r(t), \\ r(t+1) = a_2 c(t) + b_2 r(t) \end{cases}$$

that describes the dynamic of two populations as time changes (say, a **predator-prey model**).

$$\Rightarrow 0 < a_1, b_1 < 1, b_2 > 1, a_2 < 0$$

$$a_1 = 0.86, b_1 = 0.08, a_2 = -0.12, b_2 = 1.14$$

$$\bar{x}(t) = \begin{pmatrix} c(t) \\ r(t) \end{pmatrix} \Rightarrow \bar{x}(t+1) = A\bar{x}(t), A = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix}$$

$$\bar{x}_0(t) = \begin{pmatrix} c_0 \\ r_0 \end{pmatrix} \xrightarrow{A} \bar{x}_1 = \bar{x}(1) = \begin{pmatrix} c(1) \\ r(1) \end{pmatrix} \xrightarrow{A} \dots \xrightarrow{A} \bar{x}(t) \xrightarrow{A} \dots$$

$$\bar{x}_1 = A\bar{x}_0, \bar{x}_2 = A\bar{x}_1 = A^2\bar{x}_0, \dots, \boxed{\bar{x}(t) = A^t\bar{x}_0}$$

Discrete Dynamical Systems

Case 1: Let $c_0 = 100$, $r_0 = 300$.

Discrete Dynamical Systems

Case 1: Let $c_0 = 100$, $r_0 = 300$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 100 \\ 100 \end{pmatrix} = \begin{pmatrix} 110 \\ 330 \end{pmatrix} =$$

Discrete Dynamical Systems

Case 1: Let $c_0 = 100$, $r_0 = 300$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 100 \\ 100 \end{pmatrix} = \begin{pmatrix} 110 \\ 330 \end{pmatrix} = 1.1\bar{x}_0 \Rightarrow$$

Discrete Dynamical Systems

Case 1: Let $c_0 = 100$, $r_0 = 300$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 100 \\ 100 \end{pmatrix} = \begin{pmatrix} 110 \\ 330 \end{pmatrix} = 1.1\bar{x}_0 \Rightarrow$$

$$\bar{x}(t) = (1.1)^t \bar{x}_0 \Rightarrow$$

Discrete Dynamical Systems

Case 1: Let $c_0 = 100$, $r_0 = 300$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 100 \\ 100 \end{pmatrix} = \begin{pmatrix} 110 \\ 330 \end{pmatrix} = 1.1\bar{x}_0 \Rightarrow$$

$$\bar{x}(t) = (1.1)^t \bar{x}_0 \Rightarrow \begin{matrix} c(t) = (1.1)^t 100 \\ r(t) = (1.1)^t 300 \end{matrix} \Rightarrow 10\% \text{ growth}$$

Discrete Dynamical Systems

Case 1: Let $c_0 = 100$, $r_0 = 300$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 100 \\ 100 \end{pmatrix} = \begin{pmatrix} 110 \\ 330 \end{pmatrix} = 1.1\bar{x}_0 \Rightarrow$$

$$\bar{x}(t) = (1.1)^t \bar{x}_0 \Rightarrow \begin{matrix} c(t) = (1.1)^t 100 \\ r(t) = (1.1)^t 300 \end{matrix} \Rightarrow 10\% \text{ growth}$$

Case 2: Let $c_0 = 200$, $r_0 = 100$.

Discrete Dynamical Systems

Case 1: Let $c_0 = 100$, $r_0 = 300$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 100 \\ 100 \end{pmatrix} = \begin{pmatrix} 110 \\ 330 \end{pmatrix} = 1.1\bar{x}_0 \Rightarrow$$

$$\bar{x}(t) = (1.1)^t \bar{x}_0 \Rightarrow \begin{matrix} c(t) = (1.1)^t 100 \\ r(t) = (1.1)^t 300 \end{matrix} \Rightarrow 10\% \text{ growth}$$

Case 2: Let $c_0 = 200$, $r_0 = 100$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 200 \\ 100 \end{pmatrix} = \begin{pmatrix} 180 \\ 90 \end{pmatrix} =$$

Discrete Dynamical Systems

Case 1: Let $c_0 = 100$, $r_0 = 300$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 100 \\ 100 \end{pmatrix} = \begin{pmatrix} 110 \\ 330 \end{pmatrix} = 1.1\bar{x}_0 \Rightarrow$$

$$\bar{x}(t) = (1.1)^t \bar{x}_0 \Rightarrow \begin{matrix} c(t) = (1.1)^t 100 \\ r(t) = (1.1)^t 300 \end{matrix} \Rightarrow 10\% \text{ growth}$$

Case 2: Let $c_0 = 200$, $r_0 = 100$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 200 \\ 100 \end{pmatrix} = \begin{pmatrix} 180 \\ 90 \end{pmatrix} = 0.9\bar{x}_0 \Rightarrow$$

Discrete Dynamical Systems

Case 1: Let $c_0 = 100$, $r_0 = 300$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 100 \\ 100 \end{pmatrix} = \begin{pmatrix} 110 \\ 330 \end{pmatrix} = 1.1\bar{x}_0 \Rightarrow$$

$$\bar{x}(t) = (1.1)^t \bar{x}_0 \Rightarrow \begin{matrix} c(t) = (1.1)^t 100 \\ r(t) = (1.1)^t 300 \end{matrix} \Rightarrow 10\% \text{ growth}$$

Case 2: Let $c_0 = 200$, $r_0 = 100$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 200 \\ 100 \end{pmatrix} = \begin{pmatrix} 180 \\ 90 \end{pmatrix} = 0.9\bar{x}_0 \Rightarrow$$

$$\bar{x}(t) = (0.9)^t \bar{x}_0 \Rightarrow$$

Discrete Dynamical Systems

Case 1: Let $c_0 = 100$, $r_0 = 300$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 100 \\ 100 \end{pmatrix} = \begin{pmatrix} 110 \\ 330 \end{pmatrix} = 1.1\bar{x}_0 \Rightarrow$$

$$\bar{x}(t) = (1.1)^t \bar{x}_0 \Rightarrow \begin{matrix} c(t) = (1.1)^t 100 \\ r(t) = (1.1)^t 300 \end{matrix} \Rightarrow 10\% \text{ growth}$$

Case 2: Let $c_0 = 200$, $r_0 = 100$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 200 \\ 100 \end{pmatrix} = \begin{pmatrix} 180 \\ 90 \end{pmatrix} = 0.9\bar{x}_0 \Rightarrow$$

$$\bar{x}(t) = (0.9)^t \bar{x}_0 \Rightarrow \begin{matrix} c(t) = (0.9)^t 200 \\ r(t) = (0.9)^t 100 \end{matrix} \Rightarrow 10\% \text{ declination}$$

Discrete Dynamical Systems

Case 3: Let $c_0 = 1000$, $r_0 = 1000$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = \begin{pmatrix} 940 \\ 1020 \end{pmatrix} = ? C \bar{x}_0$$

Discrete Dynamical Systems

Case 3: Let $c_0 = 1000$, $r_0 = 1000$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = \begin{pmatrix} 940 \\ 1020 \end{pmatrix} = ? C \bar{x}_0$$

Since it worked in Cases 1,2, so let

$$\bar{x}_0 = \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = C_1 \underbrace{\begin{pmatrix} 100 \\ 300 \end{pmatrix}}_{\tilde{v}_1} + C_2 \underbrace{\begin{pmatrix} 200 \\ 100 \end{pmatrix}}_{\tilde{v}_2} =$$

Discrete Dynamical Systems

Case 3: Let $c_0 = 1000$, $r_0 = 1000$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = \begin{pmatrix} 940 \\ 1020 \end{pmatrix} = ? C \bar{x}_0$$

Since it worked in Cases 1,2, so let

$$\bar{x}_0 = \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = C_1 \underbrace{\begin{pmatrix} 100 \\ 300 \end{pmatrix}}_{\bar{v}_1} + C_2 \underbrace{\begin{pmatrix} 200 \\ 100 \end{pmatrix}}_{\bar{v}_2} = 2\bar{v}_1 + 4\bar{v}_2$$

Discrete Dynamical Systems

Case 3: Let $c_0 = 1000$, $r_0 = 1000$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = \begin{pmatrix} 940 \\ 1020 \end{pmatrix} = ? C \bar{x}_0$$

Since it worked in Cases 1,2, so let

$$\bar{x}_0 = \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = C_1 \underbrace{\begin{pmatrix} 100 \\ 300 \end{pmatrix}}_{\tilde{v}_1} + C_2 \underbrace{\begin{pmatrix} 200 \\ 100 \end{pmatrix}}_{\tilde{v}_2} = 2\tilde{v}_1 + 4\tilde{v}_2$$

$$\bar{x}(1) = A\bar{x}_0 =$$

Discrete Dynamical Systems

Case 3: Let $c_0 = 1000$, $r_0 = 1000$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = \begin{pmatrix} 940 \\ 1020 \end{pmatrix} = ? C \bar{x}_0$$

Since it worked in Cases 1,2, so let

$$\bar{x}_0 = \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = C_1 \underbrace{\begin{pmatrix} 100 \\ 300 \end{pmatrix}}_{\bar{v}_1} + C_2 \underbrace{\begin{pmatrix} 200 \\ 100 \end{pmatrix}}_{\bar{v}_2} = 2\bar{v}_1 + 4\bar{v}_2$$

$$\bar{x}(1) = A\bar{x}_0 = A(2\bar{v}_1 + 4\bar{v}_2) =$$

Discrete Dynamical Systems

Case 3: Let $c_0 = 1000$, $r_0 = 1000$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = \begin{pmatrix} 940 \\ 1020 \end{pmatrix} = ? C \bar{x}_0$$

Since it worked in Cases 1,2, so let

$$\bar{x}_0 = \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = C_1 \underbrace{\begin{pmatrix} 100 \\ 300 \end{pmatrix}}_{\bar{v}_1} + C_2 \underbrace{\begin{pmatrix} 200 \\ 100 \end{pmatrix}}_{\bar{v}_2} = 2\bar{v}_1 + 4\bar{v}_2$$

$$\bar{x}(1) = A\bar{x}_0 = A(2\bar{v}_1 + 4\bar{v}_2) = 2(1.1)\bar{v}_1 + 4(0.9)\bar{v}_2 \Rightarrow$$

Discrete Dynamical Systems

Case 3: Let $c_0 = 1000$, $r_0 = 1000$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = \begin{pmatrix} 940 \\ 1020 \end{pmatrix} = ? C \bar{x}_0$$

Since it worked in Cases 1,2, so let

$$\bar{x}_0 = \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = C_1 \underbrace{\begin{pmatrix} 100 \\ 300 \end{pmatrix}}_{\bar{v}_1} + C_2 \underbrace{\begin{pmatrix} 200 \\ 100 \end{pmatrix}}_{\bar{v}_2} = 2\bar{v}_1 + 4\bar{v}_2$$

$$\bar{x}(1) = A\bar{x}_0 = A(2\bar{v}_1 + 4\bar{v}_2) = 2(1.1)\bar{v}_1 + 4(0.9)\bar{v}_2 \Rightarrow$$

$$\bar{x}(t) = 2(1.1)^t \bar{v}_1 + 4(0.9)^t \bar{v}_2$$

Discrete Dynamical Systems

Case 3: Let $c_0 = 1000$, $r_0 = 1000$.

$$\bar{x}(1) = \begin{pmatrix} 0.86 & 0.08 \\ -0.12 & 1.14 \end{pmatrix} \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = \begin{pmatrix} 940 \\ 1020 \end{pmatrix} = ? C \bar{x}_0$$

Since it worked in Cases 1,2, so let

$$\bar{x}_0 = \begin{pmatrix} 1000 \\ 1000 \end{pmatrix} = C_1 \underbrace{\begin{pmatrix} 100 \\ 300 \end{pmatrix}}_{\bar{v}_1} + C_2 \underbrace{\begin{pmatrix} 200 \\ 100 \end{pmatrix}}_{\bar{v}_2} = 2\bar{v}_1 + 4\bar{v}_2$$

$$\bar{x}(1) = A\bar{x}_0 = A(2\bar{v}_1 + 4\bar{v}_2) = 2(1.1)\bar{v}_1 + 4(0.9)\bar{v}_2 \Rightarrow$$

$$\bar{x}(t) = 2(1.1)^t \bar{v}_1 + 4(0.9)^t \bar{v}_2$$

$$c(t) = 200(1.1)^t + 800(0.9)^t$$

$$r(t) = 600(1.1)^t + 400(0.9)^t$$

Eigenvectors and Eigenvalues

Definition: A non-zero vector $\vec{v} \in \mathbb{R}^n$ is called an **eigenvector** of the matrix $A_{n \times n}$ if

$$\exists \lambda \in \mathbb{K}: A\vec{v} = \lambda\vec{v}$$

Eigenvectors and Eigenvalues

Definition: A non-zero vector $\vec{v} \in \mathbb{R}^n$ is called an **eigenvector** of the matrix $A_{n \times n}$ if

$$\exists \lambda \in \mathbb{K}: A\vec{v} = \lambda\vec{v}$$

The scalar λ is called the **eigenvalue associated with the eigenvector** \vec{v} .

Eigenvectors and Eigenvalues

Definition: A non-zero vector $\vec{v} \in \mathbb{R}^n$ is called an **eigenvector** of the matrix $A_{n \times n}$ if

$$\exists \lambda \in \mathbb{K}: A\vec{v} = \lambda\vec{v}$$

The scalar λ is called the **eigenvalue associated with the eigenvector** \vec{v} .

Remarks:

1. $A\vec{v} \parallel \vec{v} \Rightarrow \vec{v}$ is the eigenvector.

Eigenvectors and Eigenvalues

Definition: A non-zero vector $\vec{v} \in \mathbb{R}^n$ is called an **eigenvector** of the matrix $A_{n \times n}$ if

$$\exists \lambda \in \mathbb{K}: A\vec{v} = \lambda\vec{v}$$

The scalar λ is called the **eigenvalue associated with the eigenvector** \vec{v} .

Remarks:

1. $A\vec{v} \parallel \vec{v} \Rightarrow \vec{v}$ is the eigenvector.
2. \vec{v} is the eigenvector of $A \Rightarrow \vec{v}$ is the eigenvector of A^2, A^3, \dots

Eigenvectors and Eigenvalues

Definition: A non-zero vector $\vec{v} \in \mathbb{R}^n$ is called an **eigenvector** of the matrix $A_{n \times n}$ if

$$\exists \lambda \in \mathbb{K}: A\vec{v} = \lambda\vec{v}$$

The scalar λ is called the **eigenvalue associated with the eigenvector** \vec{v} .

Remarks:

1. $A\vec{v} \parallel \vec{v} \Rightarrow \vec{v}$ is the eigenvector.
2. \vec{v} is the eigenvector of $A \Rightarrow \vec{v}$ is the eigenvector of A^2, A^3, \dots
3. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the orthogonal projection onto a line L

Eigenvectors and Eigenvalues

Definition: A non-zero vector $\vec{v} \in \mathbb{R}^n$ is called an **eigenvector** of the matrix $A_{n \times n}$ if

$$\exists \lambda \in \mathbb{K}: A\vec{v} = \lambda\vec{v}$$

The scalar λ is called the **eigenvalue associated with the eigenvector** \vec{v} .

Remarks:

1. $A\vec{v} \parallel \vec{v} \Rightarrow \vec{v}$ is the eigenvector.
2. \vec{v} is the eigenvector of $A \Rightarrow \vec{v}$ is the eigenvector of A^2, A^3, \dots
3. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the orthogonal projection onto a line L

$$\Rightarrow \vec{w} \perp L, \vec{v} \in L \quad \text{are eigenvectors } (\lambda = 0, \lambda = 1)$$

Eigenvectors and Eigenvalues

Definition: A non-zero vector $\vec{v} \in \mathbb{R}^n$ is called an **eigenvector** of the matrix $A_{n \times n}$ if

$$\exists \lambda \in \mathbb{K}: A\vec{v} = \lambda\vec{v}$$

The scalar λ is called the **eigenvalue associated with the eigenvector** \vec{v} .

Remarks:

1. $A\vec{v} \parallel \vec{v} \Rightarrow \vec{v}$ is the eigenvector.
2. \vec{v} is the eigenvector of $A \Rightarrow \vec{v}$ is the eigenvector of A^2, A^3, \dots
3. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the orthogonal projection onto a line L

$$\Rightarrow \vec{w} \perp L, \vec{v} \in L \quad \text{are eigenvectors } (\lambda = 0, \lambda = 1)$$

4. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the counterclockwise rotation through $\frac{\pi}{2}$

Eigenvectors and Eigenvalues

Definition: A non-zero vector $\vec{v} \in \mathbb{R}^n$ is called an **eigenvector** of the matrix $A_{n \times n}$ if

$$\exists \lambda \in \mathbb{K}: A\vec{v} = \lambda\vec{v}$$

The scalar λ is called the **eigenvalue associated with the eigenvector** \vec{v} .

Remarks:

1. $A\vec{v} \parallel \vec{v} \Rightarrow \vec{v}$ is the eigenvector.
2. \vec{v} is the eigenvector of $A \Rightarrow \vec{v}$ is the eigenvector of A^2, A^3, \dots
3. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the orthogonal projection onto a line L

$$\Rightarrow \vec{w} \perp L, \vec{v} \in L \quad \text{are eigenvectors } (\lambda = 0, \lambda = 1)$$

4. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the counterclockwise rotation through $\frac{\pi}{2}$

\Rightarrow there are no real eigenvalues and eigenvectors

Eigenvectors and Eigenvalues

Definition: A non-zero vector $\vec{v} \in \mathbb{R}^n$ is called an **eigenvector** of the matrix $A_{n \times n}$ if

$$\exists \lambda \in \mathbb{K}: A\vec{v} = \lambda\vec{v}$$

The scalar λ is called the **eigenvalue associated with the eigenvector** \vec{v} .

Remarks:

1. $A\vec{v} \parallel \vec{v} \Rightarrow \vec{v}$ is the eigenvector.
2. \vec{v} is the eigenvector of $A \Rightarrow \vec{v}$ is the eigenvector of A^2, A^3, \dots
3. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the orthogonal projection onto a line L

$$\Rightarrow \vec{w} \perp L, \vec{v} \in L \quad \text{are eigenvectors } (\lambda = 0, \lambda = 1)$$

4. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the counterclockwise rotation through $\frac{\pi}{2}$
 \Rightarrow there are no real eigenvalues and eigenvectors

5. $A_{n \times n}$ is orthogonal \Rightarrow

Eigenvectors and Eigenvalues

Definition: A non-zero vector $\vec{v} \in \mathbb{R}^n$ is called an **eigenvector** of the matrix $A_{n \times n}$ if

$$\exists \lambda \in \mathbb{K}: A\vec{v} = \lambda\vec{v}$$

The scalar λ is called the **eigenvalue associated with the eigenvector** \vec{v} .

Remarks:

1. $A\vec{v} \parallel \vec{v} \Rightarrow \vec{v}$ is the eigenvector.
2. \vec{v} is the eigenvector of $A \Rightarrow \vec{v}$ is the eigenvector of A^2, A^3, \dots
3. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the orthogonal projection onto a line L

$$\Rightarrow \vec{w} \perp L, \vec{v} \in L \quad \text{are eigenvectors } (\lambda = 0, \lambda = 1)$$

4. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the counterclockwise rotation through $\frac{\pi}{2}$

\Rightarrow there are no real eigenvalues and eigenvectors

5. $A_{n \times n}$ is orthogonal \Rightarrow the eigenvalues are ± 1

Discrete Dynamical Systems

Definition: If a state $\bar{x}(t)$ of a physical system at any given time t is described by n values $x_1(t), \dots, x_n(t)$, then the *law of change* of the state of the system from time t to time $t + 1$ in the linear form

$$\bar{x}(t + 1) = A_{n \times n} \bar{x}(t)$$

is called a **discrete dynamical system**.

Discrete Dynamical Systems

Definition: If a state $\bar{x}(t)$ of a physical system at any given time t is described by n values $x_1(t), \dots, x_n(t)$, then the *law of change* of the state of the system from time t to time $t + 1$ in the linear form

$$\bar{x}(t + 1) = A_{n \times n} \bar{x}(t)$$

is called a **discrete dynamical system**.

Remarks:

1. $\bar{x}(t) = A^t \bar{x}_0$ if the initial state \bar{x}_0 is known.

Discrete Dynamical Systems

Definition: If a state $\bar{x}(t)$ of a physical system at any given time t is described by n values $x_1(t), \dots, x_n(t)$, then the *law of change* of the state of the system from time t to time $t + 1$ in the linear form

$$\bar{x}(t + 1) = A_{n \times n} \bar{x}(t)$$

is called a **discrete dynamical system**.

Remarks:

1. $\bar{x}(t) = A^t \bar{x}_0$ if the initial state \bar{x}_0 is known.
2. If a basis $\{\bar{v}_1, \dots, \bar{v}_n\} \subset \mathbb{R}^n$ consists of eigenvalues of A :

$$A\bar{v}_i = \lambda_i \bar{v}_i, \quad i = 1..n,$$

then

Discrete Dynamical Systems

Definition: If a state $\bar{x}(t)$ of a physical system at any given time t is described by n values $x_1(t), \dots, x_n(t)$, then the *law of change* of the state of the system from time t to time $t + 1$ in the linear form

$$\bar{x}(t + 1) = A_{n \times n} \bar{x}(t)$$

is called a **discrete dynamical system**.

Remarks:

1. $\bar{x}(t) = A^t \bar{x}_0$ if the initial state \bar{x}_0 is known.
2. If a basis $\{\bar{v}_1, \dots, \bar{v}_n\} \subset \mathbb{R}^n$ consists of eigenvalues of A :

$$A\bar{v}_i = \lambda_i \bar{v}_i, \quad i = 1..n,$$

then

$$\bar{x}(t) = \underbrace{(\bar{v}_1 \quad \bar{v}_2 \dots \bar{v}_n)}_S \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}^t S^{-1} \bar{x}_0$$

Lilac Bush Growth Model

Lilac Bush Growth Model

Year 1



Year 2



Year 3




Year 4



Year 5




Lilac Bush Growth Model



$n(t)$ is the number of new branches and
 $a(t)$ is the number of old branches


Lilac Bush Growth Model



$n(t)$ is the number of new branches and
 $a(t)$ is the number of old branches

$$a(t+1) = n(t) + a(t),$$


Lilac Bush Growth Model



$n(t)$ is the number of new branches and
 $a(t)$ is the number of old branches

$$a(t+1) = n(t) + a(t), \quad n(t+1) = 2a(t)$$

Lilac Bush Growth Model




$n(t)$ is the number of new branches and
 $a(t)$ is the number of old branches

$$a(t+1) = n(t) + a(t), \quad n(t+1) = 2a(t)$$

$$\begin{pmatrix} n(t+1) \\ a(t+1) \end{pmatrix} = \underbrace{\begin{pmatrix} 0 & 2 \\ 1 & 1 \end{pmatrix}}$$

Lilac Bush Growth Model



$n(t)$ is the number of new branches and
 $a(t)$ is the number of old branches

$$a(t+1) = n(t) + a(t), \quad n(t+1) = 2a(t)$$

$$\begin{pmatrix} n(t+1) \\ a(t+1) \end{pmatrix} = \underbrace{\begin{pmatrix} 0 & 2 \\ 1 & 1 \end{pmatrix}}_A \begin{pmatrix} n(t) \\ a(t) \end{pmatrix}$$

Lilac Bush Growth Model

$$A \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix},$$

Lilac Bush Growth Model

$$A \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad A \begin{pmatrix} 2 \\ -1 \end{pmatrix} = -1 \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

Lilac Bush Growth Model

$$A \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad A \begin{pmatrix} 2 \\ -1 \end{pmatrix} = -1 \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

Since $n(0) = 1$ and $a(0) = 0$, so

Lilac Bush Growth Model

$$A \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad A \begin{pmatrix} 2 \\ -1 \end{pmatrix} = -1 \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

Since $n(0) = 1$ and $a(0) = 0$, so

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} =$$

Lilac Bush Growth Model

$$A \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad A \begin{pmatrix} 2 \\ -1 \end{pmatrix} = -1 \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

Since $n(0) = 1$ and $a(0) = 0$, so

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} = c_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} + c_2 \begin{pmatrix} 2 \\ -1 \end{pmatrix} \Rightarrow$$

Lilac Bush Growth Model

$$A \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad A \begin{pmatrix} 2 \\ -1 \end{pmatrix} = -1 \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

Since $n(0) = 1$ and $a(0) = 0$, so

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} = c_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} + c_2 \begin{pmatrix} 2 \\ -1 \end{pmatrix} \Rightarrow c_1 = c_2 = \frac{1}{3}$$

Lilac Bush Growth Model

$$A \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad A \begin{pmatrix} 2 \\ -1 \end{pmatrix} = -1 \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

Since $n(0) = 1$ and $a(0) = 0$, so

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} = c_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} + c_2 \begin{pmatrix} 2 \\ -1 \end{pmatrix} \Rightarrow c_1 = c_2 = \frac{1}{3}$$

$$\begin{pmatrix} n(t+1) \\ a(t+1) \end{pmatrix} = \frac{2^t}{3} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \frac{(-1)^t}{3} \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

Invariant Subspaces

Definition: Let $A_{n \times n}$. A linear subspace $V \subset \mathbb{R}^n$ is **A-invariant** if

$$A\vec{v} \in V \quad \forall \vec{v} \in V$$

Invariant Subspaces

Definition: Let $A_{n \times n}$. A linear subspace $V \subset \mathbb{R}^n$ is **A-invariant** if

$$A\vec{v} \in V \quad \forall \vec{v} \in V$$

Let $\dim V = 1$.

Invariant Subspaces

Definition: Let $A_{n \times n}$. A linear subspace $V \subset \mathbb{R}^n$ is **A-invariant** if

$$A\vec{v} \in V \quad \forall \vec{v} \in V$$

Let $\dim V = 1$.

1. If $\vec{v} \in V$, $\vec{v} \neq 0$, then

Invariant Subspaces

Definition: Let $A_{n \times n}$. A linear subspace $V \subset \mathbb{R}^n$ is **A-invariant** if

$$A\vec{v} \in V \quad \forall \vec{v} \in V$$

Let $\dim V = 1$.

1. If $\vec{v} \in V$, $\vec{v} \neq 0$, then

$$A\vec{v} \in V \Rightarrow$$

Invariant Subspaces

Definition: Let $A_{n \times n}$. A linear subspace $V \subset \mathbb{R}^n$ is **A-invariant** if

$$A\bar{v} \in V \quad \forall v \in V$$

Let $\dim V = 1$.

1. If $\bar{v} \in V$, $\bar{v} \neq 0$, then

$$A\bar{v} \in V \Rightarrow A\bar{v} \in \text{span } \bar{v} \Rightarrow$$

Invariant Subspaces

Definition: Let $A_{n \times n}$. A linear subspace $V \subset \mathbb{R}^n$ is **A-invariant** if

$$A\bar{v} \in V \quad \forall v \in V$$

Let $\dim V = 1$.

1. If $\bar{v} \in V$, $\bar{v} \neq 0$, then

$$A\bar{v} \in V \Rightarrow A\bar{v} \in \text{span } \bar{v} \Rightarrow A\bar{v} = \lambda\bar{v}$$

Invariant Subspaces

Definition: Let $A_{n \times n}$. A linear subspace $V \subset \mathbb{R}^n$ is **A-invariant** if

$$A\bar{v} \in V \quad \forall v \in V$$

Let $\dim V = 1$.

1. If $\bar{v} \in V$, $\bar{v} \neq 0$, then

$$A\bar{v} \in V \Rightarrow A\bar{v} \in \text{span } \bar{v} \Rightarrow A\bar{v} = \lambda\bar{v}$$

$\Rightarrow \bar{v}$ is an eigenvector of A .

Invariant Subspaces

Definition: Let $A_{n \times n}$. A linear subspace $V \subset \mathbb{R}^n$ is **A-invariant** if

$$A\bar{v} \in V \quad \forall v \in V$$

Let $\dim V = 1$.

1. If $\bar{v} \in V$, $\bar{v} \neq 0$, then

$$A\bar{v} \in V \Rightarrow A\bar{v} \in \text{span } \bar{v} \Rightarrow A\bar{v} = \lambda\bar{v}$$

$\Rightarrow \bar{v}$ is an eigenvector of A .

2. If $A\bar{v} = \lambda\bar{v}$ then

$$A\bar{v} = \lambda\bar{v} \in \text{span } \bar{v} =$$

Invariant Subspaces

Definition: Let $A_{n \times n}$. A linear subspace $V \subset \mathbb{R}^n$ is **A-invariant** if

$$A\bar{v} \in V \quad \forall v \in V$$

Let $\dim V = 1$.

1. If $\bar{v} \in V$, $\bar{v} \neq 0$, then

$$A\bar{v} \in V \Rightarrow A\bar{v} \in \text{span } \bar{v} \Rightarrow A\bar{v} = \lambda\bar{v}$$

$\Rightarrow \bar{v}$ is an eigenvector of A .

2. If $A\bar{v} = \lambda\bar{v}$ then

$$A\bar{v} = \lambda\bar{v} \in \text{span } \bar{v} = V$$

\Rightarrow as $\dim V = 1$, so V is A -invariant.

Therefore, one-dimensional A -invariant subspaces V of \mathbb{R}^n are

$$V = \{\text{span } \bar{v} : A\bar{v} = \lambda\bar{v}\}$$

Eigenvalues and eigenfunctions

Definition: A scalar λ is called an **eigenvalue** of a linear operator $T: V \rightarrow V$ if

$$\exists f \in V, f \neq 0: Tf = \lambda f.$$

Eigenvalues and eigenfunctions

Definition: A scalar λ is called an **eigenvalue** of a linear operator $T: V \rightarrow V$ if

$$\exists f \in V, f \neq 0: Tf = \lambda f.$$

The element $f \in V$ is called an **eigenfunction**.

Eigenvalues and eigenfunctions

Definition: A scalar λ is called an **eigenvalue** of a linear operator $T: V \rightarrow V$ if

$$\exists f \in V, f \neq 0: Tf = \lambda f.$$

The element $f \in V$ is called an **eigenfunction**.

Examples:

1. $T(x_1, x_2, x_3, \dots) = (x_1, x_3, x_5, x_7, \dots)$

Eigenvalues and eigenfunctions

Definition: A scalar λ is called an **eigenvalue** of a linear operator $T: V \rightarrow V$ if

$$\exists f \in V, f \neq 0: Tf = \lambda f.$$

The element $f \in V$ is called an **eigenfunction**.

Examples:

1. $T(x_1, x_2, x_3, \dots) = (x_1, x_3, x_5, x_7, \dots)$

$$T(\bar{x}) = (x_1, x_3, x_5, x_7, \dots) = \lambda(x_1, x_2, x_3, x_4, x_5, \dots)$$

Eigenvalues and eigenfunctions

Definition: A scalar λ is called an **eigenvalue** of a linear operator $T: V \rightarrow V$ if

$$\exists f \in V, f \neq 0: Tf = \lambda f.$$

The element $f \in V$ is called an **eigenfunction**.

Examples:

1. $T(x_1, x_2, x_3, \dots) = (x_1, x_3, x_5, x_7, \dots)$

$$T(\bar{x}) = (x_1, x_3, x_5, x_7, \dots) = \lambda(x_1, x_2, x_3, x_4, x_5, \dots)$$

$$\bar{x} = (x_1 = \lambda x_1, x_2, x_3 = \lambda x_2, x_4, x_5 = \lambda x_3, \dots)$$

Eigenvalues and eigenfunctions

Definition: A scalar λ is called an **eigenvalue** of a linear operator $T: V \rightarrow V$ if

$$\exists f \in V, f \neq 0: Tf = \lambda f.$$

The element $f \in V$ is called an **eigenfunction**.

Examples:

1. $T(x_1, x_2, x_3, \dots) = (x_1, x_3, x_5, x_7, \dots)$

$$T(\bar{x}) = (x_1, x_3, x_5, x_7, \dots) = \lambda(x_1, x_2, x_3, x_4, x_5, \dots)$$

$$\bar{x} = (x_1 = \lambda x_1, x_2, x_3 = \lambda x_2, x_4, x_5 = \lambda x_3, \dots)$$

$$x_{2n} \text{ is arbitrary, } x_{2n+1} = \lambda x_{n+1}$$

Eigenvalues and eigenfunctions

Definition: A scalar λ is called an **eigenvalue** of a linear operator $T: V \rightarrow V$ if

$$\exists f \in V, f \neq 0: Tf = \lambda f.$$

The element $f \in V$ is called an **eigenfunction**.

Examples:

1. $T(x_1, x_2, x_3, \dots) = (x_1, x_3, x_5, x_7, \dots)$

$$T(\bar{x}) = (x_1, x_3, x_5, x_7, \dots) = \lambda(x_1, x_2, x_3, x_4, x_5, \dots)$$

$$\bar{x} = (x_1 = \lambda x_1, x_2, x_3 = \lambda x_2, x_4, x_5 = \lambda x_3, \dots)$$

$$x_{2n} \text{ is arbitrary, } x_{2n+1} = \lambda x_{n+1}$$

$$\lambda = 5 \Rightarrow \bar{x} = (0, x_2, 5x_2, x_4, 25x_2, x_5, \dots)$$

Eigenvalues and eigenfunctions

Definition: A scalar λ is called an **eigenvalue** of a linear operator $T: V \rightarrow V$ if

$$\exists f \in V, f \neq 0: Tf = \lambda f.$$

The element $f \in V$ is called an **eigenfunction**.

Examples:

1. $T(x_1, x_2, x_3, \dots) = (x_1, x_3, x_5, x_7, \dots)$

$$T(\bar{x}) = (x_1, x_3, x_5, x_7, \dots) = \lambda(x_1, x_2, x_3, x_4, x_5, \dots)$$

$$\bar{x} = (x_1 = \lambda x_1, x_2, x_3 = \lambda x_2, x_4, x_5 = \lambda x_3, \dots)$$

$$x_{2n} \text{ is arbitrary, } x_{2n+1} = \lambda x_{n+1}$$

$$\lambda = 5 \Rightarrow \bar{x} = (0, x_2, 5x_2, x_4, 25x_2, x_5, \dots)$$

All real λ are eigenvalues with infinite number of eigensequences .

Eigenvalues and eigenfunctions

2. $T(x_1, x_2, x_3, \dots) = (x_2, x_3, x_4, \dots)$

Eigenvalues and eigenfunctions

2. $T(x_1, x_2, x_3, \dots) = (x_2, x_3, x_4, \dots)$

$$T(\bar{x}) = (x_2, x_3, x_4, x_5, \dots) = (\lambda x_1, \lambda x_2, \lambda x_3, \lambda x_4, \lambda x_5, \dots)$$

Eigenvalues and eigenfunctions

2. $T(x_1, x_2, x_3, \dots) = (x_2, x_3, x_4, \dots)$

$$T(\bar{x}) = (x_2, x_3, x_4, x_5, \dots) = (\lambda x_1, \lambda x_2, \lambda x_3, \lambda x_4, \lambda x_5, \dots)$$

$$x_2 = \lambda x_1, \quad x_3 = \lambda x_2 = \lambda^2 x_1, \quad x_4 = \lambda x_3 = \lambda^3 x_1, \dots$$

Eigenvalues and eigenfunctions

2. $T(x_1, x_2, x_3, \dots) = (x_2, x_3, x_4, \dots)$

$$T(\bar{x}) = (x_2, x_3, x_4, x_5, \dots) = (\lambda x_1, \lambda x_2, \lambda x_3, \lambda x_4, \lambda x_5, \dots)$$

$$x_2 = \lambda x_1, \quad x_3 = \lambda x_2 = \lambda^2 x_1, \quad x_4 = \lambda x_3 = \lambda^3 x_1, \dots$$

$$\bar{x} = (x_1, x_2, x_3, \dots) = (x_1, \lambda x_1, \lambda^2 x_1, \dots) = x_1(1, \lambda, \lambda^2, \dots)$$

Eigenvalues and eigenfunctions

2. $T(x_1, x_2, x_3, \dots) = (x_2, x_3, x_4, \dots)$

$$T(\bar{x}) = (x_2, x_3, x_4, x_5, \dots) = (\lambda x_1, \lambda x_2, \lambda x_3, \lambda x_4, \lambda x_5, \dots)$$

$$x_2 = \lambda x_1, \quad x_3 = \lambda x_2 = \lambda^2 x_1, \quad x_4 = \lambda x_3 = \lambda^3 x_1, \dots$$

$$\bar{x} = (x_1, x_2, x_3, \dots) = (x_1, \lambda x_1, \lambda^2 x_1, \dots) = x_1(1, \lambda, \lambda^2, \dots)$$

All real λ are eigenvalues and $(1, \lambda, \lambda^2, \dots)$ is the basis in the space of eigensequences.

Eigenvalues of a Matrix

Theorem: A scalar $\lambda \in \mathbb{R}$ is an eigenvalue of the matrix $A_{n \times n}$ iff

$$\det(A - \lambda I_n) = 0.$$

The equation $f_A(\lambda) = \det(A - \lambda I_n) = 0$ is called the **characteristic equation** of the matrix A .

Remarks:

1. The eigenvalues of a diagonal matrix $A = \text{diag}(d_1, \dots, d_n)$ are d_1, \dots, d_n .

Eigenvalues of a Matrix

Theorem: A scalar $\lambda \in \mathbb{R}$ is an eigenvalue of the matrix $A_{n \times n}$ iff

$$\det(A - \lambda I_n) = 0.$$

The equation $f_A(\lambda) = \det(A - \lambda I_n) = 0$ is called the **characteristic equation** of the matrix A .

Remarks:

1. The eigenvalues of a diagonal matrix $A = \text{diag}(d_1, \dots, d_n)$ are d_1, \dots, d_n .
2. $\det(A - \lambda I_n) = (-\lambda)^n + (\text{tr } A)(-\lambda)^{n-1} + \dots + \det A$

Eigenvalues of a Matrix

Theorem: A scalar $\lambda \in \mathbb{R}$ is an eigenvalue of the matrix $A_{n \times n}$ iff

$$\det(A - \lambda I_n) = 0.$$

The equation $f_A(\lambda) = \det(A - \lambda I_n) = 0$ is called the **characteristic equation** of the matrix A .

Remarks:

1. The eigenvalues of a diagonal matrix $A = \text{diag}(d_1, \dots, d_n)$ are d_1, \dots, d_n .
2. $\det(A - \lambda I_n) = (-\lambda)^n + (\text{tr } A)(-\lambda)^{n-1} + \dots + \det A$
3. There are n eigenvalues of a $n \times n$ matrix A .

Eigenvalues of a Matrix

Theorem: A scalar $\lambda \in \mathbb{R}$ is an eigenvalue of the matrix $A_{n \times n}$ iff

$$\det(A - \lambda I_n) = 0.$$

The equation $f_A(\lambda) = \det(A - \lambda I_n) = 0$ is called the **characteristic equation** of the matrix A .

Remarks:

1. The eigenvalues of a diagonal matrix $A = \text{diag}(d_1, \dots, d_n)$ are d_1, \dots, d_n .
2. $\det(A - \lambda I_n) = (-\lambda)^n + (\text{tr } A)(-\lambda)^{n-1} + \dots + \det A$
3. There are n eigenvalues of a $n \times n$ matrix A .
4. For $A_{n \times n}$, with eigenvalues $\lambda_1, \dots, \lambda_n$ listed with their algebraic multiplicities

$$\det A = \lambda_1 \cdot \lambda_2 \cdots \lambda_n, \quad \text{tr } A = \lambda_1 + \lambda_2 + \dots + \lambda_n$$

Algebraic Multiplicity

Definition: An eigenvalue λ_0 of a square matrix A has **algebraic multiplicity** k if λ_0 is the root of multiplicity k of the characteristic polynomial $f_A(\lambda) \Rightarrow f_A(\lambda) = (\lambda - \lambda_0)^k g(\lambda)$, $g(\lambda_0) \neq 0$

Algebraic Multiplicity

Definition: An eigenvalue λ_0 of a square matrix A has **algebraic multiplicity** k if λ_0 is the root of multiplicity k of the characteristic polynomial $f_A(\lambda) \Rightarrow f_A(\lambda) = (\lambda - \lambda_0)^k g(\lambda)$, $g(\lambda_0) \neq 0$

Examples:

$$1. \quad A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \Rightarrow \det(A - \lambda I_3) = 0 \Rightarrow \lambda_{1,2} = 0, \lambda_3 = 3$$

Algebraic Multiplicity

Definition: An eigenvalue λ_0 of a square matrix A has **algebraic multiplicity** k if λ_0 is the root of multiplicity k of the characteristic polynomial $f_A(\lambda) \Rightarrow f_A(\lambda) = (\lambda - \lambda_0)^k g(\lambda)$, $g(\lambda_0) \neq 0$

Examples:

$$1. \quad A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \Rightarrow \det(A - \lambda I_3) = 0 \Rightarrow \lambda_{1,2} = 0, \lambda_3 = 3$$

$$2. \quad A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \det(A - \lambda I_2) = 0$$

$$\Rightarrow \lambda^2 - \underbrace{(1+3)}_{\text{tr } A} \lambda + \underbrace{1 \cdot 3 - 2 \cdot 4}_{\det A} = 0$$

Algebraic Multiplicity

Definition: An eigenvalue λ_0 of a square matrix A has **algebraic multiplicity** k if λ_0 is the root of multiplicity k of the characteristic polynomial $f_A(\lambda) \Rightarrow f_A(\lambda) = (\lambda - \lambda_0)^k g(\lambda)$, $g(\lambda_0) \neq 0$

Examples:

$$1. \quad A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \Rightarrow \det(A - \lambda I_3) = 0 \Rightarrow \lambda_{1,2} = 0, \lambda_3 = 3$$

$$2. \quad A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \det(A - \lambda I_2) = 0$$

$$\Rightarrow \lambda^2 - \underbrace{(1+3)}_{\text{tr } A} \lambda + \underbrace{1 \cdot 3 - 2 \cdot 4}_{\det A} = 0$$

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the eigenspace E_λ associated with λ :

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the eigenspace E_λ associated with λ :

$$E_\lambda = \text{Ker}(A - \lambda I_n) = \{\bar{v} \in \mathbb{R}^n : A\bar{v} = \lambda\bar{v}, \bar{v} \neq 0\}$$

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the **eigenspace** E_λ associated with λ :

$$E_\lambda = \text{Ker}(A - \lambda I_n) = \{\bar{v} \in \mathbb{R}^n : A\bar{v} = \lambda\bar{v}, \bar{v} \neq 0\}$$

Examples:

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the **eigenspace** E_λ associated with λ :

$$E_\lambda = \text{Ker}(A - \lambda I_n) = \{\bar{v} \in \mathbb{R}^n : A\bar{v} = \lambda\bar{v}, \bar{v} \neq 0\}$$

Examples:

1. $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is an orthogonal projection onto a plane V in \mathbb{R}^3

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the eigenspace E_λ associated with λ :

$$E_\lambda = \text{Ker}(A - \lambda I_n) = \{\bar{v} \in \mathbb{R}^n : A\bar{v} = \lambda\bar{v}, \bar{v} \neq 0\}$$

Examples:

1. $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is an orthogonal projection onto a plane V in \mathbb{R}^3

$$\lambda = 1 \Rightarrow E_1 = V \quad \text{OR} \quad \lambda = 0 \Rightarrow E_0 = V^\perp$$

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the **eigenspace** E_λ associated with λ :

$$E_\lambda = \text{Ker}(A - \lambda I_n) = \{\bar{v} \in \mathbb{R}^n : A\bar{v} = \lambda\bar{v}, \bar{v} \neq 0\}$$

Examples:

1. $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is an orthogonal projection onto a plane V in \mathbb{R}^3

$$\lambda = 1 \Rightarrow E_1 = V \quad \text{OR} \quad \lambda = 0 \Rightarrow E_0 = V^\perp$$

2. $A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow$

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the eigenspace E_λ associated with λ :

$$E_\lambda = \text{Ker}(A - \lambda I_n) = \{\bar{v} \in \mathbb{R}^n : A\bar{v} = \lambda\bar{v}, \bar{v} \neq 0\}$$

Examples:

1. $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is an orthogonal projection onto a plane V in \mathbb{R}^3

$$\lambda = 1 \Rightarrow E_1 = V \quad \text{OR} \quad \lambda = 0 \Rightarrow E_0 = V^\perp$$

2. $A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = 5, \lambda_2 = -1$

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the **eigenspace** E_λ associated with λ :

$$E_\lambda = \text{Ker}(A - \lambda I_n) = \{\bar{v} \in \mathbb{R}^n : A\bar{v} = \lambda\bar{v}, \bar{v} \neq 0\}$$

Examples:

1. $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is an orthogonal projection onto a plane V in \mathbb{R}^3

$$\lambda = 1 \Rightarrow E_1 = V \quad \text{OR} \quad \lambda = 0 \Rightarrow E_0 = V^\perp$$

2. $A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = 5, \lambda_2 = -1$

$$E_5 = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = 5\bar{v}\} =$$

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the **eigenspace** E_λ associated with λ :

$$E_\lambda = \text{Ker}(A - \lambda I_n) = \{\bar{v} \in \mathbb{R}^n : A\bar{v} = \lambda\bar{v}, \bar{v} \neq 0\}$$

Examples:

1. $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is an orthogonal projection onto a plane V in \mathbb{R}^3

$$\lambda = 1 \Rightarrow E_1 = V \quad \text{OR} \quad \lambda = 0 \Rightarrow E_0 = V^\perp$$

2. $A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = 5, \lambda_2 = -1$

$$E_5 = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = 5\bar{v}\} = \text{Ker} \begin{pmatrix} -4 & 2 \\ 4 & -2 \end{pmatrix} =$$

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the **eigenspace** E_λ associated with λ :

$$E_\lambda = \text{Ker}(A - \lambda I_n) = \{\bar{v} \in \mathbb{R}^n : A\bar{v} = \lambda\bar{v}, \bar{v} \neq 0\}$$

Examples:

1. $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is an orthogonal projection onto a plane V in \mathbb{R}^3

$$\lambda = 1 \Rightarrow E_1 = V \quad \text{OR} \quad \lambda = 0 \Rightarrow E_0 = V^\perp$$

2. $A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = 5, \lambda_2 = -1$

$$E_5 = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = 5\bar{v}\} = \text{Ker} \begin{pmatrix} -4 & 2 \\ 4 & -2 \end{pmatrix} = \text{span} \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the **eigenspace** E_λ associated with λ :

$$E_\lambda = \text{Ker}(A - \lambda I_n) = \{\bar{v} \in \mathbb{R}^n : A\bar{v} = \lambda\bar{v}, \bar{v} \neq 0\}$$

Examples:

1. $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is an orthogonal projection onto a plane V in \mathbb{R}^3

$$\lambda = 1 \Rightarrow E_1 = V \quad \text{OR} \quad \lambda = 0 \Rightarrow E_0 = V^\perp$$

2. $A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = 5, \lambda_2 = -1$

$$E_5 = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = 5\bar{v}\} = \text{Ker} \begin{pmatrix} -4 & 2 \\ 4 & -2 \end{pmatrix} = \text{span} \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$E_{-1} = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = -\bar{v}\} =$$

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the **eigenspace** E_λ associated with λ :

$$E_\lambda = \text{Ker}(A - \lambda I_n) = \{\bar{v} \in \mathbb{R}^n : A\bar{v} = \lambda\bar{v}, \bar{v} \neq 0\}$$

Examples:

1. $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is an orthogonal projection onto a plane V in \mathbb{R}^3

$$\lambda = 1 \Rightarrow E_1 = V \quad \text{OR} \quad \lambda = 0 \Rightarrow E_0 = V^\perp$$

2. $A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = 5, \lambda_2 = -1$

$$E_5 = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = 5\bar{v}\} = \text{Ker} \begin{pmatrix} -4 & 2 \\ 4 & -2 \end{pmatrix} = \text{span} \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$E_{-1} = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = -\bar{v}\} = \text{Ker} \begin{pmatrix} 2 & 2 \\ 4 & 4 \end{pmatrix} =$$

Eigenspace

Definition: The kernel of the matrix $A - \lambda I_n$ is called the **eigenspace** E_λ associated with λ :

$$E_\lambda = \text{Ker}(A - \lambda I_n) = \{\bar{v} \in \mathbb{R}^n : A\bar{v} = \lambda\bar{v}, \bar{v} \neq 0\}$$

Examples:

1. $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is an orthogonal projection onto a plane V in \mathbb{R}^3

$$\lambda = 1 \Rightarrow E_1 = V \quad \text{OR} \quad \lambda = 0 \Rightarrow E_0 = V^\perp$$

2. $A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = 5, \lambda_2 = -1$

$$E_5 = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = 5\bar{v}\} = \text{Ker} \begin{pmatrix} -4 & 2 \\ 4 & -2 \end{pmatrix} = \text{span} \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$E_{-1} = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = -\bar{v}\} = \text{Ker} \begin{pmatrix} 2 & 2 \\ 4 & 4 \end{pmatrix} = \text{span} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

Eigenspace

3. $A = \begin{pmatrix} -2 & -1 \\ 5 & 2 \end{pmatrix} \Rightarrow \lambda^2 = -1 \Rightarrow$

Eigenspace

3. $A = \begin{pmatrix} -2 & -1 \\ 5 & 2 \end{pmatrix} \Rightarrow \lambda^2 = -1 \Rightarrow \lambda_{1,2} = \pm i$

$$E_i = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = i\bar{v}\} =$$

Eigenspace

$$3. A = \begin{pmatrix} -2 & -1 \\ 5 & 2 \end{pmatrix} \Rightarrow \lambda^2 = -1 \Rightarrow \lambda_{1,2} = \pm i$$

$$E_i = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = i\bar{v}\} = \text{Ker} \begin{pmatrix} -2-i & -1 \\ 5 & 2-i \end{pmatrix}$$

Eigenspace

$$3. A = \begin{pmatrix} -2 & -1 \\ 5 & 2 \end{pmatrix} \Rightarrow \lambda^2 = -1 \Rightarrow \lambda_{1,2} = \pm i$$

$$E_i = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = i\bar{v}\} = \text{Ker} \begin{pmatrix} -2-i & -1 \\ 5 & 2-i \end{pmatrix}$$

$$\begin{pmatrix} -2-i & -1 \\ 5 & 2-i \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Eigenspace

$$3. A = \begin{pmatrix} -2 & -1 \\ 5 & 2 \end{pmatrix} \Rightarrow \lambda^2 = -1 \Rightarrow \lambda_{1,2} = \pm i$$

$$E_i = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = i\bar{v}\} = \text{Ker} \begin{pmatrix} -2-i & -1 \\ 5 & 2-i \end{pmatrix}$$

$$\begin{pmatrix} -2-i & -1 \\ 5 & 2-i \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$-(2+i)v_1 - v_2 = 0 \Rightarrow \bar{v} = \begin{pmatrix} -1 \\ 2+i \end{pmatrix}$$

Eigenspace

$$3. A = \begin{pmatrix} -2 & -1 \\ 5 & 2 \end{pmatrix} \Rightarrow \lambda^2 = -1 \Rightarrow \lambda_{1,2} = \pm i$$

$$E_i = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = i\bar{v}\} = \text{Ker} \begin{pmatrix} -2-i & -1 \\ 5 & 2-i \end{pmatrix}$$

$$\begin{pmatrix} -2-i & -1 \\ 5 & 2-i \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$-(2+i)v_1 - v_2 = 0 \Rightarrow \bar{v} = \begin{pmatrix} -1 \\ 2+i \end{pmatrix}$$

$$E_i = \text{span} \left(\begin{pmatrix} -1 \\ 2+i \end{pmatrix} \right)$$

Eigenspace

$$3. A = \begin{pmatrix} -2 & -1 \\ 5 & 2 \end{pmatrix} \Rightarrow \lambda^2 = -1 \Rightarrow \lambda_{1,2} = \pm i$$

$$E_i = \{\bar{v} \in \mathbb{R}^2 : A\bar{v} = i\bar{v}\} = \text{Ker} \begin{pmatrix} -2-i & -1 \\ 5 & 2-i \end{pmatrix}$$

$$\begin{pmatrix} -2-i & -1 \\ 5 & 2-i \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$-(2+i)v_1 - v_2 = 0 \Rightarrow \bar{v} = \begin{pmatrix} -1 \\ 2+i \end{pmatrix}$$

$$E_i = \text{span} \left(\begin{pmatrix} -1 \\ 2+i \end{pmatrix} \right)$$

Remark: If λ is an eigenvalue of a real matrix $A_{n \times n}$ with the associated eigenvector \bar{v} , then $\bar{\lambda}$ is also an eigenvalue of A whose associated eigenvector \bar{v}^* is the complex conjugate of \bar{v} .

Geometric Multiplicity

4. $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow$

Geometric Multiplicity

4. $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_{1,2} = 1, \lambda_3 = 0$

Geometric Multiplicity

4. $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_{1,2} = 1, \lambda_3 = 0 \Rightarrow AM_1 = 2, AM_0 = 1$

Geometric Multiplicity

4. $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_{1,2} = 1, \lambda_3 = 0 \Rightarrow AM_1 = 2, AM_0 = 1$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & 1 & 1 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \end{pmatrix} =$$

Geometric Multiplicity

$$4. \quad A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_{1,2} = 1, \lambda_3 = 0 \Rightarrow AM_1 = 2, AM_0 = 1$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & 1 & 1 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \end{pmatrix} = \text{span} \left(\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right) \Rightarrow$$

Geometric Multiplicity

$$4. \quad A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_{1,2} = 1, \lambda_3 = 0 \Rightarrow AM_1 = 2, AM_0 = 1$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & 1 & 1 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \end{pmatrix} = \text{span} \left(\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right) \Rightarrow \dim E_1 = 1$$

Geometric Multiplicity

$$4. \quad A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_{1,2} = 1, \lambda_3 = 0 \Rightarrow AM_1 = 2, AM_0 = 1$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & 1 & 1 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \end{pmatrix} = \text{span} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \Rightarrow \dim E_1 = 1$$

$$E_0 = \text{Ker} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} =$$

Geometric Multiplicity

$$4. \quad A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_{1,2} = 1, \lambda_3 = 0 \Rightarrow AM_1 = 2, AM_0 = 1$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & 1 & 1 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \end{pmatrix} = \text{span} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \Rightarrow \dim E_1 = 1$$

$$E_0 = \text{Ker} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} = \text{span} \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} \Rightarrow$$

Geometric Multiplicity

$$4. \quad A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_{1,2} = 1, \lambda_3 = 0 \Rightarrow AM_1 = 2, AM_0 = 1$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & 1 & 1 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \end{pmatrix} = \text{span} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \Rightarrow \dim E_1 = 1$$

$$E_0 = \text{Ker} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} = \text{span} \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} \Rightarrow \dim E_0 = 1$$

Geometric Multiplicity

$$4. \quad A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_{1,2} = 1, \lambda_3 = 0 \Rightarrow AM_1 = 2, AM_0 = 1$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & 1 & 1 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \end{pmatrix} = \text{span} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \Rightarrow \dim E_1 = 1$$

$$E_0 = \text{Ker} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} = \text{span} \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} \Rightarrow \dim E_0 = 1$$

Definition: The dimension of eigenspace E_λ is called the **geometric multiplicity** of the eigenvalue λ .

$$G.M. = n - \text{rank}(A - \lambda I_n)$$

Eigenbasis

Theorem:

1. The system $\bar{v}_1, \dots, \bar{v}_s$ consisting of all basis vectors of each eigenspace of $A_{n \times n}$ is **linearly independent**; s is the sum of all geometric multiplicities of eigenvalues of A .
2. There exists an eigenbasis for $A_{n \times n}$ iff the geometric multiplicities add up to $n \Rightarrow s = n$.

Eigenbasis

Theorem:

1. The system $\bar{v}_1, \dots, \bar{v}_s$ consisting of all basis vectors of each eigenspace of $A_{n \times n}$ is **linearly independent**; s is the sum of all geometric multiplicities of eigenvalues of A .
2. There exists an eigenbasis for $A_{n \times n}$ iff the geometric multiplicities add up to $n \Rightarrow s = n$.

Remark: If $A_{n \times n}$ has n distinct eigenvalues then there exists an eigenbasis for A .

Eigenbasis

Theorem:

1. The system $\vec{v}_1, \dots, \vec{v}_s$ consisting of all basis vectors of each eigenspace of $A_{n \times n}$ is **linearly independent**; s is the sum of all geometric multiplicities of eigenvalues of A .
2. There exists an eigenbasis for $A_{n \times n}$ iff the geometric multiplicities add up to $n \Rightarrow s = n$.

Remark: If $A_{n \times n}$ has n distinct eigenvalues then there exists an eigenbasis for A .

Example: Let

$$T: \mathbb{R}^2 \rightarrow \mathbb{R}^2, T\vec{x} = T \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_1 + x_2 \\ x_2 \end{pmatrix}$$

Eigenbasis

Theorem:

1. The system $\vec{v}_1, \dots, \vec{v}_s$ consisting of all basis vectors of each eigenspace of $A_{n \times n}$ is **linearly independent**; s is the sum of all geometric multiplicities of eigenvalues of A .
2. There exists an eigenbasis for $A_{n \times n}$ iff the geometric multiplicities add up to $n \Rightarrow s = n$.

Remark: If $A_{n \times n}$ has n distinct eigenvalues then there exists an eigenbasis for A .

Example: Let

$$T: \mathbb{R}^2 \rightarrow \mathbb{R}^2, T\vec{x} = T \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_1 + x_2 \\ x_2 \end{pmatrix}$$

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \Rightarrow \lambda_{1,2} = 1 \quad \text{BUT} \quad \text{Ker}(A - \lambda I_2) = \text{span} \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix} \right)$$

$$\dim E_1 = 1 \Rightarrow G.M. = 1 < 2 = n$$

Similar Matrices

Theorem: Let $B = S^{-1}AS$, i.e. A, B be similar matrices. Then

1. $f_A(\lambda) = f_B(\lambda)$
2. $\text{rank}(A) = \text{rank}(B)$, $\text{nullity}(A) = \text{nullity}(B)$
3. A, B have the same eigenvalues with the same AM, GM
4. $\det A = \det B$, $\text{tr}(A) = \text{tr}(B)$

Proof:

1. $f_B(\lambda) = \det(B - \lambda I) =$

Similar Matrices

Theorem: Let $B = S^{-1}AS$, i.e. A, B be similar matrices. Then

1. $f_A(\lambda) = f_B(\lambda)$
2. $\text{rank}(A) = \text{rank}(B)$, $\text{nullity}(A) = \text{nullity}(B)$
3. A, B have the same eigenvalues with the same AM, GM
4. $\det A = \det B$, $\text{tr}(A) = \text{tr}(B)$

Proof:

$$\begin{aligned} 1. \quad f_B(\lambda) &= \det(B - \lambda I) = \det(S^{-1}AS - \lambda S^{-1}S) \\ &= \end{aligned}$$

Similar Matrices

Theorem: Let $B = S^{-1}AS$, i.e. A, B be similar matrices. Then

1. $f_A(\lambda) = f_B(\lambda)$
2. $\text{rank}(A) = \text{rank}(B)$, $\text{nullity}(A) = \text{nullity}(B)$
3. A, B have the same eigenvalues with the same AM, GM
4. $\det A = \det B$, $\text{tr}(A) = \text{tr}(B)$

Proof:

$$\begin{aligned} 1. \quad f_B(\lambda) &= \det(B - \lambda I) = \det(S^{-1}AS - \lambda S^{-1}S) \\ &= \det(S^{-1}(A - \lambda I)S) = \end{aligned}$$

Similar Matrices

Theorem: Let $B = S^{-1}AS$, i.e. A, B be similar matrices. Then

1. $f_A(\lambda) = f_B(\lambda)$
2. $\text{rank}(A) = \text{rank}(B)$, $\text{nullity}(A) = \text{nullity}(B)$
3. A, B have the same eigenvalues with the same AM, GM
4. $\det A = \det B$, $\text{tr}(A) = \text{tr}(B)$

Proof:

$$\begin{aligned} 1. \quad f_B(\lambda) &= \det(B - \lambda I) = \det(S^{-1}AS - \lambda S^{-1}S) \\ &= \det(S^{-1}(A - \lambda I)S) = \det S^{-1} \det(A - \lambda I) \det S = \end{aligned}$$

Similar Matrices

Theorem: Let $B = S^{-1}AS$, i.e. A, B be similar matrices. Then

1. $f_A(\lambda) = f_B(\lambda)$
2. $\text{rank}(A) = \text{rank}(B)$, $\text{nullity}(A) = \text{nullity}(B)$
3. A, B have the same eigenvalues with the same AM, GM
4. $\det A = \det B$, $\text{tr}(A) = \text{tr}(B)$

Proof:

$$\begin{aligned} 1. \quad f_B(\lambda) &= \det(B - \lambda I) = \det(S^{-1}AS - \lambda S^{-1}S) \\ &= \det(S^{-1}(A - \lambda I)S) = \det S^{-1} \det(A - \lambda I) \det S = f_A(\lambda) \end{aligned}$$

Similar Matrices

Theorem: Let $B = S^{-1}AS$, i.e. A, B be similar matrices. Then

1. $f_A(\lambda) = f_B(\lambda)$
2. $\text{rank}(A) = \text{rank}(B)$, $\text{nullity}(A) = \text{nullity}(B)$
3. A, B have the same eigenvalues with the same AM, GM
4. $\det A = \det B$, $\text{tr}(A) = \text{tr}(B)$

Proof:

1. $f_B(\lambda) = \det(B - \lambda I) = \det(S^{-1}AS - \lambda S^{-1}S)$
 $= \det(S^{-1}(A - \lambda I)S) = \det S^{-1} \det(A - \lambda I) \det S = f_A(\lambda)$
2. Let $\bar{x} \in \text{Ker } B \Rightarrow B\bar{x} = \vec{0} \Rightarrow$

Similar Matrices

Theorem: Let $B = S^{-1}AS$, i.e. A, B be similar matrices. Then

1. $f_A(\lambda) = f_B(\lambda)$
2. $\text{rank}(A) = \text{rank}(B)$, $\text{nullity}(A) = \text{nullity}(B)$
3. A, B have the same eigenvalues with the same AM, GM
4. $\det A = \det B$, $\text{tr}(A) = \text{tr}(B)$

Proof:

1. $f_B(\lambda) = \det(B - \lambda I) = \det(S^{-1}AS - \lambda S^{-1}S)$
 $= \det(S^{-1}(A - \lambda I)S) = \det S^{-1} \det(A - \lambda I) \det S = f_A(\lambda)$
2. Let $\bar{x} \in \text{Ker } B \Rightarrow B\bar{x} = \vec{0} \Rightarrow S^{-1}AS\bar{x} = \vec{0} \Rightarrow$

Similar Matrices

Theorem: Let $B = S^{-1}AS$, i.e. A, B be similar matrices. Then

1. $f_A(\lambda) = f_B(\lambda)$
2. $\text{rank}(A) = \text{rank}(B)$, $\text{nullity}(A) = \text{nullity}(B)$
3. A, B have the same eigenvalues with the same AM, GM
4. $\det A = \det B$, $\text{tr}(A) = \text{tr}(B)$

Proof:

1. $f_B(\lambda) = \det(B - \lambda I) = \det(S^{-1}AS - \lambda S^{-1}S)$
 $= \det(S^{-1}(A - \lambda I)S) = \det S^{-1} \det(A - \lambda I) \det S = f_A(\lambda)$
2. Let $\bar{x} \in \text{Ker } B \Rightarrow B\bar{x} = \vec{0} \Rightarrow S^{-1}AS\bar{x} = \vec{0} \Rightarrow S\bar{x} \in \text{Ker } A$

$$T : \text{Ker } B \rightarrow \text{Ker } A \quad T\bar{x} = S\bar{x}$$

T is linear, one-to-one and onto $\Rightarrow T$ is an isomorphism.

Similar Matrices

Theorem: Let $B = S^{-1}AS$, i.e. A, B be similar matrices. Then

1. $f_A(\lambda) = f_B(\lambda)$
2. $\text{rank}(A) = \text{rank}(B)$, $\text{nullity}(A) = \text{nullity}(B)$
3. A, B have the same eigenvalues with the same AM, GM
4. $\det A = \det B$, $\text{tr}(A) = \text{tr}(B)$

Proof:

1. $f_B(\lambda) = \det(B - \lambda I) = \det(S^{-1}AS - \lambda S^{-1}S)$
 $= \det(S^{-1}(A - \lambda I)S) = \det S^{-1} \det(A - \lambda I) \det S = f_A(\lambda)$
2. Let $\bar{x} \in \text{Ker } B \Rightarrow B\bar{x} = \vec{0} \Rightarrow S^{-1}AS\bar{x} = \vec{0} \Rightarrow S\bar{x} \in \text{Ker } A$

$$T : \text{Ker } B \rightarrow \text{Ker } A \quad T\bar{x} = S\bar{x}$$

T is linear, one-to-one and onto $\Rightarrow T$ is an isomorphism.

$$\Rightarrow \dim \text{Ker } B = \dim \text{Ker } A$$

Similar Matrices

Theorem: Let $B = S^{-1}AS$, i.e. A, B be similar matrices. Then

1. $f_A(\lambda) = f_B(\lambda)$
2. $\text{rank}(A) = \text{rank}(B)$, $\text{nullity}(A) = \text{nullity}(B)$
3. A, B have the same eigenvalues with the same AM, GM
4. $\det A = \det B$, $\text{tr}(A) = \text{tr}(B)$

Proof:

1. $f_B(\lambda) = \det(B - \lambda I) = \det(S^{-1}AS - \lambda S^{-1}S)$
 $= \det(S^{-1}(A - \lambda I)S) = \det S^{-1} \det(A - \lambda I) \det S = f_A(\lambda)$
2. Let $\bar{x} \in \text{Ker } B \Rightarrow B\bar{x} = \vec{0} \Rightarrow S^{-1}AS\bar{x} = \vec{0} \Rightarrow S\bar{x} \in \text{Ker } A$

$$T : \text{Ker } B \rightarrow \text{Ker } A \quad T\bar{x} = S\bar{x}$$

T is linear, one-to-one and onto $\Rightarrow T$ is an isomorphism.

$$\Rightarrow \dim \text{Ker } B = \dim \text{Ker } A$$

Also, if $A_{n \times n}$, $B_{n \times n}$, then $n = \dim \text{Ker } A + \dim \text{Im } A$

Motivation

$$A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = -1, \lambda_2 = 5 \Rightarrow \bar{v}_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \bar{v}_2 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$\Rightarrow \mathfrak{B} = \{\bar{v}_1, \bar{v}_2\}$ is the eigenbasis for A .

Motivation

$$A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = -1, \lambda_2 = 5 \Rightarrow \bar{v}_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \bar{v}_2 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$\Rightarrow \mathfrak{B} = \{\bar{v}_1, \bar{v}_2\}$ is the eigenbasis for A .

$$\bar{x} = \alpha_1 \bar{v}_1 + \alpha_2 \bar{v}_2 \xrightarrow{A} T\bar{x} = -\alpha_1 \bar{v}_1 + 5\alpha_2 \bar{v}_2$$

Motivation

$$A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = -1, \lambda_2 = 5 \Rightarrow \bar{v}_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \bar{v}_2 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$\Rightarrow \mathfrak{B} = \{\bar{v}_1, \bar{v}_2\}$ is the eigenbasis for A .

$$\begin{array}{ccc} \bar{x} = \alpha_1 \bar{v}_1 + \alpha_2 \bar{v}_2 & \xrightarrow{A} & T\bar{x} = -\alpha_1 \bar{v}_1 + 5\alpha_2 \bar{v}_2 \\ \downarrow & & \downarrow \\ \bar{x}_{\mathfrak{B}} = \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix} & \xrightarrow{B} & (T\bar{x})_{\mathfrak{B}} = \begin{pmatrix} -\alpha_1 \\ 5\alpha_2 \end{pmatrix} \end{array}$$

Motivation

$$A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = -1, \lambda_2 = 5 \Rightarrow \bar{v}_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \bar{v}_2 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$\Rightarrow \mathfrak{B} = \{\bar{v}_1, \bar{v}_2\}$ is the eigenbasis for A .

$$\bar{x} = \alpha_1 \bar{v}_1 + \alpha_2 \bar{v}_2 \xrightarrow{A} T\bar{x} = -\alpha_1 \bar{v}_1 + 5\alpha_2 \bar{v}_2$$

$$\begin{array}{ccc} \downarrow & & \downarrow \\ \bar{x}_{\mathfrak{B}} = \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix} & \xrightarrow{B} & (T\bar{x})_{\mathfrak{B}} = \begin{pmatrix} -\alpha_1 \\ 5\alpha_2 \end{pmatrix} \end{array}$$

$$B = \begin{pmatrix} -1 & 0 \\ 0 & 5 \end{pmatrix}$$

B is diagonal \Rightarrow we denote $B = D$, $\mathfrak{D} = \{\bar{v}_1, \bar{v}_2\}$

Diagonalizable Matrices

Theorem: Consider a linear transformation $T\bar{x} = A\bar{x}$, $A_{n \times n}$. Let $\mathfrak{D} = \{\bar{v}_1, \bar{v}_2, \dots, \bar{v}_n\}$ be an eigenbasis for T : $A\bar{v}_i = \lambda_i\bar{v}_i$. Then the \mathfrak{D} -matrix D of T is

$$D = S^{-1}AS = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}$$

Diagonalizable Matrices

Theorem: Consider a linear transformation $T\bar{x} = A\bar{x}$, $A_{n \times n}$. Let $\mathfrak{D} = \{\bar{v}_1, \bar{v}_2, \dots, \bar{v}_n\}$ be an eigenbasis for T : $A\bar{v}_i = \lambda_i\bar{v}_i$. Then the \mathfrak{D} -matrix D of T is

$$D = S^{-1}AS = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}$$

Definition: An $n \times n$ matrix A is called **diagonalizable** if A is similar to a diagonal matrix D :

$$\exists \text{ invertible } S_{n \times n}: D = S^{-1}AS$$

Diagonalizable Matrices

Theorem: Consider a linear transformation $T\bar{x} = A\bar{x}$, $A_{n \times n}$. Let $\mathfrak{D} = \{\bar{v}_1, \bar{v}_2, \dots, \bar{v}_n\}$ be an eigenbasis for T : $A\bar{v}_i = \lambda_i\bar{v}_i$. Then the \mathfrak{D} -matrix D of T is

$$D = S^{-1}AS = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}$$

Definition: An $n \times n$ matrix A is called **diagonalizable** if A is similar to a diagonal matrix D :

$$\exists \text{ invertible } S_{n \times n}: D = S^{-1}AS$$

Remark: The matrix $S^{-1}AS$ is diagonal iff the columns of S form an eigenbasis of A .

Theorem

1. A matrix A is diagonalizable iff there exists an eigenbasis for A .
2. If $A_{n \times n}$ has n distinct eigenvalues then A is diagonalizable.

Matrix Diagonalization: Example 1

$$A = \begin{pmatrix} 1 & a & b \\ 0 & 0 & c \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_1 = 0, \lambda_{2,3} = 1$$

Matrix Diagonalization: Example 1

$$A = \begin{pmatrix} 1 & a & b \\ 0 & 0 & c \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_1 = 0, \lambda_{2,3} = 1$$

$$E_0 = \text{Ker } A \Rightarrow \dim E_0 = \dim \text{Ker } A = 1 \text{ with } \bar{v}_1 = \begin{pmatrix} -a \\ 1 \\ 0 \end{pmatrix},$$

Matrix Diagonalization: Example 1

$$A = \begin{pmatrix} 1 & a & b \\ 0 & 0 & c \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_1 = 0, \lambda_{2,3} = 1$$

$$E_0 = \text{Ker } A \Rightarrow \dim E_0 = \dim \text{Ker } A = 1 \text{ with } \bar{v}_1 = \begin{pmatrix} -a \\ 1 \\ 0 \end{pmatrix},$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & a & b \\ 0 & -1 & c \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow$$

Matrix Diagonalization: Example 1

$$A = \begin{pmatrix} 1 & a & b \\ 0 & 0 & c \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_1 = 0, \lambda_{2,3} = 1$$

$$E_0 = \text{Ker } A \Rightarrow \dim E_0 = \dim \text{Ker } A = 1 \text{ with } \bar{v}_1 = \begin{pmatrix} -a \\ 1 \\ 0 \end{pmatrix},$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & a & b \\ 0 & -1 & c \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \dim E_1 = 2 \text{ iff } b = -ac \text{ with}$$

$$\bar{v}_2 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix},$$

Matrix Diagonalization: Example 1

$$A = \begin{pmatrix} 1 & a & b \\ 0 & 0 & c \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_1 = 0, \lambda_{2,3} = 1$$

$$E_0 = \text{Ker } A \Rightarrow \dim E_0 = \dim \text{Ker } A = 1 \text{ with } \bar{v}_1 = \begin{pmatrix} -a \\ 1 \\ 0 \end{pmatrix},$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & a & b \\ 0 & -1 & c \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \dim E_1 = 2 \text{ iff } b = -ac \text{ with}$$

$$\bar{v}_2 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \bar{v}_3 = \begin{pmatrix} 0 \\ c \\ 1 \end{pmatrix}$$

$$\dim E_0 + \dim E_1 = 3 = n \Rightarrow A \text{ is diagonalizable}$$

Matrix Diagonalization: Example 1

$$A = \begin{pmatrix} 1 & a & b \\ 0 & 0 & c \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_1 = 0, \lambda_{2,3} = 1$$

$$E_0 = \text{Ker } A \Rightarrow \dim E_0 = \dim \text{Ker } A = 1 \text{ with } \bar{v}_1 = \begin{pmatrix} -a \\ 1 \\ 0 \end{pmatrix},$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & a & b \\ 0 & -1 & c \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \dim E_1 = 2 \text{ iff } b = -ac \text{ with}$$

$$\bar{v}_2 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \bar{v}_3 = \begin{pmatrix} 0 \\ c \\ 1 \end{pmatrix}$$

$$\dim E_0 + \dim E_1 = 3 = n \Rightarrow A \text{ is diagonalizable}$$

$$S = \begin{pmatrix} -a & 1 & 0 \\ 1 & 0 & c \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow$$

Matrix Diagonalization: Example 1

$$A = \begin{pmatrix} 1 & a & b \\ 0 & 0 & c \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow \lambda_1 = 0, \lambda_{2,3} = 1$$

$$E_0 = \text{Ker } A \Rightarrow \dim E_0 = \dim \text{Ker } A = 1 \text{ with } \bar{v}_1 = \begin{pmatrix} -a \\ 1 \\ 0 \end{pmatrix},$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & a & b \\ 0 & -1 & c \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \dim E_1 = 2 \text{ iff } b = -ac \text{ with}$$

$$\bar{v}_2 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \bar{v}_3 = \begin{pmatrix} 0 \\ c \\ 1 \end{pmatrix}$$

$$\dim E_0 + \dim E_1 = 3 = n \Rightarrow A \text{ is diagonalizable}$$

$$S = \begin{pmatrix} -a & 1 & 0 \\ 1 & 0 & c \\ 0 & 0 & 1 \end{pmatrix} \Rightarrow D = S^{-1}AS = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Matrix Diagonalization: Example 2

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow$$

Matrix Diagonalization: Example 2

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \lambda_1 = 1, \lambda_{2,3} = 0$$

Matrix Diagonalization: Example 2

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \lambda_1 = 1, \lambda_{2,3} = 0$$

$E_0 = \text{Ker } A \Rightarrow \dim E_0 = \dim \text{Ker } A = 2$ with

$$\bar{v}_1 = \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix},$$

Matrix Diagonalization: Example 2

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \lambda_1 = 1, \lambda_{2,3} = 0$$

$E_0 = \text{Ker } A \Rightarrow \dim E_0 = \dim \text{Ker } A = 2$ with

$$\bar{v}_1 = \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}, \bar{v}_2 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

Matrix Diagonalization: Example 2

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \lambda_1 = 1, \lambda_{2,3} = 0$$

$$E_0 = \text{Ker } A \Rightarrow \dim E_0 = \dim \text{Ker } A = 2 \text{ with}$$

$$\bar{v}_1 = \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}, \bar{v}_2 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & 1 & 1 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \Rightarrow$$

Matrix Diagonalization: Example 2

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \lambda_1 = 1, \lambda_{2,3} = 0$$

$$E_0 = \text{Ker } A \Rightarrow \dim E_0 = \dim \text{Ker } A = 2 \text{ with}$$

$$\bar{v}_1 = \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}, \bar{v}_2 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & 1 & 1 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \Rightarrow E_1 = \text{span} \underbrace{\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}}_{\bar{v}_3} \Rightarrow$$

Matrix Diagonalization: Example 2

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \lambda_1 = 1, \lambda_{2,3} = 0$$

$$E_0 = \text{Ker } A \Rightarrow \dim E_0 = \dim \text{Ker } A = 2 \text{ with}$$

$$\bar{v}_1 = \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}, \bar{v}_2 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & 1 & 1 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \Rightarrow E_1 = \text{span} \underbrace{\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}}_{\bar{v}_3} \Rightarrow \dim E_1 = 1$$

$$\Rightarrow \dim E_0 + \dim E_1 = 3 = n \Rightarrow A \text{ is diagonalizable}$$

$$S = \begin{pmatrix} -1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \Rightarrow$$

Matrix Diagonalization: Example 2

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \lambda_1 = 1, \lambda_{2,3} = 0$$

$$E_0 = \text{Ker } A \Rightarrow \dim E_0 = \dim \text{Ker } A = 2 \text{ with}$$

$$\bar{v}_1 = \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}, \bar{v}_2 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

$$E_1 = \text{Ker} \begin{pmatrix} 0 & 1 & 1 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \Rightarrow E_1 = \text{span} \underbrace{\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}}_{\bar{v}_3} \Rightarrow \dim E_1 = 1$$

$$\Rightarrow \dim E_0 + \dim E_1 = 3 = n \Rightarrow A \text{ is diagonalizable}$$

$$S = \begin{pmatrix} -1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \Rightarrow D = S^{-1}AS = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Remark

$A_{n \times n}$ is diagonalizable $\Rightarrow \exists$ invertible $S_{n \times n}$: $D = S^{-1}AS$

$$A = SDS^{-1}$$

\Downarrow

$$A^t = (SDS^{-1})^t = SDS^{-1}SDS^{-1} \dots SDS^{-1} = SD^T S^{-1}$$

\Downarrow

$$A^t = SD^t S^{-1}$$

Example

Let

$$A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = 5, \lambda_2 = -1 \Rightarrow A \text{ is diagonalizable}$$

$$E_5 = \text{Ker} \begin{pmatrix} -4 & 2 \\ 4 & -2 \end{pmatrix} \Rightarrow \bar{v}_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \dim E_5 = 1$$

$$E_{-1} = \text{Ker} \begin{pmatrix} 2 & 2 \\ 4 & 4 \end{pmatrix} \Rightarrow \bar{v}_1 = \begin{pmatrix} -1 \\ 1 \end{pmatrix}, \dim E_{-1} = 1$$

Example

Let

$$A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = 5, \lambda_2 = -1 \Rightarrow A \text{ is diagonalizable}$$

$$E_5 = \text{Ker} \begin{pmatrix} -4 & 2 \\ 4 & -2 \end{pmatrix} \Rightarrow \bar{v}_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \dim E_5 = 1$$

$$E_{-1} = \text{Ker} \begin{pmatrix} 2 & 2 \\ 4 & 4 \end{pmatrix} \Rightarrow \bar{v}_1 = \begin{pmatrix} -1 \\ 1 \end{pmatrix}, \dim E_{-1} = 1$$

$$A = \underbrace{\begin{pmatrix} 1 & -1 \\ 2 & 1 \end{pmatrix}}_S \underbrace{\begin{pmatrix} 5 & 0 \\ 0 & -1 \end{pmatrix}}_D \underbrace{\frac{1}{3} \begin{pmatrix} 1 & 1 \\ -2 & 1 \end{pmatrix}}_{S^{-1}}$$

Example

Let

$$A = \begin{pmatrix} 1 & 2 \\ 4 & 3 \end{pmatrix} \Rightarrow \lambda_1 = 5, \lambda_2 = -1 \Rightarrow A \text{ is diagonalizable}$$

$$E_5 = \text{Ker} \begin{pmatrix} -4 & 2 \\ 4 & -2 \end{pmatrix} \Rightarrow \bar{v}_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \dim E_5 = 1$$

$$E_{-1} = \text{Ker} \begin{pmatrix} 2 & 2 \\ 4 & 4 \end{pmatrix} \Rightarrow \bar{v}_1 = \begin{pmatrix} -1 \\ 1 \end{pmatrix}, \dim E_{-1} = 1$$

$$A = \underbrace{\begin{pmatrix} 1 & -1 \\ 2 & 1 \end{pmatrix}}_S \underbrace{\begin{pmatrix} 5 & 0 \\ 0 & -1 \end{pmatrix}}_D \underbrace{\frac{1}{3} \begin{pmatrix} 1 & 1 \\ -2 & 1 \end{pmatrix}}_{S^{-1}}$$

$$\begin{aligned} A^t &= \begin{pmatrix} 1 & -1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 5^t & 0 \\ 0 & (-1)^t \end{pmatrix} \frac{1}{3} \begin{pmatrix} 1 & 1 \\ -2 & 1 \end{pmatrix} \\ &= \frac{1}{3} \begin{pmatrix} 5^t + 2(-1)^t & 5^t + (-1)^{t+1} \\ 2 \cdot 5^t - 2(-1)^t & 2 \cdot 5^t + (-1)^t \end{pmatrix} \end{aligned}$$

Diagonalizable Linear Operators

Definition: A scalar λ is called an **eigenvalue** of a linear operator $T: V \rightarrow V$ if

$$\exists f \in V, f \neq 0: Tf = \lambda f.$$

The element $f \in V$ is called an **eigenfunction**.

If $\dim V < +\infty$, then a basis \mathcal{D} consisting of eigenfunctions of T is called an **eigenbasis** for T .

A linear operator T is diagonalizable if the matrix of T w.r.t. some basis is diagonal \Leftrightarrow there exists an eigenbasis for T .

Diagonalizable Linear Operators: Example 1

$$T: P_2(\mathbb{R}) \rightarrow P_2(\mathbb{R}), \quad Tp(t) = p(2t - 1), \quad \mathfrak{B} = \{1, t, t^2\}$$

Diagonalizable Linear Operators: Example 1

$$T: P_2(\mathbb{R}) \rightarrow P_2(\mathbb{R}), \quad Tp(t) = p(2t - 1), \quad \mathfrak{B} = \{1, t, t^2\}$$

$$a + bt + ct^2 \xrightarrow{T} a - b + c + (2b - 4c)t + 4ct^2$$

Diagonalizable Linear Operators: Example 1

$$T: P_2(\mathbb{R}) \rightarrow P_2(\mathbb{R}), \quad Tp(t) = p(2t - 1), \quad \mathfrak{B} = \{1, t, t^2\}$$

$$\begin{array}{ccc} a + bt + ct^2 & \xrightarrow{T} & a - b + c + (2b - 4c)t + 4ct^2 \\ \downarrow L_{\mathfrak{B}} & & \downarrow L_{\mathfrak{B}} \\ \begin{pmatrix} a \\ b \\ c \end{pmatrix} & \xrightarrow{A} & \begin{pmatrix} a - b + c \\ 2b - 4c \\ 4c \end{pmatrix} \end{array}$$

$$A = \begin{pmatrix} 1 & -1 & 1 \\ 0 & 2 & -4 \\ 0 & 0 & 4 \end{pmatrix} \Rightarrow$$

Diagonalizable Linear Operators: Example 1

$$T: P_2(\mathbb{R}) \rightarrow P_2(\mathbb{R}), \quad Tp(t) = p(2t - 1), \quad \mathfrak{B} = \{1, t, t^2\}$$

$$\begin{array}{ccc} a + bt + ct^2 & \xrightarrow{T} & a - b + c + (2b - 4c)t + 4ct^2 \\ \downarrow L_{\mathfrak{B}} & & \downarrow L_{\mathfrak{B}} \\ \begin{pmatrix} a \\ b \\ c \end{pmatrix} & \xrightarrow{A} & \begin{pmatrix} a - b + c \\ 2b - 4c \\ 4c \end{pmatrix} \end{array}$$

$$A = \begin{pmatrix} 1 & -1 & 1 \\ 0 & 2 & -4 \\ 0 & 0 & 4 \end{pmatrix} \Rightarrow \lambda_1 = 1, \lambda_2 = 2, \lambda_3 = 4$$

Diagonalizable Linear Operators: Example 1

$$T: P_2(\mathbb{R}) \rightarrow P_2(\mathbb{R}), \quad Tp(t) = p(2t - 1), \quad \mathfrak{B} = \{1, t, t^2\}$$

$$\begin{array}{ccc} a + bt + ct^2 & \xrightarrow{T} & a - b + c + (2b - 4c)t + 4ct^2 \\ \downarrow L_{\mathfrak{B}} & & \downarrow L_{\mathfrak{B}} \\ \begin{pmatrix} a \\ b \\ c \end{pmatrix} & \xrightarrow{A} & \begin{pmatrix} a - b + c \\ 2b - 4c \\ 4c \end{pmatrix} \end{array}$$

$$A = \begin{pmatrix} 1 & -1 & 1 \\ 0 & 2 & -4 \\ 0 & 0 & 4 \end{pmatrix} \Rightarrow \lambda_1 = 1, \lambda_2 = 2, \lambda_3 = 4$$

$\Rightarrow A$ is diagonalizable

Diagonalizable Linear Operators: Example 1

$$T: P_2(\mathbb{R}) \rightarrow P_2(\mathbb{R}), \quad Tp(t) = p(2t - 1), \quad \mathfrak{B} = \{1, t, t^2\}$$

$$\begin{array}{ccc} a + bt + ct^2 & \xrightarrow{T} & a - b + c + (2b - 4c)t + 4ct^2 \\ \downarrow L_{\mathfrak{B}} & & \downarrow L_{\mathfrak{B}} \\ \begin{pmatrix} a \\ b \\ c \end{pmatrix} & \xrightarrow{A} & \begin{pmatrix} a - b + c \\ 2b - 4c \\ 4c \end{pmatrix} \end{array}$$

$$A = \begin{pmatrix} 1 & -1 & 1 \\ 0 & 2 & -4 \\ 0 & 0 & 4 \end{pmatrix} \Rightarrow \lambda_1 = 1, \lambda_2 = 2, \lambda_3 = 4$$

$\Rightarrow A$ is diagonalizable

$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix} \text{ is the eigebasis for } T$$

Diagonalizable Linear Operators: Example 1

$$S_{\mathfrak{D} \rightarrow \mathfrak{B}} = \begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{pmatrix}$$

Diagonalizable Linear Operators: Example 1

$$S_{\mathfrak{D} \rightarrow \mathfrak{B}} = \begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\Rightarrow p_1 = 1, p_2 = -1 + t, p_3 = 1 - 2t + t^2$$

Diagonalizable Linear Operators: Example 1

$$S_{\mathfrak{D} \rightarrow \mathfrak{B}} = \begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\Rightarrow p_1 = 1, p_2 = -1 + t, p_3 = 1 - 2t + t^2$$

$$T(1) = \underbrace{1}_{\lambda_1}, T(t-1) = \underbrace{2}_{\lambda_2}(t-1), T(t-1)^2 = \underbrace{4}_{\lambda_3}(t-1)^2$$

Diagonalizable Linear Operators: Example 1

$$S_{\mathfrak{D} \rightarrow \mathfrak{B}} = \begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\Rightarrow p_1 = 1, p_2 = -1 + t, p_3 = 1 - 2t + t^2$$

$$T(1) = \underbrace{1}_{\lambda_1}, T(t-1) = \underbrace{2}_{\lambda_2}(t-1), T(t-1)^2 = \underbrace{4}_{\lambda_3}(t-1)^2$$

$$D = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 4 \end{pmatrix}$$

Diagonalizable Linear Operators: Example 2

Consider a linear operator $L: \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ defined by

$$L(A) = A - A^T.$$

Diagonalizable Linear Operators: Example 2

Consider a linear operator $L: \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ defined by

$$L(A) = A - A^T.$$

$$L(A) = \lambda A$$

Diagonalizable Linear Operators: Example 2

Consider a linear operator $L: \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ defined by

$$L(A) = A - A^T.$$

$$L(A) = \lambda A$$

A. A is symmetric \Rightarrow

Diagonalizable Linear Operators: Example 2

Consider a linear operator $L: \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ defined by

$$L(A) = A - A^T.$$

$$L(A) = \lambda A$$

A. A is symmetric $\Rightarrow A^T = A \Rightarrow$

Diagonalizable Linear Operators: Example 2

Consider a linear operator $L: \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ defined by

$$L(A) = A - A^T.$$

$$L(A) = \lambda A$$

A. A is symmetric $\Rightarrow A^T = A \Rightarrow L(A) = 0 = 0 \cdot A \Rightarrow \lambda = 0$

Diagonalizable Linear Operators: Example 2

Consider a linear operator $L: \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ defined by

$$L(A) = A - A^T.$$

$$L(A) = \lambda A$$

A. A is symmetric $\Rightarrow A^T = A \Rightarrow L(A) = 0 = 0 \cdot A \Rightarrow \lambda = 0$

$$A = \begin{pmatrix} a & b \\ b & c \end{pmatrix} = a \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + b \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + c \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

Diagonalizable Linear Operators: Example 2

Consider a linear operator $L: \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ defined by

$$L(A) = A - A^T.$$

$$L(A) = \lambda A$$

A. A is symmetric $\Rightarrow A^T = A \Rightarrow L(A) = 0 = 0 \cdot A \Rightarrow \lambda = 0$

$$A = \begin{pmatrix} a & b \\ b & c \end{pmatrix} = a \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + b \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + c \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\dim E_0 = 3$$

Diagonalizable Linear Operators: Example 2

Consider a linear operator $L: \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ defined by

$$L(A) = A - A^T.$$

$$L(A) = \lambda A$$

A. A is symmetric $\Rightarrow A^T = A \Rightarrow L(A) = 0 = 0 \cdot A \Rightarrow \lambda = 0$

$$A = \begin{pmatrix} a & b \\ b & c \end{pmatrix} = a \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + b \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + c \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\dim E_0 = 3$$

B. A is skew-symmetric \Rightarrow

Diagonalizable Linear Operators: Example 2

Consider a linear operator $L: \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ defined by

$$L(A) = A - A^T.$$

$$L(A) = \lambda A$$

A. A is symmetric $\Rightarrow A^T = A \Rightarrow L(A) = 0 = 0 \cdot A \Rightarrow \lambda = 0$

$$A = \begin{pmatrix} a & b \\ b & c \end{pmatrix} = a \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + b \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + c \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\dim E_0 = 3$$

B. A is skew-symmetric $\Rightarrow A^T = -A \Rightarrow$

Diagonalizable Linear Operators: Example 2

Consider a linear operator $L: \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ defined by

$$L(A) = A - A^T.$$

$$L(A) = \lambda A$$

A. A is symmetric $\Rightarrow A^T = A \Rightarrow L(A) = 0 = 0 \cdot A \Rightarrow \lambda = 0$

$$A = \begin{pmatrix} a & b \\ b & c \end{pmatrix} = a \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + b \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + c \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\dim E_0 = 3$$

B. A is skew-symmetric $\Rightarrow A^T = -A \Rightarrow L(A) = 2A \Rightarrow \lambda = 2$

Diagonalizable Linear Operators: Example 2

Consider a linear operator $L: \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ defined by

$$L(A) = A - A^T.$$

$$L(A) = \lambda A$$

A. A is symmetric $\Rightarrow A^T = A \Rightarrow L(A) = 0 = 0 \cdot A \Rightarrow \lambda = 0$

$$A = \begin{pmatrix} a & b \\ b & c \end{pmatrix} = a \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + b \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + c \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\dim E_0 = 3$$

B. A is skew-symmetric $\Rightarrow A^T = -A \Rightarrow L(A) = 2A \Rightarrow \lambda = 2$

$$A = \begin{pmatrix} 0 & a \\ -a & 0 \end{pmatrix} = a \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \Rightarrow$$

Diagonalizable Linear Operators: Example 2

Consider a linear operator $L: \mathbb{M}_{2 \times 2} \rightarrow \mathbb{M}_{2 \times 2}$ defined by

$$L(A) = A - A^T.$$

$$L(A) = \lambda A$$

A. A is symmetric $\Rightarrow A^T = A \Rightarrow L(A) = 0 = 0 \cdot A \Rightarrow \lambda = 0$

$$A = \begin{pmatrix} a & b \\ b & c \end{pmatrix} = a \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + b \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + c \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$


$$\dim E_0 = 3$$

B. A is skew-symmetric $\Rightarrow A^T = -A \Rightarrow L(A) = 2A \Rightarrow \lambda = 2$

$$A = \begin{pmatrix} 0 & a \\ -a & 0 \end{pmatrix} = a \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \Rightarrow \dim E_2 = 1$$

Ranking Problem


- * Consider the results of a tournament



where " $A \rightarrow B$ " means A defeated B .

Ranking Problem

- * Consider the results of a tournament




where " $A \rightarrow B$ " means A defeated B .

How to rank the players fairly?

Ranking Problem

- * Consider the results of a tournament




where " $A \rightarrow B$ " means A defeated B .

How to rank the players fairly?

- * The Page-Brin idea:

Ranking Problem

- * Consider the results of a tournament




where " $A \rightarrow B$ " means A defeated B .

How to rank the players fairly?

- * The Page-Brin idea: it should *be worth more* to defeat a *better player*

Ranking Problem

- * Consider the results of a tournament




where " $A \rightarrow B$ " means A defeated B .

How to rank the players fairly?

- * The Page-Brin idea: it should *be worth more* to defeat a *better player* \Leftarrow based on Perron (1907)-Frobenius (1912) Theorem

Ranking Problem

- * Consider the results of a tournament



where " $A \rightarrow B$ " means A defeated B .


How to rank the players fairly?

- * The Page-Brin idea: it should *be worth more* to defeat a *better player* \Leftarrow based on Perron (1907)-Frobenius (1912) Theorem

How do we know who is better before ranking them?

Ranking Problem

- * Consider the results of a tournament



where " $A \rightarrow B$ " means A defeated B .

How to rank the players fairly?

- * The Page-Brin idea: it should *be worth more* to defeat a *better player* \Leftarrow based on Perron (1907)-Frobenius (1912) Theorem

How do we know who is better before ranking them?

- * Define recursion!

Ranking Problem

- * Give everyone the initial score of 1

$$\bar{x}_0 = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

Ranking Problem

- * Give everyone the initial score of 1

$$\bar{x}_0 = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

- * Define for all $n \geq 0$

$$\bar{x}_{n+1} = A\bar{x}_n,$$

where

$$A = \begin{matrix} & \begin{matrix} A & B & C & D \end{matrix} \\ \begin{matrix} A \\ B \\ C \\ D \end{matrix} & \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix} \end{matrix}$$

Ranking Problem

$$\bar{x}_1 = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 1 \\ 1 \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

Ranking Problem

$$\bar{x}_1 = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 1 \\ 1 \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

$$\bar{x}_2 = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ 2 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 1 \\ 2 \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

Ranking Problem

$$\bar{x}_1 = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 1 \\ 1 \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

$$\bar{x}_2 = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ 2 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 1 \\ 2 \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

$$\bar{x}_3 = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 3 \\ 2 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 2 \\ 3 \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

Ranking Problem

$$\bar{x}_1 = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 1 \\ 1 \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

$$\bar{x}_2 = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ 2 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 1 \\ 2 \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

$$\bar{x}_3 = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 3 \\ 2 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 2 \\ 3 \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

The $(n+1)$ th score of a player A is the sum of the n th scores of the players that the player A defeated.

Ranking Problem

$$\bar{x}_5 = \begin{pmatrix} 8 \\ 6 \\ 3 \\ 5 \end{pmatrix}, \bar{x}_{10} = \begin{pmatrix} 35 \\ 34 \\ 21 \\ 26 \end{pmatrix}, \bar{x}_{100} = \begin{pmatrix} 1037 \\ 933 \\ 547 \\ 731 \end{pmatrix}$$

Ranking Problem

$$\bar{x}_5 = \begin{pmatrix} 8 \\ 6 \\ 3 \\ 5 \end{pmatrix}, \bar{x}_{10} = \begin{pmatrix} 35 \\ 34 \\ 21 \\ 26 \end{pmatrix}, \bar{x}_{100} = \begin{pmatrix} 1037 \\ 933 \\ 547 \\ 731 \end{pmatrix}$$

Is $A > B > D > C? \Rightarrow$

Ranking Problem

$$\bar{x}_5 = \begin{pmatrix} 8 \\ 6 \\ 3 \\ 5 \end{pmatrix}, \bar{x}_{10} = \begin{pmatrix} 35 \\ 34 \\ 21 \\ 26 \end{pmatrix}, \bar{x}_{100} = \begin{pmatrix} 1037 \\ 933 \\ 547 \\ 731 \end{pmatrix}$$

Is $A > B > D > C$? \Rightarrow analyze $\bar{x}_n = A^n \bar{x}_0$ as $n \rightarrow \infty$

Ranking Problem

$$\bar{x}_5 = \begin{pmatrix} 8 \\ 6 \\ 3 \\ 5 \end{pmatrix}, \bar{x}_{10} = \begin{pmatrix} 35 \\ 34 \\ 21 \\ 26 \end{pmatrix}, \bar{x}_{100} = \begin{pmatrix} 1037 \\ 933 \\ 547 \\ 731 \end{pmatrix}$$

Is $A > B > D > C$? \Rightarrow analyze $\bar{x}_n = A^n \bar{x}_0$ as $n \rightarrow \infty$

Theorem (Perron-Frobenius): There exists a *largest positive* eigenvalue λ_{PF} for a nonnegative matrix A such that the rescaled system

$$\bar{x}_n = \left(\frac{1}{\lambda_{PF}} A \right)^n \bar{x}_0$$

converges to an equilibrium state \bar{x}_∞ .

$$\bar{x}_\infty = \bar{x}_{\infty+1} = \frac{1}{\lambda_{PF}} A \bar{x}_\infty \Rightarrow A \bar{x}_\infty = \lambda_{PF} \bar{x}_\infty$$

The equilibrium state is the eigenvector associated with λ_{PF} !!!

Ranking Problem

The largest positive eigenvalue is

$$\lambda_{PF} = 1.3953369\dots$$

and

$$\bar{x}_{\infty} = \begin{pmatrix} 0.321\dots \\ 0.288\dots \\ 0.165\dots \\ 0.230\dots \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

Ranking Problem

The largest positive eigenvalue is

$$\lambda_{PF} = 1.3953369\dots$$

and

$$\bar{x}_{\infty} = \begin{pmatrix} 0.321\dots \\ 0.288\dots \\ 0.165\dots \\ 0.230\dots \end{pmatrix} \begin{matrix} A \\ B \\ C \\ D \end{matrix}$$

$$A > B > D > C$$