

# vv214: Introduction: systems of linear equations, graphs, even/oddtown, Markov chains.

Dr.Olga Danilkina

UM-SJTU Joint Institute



May 16, 2019

# This week

## Today

1. Even/oddtown problems.
2. Card shuffling/Markov processes.
3. Graphs.
4. Systems of linear equations: motivation with two historical problems, solutions of a SLE, geometrical interpretations of linear systems.
5. Gauss-Jordan elimination.
6. Simplest input-output Leontief models.

# This week

## Today

1. Even/oddtown problems.
2. Card shuffling/Markov processes.
3. Graphs.
4. Systems of linear equations: motivation with two historical problems, solutions of a SLE, geometrical interpretations of linear systems.
5. Gauss-Jordan elimination.
6. Simplest input-output Leontief models.

## Next class

1. Matrices: coefficient, augmented, square, upper and lower triangular, identity.
2. Elementary transformations of a matrix.
3. Reduced row-echelon form of a matrix.
4. Rank of a matrix.

# Eventown Problem

- Consider a town with  $n$  residents.

# Eventown Problem

- Consider a town with  $n$  residents. They like forming clubs!

# Eventown Problem

- Consider a town with  $n$  residents. They like forming clubs!

$$P = \{p_1, \dots, p_n\}$$

# Eventown Problem

- Consider a town with  $n$  residents. They like forming clubs!

$$P = \{p_1, \dots, p_n\}$$

- Q: What is the potential number  $m$  of clubs  $C_1, \dots, C_m$ ?

# Eventown Problem

- Consider a town with  $n$  residents. They like forming clubs!

$$P = \{p_1, \dots, p_n\}$$

- Q: What is the potential number  $m$  of clubs  $C_1, \dots, C_m$ ?  
*Allow the empty club and assume that the same residents form only one club*



# Eventown Problem

- Consider a town with  $n$  residents. They like forming clubs!

$$P = \{p_1, \dots, p_n\}$$

- Q: What is the potential number  $m$  of clubs  $C_1, \dots, C_m$ ?  
*Allow the empty club and assume that the same residents form only one club* A:  $m = 2^n$

# Eventown Problem

- Consider a town with  $n$  residents. They like forming clubs!

$$P = \{p_1, \dots, p_n\}$$

- Q: What is the potential number  $m$  of clubs  $C_1, \dots, C_m$ ?  
*Allow the empty club and assume that the same residents form only one club* A:  $m = 2^n$
- *But the same residents can form as many clubs as they want!*  $\Rightarrow$

# Eventown Problem

- Consider a town with  $n$  residents. They like forming clubs!

$$P = \{p_1, \dots, p_n\}$$

- Q: What is the potential number  $m$  of clubs  $C_1, \dots, C_m$ ?  
*Allow the empty club and assume that the same residents form only one club* A:  $m = 2^n$
- *But the same residents can form as many clubs as they want!*  $\Rightarrow$  infinitely many clubs

# Eventown Problem

- Consider a town with  $n$  residents. They like forming clubs!

$$P = \{p_1, \dots, p_n\}$$

- Q: What is the potential number  $m$  of clubs  $C_1, \dots, C_m$ ?  
*Allow the empty club and assume that the same residents form only one club* A:  $m = 2^n$
- But the same residents can form as many clubs as they want!*  $\Rightarrow$  infinitely many clubs
- Club rule 1:** No two clubs have exactly the same membership

$$\forall i \neq j \quad C_i \neq C_j \quad (\Rightarrow m \leq 2^n)$$

- Club rule 2:** Every club must have an even number of people

$$\forall i \quad |C_i| = \text{even}$$

# Eventown Problem

- **Club rule 3:** Every two clubs must have an even number of common members

$$\forall i \neq j \quad |C_i \cap C_j| = \text{even}$$

# Eventown Problem

- **Club rule 3:** Every two clubs must have an even number of common members

$$\forall i \neq j \quad |C_i \cap C_j| = \text{even}$$

- **Married-couple solutions:**

# Eventown Problem

- **Club rule 3:** Every two clubs must have an even number of common members

$$\forall i \neq j \quad |C_i \cap C_j| = \text{even}$$

- **Married-couple solutions:** Pair up the residents and consider the groups of two as units: a resident can join a club if and only if the partner joins as well  $\Rightarrow$

# Eventown Problem

- **Club rule 3:** Every two clubs must have an even number of common members

$$\forall i \neq j \quad |C_i \cap C_j| = \text{even}$$

- **Married-couple solutions:** Pair up the residents and consider the groups of two as units: a resident can join a club if and only if the partner joins as well  $\Rightarrow m = 2^{\lfloor \frac{n}{2} \rfloor}$



# Eventown Problem

- **Club rule 3:** Every two clubs must have an even number of common members

$$\forall i \neq j \quad |C_i \cap C_j| = \text{even}$$

- **Married-couple solutions:** Pair up the residents and consider the groups of two as units: a resident can join a club if and only if the partner joins as well  $\Rightarrow m = 2^{\lfloor \frac{n}{2} \rfloor}$

**Theorem (Eventown-Berlekamp 1969):** If there are  $n$  residents and  $m$  clubs in Eventown then

$$m \leq 2^{\lfloor \frac{n}{2} \rfloor}$$

# Eventown Problem

- **Club rule 3:** Every two clubs must have an even number of common members

$$\forall i \neq j \quad |C_i \cap C_j| = \text{even}$$

- **Married-couple solutions:** Pair up the residents and consider the groups of two as units: a resident can join a club if and only if the partner joins as well  $\Rightarrow m = 2^{\lfloor \frac{n}{2} \rfloor}$

**Theorem (Eventown-Berlekamp 1969):** If there are  $n$  residents and  $m$  clubs in Eventown then

$$m \leq 2^{\lfloor \frac{n}{2} \rfloor}$$

**Example:** Let  $n = 16$ . Then 16 residents can form  $255 = 2^8 - 1$  clubs in an eventown.

# Oddtown problem

- **Club rule 1:** No two clubs have exactly the same membership

$$\forall i \neq j \quad C_i \neq C_j \quad (\Rightarrow m \leq 2^n)$$

- **Club rule 2:** Every club must have an odd number of people

$$\forall i \quad |C_i| = \text{odd}$$

- **Club rule 3:** Every two clubs must have an even number of common members

$$\forall i \neq j \quad |C_i \cap C_j| = \text{even}$$

# Oddtown problem

- **Club rule 1:** No two clubs have exactly the same membership

$$\forall i \neq j \quad C_i \neq C_j \quad (\Rightarrow m \leq 2^n)$$

- **Club rule 2:** Every club must have an odd number of people

$$\forall i \quad |C_i| = \text{odd}$$

- **Club rule 3:** Every two clubs must have an even number of common members

$$\forall i \neq j \quad |C_i \cap C_j| = \text{even}$$

- ▶ **Single-clubs solution** has  $m = n$  clubs.

# Oddtown problem

- **Club rule 1:** No two clubs have exactly the same membership

$$\forall i \neq j \quad C_i \neq C_j \quad (\Rightarrow m \leq 2^n)$$

- **Club rule 2:** Every club must have an odd number of people

$$\forall i \quad |C_i| = \text{odd}$$

- **Club rule 3:** Every two clubs must have an even number of common members

$$\forall i \neq j \quad |C_i \cap C_j| = \text{even}$$

► **Single-clubs solution** has  $m = n$  clubs.

**Theorem(Oddtown-Berlekamp 1969):** If there are  $n$  residents and  $m$  clubs in Oddtown then

$$m \leq n$$

# Markov Chains

Q: How many times do we need to shuffle a deck of cards before it gets reasonably well-shuffled?

# Markov Chains

Q: How many times do we need to shuffle a deck of cards before it gets reasonably well-shuffled?

A: The deck has 52 cards, the state of a deck is how cards are ordered  $\Rightarrow 52!$  states.

# Markov Chains

Q: How many times do we need to shuffle a deck of cards before it gets reasonably well-shuffled?

A: The deck has 52 cards, the state of a deck is how cards are ordered  $\Rightarrow 52!$  states.

A shuffling move is a bijective function from the set of states to itself, we shall call this bijection as a **permutation**.



# Markov Chains

Q: How many times do we need to shuffle a deck of cards before it gets reasonably well-shuffled?

A: The deck has 52 cards, the state of a deck is how cards are ordered  $\Rightarrow 52!$  states.

A shuffling move is a bijective function from the set of states to itself, we shall call this bijection as a [permutation](#).

There are  $52!$  possible transitions. Some states are more likely than the others, and some may have essentially zero probability.

# Markov Chains

Q: How many times do we need to shuffle a deck of cards before it gets reasonably well-shuffled?

A: The deck has 52 cards, the state of a deck is how cards are ordered  $\Rightarrow 52!$  states.

A shuffling move is a bijective function from the set of states to itself, we shall call this bijection as a [permutation](#).

There are  $52!$  possible transitions. Some states are more likely than the others, and some may have essentially zero probability.

$\Rightarrow$  We are interested in probability distribution over all possible transitions.

# Markov Chains

Q: How many times do we need to shuffle a deck of cards before it gets reasonably well-shuffled?

A: The deck has 52 cards, the state of a deck is how cards are ordered  $\Rightarrow 52!$  states.

A shuffling move is a bijective function from the set of states to itself, we shall call this bijection as a [permutation](#).

There are  $52!$  possible transitions. Some states are more likely than the others, and some may have essentially zero probability.

$\Rightarrow$  We are interested in probability distribution over all possible transitions. Ideally, we want all states have the same probability  $1/52! \Rightarrow$

# Markov Chains

Q: How many times do we need to shuffle a deck of cards before it gets reasonably well-shuffled?

A: The deck has 52 cards, the state of a deck is how cards are ordered  $\Rightarrow 52!$  states.

A shuffling move is a bijective function from the set of states to itself, we shall call this bijection as a **permutation**.

There are  $52!$  possible transitions. Some states are more likely than the others, and some may have essentially zero probability.

$\Rightarrow$  We are interested in probability distribution over all possible transitions. Ideally, we want all states have the same probability  $1/52! \Rightarrow$  **uniform** distribution.

# Markov Chains

Denote  $p_{ij}$  the **transition probability** of going from state  $i$  to state  $j$  ( $i, j = 1, 2, \dots, 52!$ )

**Definition:** The **transition matrix**  $T$  is a  $52! \times 52!$  matrix  $T = (p_{ij})$ .

# Markov Chains

Denote  $p_{ij}$  the **transition probability** of going from state  $i$  to state  $j$  ( $i, j = 1, 2, \dots, 52!$ )

**Definition:** The **transition matrix**  $T$  is a  $52! \times 52!$  matrix  $T = (p_{ij})$ .

**Definition:** If the transition matrix does not depend on time, then we have a **Markov Chain**.

# Markov Chains

Denote  $p_{ij}$  the **transition probability** of going from state  $i$  to state  $j$  ( $i, j = 1, 2, \dots, 52!$ )

**Definition:** The **transition matrix**  $T$  is a  $52! \times 52!$  matrix  $T = (p_{ij})$ .

**Definition:** If the transition matrix does not depend on time, then we have a **Markov Chain**.

**Q:** What happens if we shuffle the deck multiple times?

# Graphs

**Definition:** A **graph**  $G$  is a collection of nodes  $V$  called **vertices** and links between nodes  $E \subset V \times V$  called **edges**.



# Graphs

**Definition:** A **graph**  $G$  is a collection of nodes  $V$  called **vertices** and links between nodes  $E \subset V \times V$  called **edges**. We shall consider the model with only one edge between any two vertices and no edge connects a vertex to itself.

# Graphs

**Definition:** A **graph**  $G$  is a collection of nodes  $V$  called **vertices** and links between nodes  $E \subset V \times V$  called **edges**.

We shall consider the model with only one edge between any two vertices and no edge connects a vertex to itself.

**Definition:** Two vertices are said to be **adjacent** if there is an edge connecting them.

# Graphs

**Definition:** A **graph**  $G$  is a collection of nodes  $V$  called **vertices** and links between nodes  $E \subset V \times V$  called **edges**.

We shall consider the model with only one edge between any two vertices and no edge connects a vertex to itself.

**Definition:** Two vertices are said to be **adjacent** if there is an edge connecting them.

**Definition:** A **degree**  $\deg(v)$  of a vertex  $v$  is the number of adjacent vertices OR the number of edges extending from the vertex  $v$ .

# Graphs

**Definition:** A **graph**  $G$  is a collection of nodes  $V$  called **vertices** and links between nodes  $E \subset V \times V$  called **edges**.

We shall consider the model with only one edge between any two vertices and no edge connects a vertex to itself.

**Definition:** Two vertices are said to be **adjacent** if there is an edge connecting them.

**Definition:** A **degree**  $\deg(v)$  of a vertex  $v$  is the number of adjacent vertices OR the number of edges extending from the vertex  $v$ .

The **degree**  $\deg(G)$  of a graph  $G$  is defined by

$$\deg(G) = \max_{v \in V} \deg(v)$$

# Graphs

**Definition:** A **graph**  $G$  is a collection of nodes  $V$  called **vertices** and links between nodes  $E \subset V \times V$  called **edges**.

We shall consider the model with only one edge between any two vertices and no edge connects a vertex to itself.

**Definition:** Two vertices are said to be **adjacent** if there is an edge connecting them.

**Definition:** A **degree**  $\deg(v)$  of a vertex  $v$  is the number of adjacent vertices OR the number of edges extending from the vertex  $v$ .

The **degree**  $\deg(G)$  of a graph  $G$  is defined by

$$\deg(G) = \max_{v \in V} \deg(v)$$

**Definition:** The **adjacency matrix** of the graph  $G$  with vertices  $v_1, \dots, v_n$  is an  $n \times n$  matrix  $A = A_G = (a_{ij})_{n \times n}$  defined by

$$a_{ij} = \begin{cases} 1 & v_i \sim v_j \\ 0 & v_i \not\sim v_j \end{cases}$$

# Graphs

**Theorem(Handshake theorem):** The number of people who have made an odd number of hand shakes must be even.

$$\sum_{v_i \in V} \deg(v_i) = 2|E|$$

# Graphs

**Theorem(Handshake theorem):** The number of people who have made an odd number of hand shakes must be even.

$$\sum_{v_i \in V} \deg(v_i) = 2|E|$$

**Definition:** A coloring  $c: V \rightarrow \{\text{colors}\}$  is **legal** if

$$v_i \sim v_j \Rightarrow c(v_i) \neq c(v_j)$$

# Graphs

**Theorem(Handshake theorem):** The number of people who have made an odd number of hand shakes must be even.

$$\sum_{v_i \in V} \deg(v_i) = 2|E|$$

**Definition:** A coloring  $c: V \rightarrow \{\text{colors}\}$  is **legal** if

$$v_i \sim v_j \Rightarrow c(v_i) \neq c(v_j)$$

A graph  $G$  is **properly colored** with  $k$  colors if  $k$  colors suffice for a legal coloring.



# Graphs

**Theorem(Handshake theorem):** The number of people who have made an odd number of hand shakes must be even.

$$\sum_{v_i \in V} \deg(v_i) = 2|E|$$

**Definition:** A coloring  $c: V \rightarrow \{\text{colors}\}$  is **legal** if

$$v_i \sim v_j \Rightarrow c(v_i) \neq c(v_j)$$

A graph  $G$  is **properly colored** with  $k$  colors if  $k$  colors suffice for a legal coloring.

**Theorem (Graph Coloring):** If  $\deg(G) \leq d$ , then  $G$  can be colored with  $d + 1$  colors.

# Historical Problems

**Isaac Newton, Arithmetica Universalis, 1707**

# Historical Problems

**Isaac Newton, Arithmetica Universalis, 1707**

3 cows graze 1 field bare in 2 days,

# Historical Problems

## Isaac Newton, *Arithmetica Universalis*, 1707

3 cows graze 1 field bare in 2 days,

7 cows graze 4 fields bare in 4 days, and

# Historical Problems

## Isaac Newton, *Arithmetica Universalis*, 1707

3 cows graze 1 field bare in 2 days,

7 cows graze 4 fields bare in 4 days, and

3 cows graze 2 fields bare in 5 days.

# Historical Problems

## Isaac Newton, *Arithmetica Universalis*, 1707

3 cows graze 1 field bare in 2 days,

7 cows graze 4 fields bare in 4 days, and

3 cows graze 2 fields bare in 5 days.

It is assumed that each field initially provides the same amount  $x$  of grass that the daily growth  $y$  of the fields remains constant, and that all the cows eat the same amount  $z$  each day. Quantities  $x$ ,  $y$  and  $z$  are measured by weight.

Find all the solutions  $x$ ,  $y$ ,  $z$  of this problem.

# Historical Problems

## Isaac Newton, *Arithmetica Universalis*, 1707

3 cows graze 1 field bare in 2 days,

7 cows graze 4 fields bare in 4 days, and

3 cows graze 2 fields bare in 5 days.

It is assumed that each field initially provides the same amount  $x$  of grass that the daily growth  $y$  of the fields remains constant, and that all the cows eat the same amount  $z$  each day. Quantities  $x$ ,  $y$  and  $z$  are measured by weight.

Find all the solutions  $x$ ,  $y$ ,  $z$  of this problem.

**Hint:** Initial amount of grass + grass growth = grass consumed by cows

# Historical Problems

## Isaac Newton, *Arithmetica Universalis*, 1707

3 cows graze 1 field bare in 2 days,

7 cows graze 4 fields bare in 4 days, and

3 cows graze 2 fields bare in 5 days.

It is assumed that each field initially provides the same amount  $x$  of grass that the daily growth  $y$  of the fields remains constant, and that all the cows eat the same amount  $z$  each day. Quantities  $x$ ,  $y$  and  $z$  are measured by weight.

Find all the solutions  $x$ ,  $y$ ,  $z$  of this problem.

**Hint:** Initial amount of grass + grass growth = grass consumed by cows

$$N_{fields}x + N_{fields}N_{days}y = N_{cows}N_{days}z$$



# Gauss-Jordan Elimination

1. Proceed from equation to equation, from top to bottom.
2. For the  $i$ th equation: let  $x_j$  be the leading (pivot) variable. If  $x_j$  does not appear in the  $i$ th equation, swap the  $i$ th equation with the first equation below that does contain  $x_j$ .
3. Multiply the  $i$ th equation by the appropriate scalar so that the coefficient of the leading variable becomes 1.
4. Eliminate  $x_j$  from all the other equations, above and below the  $i$ th by subtracting suitable multiples of the  $i$ th equation from the others.
5. Now proceed to the next equation.
6. If an equation  $0 = \text{nonzero}$  emerges in this process, then the system fails to have solutions; the system is inconsistent.

# Elementary Row Operations

Recall, that the following algebraic operations do not affect the solution(s) of systems of linear equations:

1. Swap equations.
2. Divide/multiply an equation by a nonzero scalar.
3. Add/subtract a multiple of an equation to/from another one.

⇒ we can define the following **elementary row operations**:

1. Interchange two rows.
2. Multiply a row by a nonzero constant,
3. Add a multiple of one row to another one.

# Historical Problems

## **Metrodorus, Greek Anthology, 6th century A.D.**

“Make me a crown weighing 60 minae, mixing gold, bronze, tin, and wrought iron. Let the gold and bronze together form two-thirds, the gold and tin together three-fourths, and the gold and iron three-fifths. Tell me how much gold, tin, bronze, and iron you must put in.”

# Historical Problems

## Metrodorus, Greek Anthology, 6th century A.D.

“Make me a crown weighing 60 minae, mixing gold, bronze, tin, and wrought iron. Let the gold and bronze together form two-thirds, the gold and tin together three-fourths, and the gold and iron three-fifths. Tell me how much gold, tin, bronze, and iron you must put in.”

**Hint:**

$$\begin{cases} x + y & & & = \frac{2}{3}60 \\ x + & z & & = \frac{3}{4}60 \\ x + & & t & = \frac{3}{5}60 \\ x + y + z + t & = 60 \end{cases}$$

# Simplest Leontief input-output model



Consider an economy with several, say 2, industries.

Q: What output should each of the industries in an economy produce to satisfy the total demand for all products?

# Simplest Leontief input-output model

Consider an economy with several, say 2, industries.

Q: What output should each of the industries in an economy produce to satisfy the total demand for all products?



Assume that the consumer demand for their products is, respectively, 1,000 and 780, in millions of dollars per year. What outputs  $a$  and  $b$  (in millions of dollars per year) should the two industries generate to satisfy the demand?

# Reduced Row-Echelon Form. Rank of a Matrix

**Definition:** A matrix is in **reduced row-echelon form *rref*** if it satisfies all of the following conditions:

1. If a row has nonzero entries, then the first nonzero entry is a 1, called the **leading 1** in this row.
2. If a column contains a leading 1, then all the other entries in that column are 0.
3. If a row contains a leading 1, then each row above it contains a leading 1 further to the left  
⇒ rows of 0's, if any, appear at the bottom of the matrix.

# Reduced Row-Echelon Form. Rank of a Matrix

**Definition:** A matrix is in **reduced row-echelon form *rref*** if it satisfies all of the following conditions:

1. If a row has nonzero entries, then the first nonzero entry is a 1, called the **leading 1** in this row.
2. If a column contains a leading 1, then all the other entries in that column are 0.
3. If a row contains a leading 1, then each row above it contains a leading 1 further to the left  
 $\Rightarrow$  rows of 0's, if any, appear at the bottom of the matrix.

**Definition:** The number of leading 1's in the *rref*  $A$  is called the **rank** of the matrix  $A$ .



# Reduced Row-Echelon Form. Rank of a Matrix

$$1. A = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \sim$$

## Reduced Row-Echelon Form. Rank of a Matrix

$$1. A = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & -3 \\ 0 & -6 \end{pmatrix} \sim$$

## Reduced Row-Echelon Form. Rank of a Matrix

$$1. A = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & -3 \\ 0 & -6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$\Rightarrow rref A = \begin{pmatrix} \textcircled{1} & 0 \\ 0 & \textcircled{1} \\ 0 & 0 \end{pmatrix} \Rightarrow$$

## Reduced Row-Echelon Form. Rank of a Matrix

$$1. A = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & -3 \\ 0 & -6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$\Rightarrow \text{rref } A = \begin{pmatrix} \textcircled{1} & 0 \\ 0 & \textcircled{1} \\ 0 & 0 \end{pmatrix} \Rightarrow \text{rank } A = 2$$

$$2. A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$

$$A \sim$$

## Reduced Row-Echelon Form. Rank of a Matrix

$$1. A = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & -3 \\ 0 & -6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$\Rightarrow \text{rref } A = \begin{pmatrix} \textcircled{1} & 0 \\ 0 & \textcircled{1} \\ 0 & 0 \end{pmatrix} \Rightarrow \text{rank } A = 2$$

$$2. A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$

$$A \sim \begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & -6 & -12 \end{pmatrix} \sim$$

## Reduced Row-Echelon Form. Rank of a Matrix

$$1. A = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & -3 \\ 0 & -6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$\Rightarrow \text{rref } A = \begin{pmatrix} \textcircled{1} & 0 \\ 0 & \textcircled{1} \\ 0 & 0 \end{pmatrix} \Rightarrow \text{rank } A = 2$$

$$2. A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$

$$A \sim \begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & -6 & -12 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \sim$$

## Reduced Row-Echelon Form. Rank of a Matrix

$$1. A = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & -3 \\ 0 & -6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$\Rightarrow \text{rref } A = \begin{pmatrix} \textcircled{1} & 0 \\ 0 & \textcircled{1} \\ 0 & 0 \end{pmatrix} \Rightarrow \text{rank } A = 2$$

$$2. A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$

$$A \sim \begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & -6 & -12 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\Rightarrow \text{rref } A = \begin{pmatrix} \textcircled{1} & 0 & -1 \\ 0 & \textcircled{1} & 2 \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow$$

## Reduced Row-Echelon Form. Rank of a Matrix

$$1. A = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & -3 \\ 0 & -6 \end{pmatrix} \sim \begin{pmatrix} 1 & 4 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$\Rightarrow rref A = \begin{pmatrix} \textcircled{1} & 0 \\ 0 & \textcircled{1} \\ 0 & 0 \end{pmatrix} \Rightarrow rank A = 2$$

$$2. A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$

$$A \sim \begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & -6 & -12 \end{pmatrix} \sim \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\Rightarrow rref A = \begin{pmatrix} \textcircled{1} & 0 & -1 \\ 0 & \textcircled{1} & 2 \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow rank A = 2$$



# Questions

1. How many solutions may a system of linear equations have? What is a consistent system?
2. What is a matrix? What is a vector?
3. What are the elementary transformations of matrices?
4. When do we say that a matrix is in reduced row-echelon form?
5. What is the Gauss-Jordan elimination?
6. What is a rank of a matrix?
7. How is the rank of a coefficient matrix related to the number of solutions of a linear system?

# The number of solutions and the rank of the coefficient matrix

Consider a linear system of equations  $n$  with  $m$  variables  $\Rightarrow$  the coefficient matrix  $A$  of the system is  $A_{n \times m}$ .

1.  $\text{rank } A \leq n, \text{rank } A \leq m$
2. If  $\text{rank } A = n$  then the system is consistent.
3. If  $\text{rank } A = m$  then the system has at most one solution.
4. If  $\text{rank } A < m$  then the system either has infinitely many solutions OR inconsistent.

## Remarks:

1. If  $n < m$  then  $\text{rank } A \leq n < m \Rightarrow$  infinitely many OR no solutions
2. If  $n = m$  and
  - a.  $\text{rank } A = n \Rightarrow$  there exists a unique solution
  - b.  $\text{rank } A < n \Rightarrow$  infinitely many OR no solutions

# Matrix Algebra

1. The sum of two matrices  $A_{n \times m}$  and  $B_{n \times m}$  is the matrix  $C_{n \times m}$  s.t.

$$c_{ij} = a_{ij} + b_{ij}, i = \overline{1, n}, j = \overline{1, m}.$$

2. The scalar product  $\alpha A_{n \times m} = (\alpha a_{ij}), i = \overline{1, n}, j = \overline{1, m}$ .

3. The product of a row-matrix  $(a_1 \ a_2 \ a_3)$  and a

column-matrix  $\begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$  is

$$(a_1 \ a_2 \ a_3) \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} = a_1 b_1 + a_2 b_2 + a_3 b_3$$

# Matrix Algebra

4. The product of a matrix  $A_{n \times m}$  and a vector  $\bar{x} \in \mathbb{R}^m$  is

$$A_{n \times m} \bar{x} = \begin{pmatrix} \bar{w}_1 \\ \bar{w}_2 \\ \vdots \\ \bar{w}_n \end{pmatrix} \bar{x} \stackrel{(def)}{=} \begin{pmatrix} (\bar{w}_1, \bar{x}) \\ (\bar{w}_2, \bar{x}) \\ \vdots \\ (\bar{w}_n, \bar{x}) \end{pmatrix}$$

$$\stackrel{(prop)}{=} (\bar{a}_1 \quad \bar{a}_2 \dots \bar{a}_m) \bar{x} = x_1 \bar{a}_1 + x_2 \bar{a}_2 + \dots + x_m \bar{a}_m$$

# Next Week

- ▶ More practice with linear systems/rref/rank.
- ▶ Matrix Algebra.
- ▶ Number fields and linear spaces.
- ▶ Abelian groups and linear spaces. Cyclic groups.
- ▶ Linear combinations and linear dependence/independence.
- ▶ Structure of a linear space: basis, dimension.
- ▶ Lots of examples!