

Second Recitation Class

Linear Algebra

Yuzhou Li

UM-SJTU Joint Institute

May 29, 2019

Table of contents

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace

Direct Sum

Basis

Exercise

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace

Direct Sum

Basis

Exercise

Compute A_G^2 and $\text{trace}(A_G^2)$

1. Let $A_G = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$ be the adjacency matrix of a graph G . Compute $\text{trace}(A)$.

$$\text{trace}(A) = a_{11} + a_{22} + a_{33} = 0$$

2. Compute A_G^2 and $\text{trace}(A_G^2)$. Find the good interpretation for $\text{trace}(A_G^2)$ —the question is still open!

$$(A_G^2)_{ij} = \sum_{k=1}^3 a_{ik} a_{kj} \Rightarrow \text{we count only terms with } a_{ik} a_{kj} \neq 0$$

$$\Rightarrow a_{ik} \neq 0, a_{kj} \neq 0 \iff \underbrace{a_{ik} = a_{kj} = 1}_{v_i \sim v_k \sim v_j} \text{ we count w.r.t } k$$

$$(A_G^2)_{ij} = \text{the number of common neighbors of } v_i \text{ and } v_j$$

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace

Direct Sum

Basis

Exercise

Compute A_G^3

3 Compute A_G^3 . What are the entries $(A_G^3)_{ij}$?

$$(A_G^3)_{ij} = \sum_{k=1}^3 \sum_{m=1}^3 a_{ik} a_{km} a_{mj} \Rightarrow a_{ik} a_{km} a_{mj} \neq 0$$

$$\iff \underbrace{a_{ik} = a_{km} = a_{mj} = 1}_{v_i \sim v_k \sim v_m \sim v_j}$$

$(A_G^3)_{ij}$ is the number of walks of the length 3 from v_i to v_j

$(A_G^m)_{ij}$ is the number of walks of the length m from v_i to v_j

Remark: Recall, that a **walk** from v_i to v_j in a graph is a sequence of vertices

$$v_i - - v_p - - v_k - - v_r - - v_s - - \dots - - v_j$$

The length of a walk is the number of edges in the walk.

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace

Direct Sum

Basis

Exercise

Remark

1. According to our definition of graphs, $\text{trace}(A_G) = 0$.
2. $(A_G^m)_{ij}$ is the number of walks of the length m from v_i to v_j .
3. In the definition of walks, one can "return" to the place where it comes from.
4. Between two adjacent points i and j , there always exists a walk of the length $(2n + 1)$ for some $n \in \mathbb{N}$ from v_i to v_j .
5. We can always directly write out A_G^n for any $n \in \mathbb{N}$ since we know the definition of elements in such matrix.

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace


Direct Sum

Basis

Exercise

Exercise

Directly write out the adjacency matrix A_G , A_G^2 of this graph.



Linear Combination

Definition: If $\bar{y} = \alpha_1 \bar{x}_1 + \dots + \alpha_k \bar{x}_k$, then \bar{y} is called a **linear combination** of $\bar{x}_1, \dots, \bar{x}_k$ OR we say that \bar{y} **is spanned by** $\bar{x}_1, \dots, \bar{x}_k$ and denote

$$\bar{y} = \text{span}(\bar{x}_1, \dots, \bar{x}_k)$$

Linear Independence

Definition: Elements $v_1, v_2, \dots, v_n \in V$ are said to be **linearly independent** if

$$\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n = 0 \Rightarrow \alpha_1 = \dots = \alpha_n = 0$$

Remark: An infinite set of vectors is said to be linearly independent if *every finite subset is linearly independent*.

Definition: If

$$\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n = 0 \Rightarrow \exists \alpha_i \neq 0$$

then the elements $v_1, v_2, \dots, v_n \in V$ are said to be **linearly dependent**.

Definition: A vector v_i is said to be redundant if it is represented as a linear combination of preceding vectors

$$v_i = \alpha_1 v_1 + \dots + \alpha_{i-1} v_{i-1}$$

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace

Direct Sum

Basis

Exercise

Exercise

Let $A = \begin{bmatrix} 0 & 1 \\ 2 & 3 \end{bmatrix}$. Show that $A^2 = \begin{bmatrix} 2 & 3 \\ 6 & 11 \end{bmatrix}$ is a linear combination of A and I_2 .

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace

Direct Sum

Basis

Exercise

Answer

Solution

We have to find scalars c_1 and c_2 such that

$$A^2 = c_1 A + c_2 I_2,$$

or

$$\begin{bmatrix} 2 & 3 \\ 6 & 11 \end{bmatrix} = c_1 \begin{bmatrix} 0 & 1 \\ 2 & 3 \end{bmatrix} + c_2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

$$c_1=3, c_2=2.$$

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace

Direct Sum

Basis

Exercise

Definition

Linear spaces (or vector spaces)

A *linear space*³ V is a set endowed with a rule for addition (if f and g are in V , then so is $f + g$) and a rule for scalar multiplication (if f is in V and k in \mathbb{R} , then kf is in V) such that these operations satisfy the following eight rules⁴ (for all f, g, h in V and all c, k in \mathbb{R}):

1. $(f + g) + h = f + (g + h)$.
2. $f + g = g + f$.
3. There exists a *neutral element* n in V such that $f + n = f$, for all f in V . This n is unique and denoted by 0 .
4. For each f in V there exists a g in V such that $f + g = 0$. This g is unique and denoted by $(-f)$.
5. $k(f + g) = kf + kg$.
6. $(c + k)f = cf + kf$.
7. $c(kf) = (ck)f$.
8. $1f = f$.

Figure: Definition of Linear Space

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace

Direct Sum

Basis

Exercise

Definition

Definition: Let X be a linear space over a scalar field \mathbb{K} . A real-valued function $\|\cdot\|: X \rightarrow \mathbb{R}$ defined on X is called a **norm** provided

1. $\|x\| \geq 0 \quad \forall x \in X$ and $\|x\| = 0$ iff $x = 0$
2. $\|\alpha x\| = |\alpha| \|x\| \quad \forall x \in X \forall \alpha \in \mathbb{K}$
3. $\|x + y\| \leq \|x\| + \|y\| \quad \forall x, y \in X$

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace

Direct Sum

Basis

Exercise

Definition

Definition: Let V be a linear space over \mathbb{K} .

If $U \subset V$ and U is also a linear space closed w.r.t binary operations defined for V , then we say that U is a **linear subspace** of V :

1. $0_U = 0_V \in U$
2. $u_1, u_2 \in U \rightarrow u_1 + u_2 \in U$
3. $\alpha \in \mathbb{K}, u \in U \rightarrow \alpha u \in U$

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace

Direct Sum

Basis

Exercise

Definition

Definition: Let U_1, \dots, U_m be linear subspaces of V .
The **direct sum** $U_1 \oplus \dots \oplus U_m$ of U_1, \dots, U_m is a linear space s.t.
any of its elements can be *uniquely* represented as

$$u_1 + \dots + u_m, \quad u_i \in U_i, \quad i = 1..m$$

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace

Direct Sum

Basis

Exercise

Definition

Definition: In a lin. space V , elements v_1, \dots, v_m form a **basis** if

1. $V = \text{span}(v_1, \dots, v_m)$, and
2. v_1, \dots, v_m are linear independent.

Remark: If $v_1, v_2, \dots, v_m \in V$ for a basis of V then $\forall x \in V$ there exists a unique representation

$$x = c_1 v_1 + c_2 v_2 + \dots + c_m v_m, \quad c_1, \dots, c_m \in \mathbb{K}.$$

Definition: The scalars c_1, \dots, c_m are called **coordinates** of $x \in V$ in the basis v_1, \dots, v_m .

Examples: In class.

Definition: The number of elements in the basis is called the **dimension** of a linear space.

Matrix Product: Exercise of adjacent matrix

Compute A_G^2 and $\text{trace}(A_G^2)$

Compute A_G^3

Remark

Exercise

Linear Combination

Linear Independence

Exercise

Answer

Linear Space

Definition of Linear Space

Normed Linear Space

Linear Subspace

Direct Sum

Basis

Exercise

Problem 1

Use the matrix A , along with its reduced row-echelon form, $rref(A)$, to answer the problems below:

$$A = \begin{bmatrix} 2 & 2 & 6 & 4 \\ 0 & 1 & 2 & 1 \\ 0 & 4 & 8 & 4 \end{bmatrix}, \quad rref(A) = \begin{bmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

- (a) (2 points) What is the rank of A ?
- (b) (5 points) Give a formula that describes all solutions to $A\vec{x} = \vec{0}$.
- (c) (4 points) Let V denote the set of all solutions from the previous part. Is V a subspace of \mathbb{R}^4 ? Explain.
- (d) (5 points) Find a basis for the image of A . What is the dimension of $image(A)$?
- (e) (4 points) Is $\begin{bmatrix} 10 \\ 1 \\ 4 \end{bmatrix}$ in the image of A ? Explain.

Answer

- (a) (2 points) What is the rank of A ?

Solution:

$\text{rank}(A) = 2$ because there are two leading ones in $\text{rref}(A)$.

- (b) (5 points) Give a formula parametrizing all $\vec{x} \in \mathbb{R}^4$ satisfying $A\vec{x} = \vec{0}$.

Solution:

$$\vec{x} = s \begin{bmatrix} -1 \\ -2 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1 \\ -1 \\ 0 \\ 1 \end{bmatrix}.$$

Answer

- (c) (4 points) Let V denote the set of all solutions from part (b). Is V a subspace of \mathbb{R}^4 ? Explain.

Solution:

Yes. The set of all solutions to $A\vec{x} = \vec{0}$ form the kernel, and the kernel of a matrix with four columns is a subspace of \mathbb{R}^4 . Alternatively, the set of all solutions is the span of the vectors $(-1, -2, 1, 0)$ and $(-1, -1, 0, 1)$, and the span of a set of vectors is a subspace. Or, you could check the conditions defining a subspace directly: Let $\vec{v}_1 = (-1, -2, 1, 0)$ and $\vec{v}_2 = (-1, -1, 0, 1)$.

- (i) If $s = t = 0$, then $\vec{0} = s\vec{v}_1 + t\vec{v}_2$, so $\vec{0}$ is an element of V .
- (ii) If \vec{x}_1 and \vec{x}_2 are in V , there are scalars s_1, t_1, s_2, t_2 such that $\vec{x}_1 = s_1\vec{v}_1 + t_1\vec{v}_2$ and $\vec{x}_2 = s_2\vec{v}_1 + t_2\vec{v}_2$. Then $\vec{x}_1 + \vec{x}_2$ is equal to $(s_1 + s_2)\vec{v}_1 + (t_1 + t_2)\vec{v}_2$, and so by (b) is also in V .
- (iii) If $\vec{x} = s\vec{v}_1 + t\vec{v}_2$ is any vector in V , and k is a scalar, then we have $k\vec{x} = ks\vec{v}_1 + kt\vec{v}_2$, and so $k\vec{x}$ is in V .

- (d) (5 points) Find a basis for the image of A . What is the dimension of $\text{image}(A)$?

Solution:

$\mathcal{B} = \left\{ \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix} \right\}$ is a basis for $\text{image}(A)$ because the columns of A span the image, but $\text{rref}(A)$ indicates the 3rd and 4th column are redundant. The dimension is 2 because the basis we have given has two elements.

Answer

(e) (4 points) Is $\begin{bmatrix} 10 \\ 1 \\ 4 \end{bmatrix}$ in the image of A ? Explain.

Solution:

Yes: The vector $\begin{bmatrix} 10 \\ 1 \\ 4 \end{bmatrix}$ can be written as $4 \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix}$, and therefore is a linear combination of the two basis vectors of the image of A , which means that it is itself in the image of A .

Problem 2

- (a) (5 points) Determine if the vectors below are linearly independent.

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \quad \vec{v}_2 = \begin{bmatrix} 1 \\ -1 \\ 4 \\ 2 \end{bmatrix}, \quad \vec{v}_3 = \begin{bmatrix} 2 \\ 0 \\ 1 \\ 2 \end{bmatrix}$$

- (b) (6 points) Let \vec{w} be the vector below, and let \vec{v}_1 and \vec{v}_3 be as in part (a). For which value(s) of b are the vectors \vec{v}_1 , \vec{w} , and \vec{v}_3 linearly *dependent*?

$$\vec{w} = \begin{bmatrix} 1 \\ -1 \\ b \\ 2 \end{bmatrix}$$

Answer

(a) (5 points) Determine if the vectors below are linearly independent.

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \quad \vec{v}_2 = \begin{bmatrix} 1 \\ -1 \\ 4 \\ 2 \end{bmatrix}, \quad \vec{v}_3 = \begin{bmatrix} 2 \\ 0 \\ 1 \\ 2 \end{bmatrix}$$

Solution:

They are linearly independent because the reduced row echelon form of the matrix $[\vec{v}_1 \ \vec{v}_2 \ \vec{v}_3]$ is:

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

Since there is a pivot in every column, none of \vec{v}_1 , \vec{v}_2 , or \vec{v}_3 are redundant.

Answer

- (b) (6 points) Let \vec{w} be the vector below, and let \vec{v}_1 and \vec{v}_3 be as in part (a). For which value(s) of b are the vectors \vec{v}_1 , \vec{w} , and \vec{v}_3 linearly *dependent*?

$$\vec{w} = \begin{bmatrix} 1 \\ -1 \\ b \\ 2 \end{bmatrix}$$

Solution:

We perform a few steps of row reduction:

$$[\vec{v}_1 \ \vec{w} \ \vec{v}_3] = \begin{bmatrix} 1 & 1 & 2 \\ 1 & -1 & 0 \\ 1 & b & 1 \\ 0 & 2 & 2 \end{bmatrix} \xrightarrow[R_3 - R_1]{R_2 - R_1} \begin{bmatrix} 1 & 1 & 2 \\ 0 & -2 & -2 \\ 0 & b-1 & -1 \\ 0 & 2 & 2 \end{bmatrix} \xrightarrow[\frac{-1}{2}R_2]{R_4 + R_2} \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & 1 \\ 0 & b-1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

From here we can see that the columns are linearly dependent exactly when rows 2 and 3 are scalar multiples of each other, and that this occurs only when $b = 0$.