

Vv214 Linear Algebra

First Midterm Exam - Review class

Li Yuzhou (Cr. Du Yang)

SJTU-UM Joint Institute
Shanghai Jiao-Tong University

June 15, 2019

Table of Contents

Linear Equations and Matrix

- Linear Equation

- Linear equation to Matrix

- Reduced Row-Echelon Form

- Elementary Row Operation

Problem

Span, Linear Independence and Basis

- Span

- Linear Independence

- Subspace and Basis

Kernel, Image and Dimension Formula

- Kernel and Image

- Dimension Formula

Linear Transformation

- Linear Operation

- Column Vectors of a Matrix

- Inverse of Matrices

- Geometric Meaning

Field

Summary

Table of Contents

Linear Equations and Matrix

Linear Equation

Linear equation to Matrix

Reduced Row-Echelon Form

Elementary Row Operation

Problem

Span, Linear Independence and Basis

Span

Linear Independence

Subspace and Basis

Kernel, Image and Dimension Formula

Kernel and Image

Dimension Formula

Linear Transformation

Linear Operation

Column Vectors of a Matrix

Inverse of Matrices

Geometric Meaning

Field

Summary

Linear Equation

Definition

In mathematics, a **linear equation** is an equation that may be put in the form

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n + b = 0,$$

where x_1, \cdots, x_n are the variables (or unknowns or indeterminates), and b, a_1, \cdots, a_n are the coefficients, which are often real numbers.

System of linear equations

Definition

In mathematics, a **system of linear equations** has the form

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1m}x_m = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2m}x_m = b_2 \\ \vdots \\ a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nm}x_m = b_n \end{cases}$$

where here a_{ij} , b_i are coefficients and x_i are unknowns.

Matrix

We can write a system of linear equations in to a matrix form.

Coefficient Matrix

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1m} \\ a_{21} & a_{22} & \cdots & a_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nm} \end{bmatrix} \in \mathbb{R}^{n \times m}$$

Augmented Matrix

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1m} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2m} & b_2 \\ \vdots & \vdots & \ddots & \vdots & \cdots \\ a_{n1} & a_{n2} & \cdots & a_{nm} & b_n \end{bmatrix} \in \mathbb{R}^{n \times (m+1)}$$

Remark

1. For a specific system of linear equations, we use the augmented matrix.
2. Know how to judge the number of solutions to an augmented matrix.

Reduced Row-Echelon Form

A matrix is in **reduced row-echelon form (rref)** if it satisfies all of the following conditions:

- ▶ If a row has nonzero entries, then the first nonzero entry is a 1, called the **leading 1** (or **pivot**) in this row.
- ▶ If a column contains a leading 1, then all the other entries in that column are 0.
- ▶ If a row contains a leading 1, then each row above it contains a leading 1 further to the left.

Definition

The **rank** of a matrix A is the number of leading 1's in $\text{rref}(A)$.

Elementary Row Operation

Types of elementary row operations

- ▶ Divide a row by a nonzero scalar.
- ▶ Subtract a multiple of a row from another row.
- ▶ Swap two rows.

Remarks

- ▶ The elementary row operations will NOT change the rank of a matrix, and will NOT change the solution of a system of linear equations.
- ▶ $\text{Rank}(A) = \text{Max number of independent row vectors of } A = \text{Max number of independent column vectors of } A.$

Table of Contents

Linear Equations and Matrix

Linear Equation

Linear equation to Matrix

Reduced Row-Echelon Form

Elementary Row Operation

Problem

Span, Linear Independence and Basis

Span

Linear Independence

Subspace and Basis

Kernel, Image and Dimension Formula

Kernel and Image

Dimension Formula

Linear Transformation

Linear Operation

Column Vectors of a Matrix

Inverse of Matrices

Geometric Meaning

Field

Summary

Problem

Let A be the following 3×3 matrix:

$$A = \begin{bmatrix} 1 & 0 & 3 \\ 3 & 0 & a \\ 0 & 3 & -a \end{bmatrix}.$$

(a) (6 points) Find all values of a such that the system $A\vec{x} = \vec{0}$ has a unique solution.

(b) (6 points) Find all pairs of values (a, k) such that the system

$$A\vec{x} = \begin{bmatrix} 3 \\ k \\ 9 \end{bmatrix}$$

has infinitely many solutions.

(c) (6 points) For the pairs (a, k) that you found in part (b), find all solutions to the equation

$$A\vec{x} = \begin{bmatrix} 3 \\ k \\ 9 \end{bmatrix}.$$

Express your answer in parametrized form, i.e., as $\{\vec{u} + t\vec{v} : t \in \mathbb{R}\}$.

Answer

- (a) (6 points) Find all values of a such that the system $A\vec{x} = \vec{0}$ has a unique solution.

Solution: We row-reduce the augmented matrix:

$$\begin{aligned} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 3 & 0 & a & 0 \\ 0 & 3 & -a & 0 \end{array} \right] &\xrightarrow{\text{III} + \text{II}, \text{II} - 3\text{I}} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 0 & a-9 & 0 \\ 3 & 3 & 0 & 0 \end{array} \right] \xrightarrow{\text{II} \leftrightarrow \text{III}, \text{II}/3} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & a-9 & 0 \end{array} \right] \\ &\xrightarrow{\text{II} - \text{I}} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & -3 & 0 \\ 0 & 0 & a-9 & 0 \end{array} \right] \end{aligned}$$

From this we see that the system has a unique solution if and only if $a \neq 9$.

- (b) (6 points) Find all pairs of values (a, k) such that the system

$$A\vec{x} = \begin{bmatrix} 3 \\ k \\ 9 \end{bmatrix}$$

has infinitely many solutions.

Solution: For this system to have infinitely many solutions, first of all, we must have $\text{rank}(A) < 3$. From the computation in the previous subpart, this can only happen when $a = 9$. So we may simply consider the case $a = 9$. We row-reduce the augmented matrix:

$$\left[\begin{array}{ccc|c} 1 & 0 & 3 & 3 \\ 3 & 0 & 9 & k \\ 0 & 3 & -9 & 9 \end{array} \right] \xrightarrow{\text{II} - 3\text{I}, \text{III}/3} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 3 \\ 0 & 0 & 0 & k-9 \\ 0 & 1 & -3 & 3 \end{array} \right] \xrightarrow{\text{II} \leftrightarrow \text{III}} \left[\begin{array}{ccc|c} 1 & 0 & 3 & 3 \\ 0 & 1 & -3 & 3 \\ 0 & 0 & 0 & k-9 \end{array} \right].$$

From this we see that to have infinitely many solutions, we must have $k = 9$ as well. Thus the only pair (a, k) for which we get infinitely many solutions is $(a, k) = (9, 9)$.

Answer

(c) (6 points) For the pairs (a, k) that you found in part (b), find all solutions to the equation

$$A\vec{x} = \begin{bmatrix} 3 \\ k \\ 9 \end{bmatrix}.$$

Express your answer in parametrized form, i.e., as $\{\vec{u} + t\vec{v} : t \in \mathbb{R}\}$.

Solution: From the previous subpart we may assume that $(a, k) = (9, 9)$. The augmented matrix in this case is:

$$\left[\begin{array}{ccc|c} 1 & 0 & 3 & 3 \\ 0 & 1 & -3 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

There is one “independent” variable, namely x_3 and two “dependent” variables x_1, x_2 . The general solution is:

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3 - 3t \\ 3 + 3t \\ t \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \\ 0 \end{bmatrix} + t \begin{bmatrix} -3 \\ 3 \\ 1 \end{bmatrix}.$$

Table of Contents

Linear Equations and Matrix

Linear Equation

Linear equation to Matrix

Reduced Row-Echelon Form

Elementary Row Operation

Problem

Span, Linear Independence and Basis

Span

Linear Independence

Subspace and Basis

Kernel, Image and Dimension Formula

Kernel and Image

Dimension Formula

Linear Transformation

Linear Operation

Column Vectors of a Matrix

Inverse of Matrices

Geometric Meaning

Field

Summary

Span

Definition

Consider the vectors v_1, \dots, v_m in \mathbb{R}^n . The set of all linear combinations of the vectors v_1, \dots, v_m is called their **span**:

$$\text{span}(v_1, \dots, v_m) = \{c_1 v_1 + \dots + c_m v_m : c_1, \dots, c_m \in \mathbb{R}\}.$$

Linear Independence

Definition

Consider vectors v_1, \dots, v_m in \mathbb{R}^n .

- ▶ We say that a vector v_i in the list v_1, \dots, v_m is **redundant** if v_i is a linear combination of the preceding vectors v_1, \dots, v_{i-1} .
- ▶ The vectors v_1, \dots, v_m are called **linearly independent** if none of them is redundant. Otherwise, the vectors are called **linearly dependent** (meaning that at least one of them is redundant).

Remark

The vectors v_1, \dots, v_m are linearly independent if and only if

$$c_1 v_1 + \dots + c_m v_m = 0 \quad \Rightarrow \quad c_1 = \dots = c_m = 0.$$

Subspace of \mathbb{R}^n

Definition

A subset W of the vector space \mathbb{R}^n is called a **(linear) subspace** of \mathbb{R}^n if it has the following three properties:

1. W contains the zero vector in \mathbb{R}^n .
2. W is closed under addition: If w_1 and w_2 are both in W , then so is $w_1 + w_2$.
3. W is closed under scalar multiplication: If w is in W and k is an arbitrary scalar, then kw is in W .

Basis

Definition

We say that the vectors v_1, \dots, v_m form a **basis** of a subspace V of \mathbb{R}^n if they span V **and** are linearly independent. (Also, it is required that vectors v_1, \dots, v_m be in V .)

Unique representation

Every vector v in V can be expressed **uniquely** as a linear combination of basis,

$$v = c_1 v_1 + \dots + c_m v_m.$$

Dimension

Consider a subspace V of \mathbb{R}^n . The number of vectors in a basis of V is called the **dimension** of V , denoted by $\dim(V)$,

Table of Contents

Linear Equations and Matrix

Linear Equation

Linear equation to Matrix

Reduced Row-Echelon Form

Elementary Row Operation

Problem

Span, Linear Independence and Basis

Span

Linear Independence

Subspace and Basis

Kernel, Image and Dimension Formula

Kernel and Image

Dimension Formula

Linear Transformation

Linear Operation

Column Vectors of a Matrix

Inverse of Matrices

Geometric Meaning

Field

Summary

Kernel and Image

Image

The **image** of a function (not necessarily linear) consists of all the values the function takes in its target space. If f is a function from X to Y , then

$$\text{image}(f) = \{f(x) : x \in X\}.$$

Kernel

The **kernel** of a linear transformation (matrix) A from \mathbb{R}^m to \mathbb{R}^n consists of all zeros of the transformation, that is, the solutions of the equation $Ax = 0$.

Kernel and Image

Remarks

- ▶ The image of a linear transformation A is the span of the column vectors of A .
- ▶ If A is a linear transformation from \mathbb{R}^m to \mathbb{R}^n , then $\ker(A)$ is a subspace of \mathbb{R}^m and $\text{image}(A)$ is a subspace of \mathbb{R}^n .

Dimension Formula

Remarks

- ▶ $\dim(\text{image}(A)) = \text{rank}(A)$.

Rank-Nullity Theorem

For any $n \times m$ matrix A , or equivalently a linear transform A from \mathbb{R}^m to \mathbb{R}^n , we always have

$$\dim(\ker(A)) + \dim(\text{image}(A)) = m.$$

Problem

We give the matrix M and its row reduction. For each question, make clear how you have computed the answer.

$$M = \begin{bmatrix} 1 & 3 & 1 & 2 & 10 & 4 & 4 \\ 5 & 15 & 5 & 2 & 26 & 4 & 4 \\ 4 & 12 & 4 & 1 & 19 & 3 & 3 \\ 5 & 15 & 5 & 3 & 29 & 1 & 1 \end{bmatrix} \quad \text{rref}(M) = \begin{bmatrix} 1 & 3 & 1 & 0 & 4 & 0 & 0 \\ 0 & 0 & 0 & 1 & 3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

- (a) (3 points) What is the rank of M ? Why?
- (b) (4 points) Give a basis for the image of M and make it clear how you have computed the answer.
- (c) (6 points) Give a basis for the kernel of M and make it clear how you have computed the answer.
- (d) (5 points) For which value(s) of x and y will the vector $\begin{bmatrix} \frac{1}{2} \\ x \\ y \\ 0 \\ 0 \end{bmatrix}$ be in the kernel of M ?
- (e) (2 points) Give any basis for the image of M **other than** the one you gave in part (b). This is meant to be easy, if you understand what a basis is.

Answer

- (a) (3 points) What is the rank of M ?

Solution: The rank of M is the number of pivot columns in $\text{rref}(M)$, i.e. the number of columns with leading ones in $\text{rref}(M)$. In our case the rank of M is 3.

- (b) (4 points) Give a basis for the image of M .

Solution: A basis for the image of M is given by the column vectors in M that become pivot columns in $\text{rref}(M)$. So in our case, a basis for the image of M is given by

$$\begin{bmatrix} 1 \\ 5 \\ 4 \\ 5 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 1 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 4 \\ 3 \\ 1 \end{bmatrix}.$$

Answer

- (c) (6 points) Give a basis for the kernel of M .

Solution: Solving the system of linear equations $Mx = 0$, we obtain

$$x_1 + 3x_2 + x_3 + 4x_5 = 0$$

$$x_4 + 3x_5 = 0$$

$$x_6 + x_7 = 0$$

So the general solution is

$$x_2 \begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -1 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -4 \\ 0 \\ 0 \\ -3 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_7 \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ -1 \\ 1 \end{bmatrix}$$

where x_2, x_3, x_5, x_7 are arbitrary real numbers. A basis for the kernel of M is given by the list of vectors

$$\begin{bmatrix} -3 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -4 \\ 0 \\ 0 \\ -3 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ -1 \\ 1 \end{bmatrix}.$$

Answer

- (d) (5 points) For which value(s) of x and y will the vector $\begin{bmatrix} 1 \\ 2 \\ 1 \\ x \\ y \\ 0 \\ 0 \end{bmatrix}$ be in the kernel of M ?

Solution: The kernel of the row reduced matrix is the same as the kernel of the original matrix, and this computation is much easier with the row reduced matrix. We have

$$\begin{bmatrix} 1 & 3 & 1 & 0 & 4 & 0 & 0 \\ 0 & 0 & 0 & 1 & 3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 1 \\ x \\ y \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 8 + 4y \\ x + 3y \\ 0 \\ 0 \end{bmatrix}.$$

So this is 0 when $8 + 4y = x + 3y = 0$. In other words, $y = -2$ and $x = 6$.

- (e) (2 points) Give any basis for the image of M **other than** the one you gave in part (b). This is meant to be easy, if you understand what a basis is.

Solution: There are many solutions. Perhaps the easiest is to multiply one of the vectors by a scalar, such as 2:

$$\begin{bmatrix} 2 \\ 10 \\ 8 \\ 10 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 1 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 4 \\ 3 \\ 1 \end{bmatrix}.$$

Table of Contents

Linear Equations and Matrix

Linear Equation

Linear equation to Matrix

Reduced Row-Echelon Form

Elementary Row Operation

Problem

Span, Linear Independence and Basis

Span

Linear Independence

Subspace and Basis

Kernel, Image and Dimension Formula

Kernel and Image

Dimension Formula

Linear Transformation

Linear Operation

Column Vectors of a Matrix

Inverse of Matrices

Geometric Meaning

Field

Summary

Linear Operation

Definition

A function T from \mathbb{R}^m to \mathbb{R}^n is called a **linear transformation** if there exists an $n \times m$ matrix A such that

$$T(x) = Ax$$

for all x in the vector space \mathbb{R}^m .

Properties

If A is an $n \times m$ matrix; x and y are vectors in \mathbb{R}^m and k is a scalar, then

1. $A(x + y) = Ax + Ay$, and
2. $A(kx) = k(Ax)$.

Column Vectors of a Matrix

Remarks

- ▶ The i^{th} column vector of the **identical matrix** in \mathbb{R}^n is called **the i^{th} vector of the elementary basis**, denoted by e_i .

- ▶ For example, e_2 in \mathbb{R}^4 is $\begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$.

- ▶ A matrix is a linear transformation which maps e_i to the i^{th} column vector of the matrix.

Inverse of Matrices

Definition

A $n \times n$ matrix A is **invertible** if and only if

- ▶ $\text{rref}(A) = I_n$ or
- ▶ $\text{rank}(A) = n$ or
- ▶ $\det(A) \neq 0$.

Inverse

A matrix A^{-1} is the **inverse** of A if $AA^{-1} = A^{-1}A = I$.

Theorem

$$(AB)^{-1} = B^{-1}A^{-1}.$$

Find the Inverse

Gauss-Jordan method

$$\begin{bmatrix} A & I \end{bmatrix} \xrightarrow{\text{row elimination}} \begin{bmatrix} I & A^{-1} \end{bmatrix}$$

Adjugate matrix method

$A^* := (\text{cof } A)^T$, the transpose of the cofactor matrix of A is called an **adjugate matrix** of A . Then

$$A^{-1} = \frac{1}{\det(A)} A^*.$$

See the slide of RC3.

Geometric Meaning

Orthogonal Projection Matrix

- ▶ $A^2 = A$.
- ▶ Column vectors are on a line.

Reflection Matrix


- ▶ $A^2 = I$, where I is the identity.
- ▶ $A = A^{-1}$.
- ▶ The eigenvalues of A equal ± 1 .

Rotation, Scaling, Shear



- ▶ Recall the general form of the Rotation matrix, Scaling Matrix, Shear and their combination. Refer to the slide in RC3.

Problem

Consider the triangle Δ with vertices $(0,0)$, $(2,0)$, and $(2,1)$. We have drawn this triangle below.



We have drawn the image of Δ , and each of its vertices, under two linear transformations, S and T




(a) (4 points) Give the matrix of S .



(b) (4 points) Give the matrix of T .

Answer

Consider the triangle Δ with vertices $(0,0)$, $(2,0)$, and $(2,1)$. We have drawn this triangle below.



We have drawn the image of Δ , and each of its vertices, under two linear transformations, S and T



(a) (4 points) Give the matrix of S .

Solution: We have

$$S\begin{bmatrix} 1 \\ 0 \end{bmatrix} = \frac{1}{2} \cdot S\begin{bmatrix} 2 \\ 0 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 0 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$S\begin{bmatrix} 0 \\ 1 \end{bmatrix} = S\left(\begin{bmatrix} 2 \\ 1 \end{bmatrix} - \begin{bmatrix} 2 \\ 0 \end{bmatrix}\right) = S\begin{bmatrix} 2 \\ 1 \end{bmatrix} - S\begin{bmatrix} 2 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix} - \begin{bmatrix} 0 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \end{bmatrix}.$$

Therefore, S is represented by

$$\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

Alternatively, S is rotation by $\pi/2$, therefore S is represented by

$$\begin{bmatrix} \cos(\pi/2) & -\sin(\pi/2) \\ \sin(\pi/2) & \cos(\pi/2) \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

Answer

- (b) (4 points) Give the matrix of T .

Solution: We have

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \frac{1}{2} \cdot T \begin{bmatrix} 2 \\ 0 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 2 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$T \begin{bmatrix} 0 \\ 1 \end{bmatrix} = T \left(\begin{bmatrix} 2 \\ 1 \end{bmatrix} - \begin{bmatrix} 2 \\ 0 \end{bmatrix} \right) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \begin{bmatrix} 2 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

Therefore, T is represented by $\begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$.

Remark

Definition 2.2.1 Orthogonal Projections

Consider a line L in the coordinate plane, running through the origin. Any vector \vec{x} in \mathbb{R}^2 can be written uniquely as

$$\vec{x} = \vec{x}^{\parallel} + \vec{x}^{\perp},$$

where \vec{x}^{\parallel} is parallel to line L , and \vec{x}^{\perp} is perpendicular to L .

The transformation $T(\vec{x}) = \vec{x}^{\parallel}$ from \mathbb{R}^2 to \mathbb{R}^2 is called the *orthogonal projection of \vec{x} onto L* , often denoted by $\text{proj}_L(\vec{x})$. If \vec{w} is a nonzero vector parallel to L , then

$$\text{proj}_L(\vec{x}) = \left(\frac{\vec{x} \cdot \vec{w}}{\vec{w} \cdot \vec{w}} \right) \vec{w}.$$

In particular, if $\vec{u} = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$ is a *unit* vector parallel to L , then

$$\text{proj}_L(\vec{x}) = (\vec{x} \cdot \vec{u}) \vec{u}.$$

The transformation $T(\vec{x}) = \text{proj}_L(\vec{x})$ is linear, with matrix

$$\frac{1}{w_1^2 + w_2^2} \begin{bmatrix} w_1^2 & w_1 w_2 \\ w_1 w_2 & w_2^2 \end{bmatrix} = \begin{bmatrix} u_1^2 & u_1 u_2 \\ u_1 u_2 & u_2^2 \end{bmatrix}.$$

In a similar way, we can derive the formula for orthogonal projection and reflection in R^3 .

Table of Contents

Linear Equations and Matrix

Linear Equation

Linear equation to Matrix

Reduced Row-Echelon Form

Elementary Row Operation

Problem

Span, Linear Independence and Basis

Span

Linear Independence

Subspace and Basis

Kernel, Image and Dimension Formula

Kernel and Image

Dimension Formula

Linear Transformation

Linear Operation

Column Vectors of a Matrix

Inverse of Matrices

Geometric Meaning

Field

Summary

Concept

Abelian Groups

Definition: A set A with a binary operation $+: A \times A \rightarrow A$, i.e. $x, y \rightarrow x + y$, is called an **Abelian group** provided

1. $x + y = y + x \quad \forall x, y \in A$ (commutativity)
2. $x + (y + z) = (x + y) + z \quad \forall x, y, z \in A$ (associativity)
3. $\exists 0$ s.t. $x + 0 = 0 + x = x \quad \forall x \in A$ (identity)
4. $\forall x \in A \exists (-x)$ s.t. $x + (-x) = 0$ (inverse)

Examples:

- ▶ $(\mathbb{N}, +)$ no additive inverses
- ▶ $(\mathbb{Z}, +)$ Yes $(\mathbb{Q}, +)$ Yes $(\mathbb{R}, +)$ Yes.
- ▶ $(\mathbb{Z}, \cdot), (\mathbb{R}, \cdot)$ no multiplicative inverses
- ▶ $(\mathbb{Q} \setminus \{0\}, \cdot)$ Yes $(\mathbb{R} \setminus \{0\}, \cdot)$ Yes

Concept

Cyclic Groups

Definition: We say that $a, b \in \mathbb{Z}$ are **congruent modulo m** , $m \in \mathbb{Z}$ and write $a \equiv b \pmod{m}$ if

$$m \mid a - b \Rightarrow a - b = k \cdot m$$

Example: $7 + 3 \pmod{6} = 4$, $12 \pmod{7} = 5$

Definition: We denote \mathbb{Z}_n all integers modulo n ($n > 0$).

Example: $\mathbb{Z}_5 = \{0, 1, 2, 3, 4\}$

Definition: An Abelian group is **cyclic** if A is **generated** by an element: $\exists a \in A: A = \langle a \rangle, \quad \langle a \rangle = \{na, n \in \mathbb{Z}\}$

Example: $\mathbb{Z}_7 = \{0, 1, 2, 3, 4, 5, 6\} \Rightarrow \langle 0 \rangle = \{0\}, \langle 1 \rangle = \mathbb{Z}_7,$

$$\langle 2 \rangle = \{0, 2, 4, 6, 1, 3, 5\}, \langle 3 \rangle = \{0, 3, 6, 2, 5, 1, 4\} = \mathbb{Z}_7$$

1, 2, 3 are the **generators** of \mathbb{Z}_7 . Other generators of \mathbb{Z}_7 ?

Concept

Is \mathbb{Z}_n a linear space?

- ▶ Well, scalar multiplication in \mathbb{Z}_n over **number fields** is not defined. We define a general field:
- ▶ **Definition:** A set \mathbb{F} is called a **field** provided it is an additive Abelian group with the additive inverse 0 and nonzero elements of \mathbb{F} form a multiplicative Abelian group with the multiplicative inverse 1. AND Multiplication distributes addition

Is \mathbb{Z}_n a field?

- ▶ Consider $\mathbb{Z}_4 = \{0, 1, 2, 3\}$.
In \mathbb{Z}_4 , $2 \cdot 2 = 4 = 0 \pmod{4} \Rightarrow 2$ does not have a multiplicative inverse.

\mathbb{Z}_n is a field if $n = p$ is a prime number.

$\mathbb{F}_2 = \mathbb{Z}_2 = \{0, 1\}$ is an important example of a finite field.

Table of Contents

Linear Equations and Matrix

Linear Equation

Linear equation to Matrix

Reduced Row-Echelon Form

Elementary Row Operation

Problem

Span, Linear Independence and Basis

Span

Linear Independence

Subspace and Basis

Kernel, Image and Dimension Formula

Kernel and Image

Dimension Formula

Linear Transformation

Linear Operation

Column Vectors of a Matrix

Inverse of Matrices

Geometric Meaning

Field

Summary

Summary

Go over

- ▶ The textbook,
- ▶ Homework 1-3,
- ▶ Slides and exercises on recitation classes.

For exam

- ▶ No calculator.
- ▶ Good handwriting and clear steps shown contribute to partial credits.
- ▶ Make sure your answer is in clear position.