

MATH 342 Notes

Matthew Hinton

October 9, 2026

1 First Order Vector-Valued ODE

1.1 Repeated eigenvalues

1. Example: Find a fundamental set for $\dot{x} = \begin{bmatrix} 1 & -1 \\ 4 & 5 \end{bmatrix} x$.

$$\lambda = 3, \quad \xi = [1, -2]^T \quad (1)$$

$$(d/dt - \lambda)^2 x = 0 \text{ has solns } Ce^{\lambda t}, Cte^{\lambda t} \quad (2)$$

$$(3)$$

Propose

$$x = te^{\lambda t} \xi ? \quad (4)$$

but then

$$\dot{x} - Ax = e^{\lambda t} \xi \neq 0 \quad (5)$$

so x is not a solution.

However, we can cancel off the nonhomogeneous part with an ansatz solution $e^{\lambda t} \eta$, $\eta \in \mathbb{R}^2$, to the nonhomogeneous equation $\dot{x} - Ax = e^{\lambda t} \xi$. (Compare with method of undetermined coefficients.)

So

$$x = te^{\lambda t} \xi - e^{\lambda t} \eta. \quad (6)$$

Then $\dot{x} - Ax = 0$ gives

$$0 = e^{\lambda t} \xi + (\lambda e^{\lambda t} \eta - Ae^{\lambda t} \eta), \quad (7)$$

and therefore

$$e^{\lambda t} (\xi + \lambda \eta - A\eta) = 0 \implies \xi + \lambda \eta - A\eta = 0 \quad (8)$$

$$\implies \xi = (A - \lambda I)\eta. \quad (9)$$

Notice that $A - \lambda I$ is not invertible. But concretely we want to solve

$$\begin{bmatrix} -2 & -1 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} \eta_1 \\ \eta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}. \quad (10)$$

Is ξ always in the column space of $A - \lambda I$?

2. Summary (choice of a fundamental set).

In general when A has repeated eigenvalues, and A is 2×2 ,

$$x^1 = e^{\lambda t} \xi, \quad x^2 = te^{\lambda t} \xi + e^{\lambda t} \eta, \quad (11)$$

where η is a solution to $(A - \lambda I)\eta = \xi$.

η is called a *generalised eigenvector*.

Highly important result. If η is *any* vector independent of ξ , we will have $(A - \lambda I)\eta = c\xi$ for some $c \neq 0$. (Why?)

1.2 Nonhomogeneous system

1. The system of interest is

$$\dot{x} = P(t)x + g(t), \quad P(t) \in M_{n \times n}(\mathbb{R}), \quad x(t), g(t) \in M_{n \times 1}(\mathbb{R}) \quad (12)$$

Solution is given as a sum of a particular solution with a general solution of the corresponding homogeneous system (which we have already done). We will focus on finding particular solutions x_p to (12).

2. Methods:

- (a) Undetermined coefficients for special $g(t)$
- (b) Laplace transforms
- (c) Decoupled systems for diagonalizable A
- (d) **Variation of parameters** (we will use this one)

Example: $\dot{x} = \begin{bmatrix} 1 & -1 \\ 4 & 5 \end{bmatrix} x + \begin{bmatrix} e^t \\ 0 \end{bmatrix}.$

3. Proposed solutions (VOP):

$$x(t) = u_1(t)x^1(t) + u_2(t)x^2(t) \quad (\text{undetermined coefficients}) \quad (13)$$

where u_1, u_2 are scalar valued and x^1, x^2 solve the homogeneous equation. The goal is to find u_1, u_2 .

- (a) Rewrite

$$x(t) = \begin{bmatrix} x^1(t) & x^2(t) \end{bmatrix} \begin{bmatrix} u_1(t) \\ u_2(t) \end{bmatrix} \quad (14)$$

$$= \Psi(t)u(t) \quad (15)$$

$$= \begin{bmatrix} e^t & e^{5t} \\ -3e^t & e^{5t} \end{bmatrix} \begin{bmatrix} u_1(t) \\ u_2(t) \end{bmatrix} \quad (16)$$

- (b) Apply $x(t) = \Psi(t)u(t)$ to $\dot{x}(t) = Ax(t) + g(t)$ to get

$$\Psi'(t)u(t) + \Psi(t)u'(t) = A\Psi(t)u(t) + g(t) \quad (17)$$

- (c) Remember $\Psi(t)$ is invertible, and also explicitly

$$\Psi(t) = \begin{bmatrix} x^1(t) & x^2(t) \end{bmatrix} \quad (18)$$

$$\Psi'(t) = \begin{bmatrix} Ax^1(t) & Ax^2(t) \end{bmatrix} \quad (19)$$

$$= A\Psi(t). \quad (20)$$

Ψ solves a “matrix version” of the original homogeneous equation. We use this result to simplify (17):

$$\Psi(t)u'(t) = g(t). \quad (21)$$

(d) We have

$$\begin{bmatrix} e^t & e^{5t} \\ -3e^t & e^{5t} \end{bmatrix} \begin{bmatrix} u_1'(t) & u_2'(t) \end{bmatrix} = \begin{bmatrix} e^t & 0 \end{bmatrix}, \quad (22)$$

so

$$\begin{cases} e^t u_1'(t) + e^{5t} u_2'(t) = e^t \\ -3e^t u_1'(t) + e^{5t} u_2'(t) = 0 \end{cases} \quad (23)$$

A solution is

$$\begin{cases} u_1'(t) = 1/4 \\ u_2'(t) = \frac{3}{4}e^{-4t} \end{cases} \quad (24)$$

(e) Recover $u(t)$ from $u'(t)$.

$$u(t) = \int u'(t)dt = \int \begin{bmatrix} \frac{1}{4} \\ \frac{3}{4}e^{-4t} \end{bmatrix} dt = \begin{bmatrix} \frac{1}{4}t \\ -\frac{3}{16}e^{-4t} \end{bmatrix} + c \quad (25)$$

(f) One particular solution is

$$x(t) = \left(\int_0^t \frac{1}{4} ds \right) \begin{bmatrix} e^t \\ -3e^t \end{bmatrix} + \left(\int_0^t \frac{3}{4} e^{-4s} ds \right) \begin{bmatrix} e^{5t} \\ e^{5t} \end{bmatrix}. \quad (26)$$

4. *Solution method in general.*

Problem

$$\dot{x}(t) = P(t)x(t) + g(t).$$

Corresponding homogeneous system

$$\dot{x}(t) = P(t)x(t),$$

solved by

$$\dot{x}_h = P(t)x_h.$$

Solution to the nonhomogeneous system

$$x(t) = \Psi(t)u(t)$$

(solve for $u(t)$).

Equation for $u(t)$

$$\Psi'(t)u(t) + \Psi(t)u'(t) = P(t)\Psi(t)u(t) + g(t).$$

OBSERVE that $\Psi'(t) = P(t)\Psi(t)$ (even in the nonconstant $P(t)$ case), so that

$$\Psi(t)u(t) = g(t).$$

Furthermore, Ψ is invertible. This is because of the whole Wronskian nonzero $\implies$ nonzero everywhere / diffeq's do not collapse the dimension of subspaces spanned by solutions thing. Thus

$$u(t) = \int \Psi(t)^{-1} g(t) dt \quad (27)$$

$$\implies x_p(t) = \Psi(t) \left(c + \int_0^t \Psi(s)^{-1} g(s) ds \right) \quad (28)$$

$$= \Psi(t)c + \int_0^t \Psi(t)\Psi(s)^{-1} g(s) ds, \quad (29)$$

(which is even the general solution to the original equation, with degree of freedom c to be determined by initial conditions).

Special case: $P(t) = A$ (constant). Then

$$x(t) = e^{tA}x(0) = \int_0^t e^{(t-s)A} g(s) ds \quad (\text{Duhamel's formula}) \quad (30)$$

2 Power Series Solutions to ODE

2.1 Basic power series tools

1. Example.

$$f(x) = \sum_{n=1}^{\infty} a_n x^n = \sum_{n=1}^{\infty} \frac{2^n}{n} x^n. \quad (31)$$

(a) Find the radius of convergence.

The radius of convergence of a power series $f(x) = \sum_{n=1}^{\infty} a_n (x - x_0)^n$ is a number $R \geq 0$ such that

- i. $\forall x$ satisfying $|x - x_0| < R$, $f(x)$ converges absolutely (as in, $\sum_{n=1}^{\infty} |a_n| |x - x_0|^n$ converges).
- ii. $\sum_{n=1}^{\infty} a_n (x - x_0)^n$ diverges whenever $|x - x_0| > R$.

R is called the *radius of convergence*.

(Here $R = 1/2$ by the Ratio Test.)

Tools for finding the radius of convergence

The Root Test: If $\lim_{n \rightarrow \infty} |a_n|^{1/n}$ converges, then $R = \frac{1}{\lim_{n \rightarrow \infty} |a_n|^{1/n}}$.

For instance, (31) gives

$$R = \frac{1}{\lim_{n \rightarrow \infty} (2^n/n)^{1/n}} = \frac{1}{\lim_{n \rightarrow \infty} 2/n^{1/n}} = 1/2 \quad \text{since } n^{1/n} \rightarrow 1 \quad (32)$$

The Ratio Test: If $\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}$ converges, then $R = \frac{1}{\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}}$.

In our example,

$$R = \frac{1}{\lim_{n \rightarrow \infty} \frac{2^{n+1}/(n+1)}{2^n/n}} = \frac{1}{\lim_{n \rightarrow \infty} 2n/n+1} = 1/2. \quad (33)$$

(b) Find $f'(x)$ (for meaningful x)

As usual, differentiate term by term. If $f(x)$ is given by a power series $\sum_{n=0}^{\infty} a_n x^n$, then

$$f'(x) = \sum_{n=0}^{\infty} n a_n x^{n-1} = \sum_{n=1}^{\infty} n a_n x^{n-1} \quad (34)$$

By making a change of variable $k = n - 1$, and noticing that the first term drops out of the above series (because $n = 0$), we have

$$f'(x) = \sum_{k=0}^{\infty} (k+1) a_{k+1} x^k. \quad (35)$$

This procedure for getting the power series into the desired powers of x is called *index shifting*. It is important to keep track of the initial value for k and n in the sum. This is analagous to changing the bounds of integration when performing a change of variable.

A remarkable feature of the radius of convergence is that it is *invariant under differentiation or integration of the power series*. Consider our example. Differentiating term by term or using (35), the derivative series is

$$f'(x) = \sum_{k=0}^{\infty} (k+1) \frac{2^{k+1}}{k+1} x^k = \sum_{n=0}^{\infty} 2^{n+1} x^n. \quad (36)$$

By the ratio test, the radius of convergence of $f'(x)$ is

$$R = \frac{1}{\lim_{n \rightarrow \infty} \frac{2^{n+2}}{2^{n+1}}} = \frac{1}{2}, \quad (37)$$

which is the same as the radius of convergence for $f(x)$. The radius of convergence of the integral power series $\int dx f(x)$ is a fun exercise (it should also be $1/2$).

(c) Some notions. Let

$$f(x) = \sum_{n=0}^{\infty} a_n (x - x_0)^n \quad \text{with} \quad R > 0. \quad (38)$$

- i. $f(x)$ is *analytic* at $x = x_0$. (This is what it means to be analytic at x_0 ; there exists a convergent power series equal to the function on a neighborhood of x_0 .)
- ii. $a_n = f^{(n)}(x_0)/n!$, and $f(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)^2 + \dots$, and in particular $f(x_0) = a_0$.
- iii. Taylor series expansion: When f is analytic at x_0 , it has a particular power series representation for $f(x)$, given by

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n \quad (39)$$

For example, the exponential function is its own derivative and $e^0 = 1$. This is enough to define the exponential everywhere by its Taylor series at $x_0 = 0$:

$$e^x = \sum_{n=0}^{\infty} \frac{(e^x)^{(n)}|_{x=0}}{n!} x^n = \sum_{n=0}^{\infty} \frac{e^0}{n!} x^n = \sum_{n=0}^{\infty} \frac{1}{n!} x^n, \quad (40)$$

which should look familiar.


2. Analytic vs infinitely differentiable functions.

Question: Every function that is analytic at $x = 0$ is infinitely differentiable at $x = 0$. Is the converse true? NO!

Example: A classic example is the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) = \begin{cases} e^{-1/x^2}, & x > 0 \\ 0, & x \leq 0 \end{cases} \quad (41)$$

This function is infinitely differentiable at $x_0 = 0$ (why?) but is not analytic there. For if it were analytic, its Taylor series would be identically zero since $f^{(n)}(0) = 0$ for all n (look at the value of f for negative x arbitrarily close to zero).



Fact: $\sum_{n=0}^{\infty} a_n x^n = 0 \implies a_n = 0$ for all n . This is just a restatement of what we said above. (If you define $f(x)$ by that power series then we identify the a_n with the derivatives of f , which are all zero since f is by assumption identically zero.)

2.2 Homogeneous DE with variable coefficients

Consider the general type of ordinary differential equation

$$P(x)y'' + Q(x)y' + R(x)y = 0 \quad (y = y(x) \text{ is real valued}) \quad (42)$$

We will make the following assumption.

Assumption: P, Q, R are polynomials and there is no scalar c such that $P(x), Q(x), R(x)$ are all divisible by $(x - c)$ (they are essentially “coprime” as elements of the ring of polynomials over the real or complex numbers).

1. Well known equations.

(a) Airy’s equation:

$$y'' - xy = 0, \quad P(x) = 1, Q = 0, R(x) = -x. \quad (43)$$

(b) Bessel's equation:

$$x^2 y'' + xy' + (x^2 - \nu^2)y = 0, \quad P(x) = x^2, Q(x) = x, R(x) = x^2 - \nu^2 \quad (44)$$

where ν (the Greek letter “nu”) is a parameter in the equation.

If $\nu \neq 0$, then our assumption (about the coprime nature of P, Q, R) is satisfied, but not if $\nu = 0$. In the case $\nu = 0$, the equation simplifies a fair bit.

(c) Legendre's equation:

$$(1 - x^2)y'' - 2xy' + \alpha(\alpha + 1)y = 0, \quad (45)$$

where α is a parameter.

2. Solution methods.

Let us restate our DE in **standard form**.

$$y'' + \frac{Q}{P}(x)y' + \frac{R}{P}(x)y = 0 \quad (46)$$

or

$$y'' + p(x)y' + q(x)y = 0, \quad (47)$$

where $p(x) = \frac{Q}{P}(x)$, and $q(x) = \frac{R}{P}(x)$.

But wait! We need to verify that it is okay to divide by $P(x)$ (what if we are dividing by zero?). If $P(x_0) \neq 0$, then $P(x) \neq 0$ for all x near x_0 , and the standard form is valid near x_0 . In this situation, x_0 is called an **ordinary point**.

3. Example: Airy's equation.

$$y'' - xy = 0 \quad (x \in \mathbb{R}) \quad (48)$$

We can do some quantitative analysis on Airy's equation and make some rough sketches of various solution curves to (48). This essentially boils down to using the equation to guess the sign and magnitude of y'' at different points on the sketch, and deducing the shape of the solution by forcing it to be consistent with this data. Unfortunately I can't draw in L^AT_EX, so refer to the course notes for details.

To proceed with the solution we use a power series ansatz

$$y(x) = \sum_{n=0}^{\infty} a_n(x - x_0)^n = \sum_{n=0}^{\infty} a_n x^n \quad (49)$$

where $x_0 = 0$.

Solving the (48) amounts to determining the coefficients a_n . To do this we need to plug $y(x)$ into (48). Let's calculate the derivatives (see the short section on power series):

$$y'(x) = \sum_{n=0}^{\infty} \frac{d}{dx} [a_n x^n] \quad (50)$$

$$= \sum_{n=1}^{\infty} n a_n x^{n-1} \quad (51)$$

$$y''(x) = \sum_{n=1}^{\infty} \frac{d}{dx} [n a_n x^{n-1}] \quad (52)$$

$$= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}. \quad (53)$$

The lower limit of the sums is *very important*. We drop terms when they are zero so that we don't end up with x to negative powers in the derivative (e.g. $na_n = 0$ when $n = 0$).

Plugging into the differential equation, we have

$$\sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} - x \sum_{n=0}^{\infty} a_n x^n = 0 \quad (54)$$

$$\implies \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} - \sum_{n=0}^{\infty} a_n x^{n+1} = 0 \quad (55)$$

Doing some index shifting with $k = n - 2$ in the first sum and $k = n + 1$ in the second sum, we get

$$\sum_{k=0}^{\infty} (k+2)(k+1)a_{k+2}x^k - \sum_{k=1}^{\infty} a_{k-1}x^k = 0. \quad (56)$$

Writing out the first few terms, you can convince yourself that we have

$$(2a_2 + 6a_3x + 12a_4x^2 + 20a_5x^3 + \dots) - (a_0x + a_1x^2 + a_2x^3 + a_3x^4 + \dots) = 0. \quad (57)$$

Now we can simply collect terms with the same powers of x and require that all of the coefficients are zero. We can do this because $1, x, x^2, x^3, \dots$ are linearly independent functions on an *interval* (they are naturally called the Taylor basis for polynomials and power series). Since we know that the resulting power series is zero for x in an interval (this technically comes from our hypothesis that the ansatz $y(x)$ is a solution), we can use this to determine a_n for all n .

We could use (57) to determine $a_0, a_1, a_2, \dots$, but we would never find them all. So let's do it formally. From (56), we have

$$2a_2 + \sum_{k=1}^{\infty} [a_{k+2}(k+2)(k+1) - a_{k-1}]x^k = 0, \quad (58)$$

which gives us the equations

$$\begin{cases} 2a_2 = 0, \\ a_{k+2}(k+2)(k+1) - a_{k-1} = 0 \quad \forall k \geq 1 \end{cases} \quad (59)$$

The first equation gives $a_2 = 0$, and the second can be arranged to be

$$a_{k+2} = -\frac{a_{k-1}}{(k+2)(k+1)} \quad \forall k \geq 1 \quad (60)$$

$$\implies a_{k+3} = -\frac{a_k}{(k+3)(k+2)} \quad \forall k \geq 1. \quad (61)$$

So any coefficient determines the coefficient three terms later in the series $f(x)$. Once we have a_0 , we can generate a_3 , and then a_6 , and so on. Naturally, the coefficients form three decoupled sequences or "chains"

$$a_0, a_3, a_6, a_9, \dots, a_{3n}, \dots \quad (62)$$

$$a_1, a_4, a_7, a_{10}, \dots, a_{3n+1}, \dots \quad (63)$$

$$a_2, a_5, a_8, a_{11}, \dots, a_{3n+2}, \dots = 0 \quad (64)$$

The last of these is constant zero because $a_2 = 0$. The other two are controlled by two free parameters a_0 and a_1 . By computing a few of these a_n s you will quickly realise a pattern, which you can write down as an explicit formula for coefficients in the first two chains:

$$a_{3n} = \frac{a_0}{\prod_{k=1}^n (3k-1)(3k)} \quad (65)$$

$$a_{3n+1} = \frac{a_1}{\prod_{k=1}^n 3k(3k+1)} \quad (66)$$

(recall the notation $\prod_{k=1}^n q_k = q_1 \cdot q_2 \cdots q_n$).

With this information we can simplify our ansatz and obtain the general solution.

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \quad (67)$$

$$= \sum_{n=0}^{\infty} a_{3n} x^{3n} + \sum_{n=0}^{\infty} a_{3n+1} x^{3n+1} \quad (68)$$

$$= \sum_{n=0}^{\infty} \frac{a_0}{\prod_{k=1}^n (3k-1)(3k)} x^{3n} + \sum_{n=0}^{\infty} \frac{a_1}{\prod_{k=1}^n 3k(3k+1)} x^{3n+1} \quad (69)$$

$$= a_0 y_1(x) + a_1 y_2(x), \quad (70)$$

where $y_1(x)$ and $y_2(x)$ are the two linearly independent solutions given by the power series in the previous line. This completes the derivation of the general solution to Airy's equation.

Radius of convergence Remember that power series functions are only defined within their radius of convergence. Let's check where our solution to the Airy's equation is actually defined!

For $y_1(x)$,

$$b_n = \frac{a_0}{\prod_{k=1}^n (3k-1)(3k)} \quad (71)$$

and so by the Ratio test

$$R = \frac{1}{\lim_{n \rightarrow \infty} \frac{|b_{n+1}|}{|b_n|}} \quad (72)$$

$$= \frac{1}{\lim_{n \rightarrow \infty} \frac{\prod_{k=1}^n (3k-1)(3k)}{\prod_{k=1}^{n+1} (3k-1)(3k)}} \quad (73)$$

$$= \frac{1}{\lim_{n \rightarrow \infty} \frac{1}{(3n-1)(3n)}} \quad (74)$$

$$= \infty. \quad (75)$$

This means our power series converges everywhere and we have a solution on the whole real line.

As a final remark, we can imagine expanding the solution around some other point $x_0 \neq 0$. If we do this, we get faster convergence around the new x_0 , which is important for numerical applications. In addition, we can imagine situations in which $R < \infty$, in which case it would be necessary to find power series solutions around a collection of points; these solutions can in some cases be “patched together” to get a solution on a larger domain than just $[-R, R]$.