

UNIVERSITY OF COPENHAGEN
Faculty of Life Sciences

Introduction to Bayesian Networks

Advanced Herd Management

28th of september 2009





Tina Birk Jensen


UNIVERSITY OF COPENHAGEN
FACULTY OF LIFE SCIENCES

What is it all about?

Method to reasoning under uncertainty

➡ Where we reason using probabilities





UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Text books/literature

1. Bayesian Networks and Decision Graphs

A *general* textbook on Bayesian networks and decision graphs.

Written by professor Finn Verner Jensen from Ålborg University – one of the leading research centers for Bayesian networks.

2. Bayesian Networks without Tears

Article written by Eugene Charniak

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Software

Esthaug LIMID Software System

www.esthaug.dk

You can download the software from the course homepage

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Outline

Today (28th of september)
General introduction to Bayesian networks:

- What is a Bayesian network?
- Exercise 11.1
- Transmission of evidence
- Exercise 11.2

Tuesday (29th of september)
Building Bayesian network models

Friday (2nd of October)
Case example

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Bayesian networks (in general)

- Graphical model with some restrictions (next slide)
- Basically a static method ("here and now" imagine)
- A static version of data filtering
- All parameters are probabilities

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES







UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Bayesian networks - definition

Bayesian networks consist of:

- A set of variables and a set of directed edges between variables
- Each variable has a finite set of mutually exclusive states

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES



UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Baye's Theorem

$$P(A_i | B_j) = \frac{P(B_j | A_i)P(A_i)}{P(B_j | A_1)p(A_1) + P(B_j | A_2)P(A_2) + \dots + P(B_j | A_n)P(A_n)}$$

$$P(B_j) = \sum_{i=1}^n P(B_j | A_i)P(A_i)$$

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Now an example!!

A small Bayesian network: Pregnancy and heat detection in cows

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Pregnancy and heat detection in cows

What is the probability that a farmer **observes** a particular cow in heat during a 3-week period?

- $P(\text{Heat} = \text{"yes"}) = a$
- $P(\text{Heat} = \text{"no"}) = b$
- $a + b = 1$ (no other options)

What is the probability that the cow is pregnant?

- $P(\text{Pregnant} = \text{"yes"}) = c$
- $P(\text{Pregnant} = \text{"no"}) = d$
- $c + d = 1$ (no other options)

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Conditional probabilities

Now, assume that the cow is pregnant.
What is the conditional probability that the farmer observes it in heat?

- $P(\text{Heat} = \text{"yes"} | \text{Pregnant} = \text{"yes"}) = a_{p+}$
- $P(\text{Heat} = \text{"no"} | \text{Pregnant} = \text{"yes"}) = b_{p+}$
- Again, $a_{p+} + b_{p+} = 1$

Now, assume that the cow is *not* pregnant.
Accordingly:

- $P(\text{Heat} = \text{"yes"} | \text{Pregnant} = \text{"no"}) = a_{p-}$
- $P(\text{Heat} = \text{"no"} | \text{Pregnant} = \text{"no"}) = b_{p-}$
- Again, $a_{p-} + b_{p-} = 1$

Each value of *Pregnant* defines a full probability distribution for *Heat*.
Such a distribution is called **conditional**

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

A small Bayesian net

	<i>Pregnant</i> = "yes"	<i>Pregnant</i> = "no"
	$c = 0.5$	$d = 0.5$

	<i>Heat</i> = "yes"	<i>Heat</i> = "no"
<i>Pregnant</i> = "yes"	$a_{p+} = 0.02$	$b_{p+} = 0.98$
<i>Pregnant</i> = "no"	$a_{p-} = 0.60$	$b_{p-} = 0.40$

Let us build the net!

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Experience with the net: Evidence

By entering information on an observed value of *Heat* we can revise our belief in the value of the **unobservable** variable *Pregnant*.

The observed value of a variable is called **evidence**.

The revision of beliefs is done by use of **Baye's Theorem**:

$$P(A_i | B_j) = \frac{P(B_j | A_i)P(A_i)}{P(B_j | A_1)p(A_1) + P(B_j | A_2)P(A_2) + \dots + P(B_j | A_n)P(A_n)}$$

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Baye's Theorem for our net

How do we use Bayes formula to calculate:
 $P(\text{Pregnant} = \text{"yes"} | \text{Heat} = \text{"yes"})$

$$P(A_i | B_j) = \frac{P(B_j | A_i)P(A_i)}{P(B_j | A_1)p(A_1) + P(B_j | A_2)P(A_2) + \dots + P(B_j | A_n)P(A_n)}$$

$$P(\text{Pregnant} = \text{yes} | \text{Heat} = \text{yes}) = \frac{P(\text{Heat} = \text{yes} | \text{Pregnant} = \text{yes})P(\text{Pregnant} = \text{yes})}{P(\text{Heat} = \text{yes} | \text{Pregnant} = \text{yes})P(\text{Pregnant} = \text{yes}) + P(\text{Heat} = \text{yes} | \text{Pregnant} = \text{no})P(\text{Pregnant} = \text{no})} = \frac{0.02 \times 0.5}{0.02 \times 0.5 + 0.6 \times 0.5} = 0.032258$$

	Heat = "yes"	Heat = "no"
Pregnant = "yes"	$a_{p+} = 0.02$	$b_{p+} = 0.98$
Pregnant = "no"	$a_{p-} = 0.60$	$b_{p-} = 0.40$

Pregnant = "yes" Pregnant = "no"
 $c = 0.5$ $d = 0.5$

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Baye's Theorem for our net

How do we use Bayes formula to calculate:
 $P(\text{Pregnant} = \text{"yes"} | \text{Heat} = \text{"no"})$


$$P(\text{Pregnant} = \text{yes} | \text{Heat} = \text{no}) = \frac{P(\text{Heat} = \text{no} | \text{Pregnant} = \text{yes})P(\text{Pregnant} = \text{yes})}{P(\text{Heat} = \text{no} | \text{Pregnant} = \text{yes})P(\text{Pregnant} = \text{yes}) + P(\text{Heat} = \text{no} | \text{Pregnant} = \text{no})P(\text{Pregnant} = \text{no})} = \frac{0.98 \times 0.5}{0.98 \times 0.5 + 0.4 \times 0.5} = 0.7101449$$

	Heat = "yes"	Heat = "no"
Pregnant = "yes"	$a_{p+} = 0.02$	$b_{p+} = 0.98$
Pregnant = "no"	$a_{p-} = 0.60$	$b_{p-} = 0.40$

Pregnant = "yes" Pregnant = "no"
 $c = 0.5$ $d = 0.5$

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES


Now time for exercise 11.1!



UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Advantages of Bayesian networks

- Consistent combination of information from various sources
- Can estimate certainties for the values of variables that are not observable (or very costly to observe). These variables are called "hypothesis variables".
- These estimates are obtained by entering evidence in "information variables" that
 - Influence the hypothesis variable
 - Depend on the hypothesis variable



UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Transmission of evidence

Serial connections

```

graph LR
    Feed((Feed)) --> Colic((Colic))
    Colic --> Death((Death))
  
```

- If "Colic" is observed, there will be no connection between "Feed" and "Death"
- "Feed" and "Death" are d-separated given "Colic"
- Evidence may be transmitted through a serial connection unless, the state of the intermediate variable is known

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Transmission of evidence

Diverging connection

```

graph TD
    Breed((Breed)) --> Litter_size((Litter size))
    Breed --> Color((Color))
  
```

- If "Breed" is observed, there will be no influence of "Color" on "Litter size"
- "Litter size" and "Color" are d-separated given "Breed"
- Evidence may be transmitted through a diverging connection unless, the state of the intermediate variable is known

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

Transmission of evidence

Converging connection

```

graph TD
    Mastitis((Mastitis)) --> Temp((Temp))
    Heat((Heat)) --> Temp
  
```

- If "Temp" is observed, the information that a cow is not in heat will influence the belief that the cow has mastitis
- Evidence may only be transmitted through a converging connection if a connecting variable (or descendant is observed)

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

The previous example – d-separation

```

graph TD
    Gastro[Gastro Intestinal disorders] --> Diarrhea((Diarrhea))
    Gastro --> ReducedWeightGain((Reduced Weight gain))
    Gastro --> Coughing((Coughing))
    Pneumonia((Pneumonia)) --> ReducedWeightGain
    Pneumonia --> Coughing
  
```

UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

The previous example – d-separation

```

graph TD
    Age((Age)) --> Gastro[Gastro Intestinal disorders]
    Age --> Pneumonia((Pneumonia))
    Season((Season)) --> Gastro
    Season --> Pneumonia
    Gastro --> Diarrhea((Diarrhea))
    Gastro --> ReducedWeightGain((Reduced Weight gain))
    Gastro --> Coughing((Coughing))
    Pneumonia --> ReducedWeightGain
    Pneumonia --> Coughing
  
```


UNIVERSITY OF COPENHAGEN FACULTY OF LIFE SCIENCES

The previous example – d-separation

```

graph TD
    Age((Age)) --> Gastro[Gastro Intestinal disorders]
    Age --> Pneumonia((Pneumonia))
    Season((Season)) --> Gastro
    Season --> Pneumonia
    Gastro --> Diarrhea((Diarrhea))
    Gastro --> ReducedWeightGain((Reduced Weight gain))
    Gastro --> Coughing((Coughing))
    Pneumonia --> ReducedWeightGain
    Pneumonia --> Coughing
  
```

Exercise: Mastitis detection



Compilation of Bayesian networks

Cursory

Compilation:

- Create a **moral graph**
 - Add edges between all pairs of nodes having a common child.
 - Remove all directions
- **Triangulate the moral graph**
 - Add edges until all cycles of more than 3 nodes have a chord
- Identify the cliques of the triangulated graph and organize them into a **junction tree**.

The software system does it automatically (and can show all intermediate stages).



Sum up

Bayesian networks

- Reasoning under uncertainty
- Graphical model with some restrictions
 - Variables and nodes form a DAG
 - All interdependencies are described using conditional probability distributions
 - Can reason against the causal direction
- Consistent combination of information from various sources
- Can estimate certainties for hypothesis variables



Next time (29th of september)

Building Bayesian networks

- Determining the graphical structure
- Determining the conditional probabilities
- Modeling tricks and tips

