

Triplos: Isometric Coordinates and the Pythagorean Theorem

Zane Billings

Department of Mathematics and Computer Science
Western Carolina University

December 9, 2019

Outline

- 1 Triphosian Numbers
- 2 Operations and Distance
 - Algebra Definitions
 - Algebraic Structure of Triphos
 - Distance in Triphos
- 3 Some results in Triphosian geometry
 - Calculating Pi in Triphos
 - The Triphosian Pythagorean Theorem
- 4 Conclusion and Further Questions

Outline

- 1 Triphosian Numbers
- 2 Operations and Distance
 - Algebra Definitions
 - Algebraic Structure of Triphos
 - Distance in Triphos
- 3 Some results in Triphosian geometry
 - Calculating Pi in Triphos
 - The Triphosian Pythagorean Theorem
- 4 Conclusion and Further Questions

Motivation

- What if we used three axes to plot two-dimensional numbers?
- Additive coloring models work similarly: e.g., the RGB model of light.




Figure: The RGB model of light is an additive coloring model which inspired the discovery of Triphos. [1] Source: https://en.wikipedia.org/wiki/File:RGB_illumination.jpg

Geometric Interpretation




Figure: Triphos has three axes, equally spaced 120 degrees apart. We call these axes the red, green, and blue axes, due the motivation behind Triphos. Figure from [2].

Geometric Interpretation




Figure: Each point in Triphosian coordinates is not uniquely determined by an ordered pair, unlike Cartesian coordinates. Left: $P = (3, 4, 1)$, Right: $P = (2, 3, 0)$. Figure from [1].

Formal Definition

- Consider the set of all triples (r, g, b) with real entries.

Formal Definition

- Consider the set of all triples (r, g, b) with real entries.
- We define the following relation: $(r_1, g_1, b_1) \sim (r_2, g_2, b_2)$ if there exists some $c \in \mathbb{R}$ such that

$$(r_2, g_2, b_2) = (r_1 + c, g_1 + c, b_1 + c).$$

Formal Definition

- Consider the set of all triples (r, g, b) with real entries.
- We define the following relation: $(r_1, g_1, b_1) \sim (r_2, g_2, b_2)$ if there exists some $c \in \mathbb{R}$ such that

$$(r_2, g_2, b_2) = (r_1 + c, g_1 + c, b_1 + c).$$

This is an equivalence relation. [2]

Formal Definition

- Consider the set of all triples (r, g, b) with real entries.
- We define the following relation: $(r_1, g_1, b_1) \sim (r_2, g_2, b_2)$ if there exists some $c \in \mathbb{R}$ such that

$$(r_2, g_2, b_2) = (r_1 + c, g_1 + c, b_1 + c).$$

This is an equivalence relation. [2]

- We define the equivalence class $[(r, g, b)]$ to be the set of all triples related to (r, g, b) with this relation.

Formal Definition

- Consider the set of all triples (r, g, b) with real entries.
- We define the following relation: $(r_1, g_1, b_1) \sim (r_2, g_2, b_2)$ if there exists some $c \in \mathbb{R}$ such that

$$(r_2, g_2, b_2) = (r_1 + c, g_1 + c, b_1 + c).$$

This is an equivalence relation. [2]

- We define the equivalence class $[(r, g, b)]$ to be the set of all triples related to (r, g, b) with this relation.
- The **Triphosian reals**, denoted \mathbb{R}_T , refers to the set of all such equivalence classes.

Formal Definition

- Consider the set of all triples (r, g, b) with real entries.
- We define the following relation: $(r_1, g_1, b_1) \sim (r_2, g_2, b_2)$ if there exists some $c \in \mathbb{R}$ such that

$$(r_2, g_2, b_2) = (r_1 + c, g_1 + c, b_1 + c).$$

This is an equivalence relation. [2]

- We define the equivalence class $[(r, g, b)]$ to be the set of all triples related to (r, g, b) with this relation.
- The **Triphosian reals**, denoted \mathbb{R}_T , refers to the set of all such equivalence classes.
- All triples in the same equivalence class represent the same geometric point in Triphosian coordinates.

Equivalence and Standard Form

- Every equivalence class of a Triphosian point has infinitely many members. We designate the standard form triple to represent all members of each equivalence class.

Equivalence and Standard Form

- Every equivalence class of a Triphosian point has infinitely many members. We designate the standard form triple to represent all members of each equivalence class.

Definition

A Triphosian real number, r, g, b , is said to be in **standard form** if all coordinates are nonnegative and at least one coordinate is equal to 0.

Equivalence and Standard Form

- Every equivalence class of a Triphosian point has infinitely many members. We designate the standard form triple to represent all members of each equivalence class.

Definition

A Triphosian real number, r, g, b , is said to be in **standard form** if all coordinates are nonnegative and at least one coordinate is equal to 0.

Example

Let $T = [(-2, 1, 2)]$. We convert to standard form by calculating $T^* = [(-2 + 2, 1 + 2, 2 + 2)] = [(0, 3, 4)]$.

Equivalence and Standard Form

- Every equivalence class of a Triphosian point has infinitely many members. We designate the standard form triple to represent all members of each equivalence class.

Definition

A Triphosian real number, r, g, b , is said to be in **standard form** if all coordinates are nonnegative and at least one coordinate is equal to 0.

Example

Let $T = [(-2, 1, 2)]$. We convert to standard form by calculating $T^* = [(-2 + 2, 1 + 2, 2 + 2)] = [(0, 3, 4)]$.

- If two Triphosian numbers are in standard form, they can only be equivalent if all three coordinates match. [2]

Outline

- 1 Triphosian Numbers
- 2 Operations and Distance
 - Algebra Definitions
 - Algebraic Structure of Triphos
 - Distance in Triphos
- 3 Some results in Triphosian geometry
 - Calculating Pi in Triphos
 - The Triphosian Pythagorean Theorem
- 4 Conclusion and Further Questions

Outline

- 1 Triphosian Numbers
- 2 Operations and Distance
 - Algebra Definitions
 - Algebraic Structure of Triphos
 - Distance in Triphos
- 3 Some results in Triphosian geometry
 - Calculating Pi in Triphos
 - The Triphosian Pythagorean Theorem
- 4 Conclusion and Further Questions

Definition

An **Abelian group** is a nonempty set G equipped with a binary operation $*$ for which the following properties hold.

Definition

An **Abelian group** is a nonempty set G equipped with a binary operation $*$ for which the following properties hold.

- 1 G is closed under $*$: if $x, y \in G$, then $x * y \in G$.

Definition

An **Abelian group** is a nonempty set G equipped with a binary operation $*$ for which the following properties hold.

- 1 G is closed under $*$: if $x, y \in G$, then $x * y \in G$.
- 2 G is associative under $*$: if $x, y, z \in G$, then $(x * y) * z = x * (y * z)$.

Definition

An **Abelian group** is a nonempty set G equipped with a binary operation $*$ for which the following properties hold.

- 1 G is closed under $*$: if $x, y \in G$, then $x * y \in G$.
- 2 G is associative under $*$: if $x, y, z \in G$, then $(x * y) * z = x * (y * z)$.
- 3 G is commutative under $*$: if $x, y \in G$, then $x * y = y * x$.

Definition

An **Abelian group** is a nonempty set G equipped with a binary operation $*$ for which the following properties hold.

- 1 G is closed under $*$: if $x, y \in G$, then $x * y \in G$.
- 2 G is associative under $*$: if $x, y, z \in G$, then $(x * y) * z = x * (y * z)$.
- 3 G is commutative under $*$: if $x, y \in G$, then $x * y = y * x$.
- 4 G contains an identity element e : $x * e = e * x = x$ for all $x \in G$.

Definition

An **Abelian group** is a nonempty set G equipped with a binary operation $*$ for which the following properties hold.

- 1 G is closed under $*$: if $x, y \in G$, then $x * y \in G$.
- 2 G is associative under $*$: if $x, y, z \in G$, then $(x * y) * z = x * (y * z)$.
- 3 G is commutative under $*$: if $x, y \in G$, then $x * y = y * x$.
- 4 G contains an identity element e : $x * e = e * x = x$ for all $x \in G$.
- 5 G contains an inverse element for every element: if $x \in G$, then there is some $y \in G$ such that $x * y = y * x = e$.

Algebra Terminology [3]

Definition

An **Abelian group** is a nonempty set G equipped with a binary operation $*$ for which the following properties hold.

- 1 G is closed under $*$: if $x, y \in G$, then $x * y \in G$.
- 2 G is associative under $*$: if $x, y, z \in G$, then $(x * y) * z = x * (y * z)$.
- 3 G is commutative under $*$: if $x, y \in G$, then $x * y = y * x$.
- 4 G contains an identity element e : $x * e = e * x = x$ for all $x \in G$.
- 5 G contains an inverse element for every element: if $x \in G$, then there is some $y \in G$ such that $x * y = y * x = e$.

Example

- \mathbb{Z} or \mathbb{R} with addition.
- $\mathbb{R} - \{0\}$ with multiplication.
- $\{0, 1\}$ with multiplication modulo 2.

Definition

A **field** is a set F equipped with two operations, addition $(+)$ and multiplication (\cdot) , in which the following properties hold:

Definition

A **field** is a set F equipped with two operations, addition $(+)$ and multiplication (\cdot) , in which the following properties hold:

- 1 F is an Abelian group under addition;

Definition

A **field** is a set F equipped with two operations, addition $(+)$ and multiplication (\cdot) , in which the following properties hold:

- 1 F is an Abelian group under addition;
- 2 F , excluding the additive identity, is an Abelian group under multiplication;

Definition

A **field** is a set F equipped with two operations, addition $(+)$ and multiplication (\cdot) , in which the following properties hold:

- 1 F is an Abelian group under addition;
- 2 F , excluding the additive identity, is an Abelian group under multiplication;
- 3 Addition distributes over multiplication.

Algebra Terminology [3]

Definition

A **field** is a set F equipped with two operations, addition $(+)$ and multiplication (\cdot) , in which the following properties hold:

- 1 F is an Abelian group under addition;
- 2 F , excluding the additive identity, is an Abelian group under multiplication;
- 3 Addition distributes over multiplication.

Example

- \mathbb{Q}, \mathbb{R} , and \mathbb{C} with addition and multiplication.
- $\{0, 1\}$ with addition and multiplication both modulo 2.

Outline

- 1 Triphosian Numbers
- 2 Operations and Distance
 - Algebra Definitions
 - Algebraic Structure of Triphos
 - Distance in Triphos
- 3 Some results in Triphosian geometry
 - Calculating Pi in Triphos
 - The Triphosian Pythagorean Theorem
- 4 Conclusion and Further Questions

Triphosian Addition

Addition definition

$$[(r_1, g_1, b_1)] + [(r_2, g_2, b_2)] = [(r_1 + r_2, g_1 + g_2, b_1 + b_2)]$$

Properties [2]

Triphosian Addition

Addition definition

$$[(r_1, g_1, b_1)] + [(r_2, g_2, b_2)] = [(r_1 + r_2, g_1 + g_2, b_1 + b_2)]$$

Properties [2]

- Well-defined on the Triphosian reals.

Triphosian Addition

Addition definition

$$[(r_1, g_1, b_1)] + [(r_2, g_2, b_2)] = [(r_1 + r_2, g_1 + g_2, b_1 + b_2)]$$

Properties [2]

- Well-defined on the Triphosian reals.
- The Triphosian reals form an Abelian group under addition (closed, associative, and commutative).

Triphosian Addition

Addition definition

$$[(r_1, g_1, b_1)] + [(r_2, g_2, b_2)] = [(r_1 + r_2, g_1 + g_2, b_1 + b_2)]$$

Properties [2]

- Well-defined on the Triphosian reals.
- The Triphosian reals form an Abelian group under addition (closed, associative, and commutative).
- Additive Identity: $[(0, 0, 0)]$

Triphosian Addition

Addition definition

$$[(r_1, g_1, b_1)] + [(r_2, g_2, b_2)] = [(r_1 + r_2, g_1 + g_2, b_1 + b_2)]$$

Properties [2]

- Well-defined on the Triphosian reals.
- The Triphosian reals form an Abelian group under addition (closed, associative, and commutative).
- Additive Identity: $[(0, 0, 0)]$
- Additive Inverse: $[(-r, -g, -b)] = [(g + b, r + b, r + g)] [1]$

Geometry of Addition




Figure: Adding in Triphos corresponds to Euclidean vector addition. The above figure from [1] shows the sum of $P = (2, 3, 0)$ and $Q = (0, 1, 4)$, which gives $P + Q = (2, 4, 4)$.

Triphosian Multiplication

Multiplication definition

$$[(r_1, g_1, b_1)] \cdot [(r_2, g_2, b_2)] = \\ [(r_1 r_2 + g_1 b_2 + b_1 g_2, r_1 g_2 + g_1 r_2 + b_1 b_2, r_1 b_2 + g_1 g_2 + b_1 r_2)]$$

Properties [1, 2]

Triphosian Multiplication

Multiplication definition

$$[(r_1, g_1, b_1)] \cdot [(r_2, g_2, b_2)] = \\ [(r_1 r_2 + g_1 b_2 + b_1 g_2, r_1 g_2 + g_1 r_2 + b_1 b_2, r_1 b_2 + g_1 g_2 + b_1 r_2)]$$

Properties [1, 2]

- Multiplication is well-defined over the Triphosian reals, and is closed, associative, and commutative.

Triphosian Multiplication

Multiplication definition

$$[(r_1, g_1, b_1)] \cdot [(r_2, g_2, b_2)] = \\ [(r_1 r_2 + g_1 b_2 + b_1 g_2, r_1 g_2 + g_1 r_2 + b_1 b_2, r_1 b_2 + g_1 g_2 + b_1 r_2)]$$

Properties [1, 2]

- Multiplication is well-defined over the Triphosian reals, and is closed, associative, and commutative.
- The Triphosian reals form an Abelian group under multiplication, if $[(0, 0, 0)]$ is excluded.

Triphosian Multiplication

Multiplication definition

$$[(r_1, g_1, b_1)] \cdot [(r_2, g_2, b_2)] = \\ [(r_1 r_2 + g_1 b_2 + b_1 g_2, r_1 g_2 + g_1 r_2 + b_1 b_2, r_1 b_2 + g_1 g_2 + b_1 r_2)]$$

Properties [1, 2]

- Multiplication is well-defined over the Triphosian reals, and is closed, associative, and commutative.
- The Triphosian reals form an Abelian group under multiplication, if $[(0, 0, 0)]$ is excluded.
- Multiplicative identity: $[(1, 0, 0)]$

Triphosian Multiplication

Multiplication definition

$$[(r_1, g_1, b_1)] \cdot [(r_2, g_2, b_2)] = \\ [(r_1 r_2 + g_1 b_2 + b_1 g_2, r_1 g_2 + g_1 r_2 + b_1 b_2, r_1 b_2 + g_1 g_2 + b_1 r_2)]$$

Properties [1, 2]

- Multiplication is well-defined over the Triphosian reals, and is closed, associative, and commutative.
- The Triphosian reals form an Abelian group under multiplication, if $[(0, 0, 0)]$ is excluded.
- Multiplicative identity: $[(1, 0, 0)]$
- Multiplicative inverse: $1/n \cdot [(r, g, b)]$ where $n = r^2 + g^2 + b^2 - rb - rg - gb$ and $[(r, g, b)] \neq [(0, 0, 0)]$.

Geometry of Multiplication




Figure: Triphosian multiplication looks messy, but geometrically, *angles add and magnitudes multiply*.

The Triphosian reals form a field

Distributive Property [2]

Given three Triphosian numbers, $[(r_1, g_1, b_1)]$, $[(r_2, g_2, b_2)]$ and $[(r_3, g_3, b_3)]$, the distributive property holds, i.e.

$$[(r_1, g_1, b_1)] \cdot ([[(r_2, g_2, b_2)] + [(r_3, g_3, b_3)]] = \\ [[(r_1, g_1, b_1)] \cdot [(r_2, g_2, b_2)] + [(r_1, g_1, b_1)] \cdot [(r_3, g_3, b_3)].$$

Triphos is a field

Since \mathbb{R}_T is an Abelian group under addition, $\mathbb{R}_T - \{[(0, 0, 0)]\}$ is an Abelian group under multiplication, and addition distributes over multiplication, \mathbb{R}_T is a field.

Outline

- 1 Triphosian Numbers
- 2 Operations and Distance
 - Algebra Definitions
 - Algebraic Structure of Triphos
 - Distance in Triphos
- 3 Some results in Triphosian geometry
 - Calculating Pi in Triphos
 - The Triphosian Pythagorean Theorem
- 4 Conclusion and Further Questions

The Hexa-Metric Function

Definition: The Hexa-Metric function [2]

Let $T_1 = [(r_1, g_1, b_1)]$, $T_2 = [(r_2, g_2, b_2)]$ be Triphosian real numbers. The Hexa-Metric function, H , is defined as

$$H(T_1, T_2) = \min \{ |(r_1 - b_1) - (r_2 - b_2)| + |(g_1 - b_1) - (g_2 - b_2)|, \\ |(r_1 - g_1) - (r_2 - g_2)| + |(b_1 - g_1) - (b_2 - g_2)|, \\ |(g_1 - r_1) - (g_2 - r_2)| + |(b_1 - r_1) - (b_2 - r_2)| \}.$$

Hexa-Metric Properties

- Distance is non-negative: $H(T_1, T_2) \geq 0$ for all $T_1, T_2 \in \mathbb{R}_T$.
- The distance between a point and itself is zero, and no two distinct points have distance zero: $H(T_1, T_2) = 0 \iff T_1 = T_2$.
- Distance is symmetric: $H(T_1, T_2) = H(T_2, T_1)$.
- The Triangle Inequality holds: $H(T_1, T_2) \leq H(T_1, T_3) + H(T_3, T_2)$.

The Hexa-Metric Function




Figure: The Hexa-Metric only allows us to use paths parallel to the axes. Interestingly, there is no unique geodesic like we are used to in Euclidean space. In this figure from [2], all three paths result in a hexa-metric distance of 7 between the two points shown..

Outline

- 1 Triphosian Numbers
- 2 Operations and Distance
 - Algebra Definitions
 - Algebraic Structure of Triphos
 - Distance in Triphos
- 3 Some results in Triphosian geometry
 - Calculating Pi in Triphos
 - The Triphosian Pythagorean Theorem
- 4 Conclusion and Further Questions

Outline

- 1 Triphosian Numbers
- 2 Operations and Distance
 - Algebra Definitions
 - Algebraic Structure of Triphos
 - Distance in Triphos
- 3 Some results in Triphosian geometry
 - Calculating Pi in Triphos
 - The Triphosian Pythagorean Theorem
- 4 Conclusion and Further Questions

The Triphosian Unit Circle and Pi




Figure: A plot of the Triphosian unit circle, generated with MATLAB.

The Triphosian Unit Circle and Pi




Figure: These points (in Triphosian coordinates) are the vertices of the unit circle.

The Triphosian Unit Circle and Pi




Figure: We compute the diameter of the unit circle, as well as the length of each side. Taking the ratio of the circumference to the diameter, we get $\pi_T = 3$.

Outline

- 1 Triphosian Numbers
- 2 Operations and Distance
 - Algebra Definitions
 - Algebraic Structure of Triphos
 - Distance in Triphos
- 3 Some results in Triphosian geometry
 - Calculating Pi in Triphos
 - The Triphosian Pythagorean Theorem
- 4 Conclusion and Further Questions

The Euclidean Pythagorean Theorem




Figure: Suppose we have an arbitrary right triangle. I have rotated it this way for convenience, but any right triangle can be rotated to this orientation.

The Euclidean Pythagorean Theorem




Figure: If we have any triangle with a right angle, we can impose the Euclidean axes over the right angle. We place the origin at the vertex of the right angle, so we can use the distance function to get the length of two legs.

The Euclidean Pythagorean Theorem




Figure: Then, we get the distance of the hypotenuse by taking the distance between the ends of both legs.

Deriving a Pythagorean Theorem in Tripos




Figure: Suppose that we have an arbitrary triangle where one of the angles is 60° .

Deriving a Pythagorean Theorem in Triphos




Figure: Due to the spacing of the three Triphosian axes, we can place the origin at the vertex of the 60° angle and impose Triphosian coordinates. We then know how long the two legs are.

Deriving a Pythagorean Theorem in Triphos




Figure: So, we know the coordinates of the points defining the “hypotenuse”, and we can evaluate the length of the other side.

Triphosian Pythagorean Theorem

At this point, I enlisted Mathematica to assist in computations.

```
HM[r1_, g1_, b1_, r2_, g2_, b2_] :=  
  Min[Abs[(r1 - b1) - (r2 - b2)] + Abs[(g1 - b1) - (g2 - b2)], Abs[(r1 - g1) - (r2 - g2)] + Abs[(b1 - g1) - (b2 - g2)],  
    Abs[(g1 - r1) - (g2 - r2)] + Abs[(b1 - r1) - (b2 - r2)]]  
  
d1 = HM[r1, g1, b1, 0, 0, 0]  
d2 = HM[r2, g2, b2, 0, 0, 0]  
  
Min[Abs[b1 - r1] + Abs[g1 - r1], Abs[-b1 + g1] + Abs[-b1 + r1], Abs[b1 - g1] + Abs[-g1 + r1]]  
  
Min[Abs[b2 - r2] + Abs[g2 - r2], Abs[-b2 + g2] + Abs[-b2 + r2], Abs[b2 - g2] + Abs[-g2 + r2]]  
  
PT = HM[d1, 0, 0, d2, d2, 0]  
  
Min[Abs[Min[Abs[b1 - r1] + Abs[g1 - r1], Abs[-b1 + g1] + Abs[-b1 + r1], Abs[b1 - g1] + Abs[-g1 + r1]] +  
  Abs[Min[Abs[b2 - r2] + Abs[g2 - r2], Abs[-b2 + g2] + Abs[-b2 + r2], Abs[b2 - g2] + Abs[-g2 + r2]]],  
  Abs[Min[Abs[b1 - r1] + Abs[g1 - r1], Abs[-b1 + g1] + Abs[-b1 + r1], Abs[b1 - g1] + Abs[-g1 + r1]] -  
    Min[Abs[b2 - r2] + Abs[g2 - r2], Abs[-b2 + g2] + Abs[-b2 + r2], Abs[b2 - g2] + Abs[-g2 + r2]]] +  
  Abs[Min[Abs[b2 - r2] + Abs[g2 - r2], Abs[-b2 + g2] + Abs[-b2 + r2], Abs[b2 - g2] + Abs[-g2 + r2]]],  
  Abs[Min[Abs[b1 - r1] + Abs[g1 - r1], Abs[-b1 + g1] + Abs[-b1 + r1], Abs[b1 - g1] + Abs[-g1 + r1]] +  
    Abs[-Min[Abs[b1 - r1] + Abs[g1 - r1], Abs[-b1 + g1] + Abs[-b1 + r1], Abs[b1 - g1] + Abs[-g1 + r1]] +  
      Min[Abs[b2 - r2] + Abs[g2 - r2], Abs[-b2 + g2] + Abs[-b2 + r2], Abs[b2 - g2] + Abs[-g2 + r2]]]]]
```

Triphosian Pythagorean Theorem

By simplification, we obtain an equation which is (slightly) less messy.

SPT = FullSimplify[PT]

```
Min[Abs[Min[Abs[b1 - g1] + Abs[b1 - r1], Abs[b1 - g1] + Abs[g1 - r1], Abs[b1 - r1] + Abs[g1 - r1]] -  
  Min[Abs[b2 - g2] + Abs[b2 - r2], Abs[b2 - g2] + Abs[g2 - r2], Abs[b2 - r2] + Abs[g2 - r2]]] +  
  Min[Abs[b1 - g1] + Abs[b1 - r1], Abs[b1 - g1] + Abs[g1 - r1], Abs[b1 - r1] + Abs[g1 - r1]],  
Abs[Min[Abs[b1 - g1] + Abs[b1 - r1], Abs[b1 - g1] + Abs[g1 - r1], Abs[b1 - r1] + Abs[g1 - r1]] -  
  Min[Abs[b2 - g2] + Abs[b2 - r2], Abs[b2 - g2] + Abs[g2 - r2], Abs[b2 - r2] + Abs[g2 - r2]]] +  
  Min[Abs[b2 - g2] + Abs[b2 - r2], Abs[b2 - g2] + Abs[g2 - r2], Abs[b2 - r2] + Abs[g2 - r2]],  
Min[Abs[b1 - g1] + Abs[b1 - r1], Abs[b1 - g1] + Abs[g1 - r1], Abs[b1 - r1] + Abs[g1 - r1]] +  
  Min[Abs[b2 - g2] + Abs[b2 - r2], Abs[b2 - g2] + Abs[g2 - r2], Abs[b2 - r2] + Abs[g2 - r2]]]
```

So, given a triangle with a 60° angle, we can rotate it, impose coordinates, and use this formula to find the length of the third side.

Outline

- 1 Triphosian Numbers
- 2 Operations and Distance
 - Algebra Definitions
 - Algebraic Structure of Triphos
 - Distance in Triphos
- 3 Some results in Triphosian geometry
 - Calculating Pi in Triphos
 - The Triphosian Pythagorean Theorem
- 4 Conclusion and Further Questions

Further Research Areas

- My next step would be to develop a notion of the trigonometric functions in Triphos.

Further Research Areas

- My next step would be to develop a notion of the trigonometric functions in Triphos.
- Is there a bijection between Triphosian equivalence classes and \mathbb{R}^2 (or equivalently, \mathbb{C}), and if so, is it an isomorphism?

Further Research Areas

- My next step would be to develop a notion of the trigonometric functions in Triphos.
- Is there a bijection between Triphosian equivalence classes and \mathbb{R}^2 (or equivalently, \mathbb{C}), and if so, is it an isomorphism?
- Dr. Penland and Dr. Lawson have also posed questions to me regarding the topology and curvature of Triphos respectively.

Further Research Areas

- My next step would be to develop a notion of the trigonometric functions in Triphos.
- Is there a bijection between Triphosian equivalence classes and \mathbb{R}^2 (or equivalently, \mathbb{C}), and if so, is it an isomorphism?
- Dr. Penland and Dr. Lawson have also posed questions to me regarding the topology and curvature of Triphos respectively.
- The authors of [2] and [1] propose interesting open problems regarding calculus, fractals, and functions in Triphos.

Conclusions

- Triphos is an interesting coordinate system which forms a field and a measure space when equipped with certain operations and a metric function. [1, 2]

Conclusions

- Triphos is an interesting coordinate system which forms a field and a measure space when equipped with certain operations and a metric function. [1, 2]

Conclusions

- Triphos is an interesting coordinate system which forms a field and a measure space when equipped with certain operations and a metric function. [1, 2]
- I proved that the field of Triphosian reals has characteristic 0 (not included in this presentation).

Conclusions

- Triphos is an interesting coordinate system which forms a field and a measure space when equipped with certain operations and a metric function. [1, 2]
- I proved that the field of Triphosian reals has characteristic 0 (not included in this presentation).
- I derived the unit circle in Triphos and computed $\pi_T = 3$.

Conclusions

- Triphos is an interesting coordinate system which forms a field and a measure space when equipped with certain operations and a metric function. [1, 2]
- I proved that the field of Triphosian reals has characteristic 0 (not included in this presentation).
- I derived the unit circle in Triphos and computed $\pi_{\mathcal{T}} = 3$.
- Finally, I derived a notion of a “Pythagorean theorem” in Triphos.

Conclusions

- Triphos is an interesting coordinate system which forms a field and a measure space when equipped with certain operations and a metric function. [1, 2]
- I proved that the field of Triphosian reals has characteristic 0 (not included in this presentation).
- I derived the unit circle in Triphos and computed $\pi_{\mathcal{T}} = 3$.
- Finally, I derived a notion of a “Pythagorean theorem” in Triphos.
- There are a lot of open questions remaining about Triphos!

References



Keely Grossnickle, Brian Hollenbeck, Jeana Johnson, and Zhihao Sun.
Triphos: A world without subtraction.
Mathematics Magazine, 92(4):272–285, 2019.



Paula Egging and Erica Johnson.
Triphos: An alternative coordinate system.
The Pentagon, Fall:16–45, 2015.



Serge Lang.
Algebra.
Springer, third edition, 2002.