

Stochastic Methods in Asset Pricing

The MIT Press (2017)

The Martingale Solution to Merton's Problem

(section 14.3)

Andrew Lyasoff

© 2020 by Andrew Lyasoff | www.andrewlyasoff.tech

All rights are reserved.

Prerequisites

Merton's problem

Martingale solution

Prerequisites from Convex Analysis

Merton's Investment-Consumption Problem

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Prerequisites from Convex Analysis

Merton's Investment-Consumption Problem

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Prerequisites from Convex Analysis

Merton's Investment-Consumption Problem

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Prerequisites

Merton's problem

Martingale solution

Prerequisites

Merton's problem

Martingale solution

Given $f : \mathbb{R} \mapsto]-\infty, \infty]$, $\text{Epi}(f) \stackrel{\text{def}}{=} \{(x, y) \in \mathbb{R}^2 : y \geq f(x), x \in \mathbb{R}\}.$

Prerequisites

Merton's problem

Martingale solution

Given $f : \mathbb{R} \mapsto]-\infty, \infty]$, $\text{Epi}(f) \stackrel{\text{def}}{=} \{(x, y) \in \mathbb{R}^2 : y \geq f(x), x \in \mathbb{R}\}$.

f is said to be convex if $\text{Epi}(f)$ is a nonempty closed convex set.

Prerequisites

Merton's problem

Martingale solution

Given $f : \mathbb{R} \mapsto]-\infty, \infty]$, $\text{Epi}(f) \stackrel{\text{def}}{=} \{(x, y) \in \mathbb{R}^2 : y \geq f(x), x \in \mathbb{R}\}$.

f is said to be convex if $\text{Epi}(f)$ is a nonempty closed convex set.

Given $a \in \mathbb{R}$, what is the smallest value for $b \in \mathbb{R}$ such that $f(x) \geq ax - b$ for all $x \in \mathbb{R}$?

Given $f : \mathbb{R} \mapsto]-\infty, \infty]$, $\text{Epi}(f) \stackrel{\text{def}}{=} \{(x, y) \in \mathbb{R}^2 : y \geq f(x), x \in \mathbb{R}\}$.

f is said to be convex if $\text{Epi}(f)$ is a nonempty closed convex set.

Given $a \in \mathbb{R}$, what is the smallest value for $b \in \mathbb{R}$ such that $f(x) \geq ax - b$ for all $x \in \mathbb{R}$?

FENCHEL-LEGENDRE TRANSFORM of f : $\mathbb{R} \ni a \rightsquigarrow f^*(a) \stackrel{\text{def}}{=} \sup_{x \in \mathbb{R}} (ax - f(x)) \in \mathbb{R}$.

Given $f : \mathbb{R} \mapsto]-\infty, \infty]$, $\text{Epi}(f) \stackrel{\text{def}}{=} \{(x, y) \in \mathbb{R}^2 : y \geq f(x), x \in \mathbb{R}\}$.

f is said to be convex if $\text{Epi}(f)$ is a nonempty closed convex set.

Given $a \in \mathbb{R}$, what is the smallest value for $b \in \mathbb{R}$ such that $f(x) \geq ax - b$ for all $x \in \mathbb{R}$?

FENCHEL–LEGENDRE TRANSFORM of f : $\mathbb{R} \ni a \rightsquigarrow f^*(a) \stackrel{\text{def}}{=} \sup_{x \in \mathbb{R}} (ax - f(x)) \in \mathbb{R}$.

N.B. The function f^* is convex and $(f^*)^* = f$ (this is a theorem).

Prerequisites

Merton's problem

Martingale solution

Given $f : \mathbb{R} \mapsto]-\infty, \infty]$, $\text{Epi}(f) \stackrel{\text{def}}{=} \{(x, y) \in \mathbb{R}^2 : y \geq f(x), x \in \mathbb{R}\}$.

f is said to be convex if $\text{Epi}(f)$ is a nonempty closed convex set.

Given $a \in \mathbb{R}$, what is the smallest value for $b \in \mathbb{R}$ such that $f(x) \geq ax - b$ for all $x \in \mathbb{R}$?

FENCHEL–LEGENDRE TRANSFORM of f : $\mathbb{R} \ni a \rightsquigarrow f^*(a) \stackrel{\text{def}}{=} \sup_{x \in \mathbb{R}} (ax - f(x)) \in \mathbb{R}$.

N.B. The function f^* is convex and $(f^*)^* = f$ (this is a theorem).

N.B. $ax - f(x) \leq f^*(a)$ for all $a, x \in \mathbb{R} \iff -ax - f(x) \leq f^*(-a)$ for all $a, x \in \mathbb{R}$,
(if ∂f exists and is continuous, then equality is attained only with $a = \partial f(x)$).

Given $f : \mathbb{R} \mapsto]-\infty, \infty]$, $\text{Epi}(f) \stackrel{\text{def}}{=} \{(x, y) \in \mathbb{R}^2 : y \geq f(x), x \in \mathbb{R}\}$.

f is said to be convex if $\text{Epi}(f)$ is a nonempty closed convex set.

Given $a \in \mathbb{R}$, what is the smallest value for $b \in \mathbb{R}$ such that $f(x) \geq ax - b$ for all $x \in \mathbb{R}$?

FENCHEL–LEGENDRE TRANSFORM of f : $\mathbb{R} \ni a \rightsquigarrow f^*(a) \stackrel{\text{def}}{=} \sup_{x \in \mathbb{R}} (ax - f(x)) \in \mathbb{R}$.

N.B. The function f^* is convex and $(f^*)^* = f$ (this is a theorem).

N.B. $ax - f(x) \leq f^*(a)$ for all $a, x \in \mathbb{R} \iff -ax - f(x) \leq f^*(-a)$ for all $a, x \in \mathbb{R}$,

(if ∂f exists and is continuous, then equality is attained only with $a = \partial f(x)$).

The domain of any utility function $U : \mathbb{R}_{++} \mapsto \mathbb{R}$ can be extended to \mathbb{R} by postulating $U(x) = -\infty$ for all $x \in]-\infty, 0]$.

Prerequisites

Merton's problem

Martingale solution

Given $f : \mathbb{R} \mapsto]-\infty, \infty]$, $\text{Epi}(f) \stackrel{\text{def}}{=} \{(x, y) \in \mathbb{R}^2 : y \geq f(x), x \in \mathbb{R}\}$.

f is said to be convex if $\text{Epi}(f)$ is a nonempty closed convex set.

Given $a \in \mathbb{R}$, what is the smallest value for $b \in \mathbb{R}$ such that $f(x) \geq ax - b$ for all $x \in \mathbb{R}$?

FENCHEL–LEGENDRE TRANSFORM of f : $\mathbb{R} \ni a \rightsquigarrow f^*(a) \stackrel{\text{def}}{=} \sup_{x \in \mathbb{R}} (ax - f(x)) \in \mathbb{R}$.

N.B. The function f^* is convex and $(f^*)^* = f$ (this is a theorem).

N.B. $ax - f(x) \leq f^*(a)$ for all $a, x \in \mathbb{R} \iff -ax - f(x) \leq f^*(-a)$ for all $a, x \in \mathbb{R}$,

(if ∂f exists and is continuous, then equality is attained only with $a = \partial f(x)$).

The domain of any utility function $U : \mathbb{R}_{++} \mapsto \mathbb{R}$ can be extended to \mathbb{R} by postulating $U(x) = -\infty$ for all $x \in]-\infty, 0]$.

INADA'S CONDITION: ∂U is continuous, strictly decreasing on \mathbb{R}_{++} , and such that $\lim_{x \searrow 0} \partial U(x) = \infty$ and $\lim_{x \nearrow \infty} \partial U(x) = 0$.

Prerequisites

Merton's problem

Martingale solution

Given $f : \mathbb{R} \mapsto]-\infty, \infty]$, $\text{Epi}(f) \stackrel{\text{def}}{=} \{(x, y) \in \mathbb{R}^2 : y \geq f(x), x \in \mathbb{R}\}$.

f is said to be convex if $\text{Epi}(f)$ is a nonempty closed convex set.

Given $a \in \mathbb{R}$, what is the smallest value for $b \in \mathbb{R}$ such that $f(x) \geq ax - b$ for all $x \in \mathbb{R}$?

FENCHEL–LEGENDRE TRANSFORM of f : $\mathbb{R} \ni a \rightsquigarrow f^*(a) \stackrel{\text{def}}{=} \sup_{x \in \mathbb{R}} (ax - f(x)) \in \mathbb{R}$.

N.B. The function f^* is convex and $(f^*)^* = f$ (this is a theorem).

N.B. $ax - f(x) \leq f^*(a)$ for all $a, x \in \mathbb{R} \iff -ax - f(x) \leq f^*(-a)$ for all $a, x \in \mathbb{R}$,
(if ∂f exists and is continuous, then equality is attained only with $a = \partial f(x)$).

The domain of any utility function $U : \mathbb{R}_{++} \mapsto \mathbb{R}$ can be extended to \mathbb{R} by postulating $U(x) = -\infty$ for all $x \in]-\infty, 0]$.

INADA'S CONDITION: ∂U is continuous, strictly decreasing on \mathbb{R}_{++} , and such that $\lim_{x \searrow 0} \partial U(x) = \infty$ and $\lim_{x \nearrow \infty} \partial U(x) = 0$. N.B. $-U : \mathbb{R} \mapsto]-\infty, \infty]$, is convex and (from the above)

$$U(x) - ax \leq (-U)^*(-a) \quad \text{for all } a, x \in \mathbb{R},$$

with identities possible for $x \in \mathbb{R}_{++}$ if and only if $\partial U(x) = a \iff x = (\partial U)^{-1}(a)$.

Prerequisites

Merton's problem

Martingale solution

Given $f : \mathbb{R} \mapsto]-\infty, \infty]$, $\text{Epi}(f) \stackrel{\text{def}}{=} \{(x, y) \in \mathbb{R}^2 : y \geq f(x), x \in \mathbb{R}\}$.

f is said to be convex if $\text{Epi}(f)$ is a nonempty closed convex set.

Given $a \in \mathbb{R}$, what is the smallest value for $b \in \mathbb{R}$ such that $f(x) \geq ax - b$ for all $x \in \mathbb{R}$?

FENCHEL–LEGENDRE TRANSFORM of f : $\mathbb{R} \ni a \rightsquigarrow f^*(a) \stackrel{\text{def}}{=} \sup_{x \in \mathbb{R}} (ax - f(x)) \in \mathbb{R}$.

N.B. The function f^* is convex and $(f^*)^* = f$ (this is a theorem).

N.B. $ax - f(x) \leq f^*(a)$ for all $a, x \in \mathbb{R} \iff -ax - f(x) \leq f^*(-a)$ for all $a, x \in \mathbb{R}$,
(if ∂f exists and is continuous, then equality is attained only with $a = \partial f(x)$).

The domain of any utility function $U : \mathbb{R}_{++} \mapsto \mathbb{R}$ can be extended to \mathbb{R} by postulating $U(x) = -\infty$ for all $x \in]-\infty, 0]$.

INADA'S CONDITION: ∂U is continuous, strictly decreasing on \mathbb{R}_{++} , and such that $\lim_{x \searrow 0} \partial U(x) = \infty$ and $\lim_{x \nearrow \infty} \partial U(x) = 0$. **N.B.** $-U : \mathbb{R} \mapsto]-\infty, \infty]$, is convex and (from the above)

$$U(x) - ax \leq (-U)^*(-a) \quad \text{for all } a, x \in \mathbb{R},$$

with identities possible for $x \in \mathbb{R}_{++}$ if and only if $\partial U(x) = a \iff x = (\partial U)^{-1}(a)$.

N.B. $\partial U : \mathbb{R}_{++} \mapsto \mathbb{R}_{++}$ is one-to-one and invertible due to Inada's condition.

Prerequisites

Merton's problem

Martingale solution

Given $f: \mathbb{R} \mapsto]-\infty, \infty]$, $\text{Epi}(f) \stackrel{\text{def}}{=} \{(x, y) \in \mathbb{R}^2 : y \geq f(x), x \in \mathbb{R}\}$.

f is said to be convex if $\text{Epi}(f)$ is a nonempty closed convex set.

Given $a \in \mathbb{R}$, what is the smallest value for $b \in \mathbb{R}$ such that $f(x) \geq ax - b$ for all $x \in \mathbb{R}$?

FENCHEL–LEGENDRE TRANSFORM of $f: \mathbb{R} \ni a \rightsquigarrow f^*(a) \stackrel{\text{def}}{=} \sup_{x \in \mathbb{R}} (ax - f(x)) \in \mathbb{R}$.

N.B. The function f^* is convex and $(f^*)^* = f$ (this is a theorem).

N.B. $ax - f(x) \leq f^*(a)$ for all $a, x \in \mathbb{R} \iff -ax - f(x) \leq f^*(-a)$ for all $a, x \in \mathbb{R}$,

(if ∂f exists and is continuous, then equality is attained only with $a = \partial f(x)$).

The domain of any utility function $U: \mathbb{R}_{++} \mapsto \mathbb{R}$ can be extended to \mathbb{R} by postulating $U(x) = -\infty$ for all $x \in]-\infty, 0]$.

INADA'S CONDITION: ∂U is continuous, strictly decreasing on \mathbb{R}_{++} , and such that $\lim_{x \searrow 0} \partial U(x) = \infty$ and $\lim_{x \nearrow \infty} \partial U(x) = 0$. N.B. $-U: \mathbb{R} \mapsto]-\infty, \infty]$, is convex and (from the above)

$$U(x) - ax \leq (-U)^*(-a) \quad \text{for all } a, x \in \mathbb{R},$$

with identities possible for $x \in \mathbb{R}_{++}$ if and only if $\partial U(x) = a \iff x = (\partial U)^{-1}(a)$.

N.B. $\partial U: \mathbb{R}_{++} \mapsto \mathbb{R}_{++}$ is one-to-one and invertible due to Inada's condition.

Prerequisites

Merton's problem

Martingale solution

Prerequisites

Merton's problem

Martingale solution

Prerequisites

Merton's problem

Martingale solution

An agent has initial wealth x^* , utility from intertemporal consumption U , and utility from final (on date $T > 0$) wealth \bar{U} (Inada's conditions are assumed).

Prerequisites

Merton's problem

Martingale solution

An agent has initial wealth x^* , utility from intertemporal consumption U , and utility from final (on date $T > 0$) wealth \bar{U} (Inada's conditions are assumed). The agent must choose their trading strategy $\pi = (\pi_t)_{t \in [0, T]}$ and a consumption plan $c = (c_t)_{t \in [0, T]}$, so as to maximize their time-separable expected utility from consumption and final wealth:

$$\mathbb{E} \left[\left(e^{-\beta t} U(c) \right) \cdot \iota_T + e^{-\beta T} \bar{U}(V_T) \right],$$

where $V_t \equiv V_t^{x^*, \pi, c}$, $t \in [0, T]$, is the wealth process generated by the strategy (π, c) and the initial endowment x^* .

Prerequisites

Merton's problem

Martingale solution

An agent has initial wealth x^* , utility from intertemporal consumption U , and utility from final (on date $T > 0$) wealth \bar{U} (Inada's conditions are assumed). The agent must choose their trading strategy $\pi = (\pi_t)_{t \in [0, T]}$ and a consumption plan $c = (c_t)_{t \in [0, T]}$, so as to maximize their time-separable expected utility from consumption and final wealth:

$$\mathbb{E} \left[\left(e^{-\beta t} U(c) \right) \cdot \iota_T + e^{-\beta T} \bar{U}(V_T) \right],$$

where $V_t \equiv V_t^{x^*, \pi, c}$, $t \in [0, T]$, is the wealth process generated by the strategy (π, c) and the initial endowment x^* .

The market is complete and the agent's only constraint is:

$$\mathbb{E}^Q \left[\left((S^\circ)^{-1} c \right) \cdot \iota_T + (S_T^\circ)^{-1} V_T \right] = x^*.$$

Prerequisites

Merton's problem

Martingale solution

An agent has initial wealth x^* , utility from intertemporal consumption U , and utility from final (on date $T > 0$) wealth \bar{U} (Inada's conditions are assumed). The agent must choose their trading strategy $\pi = (\pi_t)_{t \in [0, T]}$ and a consumption plan $c = (c_t)_{t \in [0, T]}$, so as to maximize their time-separable expected utility from consumption and final wealth:

$$\mathbb{E} \left[\left(e^{-\beta t} U(c) \right) \cdot \iota_T + e^{-\beta T} \bar{U}(V_T) \right],$$

where $V_t \equiv V_t^{x^*, \pi, c}$, $t \in [0, T]$, is the wealth process generated by the strategy (π, c) and the initial endowment x^* .

The market is complete and the agent's only constraint is:

$$\mathbb{E}^Q \left[\left((S^\circ)^{-1} c \right) \cdot \iota_T + (S_T^\circ)^{-1} V_T \right] = x^*.$$

Introduce the martingale $R_t \stackrel{\text{def}}{=} \frac{d(Q \upharpoonright \mathcal{G}_t)}{d(P \upharpoonright \mathcal{G}_t)}$, $t \in [0, T]$, and write the constraint as

$$\mathbb{E} \left[\left((S^\circ)^{-1} R c \right) \cdot \iota_T + (S_T^\circ)^{-1} R_T V_T \right] = x^*.$$

N.B. Recall that \mathcal{G} stands for the market (observation) filtration.

Prerequisites

Merton's problem

Martingale solution

An agent has initial wealth x^* , utility from intertemporal consumption U , and utility from final (on date $T > 0$) wealth \bar{U} (Inada's conditions are assumed). The agent must choose their trading strategy $\pi = (\pi_t)_{t \in [0, T]}$ and a consumption plan $c = (c_t)_{t \in [0, T]}$, so as to maximize their time-separable expected utility from consumption and final wealth:

$$\mathbb{E} \left[\left(e^{-\beta t} U(c) \right) \cdot \iota_T + e^{-\beta T} \bar{U}(V_T) \right],$$

where $V_t \equiv V_t^{x^*, \pi, c}$, $t \in [0, T]$, is the wealth process generated by the strategy (π, c) and the initial endowment x^* .

The market is complete and the agent's only constraint is:

$$\mathbb{E}^Q \left[\left((S^\circ)^{-1} c \right) \cdot \iota_T + (S_T^\circ)^{-1} V_T \right] = x^*.$$

Introduce the martingale $R_t \stackrel{\text{def}}{=} \frac{d(Q \upharpoonright \mathcal{G}_t)}{d(P \upharpoonright \mathcal{G}_t)}$, $t \in [0, T]$, and write the constraint as (why?)

$$\mathbb{E} \left[\left((S^\circ)^{-1} R c \right) \cdot \iota_T + (S_T^\circ)^{-1} R_T V_T \right] = x^*.$$

N.B. Recall that \mathcal{G} stands for the market (observation) filtration.

Prerequisites

Merton's problem

Martingale solution

An agent has initial wealth x^* , utility from intertemporal consumption U , and utility from final (on date $T > 0$) wealth \bar{U} (Inada's conditions are assumed). The agent must choose their trading strategy $\pi = (\pi_t)_{t \in [0, T]}$ and a consumption plan $c = (c_t)_{t \in [0, T]}$, so as to maximize their time-separable expected utility from consumption and final wealth:

$$\mathbb{E} \left[\left(e^{-\beta t} U(c) \right) \cdot \iota_T + e^{-\beta T} \bar{U}(V_T) \right],$$

where $V_t \equiv V_t^{x^*, \pi, c}$, $t \in [0, T]$, is the wealth process generated by the strategy (π, c) and the initial endowment x^* .

The market is complete and the agent's only constraint is:

$$\mathbb{E}^Q \left[\left((S^\circ)^{-1} c \right) \cdot \iota_T + (S_T^\circ)^{-1} V_T \right] = x^*.$$

Introduce the martingale $R_t \stackrel{\text{def}}{=} \frac{d(Q \upharpoonright \mathcal{G}_t)}{d(P \upharpoonright \mathcal{G}_t)}$, $t \in [0, T]$, and write the constraint as (why?)

$$\mathbb{E} \left[\left((S^\circ)^{-1} R c \right) \cdot \iota_T + (S_T^\circ)^{-1} R_T V_T \right] = x^*.$$

N.B. Recall that \mathcal{G} stands for the market (observation) filtration.

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Develop a solution in terms of the method of Lagrange:

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Develop a solution in terms of the method of Lagrange: transform the constrained problem into the following unconstrained one: (N.B. ϕ is the shadow price of initial wealth so that $\phi \in \mathbb{R}_{++}$.)

$$\inf_{\phi \in \mathbb{R}_{++}} \sup_{c, \pi} L(\pi, c, \phi),$$

where

$$\begin{aligned} L(\pi, c, \phi) = & \mathbb{E} \left[e^{-\beta t} U(c) \cdot \iota_T + e^{-\beta T} \bar{U}(V_T) \right] \\ & + \phi \left(x^* - \mathbb{E} \left[((S^\circ)^{-1} R c) \cdot \iota_T + (S_T^\circ)^{-1} R_T V_T \right] \right). \end{aligned}$$

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Develop a solution in terms of the method of Lagrange: transform the constrained problem into the following unconstrained one: (N.B. ϕ is the shadow price of initial wealth so that $\phi \in \mathbb{R}_{++}$.)

$$\inf_{\phi \in \mathbb{R}_{++}} \sup_{c, \pi} L(\pi, c, \phi),$$

where

$$\begin{aligned} L(\pi, c, \phi) = & \mathbb{E} \left[e^{-\beta t} U(c) \cdot \iota_T + e^{-\beta T} \bar{U}(V_T) \right] \\ & + \phi \left(x^* - \mathbb{E} \left[((S^\circ)^{-1} R c) \cdot \iota_T + (S_T^\circ)^{-1} R_T V_T \right] \right). \end{aligned}$$

By the estimates from the convex analysis prerequisites:

$$\begin{aligned} L(\pi, c, \phi) &= \phi x^* + \mathbb{E} \left[e^{-\beta t} \left(U(c) - \phi e^{\beta t} (S^\circ)^{-1} R c \right) \cdot \iota_T + e^{-\beta T} \left(\bar{U}(V_T) - \phi e^{\beta T} (S_T^\circ)^{-1} R_T V_T \right) \right] \\ &\leq \phi x^* + \mathbb{E} \left[e^{-\beta t} \left((-U)^* \left(-\phi e^{\beta t} (S^\circ)^{-1} R \right) \right) \cdot \iota_T + e^{-\beta T} \left(-\bar{U} \right)^* \left(-\phi e^{\beta T} (S_T^\circ)^{-1} R_T \right) \right]. \end{aligned}$$

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Develop a solution in terms of the method of Lagrange: transform the constrained problem into the following unconstrained one: (N.B. ϕ is the shadow price of initial wealth so that $\phi \in \mathbb{R}_{++}$.)

$$\inf_{\phi \in \mathbb{R}_{++}} \sup_{c, \pi} L(\pi, c, \phi),$$

where

$$\begin{aligned} L(\pi, c, \phi) = & \mathbb{E} \left[e^{-\beta t} U(c) \cdot \iota_T + e^{-\beta T} \bar{U}(V_T) \right] \\ & + \phi \left(x^* - \mathbb{E} \left[((S^\circ)^{-1} R c) \cdot \iota_T + (S_T^\circ)^{-1} R_T V_T \right] \right). \end{aligned}$$

By the estimates from the convex analysis prerequisites:

$$\begin{aligned} L(\pi, c, \phi) &= \phi x^* + \mathbb{E} \left[e^{-\beta t} \left(U(c) - \phi e^{\beta t} (S^\circ)^{-1} R c \right) \cdot \iota_T + e^{-\beta T} \left(\bar{U}(V_T) - \phi e^{\beta T} (S_T^\circ)^{-1} R_T V_T \right) \right] \\ &\leq \phi x^* + \mathbb{E} \left[e^{-\beta t} \left((-U)^* \left(-\phi e^{\beta t} (S^\circ)^{-1} R \right) \right) \cdot \iota_T + e^{-\beta T} \left(-\bar{U} \right)^* \left(-\phi e^{\beta T} (S_T^\circ)^{-1} R_T \right) \right]. \end{aligned}$$

N.B. $\sup_{c, \pi}$ can be attained (for fixed ϕ) only with

$$c_t^\phi = (\partial U)^{-1} \left(\phi e^{\beta t} (S_t^\circ)^{-1} R_t \right) \quad \text{and} \quad V_T^\phi = (\partial \bar{U})^{-1} \left(\phi e^{\beta T} (S_T^\circ)^{-1} R_T \right).$$

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Exercise: If (x^*, π, c) is *any* Q -tame investment-consumption strategy, and if \mathcal{V} is the associated wealth process, then

$$U(c_t^\phi) \geq U(c_t) + \phi e^{\beta t} (S_t^\circ)^{-1} R_t (c_t^\phi - c_t) \quad \text{for all } t \in [0, T],$$

and

$$\bar{U}(V_T^\phi) \geq \bar{U}(\mathcal{V}_T) + \phi e^{\beta T} (S_T^\circ)^{-1} R_T (V_T^\phi - \mathcal{V}_T).$$

Prerequisites

Merton's problem

Martingale solution

Exercise: If (x^*, π, c) is *any* Q -tame investment-consumption strategy, and if \mathcal{V} is the associated wealth process, then

$$U(c_t^\phi) \geq U(c_t) + \phi e^{\beta t} (S_t^\circ)^{-1} R_t (c_t^\phi - c_t) \quad \text{for all } t \in [0, T],$$

and

$$\bar{U}(V_T^\phi) \geq \bar{U}(\mathcal{V}_T) + \phi e^{\beta T} (S_T^\circ)^{-1} R_T (V_T^\phi - \mathcal{V}_T).$$

HINT: $U(c_t^\phi) - (\phi e^{\beta t} (S_t^\circ)^{-1} R_t) c_t^\phi \geq U(c_t) - (\phi e^{\beta t} (S_t^\circ)^{-1} R_t) c_t.$

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Exercise: If (x^*, π, c) is *any* Q -tame investment-consumption strategy, and if \mathcal{V} is the associated wealth process, then

$$U(c_t^\phi) \geq U(c_t) + \phi e^{\beta t} (S_t^\circ)^{-1} R_t (c_t^\phi - c_t) \quad \text{for all } t \in [0, T],$$

and

$$\bar{U}(V_T^\phi) \geq \bar{U}(\mathcal{V}_T) + \phi e^{\beta T} (S_T^\circ)^{-1} R_T (V_T^\phi - \mathcal{V}_T).$$

HINT: $U(c_t^\phi) - (\phi e^{\beta t} (S_t^\circ)^{-1} R_t) c_t^\phi \geq U(c_t) - (\phi e^{\beta t} (S_t^\circ)^{-1} R_t) c_t.$

Can the budget constraint be satisfied with $c = c^\phi$ and $V_T = V_T^\phi$, chosen as in the previous slide (as explicit functions of ϕ), for a particular ϕ ?

Prerequisites

Merton's problem

Martingale solution

Exercise: If (x^*, π, c) is *any* Q -tame investment-consumption strategy, and if \mathcal{V} is the associated wealth process, then

$$U(c_t^\phi) \geq U(c_t) + \phi e^{\beta t} (S_t^\circ)^{-1} R_t (c_t^\phi - c_t) \quad \text{for all } t \in [0, T],$$

and

$$\bar{U}(V_T^\phi) \geq \bar{U}(\mathcal{V}_T) + \phi e^{\beta T} (S_T^\circ)^{-1} R_T (V_T^\phi - \mathcal{V}_T).$$

HINT: $U(c_t^\phi) - (\phi e^{\beta t} (S_t^\circ)^{-1} R_t) c_t^\phi \geq U(c_t) - (\phi e^{\beta t} (S_t^\circ)^{-1} R_t) c_t.$

Can the budget constraint be satisfied with $c = c^\phi$ and $V_T = V_T^\phi$, chosen as in the previous slide (as explicit functions of ϕ), for a particular ϕ ? We must solve for ϕ from

$$\mathbb{E} \left[((S^\circ)^{-1} R c^\phi) \cdot \iota_T + (S_T^\circ)^{-1} R_T V_T^\phi \right] = x^*.$$

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Exercise: If (x^*, π, c) is *any* Q -tame investment-consumption strategy, and if \mathcal{V} is the associated wealth process, then

$$U(c_t^\phi) \geq U(c_t) + \phi e^{\beta t} (S_t^\circ)^{-1} R_t (c_t^\phi - c_t) \quad \text{for all } t \in [0, T],$$

and

$$\bar{U}(V_T^\phi) \geq \bar{U}(\mathcal{V}_T) + \phi e^{\beta T} (S_T^\circ)^{-1} R_T (V_T^\phi - \mathcal{V}_T).$$

HINT: $U(c_t^\phi) - (\phi e^{\beta t} (S_t^\circ)^{-1} R_t) c_t^\phi \geq U(c_t) - (\phi e^{\beta t} (S_t^\circ)^{-1} R_t) c_t.$

Can the budget constraint be satisfied with $c = c^\phi$ and $V_T = V_T^\phi$, chosen as in the previous slide (as explicit functions of ϕ), for a particular ϕ ? We must solve for ϕ from

$$\mathbb{E} \left[((S^\circ)^{-1} R c^\phi) \cdot \iota_T + (S_T^\circ)^{-1} R_T V_T^\phi \right] = x^*.$$

If this equation has a solution $\phi = \phi_*$, then we must find a Q -tame portfolio process π^* , which, together with the consumption process c^{ϕ_*} and the endowment x^* , generates a wealth process with final value that exactly matches $V_T^{\phi_*}$.

Prerequisites

Merton's problem

Martingale solution

Exercise: If (x^*, π, c) is *any* Q -tame investment-consumption strategy, and if \mathcal{V} is the associated wealth process, then

$$U(c_t^\phi) \geq U(c_t) + \phi e^{\beta t} (S_t^\circ)^{-1} R_t (c_t^\phi - c_t) \quad \text{for all } t \in [0, T],$$

and

$$\bar{U}(V_T^\phi) \geq \bar{U}(\mathcal{V}_T) + \phi e^{\beta T} (S_T^\circ)^{-1} R_T (V_T^\phi - \mathcal{V}_T).$$

HINT: $U(c_t^\phi) - (\phi e^{\beta t} (S_t^\circ)^{-1} R_t) c_t^\phi \geq U(c_t) - (\phi e^{\beta t} (S_t^\circ)^{-1} R_t) c_t.$

Can the budget constraint be satisfied with $c = c^\phi$ and $V_T = V_T^\phi$, chosen as in the previous slide (as explicit functions of ϕ), for a particular ϕ ? We must solve for ϕ from

$$\mathbb{E} \left[((S^\circ)^{-1} R c^\phi) \cdot \iota_T + (S_T^\circ)^{-1} R_T V_T^\phi \right] = x^*.$$

If this equation has a solution $\phi = \phi_*$, then we must find a Q -tame portfolio process π^* , which, together with the consumption process c^{ϕ_*} and the endowment x^* , generates a wealth process with final value that exactly matches $V_T^{\phi_*}$.

N.B. This step is the same as the replication of an European-style contingent claim with intertemporal payoff rate $\varphi = c^{\phi_*}$ and with final (closing) payoff $\Phi = V_T^{\phi_*}$.

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Exercise:

Suppose (x^*, π^*, c^{ϕ_*}) is chosen as in the previous slide, and let (x^*, π, c) be *any* other Q -tame investment consumption strategy that is efficient for the initial investment x^* , that is, (x^*, π, c) generates wealth process \mathcal{V} such that $\mathbb{E} \left[((S^\circ)^{-1} R c) \cdot \iota_T + (S_T^\circ)^{-1} R_T \mathcal{V}_T \right] = x^*$. Prove that

$\mathbb{E} \left[(e^{-\beta t} U(c^{\phi_*})) \cdot \iota_T + e^{-\beta T} \bar{U}(V_T^{\phi_*}) \right] \geq \mathbb{E} \left[(e^{-\beta t} U(c)) \cdot \iota_T + e^{-\beta T} \bar{U}(\mathcal{V}_T) \right]$, and conclude that the strategy (x^*, π^*, c^{ϕ_*}) is indeed the optimal strategy.

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Exercise:

Suppose (x^*, π^*, c^{ϕ_*}) is chosen as in the previous slide, and let (x^*, π, c) be *any* other Q -tame investment consumption strategy that is efficient for the initial investment x^* , that is, (x^*, π, c) generates wealth process \mathcal{V} such that $\mathbb{E} \left[((S^\circ)^{-1} R c) \cdot \iota_T + (S_T^\circ)^{-1} R_T \mathcal{V}_T \right] = x^*$. Prove that

$\mathbb{E} \left[(e^{-\beta t} U(c^{\phi_*})) \cdot \iota_T + e^{-\beta T} \bar{U}(V_T^{\phi_*}) \right] \geq \mathbb{E} \left[(e^{-\beta t} U(c)) \cdot \iota_T + e^{-\beta T} \bar{U}(\mathcal{V}_T) \right]$, and conclude that the strategy (x^*, π^*, c^{ϕ_*}) is indeed the optimal strategy.

HINT: From the previous exercise

$$(e^{-\beta t} (U(c^{\phi_*}) - U(c))) \cdot \iota_T + e^{-\beta T} (\bar{U}(V_T^{\phi_*}) - \bar{U}(\mathcal{V}_T)) \geq \phi \left(((S^\circ)^{-1} R(c^{\phi_*} - c)) \cdot \iota_T + (S_T^\circ)^{-1} R_T (V_T^{\phi_*} - \mathcal{V}_T) \right),$$

where the expectation of the right side of the inequality is 0.

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Exercise:

Suppose (x^*, π^*, c^{ϕ_*}) is chosen as in the previous slide, and let (x^*, π, c) be *any* other Q -tame investment consumption strategy that is efficient for the initial investment x^* , that is, (x^*, π, c) generates wealth process \mathcal{V} such that $\mathbb{E} \left[((S^\circ)^{-1} R c) \cdot \iota_T + (S_T^\circ)^{-1} R_T \mathcal{V}_T \right] = x^*$. Prove that

$\mathbb{E} \left[(e^{-\beta t} U(c^{\phi_*})) \cdot \iota_T + e^{-\beta T} \bar{U}(V_T^{\phi_*}) \right] \geq \mathbb{E} \left[(e^{-\beta t} U(c)) \cdot \iota_T + e^{-\beta T} \bar{U}(\mathcal{V}_T) \right]$, and conclude that the strategy (x^*, π^*, c^{ϕ_*}) is indeed the optimal strategy.

HINT: From the previous exercise

$$(e^{-\beta t} (U(c^{\phi_*}) - U(c))) \cdot \iota_T + e^{-\beta T} (\bar{U}(V_T^{\phi_*}) - \bar{U}(\mathcal{V}_T)) \geq \phi \left(((S^\circ)^{-1} R(c^{\phi_*} - c)) \cdot \iota_T + (S_T^\circ)^{-1} R_T (V_T^{\phi_*} - \mathcal{V}_T) \right),$$

where the expectation of the right side of the inequality is 0 (why?).

Martingale solution to Merton's Investment-Consumption Problem

Prerequisites

Merton's problem

Martingale solution

Exercise:

Suppose (x^*, π^*, c^{ϕ_*}) is chosen as in the previous slide, and let (x^*, π, c) be *any* other Q -tame investment consumption strategy that is efficient for the initial investment x^* , that is, (x^*, π, c) generates wealth process \mathcal{V} such that $\mathbb{E}\left[\left((S^\circ)^{-1} R c\right) \cdot \iota_T + (S_T^\circ)^{-1} R_T \mathcal{V}_T\right] = x^*$. Prove that

$\mathbb{E}\left[\left(e^{-\beta t} U(c^{\phi_*})\right) \cdot \iota_T + e^{-\beta T} \bar{U}(V_T^{\phi_*})\right] \geq \mathbb{E}\left[\left(e^{-\beta t} U(c)\right) \cdot \iota_T + e^{-\beta T} \bar{U}(\mathcal{V}_T)\right]$, and conclude that the strategy (x^*, π^*, c^{ϕ_*}) is indeed the optimal strategy.

HINT: From the previous exercise

$$\left(e^{-\beta t} \left(U(c^{\phi_*}) - U(c)\right)\right) \cdot \iota_T + e^{-\beta T} \left(\bar{U}(V_T^{\phi_*}) - \bar{U}(\mathcal{V}_T)\right) \geq \phi\left(\left((S^\circ)^{-1} R(c^{\phi_*} - c)\right) \cdot \iota_T + (S_T^\circ)^{-1} R_T (V_T^{\phi_*} - \mathcal{V}_T)\right),$$

where the expectation of the right side of the inequality is 0.