

Lecture 1, MATH 239 - Introduction to Combinatorics

Graph Theory 1 - Graph, Vertex, Edge, Complete Graph, Path, Cycle, Complete Bipartite Graph

Professor Luke Postle


Notes by Dadi Zhang (dzed.me)


Faculty of Mathematics, University of Waterloo

January 4, 2017


DEFINITION: A **graph** is a set of elements called **vertices** and a set of pairs of distinct vertices called **edges**. If G is a graph, we let $V(G)$ denote the set of vertices and $E(G)$ denote the set of edges.

We tend to draw graphs where vertices are points and edges are lines/curves connecting the pairs of vertices.



Example: $V = \{a\}, E = \emptyset$ 

Example: $V = \{x, y, z\}, E = \{xy, yz\}$ 

Example: $V = \emptyset, E = \emptyset$ known as the empty graph.

Example: $V = \{1, 2, 3, 4, 5\}, E = \{12, 34, 35, 45\}$ 

DEFINITION: K_n denotes the complete graph on n vertices where complete means all pairs of vertices are edges.

Example: K_3  K_4 

DEFINITION: C_n denotes the cycle on n vertices. P_n denotes the path (or path graph) on n vertices.

DEFINITION: $K_{m,n}$ denotes a **complete bipartite graph** on m and n vertices. A complete bipartite graph is a special kind of bipartite graph where every vertex of the first set is connected to every vertex of the second set. $V = \{x_1, x_2, \dots, x_m, y_1, y_2, \dots, y_n\}$

Example: 