

## Chapter 2: Cardiology

Acute Rheumatic Fever and Rheumatic Heart Disease .....	3
Definition .....	3
Risk factors .....	3
Prevention/promotion .....	4
Signs and symptoms .....	5
Investigations .....	6
Differential diagnosis .....	6
Management .....	6
Myocarditis .....	8
Definition .....	8
Risk factors/causes .....	8
Prevention/promotion .....	8
Signs/symptoms .....	8
Investigations .....	8
Management .....	8
Follow up .....	9
Dilated Cardiomyopathy .....	10
Definition .....	10
Risk factors/causes .....	10
Prevention/promotion .....	10
Investigations .....	10
Management .....	10
Pulmonary Hypertension .....	12
Definition .....	12
Risk factors/causes .....	12
Prevention/promotion .....	12
Signs and symptoms .....	12
Investigations .....	13
Management .....	13
Congenital heart diseases .....	14
Definition .....	14
Risk factors/causes .....	14

## Chapter 2: Cardiology

Prevention/promotion .....	14
Signs and symptoms .....	14
Investigations.....	14
Management.....	17
Follow up .....	17
Hypercyanotic spell in tetralogy of fallot.....	18
Definition.....	18
Risk factors for spells.....	18
Prevention/promotion.....	18
Signs and symptoms (clinical features).....	18
Management.....	19
Follow up .....	19
Congestive heart failure.....	21
Definition.....	21
Risk factors/causes .....	21
Prevention/promotion.....	21
Signs and symptoms .....	21
Investigations.....	21
Differential diagnosis.....	22
Management.....	22
Follow up .....	24
Paediatric arrhythmias .....	25
Atrioventricular blocks.....	25
Definition.....	25
Criteria for bradycardia.....	25
Risk factors/causes .....	27
Prevention/promotion.....	27
Signs and symptoms .....	27
Investigations.....	27
Differential diagnosis.....	27
Management.....	28
Primary level.....	28
See the secondary-level guidance below.....	28

Secondary level .....	28
Supraventricular tachycardia .....	29
Definition .....	29
Causes/risk factors.....	29
Signs and symptoms/clinical features.....	29
Management.....	29
Chest pain .....	31
Causes/risk factors (non-cardiac causes):.....	31
Causes/risk factors (cardiac causes):.....	31
Health promotion .....	32
Investigations.....	32
Differential diagnosis .....	32
Management.....	32
Syncope.....	34
Definition .....	34
Causes/risk factors.....	34
Investigations.....	34
Differential diagnosis .....	35
Management.....	35
References .....	36

## Acute Rheumatic Fever and Rheumatic Heart Disease

### Definition

Acute Rheumatic Fever (ARF) is an inflammatory disease of childhood resulting from *Streptococcus pyogenes* (group A streptococcus) pharyngeal or skin infections. Rheumatic Heart Disease (RHD) results from recurrent attacks of acute rheumatic fever causing scarring of mitral and aortic valves. It commonly presents with congestive cardiac failure (CCF) and mitral regurgitation. It commonly occurs between the ages of 5 to 18 years, with a female predominance.

### Risk factors

- Over crowding
- Low social economic status
- Genetic predisposition

### **Prevention/promotion**

- Four weekly intramuscular benzathine penicillin prophylaxis up to 21 years of age in mild cases of rheumatic heart disease and lifelong in moderate to severe cases
- Oral penicillin VK or macrolides in people with elevated risk of adverse reactions i.e.
  - Severe aortic insufficiency,
  - Severe mitral stenosis,
  - Severe aortic stenosis,
  - Ventricular dysfunction (EF <50%), or
  - Severe symptoms (NYHA class III or IV))
- Health education & advocacy
- Screening (as health system permits)

**Signs and symptoms**

<b>Acute Rheumatic Fever*</b>	<b>Rheumatic Heart Disease</b>
<ul style="list-style-type: none"> <li>• Fever</li> <li>• Carditis (presence of a murmur)</li> <li>• Arthritis or arthralgia (joint pain/swelling)</li> <li>• Skin rash</li> <li>• Nodules</li> <li>• Chorea (abnormal movement)</li> </ul>	<ul style="list-style-type: none"> <li>• Clinical features of cardiac failure               <ul style="list-style-type: none"> <li>• Respiratory distress</li> <li>• Tachypnoea</li> <li>• Hepatomegaly</li> </ul> </li> </ul>

\*Also see Modified Jones Criteria below

Diagnosis of Acute Rheumatic Fever

Modified 2015 Jones criteria for High-Risk population

<b>Major Criteria</b>
<ul style="list-style-type: none"> <li>• Carditis (clinical or subclinical)</li> <li>• Arthritis (Monoarthritis or Polyarthritis or polyarthralgia)</li> <li>• Sydenham's chorea</li> <li>• Erythema marginatum</li> <li>• Subcutaneous nodules</li> </ul>
<b>Minor Criteria</b>
<ul style="list-style-type: none"> <li>• Monoarthralgia</li> <li>• Fever (<math>\geq 38.0^{\circ}\text{C}</math>)</li> <li>• ESR <math>\geq 30\text{mm/h}</math> or CRP <math>\geq 3.0\text{mg/dl}</math></li> <li>• Prolonged PR interval (after taking into account the differences related to age, if there is no carditis as a major criterion)</li> </ul>
<b>Making a Diagnosis</b>
<ol style="list-style-type: none"> <li>1. Initial episode of ARF: Evidence of preceding group A <math>\beta</math>-haemolytic streptococcal infection + 2 major criteria or 1 major and 2 minor criteria</li> <li>2. Recurrent attack of ARF (Past ARF or known RHD): 2 major criteria or 1 major and 2 minor criteria or 3 minor criteria</li> </ol>

**In the Malawi setting, it is reasonable to make a diagnosis of ARF without evidence of preceding group A strep infection or supporting lab criteria given the lack of access to lab tests.**

### Investigations

- Acute Rheumatic fever
  - C-reactive protein (CRP)
  - Erythrocyte sedimentation rate (ESR)
  - Antistreptolysin O Titre (ASOT)
  - Throat swab
  - Cardiac echo
  - Electrocardiogram (ECG)
- Rheumatic Heart Disease
  - Electrocardiogram (ECG)
  - Cardiac echo

### Differential diagnosis

- Juvenile idiopathic arthritis
- Septic arthritis
- Post streptococcal reactive arthritis
- Malaria

### Management

#### Primary level

- Give a stat dose of I.M. benzathine penicillin, stabilize and refer to secondary level

#### Secondary level

- Anti-streptococcal therapy
  - STAT dose IM benzathine Penicillin (600,000 IU IM < 25kg; 1,200,000 IU IM > 25kg) then 4-weekly
- Ibuprofen 10 mg/kg TDS for 7 days (until joint pains resolve)
- Chorea
  - Severe/ distressing chorea: Haloperidol 0.02mg/kg once daily (Max 1mg) or sodium valproate 5mg/kg twice daily (Max 600mg/day). Mild cases of chorea do not require specific treatment.
  - Steroids e.g. prednisone can be added in severe/distressing chorea (prednisone 2mg/ kg daily for 2 weeks then taper over 1 to 2 weeks)
- If signs of heart failure, give Furosemide 1 mg/kg BD

### Duration of prophylaxis

21 years of age in mild cases of rheumatic heart diseases and lifelong in moderate to severe cases.

**Note:** All patients should be referred to a tertiary facility

### **Tertiary level**

- Treat as above
- Referral for cardiology review

### **Follow up**

- Regular follow up for benzathine penicillin prophylaxis and management of cardiac failure in PEN-Plus Clinic

## Myocarditis

### Definition

Inflammation of myocardium causing poor contractility of heart muscle.

### Risk factors/causes

- Idiopathic
- Viruses e.g. Adenovirus, Coxsackie
- Bacterial infections
- Fungal infections

### Prevention/promotion

- Health education

### Signs/symptoms

- Signs of heart failure
- A low-grade fever
- Other features of a viral infection, e.g. rhinorrhoea
- Regurgitant murmurs (e.g. pansystolic murmur of mitral regurgitation)
- Myocarditis in children presents with unexplained shortness of breath

### Investigations

- Chest X-Ray
- Echocardiogram
- Electrocardiogram
- HIV test
- Viral screening
- Blood culture

### Management

#### Primary level

- Stabilise the patient and refer

#### Secondary level

- Stabilise the patient
- Treat as heart failure if symptoms present
- Start antibiotics, if any signs of bacterial infection
- Refer to tertiary level



### **Tertiary level**

- Stabilise the patient
- Treat as heart failure if symptoms present.
- Start antibiotics, if any signs of bacterial infection
- Cardiology review

### **Follow up**

- Medication
- Growth
- Review symptoms

## Dilated Cardiomyopathy

### Definition

Dilated cardiomyopathy (DCMO) is defined as left ventricular dilation with systolic dysfunction

### Risk factors/causes

- Idiopathic
- Drugs e.g. Doxorubicin, HAART
- Infections e.g. HIV and other viruses
- Malnutrition
- Hypertension
- Genetic causes

### Prevention/promotion

- Health education
- Signs and symptoms
- Signs of heart failure
- Regurgitant murmurs (e.g. pansystolic murmur of mitral regurgitation)

### Investigations

- Chest X-Ray
- Echocardiogram
- Electrocardiogram
- HIV test

### Management

<b>Primary level</b> <ul style="list-style-type: none"> <li>• Stabilize the patient and refer</li> </ul>
<b>Secondary level</b> <ul style="list-style-type: none"> <li>• Stabilize the patient</li> <li>• Treat as heart failure if symptoms present. See section on heart failure</li> <li>• Refer to tertiary level</li> </ul>
<b>Tertiary level</b> <ul style="list-style-type: none"> <li>• Stabilize the patient</li> <li>• Treat as heart failure if symptoms present. See section on heart failure</li> </ul>

- Cardiology review

### **Follow up**

- Medication
- Growth
- Review of symptoms

## Pulmonary Hypertension

### Definition

Pulmonary hypertension (PH) is a disease characterized by elevated pulmonary arterial pressure, which can result in right ventricular (RV) failure

### Risk factors/causes

The NICE (2013) Updated Classification of Pulmonary Hypertension

<b>1. Pulmonary Arterial Hypertension (PAH)</b> <ul style="list-style-type: none"> <li>• Idiopathic PAH</li> <li>• Heritable PAH</li> <li>• Drug and toxin induced</li> <li>• Associated with connective tissue disease, HIV infection, congenital heart diseases, schistosomiasis</li> </ul>
<b>2. PH due to left heart disease</b> <ul style="list-style-type: none"> <li>• Left ventricular systolic dysfunction</li> <li>• Left ventricular diastolic dysfunction</li> <li>• Valvular disease</li> </ul>
<b>3. Chronic thromboembolic PH</b>
<b>4. PH with unclear multifactorial mechanisms</b> <ul style="list-style-type: none"> <li>• Haematological disorders e.g. chronic haemolytic anaemia</li> <li>• Metabolic disorders e.g. thyroid diseases</li> <li>• Systemic diseases and Others e.g. chronic renal failure</li> </ul>

### Prevention/promotion

- Appropriate management of treatable diseases
- Surgical management of congenital heart diseases in children
- Health awareness and advocacy

### Signs and symptoms

The presentation of PH varies considerably based upon the following; age of the patient, presence and absence of associated medical conditions, and severity of PH and right ventricular function. Signs and Symptoms include:

- Difficulty feeding or shortness of breath on activity
- Fatigue
- Syncope
- Cyanosis

- Failure to thrive
- Cough
- Chest Pains
- Signs of heart Failure
- Right ventricular dilatation/dysfunction: Loud P2, left parasternal heave, systolic/diastolic murmurs, hepatomegaly

### Investigations

Investigations are aimed at identifying the underlying cause and include:

- Echocardiogram: Most helpful in the initial assessment and follow-up of PH
- Chest X-ray
- ECG

### Management

#### Primary level

- Stabilize patient
- Refer to secondary/tertiary facility for pulmonologist and/or Cardiologist review

#### Secondary level

- Stabilize the patient
- Bedrest: avoid strenuous exercise
- Supplemental oxygen
- If presents with signs of heart failure, manage appropriately (see section on Heart failure Management)
- Treat underlying cause of pulmonary hypertension if possible
- Refer to tertiary facility for pulmonology and/or cardiology review

#### Tertiary level

- Manage as above
- Pulmonologist and/or Cardiologist review
- Treat underlying cause of pulmonary hypertension if possible
- Vasodilators e.g. Sildenafil can be considered with pulmonology and/or cardiology consultation
- Involve palliative care

### Follow up

- Regular pulmonology and/or cardiology review

### **Congenital heart diseases**

#### **Definition**

Congenital heart diseases (CHD) encompass a spectrum of structural abnormalities of the heart or intrathoracic vessels.

#### **Risk factors/causes**

- Majority of the cases occur sporadically
- Exposure to teratogenic drugs (e.g. warfarin, sodium valproate)
- Maternal illness: diabetes mellitus and rubella
- Advanced maternal age
- Family history
- Genetic syndromes

#### **Prevention/promotion**

- Avoid drugs that predispose to development of heart defects during pregnancy
- Antenatal screening
- Health education
- Increased awareness of common presentations of congenital heart disease

#### **Signs and symptoms**

- Sweating and/or difficulties breast feeding
- Respiratory distress
- Wheeze/crepitations
- Poor weight gain
- Squatting
- Recurrent chest infections
- Cyanosis
- Effort intolerance, e.g. during play.
- Finger clubbing
- Chest deformity
- Differential saturations ( $\text{SpO}_2$  lower limbs < upper limbs)
- Oedema
- Weak or unequal pulses
- Heart murmur
- Hepatomegaly

#### **Investigations**

- Full blood count


- Chest X-ray
- Electrocardiogram
- Echocardiography

### Algorithm for New-born screening

#### **Newborn screening**

Congenital Heart Defects (CHD) are the most common and severe types of congenital anomalies. Pulse oximetry is an effective and reliable tool for screening and early detection of cyanotic congenital cardiac defects with a sensitivity of 75-83% and specificity of 99%. Pulse oximetry may not detect severe acyanotic lesions in the new-born period. Cardiac auscultation may detect these lesions if a murmur is present.

## Algorithm for new-born screening



RH = right hand

A failed pulse oximetry screen should be referred to a secondary/tertiary level facility for further evaluation. Presence of a murmur on auscultation should prompt referral to secondary/tertiary level facility for further evaluation.



**Management****Primary level**

- Stabilize using ABCDE
- Refer to secondary level

**Secondary level**

- Stabilize using ABCDE
- Treat heart failure if present (see section on heart failure)
  - Furosemide 1-2 mg/kg/dose IV BD (use oral furosemide if IV not available)**and**
  - Spironolactone 1-2 mg/kg/dose PO BD (maximum 25mg/dose)
- Virtual consultation and refer to tertiary centre

**Tertiary level**

- Stabilize using ABCDE
- Can use CPAP if severely distressed/hypoxic
- Treat heart failure if present (see section on heart failure)
  - Furosemide 1-2 mg/kg/dose IV BD (use oral furosemide if IV not available)**and**
  - Spironolactone 1-2 mg/kg/dose PO BD (maximum 25mg/dose)
- Nutritionist/Dietician review
- Cardiologist review and possible corrective surgery referral
- Palliative care team when necessary

**Follow up**

- Once diagnosis is confirmed, the child must be followed up in a PEN-Plus clinic regularly
- Ensure child has adequate stock of medications for home use at each visit
- Monitor growth and development
- Ongoing counselling and education

## **Hypercyanotic spell in tetralogy of fallot**

### **Definition**

Sudden severe episodes of intense cyanosis caused by reduction of pulmonary flow in patients with underlying Tetralogy of Fallot or other cyanotic heart lesions.

### **Risk factors for spells**

- Hypovolaemia, e.g. diarrhoea and vomiting
- Febrile illness
- Pain/agitation
- Anaemia
- Crying, feeding, defaecation
- Cold weather

### **Prevention/promotion**

- Educate parent/guardian on home measures when child is in spell (knee chest position)
- Early health seeking behaviour if acutely unwell
- Avoid unfavourable conditions (pain, cold, agitation, dehydration)
  - Adherence to prescribed medications

### **Signs and symptoms (clinical features)**

- Severe hypoxia/cyanosis
- Hyperpnoea
- Syncope
- Seizure
- Stroke
- Reduced intensity or absence of systolic murmur during spell

**Management****Primary level**

- Keep the child with the parent/guardian and place in a knee chest position
- Place the baby on the mother's shoulder with the knees tucked up underneath.
  - This provides a calming effect, reduces systemic venous return and increases systemic vascular resistance
- Stabilize with ABCDE approach
- Place on oxygen (preferably 100% oxygen)
- Give bolus RL or NS 20 ml/kg rapid IV over 30 minutes to increase preload. Can repeat another bolus if not improving (max 2 boluses)
- Consult and refer to secondary level

**Secondary level**

- As above
- Give PO morphine 0.1mg/kg to reduce distress and hyperpnoea.
- Propranolol 0.5 - 1mg/kg/dose PO TDS
- Consult and refer to tertiary level

**Tertiary level**

- Put on knee chest position
- Place on oxygen (preferably 100% oxygen)
- Give bolus RL or NS 20 ml/kg rapid IV over 30 minutes to increase preload. Can repeat another bolus if not improving (max 3 boluses)
- Give PO/IV morphine 0.1 mg/kg to reduce distress and hyperpnoea
- Do arterial blood gas. If there is severe metabolic acidosis:
  - Give IV sodium bicarbonate 1 mEq/kg STAT to correct
- In resistant cases consider PICU admission for IV Phenylephrine / IV Ketamine in consultation with cardiologist
- Long Term medical management
  - Propranolol 0.5 - 1 mg/kg/dose PO TDS
  - Aspirin 3-5 mg/kg PO daily
  - Ferrous Sulphate and Folic Acid (Fefol) supplement according to weight
- Cardiology and dietician/nutritionist review

Refer for surgical repair if appropriate

**Follow up**

- Once diagnosis is confirmed, the child must be followed up in a PEN-Plus clinic regularly

## Chapter 2: Cardiology

- Ensure child has adequate stock of medications for home use at each visit
- Monitor growth and development
- Ongoing counselling and education

### **Congestive heart failure**

#### **Definition**

Congestive heart failure (CHF) is a clinical condition in which the heart is unable to pump enough blood to the body to meet its needs, to dispose of systemic or pulmonary venous return adequately, or a combination of the two.

#### **Risk factors/causes**

- Cardiac causes
  - Congenital heart diseases
  - Acquired heart diseases
  - Arrhythmias
- Non-cardiac causes:
  - Anaemia
  - Sepsis
  - Chronic kidney disease
  - Auto-immune

#### **Prevention/promotion**

- Health education on early health seeking behaviour
- Early referral of patients with suspected heart disease with or without heart failure.
- Early detection and treatment of anaemia

#### **Signs and symptoms**

- In infants, tachycardia, tachypnoea and hepatomegaly are the most common signs
- Respiratory distress
- Cardiomegaly (displaced apex beat)
- Exertional dyspnoea/sweating during breastfeeding in infants
- Presence of bibasal crepitations
- Signs of shock; reduced pulse volume, delayed capillary refill time, tachycardia, hypotension
- Gallop rhythm
- Raised JVP (older child)
- Peripheral/sacral oedema (oedema will be in dependant area)
- Failure to thrive/severe malnutrition

#### **Investigations**

- Imaging

- Chest x-ray
- Echocardiogram
- ECG
- Blood tests
  - HIV test
  - Full blood count
  - Urea and electrolytes
  - Liver function tests
- Urine dipstick
- Other investigations according to suspected aetiology

### Differential diagnosis

- Pneumonia
- Bronchiolitis
- Asthma
- Nephrotic/nephritic syndrome
- Renal failure
- Chronic liver failure

### Management

#### Primary level

- Stabilize using ABCDE
  - If available give a single dose of Frusemide 1mg/kg IV stat (use oral furosemide if IV not available)
- Refer to secondary level

#### Secondary level

- Stabilize using ABCDE
- Treat heart failure if present (see section on heart failure)
  - Furosemide 1-2 mg/kg/dose IV BD (use oral furosemide if IV not available)
  - and**
  - Spironolactone 1-2 mg/kg/dose PO BD (maximum 25mg/dose)
- Virtual consultation and refer to tertiary centre

#### Tertiary level

- Stabilize using ABCDE
- Can use CPAP if severely distressed/hypoxic
- Treat heart failure if present (see section on heart failure)
  - Furosemide 1-2 mg/kg/dose IV BD (use oral furosemide if IV not available)
  - and**

- Spironolactone 1-2 mg/kg/dose PO BD (maximum 25mg/dose)
- Nutritionist/Dietician review
- Cardiologist review and possible corrective surgery referral
- Palliative care team when necessary

### **Spironolactone (potassium sparing diuretic, modest diuretic effect)**

- Dose: 1 mg/kg/dose BD

#### Afterload reduction: second line

(Discuss with Paediatrician before use)

### **Angiotensin converting enzyme inhibitor**

Enalapril 0.1mg/kg/dose PO OD

**or**

Captopril 0.1mg/kg/dose PO TDS, gradually increase up to 1 mg/kg/dose TDS

Monitor blood pressure and potassium level (risk of hyperkalaemia)

#### Third line drugs

Digoxin (Discuss with Paediatrician/Cardiologist before use)

- Useful in heart failure with excessive tachycardia, supraventricular tachyarrhythmias.
- In practice: Don't "digitalize" with high-dose digoxin. Start with maintenance dose:
  - 2-17 years: 10mcg/kg/day OD (maximum 0.25 mg daily)

**IV inotropic agents** - Adrenaline, dobutamine (discuss with cardiologist)

**Beta blockers (Carvedilol)** - To be considered in acute heart failure with cardiogenic shock (discuss with cardiologist)

- **Pericardiocentesis** - May be necessary if:
  - Large pericardial effusion (>3cm on echo) and/or
  - Signs of cardiac tamponade (Tachycardia, elevated JVP, hypotension, pulsus paradoxus muffled heart sounds)
  - Pericardiocentesis should only be done in consultation with the cardiologist
- Nutritionist/dietician review
- Cardiology review and possible corrective surgery referral
- Palliative care team when necessary
  - Palliative care involvement early if in terminal chronic cardiac failure
  - Morphine is a venodilator reducing preload in children in cardiac failure and improves pulmonary oedema (Morphine dosing: 200-400 mcg/kg)

every 4 hours)

### **Follow up**

- Once diagnosis is confirmed, the child must be followed up in a PEN-Plus clinic regularly
- Ensure child has adequate stock of medications for home use at each visit
- Monitor for complications of medications
- Monitor growth and development
- Ongoing counselling and education



## Paediatric arrhythmias

### Atrioventricular blocks

#### Definition


An interruption or delay of electrical conduction from the atria to the ventricles leading to abnormally slow heart rate and irregular rhythm.

#### Criteria for bradycardia

Age Group	Heart Rate
Infants to < 3 years	< 100 bpm
Children 3-9 years	< 60 bpm
Children 9-16 years	< 50 bpm
Adolescents > 16years	< 40bpm


#### Types of conduction defects

1st degree - prolonged PR interval




2nd degree

- Mobitz type 1 (Wenckebach): progressive PR prolongation before
  - Dropped AV conduction
  - Generally not pathologic




PR	.18	.23	.26	.27	.18
Increment		.05	.03	.01	
RR		.57	.54	.53	.96

- Mobitz type 2: abrupt failure of AV conduction without prior PR prolongation



- 3rd degree (complete heart block): AV dissociation with no atrial impulses conducted to ventricles



**Note:** 2nd degree (Type 2 and above) and 3rd degree heart block are always pathological

### **Risk factors/causes**

- Hypoxia
- Sepsis
- Acidosis
- Hypothyroidism
- Electrolyte abnormalities i.e. hypokalaemia, hypocalcaemia
- Maternal Lupus (anti Ro/La positive); mother frequently asymptomatic
- Structural heart diseases: AVSD, congenital corrected transposition of great arteries (L-TGA), left atrial isomerism
- Long QT syndrome
- Surgical trauma: Especially in VSD closure, TOF repair, AVSD repair
- Infection: Rheumatic fever, endocarditis, viral myocarditis
- Drugs e.g. quinine

### **Prevention/promotion**

- Education and awareness of risk factors
- Screening of immediate family members of a person who suffered sudden cardiac death

### **Signs and symptoms**

- Most patients are asymptomatic
- Syncope
- Hypotension
- Exercise intolerance
- Fatigue
- Dyspnoea
- Dizziness
- Palpitations
- Cardiac arrest

### **Investigations**

- ECG
- Chest X Ray
- Cardiac Echo
- Urea and Electrolytes
- Other tests as clinically indicated

### **Differential diagnosis**

- See causes

**Management****Primary level**

See the secondary-level guidance below

**Secondary level**

- If unstable, manage ABC then refer to a tertiary centre
- Hemodynamically stable patients – Refer

**Tertiary level**

- Treat the underlying systemic causes of bradycardia.
- Hemodynamically stable patient with normal blood pressure:
  - No medical management for the bradycardia
  - Refer to a cardiology specialist
- Symptomatic bradycardia with haemodynamic instability
- Manage ABCDE
  - Cautious fluid resuscitation. Give 10mls/kg bolus if in shock or hypotensive with close monitoring for signs of fluid overload
- Medical management of bradycardia:
  - IV Atropine 0.02mg/kg/dose (maximum 0.6mg) - can repeat doses if remains bradycardic (maximum 3mg)
  - If atropine not working, add IV Adrenaline infusion at 0.02 mcg/kg/min and/or dopamine 2-10mcg/kg/min
- Cardiologist review

**Follow up**

- Once diagnosis is confirmed, the child must be followed up in a PEN-Plus clinic regularly
- Monitor growth and development
- Ongoing counselling and education

## Supraventricular tachycardia

### Definition

Abnormally rapid heart rhythm originating above the ventricles, often (but not always) with a narrow QRS complex

### Causes/risk factors

- Wolff-Parkinson-White (WPW)
- Congenital heart disease
- Pericarditis
- Idiopathic
- Infections (e.g. sepsis, pneumonia)

### Signs and symptoms/clinical features

- Severe tachycardia (> 220bpm)
- Heart palpitations
- Respiratory distress
- Shock

### Management

#### Primary level

- Manage ABCDE
- Consult/refer to secondary level

#### Secondary level

- Manage ABCDE
- Haemodynamically stable patients, do:
  - Vagal manoeuvres:
- Icepack/iced water for infants: apply to face for a maximum of 30 seconds.
- Valsalva manoeuvres if the child is old enough (blow into a pinched straw)
- Refer to tertiary hospital

#### Tertiary level

- **Haemodynamically stable patients**
  - Vagal manoeuvres:
    - Icepack/iced water for infants: apply to face for a maximum of 30 seconds.
    - Valsalva manoeuvres if child is old enough (blow into a pinched straw).

- For the performance of the following call the paediatric consultant or the cardiologist
  - IV Adenosine: 0.1mg/kg (maximum 6mg) rapid push into a big vein and follow with a flush. Increase by 0.1mg/kg every 2 minutes until tachycardia terminated or up to a maximum of 0.5mg/kg/dose (maximum: 18 mg total)
  - IV Amiodarone: 25mcg/kg/minute for 4 hours then 5 - 15mcg/kg/minute until conversion
  - Patient needs to be on a continuous cardiac monitoring in HDU/ICU
- **Haemodynamically unstable (in shock)**
  - Synchronized DC conversion at 0.5 to 1 joule/kg
  - Cardiologist review

### Follow up

- Once diagnosis is confirmed, the child must be followed up in a PEN-Plus clinic regularly if SVT persistent
- Ensure child has adequate stock of medications for home use at each visit
- Ongoing counselling and education

### Chest pain

A complaint of chest pain is frequently encountered in children. Although chest pain does not indicate serious disease of the heart or other systems in most paediatric patients, the clinician should be aware of the differential diagnosis for chest pain

#### **Causes/risk factors (non-cardiac causes):**

##### Psychogenic

- Life stressor (death in family, family discord, divorce, failure in school, nonacceptance from peers, sexual molestation)

##### Musculoskeletal e.g.

- Costochondritis,
- Trauma to chest wall
- Muscle strains

##### Abnormalities of the rib cage or thoracic spine

##### Respiratory e.g.

- Reactive airway disease
- Pneumonia
- Pleural irritation
- Pneumothorax or pneumomediastinum
- Foreign bodies in the airway

##### Gastrointestinal e.g.

- Gastroesophageal reflux
- Peptic ulcer disease
- Esophagitis
- Gastritis
- Foreign bodies (e.g. coins)
- Cholecystitis

#### **Causes/risk factors (cardiac causes):**

- Structural abnormalities of the heart
  - Severe aortic or pulmonary stenosis, hypertrophic obstructive cardiomyopathy, Eisenmenger's syndrome, MVP
- Coronary artery abnormalities
  - Previous Kawasaki disease, congenital anomaly, coronary heart disease, hypertension, sickle cell disease

- Inflammatory conditions
  - Pericarditis, Myocarditis, Kawasaki disease

### Health promotion

- Health education
- Early health seeking behaviour for persistent chest pain

### Investigations

The investigations should be done according to the suspected underlying cause. If cardiac cause is suspected do the following:

- Electrocardiogram
- Chest Xray
- Cardiac Echo (There is no need to do an echo if a cardiac cause is not suspected by history and physical examination)

### Differential diagnosis

- See causes

### Management

#### Primary level

#### History and physical examination

The initial history should be directed at determining the probable cause and should include the following:

- Nature of the pain, in terms of the duration, intensity, frequency, location, and points of radiation
- Physical examination should include a complete cardiovascular examination to rule out probable cardiac causes

Refer to secondary level:

- When history reveals that chest pain is triggered or worsened by physical activities
- When there are abnormal findings in the cardiac examination

#### Secondary level

See the tertiary-level guidance below



### **Tertiary level**

- History and physical examination as above.
- Refer to a cardiologist:
  - When history reveals that chest pain is triggered or worsened by physical activities
  - When there are abnormal findings in the cardiac examination or when abnormalities occur in the chest radiographs or ECG, cardiology referral is clearly indicated
  - When there is a positive family history for cardiomyopathy, long QT syndrome, sudden unexpected death, or other hereditary diseases commonly associated with cardiac abnormalities

### **Follow up**

- If cardiac cause, patient should be followed up in a PEN-Plus clinic.
- Medication use and side effects

## Syncope

### Definition

Syncope is a transient loss of consciousness and muscle tone that results from inadequate cerebral perfusion. Presyncope is the feeling that one is about to pass out but remains conscious with a transient loss of postural tone. It is usually less serious than syncope and is often a manifestation of a benign condition. Dizziness is the most common prodromal symptom of syncope

### Causes/risk factors

#### Autonomic

- Vasovagal syncope
- Orthostatic (postural) hypotension
- Situational syncope
- Postural orthostatic tachycardia syndrome (POTS)

#### Cardiac arrhythmias

- Tachycardia: SVT, atrial flutter or fibrillation, ventricular tachycardia
- Bradycardia: Sinus bradycardia, asystole, complete heart block

#### Cardiac obstructive lesions

- Outflow obstruction: Severe aortic/pulmonary stenosis, hypertrophic cardiomyopathy, pulmonary hypertension
- Inflow obstruction: Severe mitral stenosis, cardiac tamponade, constrictive pericarditis

#### Metabolic

- Dehydration
- Hypoglycaemia
- Electrolyte disorders

#### Pregnancy

### Investigations

- Investigate for underlying conditions.
- Measure blood pressure in the supine and upright position to check for postural hypotension and heart rate changes
- ECG
- Echocardiogram

### Differential diagnosis

- Neuropsychiatric disorders
  - Anxiety disorders: panic disorders, phobia
  - Hyperventilation
  - Seizure disorders
  - Brain tumours
  - Conversion disorders (malingering, hysteria etc)

### Management

<b>Primary level</b> Refer to tertiary level for cardiology/neurology review
<b>Secondary level</b> Refer to tertiary level for cardiology/neurology review
<b>Tertiary level</b> <ul style="list-style-type: none"><li>• Investigate for underlying cause and manage as appropriate</li><li>• Consult cardiologist</li></ul>

### Follow up

- If cardiac cause, patient should be followed up in PenPlus clinic

### References

Hussain Imam Hj Muhammad Ismail, Ng, H. P., & Thomas, T. (n.d.). *Paediatric protocols for Malaysian hospitals* (3rd ed.).

Szczygielska, I., Hernik, E., Kołodziejczyk, B., Gazda, A., Maślińska, M., & Gietka, P. (2018). Rheumatic fever: New diagnostic criteria. *Rheumatology*, 56(1), 37–41. <https://doi.org/10.1093/rheumatology/kex456>

Park, M. K., & Salamat, M. (2021). *Park's pediatric cardiology for practitioners* (7th ed.). Elsevier.

UpToDate. (2023). *Newborn screening for critical congenital heart disease using pulse oximetry*. <https://www.uptodate.com/contents/newborn-screening-for-critical-congenital-heart-disease-using-pulse-oximetry>

Dean, S. L., & Singer, H. S. (2017). Treatment of Sydenham's chorea: A review of the current evidence. *Tremor and Other Hyperkinetic Movements*, 7, 456. <https://doi.org/10.7916/D8W95GJ2>

Simonneau, G., Gatzoulis, M. A., Adatia, I., Celermajer, D., Denton, C., Ghofrani, A., Gomez Sanchez, M. A., Krishna Kumar, R., Landzberg, M., Machado, R. F., Olschewski, H., Robbins, I. M., & Souza, R. (2013). Updated clinical classification of pulmonary hypertension. *Journal of the American College of Cardiology*, 62(25 Suppl), D34–D41. <https://doi.org/10.1016/j.jacc.2013.10.029>

Beaton, A., & Carapetis, J. (2015). The 2015 revision of the Jones criteria for the diagnosis of acute rheumatic fever: Implications for practice in low-income and middle-income countries. *Heart Asia*, 7(2), 7–11. <https://doi.org/10.1136/heartasia-2015-010648>

