

A counterexample to Haglund’s monotonic-zeros conjecture for the Riemann Ξ approximants

Kunal Tyagi*

October 1, 2026

Abstract

Haglund introduced the partial sums $\Xi_N(z) = \sum_{n \leq N} \Phi_n(z)$ of Riemann’s expansion of the Ξ -function in incomplete gamma functions, and conjectured that for every N the zeros of Ξ_N in the closed first quadrant, listed by increasing real part, have nondecreasing imaginary parts. The conjecture implies the Riemann hypothesis, and Haglund reported computations indicating that a weaker form of it holds for $N \leq 10$. We show that the conjecture is false for $N = 27$: Ξ_{27} has a real zero in $(3144.8946, 3144.8947)$ and a non-real zero $z_0 = 3143.2206824215 \dots + 0.3152587993 \dots i$ in the first quadrant, whose real part is smaller. The weak form fails as well. Both facts are certified twice, independently, in rigorous interval arithmetic — once with Arb ball arithmetic, once with outward-rounded interval arithmetic and proved truncation bounds — and z_0 is enclosed in a square of half-width 10^{-48} . The same violation occurs at $N = 24$ for the companion approximants $\xi_N = \Xi_N + c_N$. The failure has a mechanism: on the critical line $\Xi_N < \Xi < \xi_N$, so the real zeros of Ξ_N lie in those positive lobes of Ξ that rise above a positive level, and a shallow lobe followed by a deeper one near the height $4(N+1)^2$, where the approximant departs from Ξ , leaves a non-real zero below real zeros. Two statements in Haglund’s paper are corrected: the number of real zeros of Ξ_N is odd, so his table entry 32 at $N = 4$ is inconsistent (a numerical census finds 31), and the $1/x^2$ coefficient of Ξ_N on the real axis is negative for every N . The result refutes a sufficient condition for the Riemann hypothesis and says nothing about the Riemann hypothesis itself: the non-real zero belongs to the approximant, at a height where the approximant no longer tracks Ξ , and the Riemann hypothesis has been verified far beyond that height.

1 Introduction

1.1 Haglund’s approximants and his conjecture

Write $s = \frac{1}{2} + iz$ and let

$$\Xi(z) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s) = \xi(s), \quad (1)$$

Riemann’s Ξ -function, which is even, entire, and real on the real axis; the Riemann hypothesis is the statement that all its zeros are real. Let $\mathcal{Q} = \{z : \operatorname{Re} z \geq 0, \operatorname{Im} z \geq 0\}$ be the closed first quadrant. Haglund [Hag09] starts from Riemann’s representation $\Xi(z) = \int_0^\infty \cos(zt)\varphi(t) dt$ with

*`hello.jay.tyagi@gmail.com`. *Use of AI.* The mathematics in this paper — the derivations, the computations, the verification suite, and the text — was produced by Claude (Anthropic) working under the author’s direction inside a structured research program; the author set the program’s objectives, adjudicated its decisions, and is responsible for the content. Claude is a tool and is not an author. The complete working record is public: <https://github.com/x67ai/riemann-rh-program>.

$\varphi = \sum_{n \geq 1} \varphi_n$, $\varphi_n(t) = \exp(-n^2 \pi e^{2t})(8\pi^2 n^4 e^{4.5t} - 12\pi n^2 e^{2.5t})$, and asks what happens when φ is replaced by the finite sum $\sum_{n \leq N} \varphi_n$. With the “hyperbolic gamma function”

$$G(z; a, b) = 4 \int_0^\infty \cos(2zu) e^{2bu - ae^{2u}} du = \frac{\Gamma(b + iz, a)}{a^{b+iz}} + \frac{\Gamma(b - iz, a)}{a^{b-iz}} \quad (a > 0, b \in \mathbf{R}), \quad (2)$$

where $\Gamma(w, a) = \int_a^\infty e^{-t} t^{w-1} dt$ is the upper incomplete gamma function, this gives Riemann’s series $\Xi(z) = \sum_{n \geq 1} \Phi_n(z)$ and Haglund’s Ξ -approximates

$$\Xi_N(z) := \sum_{n=1}^N \Phi_n(z), \quad \Phi_n(z) := 2\pi^2 n^4 G\left(\frac{z}{2}; n^2 \pi, \frac{9}{4}\right) - 3\pi n^2 G\left(\frac{z}{2}; n^2 \pi, \frac{5}{4}\right) \quad (3)$$

([Hag09, (1)–(3), (6), (10)–(14), pp. 1–3]; page and equation numbers refer to the arXiv version). Each Ξ_N is even, entire, and real on the real axis. Haglund defines: “We say that a given function $F(z)$ has monotonic zeros in a region D of the complex plane if, when we list the zeros of F in D by increasing real part, the imaginary parts of the zeros are monotone nondecreasing”, with the side assumption “(We assume F has at most one zero on the intersection of any vertical line with D).” [Hag09, p. 3]. He then states:

Conjecture 1 (Haglund [Hag09, p. 3]). “For $N \in \mathbf{N}$, $\Xi_N(z)$ has monotonic zeros in $\mathcal{Q}$.”

His Proposition 1 reads “Conjecture 1 implies the Riemann Hypothesis” [Hag09, pp. 3–4]; the proof is a Hurwitz-type argument, using that $\Xi_N \rightarrow \Xi$ uniformly on compacta and that Ξ has infinitely many real zeros. His Remark 1 records a weak form: “A weaker form of Conjecture 1, which still implies RH, is that there are no non-real zeros of $\Xi_N(z)$ in $\mathcal{Q}$ whose real part is less than the largest real zero of $\Xi_N(z)$ ”, and reports that “Computations along these lines indicate this weaker form of Conjecture 1 is true at least for $N \leq 10$ ” [Hag09, p. 4], the computations being floating-point ones for which “no attempt has been made to establish rigorous error bounds” [Hag09, p. 3]. The paper was published as [Hag11].

1.2 The result

Theorem 1.1. (a) Ξ_{27} has a real zero $x_1 \in (3144.8946, 3144.8947)$, and a second real zero $x_2 \in (3145.5998, 3145.5999)$.

(b) Ξ_{27} has exactly one zero z_0 , which is simple, in the open square of half-width $3.7 \cdot 10^{-11}$ about $c = 3143.2206824215 + 0.3152587994i$. More precisely, $z_0 \in z^* + [-10^{-48}, 10^{-48}]^2$ with

$$\begin{aligned} \operatorname{Re} z^* &= 3143.220682421536585287281295989417800119\dots, \\ \operatorname{Im} z^* &= 0.315258799378214845382286373009271765\dots, \end{aligned}$$

where the full explicit center z^* is given in Section 6.

(c) Consequently $z_0 \in \mathcal{Q}$ is non-real, with $\operatorname{Im} z_0 > 0.3152$ and $\operatorname{Re} z_0 < 3143.2207 < x_1$. Listed by increasing real part, the zeros of Ξ_{27} in $\mathcal{Q}$ do not have nondecreasing imaginary parts: Conjecture 1 is false for $N = 27$. Since z_0 is a non-real zero in $\mathcal{Q}$ whose real part is less than the real zero x_1 , and hence less than the largest real zero, the weak form of Remark 1 is false for $N = 27$ as well.

The pair (z_0, x_1) violates the conjecture under every enumeration of the zeros by increasing real part, so Haglund’s side assumption (at most one zero on each vertical line) plays no role. Statements (a) and (b) are established by two independent computer-assisted proofs in rigorous interval arithmetic, described in Section 6: one in Arb ball arithmetic [Joh17, FLINT], one in outward-rounded interval arithmetic [mpm] with every truncation bounded by a proved remainder estimate. Each was first run on a ladder of known cases (two zeros from Haglund’s table and a non-real zero of Ξ_1 from his appendix).

1.3 Scope

Theorem 1.1 refutes a sufficient condition for the Riemann hypothesis. It says nothing about the Riemann hypothesis, and nothing about the zeros of ζ . The non-real zero z_0 is a zero of the approximant Ξ_{27} , not of Ξ , and it lies at a height where Ξ_{27} no longer tracks Ξ : there the difference $\Xi - \Xi_{27}$ is of the same size as Ξ itself (Section 5). The Riemann hypothesis has been verified rigorously far above this height: Platt and Trudgian prove that “all zeroes $\beta + i\gamma$ of the Riemann zeta-function with $0 < \gamma \leq 3 \cdot 10^{12}$ have $\beta = 1/2$ ” [PT21]. Nor does the paper decide Conjecture 1 for $N \leq 26$: the scan that located the violation at $N = 27$ (Section 5) is numerical, covers a window near one height for each N , and is not a proof that smaller N satisfy the conjecture.

1.4 The mechanism, and two corrections to Haglund’s paper

The failure is not an accident of $N = 27$. On the critical line the approximants sit below Ξ by a positive amount: for every $N \geq 1$ and real t ,

$$\Xi_N(t) = \Xi(t) - Q_N(t), \quad Q_N(t) > 0$$

(Theorem 3.3). Hence $\Xi_N < 0$ wherever $\Xi \leq 0$, and the real zeros of Ξ_N lie in the *positive lobes* of Ξ (the intervals between consecutive zeros on which $\Xi > 0$): an even number in each positive lobe (γ_j, γ_{j+1}) and an odd number in the central lobe $(0, \gamma_1)$, where $0 < \gamma_1 < \gamma_2 < \dots$ are the positive zeros of Ξ . A positive lobe carries real zeros of Ξ_N only if Ξ rises above the level Q_N there. Near the height $4(N+1)^2$ the level Q_N becomes comparable to the size of Ξ , and the heights of consecutive lobes fluctuate; a lobe that stays below the level, followed by a lobe that rises above it, leaves a pair of non-real zeros below real zeros. At $N = 27$ the positive lobe of Ξ over $(3142.98, 3143.57)$ reaches only 0.411 of the level Q_{27} and the next one, over $(3144.49, 3146.28)$, reaches 1.325 of it: the pair $z_0, \bar{z}_0$ sits over the first, and the real zeros x_1, x_2 lie in the second (Section 5; these lobe figures are floating-point computations, not part of the proof).

The same sandwich corrects two statements in [Hag09].

- (i) The number of real zeros of Ξ_N on $(0, \infty)$, counted with multiplicity, is odd for every N (Corollary 3.5). Haglund’s table [Hag09, p. 4] lists 1, 7, 15, 32, 53, 79, 113, 155, 207, 263 real zeros for $N = 1, \dots, 10$; the entry 32 at $N = 4$ is inconsistent with this, and a numerical census of Ξ_4 finds 31, the largest being 103.3679880094135, which is Haglund’s printed value.
- (ii) On the real axis $x^2 \Xi_N(x)$ tends to a negative limit for every $N \geq 1$ (Proposition 3.4), whereas [Hag09, p. 10] states “Thus the coefficient of $1/x^2$ in $\Xi_N(x)$ is positive for $k \geq 3$ and negative for $1 \leq k < 3$ ” (his k is N).

1.5 The companion approximants, and why an ordering property

Haglund’s Ξ_N differs by a constant from the more obvious truncation of Riemann’s formula,

$$\xi_N(s) := \frac{1}{2} + \frac{1}{2}s(s-1) \sum_{n \leq N} g_n(s), \quad \Xi_N(z) = \xi_N\left(\frac{1}{2} + iz\right) - c_N, \quad c_N = \sum_{n > N} (4\pi n^2 - 1)e^{-\pi n^2} > 0$$

(Section 2). On the critical line ξ_N lies *above* Ξ (Theorem 3.1), so its real zeros lie in the negative lobes of Ξ , and the same ordering property fails for it at $N = 24$ (Theorem 7.1). Neither family can satisfy the Riemann hypothesis literally: no member of the family $\xi_w = \frac{1}{2} + \frac{1}{2}s(s-1) \sum_n w_n g_n$ with $0 \leq w_n \leq 1$ and $w \not\equiv 1$, or with w finitely supported, has all its zeros on the critical line; it has finitely many zeros there and infinitely many off it (Theorem 4.1). With polynomially bounded weights, having all zeros on the line forces $w \equiv 1$ (Theorem 4.2). This is why a conjecture about approximants of this kind has to be an ordering statement such as Conjecture 1, and not real-rootedness; and any approach through real-rootedness in a window that grows with N is a height filtration of the Riemann hypothesis, with nothing to prove beyond the Riemann hypothesis on the new window (Proposition 4.3).

1.6 Prior work

Haglund’s conjecture was stated in 2009 [Hag09, Hag11] and checked by him, in its weak form, for $N \leq 10$. Ahn’s thesis [Ahn12], written under Haglund’s supervision, computes zeros of the component functions of the approximants (incomplete gamma functions, hyperbolic gamma functions, and small linear combinations of the Φ_n) and restates the check: “Computations indicate this weaker form of Conjecture 1.6 is true at least for $N \leq 10$ ” [Ahn12, pp. 10–11]; it tests no Ξ_N beyond $N = 10$. Haglund’s Conjecture 4 concerns the zeros of the pencil $\Xi_k + t\Phi_{k+1}$, $0 \leq t \leq 1$ [Hag09, p. 11]; its first case $k = 1$ is the subject of a 2026 preprint of Baccaro [Bac26], whose record states that “The result covers only the first interpolation.” We found no test of Conjecture 1 beyond $N = 10$, and no refutation of it, in the literature we could reach; a null search is not a proof of absence. Lagarias and Montague study a related Fourier-integral family whose members have only finitely many zeros on the critical line; they ask whether monotone increase of the imaginary parts of the zeros in the first quadrant would be a sufficient condition for all zeros to be on the line, and point to Haglund’s paper as a related situation “involving the ξ -function, where such a monotonicity implies the RH” [LM11, p. 24]. Truncations that keep all but finitely many zeros *on* the line are known in another setting: Ki proved that for $y \geq 1$ all but finitely many complex zeros of any partial sum of the Chowla–Selberg formula are simple and on the line, and observed that the Riemann hypothesis would follow if the words “but finitely many” could be removed for infinitely many N in the case $z = i$ [Ki06, p. 198]. The approximants here are the opposite case (Theorem 4.1). Finally, Liu, Matiyasevich, Oesterlé and Zaharescu prove, for a different family of approximants to ξ built from truncated Euler products, that “The Bounded Riemann Hypothesis is equivalent to the Riemann Hypothesis plus the Simplicity Conjecture” [LMOZ24, Theorem 1.7]; Proposition 4.3 is the analogue for the approximants of this paper.

1.7 Organization

Section 2 recalls Riemann’s identity and relates the two families. Section 3 proves the positivity and sandwich theorems and the two corrections. Section 4 gives the structural theorems. Section 5 describes the departure height and the lobe scan. Section 6 is the certificate for Theorem 1.1, and Section 7 the companion violation at $N = 24$.

2 Riemann's identity and the two families

Let $\psi(x) = \sum_{n \geq 1} e^{-\pi n^2 x}$. Splitting $\int_0^\infty \psi(x) x^{s/2-1} dx$ at $x = 1$ and using $\psi(1/x) = \sqrt{x} \psi(x) + (\sqrt{x} - 1)/2$ gives Riemann's identity [Rie59, Edw74], valid for all $s \in \mathbf{C}$:

$$\Lambda(s) := \pi^{-s/2} \Gamma(s/2) \zeta(s) = -\frac{1}{s} - \frac{1}{1-s} + \sum_{n \geq 1} g_n(s), \quad (4)$$

where

$$g_n(s) = \frac{\Gamma(s/2, X)}{X^{s/2}} + \frac{\Gamma((1-s)/2, X)}{X^{(1-s)/2}}, \quad X = \pi n^2.$$

Hence $\xi(s) = \frac{1}{2}s(s-1)\Lambda(s) = \frac{1}{2} + \frac{1}{2}s(s-1) \sum_{n \geq 1} g_n(s)$. For weights $w: \mathbf{N} \rightarrow \mathbf{R}$ put

$$\xi_w(s) := \frac{1}{2} + \frac{1}{2}s(s-1) \sum_{n \geq 1} w_n g_n(s), \quad (5)$$

and write ξ_N for $w = \mathbf{1}_{[1, N]}$. Every ξ_w satisfies $\xi_w(1-s) = \xi_w(s)$ and $\xi_w(\bar{s}) = \overline{\xi_w(s)}$, so $\xi_w(\frac{1}{2} + it)$ is real for real t ; and $|g_n(s)| \leq g_n(\operatorname{Re} s)$, since $|\Gamma(a, X)| \leq \Gamma(\operatorname{Re} a, X)$. Throughout, $s = \frac{1}{2} + iz$; on the critical line $z = t$ is real, and we write $\Xi(t) = \xi(\frac{1}{2} + it)$ and $P_w(t) := \xi_w(\frac{1}{2} + it) - \Xi(t)$.

Lemma 2.1. *For every $n \geq 1$, $\Phi_n(z) = \frac{1}{2}s(s-1)g_n(s) + (4\pi n^2 - 1)e^{-\pi n^2}$. Consequently*

$$\Xi_N(z) = \xi_N(s) - c_N, \quad c_N := \sum_{n > N} (4\pi n^2 - 1)e^{-\pi n^2} > 0.$$

Proof. Put $h(w) = X^{-w} \Gamma(w, X)$ with $X = \pi n^2$. Since $iz = s - \frac{1}{2}$, one has $\frac{9}{4} + \frac{iz}{2} = 2 + \frac{s}{2}$ and $\frac{9}{4} - \frac{iz}{2} = 2 + \frac{1-s}{2}$, and the same with 1 in place of 2 for $b = \frac{5}{4}$; so definition (3) reads $\Phi_n = 2X^2[h(\frac{s}{2} + 2) + h(\frac{1-s}{2} + 2)] - 3X[h(\frac{s}{2} + 1) + h(\frac{1-s}{2} + 1)]$. The recurrence $\Gamma(w+1, X) = w\Gamma(w, X) + X^w e^{-X}$ gives $h(w+1) = (wh(w) + e^{-X})/X$, and applying it twice, $2X^2h(a+2) - 3Xh(a+1) = (2a^2 - a)h(a) + (2a - 1 + 2X)e^{-X}$. For $a = s/2$ and for $a = (1-s)/2$ one has $2a^2 - a = \frac{1}{2}s(s-1)$, while $2a - 1$ equals $s - 1$ and $-s$ respectively; adding the two gives the first claim. Summing over $n \leq N$ and using $\sum_{n \geq 1} (4\pi n^2 - 1)e^{-\pi n^2} = \frac{1}{2}$ (differentiate $\theta(1/x) = \sqrt{x} \theta(x)$, $\theta = 1 + 2\psi$, at $x = 1$) gives the second. $\square$

Numerically $c_1 = 1.7180566258547875025 \cdot 10^{-4}$, $c_2 = 5.891258854396012012 \cdot 10^{-11}$, $c_3 = 2.9589851955455663928 \cdot 10^{-20}$, $c_4 = 2.4342009049056017 \cdot 10^{-32}$ and $c_5 = 3.443519624082649 \cdot 10^{-47}$. So Haglund's zero sets are the level sets $\{\xi_N = c_N\}$ of the truncations ξ_N , not their zero sets; the two families have different zeros, and both are studied below.

3 Positivity on the critical line

Theorem 3.1 (positivity). *For every $n \geq 1$ and real t , $g_n(\frac{1}{2} + it) > 0$. Consequently, for weights $0 \leq w_n \leq 1$,*

$$P_w(t) = \frac{1}{2} \left(\frac{1}{4} + t^2 \right) \sum_{n \geq 1} (1 - w_n) g_n\left(\frac{1}{2} + it\right) \geq 0,$$

with equality for all t only if $w \equiv 1$. In particular $\xi_1(\frac{1}{2} + it) > \xi_2(\frac{1}{2} + it) > \dots > \Xi(t)$ for every real t .

Proof. With $x = e^{2u}$,

$$g_n(\tfrac{1}{2} + it) = \int_1^\infty e^{-Xx} 2x^{1/4} \cos(\tfrac{t}{2} \log x) \frac{dx}{x} = 4 \int_0^\infty k(u) \cos(tu) du, \quad k(u) = \exp(u/2 - Xe^{2u}).$$

With $y = Xe^{2u} \geq X \geq \pi$ one has $k' = k(\frac{1}{2} - 2y) < 0$ and $k'' = k(4y^2 - 6y + \frac{1}{4}) > 0$, since the larger root of $4y^2 - 6y + \frac{1}{4}$ is $(3 + 2\sqrt{2})/4 = 1.457\dots < \pi$. So k is positive, strictly decreasing and strictly convex on $[0, \infty)$, and tends to 0. For $t \neq 0$, $\int_0^\infty k(u) \cos(tu) du = t^{-1} \int_0^\infty (-k'(u)) \sin(tu) du$; since $-k'$ is positive and strictly decreasing, the pieces of the sine integral over consecutive half-periods alternate in sign with strictly decreasing sizes, and the first is positive, so the integral is positive (this is the argument behind Pólya's criterion). At $t = 0$ the integral is $\int_0^\infty k > 0$. The formula for P_w is Riemann's identity (4) minus (5), with $\frac{1}{2}s(s-1) = -\frac{1}{2}(\frac{1}{4} + t^2)$ on the line. $\square$

Corollary 3.2 (lobe law for ξ_w). *Let $0 \leq w \leq 1$, $w \neq 1$. Every zero of ξ_w on the critical line lies in an open negative lobe of Ξ (an interval between consecutive zeros of Ξ on which $\Xi < 0$), and each negative lobe contains an even number of them, counted with multiplicity.*

Proof. At the ends of a lobe $\xi_w = P_w > 0$, and on positive lobes $\xi_w \geq \Xi > 0$. $\square$

The real-zero counts of ξ_N observed numerically are 2, 6, 16, 32, 52, 80 for $N = 1, \dots, 6$, all even. Haglund's approximants lie on the other side of Ξ . For $n \geq 1$ put $k_n(u) = \exp(u/2 - \pi n^2 e^{2u})$ and

$$\tilde{\varphi}_n := k_n'' - \tfrac{1}{4}k_n = 2y(2y - 3)e^{u/2-y}, \quad y = \pi n^2 e^{2u},$$

which equals $(4\pi^2 n^4 e^{9u/2} - 6\pi n^2 e^{5u/2}) \exp(-\pi n^2 e^{2u})$, one half of the term $\varphi_n(u)$ of Riemann's kernel (Section 1.1).

Theorem 3.3 (sandwich). *For every $N \geq 1$ and real t ,*

$$\Xi_N(t) = \Xi(t) - Q_N(t) < \Xi(t) < \Xi(t) + P_N(t) = \xi_N(\tfrac{1}{2} + it),$$

where

$$Q_N(t) = c_N - P_N(t) = 2 \int_0^\infty \sum_{n>N} \tilde{\varphi}_n(u) \cos(tu) du > 0.$$

Proof. Integrating by parts twice,

$$\tfrac{1}{2}(\tfrac{1}{4} + t^2)g_n(\tfrac{1}{2} + it) = 2(\tfrac{1}{4} + t^2) \int_0^\infty k_n \cos(tu) du = -2k_n'(0) - 2 \int_0^\infty (k_n'' - \tfrac{1}{4}k_n) \cos(tu) du,$$

and $-2k_n'(0) = (4\pi n^2 - 1)e^{-\pi n^2}$; summing over $n > N$ gives $P_N = c_N - Q_N$, and $\Xi_N = \xi_N - c_N = \Xi + P_N - c_N = \Xi - Q_N$ by Lemma 2.1. For $n \geq 2$ one has $y \geq 4\pi$, and then $\tilde{\varphi}_n > 0$, $\tilde{\varphi}_n' = e^{u/2-y} y (-8y^2 + 30y - 15) < 0$ and $\tilde{\varphi}_n'' = e^{u/2-y} y (16y^3 - 112y^2 + 165y - 37.5) > 0$ (because $16y^2(y-7) > 0$ and $165y > 37.5$). So $\sum_{n>N} \tilde{\varphi}_n$ is positive, strictly decreasing and strictly convex on $[0, \infty)$ for $N \geq 1$, and the argument of Theorem 3.1 gives $Q_N(t) > 0$. $\square$

For $n = 1$ the kernel is not monotone at $u = 0$: the factor $-8y^2 + 30y - 15$ is $+0.291$ at $y = \pi$, so $\tilde{\varphi}_1'(0) = e^{-\pi} \cdot \pi \cdot 0.291 = +0.0395 > 0$. This is why the theorem needs $N \geq 1$.

Proposition 3.4 (the $1/t^2$ tail). *For every $N \geq 1$, with $X = \pi n^2$,*

$$\lim_{t \rightarrow \infty} t^2 \Xi_N(t) = 2 \sum_{n>N} \tilde{\varphi}_n'(0) = -2 \sum_{n>N} (8X^3 - 30X^2 + 15X) e^{-X} < 0.$$

Proof. $\Xi(t)$ decays exponentially, and two integrations by parts in the integral for $Q_N(t)$ in Theorem 3.3 give $t^2 Q_N(t) \rightarrow -2 \sum_{n>N} \tilde{\varphi}'_n(0)$. At $u=0$, $\tilde{\varphi}'_n(0) = -(8X^3 - 30X^2 + 15X)e^{-X}$, and each summand $8X^3 - 30X^2 + 15X$ is positive for $n \geq 2$. $\square$

Since the full kernel $\sum_{n \geq 1} \tilde{\varphi}_n$ is even, $\sum_{n \geq 1} \tilde{\varphi}'_n(0) = 0$, and the limit is also the sum over $n \leq N$ of the coefficients of $1/t^2$ in the individual Φ_n , namely $-2\tilde{\varphi}'_n(0)$; this is how Haglund computes it. Evaluated at 40 digits the limits are -0.078997 , $-1.6530 \cdot 10^{-7}$, $-2.7831 \cdot 10^{-16}$, $-5.7375 \cdot 10^{-28}$ and $-1.7062 \cdot 10^{-42}$ for $N = 1, \dots, 5$. Haglund lists the coefficients of $1/x^2$ in Φ_1 and Φ_2 as -0.01974938206 and $.01974934121$ [Hag09, (52), p. 10]; his values are one quarter of ours (at $N = 1$, -0.01974938206 against -0.078997), a constant factor that comes from the argument $z/2$ in (3) and does not affect signs. He concludes that the coefficient of $1/x^2$ in $\Xi_N(x)$ “is positive for $k \geq 3$ and negative for $1 \leq k < 3$ ” [Hag09, p. 10]. Proposition 3.4 shows that it is negative for every N : the coefficients of Φ_1 , Φ_2 and Φ_3 nearly cancel, and ten-digit values of them cannot resolve their sum, which is of order 10^{-16} , at $N = 3$. (Haglund uses the sign change to exhibit non-monotone zeros of the pencil $t\Phi_1 + \Phi_2$; that pencil is not one of the Ξ_N , and we do not revisit the example.)

Corollary 3.5 (real zeros of Ξ_N). *Let $N \geq 1$ and let $0 < \gamma_1 < \gamma_2 < \dots$ be the positive zeros of Ξ . Every real zero of Ξ_N in $(0, \infty)$ lies in a positive lobe of Ξ . Counted with multiplicity, each positive lobe (γ_j, γ_{j+1}) contains an even number of them and the central lobe $(0, \gamma_1)$ an odd number. Consequently the number of real zeros of Ξ_N in $(0, \infty)$, counted with multiplicity, is finite and odd.*

Proof. Where $\Xi \leq 0$, $\Xi_N = \Xi - Q_N < 0$ (Theorem 3.3); in particular $\Xi_N(\gamma_j) = -Q_N(\gamma_j) < 0$ at both ends of every positive lobe. At the origin $\Xi_N(0) = \Xi(0) - Q_N(0) \geq 0.4971 - c_1 > 0$, since $Q_N = c_N - P_N \leq c_N \leq c_1$, while $\Xi_N(\gamma_1) < 0$. By Proposition 3.4, $\Xi_N < 0$ on $[T_0, \infty)$ for some T_0 , so only finitely many lobes carry zeros, and the zeros of the entire function Ξ_N in $[0, T_0]$ are finite in number. $\square$

Haglund’s table [Hag09, p. 4] gives the numbers of real zeros of Ξ_N as 1, 7, 15, 32, 53, 79, 113, 155, 207, 263 for $N = 1, \dots, 10$. The entry 32 at $N = 4$ is even; by Corollary 3.5 there cannot be 32 simple real zeros. A numerical census of Ξ_4 (real zeros by sign changes, the total by the argument principle, not interval-rigorous) finds 31 real zeros, the largest being 103.3679880094135; Haglund prints the largest zero as 103.3679880094. The other nine entries are odd.

4 Structure: no member is real-rooted

Theorem 4.1. *Let $w \neq 0$, and suppose either (i) $0 \leq w_n \leq 1$ for all n , with $w_m < 1$ for some m , or (ii) w is finitely supported with arbitrary real values. Then ξ_w has only finitely many zeros on $\operatorname{Re} s = \frac{1}{2}$ and infinitely many zeros off it.*

Proof. (a) *Infinitely many zeros.* Since $|g_n(s)| \leq g_n(\operatorname{Re} s)$,

$$|\xi_w(s)| \leq \frac{1}{2} + \frac{1}{2}|s|^2 \sum_n |w_n| g_n(\operatorname{Re} s) = \exp(O(|s| \log |s|)).$$

With n_0 the least n such that $w_n \neq 0$, $\xi_w(\sigma) \sim \frac{1}{2}\sigma^2 w_{n_0} (\pi n_0^2)^{-\sigma/2} \Gamma(\sigma/2)$ as $\sigma \rightarrow +\infty$, since $\Gamma(\sigma/2, X)/\Gamma(\sigma/2) \rightarrow 1$ and the other terms are smaller by $(n_0/n)^\sigma$. So $\log |\xi_w(\sigma)| \geq (\sigma/2) \log \sigma (1 - o(1))$, ξ_w is not $e^{a+bs} P(s)$ with P a polynomial, and by Hadamard’s theorem it has infinitely many zeros.

(b) *Finitely many on the line.* Case (i): by Theorem 3.1, $\xi_w(\frac{1}{2} + it) \geq \Xi(t) + (1 - w_m)\frac{1}{2}(\frac{1}{4} + t^2)g_m(\frac{1}{2} + it)$. From $\gamma(a, X) = X^a e^{-X} \sum_{k \geq 0} X^k/(a)_{k+1}$ and $|(a)_{k+1}| \geq (|t|/2)^{k+1}$ on the line,

$\frac{1}{2}(\frac{1}{4} + t^2)g_m(\frac{1}{2} + it) \rightarrow (4\pi m^2 - 1)e^{-\pi m^2} > 0$ as $|t| \rightarrow \infty$ (the parts involving the complete $\Gamma(a)$ are $O(|t|^{7/4}e^{-\pi|t|^{1/4}})$), while $\Xi(t) \rightarrow 0$; so $\liminf_{|t| \rightarrow \infty} \xi_w(\frac{1}{2} + it) \geq (1 - w_m)(4\pi m^2 - 1)e^{-\pi m^2} > 0$. Case (ii): integrating by parts twice as in Theorem 3.3,

$$\xi_w(\frac{1}{2} + it) = C_w + 2 \int_0^\infty \Phi_w(u) \cos(tu) du, \quad \Phi_w = K_w'' - \frac{1}{4}K_w = \sum_n w_n \tilde{\varphi}_n,$$

where $K_w(u) = e^{u/2} \sum_n w_n e^{-\pi n^2 e^{2u}}$ and $C_w = \frac{1}{2} + 2K_w'(0)$, and the integral has the asymptotic expansion $\sum_{j \geq 0} 2(-1)^{j+1} \Phi_w^{(2j+1)}(0) t^{-2j-2}$. If every coefficient of $\xi_w(\frac{1}{2} + it) \sim C_w + \sum_j a_j t^{-2j-2}$ vanished, the odd part of the real-analytic function Φ_w would have zero Taylor series at 0, so Φ_w would be even. But the expansion of $\Phi_w(u)$ as $u \rightarrow -\infty$ has as coefficient of $e^{(5/2+2k)u}$ the number $-2\pi(2k+3)(-\pi)^k/k! \cdot \sum_n w_n n^{2k+2}$, while an even Φ_w decays double-exponentially as $u \rightarrow -\infty$. So $\sum_n w_n n^{2k+2} = 0$ for all $k \geq 0$, and since only finitely many n occur, $w \equiv 0$ (Vandermonde), which is excluded. Hence some coefficient is nonzero, and $\xi_w(\frac{1}{2} + it)$ has constant sign for $|t|$ large. $\square$

For ξ_N the leading coefficient is $C_w = c_N$. The same proof applies to Haglund's $\Xi_N = \xi_N - c_N$, whose leading coefficient is 0 and whose t^{-2} coefficient is negative (Proposition 3.4): no Ξ_N is real-rooted either, and Haglund's own list of the zeros of Ξ_1 [Hag09, p. 16] shows one real zero followed by non-real ones.

Theorem 4.2. *Let $|w_n| \leq Cn^A$ for some constants C, A , and $w \not\equiv 0$. If every zero of ξ_w lies on $\text{Re } s = \frac{1}{2}$, then $w \equiv 1$.*

Proof. Part (a) of the proof of Theorem 4.1 holds for polynomially bounded w , so ξ_w has infinitely many zeros, all on the line by hypothesis. The expansion of part (b) also holds for such w (Φ_w is real-analytic on $\mathbf{R}$ and all its derivatives decay double-exponentially on $[0, \infty)$), so all its coefficients vanish, $C_w = 0$, and Φ_w is even. Then $D(u) := K_w(u) - K_w(-u)$ is odd with $D'' = D/4$, so $D = 2\alpha \sinh(u/2)$, that is, $\psi_w(1/x) = \sqrt{x} \psi_w(x) - \alpha(\sqrt{x} - 1)$ for $\psi_w(x) = \sum_n w_n e^{-\pi n^2 x}$. Differentiating at $x = 1$ gives $K_w'(0) = \alpha/2$, and $C_w = 0$ gives $\alpha = -\frac{1}{2}$: ψ_w satisfies Riemann's theta relation exactly. Then $\delta := \psi_w - \psi = \sum_n (w_n - 1) e^{-\pi n^2 x}$ satisfies $\delta(1/x) = \sqrt{x} \delta(x)$ and decays super-exponentially at 0 and at ∞ , so its Mellin transform $\pi^{-s/2} \Gamma(s/2) \sum_n (w_n - 1) n^{-s}$ is entire, symmetric under $s \mapsto 1 - s$, and bounded in vertical strips. By Hamburger's theorem [Ham21] the Dirichlet series $f(s) = \sum_n (w_n - 1) n^{-s}$ is a constant multiple of ζ ; since f is entire, the constant is 0 and $w \equiv 1$. $\square$

For bounded w , Hamburger's theorem is used in the form stated by Nakamura [Nak20, Theorem D, p. 4]: if $F(s) = \sum a(n) n^{-s}$ converges absolutely for $\sigma > 1$, $P(s)F(s)$ is entire of finite order for some polynomial P , and $\xi_F(1 - s) = \xi_F(s)$ with $\xi_F(s) := \pi^{-s/2} \Gamma(s/2) F(s)$, then $F = C\zeta$; here $F = f$ is entire of finite order. For polynomially bounded w , Burnol's extension [Bur12, pp. 2–3, Corollaire 2] needs convergence only for $\text{Re } s$ large. Some such hypothesis is needed: with absolute convergence only in *some* half-plane, Knopp exhibits infinitely many linearly independent solutions of the functional equation (as reported in [Nak20, p. 4]).

Proposition 4.3 (windowed real-rootedness is a height filtration). *Let F_N be entire with $F_N(1 - s) = F_N(s)$ and $F_N(\bar{s}) = \overline{F_N(s)}$. Suppose that for each T there is $N_0(T)$ such that for $N \geq N_0(T)$: (i) $\sup_{R_T} |F_N - \xi| < \min_{\partial D} |\xi|$ for a finite family D of small discs about the zeros of ξ in $R_T = [-1, 2] \times [-T, T]$ and for ∂R_T ; and (ii) F_N has no zeros with $|\text{Im } s| \leq T$ outside R_T . Then, by Rouché's theorem on each disc and on R_T minus the discs, the zeros of F_N in R_T are those of ξ , moved by less than the disc radii, with multiplicity, and a zero of multiplicity one near the line lies on it by the symmetry $s \mapsto 1 - \bar{s}$. Hence, if the zeros of ξ in R_T are simple, F_N is real-rooted on R_T for all large N if and only if the Riemann hypothesis holds on R_T .*

For $F_N = \xi_N$ one has $\sup_{R_T} |\xi_N - \xi| \leq \frac{1}{2}(T^2 + 4) \max_{-1 \leq \sigma \leq 2} \sum_{n > N} g_n(\sigma) \approx T^2 e^{-\pi(N+1)^2} / (\pi(N+1)^2)$, which is below the size of ξ on the line up to $T \approx 4(N+1)^2$; and (ii) holds for all large N , by comparing ξ_N with ξ for $2 \leq \operatorname{Re} s \leq \pi(N+1)^2$, with the main term $\frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2) \sum_{n \leq N} n^{-s}$ for $\operatorname{Re} s \geq \pi(N+1)^2$, and using $s \mapsto 1-s$ on the left. So in any induction “members real-rooted up to T_N implies real-rooted up to T_{N+1} ”, the only content is the Riemann hypothesis on the new window $T_N < |\operatorname{Im} s| \leq T_{N+1}$; what the family proves by itself (Theorem 3.1: real zeros are never lost as N grows) is the easy half. This is the frame in which Conjecture 1 lives: it asks for an ordering property to carry the induction, and Theorem 1.1 shows that the ordering property fails near the height at which the approximant leaves the window.

5 Departure height and the lobe scan

The figures in this section are floating-point computations in mpmath at 30 to 80 digits, cross-checked by a second formula or a second contour where zeros are reported; they explain where to look and are not part of the proof of Theorem 1.1.

Departure. On the line, $\Xi(t)$ is of size roughly $e^{-\pi t/4}$, and the corrections P_N and Q_N are of size roughly $e^{-\pi(N+1)^2}$. The two balance near $t = 4(N+1)^2$, the *departure height*: below it the approximant tracks Ξ , above it the approximant is dominated by its correction. For the truncations ξ_N this is visible in a complete census of the zeros in $0 < \operatorname{Im} s \leq 200$: the first negative lobe of Ξ that carries no real zero of ξ_N is (25.01, 30.42), (40.92, 43.33), (69.55, 72.07), (107.17, 111.03), (150.05, 150.93), (202.49, 204.19), (265.56, 266.61) for $N = 1, \dots, 7$, with ratio of lobe depth to P_N equal to 0.0070, 0.20, 0.32, 0.24, 0.12, 0.27, 0.020: the departure height is $4(N+1)^2 + 6$ to $4(N+1)^2 + 10$. Haglund’s table shows the same law: his largest real zeros 14.05, 39.53, 65.03, 103.37, $\dots$, 489.39 for $N = 1, 2, 3, 4, \dots, 10$ lie close to $4(N+1)^2$. Every off-line zero of ξ_N found below height 200 lies outside the critical strip, at $\operatorname{Re} s \geq 1.39$.

The ordering property fails near departure. By Theorem 3.3, the real zeros of Ξ_N come in pairs from the positive lobes of Ξ whose height exceeds the smooth level Q_N , and by Corollary 3.2 those of ξ_N come from the negative lobes deeper than P_N . A lobe that fails the level, followed by a lobe that passes it, leaves a pair of non-real zeros over the first lobe with real zeros above it, which violates the ordering. Lobe heights decrease on average by a factor $e^{-\pi^2/\log(t/2\pi)}$ per step between lobes of the same sign, but fluctuate by comparable factors, so such an inversion is to be expected near the departure height for some N . We scanned the windows $[4(N+1)^2 - 40, 4(N+1)^2 + 90]$ for $N = 6, \dots, 29$ (and $N = 30$), counting zeros in every window and comparing each lobe’s height with the level. The only violations are at $N = 24$ for ξ_N and at $N = 27$ for Ξ_N . The closest other approach for Haglund’s family is at $N = 21$, where a lobe beyond the first failing one reaches 0.93 of the level (a violation needs more than 1); the next closest is $N = 20$, at 0.48. The scan is not a proof that Conjecture 1 holds for $N \leq 26$.

The violation at $N = 27$. The positive lobe (3142.97945673, 3143.56749918) of Ξ has height 0.411 times Q_{27} and carries no real zero of Ξ_{27} ; the next positive lobe, (3144.49048815, 3146.28352838), has height 1.325 times Q_{27} and carries the real zeros 3144.8946622186 and 3145.5998495765. For $\operatorname{Im} s \in [3142.95, 3144.0]$ and $\operatorname{Re} s \in [-\frac{1}{2}, \frac{3}{2}]$ the argument principle counts 2.0 zeros at 30 and at 45 digits, none of them on the line: $s = 0.8152587993782148453823 + 3143.220682421536585287i$, inside the critical strip, and its mirror $1 - \bar{s}$. In Haglund’s variable $z = -i(s - \frac{1}{2})$ the mirror is the zero z_0 of Theorem 1.1, whose real part is below both real zeros.

6 The certificate

Theorem 1.1 was certified twice, by two computations written independently, neither with access to the other: *certificate A* in Arb ball arithmetic [Joh17] through python-flint 0.6.0 (FLINT 3.0.1) [FLINT], and *certificate B* in the outward-rounded interval arithmetic of mpmath 1.3.0 [mpm], with no use of Arb. Both evaluate the objects (2)–(3) exactly as Haglund defines them. Each was first run on a ladder of known cases (Table 3).

6.1 Certificate A: Arb ball arithmetic

Certificate A evaluates Ξ_{27} by two routes. *Route L* is the literal sum (3), with $\Gamma(w, a)$ from FLINT’s `acb_hypgeom_gamma_upper`. At $x = 3144.8946$ its terms fall from $|\Phi_1| \approx |\Phi_2| \approx 8.0 \cdot 10^{-9}$ to $|\Phi_{27}| \approx 1.5 \cdot 10^{-991}$ while the sum is about 10^{-1070} , so about 1062 digits cancel; route L is used only at exact points, at 4800 and 6400 bits, where the cancellation is fully resolved. *Route T* writes $\Xi_{27} = \Xi - \sum_{n=28}^{31} \Phi_n - R_{31}$, with Ξ from (1) (FLINT’s ζ and Γ), the four terms as in route L, and the remainder bounded by Lemma 6.3; near z_0 all these terms are about 10^{-1066} or smaller ($\Phi_{28} \approx \Xi \approx 1.49 \cdot 10^{-1066}$, $\Phi_{29} \approx 3 \cdot 10^{-1144}$), so route T has no catastrophic cancellation and is sharp on ball inputs at 256 bits. Route T uses Riemann’s series $\Xi = \sum_{n \geq 1} \Phi_n$ [Hag09, (12), p. 3]; route L does not.

Lemma 6.1. *For $w \in \mathbf{C}$, $a > 0$, $b \in \mathbf{R}$: $|G(w; a, b)| \leq 2\Gamma(\beta, a)/a^\beta$ with $\beta = b + |\operatorname{Im} w|$.*

Proof. For $u \geq 0$, $|\cos(x + iy)|^2 = \cos^2 x + \sinh^2 y \leq \cosh^2 y$, so $|\cos(2wu)| \leq e^{2u|\operatorname{Im} w|}$ and $|G| \leq 4 \int_0^\infty e^{2\beta u - ae^{2u}} du$; the substitution $t = ae^{2u}$ turns this into $2a^{-\beta} \int_a^\infty t^{\beta-1} e^{-t} dt$. $\square$

Lemma 6.2. *For $\beta \geq 1$ and $a > \beta - 1$: $\Gamma(\beta, a)/a^\beta \leq e^{-a}/(a - \beta + 1)$.*

Proof. With $t = ax$, $\Gamma(\beta, a)/a^\beta = \int_1^\infty x^{\beta-1} e^{-ax} dx$, and $x^{\beta-1} \leq e^{(\beta-1)(x-1)}$ for $x \geq 1$ since $\log x \leq x - 1$ and $\beta \geq 1$; integrate. $\square$

Lemma 6.3. *Let $Y \geq |\operatorname{Im} z|$, $\beta_9 = \frac{9}{4} + Y/2$, $a_n = \pi n^2$, and $M \geq 1$ with $a_{M+1} - \beta_9 + 1 > 0$. Put $U_n = 2e^{-a_n}(2\pi^2 n^4 + 3\pi n^2)/(a_n - \beta_9 + 1)$. Then $|\sum_{n>M} \Phi_n(z)| \leq \sum_{n>M} U_n \leq 2U_{M+1}$.*

Proof. In (3), $|\operatorname{Im}(z/2)| \leq Y/2$, and Lemmas 6.1 and 6.2 (with $\beta = \frac{9}{4} + Y/2$ and $\frac{5}{4} + Y/2$, both ≥ 1) give $|\Phi_n| \leq U_n$. For $n \geq 1$, $U_{n+1}/U_n \leq e^{-3\pi} \cdot 16 < 1.3 \cdot 10^{-3}$, since the ratio of the polynomial factors is at most $((n+1)/n)^4 \leq 16$ and the ratio of the denominators is below 1; so $\sum_{n>M} U_n \leq 2U_{M+1}$. $\square$

At $N = 27$, $M = 31$ and $Y \leq 0.317$, and $2U_{32} \leq 2 \cdot 10^{-1393}$. The winding numbers use the elementary fact that if a path lies in a closed axis-parallel box that excludes 0, the box lies in an open half-plane through 0 and the continuous change of argument along the path equals the principal value $\operatorname{Arg}(\gamma(1)/\gamma(0)) \in (-\pi, \pi)$. The boundary of each square is cut into pieces, a piece is bisected until its ball enclosure excludes 0, and the increments are summed; by the argument principle a total in $(2\pi(k - \frac{1}{2}), 2\pi(k + \frac{1}{2}))$ certifies k zeros inside. Every arithmetic operation and special function is Arb’s, whose contract is that the output ball contains the exact value for every input in the input balls; no floating-point value is used in any decision. The only other inputs are Lemmas 6.1–6.3, the half-plane fact, Riemann’s series for route T, the intermediate value theorem and the argument principle.

6.2 Certificate B: interval arithmetic with proved remainders

Certificate B uses mpmath's real intervals with outward rounding, complex numbers as rectangular boxes built on them, and exact rational Bernoulli numbers; every truncation adds a disk whose radius is a proved bound, each targeting 2^{-190} at a working precision of 200 bits. It writes $\Phi_n = 2X^2[h(\frac{s}{2}+2)+h(\frac{1-s}{2}+2)]-3X[h(\frac{s}{2}+1)+h(\frac{1-s}{2}+1)]$ with $h(w) = X^{-w}\Gamma(w, X) = \int_1^\infty e^{-Xu}u^{w-1}du$, as in the proof of Lemma 2.1, and evaluates $\Xi_{27} = \Xi - \Phi_{28} - \Phi_{29} - \Phi_{30} - T_{30}$ with the following bounds.

- Γ by Stirling's series with the remainder bound of [DLMF, 5.11.1, §5.11(ii)], shifting the argument when $|w| < 300$ or $\operatorname{Re} w < 0$.
- ζ by Euler–Maclaurin summation [DLMF, 2.10.1] with the remainder bound $|R_m| \leq (2 - 2^{1-2m})|B_{2m}| |(s)_{2m}| N^{1-\sigma-2m}/((2m)!(\sigma+2m-1))$ from [DLMF, 24.9.2]; at this height $N = 1012$ and $m = 94$.
- h by repeated integration by parts, $h(w) = e^{-X} \sum_{k < K} \tau_k + \rho_K$ with $\tau_k = (w-1) \cdots (w-k)/X^{k+1}$ and $|\rho_K| \leq e^{-X} |\tau_K| X/(X - \max(\beta, 0))$, $\beta = \operatorname{Re} w - 1 - K$.
- The tail: $|h(w)| \leq e^{-X}/(X - \beta')$ with $\beta' = \max(\operatorname{Re} w - 1, 0)$, so for $X \geq \max(2\beta', 12)$ (with β' maximized over the four arguments in Φ_n) one has $|\Phi_n| \leq 9Xe^{-X}$, whence $|\sum_{n > M} \Phi_n| \leq 18\pi(M+1)^2 e^{-\pi(M+1)^2}$; at $N = 27$, $M = 30$ and the bound is $3.7 \cdot 10^{-1307}$.

Near $x = 3144.8946$ one has $|\Xi| = 1.4923145335789 \cdot 10^{-1066}$ and $|\Phi_{28}| = 1.49249055304208 \cdot 10^{-1066}$, so this route resolves $\Xi_{27} \approx 10^{-1070}$ with about 50 digits to spare. As a second rigorous route for the real zeros, certificate B also evaluates the literal sum (3) at 4400 bits, with $\Phi_1, \dots, \Phi_{27}$ from the lower incomplete gamma series (remainder $\leq e^{-X} X^K/|(w)_K|$ once $\operatorname{Re} w + K \geq X + 1$), integration by parts where it converges, and Stirling's series: no ζ , no Euler–Maclaurin summation, and no use of Riemann's series.

Winding numbers come from a rigorous Taylor model. Since $f = \Xi_{27}$ is entire, its Taylor coefficients about c satisfy $|a_k| \leq M_R/R^k$, with M_R an upper bound for $|f|$ on $|z - c| = R$ obtained from box enclosures covering the circle. The discrete Fourier transform of f on m points of the circle $|z - c| = \rho$ gives $D_k = \sum_{l \geq 0} a_{k+lm} \rho^{lm}$, so $|D_k - a_k| \leq (M_R/R^k) q^m / (1 - q^m)$ with $q = \rho/R$, and for $|z - c| \leq r' < R$, with $x = r'/R$,

$$\left| f(z) - \sum_{k < K} D_k (z - c)^k \right| \leq E := M_R \left[\frac{q^m}{1 - q^m} \sum_{k < K} x^k + \frac{x^K}{1 - x} \right].$$

A boundary piece is accepted when the model's box enclosure plus $[-E, E]^2$ excludes 0 and the ratio of the end values has positive real part; the change of argument along it is then the principal value of that ratio's argument. At $N = 27$: $R = 0.1$, $M_R = 2.09255 \cdot 10^{-1065}$, $\rho = 0.002$, $m = 24$, $K = 12$, giving $E = 3.51 \cdot 10^{-1106}$ for $r \leq 10^{-5}$ and $1.36 \cdot 10^{-1087}$ at $r = 10^{-3}$ (and D_0 agrees with the direct enclosure of $\Xi_{27}(c)$ to all digits shown). For the 10^{-48} enclosure a second model is built at the center

$$\operatorname{Re} z^* = 3143.220682421536585287281295989417800119257644961983569,$$

$$\operatorname{Im} z^* = 0.3152587993782148453822863730092717652956513976685290909,$$

with $\rho = 0.001$ and $m = 40$, giving $E = 2.09 \cdot 10^{-1145}$; at $r = 10^{-48}$ the smallest lower modulus over the boundary pieces, $5.9 \cdot 10^{-1114}$, still exceeds the radii of the point enclosures, about 10^{-1120} , by a factor 10^6 . The trust base of certificate B is mpmath's primitive interval operations and the cited DLMF remainder bounds; Riemann's series is used for the tail route and checked numerically, not re-derived.

6.3 The enclosures

Tables 1–3 give the certified values. Here $c = 3143.2206824215 + 0.3152587994i$, and a square of half-width r about a center w is $w + [-r, r]^2$.

x	certificate A (ball midpoint $\pm$ radius)	certificate B (interval)
3144.8946	$-1.76019463127752 \cdot 10^{-1070} \pm 4.1 \cdot 10^{-1085}$	$-1.760194631277516 \cdot 10^{-1070}$
3144.8947	$+1.06871649226171 \cdot 10^{-1070} \pm 3.3 \cdot 10^{-1085}$	$+1.068716492261707 \cdot 10^{-1070}$
3145.5998	$+1.12975694291664 \cdot 10^{-1070} \pm 1.9 \cdot 10^{-1085}$	$+1.129756942916642 \cdot 10^{-1070}$
3145.5999	$-1.14913915362778 \cdot 10^{-1070} \pm 2.9 \cdot 10^{-1085}$	$-1.149139153627783 \cdot 10^{-1070}$

Table 1: Enclosures of $\Xi_{27}(x)$ at the ends of the two real-zero intervals. Certificate A: balls as printed, display rounding included; the Arb radii are $3.3 \cdot 10^{-1435}$ (route L, 4800 bits), $6.0 \cdot 10^{-1915}$ (route L, 6400 bits) and $5.0 \cdot 10^{-1138}$ (route T, 256 bits), and all four evaluations (route T at 256 and 1024 bits, route L at 4800 and 6400 bits) give the same signs. Certificate B: tail route, box radii at most $6 \cdot 10^{-1121}$; its literal route at 4400 bits (radius about $1\text{--}2 \cdot 10^{-1171}$) agrees to all digits shown and overlaps.

cert.	center	half-width r	boundary pieces	k
A	c	10^{-3} (256 and 512 bits)	512, 512	1
A	c	$10^{-6}, 10^{-10}, 4 \cdot 10^{-11}$	512, 663, 807	1
A	$c + 2.5 \cdot 10^{-3}$	10^{-3}	269	0
A	$c + 1.5 \cdot 10^{-10}$	10^{-10}	714	0
B	c	$10^{-3}, 10^{-5}, 10^{-7}, 10^{-9}$	32 each	1
B	c	$10^{-10}, 5 \cdot 10^{-11}, 4 \cdot 10^{-11}$	32, 32, 33	1
B	c	$3.7 \cdot 10^{-11}$	36	1
B	c	$3.5 \cdot 10^{-11}$	33	0
B	$c + 0.01$	10^{-3}	32	0
B	z^*	$10^{-30}, 10^{-40}, 10^{-45}, 10^{-48}$		1

Table 2: Certified winding numbers k of Ξ_{27} along the boundaries of squares. In certificate A every run with $k = 1$ has total change of argument $6.28318530717959 \pm 4 \cdot 10^{-15}$. In certificate B at $r = 10^{-3}$ the smallest lower modulus over the pieces is $5.89 \cdot 10^{-1069}$ against the model error $E = 1.36 \cdot 10^{-1087}$; at $r = 3.7 \cdot 10^{-11}$ it is $1.97 \cdot 10^{-1078}$ against $E = 3.51 \cdot 10^{-1106}$. The run at $r = 3.5 \cdot 10^{-11}$ returns 0 because the zero lies just outside that square, and the rows with centers $c + 2.5 \cdot 10^{-3}$, $c + 1.5 \cdot 10^{-10}$ and $c + 0.01$ are zero-free controls.

Cross-checks. In certificate A, routes L and T overlap at all 18 points compared (the center, the reference zero, the corners and midpoints of the squares of half-width 10^{-3} , the corners of the square of half-width $4 \cdot 10^{-11}$, and the four points of Table 1), with $|L - T| \leq 1.2 \cdot 10^{-1288}$ at 1024 bits. In certificate B the literal and tail routes overlap at the four points of Table 1, its enclosures of Ξ contain mpmath’s ordinary 400-bit values, and its two methods for h at $X = 784\pi$ overlap. The two certificates agree with each other to every digit printed. A third, separately written evaluation of the literal sum (3) in Arb at 4800 bits gives $\Xi_{27}(3144.8946) \in [-1.76019463128 \cdot 10^{-1070} \pm 2.49 \cdot 10^{-1082}]$ and $\Xi_{27}(3144.8947) \in [1.06871649226 \cdot 10^{-1070} \pm 1.71 \cdot 10^{-1082}]$, the same signs at 3145.5998 and 3145.5999, and Haglund’s two ladder zeros; at the point $3143.220682421536585287 + 0.3152587993782148453823i$ it gives a value of modulus about $1.7 \cdot 10^{-1084}$, against about $6.1 \cdot 10^{-1072}$.

test	certificate A	certificate B
Ξ_1 at 14.04543957 and 14.04543959 (Haglund: 14.0454395788)	+1.347060456147 $\cdot 10^{-11}$, -1.704102125315 $\cdot 10^{-11}$	+1.347060456146727 $\cdot 10^{-11}$, -1.704102125314551 $\cdot 10^{-11}$
Ξ_2 at 39.5324810797 and 39.5324810799 (Haglund: 39.5324810798)	+3.605517254360 $\cdot 10^{-21}$, -4.593418771015 $\cdot 10^{-21}$	+3.605517254360067 $\cdot 10^{-21}$, -4.5934187710153 $\cdot 10^{-21}$
Ξ_1 about Haglund's zero 20.62534600592171760132974 + 2.697151842339519632505712 i zero-free controls for Ξ_1	$k = 1$ at $r = 10^{-10}$ (1516 pieces) $k = 0$ (center shifted by 0.5, $r = 0.1$; shifted by $3 \cdot 10^{-10}$, $r = 10^{-10}$)	$k = 1$ at $r = 10^{-3}$ and 10^{-8} $k = 0$ ($17 + i$, $r = \frac{1}{4}$)

Table 3: The ladder, run before $N = 27$ with the same code: the largest real zeros of Ξ_1 and Ξ_2 from Haglund's table [Hag09, p. 4], the first non-real zero of Ξ_1 from his appendix [Hag09, p. 16], and zero-free squares.

at a point 10^{-6} to the right. Independent re-runs of certificate A's real-zero computation and of certificate B's ladder reproduce their outputs.

6.4 Proof of Theorem 1.1

(a) By (2), $G(z; a, b)$ is real for real z , so Ξ_{27} is real and continuous on $\mathbf{R}$. The enclosures of Table 1 have strictly opposite signs at the ends of each interval, in both certificates; by the intermediate value theorem each interval contains a zero.

(b) Ξ_{27} is entire, being a finite sum of entire functions. Certificate B finds winding number 1 along the boundary of the square of half-width $3.7 \cdot 10^{-11}$ about c , with every boundary enclosure excluding 0; by the argument principle the open square contains exactly one zero z_0 , counted with multiplicity, so z_0 is simple. Certificate A independently finds winding number 1 at half-width $4 \cdot 10^{-11}$. The square $z^* + [-10^{-48}, 10^{-48}]^2$ lies inside the first square, since $z^* - c = 3.66 \dots \cdot 10^{-11} - 2.18 \dots \cdot 10^{-11} i$; winding number 1 there locates the same zero z_0 .

(c) From (b), $\text{Im } z_0 \geq 0.3152587994 - 3.7 \cdot 10^{-11} > 0.3152$ and $\text{Re } z_0 \leq 3143.2206824215 + 3.7 \cdot 10^{-11} < 3143.2207 < 3144.8946 < x_1$. So $z_0 \in \mathcal{Q}$ is non-real and precedes the real zero x_1 in any listing of the zeros of Ξ_{27} in $\mathcal{Q}$ by increasing real part, while $\text{Im } z_0 > 0 = \text{Im } x_1$. $\square$

7 The companion violation at $N = 24$

For the truncations ξ_N the ordering property reads: the zeros of $z \mapsto \xi_N(\frac{1}{2} + iz)$ in $\mathcal{Q}$, listed by increasing real part, have nondecreasing imaginary parts. Since $\xi_N \rightarrow \xi$ locally uniformly, the Hurwitz argument of Haglund's Proposition 1 shows that this property for every N would also imply the Riemann hypothesis. It fails at $N = 24$.

Theorem 7.1. (a) *The real function $t \mapsto \xi_{24}(\frac{1}{2} + it)$ changes sign on $(2510.2026, 2510.2027)$ and on $(2510.7086, 2510.7087)$, so ξ_{24} has zeros on the critical line at heights in these intervals.*

(b) *The function $z \mapsto \xi_{24}(\frac{1}{2} + iz)$ has exactly one zero in the square of half-width 10^{-10} about $2508.2839748053 + 0.3159896243 i$.*

(c) Hence ξ_{24} has a zero ρ_{24} off the critical line, inside the critical strip, with $\text{Im } \rho_{24} < 2508.2839748054$, below zeros on the line: the ordering property fails for ξ_{24} .

Both certificates prove this, with the same methods as in Section 6; here route T and certificate B use Riemann's identity (4), that is, Jacobi's theta relation, which is cited rather than re-proved. The enclosures on the line are, in certificate A (both routes; route L is the literal sum at 4800 bits, cancelling about 855 digits)

$$\xi_{24}\left(\frac{1}{2} + it\right) \in \begin{cases} [+2.81317536164 \cdot 10^{-854} \pm 5 \cdot 10^{-866}], & t = 2510.2026, \\ [-1.65267129215 \cdot 10^{-854} \pm 3 \cdot 10^{-866}], & t = 2510.2027, \\ [-3.23281139505 \cdot 10^{-854} \pm 4 \cdot 10^{-866}], & t = 2510.7086, \\ [+6.14184061542 \cdot 10^{-855} \pm 4 \cdot 10^{-867}], & t = 2510.7087, \end{cases}$$

and in certificate B $+2.813175361644763 \cdot 10^{-854}$, $-1.65267129214774 \cdot 10^{-854}$, $-3.232811395053649 \cdot 10^{-854}$ and $+6.141840615423139 \cdot 10^{-855}$ at the same four points. Certificate A finds winding number 1 along the s -plane squares of half-width 10^{-3} (151 pieces) and 10^{-10} (174 pieces) about $0.8159896243 + 2508.2839748053i$, and 0 for a control square shifted by $2.5 \cdot 10^{-3}i$; certificate B finds winding number 1 for $z \mapsto \xi_{24}(\frac{1}{2} + iz)$ on squares of half-width 10^{-3} , 10^{-6} , 10^{-9} and 10^{-10} about $2508.2839748053 + 0.3159896243i$ (Taylor model with $R = 0.1$, $\rho = 0.002$, $m = 24$, $K = 12$, $M_R = 4.07 \cdot 10^{-849}$; at $r = 10^{-3}$ the smallest lower modulus is $1.16 \cdot 10^{-852}$ against $E = 2.6 \cdot 10^{-871}$), and 0 for the control square shifted by 0.01. Numerically, at 50 digits, $\rho_{24} = 0.8159896243043424688941 + 2508.283974805324153202i$, and the zero in (b) is its mirror $1 - \bar{\rho}_{24}$.

The mechanism is the one of Section 5 with the signs reversed: the negative lobe (2507.87103958, 2508.7660461) of Ξ has depth 0.595 times P_{24} and carries no real zero of ξ_{24} , while the next negative lobe (2509.65299744, 2511.66018979) has depth 1.130 times P_{24} . The off-line zero lies inside the critical strip, at $\text{Re } s = 0.8160$. As with Ξ_{27} , it is a zero of the approximant at a height where the approximant has departed from ξ ; it says nothing about the zeros of ζ .

Provenance, data and code

Method. The two certificates of Section 6 were written independently from a common written statement of what was to be certified, and each was run on the ladder of Table 3 before $N = 27$. The proofs of Sections 2–4 were re-derived line by line by an independent reader, and the corrections found there are incorporated.

Data and code availability. Every number in this paper comes from the two certificates, from the independent evaluations that checked them, or from the floating-point records behind Sections 3 and 5. All are version-controlled and available in the repository <https://github.com/x67ai/riemann-rh-program>; paths below are relative to `rh-program/` in that repository. The certificate is also available as a self-contained archive, `haglund-counterexample-certificate.zip`, with SHA-256 ZIPHASH1ZIPHASH2, which contains both certificates, the independent re-runs, the third evaluation, and a README with the exact commands and expected outputs. On one core of an Apple M4, certificate A runs in about 72 seconds (Python 3.9.6, python-flint 0.6.0) and certificate B in about 15 minutes (Python 3.9.6, mpmath 1.3.0), most of it the 4400-bit literal cross-check. The components are:

- *Certificate A* (`results/haglund-cert-s37/producer-A/`): the evaluation library `hag_core.py`, the scripts `ladder.py`, `h1.py`, `h2.py`, `xcheck.py`, `h5.py`, their logs, and `CERT.md`, with the proofs of Lemmas 6.1–6.3 and of the half-plane fact and the SHA-256 hash of every script and log. SHA-256 of `CERT.md`: `b1ecfe0a604c8b05e3439ad9ecae1c2a86c2fcdda55930cecbd1defea499513f`.
- *Certificate B* (`results/haglund-cert-s37/producer-B/`): interval complex arithmetic (`ivc.py`), Γ , ζ and h with their remainders (`specfun.py`), Φ_n , Ξ_N and the tail (`xin.py`), the Taylor model and boundary walk (`winding.py`), the scripts `ladder.py`, `cert27.py`, `crosscheck.py`, `selftest.py`, `cert_h5.py`, the logs, the DLMF pages used, and `CERT.md`, with every bound of Section 6 and its proof or source. SHA-256 of `CERT.md`: `c4a8ca6bc2d02b6f2e428d91b7cd05044272fd2d7546d37976dd1cd1648d87a3`.
- *Independent checks* (`results/haglund-cert-s37/rerun-F/` and `results/novel-wave-s36/staircase/verify-F/`): the re-runs of certificate A’s real-zero computation and of certificate B’s ladder, and the third evaluation of the literal sum in Arb.
- *The analytic and numerical record* (`results/novel-wave-s36/staircase/`): `NOTE.md`, the source of Sections 2–5; `read-F.md`, the independent re-derivation of its proofs; and `verify/`, the floating-point computations (the census, the tail coefficients, the lobe scan, and the 50-digit zeros).
- *The literature searches* (`results/haglund-cert-s37/NOVELTY-F.md` and `results/novel-wave-s36/staircase/lit/PRIOR-ART.md`) behind the prior-work paragraph.

One exception: the working corpus of fetched literature is not redistributed, for copyright reasons, and nothing here depends on it; every external claim is cited to its published source.

References

- [Ahn12] S.-H. (Shirley) Ahn, *On the zeros of component functions of the Riemann Ξ approximates*, M.A. thesis, University of Pennsylvania, 2012.
- [Bac26] M. L. Baccaro, *Haglund’s Zero-Trajectory Conjecture for the First Riemann Ξ Approximant*, preprint, version v1.0.0, Zenodo, August 22, 2026, DOI 10.5281/zenodo.22059236.
- [Bur12] J.-F. Burnol, *Deux extensions de Théorèmes de Hamburger (portant sur l’équation fonctionnelle de la fonction zêta)*, Expo. Math. **30** (2012), 295–308; arXiv:1106.4749.
- [DLMF] *NIST Digital Library of Mathematical Functions*, <https://dlmf.nist.gov/>, release date 2026-09-15; equations 2.10.1, 5.11.1, 24.9.2 and §5.11(ii).
- [Edw74] H. M. Edwards, *Riemann’s Zeta Function*, Academic Press, New York, 1974; reprinted by Dover Publications, Mineola, NY, 2001.
- [FLINT] The FLINT team, *FLINT: Fast Library for Number Theory*, version 3.0.1, <https://flintlib.org>; used through python-flint 0.6.0, <https://github.com/flintlib/python-flint>.
- [Hag09] J. Haglund, *Some conjectures on the zeros of approximates to the Riemann Ξ -function and incomplete gamma functions*, arXiv:0910.5228v1, 27 October 2009. Page and equation numbers in this paper refer to this version.

- [Hag11] J. Haglund, *Some conjectures on the zeros of approximates to the Riemann Ξ -function and incomplete gamma functions*, Cent. Eur. J. Math. **9** (2011), no. 2, 302–318, DOI 10.2478/s11533-010-0095-3.
- [Ham21] H. Hamburger, *Über die Riemannsche Funktionalgleichung der ζ -Funktion (erste Mitteilung)*, Math. Z. **10** (1921), no. 3–4, 240–254.
- [Joh17] F. Johansson, *Arb: efficient arbitrary-precision midpoint-radius interval arithmetic*, IEEE Trans. Comput. **66** (2017), no. 8, 1281–1292, DOI 10.1109/TC.2017.2690633.
- [Ki06] H. Ki, *Zeros of the constant term in the Chowla–Selberg formula*, Acta Arith. **124** (2006), no. 3, 197–204.
- [LM11] J. C. Lagarias and D. Montague, *The integral of the Riemann ξ -function*, Comment. Math. Univ. St. Pauli **60** (2011), 143–169; arXiv:1106.4348.
- [LMOZ24] D. Liu, Y. Matiyasevich, J. Oesterlé and A. Zaharescu, *Euler product sieve*, arXiv:2406.00786, 2 June 2024.
- [mpm] The mpmath development team, *mpmath: a Python library for arbitrary-precision floating-point arithmetic*, version 1.3.0, <https://mpmath.org>.
- [Nak20] T. Nakamura, *Dirichlet series with periodic coefficients, Riemann’s functional equation and real zeros of Dirichlet L -functions*, arXiv:2008.02570v4, 10 September 2021.
- [PT21] D. Platt and T. Trudgian, *The Riemann hypothesis is true up to $3 \cdot 10^{12}$* , Bull. London Math. Soc. **53** (2021), no. 3, 792–797, DOI 10.1112/blms.12460; arXiv:2004.09765.
- [Rie59] B. Riemann, *Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*, Monatsber. Königl. Preuss. Akad. Wiss. Berlin (1859), 671–680; English translation in [Edw74, Appendix].