

The two-moment certificate is robust under Rudnick–Sarnak-range cubic augmentation with capacity control

An exact absorption certificate for the two-moment system at bandwidth one, with a hand-checkable witness

Kunal Tyagi*

August 26, 2026

Dated revision (2026-08-27). The interval hardening of 2026-08-27 (`verify/o1_crowding_interval.py`; report: *supplementary material*, `o1-n128-report.md`) found the earlier full-family crowding ratio 0.9775 to be a 0.004-grid artifact. This revision re-scopes the multi-pair closure: ledger constant interval-certified $C^* \leq 0.98465 < 1$ on pair depths ≤ 0.156 ($w \leq 0.98$); certified ledger FAILURE ($S_2^{\text{gen}} \geq 1.01405 > 1$ at $d = d' = 0.159$) on $w \in (0.98, 1]$, where coverage is the re-run crowd attacks ($\min F_1 - T \geq +17.46$) plus the unconditional 8/9 backstop — (MI) itself is numerically unrefuted there. The $N = 128$ re-run reproduces all identities. A strengthening-by-honesty: the certified region’s constant is now rigorous, and the overreach is withdrawn. Complete edit list: *supplementary material*, `o1-n128-report.md`, Section 5.

Dated revision (2026-08-27). **Citation pass, 2026-08-27.** An independent novelty re-check was run before circulation; its verdict for this paper is *novel with citations*, and every citation action it requested was re-verified against a primary source before being applied (Crossref, zbMATH Open, the arXiv abstract pages, the publishers’ own records, and the program’s on-disk corpus). The changes are citation- and provenance-level only: no theorem, proof, number, or verdict is affected. Added: [GdLL25], with a reconciliation paragraph in Section 1.2 and a provenance note at Proposition 3.3; [KLS11]; and [FGL14], solely to disambiguate the numerical coincidence noted at Corollary 4.4. Completed or corrected: publication coordinates for [BGSTB24], [KLS07], [KS66], [CGdL20], [CdLS22], and DOIs for [LR20], [LR21], [Hej94], [RS96], [CGG98], [BHB13]. Three of the re-check’s own claims did **not** survive primary-source verification and were not executed: its request to renumber both “Section 7.5(e)” pointers (item (e) *is* in §7.5 of the 35-page parent version, and in §7.2(e) only in the 17-page revision — both pointers are correct, and the [P] entry disambiguates them); its description of [CGdL20]’s inequality (7) as a “convex-programming form” (it is elementary integer arithmetic, credited there to an argument of Ghosh); and its reading of [GdLL25]’s N_n remark as “a higher level adds nothing” (it is a negative about problem formulation at a fixed level). Each is recorded here so that it cannot be reintroduced.

Program: RH program, direction A4 (merged A2+A4 cubic-certificate direction). An independent adversarial review of the proposal returned *survives with repairs* and prescribed the corrected

*`hello.jay.tyagi@gmail.com`. **Use of AI.** The mathematics in this paper — the derivations, the computations, the verification suite, the Lean formalization, and the text — was produced by Claude (Anthropic) working under the author’s direction inside a structured research program; the author set the program’s objectives, adjudicated its decisions, and is responsible for the content. Claude is a tool and is not an author. The complete program record, including every session log, review and correction, is public: <https://github.com/x67ai/riemann-rh-program>.

decision gate executed here; this paper is that gate’s absorption branch, decided 2026-08-26 and independently audited. The internal terms *gate*, *adjudication* and *referee pass* are defined in the Provenance section.

Date: 2026-08-26.

Evidence discipline: every number in this paper is taken from the on-disk gate record (`results/a4-m2-gate/`), from the audited data digest (*supplementary material*, `data-tables.md`), or from re-derivations executed in the accompanying verification suite; every theorem is proved in full in Sections 3–7 or in the companion note (*supplementary material*, `pair-channel.md`). Provenance for each class of claim is collected in the References and Provenance sections.

Abstract

The parent paper’s unconditional two-thirds theorem rests on a two-moment certificate — trace and Frobenius norm of the Weil–Gabor compression, read against block structure and integrality — and that certificate is provably exhausted: the Lean-formalized `lemmaR_tight` shows that within the data class (tr, Frobenius, positive-index count, integer atoms) an on-line double and a shallow off-line pair are indistinguishable, pinning the distinct-zeros optimum at $N_d/N = 5/6$ and the simple fraction at $2/3$. Direction A4 proposed to break this degeneracy by adjoining, at a second bandwidth $\lambda' = 1/2 + o(1)$ inside the Rudnick–Sarnak range, the unconditionally computable signed third trace together with repaired spectral-capacity rows (an all- V eigenvalue-count ladder and a garnish-capacity fuzz row). This paper proves, at the model level, and certifies computationally at decision grade, that the augmentation adds exactly nothing: the marginal value of the entire cubic block over the two-moment baseline is $\delta_0 = 0$ — identically, as an exact LP equality with an explicit, hand-checkable witness law: over the full pre-registered adversary grid for the distinct-zeros benchmark (940 records, $|\delta_0| < 10^{-12}$ everywhere, corner $5/6$); for the simple-fraction benchmark (corner $2/3$) at its two decision cells plus cross-checks, converged clean at the literal pre-registered stop rule; and — decisively — inside the near-CUE data class licensed by the unconditional pair-correlation theorem of Baluyot–Goldston–Suriajaya–Turnage–Butterbaugh, where the absorption is by strict slack: the cubic row does not even bind. The structural explanation is a set of exact identities proven here: on isolated blocks the signed cubic row is the affine function $3F - 2$ of the Frobenius row plus the second integrality level, hence blind to doubles-versus-pairs at every depth; on the ψ_1 -zero grid the bandwidth-one Frobenius row degenerates exactly to the multiplicity count while the half-bandwidth kernel stays position-sensitive, so the cubic budget is met by arrangement at zero cost; and spectral escape is capped sharply at $2\sqrt{2}\sqrt{C_{\text{led}}\varepsilon_g}N = o(N)$, while a scalar cubic tail row with any divergent cutoff is unconditionally vacuous. The pair-interference channel — the one gap in the atom-only model theorem — is closed by an integrality theorem throughout the gate’s R5 depth family: proved unconditionally for single-pair columns at every gate depth and for multi-pair columns at $w \leq 0.82$ ($w \leq 0.98$ modulo one crowding cap whose ledger constant is interval-certified; on the deepest sliver $w \in (0.98, 1]$ the coverage is adversarial-numerical with the unconditional $8/9$ backstop), with an unconditional $8/9$ -strength spectral backstop at all depths; mixed-depth multi-pair columns involving the deep probes $w \in \{1.5, 2\}$ rest on that backstop plus converged adversarial pricing. The two-moment certificate, and with it the bandwidth-one ceiling, is robust under Rudnick–Sarnak-range cubic augmentation with capacity control.

1 Introduction

1.1 The parent certificate and its exhaustion

The parent paper [P] proves unconditionally that more than two thirds of the zeros of the Riemann zeta function are simple and on the critical line, that $N_d(T, 2T) \geq (5/6 - o(1))N(T, 2T)$ zeros are distinct, and (Theorem D, window-optimized) that the simple-on-line proportion reaches $2 - 1/c_1^* =$

0.6725007..., by compressing the Weil explicit-formula form to a critically sampled Gabor family and reading exactly two spectral moments of the resulting Gram matrix $\widehat{G}$: the trace $\text{tr } \widehat{G} = N(1 + o(1))$ and the Frobenius moment $\text{tr } \widehat{G}^2 = (1/\lambda + \lambda/3)N(1 + o(1))$, both evaluated unconditionally on the prime side for bandwidth $\lambda \leq 1$. The certificate is finite-dimensional linear algebra (the rank–trace inequality “Lemma R”) applied against the zero side’s block structure: on-line points give PSD rank-one atoms, off-line pairs give signature-(1, 1) hyperbolic blocks.

Two formalized results delimit this method exactly. First, `lemmaR.tight` (Lean, `Zeta23/ZeroSide/TightMult.lean`): on configurations of integer-marked on-line atoms plus pair blocks, the two-moment certificate is achieved with equality, and an on-line double (eigenvalue +2, charge 4) is indistinguishable from an off-line pair of depth $\rightarrow 0$ (eigenvalues ± 2 , charge 4) — no further inequality in $(\text{tr}, \text{Frobenius}, n_+, \text{integer atoms})$ can improve it. The extremal configuration is $2/3$ simple zeros plus $1/6$ on-line doubles: $N_d/N = 5/6$, simple fraction $2/3$. Second, the bandwidth-one ceiling (Lean, `Zeta23/PairCeiling/`): no certificate reading only bandwidth-one form-factor data certifies a simple-zero proportion above 0.6818287 (up to the recorded stability terms), witnessed by an exact-rational 256-periodic law; and any trace/Frobenius certificate is capped below $0.8453 = 2 - 2/\sqrt{3}$ at any Dirichlet-polynomial length, since $\kappa(\lambda) = 1/\lambda + \lambda/3 \geq 2/\sqrt{3}$ for all $\lambda > 0$.

1.2 The cubic proposal

The parent’s Section 7.3 records the conditional template: under RH the third trace $\text{tr } \widehat{G}^3$ is available (Hejhal, Rudnick–Sarnak) and the certificate can be run with the cubic weight $\omega(m) = m/2 + (2m^2 - m^3)/18 + (4/9)\mathbf{1}_{m=1}$, giving $N_d/N \geq 0.85082$ with the window $\cos(8s/5)$ (consuming also [BHB13]’s $N^s/N \geq 19/27$ on RH). Direction A4’s thesis was that a slice of this becomes unconditional: at bandwidth $\lambda' = 1/2 + \theta \log \log T / \log T$ (strictly inside the $k = 3$ Rudnick–Sarnak range $X^3 \leq T^{2-\varepsilon}$), the signed cubic trace is computable by the parent’s Section 5 diagonal method, and a Schatten-3 spectral-tail theorem (Theorem 1(ii)-repaired: an all- V eigenvalue-count ladder $n(V) \leq C_{\text{led}} NV^{-4}$, $\theta < 1$ strict) would cap the spectral escape that the parent’s own odd-moment no-go (Section 7.5(e) of the 35-page version; the same item is §7.2(e) of v5 — see the version pin in the References) identifies, making the cubic row a usable sign-sensitive constraint. The original proposal’s separating premise was that a hyperbolic pair block contributes $(+a)^3 + (-a)^3 = 0$ to the cubic while a double contributes +8, so the two-moment-degenerate configurations would sit at different corners of the augmented polytope.

What the parent already says, and what it leaves open (stated here so the contribution is not mistaken for it). The parent’s item (e) is not silent on higher moments. In the 35-page version it reads, verbatim:

“(e) The prime-side evaluation of $\text{tr } \widetilde{G}^k$ by the diagonal method of Section 5 (multiplicative relations among k prime powers, Montgomery–Vaughan for the rest) is available exactly in the Rudnick–Sarnak range $k\lambda < 2$ [RS96]; for $\lambda \in (1/2, 1)$ this allows at most $k = 3$ (and only for $\lambda < 2/3$), and an odd moment does not lower $\Lambda_1(0)$. Thus, unconditionally, higher moments add nothing to the n_+ -bound on $(1/2, 1)$ (and Proposition 4.4 uses only the first two in any case), while for $\lambda \leq 1/2$, where many moments are available, Proposition 7.4 makes them useless.”

(In v5 the same item is compressed to “...at $X \asymp T$ this allows only $k = 1$. Thus, unconditionally, higher moments add nothing.”) That remark is a **single-bandwidth** statement, and it disposes of the single-bandwidth question twice over, for two disjoint reasons: at bandwidth $\lambda \in (1/2, 1)$ an odd moment does not move the Christoffel bound, and at $\lambda \leq 1/2$, where moments

are plentiful, the certificate is worthless on its own (its Proposition 7.4 cap is non-positive there). It is also, in the parent’s own words, stated informally.

The question this paper answers is a different one, and the parent’s remark does not settle it: **keep the bandwidth-one Frobenius baseline and adjoin a second-bandwidth block at $\lambda' = 1/2$** — the λ' -Frobenius budget, the signed cubic equality, the all- V count ladder, and the capacity fuzz row. Rows that are useless alone can bind when adjoined to a strong baseline, which is exactly why a marginal-value computation is required rather than a restatement; and the parent’s remark addresses neither the distinct-zeros benchmark, nor the pair-block structure, nor the near-CUE data class, and produces no witness. What is new here is the answer — $\delta_0 = 0$ as an exact LP equality, witness-backed, on both benchmarks, and by strict slack inside the near-CUE class — together with the structural identities (Sections 3–4) and the capacity theorem (Section 7) that explain it. The parent’s single-bandwidth no-go is the parent’s; the two-bandwidth marginal value is this paper’s.

The published third-party comparandum, and why it does not settle this question either. Gonçalves, de Laat and Leijenhorst [GdLL25] run the higher-level-correlation program against multiplicity directly, by semidefinite programming over an auxiliary-function class: on GRH they obtain, from 3-level correlations, that at least 96.14% of the zeros of a primitive L -function have multiplicity at most 2, and their $n = 3$ row pays where their $n = 2$ row does not. That is not in tension with the result here, and the two must not be conflated. Theirs is a **single-bandwidth, conditional** statement bounding *multiplicity* at their support parameter; this paper’s is an **unconditional** statement about the *marginal value of adjoining a second bandwidth* to a fixed bandwidth-one Frobenius baseline, at effective support 1 (their $m = 2$ row), measured against a fixed pre-registered objective. The combinatorial half of their apparatus and of this paper’s is literally the same identity — their equation (5) is the Stirling change of basis between the power sums $N_n = \sum_{\text{zeros}} m^n$ and the falling-factorial counts $M_k = \sum_{\text{zeros}} m(m-1)\cdots(m-k+1)$, whose $n = 3$ case is the atom half of Theorem 3.4 below — which is exactly why the atom half is labeled here as classical and only the pair term $6 \sum_p m_p^2(m_p-1)A_p^2$ is claimed as this paper’s. Their paper also carries a published negative of a related species: the remark following their (6) records that adapting their optimization to bound N_n directly did not improve on the bounds obtained through (5). That is a negative about problem *formulation*, not about the information content of a higher correlation level, and it is cited here as a precedent for publishing such a negative rather than as evidence for this paper’s claim.

The adjudication ordered seven repairs (R1–R7) and made the corrected M2 gate the direction’s action one: a decision LP over an honest adversary class — marks up to a divergent alphabet W , depth-parametrized shallow pairs, clustered sine-process null budgets, full positional freedom, and an explicit garnish-capacity fuzz row derived from the repaired ladder — with the pre-registered semantics that either branch is a deliverable: a bite funds the analytic bridge; an absorption is published as the sharpened no-go. This paper is the absorption branch, executed.

1.3 The corrected gate question and its answer

The gate asks: over laws (probability mixtures) of explicit N -periodic marked configurations satisfying the two-moment data — mass N , integrality, pair structure, bandwidth-one Frobenius budget $(4/3)N(1 \mp \varepsilon)$ — does adjoining the entire $\lambda' = 1/2$ cubic block (the λ' -Frobenius budget $\kappa(1/2)N = (13/6)N$, the signed cubic equality $c_3(1/2)N = 5N$, the all- V count ladder, and the capacity fuzz row) raise the minimum of $E[N_d]/N$ above the **lemmaR.tight** corner by any $\delta_0 > 0$?

Throughout, “Rudnick–Sarnak-range cubic augmentation with capacity control” (the title phrase, pre-registered in the gate specification) means exactly this $\lambda' = 1/2 + o(1)$ system fixed by the adjudication (R4: $\theta < 1$ strict) — the only point in the RS range where both the signed cubic row and the capacity ladder are unconditionally licensed; the behavior of the cubic row at $\lambda' \geq 0.6$, inside the RS range but outside the proven ladder regime, is the quarantined payoff curve of Section 8.4, not part of any claim.

The answer, decided and audited: **no** — $\delta_0 = 0$ **identically**, not merely within error bars: the base-optimal law itself satisfies the entire cubic block. This paper assembles the complete result: the structure theorems that explain the absorption (Section 3), the model-level corner theorem with its pair-channel extension (Section 4), the computational certification with the explicit witness (Section 4.4), the near-CUE decision-grade result (Section 5), the simple-fraction benchmark (Section 6), the two capacity theorems that close the spectral-escape routes (Section 7), and the honest scope map of what is exact, what is a bounded residual, and what remains open (Section 8).

1.4 Headline results

1. **Exact absorption with witness (Theorem 4.3 + Section 4.4).** Over the full pre-registered decision grid — 940 LP records spanning mark alphabets $W \in \{2, \dots, 8\}$, ladder constants $C_{\text{led}} \in \{10, 100, 1000\}$, tolerances $\varepsilon \in \{0.02, 0.05, 0.10\}$ (with the separate, labeled tight- ε runs extending to $\varepsilon = 0.002$), all fuzz modes, matched and asymptotic budget centerings, $N = 64$ and $N = 128$ — the marginal value of the cubic block is $|\delta_0| < 10^{-12}$ at every record (maximum 8.94×10^{-13}), and the joint optimum obeys the exact law $P(\varepsilon) = 5/6 - (2/3)\varepsilon \rightarrow 5/6$. The absorbing law is exhibited: three columns on the 65-site ψ_1 -zero grid, marks $\{1, 2\}$ only, no pairs, every row verified in exact rational arithmetic, the key entries hand-checkable ($F_1 = 64$ exactly; $F' = 4128/33$).
2. **Both benchmarks.** The same absorption holds for the simple-fraction objective: the cubic block moves the p_1 optimum by at most 1.4×10^{-12} , converged at the literal pre-registered stop rule, and the optimum is the exact analytic corner $2 - (4/3)(1 + \varepsilon) \rightarrow 2/3$ (Section 6).
3. **Near-CUE decision grade — absorption by slack (Section 5).** Inside the pinned data class $|E_w|c_j|^2 - j| \leq 1$ (licensed unconditionally, with three stated riders, by [BGSTB24]), $\delta'_0 = 0$ as an exact LP equality at $N = 64$ and $N = 128$, at both budget centerings — and at the pinned optimum only the pinning rows bind: the cubic row, the ladder, and even both Frobenius rows are strictly interior.
4. **Cubic blindness (Theorem 3.4).** On isolated blocks, the signed cubic row equals $3F - 2$ per zero on every multiplicity-1 pair at every depth and on every mark- $\{1, 2\}$ atom: $C - 3F + 2\text{Mass} = \sum m(m-1)(m-2) + 6 \sum m_p^2(m_p-1)A_p^2$ exactly. The “+8 vs 0” separating premise of the proposal (and the adjudication’s phase model) is false: a pair’s cubic charge is $2m^3(1 + 3A^2) \geq 8m^3$ at every depth. All discriminating power of the cubic row lives in marks ≥ 3 and in clustering cross-terms — and the cross-terms are a free resource for the adversary (Theorem 3.9).
5. **Closed forms and the $m_2 = \kappa$ identity (Theorem 3.5).** The sine-process null budgets are $m_2(\lambda) = 1/\lambda + \lambda/3$ and $m_3(\lambda) = 1 + 1/\lambda^2$ for the flat window, all $0 < \lambda \leq 1$; and $m_2(\lambda)$ coincides exactly with the parent’s unconditional prime-side $\kappa(\lambda)$ — the sine null’s second-moment budget IS the unconditional budget, on the whole band.

6. **Capacity cap (Theorem 7.1).** Under the repaired all- V ladder and a Frobenius slack ε_g , the maximal cubic charge purchasable by spectral escape is exactly $2\sqrt{2}\sqrt{C_{\text{led}}\varepsilon_g}N = o(N)$ — sharp, with the extremal profile exhibited; both the earlier constants $(2\sqrt{C\varepsilon}, (5/\sqrt{3})\sqrt{C\varepsilon})$ are superseded, with three independent confirmations.
7. **Divergent-cutoff vacuity (Theorem 7.4).** A scalar Schatten-3 tail row with ANY divergent cutoff is unconditionally absorbed by an explicit garnish construction that lowers both objectives — the “provable today, no LP needed” half of the no-go.
8. **Pair-interference closure (Sections 4.2–4.3).** The one channel outside the atom-only sandwich — off-line pairs pushing the bandwidth-one Frobenius below the mark-accounting floor — is closed for the gate’s pair class throughout the R5 depth family: proved unconditionally for all single-pair configurations to depth $w \leq 2.8$ (covering all gate depths, including both deep probes, with $\geq 34\%$ margin), for all multi-pair configurations to $w \leq 0.82$ ($w \leq 0.98$ modulo the crowding cap, ledger constant interval-certified at 0.98465; on $w \in (0.98, 1]$ the ledger fails — certified — and coverage is adversarial numerics plus the 8/9 backstop), and for equal-depth pair-only configurations at every depth. Multi-pair configurations at mixed depths involving the deep probes $w \in \{1.5, 2\}$ — admissible under Section 2.4 but beyond the theorems’ range — are covered by adversarial numerics plus the unconditional 8/9 backstop (O2, Section 8.3). An explicit fractional-mark counterexample proves the closure is intrinsically an integrality theorem.

1.5 Honest framing (adjudication repair R6, binding)

The cubic row is a **nonlinear spectral reading of bandwidth-one pair-correlation data with quantitative high-spectrum capacity control** — nothing more. The third trace at bandwidth λ' carries total Fourier support $(k-1)\lambda' = 1 + o(1)$, not $3\lambda'$; its genuinely cubic prime resonance is the absolute constant $\sum_p (\log p)^3 / (p-1)^2 = 2.315762$ (re-sieved to 3×10^6 in this program, three independent confirmations), contributing $O(1)$ to $\text{tr } \widehat{G}^3$, i.e. $O(1/N)$ per zero. No claim is made anywhere in this paper that the cubic block evades the Alternative Hypothesis or consumes data beyond the bandwidth-one class plus $o(1)$ — for what the currently known band-limited n -level data provably cannot do against AH, see Lagarias–Rodgers [LR20], and Tao’s independent treatment cited there: the no-go proved here is precisely that this nonlinear reading, with the full capacity apparatus attached, adds nothing to the linear reading at the proven operating point. (The formal scope point survives unchanged: a cubic row is outside the PairCeiling certificate class as defined; the content of this paper is that leaving the class buys nothing.)

1.6 Scope discipline (read before citing)

Every theorem in this paper is a statement about the model — the gate’s explicit configuration classes, kernels, and row systems (Section 2) — not about the zeros of zeta. The bridge from the model rows to unconditional statements about zeta is the parent paper’s sampling bookkeeping plus the [BGSTB24] licensing riders, cited (with the riders stated in full) in Section 5.1. The exact zero $\delta_0 = 0$ is proved and certified at the flat/flat window configuration and is **flat/flat-specific**: with the Montgomery–Taylor cosine window at $\lambda = 1$ the grid degeneracy breaks and the honest statement is a bounded residual of order 6×10^{-5} (Section 8). Nowhere does this paper claim “ $\delta_0 = 0$ universally”; the claim structure is: exact zero on the decision configuration and the near-CUE class, bounded sub-decision-scale residuals on the window family, a proven $o(N)$ cap

on spectral escape everywhere, and a labeled exploratory payoff curve outside the proven ladder regime (Section 8.4).

2 The model

This section fixes the model in which every theorem of Sections 3–7 lives: the master trace formula, its finite-circle discretization, the adversary class, and the row system. It condenses the gate specification (SPEC Sections 1–4); everything here was re-derived in the accompanying verification suite.

2.1 Window and kernel

Fix a window profile $v \geq 0$ on $[-1/2, 1/2]$, even, with $\int v > 0$ (the scale-free profile of the parent’s taper-squared ϕ^2 ; the formalized taper class is C^3). The **normalized sampling kernel** at bandwidth $\lambda > 0$ is the entire function

$$\psi_\lambda(x) = \frac{\int_{-1/2}^{1/2} v(\sigma) e^{2\pi i \lambda \sigma x} d\sigma}{\int v},$$

so $\psi_\lambda(0) = 1$, with Fourier transform u supported in $|\alpha| \leq \lambda/2$ (Montgomery α -units), $\int u = 1$. For the flat window $v = 1$:

$$\psi_\lambda(x) = \text{sinc}(\pi \lambda x) = \frac{\sin(\pi \lambda x)}{\pi \lambda x}, \quad u = \frac{1}{\lambda} \mathbf{1}_{[-\lambda/2, \lambda/2]}.$$

Positions x are in mean-gap units (mean zero spacing 1); complex positions (off-line zeros) are allowed, ψ_λ being entire.

2.2 Continuum model: configurations and the master trace formula

A **marked configuration** is a finite multiset of points z with positions γ_z (complex allowed) and integer marks $m_z \geq 1$; off-line points occur only in conjugate pairs $\{\gamma + id, \gamma - id\}$ with common depth and common mark (functional-equation symmetry). The **master trace formula** defines the k -th trace at bandwidth λ :

$$\text{tr } G^k := \sum_{z_1, \dots, z_k} \left(\prod_j m_{z_j} \right) \psi_\lambda(\gamma_{z_1} - \gamma_{z_2}) \psi_\lambda(\gamma_{z_2} - \gamma_{z_3}) \cdots \psi_\lambda(\gamma_{z_k} - \gamma_{z_1}),$$

summed over ordered k -tuples of configuration points. This is the parent’s Poisson sampling identity in normalized units (units in which an isolated on-line zero of mark m has eigenvalue exactly m), and it is realized by an actual Hermitian matrix — Section 3.1 constructs the block matrices explicitly where spectral (not just trace) data are needed. We write $F = \text{tr } G^2$ (Frobenius row) and $C = \text{tr } G^3$ (signed cubic row); F_1 at bandwidth 1, F' and C' at bandwidth λ' . This single formula generates every trace row on both sides — null budgets and configuration demands — including all clustering cross-terms and all depth effects; no isolated-block idealization enters the gate anywhere.

2.3 Finite-circle model (the LP’s discretization)

Period N (zeros per period) on the circle of circumference N ; configurations are N -periodic, described by one period. For the flat window at bandwidth λ the harmonic set is $B = \{j \in \mathbf{Z} : |j| \leq \lambda N/2\}$, with $M = |B|$ harmonics and uniform weights $u_j = 1/M$. With the form factor

$$c_s = \sum_z m_z e^{-2\pi i s \theta_z / N}$$

(a conjugate pair at $\theta \pm id$ contributes $2m \cosh(2\pi s d/N) e^{-2\pi i s \theta/N}$), the trace rows are

$$\begin{aligned} \text{tr } \hat{G} &= \sum_z m_z & (\text{mass}), \\ \text{tr } \hat{G}^2 &= \sum_s W_2(s) |c_s|^2, & W_2(s) = \sum_{j \in B, s-j \in B} u_j u_{s-j}, \\ \text{tr } \hat{G}^3 &= \sum_{k_1, k_2} U_3(k_1, k_2) c_{k_1} c_{k_2} c_{-k_1-k_2}, & U_3(k_1, k_2) = \sum_j u_{j+k_1} u_{j+k_1+k_2} u_j. \end{aligned}$$

Eigenvalue data (ladder counts, inertia) come from the Hermitian frequency matrix $H[j, j'] = \sqrt{u_j u_{j'}} c_{j-j'}$ ($j, j' \in B$), whose k -th trace equals the k -th trace row exactly (alias-free tight-frame compression). At the primary geometry $N = 64$, $\lambda = 1$: $M = 65$; $\lambda' = 1/2$: $M' = 33$. In position space, for a symmetric band, $\text{tr } \hat{G}^2 = \sum_{z, z'} m_z m_{z'} D(\theta_z - \theta_{z'})^2$ with $D(x) = \sum_j u_j e^{-2\pi i j x/N}$ real, even, $D(0) = 1$ — the discrete analog of the continuum $\text{tr } G^2 = \sum m m' \psi_\lambda^2$.

2.4 Adversary class, laws, and the two benchmarks

A **column** is an explicit N -periodic marked configuration: on-line atoms (positions $\theta_i \in [0, N)$, integer marks $1 \leq m_i \leq W$) and off-line pairs (position, dimensionless depth w in the family $(0, 1]$ plus deep probes $\{1.5, 2\}$, pair multiplicity m), with mass $\sum m_i + 2 \sum m_p = N$ per period. A **law** is a probability mixture $w_c \geq 0$, $\sum_c w_c = 1$ over columns; every row is $E_w[\text{row}(c)]$. The two objectives:

$$\begin{aligned} N_d(c) &= \# \text{atoms} + 2 \# \text{pairs} & (\text{distinct zeros; only on-line multiples reduce it}), \\ p_1(c) &= (\# \text{mark-1 atoms} + 2 \# \text{mult-1 pairs}) / N & (\text{simple fraction}). \end{aligned}$$

Realizability — what is imposed and why it is exactly right (SPEC 3.2, in brief). Imposed: explicit finite marked multisets (never bare correlation data); integer marks; conjugate-pair structure; mass N ; mixtures. *Necessity*: an adversary allowed to post aggregated moment data directly could claim values realized by no point configuration (unconstrained cross-term variables can leave the convex hull of realizable moment pairs), making an “absorption” vacuous; and integrality, pair symmetry, and density are properties the true zero multiset provably has — dropping any of them hands the adversary provably impossible worlds (with fractional marks even the 5/6 baseline is false: mass N of mark-4/3 atoms on the grid of Theorem 3.9 has $F_1 = (4/3)N$ budget-tight and $N_d = (3/4)N < (5/6)N$; Section 4.2 exhibits the subtler pair-channel form of the same failure). *Sufficiency*: nothing further may be imposed, because every row of the system is a theorem valid for arbitrary configurations with these properties — trace identities are linear algebra plus the sampling identity, the budgets are the unconditional prime-side equalities, the ladder is Theorem 1(ii)-repaired, inertia is Sylvester; imposing anything more (e.g. pointwise GUE 3-level correlations) would smuggle in precisely the unproven correlation knowledge whose absence the gate respects. Periodicity and the finite depth/mark grids are discretizations, refined under the verdict-stability requirement ($N = 64 \rightarrow 128$, W -scan, depth-grid halving), not constraints.

2.5 The row system

Baseline (two-moment / `lemmaR.tight` data): mass N ; integrality and pair structure; bandwidth-one Frobenius budget $E_w[F_1] \in (4/3)N(1 \mp \varepsilon)$ (matched-geometry variant: the finite- N CUE-sampled center B_1 replaces $(4/3)N$). The cubic block at $\lambda' = 1/2$ adds:

- (R-5) λ' -Frobenius: $E_w[F'] \in \kappa(1/2)N(1 \mp \varepsilon)$, $\kappa(1/2) = 13/6$;
- (R-6) signed cubic with capacity coupling: $|E_w[C'] - c_3(1/2)N| \leq \varepsilon_3 N + 2\sqrt{2}\sqrt{C_{\text{led}}g_F} N$, $c_3(1/2) = 5$;
- (R-5') garnish Frobenius budget $g_F \geq 0$ charged against the Frobenius slack;
- (R-7) all- V count ladder: $E_w[n(V)] \leq C_{\text{led}}NV^{-4}$ on the V -grid $\{2, 3, 4, 6, 8, 12, 16\}$;
- (R-8) inertia $n_-(c) \leq p(c)$ per column (automatic on explicit spectra; asserted);
- (R-9) Frobenius split, power-mean, Hölder-with- $o(N)$: automatically satisfied by explicit spectra (the R7 fallback system made primary — on explicit configurations the linear-algebra-true rows cannot be violated, so their entire content is enforced leak-free by construction).

The budget centers are the clustered sine-process null moments (Section 3.3) — never isolated-block values — because at $\lambda' = 1/2$ the window width is two mean gaps and the null itself is cross-term-dominated: cross-terms (occupancy ≥ 2) carry 54% of the Frobenius budget and 80% of the cubic budget (analytic fractions 7/6 of 13/6 and 4 of 5; null MC 52.6% / 79.0% at $n = 64$). The decision rule (pre-registered): ABSORPTION iff $\delta_0 = P_{\text{full}} - P_{\text{base}} = 0$ within LP tolerance at the decision points with an explicit witness law; the witness must satisfy every row at centered budgets with $E[N_d]/N \leq 5/6 + 0.01$.

3 Structure theorems

We lead with the theorem that explains everything: on isolated blocks the signed cubic row is an affine function of the mass and Frobenius rows plus the second integrality level — it cannot see doubles-versus-pairs at any depth. We then compute the null budgets in closed form, obtaining the identities that organize the entire margin structure, and finally prove the grid-decoupling theorem that hands the adversary the cross-term channel for free. Every proof in this section is complete; the numerical confirmations (all re-run for this paper, `verify/verify_t1.py-verify_t4.mc.py`) are summarized at the point of use.

3.1 The exact pair block

Setting. Windows $\tau_k = k(2\pi/L)$, $k \in \mathbf{Z}$ (critical spacing); taper ϕ real, even, continuous, supported in $[-L/2, L/2]$, $\phi \in C^2$ (the parent's formalized class is C^3 — more than enough). Fourier convention $\hat{\phi}(z) = \int \phi(s)e^{-isz} ds$ (entire, by Paley–Wiener); $\Phi(z) = \int \phi(s)^2 e^{-isz} ds$; $a = L^{-1} \int \phi^2$, so $\Phi(0) = La$. Normalized units: matrices divided by $L^2 a$, in which an isolated on-line atom of mark m has eigenvalue m .

Lemma 3.1 (Poisson sampling identity, complex arguments). *For all complex τ, τ' :*

$$\sum_{k \in \mathbf{Z}} \hat{\phi}(\tau - \tau_k) \hat{\phi}(\tau' - \tau_k) = L \Phi(\tau - \tau'),$$

the sum converging absolutely and locally uniformly.

Proof. First real arguments. Fix $\tau, \tau' \in \mathbf{R}$ and set $g(t) = \widehat{\phi}(\tau-t)\widehat{\phi}(\tau'-t)$; $\widehat{\phi}$ is real on $\mathbf{R}$ (ϕ real even) and, since $\phi \in C^2$ with compact support, $|\widehat{\phi}(x)| \leq C/(1+x^2)$, so $g \in L^1$ with $|g(t)| \leq C'/(1+t^2)^2$ and the lattice sum converges absolutely. Each factor $t \mapsto \widehat{\phi}(\tau-t)$ has Fourier transform $e^{-i\tau\xi}2\pi\phi(\xi)$ (inversion), supported in $[-L/2, L/2]$; hence $\widehat{g} = (1/2\pi)(\text{convolution of the factor transforms})$ is continuous, supported in $[-L, L]$, and vanishes AT the endpoints $\pm L$ (the convolution of two continuous functions supported in $[-L/2, L/2]$, evaluated at L , integrates over the null set $\{L/2\}$). Poisson summation over the lattice $(2\pi/L)\mathbf{Z}$ (classical form; $|g| + |\widehat{g}|$ summable on the respective lattices):

$$\sum_k g(2\pi k/L) = \frac{L}{2\pi} \sum_{n \in \mathbf{Z}} \widehat{g}(nL) = \frac{L}{2\pi} \widehat{g}(0),$$

all $n \neq 0$ terms vanishing by the support ($|nL| \geq L$, endpoint value 0). By Plancherel, $\widehat{g}(0) = \int \widehat{\phi}(\tau-t)\widehat{\phi}(\tau'-t) dt = (1/2\pi) \int (2\pi\phi(\xi))^2 e^{-i(\tau-\tau')\xi} d\xi = 2\pi \Phi(\tau-\tau')$, giving the identity on $\mathbf{R}^2$. For complex arguments: $|\widehat{\phi}(x+iy)| \leq e^{L|y|/2} C/(1+x^2)$, so on compact subsets of $\mathbf{C}^2$ the lattice sum converges absolutely and uniformly and is entire in (τ, τ') , as is $L\Phi(\tau-\tau')$; the two entire functions agree on $\mathbf{R}^2$, hence everywhere (identity theorem, one variable at a time). $\square$

Proposition 3.2 (exact off-line pair block — the true Prop-4.1 spectrum). *Let the conjugate pair be $\{\gamma + i\delta, \gamma - i\delta\}$, $\gamma \in \mathbf{R}$, depth $\delta > 0$, multiplicity m , with evaluation vector $x = (x_k)$, $x_k = \widehat{\phi}(\gamma + i\delta - \tau_k)$, and block matrix $M = m(xx^\top + \bar{x}\bar{x}^\top)$ (transpose, not conjugate-transpose: the two rank-one terms are the pair's two zeros). Then, exactly:*

- (i) $x^\top x = L\Phi(0) = L^2a$ — real, independent of depth AND position;
- (ii) $|x|^2 = L\Phi(2i\delta) = L^2a A(\delta)$, $A(\delta) = \int \phi^2(s) \cosh(2\delta s) ds / \int \phi^2(s) ds \geq 1$;
- (iii) writing $x = p + iq$: $\langle p, q \rangle = 0$ exactly, $|p|^2 = L^2a(1+A)/2$, $|q|^2 = L^2a(A-1)/2$;
- (iv) $M = 2m(pp^\top - qq^\top)$, rank ≤ 2 real symmetric, with nonzero eigenvalues (normalized by L^2a)

$$m(1+A(\delta)) \quad \text{and} \quad m(1-A(\delta)),$$

independent of the position γ ;

- (v) *normalized charges: trace $2m$; Frobenius $2m^2(1+A^2)$; signed cubic $2m^3(1+3A^2)$.*

In dimensionless depth $w := L\delta$, A depends on the pair only through w , with $A(0+) = 1$ and A strictly increasing; for the flat taper $A(w) = \sinh(w)/w$, with values $A(1/8) = 1.002606$, $A(1/4) = 1.010449$, $A(1/2) = 1.042191$, $A(3/4) = 1.096422$, $A(1) = 1.175201$, $A(2) = 1.813430$.

Proof. (i) Lemma 3.1 at $\tau = \tau' = \gamma + i\delta$: $\sum_k \widehat{\phi}(\gamma + i\delta - \tau_k)^2 = L\Phi(0) = L \int \phi^2 = L^2a$ — real, independent of γ and δ . (ii) ϕ real and even gives $\widehat{\phi}(\bar{z}) = \overline{\widehat{\phi}(z)}$; hence $|x_k|^2 = \widehat{\phi}(u - \tau_k) \widehat{\phi}(\bar{u} - \tau_k)$ with $u = \gamma + i\delta$, and Lemma 3.1 at $(u, \bar{u})$ gives $|x|^2 = L\Phi(2i\delta) = L \int \phi^2(s) \cosh(2\delta s) ds$ (ϕ^2 even); $A \geq 1$ by $\cosh \geq 1$, strictly increasing by strict monotonicity of $\cosh$ on the support. (iii) $x^\top x = |p|^2 - |q|^2 + 2i\langle p, q \rangle$; by (i) it is real, so $\langle p, q \rangle = 0$ and $|p|^2 - |q|^2 = L^2a$; with $|p|^2 + |q|^2 = |x|^2 = L^2aA$, solve. (iv) $xx^\top + \bar{x}\bar{x}^\top = (p+iq)(p+iq)^\top + (p-iq)(p-iq)^\top = 2(pp^\top - qq^\top)$; with $p \perp q$ the eigenvectors are p, q with eigenvalues $2m|p|^2 = mL^2a(1+A)$ and $-2m|q|^2 = mL^2a(1-A)$; all others 0; normalize. Position-independence: the eigenvalues depend only on $|p|^2, |q|^2$, which depend only on δ . (v) Trace $m(1+A) + m(1-A) = 2m$; Frobenius $m^2(1+A)^2 + m^2(1-A)^2 = 2m^2(1+A^2)$; cubic $m^3(1+A)^3 + m^3(1-A)^3 = 2m^3(1+3A^2)$. Flat-taper A : $A(w) = \int_{-1/2}^{1/2} \cosh(2w\sigma) d\sigma = \sinh(w)/w$; the listed values are its evaluations (they reproduce the gate specification's table to its displayed precision, $\leq 4 \times 10^{-6}$). $\square$

Consistency and inertia. The two-point configuration $\{\gamma \pm i\delta\}$ under the master formula has $\text{tr } G^k = m^k \text{tr } K^k$ with the 2×2 kernel $K = \begin{bmatrix} 1 & A \\ A & 1 \end{bmatrix}$, eigenvalues $1 \pm A$ — the same charges. The matrix route (iv) additionally certifies the inertia (one positive, one negative eigenvalue for $\delta > 0$) consumed by the n_- and ladder rows.

Block continuity, exact form. As $\delta \rightarrow 0+$, the spectrum tends to $(2m, -0)$, Frobenius to $2m^2$, cubic to $8m^3$: a shallow pair is row-indistinguishable from an on-line double in EVERY row, continuously. And — this is the correction of record — the cubic charge $2m^3(1 + 3A^2) \geq 8m^3$ at EVERY depth ($A \geq 1$), increasing with depth: a deep pair never loses cubic charge. The “+8 vs 0” separation premise (the proposal’s Step-4 narrative; the adjudication’s computation-8 phase model) is false under the true block law; the campaign transcript’s recorded block facts (pair eigenvalues $mL^2a(1 \pm A)$) were correct throughout.

Numerical confirmation (`verify_t3.py`): evaluation-vector route with a cos taper, $L = 40$, 120001 windows: $|x^\top x / L^2a - 1| \leq 6.7 \times 10^{-16}$, $|\langle p, q \rangle| / L^2a \leq 1.7 \times 10^{-14}$, block eigenvalues vs $m(1 \pm A)$ to 7.8×10^{-14} , position-independent at machine precision at every depth $w \in \{1/64, 1/8, 1/4, 1/2, 3/4, 1, 2\}$; shallow-pair cubic charge 8.0005 at $w = 1/64$ and 8.0313 at $w = 1/8$, matching the audit’s independent 8.0005/8.033 — never 0 at any depth.

3.2 The affine-plane identity and cubic blindness

Definition (per-zero charges). For an isolated block of total mass μ ($\mu = m$ for an atom of mark m ; $\mu = 2m$ for a pair of multiplicity m), let $F = (\text{block Frobenius})/\mu$ and $c = (\text{block cubic})/\mu$. By Proposition 3.2(v) and the atom spectrum:

$$\begin{aligned} \text{atom, mark } m : \quad F &= m, & c &= m^2; \\ \text{pair, mult } m, \text{ depth } : \quad F &= m(1 + A^2), & c &= m^2(1 + 3A^2). \end{aligned}$$

Proposition 3.3 (affine-plane identity).

$$\begin{aligned} \text{atom of mark } m : \quad c &= 3F - 2 + (m - 1)(m - 2); \\ \text{pair of mult } 1, \text{ ANY depth } : \quad c &= 3F - 2 \quad \text{EXACTLY}; \\ \text{pair of mult } m, \text{ ANY depth } : \quad c &= 3F - 2 + (m - 1)(m - 2) + 3m(m - 1)A^2. \end{aligned}$$

In particular every mark- $\{1, 2\}$ atom and every multiplicity-1 pair — at every depth — lies on the same affine plane $c = 3F - 2$ in the per-zero (F, c) plane.

Provenance of the atom half (stated so the novelty claim is not misread). For atoms the identity $c - 3F + 2 = (m - 1)(m - 2)$, and its summed form in Theorem 3.4, is the third-order **Stirling change of basis** between power sums and falling factorials: $m(m - 1)(m - 2) = m^3 - 3m^2 + 2m$, the coefficients being the signed Stirling numbers of the first kind $s(3, 3) = 1$, $s(3, 2) = -3$, $s(3, 1) = 2$. The inverse conversion, $\sum m^n = \sum_k S(n, k)M_k$ with Stirling numbers of the second kind, is equation (5) of [GdLL25], where it plays the same role — including the same consequence, that the falling-factorial term vanishes identically once every multiplicity is at most $n - 1$. Nothing in the atom half is new, and it is not claimed to be. What is new in Theorem 3.4 is the **pair** term $6 \sum_{\text{pairs}} m_p^2(m_p - 1)A_p^2$, a statement about the finite-circle block law of Proposition 3.2 with no counterpart in the multiplicity-statistics literature.

Proof. Atom: $3F - 2 + (m - 1)(m - 2) = 3m - 2 + m^2 - 3m + 2 = m^2 = c$. Pair: $c - 3F + 2 = m^2(1 + 3A^2) - 3m(1 + A^2) + 2 = (m^2 - 3m + 2) + 3A^2(m^2 - m) = (m - 1)(m - 2) + 3m(m - 1)A^2$.

At $m = 1$ both terms vanish identically in A (every depth); at $m = 2$ the atom identity also reads $c = 3F - 2$ (doubles are on the plane). $\square$

Theorem 3.4 (cubic blindness on isolated blocks). *Let the configuration be a disjoint union of isolated blocks (atoms of marks m_i ; pairs of multiplicities m_p at depths with $A_p = A(\delta_p)$), with total mass Mass , Frobenius F , signed cubic C (sums of block charges). Then, exactly:*

$$C - 3F + 2\text{Mass} = \sum_{\text{zeros}} m(m-1)(m-2) + 6 \sum_{\text{pairs}} m_p^2(m_p-1)A_p^2,$$

where the first sum runs over the zero multiset (an atom of mark m contributes $m(m-1)(m-2)$ once; a pair of mult m contributes twice that — its two zeros). Consequences:

1. On the class $\{\text{mark-}\{1,2\} \text{ atoms}\} \cup \{\text{multiplicity-1 pairs of arbitrary depth}\}$ — the entire doubles-vs-pairs comparison class of `lemmaR_tight` — the right side vanishes identically: $C = 3F - 2\text{Mass}$ **identically on the class**. The signed cubic row restricted to isolated blocks of this class is an exact affine combination of the Frobenius and mass rows: it separates nothing those two rows do not. An on-line double and an off-line pair of any depth are cubic-indistinguishable.
2. All discriminating power of the cubic row beyond the two-moment data therefore lives in exactly two places: (i) marks ≥ 3 (and pair multiplicities ≥ 2), through the nonnegative integrality terms on the right; (ii) violations of block isolation — the clustering cross-terms of the master trace formula, which the identity does not constrain, and which Theorem 3.9 shows to be adversary-tunable at zero cost.
3. The premise “a hyperbolic pair block contributes $(+a)^3 + (-a)^3 = 0$ to $\text{tr } G^3$ while a double contributes $+8$ ” is **FALSE** under the true block law: the pair’s signed cubic is $2m^3(1+3A^2) \geq 8m^3$ at every depth. The sign-split “different corners of the constraint polytope” picture is superseded, as already recorded in the direction file.

Proof. Sum the per-block identities. An atom of mark m has block charges (mass m , Frobenius m^2 , cubic m^3), contributing $m^3 - 3m^2 + 2m = m(m-1)(m-2)$ to $C - 3F + 2\text{Mass}$. A pair of mult m has block charges (mass $2m$, Frobenius $2m^2(1+A^2)$, cubic $2m^3(1+3A^2)$), contributing

$$2m^3(1+3A^2) - 6m^2(1+A^2) + 4m = 2m(m-1)(m-2) + 6m^2(m-1)A^2,$$

exactly its two zeros’ worth of the integrality sum plus the multiplicity-weighted depth term. On marks $\{1,2\}$ and mult-1 pairs every term vanishes. Consequence 3 is Proposition 3.2 ($A \geq 1$). $\square$

Remark (why the gate had to be clustered). By consequence 1, an isolated-block gate over the doubles-vs-pairs class returns a tautological answer along the plane $c = 3F - 2$. The corrected gate’s clustered null and positional freedom were therefore not conservatism but necessity — and the decided outcome is that the clustering cross-terms, the only live channel, are a free resource for the adversary rather than a signal for the certificate.

Numerical confirmation (`verify_t3.py`): charges and the per-zero identity for $m \in \{1,2,3,5\}$ at all depths to $\leq 9.1 \times 10^{-13}$; atom identity exhaustive $m = 1, \dots, 29$, exact; master-formula 2×2 kernel route reproduces all three charges.

3.3 Sine-process closed forms and the doubles-plane diagnostic

The gate's null budgets are the per-zero moments of the flat-window Gram over the determinantal sine process — the process with kernel $S(x) = \sin(\pi x)/(\pi x)$, intensity 1, and correlation functions (taken as the null's definition)

$$\begin{aligned}\rho_2(x_1, x_2) &= 1 - S(x_{12})^2, \\ \rho_3(x_1, x_2, x_3) &= 1 - S(x_{12})^2 - S(x_{13})^2 - S(x_{23})^2 + 2S(x_{12})S(x_{13})S(x_{23}).\end{aligned}$$

The per-zero Gram moments at bandwidth λ (flat window, $\psi = \psi_\lambda$, $u = \hat{\psi}$):

$$\begin{aligned}m_2(\lambda) &:= 1 + \int \rho_2(0, x) \psi(x)^2 dx, \\ m_3(\lambda) &:= 1 + 3 \int \rho_2(0, x) \psi(x)^2 dx + T_3(\lambda), \\ T_3(\lambda) &:= \iint \rho_3(0, x, x+y) \psi(x) \psi(y) \psi(x+y) dx dy,\end{aligned}$$

the $N \rightarrow \infty$ limits of $E[\text{tr } G^k]/N$: the k -tuple sum of Section 2.2 splits over coincidence patterns — all-equal (value 1), exactly-two-equal (3 ordered patterns at $k = 3$, each $\psi(x)^2$ against ρ_2), all-distinct (ρ_3 against the cyclic kernel product, reparametrized by separations using the evenness of ψ). The sine process is simple, so all marks are 1.

Theorem 3.5 (flat-window closed forms). *For the flat window and every $0 < \lambda \leq 1$:*

$$m_2(\lambda) = \frac{1}{\lambda} + \frac{\lambda}{3}, \quad m_3(\lambda) = 1 + \frac{1}{\lambda^2}.$$

Proof. Step 1 (Fourier-side reduction). With $t(\alpha) = (1 - |\alpha|)_+$ (the transform of S^2), $b = \mathbf{1}_{[-1/2, 1/2]}$ ($= \hat{S}$), and the convention $\hat{f}(\alpha) = \int f(x) e^{-2\pi i \alpha x} dx$:

- (i) $\int \psi^2 dx = \int u^2 d\alpha$ [Plancherel];
- (ii) $\int S^2 \psi^2 dx = \int t(u * u) d\alpha$ [Plancherel on the product pair];
- (iii) $\iint \psi(x) \psi(y) \psi(x+y) dx dy = \int u^3 d\alpha$;
- (iv) $\iint S(x)^2 \psi(x) \psi(y) \psi(x+y) dx dy = \int (t * u) u^2 d\alpha$, and the same value for the $S(y)^2$ and $S(x+y)^2$ variants;
- (v) $\iint S(x) S(y) S(x+y) \psi(x) \psi(y) \psi(x+y) dx dy = \int (b * u)^3 d\alpha$.

For (iii), two Plancherel steps with no delta calculus: with y fixed, $g(y) := \int \psi(x) \psi(x+y) dx$ is the Fourier transform of u^2 at y ; then $\int \psi(y) g(y) dy = \int \hat{\psi} u^2 = \int u^3$ by the multiplication identity $\int f \hat{g} = \int \hat{f} g$; Fubini is licensed by $|\psi(x) \psi(y) \psi(x+y)| \leq C/((1+|x|)(1+|y|)(1+|x+y|))$, integrable on $\mathbf{R}^2$. For (iv), the same two-step Plancherel with first factor $S(x)^2 \psi(x)$ gives $\int (S^2 \psi) \hat{u}^2 = \int (t * u) u^2$; the $S(y)^2$ and $S(x+y)^2$ variants reduce to the same value under the substitutions $(x, y) \mapsto (y, x)$ and $(x, y) \mapsto (x, -x-y)$ (Jacobian 1, integrand invariant). For (v), set $h = S\psi$, $\hat{h} = b * u$, and apply (iii) to h . Hence

$$\begin{aligned}m_2 &= 1 + \int u^2 - \int t(u * u), & m_3 &= 1 + 3 \left[\int u^2 - \int t(u * u) \right] + T_3, \\ T_3 &= \int u^3 - 3 \int (t * u) u^2 + 2 \int (b * u)^3.\end{aligned}$$

Step 2 (evaluation at $u = (1/\lambda) \mathbf{1}_{[-\lambda/2, \lambda/2]}$, using $\lambda \leq 1$ exactly where indicated).

- (1) $\int u^2 = 1/\lambda$.
- (2) $(u * u)(\alpha) = (1/\lambda^2)(\lambda - |\alpha|)_+$; since $\lambda \leq 1$, $t = 1 - |\alpha|$ on this support, so $\int t(u * u) = (2/\lambda^2) \int_0^\lambda (1-a)(\lambda-a) da = (2/\lambda^2)[\lambda^2/2 - \lambda^3/6] = 1 - \lambda/3$. Hence $m_2 = 1 + 1/\lambda - (1 - \lambda/3) = 1/\lambda + \lambda/3$.
- (3) $\int u^3 = 1/\lambda^2$.
- (4) $\int (t * u)u^2 = (1/\lambda^3) \iint_{|\alpha| \leq \lambda/2, |\alpha-\beta| \leq \lambda/2} t(\beta) d\beta d\alpha = (1/\lambda^3) \int t(\beta)(\lambda - |\beta|)_+ d\beta = (1/\lambda^3)[\lambda^2 - \lambda^3/3] = 1/\lambda - 1/3$ (reusing the integral of (2); valid for $\lambda \leq 1$).
- (5) $(b * u)(\alpha)$ is the trapezoid equal to 1 on $|\alpha| \leq (1-\lambda)/2$, decaying linearly to 0 at $|\alpha| = (1+\lambda)/2$; hence $\int (b * u)^3 = (1-\lambda) + 2\lambda \int_0^1 s^3 ds = 1 - \lambda/2$.

Assembling: $T_3 = 1/\lambda^2 - 3(1/\lambda - 1/3) + 2(1 - \lambda/2) = 1/\lambda^2 - 3/\lambda + 3 - \lambda$, and $m_3 = 1 + 3[1/\lambda - 1 + \lambda/3] + T_3 = 1 + 1/\lambda^2$. $\square$

Corollary 3.6 (anchor values, the doubles-plane diagnostic, the moment margin).

$$m_2(1) = 4/3, \quad m_3(1) = 2, \quad m_2(1/2) = 13/6, \quad m_3(1/2) = 5.$$

Define $G(\lambda) := m_3 - (3m_2 - 2)$ (the null budget's excess over the isolated-block doubles/pairs plane of Theorem 3.4) and $\text{Marg}(\lambda) := 2m_2 - m_3$ (the Section-7.3-style certificate margin). Then, flat window, on $(0, 1]$:

$$\begin{aligned} G(\lambda) &= 3 + \frac{1}{\lambda^2} - \frac{3}{\lambda} - \lambda; & G(1) &= 0, & G(1/2) &= 1/2, \\ \text{Marg}(\lambda) &= \frac{2}{\lambda} + \frac{2\lambda}{3} - 1 - \frac{1}{\lambda^2}; & \text{Marg}(1) &= 2/3, & \text{Marg}(1/2) &= -2/3, \end{aligned}$$

and Marg has a unique sign change on $(0, 1]$ (the interval on which the closed forms are proved), at $\lambda_0 = 0.610511\dots$. Each value is load-bearing for the no-go:

1. $G(1) = 0$ **is the “2 = 2” saturation.** At bandwidth one the sine budget $m_3(1) = 2$ exactly equals the isolated-block cubic demand of the 5/6-extremal (2/3 simple + 1/6 doubles: (2/3)(1) + (1/3)(4) mass-weighted = 2, on the plane of Theorem 3.4). The bandwidth-one cubic row alone cannot cut the doubles corner — the budget is exactly met.
2. $G(1/2) = +1/2 > 0$. At the operating bandwidth the null budget sits strictly ABOVE the doubles plane: a doubles-saturated adversary must supply +1/2 per zero of cubic through positional cross-terms — which Theorem 3.9's grid freedom supplies at zero cost (the gate's witness does exactly this, sitting at the LOWER cubic budget edge).
3. $\text{Marg}(1/2) = -2/3 < 0$. The $\omega(m)$ -moment route's margin is negative at the proven operating point: a bite could never have come from the Section-7.3 mechanism there, only from joint positional pricing. The sign change at $\lambda_0 = 0.6105$ is precisely where the gate's exploratory scan saw δ_0 turn positive (between $\lambda' = 0.55$ and 0.60 ; Section 8.4), outside the proven ladder regime.

Proof. Substitute Theorem 3.5; the displayed formulas are algebraic identities on $(0, 1]$. $G(1) = 3 + 1 - 3 - 1 = 0$; $G(1/2) = 3 + 4 - 6 - 1/2 = 1/2$; $\text{Marg}(1) = 2 + 2/3 - 1 - 1 = 2/3$; $\text{Marg}(1/2) = 4 + 1/3 - 1 - 4 = -2/3$. Uniqueness of the root: $q(\lambda) = \lambda^2 \text{Marg}(\lambda) = 2\lambda + (2/3)\lambda^3 - \lambda^2 - 1$

has $q' = 2 + 2\lambda^2 - 2\lambda > 0$ (negative discriminant), so q is strictly increasing with $q(0.55) < 0 < q(0.66)$; numerically the root is 0.610511 (`brentq`, re-verified by `sympy nsolve`). The extremal demand in item 1: per-zero $(F, c) = (1, 1)$ on simples and $(2, 4)$ on doubles; mass-weighted cubic $(2/3)(1) + (1/3)(4) = 2 = m_3(1)$ and Frobenius $(2/3)(1) + (1/3)(2) = 4/3 = m_2(1)$ — both budgets exactly saturated at $\lambda = 1$. $\square$

Corollary 3.7 (the $m_2 = \kappa$ identity: sine null = unconditional budget). *The closed form $m_2(\lambda) = 1/\lambda + \lambda/3$ coincides, at every $\lambda \leq 1$, with the value obtained by integrating the unconditional Montgomery/[BGSTB24] pair-correlation form factor $F(\alpha) = \delta_0(\alpha) + |\alpha|$ (on $|\alpha| \leq 1$) against the flat-window pair kernel:*

$$\frac{1}{\lambda^2} \int_{-\lambda}^{\lambda} (\lambda - |\alpha|) [\delta_0(\alpha) + |\alpha|] d\alpha = \frac{\lambda + \lambda^3/3}{\lambda^2} = \frac{1}{\lambda} + \frac{\lambda}{3} = m_2(\lambda).$$

*The sine-process null's second-moment budget IS the parent paper's unconditional prime-side $\kappa(\lambda)$, exactly, on the whole band $\lambda \leq 1$ — the strongest possible identity check at second order, and the reason the gate's F' budget row is unconditional-data-backed. (Provenance: the prime-side member — $\text{tr } \hat{G}^2 \leftrightarrow \int (\lambda - |\alpha|) F(\alpha) d\alpha$ with F unconditional and pointwise on the closed band $0 \leq \alpha \leq 1$ — is the parent paper's Theorem 5.8 / Remark 5.10 bookkeeping together with [BGSTB24] Theorem 1, quoted, not re-derived; the displayed arithmetic identity is the elementary integral $\int_0^\lambda (\lambda - a) a da = \lambda^3/6$, doubled, plus the delta term λ .) **No third-order analog is claimed:** the identification $c_3(\lambda') = m_3(\lambda')$ rests on Rudnick–Sarnak-range GUE 3-level correlations and remains an open flag (F2; Section 8.3).*

Numerical confirmation (`verify_t4.py`, `verify_t4_mc.py`): Fourier-side quadrature matches both closed forms at 7 bandwidths to $\leq 1.7 \times 10^{-5}$; an independent real-space determinantal quadrature gives $m_3(1/2) = 4.9924$ (truncation $X = 40$) $\rightarrow 4.9960$ ($X = 80$), trending to 5 as the audit's independent quadrature did ($4.9896 \rightarrow 4.9932$); a fresh CUE Monte Carlo (third independent sampler) extrapolates $m_2(1/2) \rightarrow 2.1668$ and $m_3(1/2) \rightarrow 5.0014$ against $13/6$ and 5, each within $\sim 1\sigma$ of both the gate's and the auditor's samplers (full tables: Appendix B). The values $m_2(1) = 4/3$, $m_3(1) = 2$ also discharge, for $k \leq 3$, the parent's recalled sine moments $m_k(1) = 1, 4/3, 2$.

3.4 Grid Parseval decoupling

The absorption mechanism rests on one exact identity: on a specific uniform grid the entire bandwidth-one Frobenius row degenerates to the multiplicity count — for every site subset and every mark assignment — while the half-band kernel on the same grid remains position-sensitive.

Lemma 3.8 (flat-band lemma). *Let $M \geq 1$ and let the harmonic band be ANY set of M consecutive integers $B = \{j_0, \dots, j_0 + M - 1\}$, with uniform weights $u_j = 1/M$. Let the configuration be supported on the M -site uniform grid $\theta_k = kN/M$ ($k = 0, \dots, M-1$) on the circle of circumference N , with arbitrary real marks $m_k \geq 0$. Then*

$$\text{tr } \hat{G}^2 = \sum_k m_k^2 \quad \text{EXACTLY.}$$

Proof. On the grid the form factor is a DFT on $\mathbf{Z}_M$: $c_s = \sum_k m_k e^{-2\pi i s k / M}$, which depends only on $s \bmod M$ — c is M -periodic in s — and DFT Parseval gives $\sum_{r \bmod M} |c_r|^2 = M \sum_k m_k^2$ (P). The Frobenius row is $\sum_s W_2(s) |c_s|^2$ with $W_2(s) = \#\{(j_1, j_2) \in B \times B : j_1 + j_2 = s\} / M^2$; the sum-set

$B + B = \{2j_0, \dots, 2j_0 + 2M - 2\}$ carries the triangular pair counts $M - |s - (2j_0 + M - 1)|$ on $|s - (2j_0 + M - 1)| \leq M - 1$. Fix a residue class $r \bmod M$: the window $B + B$ has length $2M - 1$, so it contains either one member of the class ($t = 0$, count M) or two members t and $t \mp M$ with $0 < |t| < M$, whose counts sum to $(M - |t|) + |t| = M$ — EXACTLY M on every residue class. Hence, by the M -periodicity of $|c_s|^2$,

$$\text{tr } \widehat{G}^2 = \frac{1}{M^2} \sum_{r \bmod M} M |c_r|^2 = \frac{1}{M} \sum_r |c_r|^2 = \sum_k m_k^2$$

by (P). □

Theorem 3.9 (grid Parseval decoupling at bandwidth one). *Let N be even, flat window at $\lambda = 1$: band $B = \{|j| \leq N/2\}$, $M = N + 1$ harmonics, $u_j = 1/(N + 1)$. Let the configuration be supported on the $(N + 1)$ -site uniform grid of spacing $N/(N + 1)$ (the ψ_1 -zero grid), with arbitrary marks. Then*

$$F_1 = \text{tr } \widehat{G}_1^2 = \sum_z m_z^2 \quad \text{EXACTLY,}$$

*for every site subset and every mark assignment: the bandwidth-one reading on this grid sees only multiplicities, never positions, and the **lemmaR_tight** extremal analysis transfers verbatim to grid configurations.*

Proof. Lemma 3.8 with $M = N + 1$, $B = \{-N/2, \dots, N/2\}$ ($N + 1$ consecutive integers since N is even), grid spacing $N/M = N/(N + 1)$. □

Nomenclature. Lemma 3.8 is critical-sampling Parseval for a flat band — the discrete orthogonality of M consecutive characters on an M -site uniform grid — and Proposition 3.11 is its Shannon–Whittaker continuum form on the integer lattice. Both are classical; they are proved here in the exact finite-circle normalization the gate uses, and the Lean development machine-checks Lemma 3.8 over an arbitrary integral domain (Section 10).

Remark (why this grid). Equivalently: all pairwise site differences are nonzero multiples of $N/(N + 1)$, the zero set of the discrete bandwidth-one kernel D . The Fourier proof is the one that makes the exactness structural — assembly weights constant on residue classes — rather than a numerical coincidence, and it is the form queued for Lean.

Proposition 3.10 (the $\lambda' = 1/2$ kernel is alive on the same grid). *Same geometry, N and $N/2$ even, $\lambda' = 1/2$: band $B' = \{|j| \leq N/4\}$, $M' = N/2 + 1$. For grid-supported configurations,*

$$F' = \sum_{|s| \leq N/2} \frac{M' - |s|}{M'^2} |c_s|^2,$$

where $\{-N/2, \dots, N/2\}$ is a COMPLETE residue system mod $(N + 1)$ and c_s is the $\mathbf{Z}_{N+1}$ DFT of the mark vector. The weights are strictly positive and non-constant across residues (triangular: at $N = 64$, from $1/33$ at $s = 0$ down to $1/33^2$ at $|s| = 32$), so F' is a strictly position-sensitive functional of the grid configuration, subject only to Parseval.

Proof. $F' = \sum_s W'_2(s) |c_s|^2$ with $W'_2(s) = (M' - |s|)_+/M'^2$ supported in $|s| \leq N/2$. That range has $N + 1$ elements, hence meets each residue class mod $N + 1$ exactly once; no residue-class telescoping occurs (band length $M' = N/2 + 1$ differs from the grid size $N + 1$), so F' reads the full grid power spectrum with a non-uniform weight; distinct site subsets of equal mark statistics generically differ. □

Worked exact value (hand-checkable; the witness’s first column). $N = 64$, the “vacancy lattice”: 64 simple marks on 64 of the 65 grid sites. Then $c_0 = 64$ and $|c_s|^2 = 1$ for $s \not\equiv 0 \pmod{65}$, so

$$F_1 = 64 \quad (\text{Theorem 3.9}),$$

$$F' = \frac{33}{33^2} 64^2 + \sum_{0 < |s| \leq 32} \frac{33 - |s|}{33^2} = \frac{4096}{33} + \frac{32}{33} = \frac{4128}{33} = 125.0909 \dots,$$

using $\sum_{|s| \leq 32} (33 - |s|) = 33^2$ and subtracting the $s = 0$ term. This reproduces the shipped witness column exactly (`witness_N64.json`: $F_1 = 64$, $F' = 125.09090 \dots$; independently hand-derived in the audit). The two-bandwidth decoupling is thereby explicit: F_1 is blind to the vacancy’s position, F' is not.

Proposition 3.11 (continuum analog: the adversary lattice). *In the continuum model, flat window, configuration supported on the integer lattice $\mathbf{Z}$ with arbitrary real marks: $\text{tr } G_1^2 = \sum m_z^2$ exactly, while the half-band kernel is alive on the lattice: $\psi_{1/2}(k) = \text{sinc}(\pi k/2) = (2/(\pi k))(-1)^{(k-1)/2}$ for odd k (0 for even $k \neq 0$). For the finite- T Weil–Gabor matrix itself the identity holds as $\text{tr } \widehat{G}^2 = (\sum m^2)(1 + o(1))$, the $o(1)$ being the parent’s finite-window corrections (quoted; the campaign’s recorded “adversary lattice”).*

Proof (model part). At $k = 2$ the master formula gives $\text{tr } G_1^2 = \sum_{z, z'} m_z m_{z'} \psi_1(z - z')^2$; $\psi_1(x) = \text{sinc}(\pi x)$ vanishes at every nonzero integer and equals 1 at 0, so only the diagonal survives. The half-band values are $\sin(\pi k/2)/(\pi k/2)$ with $\sin(\pi k/2) = (-1)^{(k-1)/2}$ for odd k . $\square$

Remark. Theorem 3.9 is the exact periodization of Proposition 3.11: spacing $N/(N + 1)$ instead of 1 makes the identity exact rather than asymptotic — which is precisely why the finite- N gate could exhibit $\delta_0 = 0$ as an exact LP equality rather than a numerical near-zero. The absorption is not a finite- N artifact.

Numerical confirmation (`verify_t1.py`): grid exactness on 40 random site-subsets with random marks at $N = 64$ and $N = 128$, two independent evaluation routes (Fourier double sum; eigenvalues of H): $\max |F_1 - \sum m^2| = 9.1 \times 10^{-13} / 4.1 \times 10^{-12}$, matching the audit’s independent 4.5×10^{-13} ; general flat-band lemma on 20 random (N, M, j_0) draws including asymmetric bands: 3.2×10^{-12} ; controls — off-grid configurations give $|F_1 - \sum m^2| \in [1.3 \times 10^{-2}, 62.5]$ and on-grid half-band $|F' - \sum m^2| \in [40.9, 120.8]$ (the identity genuinely fails off the grid, and the half-band kernel is genuinely alive on it); continuum lattice exact to machine precision.

4 The absorption theorem

This section states and proves the model-level theorem the gate pins — the two-moment data force exactly the `lemmaR_tight` corner, for both benchmarks, over the honest configuration classes — and then presents the computational certification that the entire cubic block is absorbed at that corner, with the explicit witness.

4.1 The atom-only corner theorem

Fix N and $\varepsilon \geq 0$, and let $\mathcal{A}$ be the class of atom-only configurations: atoms at arbitrary positions (finite-circle or continuum model, flat windows — the proofs work verbatim in both), integer marks $m_i \geq 1$, mass $\sum m_i = N$ per period.

Lemma 4.1 (diagonal positivity). *For every atom-only configuration (any positions, any real marks $m_i \geq 0$), in either model, $F_1 \geq \sum_i m_i^2$, with equality iff every off-diagonal kernel value vanishes — in particular on the grid of Theorem 3.9 and the lattice of Proposition 3.11.*

Proof. $F_1 = \sum_{i,i'} m_i m_{i'} K(\theta_i - \theta_{i'})$ with $K = D^2$ (finite circle) or $K = \psi_1^2$ (continuum): $K \geq 0$ pointwise, $K(0) = 1$, so $F_1 = \sum m_i^2 +$ (a sum of nonnegative off-diagonal terms). $\square$

This is the one step that fails for pair-containing configurations — an off-line pair contributes cross-terms of the form $2mm' \operatorname{Re}[\psi(x+iy)^2]$, which need not be nonnegative; Sections 4.2–4.3 treat that channel.

Lemma 4.2 (integer-mark corner inequalities). *For every atom-only configuration with integer marks $m_i \geq 1$ and mass N :*

$$\begin{aligned} (a) \quad N_d = \#\text{atoms} &\geq \frac{3N - \sum_i m_i^2}{2}, \\ (b) \quad n_1 = \#\text{mark-1 atoms} &\geq 2N - \sum_i m_i^2, \end{aligned}$$

each with equality iff all marks lie in $\{1, 2\}$.

Proof. Per-atom inequalities summed over atoms. (a) For integer $m \geq 1$, $(m-1)(m-2) \geq 0$, i.e. $1 \geq (3m - m^2)/2$, equality iff $m \in \{1, 2\}$; sum. (b) $\mathbf{1}_{m=1} \geq 2m - m^2$: equality at $m = 1, 2$; at $m \geq 3$ the right side is negative; sum. $\square$

Remark. (a) is the second integrality level $(m-1)(m-2) \geq 0$ — exactly the level `lemmaR.tight` proves exhausted by the two-moment certificate, and, by Theorem 3.4, exactly the level the cubic row re-measures on isolated blocks; (b) is its simple-fraction sibling. In the parent’s terms, (a) is the doubles-optimal corner arithmetic: an atom of mark m spends $m(m-1)$ of Frobenius excess to save $m-1$ units of N_d , so doubles are the efficient coin.

Theorem 4.3 (atom-only corner theorem, both benchmarks). *Fix N even, $0 \leq \varepsilon \leq 1/2$ (finite-circle model, flat windows). Over laws w on $\mathcal{A}$ with mass N and $E_w[F_1] \leq (4/3)N(1 + \varepsilon)$:*

$$\begin{aligned} (a) \quad \min_w E_w[N_d]/N &= \frac{5}{6} - \frac{2}{3}\varepsilon, \\ (b) \quad \min_w E_w[p_1] &= 2 - \frac{4}{3}(1 + \varepsilon) = \frac{2}{3} - \frac{4}{3}\varepsilon. \end{aligned}$$

Both minima are attained by explicit laws supported on the $(N+1)$ -site grid of Theorem 3.9 with marks $\{1, 2\}$ only. Every configuration in $\mathcal{A}$ satisfies the pointwise versions with its own Frobenius excess ε_c . The same statements hold with the two-sided budget band $(4/3)N[1 - \varepsilon, 1 + \varepsilon]$ (the attaining laws sit at the upper edge), and the minima over any richer mark alphabet (any W , or unbounded marks) are the same.

Proof. Lower bounds: by Lemma 4.1, $E_w[\sum m^2] \leq E_w[F_1] \leq (4/3)N(1 + \varepsilon)$; by Lemma 4.2(a) and linearity, $E_w[N_d] \geq (3N - (4/3)N(1 + \varepsilon))/2 = N(5/6 - (2/3)\varepsilon)$; by Lemma 4.2(b), $E_w[n_1] \geq 2N - (4/3)N(1 + \varepsilon) = N(2/3 - (4/3)\varepsilon)$.

Attainment: for integer $0 \leq k \leq N/2$ let c_k be the grid configuration with k doubles and $N-2k$ simples on $N-k$ distinct grid sites (which exist: $N+1 \geq N-k$). By Theorem 3.9, $F_1(c_k) = N+2k$

exactly; $N_d(c_k) = N - k$; $n_1(c_k) = N - 2k$. Set $\bar{k} = (B - N)/2$ with $B = (4/3)N(1 + \varepsilon)$; then $0 \leq \bar{k} \leq N/2$ for $\varepsilon \leq 1/2$. The two-column law w^* mixing $c_{[\bar{k}]}$ and $c_{[\bar{k}]+1}$ with $E[k] = \bar{k}$ achieves

$$E[F_1] = N + 2\bar{k} = B \quad (\text{upper edge, exactly}),$$

$$\frac{E[N_d]}{N} = \frac{3N - B}{2N} = \frac{5}{6} - \frac{2}{3}\varepsilon, \quad E[p_1] = 2 - \frac{B}{N} = \frac{2}{3} - \frac{4}{3}\varepsilon,$$

and both lower-bound chains hold with equality (marks in $\{1, 2\}$; grid kills all cross-terms; budget saturated), so w^* is simultaneously optimal for both objectives. The two-sided band adds only the lower budget edge, which w^* satisfies. Enlarging the mark alphabet cannot lower the minimum (the chain never used a mark cap) and cannot raise it (w^* uses marks $\{1, 2\}$, inside every alphabet): the W -independence is inclusion, not extrapolation. $\square$

Corollary 4.4 (asymptotic and matched corners). *As $\varepsilon \rightarrow 0$ the two optima tend to $5/6$ and $2/3$ — the model’s bandwidth-one analogs of the Montgomery extremal values for distinct zeros and simple fraction. With the gate’s matched-geometry budget B_1 replacing $(4/3)N$, the same proof gives $\min E[N_d]/N = (3N - B_1(1 + \varepsilon))/(2N)$ and $\min E[p_1] = 2 - (B_1/N)(1 + \varepsilon)$; at the shipped matched values ($N = 64$, $\varepsilon = 0.05$, $B_1 = 84.71214548308222$) these are 0.8050957 and 0.6101914 — the run’s and follow-up’s reported optima to LP precision.*

Remark (a numerical coincidence, disambiguated). The corner value 0.8050957 rounds to 0.8051 at four decimals, and the string 0.8051 also occurs in the zeta literature as Farmer–Gonek–Lee’s RH-conditional pair-correlation lower bound for the proportion of **distinct** zeros of zeta, $N_d(T) > 0.8051 N(T)$ [FGL14, Cor. 1.4] (since superseded by 0.84665 in [BHB13] and by 0.8477 on RH / 0.8486 on GRH in [CGdL20, Cor. 3]). The two are unrelated: the value here is a finite- N LP corner of the bandwidth-one model at $N = 64$, $\varepsilon = 0.05$, $B_1 = 84.71214548308222$, and carries no information about zeta.

Numerical confirmation (`verify_t2.py`): per-atom inequalities exhaustive to $m = 50$ plus 20,000 random draws (0 violations); grid-column LP matches the closed forms at $\varepsilon \in \{0.10, 0.05, 0.02, 0.002\}$ to $\leq 3.3 \times 10^{-16}$ for both objectives; exact rational identities verified in `Fraction` arithmetic; the shipped witness reproduces $P = (3N - E[F_1])/(2N) = 0.8050957$ to 6×10^{-13} .

4.2 Integrality is necessary: the exact fractional counterexample

The extension of Theorem 4.3 beyond atoms must consume integrality — provably. Work in the gate’s primary geometry ($N = 64$, flat window, $\lambda = 1$; kernel $\psi(z) = \sum_j u_j e^{-2\pi i j z/N}$, grid $g_k = k 64/65$) and define $\bar{a}(x) := \psi(ix) = \sum_j u_j \cosh(2\pi j x/N) \geq 1$. Write $M(c)$ for total mark mass, $S_2(c) = \sum_z m_z^2$ (each pair member counted m^2), $T(c) = 3M(c) - 2N_d(c)$, and consider the **master inequality**

$$(MI) \quad F_1(c) \geq 3M(c) - 2N_d(c),$$

which is precisely what the corner consumes: (MI) per configuration gives, for every law with mass N and $E[F_1] \leq (4/3)N(1 + \varepsilon)$, BOTH corners — $E[N_d]/N \geq 5/6 - (2/3)\varepsilon$ and $E[p_1] \geq 2/3 - (4/3)\varepsilon$. (Per configuration (MI) reads $N_d \geq (3M - F_1)/2$; take expectations. For simples, $\mathbf{1}_{m=1} \geq 2 - m$ summed over the zero multiset gives $n_1 \geq 2N_d - M$, hence $E[n_1] \geq (3N - E[F_1]) - N$.) Note $S_2 \geq T$ for integer marks (per zero, $m^2 - (3m - 2) = (m - 1)(m - 2) \geq 0$), so (MI) is weaker than $F_1 \geq S_2$ — and aiming at (MI), which carries the integrality slack explicitly, is what makes the closure close.

Proposition 4.5 (fractional marks break the floor at every depth). *Let c_μ be the vacancy lattice (unit atoms on grid sites $g_1, \dots, g_{64}$) plus one pair at the hole g_0 with depth $d > 0$ and REAL mark $\mu > 0$. Then, exactly,*

$$F_1(c_\mu) - S_2(c_\mu) = 2\mu^2 \bar{a}(2d)^2 - 4\mu (\bar{a}(d)^2 - 1),$$

which is NEGATIVE for every $0 < \mu < 2(\bar{a}(d)^2 - 1)/\bar{a}(2d)^2$ — the pair harvests the hole’s missing coupling linearly in μ while paying only quadratically. For integer marks the same family is safe: at $\mu = m \in \mathbf{Z}$, the log-convexity bound $\bar{a}(d)^2 - 1 \leq (\bar{a}(2d) - 1)/2$ gives $F_1 - S_2 \geq 2m^2 \bar{a}(2d)^2 - 2m(\bar{a}(2d) - 1) > 0$.

Proof. On the vacancy lattice the atom form factor is $\varphi_0(s) = 64$ at $s \equiv 0$ and -1 otherwise; the pair adds $2\mu \cosh(\beta s)$, $\beta = 2\pi d/N$. Expanding $F_1 = \sum_s w_s |c_s|^2$ with the generating identities $\sum_s w_s \cosh(\beta s) = \bar{a}(d)^2$ and $\sum_s w_s \cosh(2\beta s) = \bar{a}(2d)^2$ (the assembly weights $w = u * u$ factor through ψ at imaginary arguments) gives the display; the numeric check at $(d, \mu) = (0.25, 0.05)$ reproduces it to 10^{-12} , with $F_1 - S_2 = -3.520 \times 10^{-2}$. The integer case uses Cauchy–Schwarz on the positive weights u_j ($\bar{a}((x+y)/2)^2 \leq \bar{a}(x)\bar{a}(y)$, hence $\bar{a}(d)^2 \leq (1 + \bar{a}(2d))/2$). $\square$

Reading. An adversary with fractional marks takes $\mu \rightarrow 0$ and wins at any depth; integer marks force $\mu \geq 1$, where the quadratic cost dominates. Any closure of the pair channel must therefore consume integrality — the channel is invisible to every convexified or soft-positivity analysis. (This is the textbook **integrality gap** in its usual form: the fractional relaxation of the marked-configuration program has a strictly smaller optimum than the integer program, so a bound proved for the relaxation is simply not a bound for the problem.) This also explains the gate’s LP phenomenology: the LP’s columns are explicit integer configurations (nothing about marks is relaxed), which is why its converged pricing found no pair column. And it is the sharp form of the realizability discussion of Section 2.4: with fractional marks even the 5/6 baseline is false.

4.3 The pair-interference channel: closed on the R5 family (multi-pair to $w \leq 0.98$), backstopped everywhere

The full closure argument, with its capacity machinery and certified constants, is the companion note (*supplementary material*, `pair-channel.md`); we state the results, the exact ledger they rest on, and the proofs that are short enough to give in full. Everything is at the gate’s decision geometry ($N = 64$, flat window, $\lambda = 1$); depths d are in circle units, $w = 2\pi d$ dimensionless (the gate’s pair family is $w \in (0, 1]$ plus deep probes $w \in \{1.5, 2\}$, i.e. $d \leq 0.3183$).

The exact interference ledger. For every admissible configuration (any atoms, any pairs, any integer marks), with $R_y(x) := \text{Re } \psi(x + iy)^2$:

$$\begin{aligned}
F_1 - T &= \sum_a (m_a - 1)(m_a - 2) + 2 \sum_p (m_p - 1)(m_p - 2) && \text{[integrality slack]} \\
&+ \sum_{a \neq a'} m_a m_{a'} \psi(x_{aa'})^2 && \text{[atom-atom} \geq 0] \\
&+ \sum_p 2m_p^2 \bar{a} (2d_p)^2 && \text{[pair diagonals]} \\
&+ \sum_p 4m_p \sum_a m_a R_{d_p}(x_{pa}) && \text{[pair-atom]} \\
&+ \sum_{p < q} 4m_p m_q [R_{d_p+d_q}(x_{pq}) + R_{|d_p-d_q|}(x_{pq})] && \text{[pair-pair]}
\end{aligned}$$

(verified on 25 random mixed configurations to 2.2×10^{-11} , the stored `pairchan_verify.out.json` maximum). Everything on the right is nonnegative EXCEPT the R -couplings, whose negative parts live only in the dips of R_y ; the question is whether the dips can outrun the pair diagonals plus the integrality slack. Quantitatively they cannot — by a factor of about 1.5 at the worst depth:

Theorem 4.6 (single-pair closure). *Every configuration with exactly one pair (arbitrary atoms, arbitrary integer marks, arbitrary positions) satisfies (MI) whenever the pair depth is $d \leq 0.45$ — i.e. $w \leq 2.8$, covering the gate’s ENTIRE pair-depth family including both deep probes, with $\geq 34\%$ capacity margin on the family itself (certified stacking-corrected capacity ratio ≤ 0.66 on the family, ≤ 0.921 out to $d = 0.45$). For mass $\leq N$ configurations (the LP class) (MI) also holds for every $d \geq 1.0$; the window $(0.45, 1.0)$ is certified numerically (raw capacity ratio ≤ 0.6823 throughout; direct adversarial searches reach at most 74% of the pair budget).*

Theorem 4.7 (multi-pair closure). *(MI) holds for every configuration all of whose pair depths satisfy $w \leq 0.82$, unconditionally; and for all depths $w \leq 0.98$ modulo one isolated cell-crowding cap whose ledger constant is interval-certified ($\sup S_2^{\text{gen}} \leq 0.98465 < 1$ on $(0, 0.156]^2$, per-cell rigorous enclosures; 2026-08-27). On the deepest sliver $w \in (0.98, 1]$ the ledger inequality fails (interval-certified $S_2^{\text{gen}} \geq 1.01405 > 1$ at $d = d' = 0.159$); there the targeted crowd-plus-sea attacks — 8 to 65 pairs, one per cell tranche, plus greedy mark-2 seas up to 130 atoms, re-run at $d = 0.158$ and 0.159 — never bring $F_1 - T$ below $+17.46$, against a floor requirement of 0, and the unconditional 8/9 backstop applies.*

Proof architecture (full details in the companion note). Partition the circle into 65 kernel-zero cells; per cell, the exploitable dip mass $\kappa_k(y) = \sup_{\text{cell}} [-R_y]_+$ sums to the capacity curve $\nu(y)$, with the exact curvature identity $\alpha := \sum_j u_j (2\pi j/N)^2 = \sum_k \psi'(g_k)^2 = 3.392677$ controlling the shallow regime ($\nu(y) = \alpha y^2 (1 + o(1))$). Integer atoms harvesting a cell’s dip pay in-cell atom-atom crosses ($\geq \psi(\text{dip-zone diameter})^2$ each) and their own integrality slack, capping the harvest at $8m_p \kappa_k$ per cell for marks ≤ 2 (with a certified stacking correction beyond); on the R5 family the closing inequality $8\nu(y) \leq 2\bar{a}(2y)^2$ follows in closed form from one harmonic-Taylor bound with the exact α and one certified finite constant (1.2% closing margin at the endpoint). Pair-pair couplings are priced through the joint kernel $R_{d+d'} + R_{|d-d'|}$ (never the two parts separately), with a one-pair-per-cell crowding cap that is self-consistent for $w \leq 0.82$ (both sides interval-certified: $2 \max \nu_{\text{joint}} \leq 0.60910 < \Phi_0 \geq 0.64371$) and whose ledger constant beyond it is interval-certified to $w \leq 0.98$; beyond $w = 0.98$ the ledger fails (certified) and the closure defers to numerics plus

the backstop. Deep single pairs ($d \geq 1$) fall to the crude bound $\bar{a}(2d)/\bar{a}(d) \geq 12.158$ nondecreasing. QED (architecture)

Theorem 4.8 (equal-depth pair-only configurations, exact, all depths). *If c consists only of pairs, all at one common depth d (any positions, any integer multiplicities), then $F_1 \geq S_2 + 2 \sum_p m_p^2$ — (MI) with a full doubling-cushion surplus at every depth.*

Proof. $c_s = 2 \cosh(2\pi s d/N) \varphi(s)$ with φ the position-multiset form factor, and $4 \cosh^2 = 2 + 2 \cosh(\text{double})$: F_1 splits as twice the atom-only Frobenius of the position multiset ($\geq 2 \sum_p m_p^2$ by diagonal positivity) plus a termwise-cosh-weighted copy ($\geq$ the same floor). $\square$

Theorem 4.9 (unconditional spectral backstop, no restrictions). *For EVERY admissible configuration — any number of pairs, any depths, any marks — $N_d \geq (4/9)(3M - F_1)$. Hence for any law with mass N and $E[F_1] \leq (4/3)N(1 + \varepsilon)$:*

$$\frac{E[N_d]}{N} \geq \frac{8}{9} \left(\frac{5}{6} - \frac{2}{3} \varepsilon \right) = \frac{20}{27} - \frac{16}{27} \varepsilon = 0.7407 \dots - 0.593 \varepsilon,$$

so the maximal conceivable pair advantage on the corner, in any regime whatever, is $\leq 5/6 - 20/27 = 5/54 = 0.0926$, unconditionally.

Proof. Let B be the Hermitian frequency matrix $B[j, j'] = \sqrt{u_j u_{j'}} c_{j-j'}$ (65×65): exactly $\text{tr } B = M$ and $\text{tr } B^2 = F_1$. B decomposes as $\sum_{\text{atoms}} m_a v_a v_a^\dagger + \sum_{\text{pairs}} B_p$ with each atom block PSD of rank 1. Each pair block is $B_p = m(pq^\dagger + qp^\dagger)$, with $p_j = \sqrt{u_j} e^{-2\pi i j \theta/N} e^{2\pi j d/N}$ and $q_j = \sqrt{u_j} e^{-2\pi i j \theta/N} e^{-2\pi j d/N}$ (the two summands of the conjugate-pair form factor of Section 2.3); on $\text{span}\{p, q\}$ its nonzero eigenvalues are $m(\text{Re}\langle q, p \rangle \pm |p||q|)$, and Cauchy–Schwarz gives $\text{Re}\langle q, p \rangle - |p||q| \leq 0$, so $n_+(B_p) \leq 1$ — the finite-circle counterpart of the signature-(1, 1) block of Proposition 3.2(iv). By subadditivity of positive inertia, $n_+(B) \leq \#\text{atoms} + \#\text{pairs} \leq N_d$. For every real t , $3t - t^2 \leq (9/4)\mathbf{1}_{t>0}$; summing over the spectrum, $3M - F_1 = \sum_i (3\lambda_i - \lambda_i^2) \leq (9/4)n_+(B) \leq (9/4)N_d$. $\square$

Remark. The $9/4$ (rather than 2) is the price of a purely spectral argument — $3t - t^2$ exceeds 2 on $(1, 2)$ with max $9/4$ at $t = 3/2$ — the same structural reason the parent’s inertia route cannot reach $5/6$ by itself. But the backstop is depth-uniform, mark-uniform, and interference-proof, which is exactly what a backstop must be.

Status and the sharp conjecture. Roughly 10^4 adversarially optimized configurations (dip attacks, pair crowds, half-gap frustrated twins, deep seas, random mixed stress; seeds fixed, independent assembly) never brought $F_1 - T$ below $+2.000000$, and the minimum of $F_1 - T - \sum_p 2(2m_p^2 - 3m_p + 2)$ over every family is 0.000000 , attained only in the depth $\rightarrow 0$ grid limit. The searches establish that the binding adversary is always shallow; that coordinated dip attacks can consume nearly the whole depth-dependent excess $2m^2(\bar{a}(2d)^2 - 1)$ (within 3% at $d = 0.1$) but never any of the depth-independent doubling cushion $2m^2$ — exactly the split the capacity lemmas predict. We record the **sharp conjecture** (numerically exhaustive, not proved): $F_1 \geq \sum_a (3m_a - 2) + 4 \sum_p m_p^2$ — every pair pays its full flattened-double cost and interference recovers nothing. Its near-tight witnesses ($+0.03$) show any proof must be capacity-sharp, which is why the theorems above target (MI), with its floor headroom of 2 per pair, instead.

Consequence. The model-level corner theorem extends from the atom-only class to the bulk of the gate’s admissible class: single-pair columns at every gate depth ($w \leq 2.8$), multi-pair columns at $w \leq 0.82$ unconditionally and $w \leq 0.98$ modulo the crowding cap (constant interval-certified), equal-depth pair-only columns at every depth — for both benchmarks, via (MI). The remaining admissible regimes — multi-pair columns at the deep-probe depths $w \in \{1.5, 2\}$ (inside the gate class but beyond the theorems’ range), multi-pair columns with a depth in $w \in (0.98, 1]$ (ledger failure certified there; coverage is the re-run attacks plus the backstop), and mass-unbounded crowds (not LP-relevant, since columns have mass N and deeper objects are garnish priced by Section 7’s ladder) — are floored at $8/9$ of the corner unconditionally and covered by converged pricing (O2); the $(0.98, 1]$ sliver’s coverage is the re-run attacks under (O1) plus the backstop. Class-level exactness of $\delta_0 = 0$ is therefore proved on the closure’s regimes and heuristic (pricing-backed, backstopped at $20/27$) on that residue. The channel could only ever have produced a bite (a pair-assisted violation would be cut by pricing, not absorption), and it produces none; the audit’s minor finding 4 is discharged for the closure’s regimes, with the honest residue stated in Section 8.3.

4.4 The computational certification: 940 records, the ε law, and the witness

The gate LP (column generation over explicit configurations, analytic-gradient multi-start pricing, exact-LP equality on the accumulated dictionary, final solutions re-verified in **Fraction** arithmetic on 10^{-12} rationalizations) was run over the full pre-registered decision grid. Verdict, machine-readable: ABSORPTION.

The grid. $N = 64$: {matched, asymptotic} budgets $\times W \in \{2, 3, 4, 6, 8\} \times C_{\text{led}} \in \{10, 100, 1000\} \times \varepsilon \in \{0.02, 0.05, 0.10\} \times \text{fuzz} \{\text{coupled}, \text{none}, \Gamma \in \{0.05, 0.10, 0.25\}\} \times \text{error bars} \{\text{off}, \text{on}\} = 900$ records. $N = 128$: {matched, asymptotic} $\times W \times \text{fuzz} \{\text{coupled}, \text{none}\} \times \text{bars} = 40$ records. Total **940**; every record solved to status success; LP duality gap $\sim 10^{-9}$.

The result. $\delta_0 = P_{\text{full}} - P_{\text{base}} = 0$ at every record: $\max |\delta_0| = 6.39 \times 10^{-13}$ over the 900 $N = 64$ records and 8.94×10^{-13} over the 40 $N = 128$ records — $|\delta_0| < 10^{-12}$ **at all** 940 (the number to cite for the full grid is 8.94×10^{-13} , the $N = 128$ maximum). Per- W values at the primary point: -3.77×10^{-13} ($W = 2$), $+1.44 \times 10^{-13}$ ($W = 3$), 0.0 exactly ($W = 4, 6, 8$); the fit $\delta_0(W) = \delta_\infty + cW^{-\kappa}$ degenerates to $\delta_\infty = 0$ with zero residual, and $d\delta_0/d\Gamma = 0$. The optima are not merely equal but achieved by identical optimal laws: the base-optimal law itself satisfies the entire cubic block.

The exact ε law. At asymptotic budgets the joint optimum obeys $P(\varepsilon) = 5/6 - (2/3)\varepsilon$ — the grid-decoupled corner of Theorem 4.3 — at every tested ε :

ε	P (asymptotic, LP)	$5/6 - (2/3)\varepsilon$	$ \delta_0 $
0.02	0.8200000000002683	0.82	2.8×10^{-13}
0.01	0.82666666666669427	0.8266666...	2.4×10^{-13}
0.005	0.83000000000002841	0.83	2.2×10^{-13}
0.002	0.83200000000003910	0.832	1.0×10^{-13}

Matched-variant values differ only through the finite-size budget center (e.g. $P = 0.8368627361$ at $\varepsilon = 0.002$, from $B_1 = 84.712 < (4/3)64 = 85.33$). Moreover $P_{\text{base}} = P_{\text{cal}}$ at every record: even the λ' -Frobenius row alone adds nothing to the two-moment baseline.

Primary decision point (flat/flat, $\lambda' = 1/2$, $N = 64$, matched budgets, $\varepsilon = 0.05$, $C_{\text{led}} = 100$, coupled fuzz): $P_{\text{full}} = P_{\text{base}} = 0.8050956815843511$, $\delta_0 = -3.8 \times 10^{-13}$, identical optimal laws; the same P at fuzz = none (the fuzz row is not load-bearing for the absorption). $N = 128$ **confirmation**: $P_{\text{full}} = P_{\text{base}} = 0.8024488905983833$ (matched) and 0.8000000000004773 (asymptotic) at $\varepsilon = 0.05$; $\delta_0 = 0$ at all 40 records.

The witness (witness_N64.json; $N = 64$, matched budgets, $\varepsilon = 0.05$, fuzz = none). Three columns, all supported on the 65-site ψ_1 -zero grid (spacing $64/65 = 0.9846153846$), marks $\{1, 2\}$ only, no pairs, $n_- = 0$:

weight	column	N_d	$F_1 (= \sum m^2, \text{ exact})$	F'	C'
0.5299906673474316	vacancy lattice: 64 simples on 64 of the 65 sites	64	64	125.0909091 = 4128/33	245.4508724
0.16040139164388625	16 doubles + 32 simples on an explicit 48-site subset	48	96	153.7604346	411.0865130
0.30960794100868216	32 doubles on an explicit 32-site subset	32	128	135.1959163	301.3541285

Row values in this table are the clean-integer-grid values (clean_checks[].*clean — the canonical hand-checkable description); the optimizer's stored law rows (law[].F1/Fp/Cp) differ from them by at most 4.1×10^{-5} per row (per column $3.4 \times 10^{-6}/1.6 \times 10^{-5}/4.1 \times 10^{-5}$), and the aggregates below are the stored law's, which saturate the budget edges exactly. Aggregates: $E[F_1] = 88.94775275723633 =$ the upper Frobenius edge $B_1(1+\varepsilon)$ exactly; $E[F'] = 132.818$ (strictly inside its band $[128.34, 141.85]$); $E[C'] = 289.32716513260516 =$ the LOWER cubic edge $B_3(1-\varepsilon)$ exactly (the doubles corner is cubic-short, per $G(1/2) > 0$, and the arrangement supplies the budget from below); $E[N_d]/N = 51.52612362142/64 = 0.8050956815846874 \leq 5/6 + 0.01$ (the pre-registered absorption bar). Active rows at the optimum: only F1_hi and Cp_lo; every ladder row slack by orders of magnitude — $E[n(2)] = 3.50$ and $E[n(3)] = 0.94$ against caps $C_{\text{led}}NV^{-4} = 400$ and 79.0 , all higher rows zero, and the vacancy column's $n(V)$ vector identically zero. Every row was re-verified in exact Fraction arithmetic (all 10 checks pass, objective deviation 5×10^{-13}), and the audit independently recomputed all three columns through a position-space path (reproduction to 10^{-6} or better, feasibility re-confirmed at $C_{\text{led}} = 100$ and $C_{\text{led}} = 10$).

The mechanism is now fully visible: $F_1 = \sum m^2$ exactly on all three columns (Theorem 3.9 — the equality case of Lemma 4.1), so the bandwidth-one reading transfers the lemmaR.tight corner verbatim, while the sub-grid arrangement (which sites, which clusters) freely meets the λ' rows (Proposition 3.10): the two doubles-subsets' occupancy-2 clustering supplies the cubic cross-terms ($C' = 411.09$ and 301.35 against isolated-block values 160 and 256) at zero cost in N_d and zero cost at $\lambda = 1$. The witness uses no pairs, no marks ≥ 3 , no spectral escape — arrangement alone.

Pricing and its honest role. No improving column (reduced cost $< -10^{-6}$) was found in $S = 200$ pricing restarts at the primary and tightest-asymptotic points ($S = 100$ at $N = 128$ — RUN-REPORT-recorded, with no stored verify block; $S = 40$ at W -scan re-checks; $\min_{\text{rc}} = 0.0$, 0 improving columns at every stored verification). The ABSORPTION verdict does not rest on

pricing optimality — the witness is exhibited and verified, and $P_{\text{full}} = P_{\text{base}}$ is an exact LP equality on the accumulated dictionary regardless; pricing convergence supports only the claim that no undiscovered configuration class outside the dictionary would create a bite (and more columns can only lower both optima together — the anti-absorption direction). The audit’s eight break attempts (independent witness recomputation; a per-column ladder at $C_{\text{led}} = 4$, barely above the null floor 3.82; a missing-row hunt; near-CUE pinning; window-degeneracy probes; the pair-channel attack; budget-center stress across ε in 0.002–0.10 and Γ to 0.25; seed and scale stability) all failed to flip the verdict; the two that produced findings (near-CUE promotion; p_1 scoping) are executed in Sections 5 and 6.

5 The near-CUE class: absorption inside the unconditional data class

The Tier-1 adversary of Section 4 is free to hold anti-correlated (non-CUE) bandwidth-one data. The strongest form of the no-go pins the adversary inside the near-CUE class — the data class of the PairCeiling formalization — and asks whether the cubic block bites there. It does not; and the way it fails is the paper’s sharpest structural finding.

5.1 The license: [BGSTB24], with three riders

The pinning rows $|E_w|c_j|^2 - j| \leq \tau_2$ for $1 \leq j < N$ at $\lambda = 1$ (edge row free) are licensed by:

Baluyot–Goldston–Suriajaya–Turnage–Butterbaugh, *An unconditional Montgomery theorem for pair correlation of zeros of the Riemann zeta function*, arXiv:2306.04799 (v1, June 2023); published Acta Arithmetica, 2024. Theorem 1 there is unconditional and pointwise in α , uniform on the CLOSED band $0 \leq \alpha \leq 1$:

$$F(\alpha) = T^{-2\alpha}(\log T + O(1)) + \alpha + O(1/\sqrt{\log T}),$$

for the form factor defined with zeros counted with multiplicity and off-line zeros entering with the depth weight $x^{\delta+\delta'}$ — exactly the cosh depth-weighting the gate’s form factor c_j applies. With $\alpha = j/N$, each open-band row is licensed pointwise per row, unconditionally, in the depth-weighted multiplicity-counted form (directly, or through the paper’s Lemma 5 admissible kernel class: even, L^1 , supported in $[-1, 1]$, Lipschitz at 0 — the bandwidth-one class); the diagonal spike is confined to the edge rows the gate leaves free, and the Poisson factor acts below the model’s $1/N$ bin resolution whenever $N \ll \log T$. Two citation corrections for the program’s literature file, read off the fetched paper: the uniform range is the closed band $0 \leq \alpha \leq 1$, not $|\alpha| < 1$ (Theorem 1 includes the [GM87, Lemma 8] improvement “up to $\alpha = 1$ with explicit error terms”); and the published venue is Acta Arithmetica 2024.

The three licensing riders (binding on every use of this section’s results):

1. **Depth-aggregation only.** The licensed statistic never separates on-line from off-line mass: the Remark after their Theorem 2 states the method “neither requires nor provides any information” about whether the zeros are on the critical line. What is pinned is the cosh-weighted aggregate $|c_j|^2$ (pairs entering with depth weights), never an on-line-only statistic. Shedding the far-off-line coupling term costs an extra hypothesis — the thin-box hypothesis or the strong zero-density hypothesis, which their Theorems 2–3 (the 61.7% simple-zeros results) consume; the row class used here needs none of that.

2. **Finite- T slack** $O(N/\sqrt{\log T})$. The licensed per-row slack at finite T is $O(N/\sqrt{\log T})$ grid units (plus the edge spike), so $\tau_2 \in \{1, 4\}$ is the fixed- N , $T \rightarrow \infty$ asymptotic reading — legitimate for the gate’s asymptotic semantics, but at honest finite T the $O(1)$ -unit pinning overstates the theorem until $\log T \gg N^2$, and this no-go says so.
3. **Second-moment scope.** [BGSTB24] stops at the second moment — pair correlation only; nothing in it licenses the cubic budget column, whose $c_3(\lambda')$ identification stays on flag F2 (Section 8.3) exactly as before.

Net: the Tier-2 rows are an unconditional asymptotic data class, pointwise on the open band in the depth-weighted form; the Tier-2 verdict’s *strength* (“absorption inside the parent ceiling’s own data class”) inherits riders 1–3 verbatim.

5.2 The decision-grade result: $\delta'_0 = 0$ exactly, both N , both centerings

Row class: all Tier-1 rows (fuzz = none — the harsher variant for absorption) plus the pinning rows at τ_2 . Four cells at $N = 64$ (matched budgets, $C_{\text{led}} = 100$, $W \leq 8$):

τ_2	ε	$P_{\text{full}}^{\text{T2}}$	$P_{\text{base}}^{\text{T2}}$	δ'_0	reading
1	0.05	0.8332334059839699	0.8332334059839699	0.0 (exact)	absorb, decision grade
1	0.02	0.8359542395410772	0.8359374423865479	1.68×10^{-5}	absorb (0 within the residual band $[-3.1 \times 10^{-5}, +1.8 \times 10^{-5}]$)
4	0.05	0.8150864520465212	0.8137831067863290	1.30×10^{-3}	bounded residual, sub-decision scale
4	0.02	0.8316328692004420	0.8301979402692543	1.43×10^{-3}	bounded residual, sub-decision scale

Decision-grade discipline at the primary cell ($\tau_2 = 1$, $\varepsilon = 0.05$): $\delta'_0 = 0$ is an exact LP equality — identical optima — at every one of 9 cumulative $S = 200$ verification passes across a $\sim 1,100$ -column dictionary enrichment; the common level fell $0.8333622 \rightarrow 0.8332334$ while the equality never broke (residual improvements are common-mode), and the reduced-cost bound for the true marginal value is $|\delta'_0| \leq 1.2 \times 10^{-5}$. Stability under budget re-centering: asymptotic variant $P_{\text{full}} = P_{\text{base}} = 0.8332332347490976$, $\delta'_0 = 0.0$ exact.

$N = 128$ **confirmation.** The pinned LP was infeasible on the raw gate dictionary and was bootstrapped with 300 CUE and doubles-decorated-CUE columns (anti-absorption-safe), then built out over 10 cumulative $S = 200$ passes to 6,457 columns at the matched cell (6,739 after the asymptotic re-centering): $P_{\text{full}} = P_{\text{base}} = 0.8374218534574983$ with $\delta'_0 = 0.0$ exact at the matched cell, and 0.8373740026259704 with $\delta'_0 = 0.0$ exact at the asymptotic re-centering. Build-out invariance: $\delta'_0 \leq 8 \times 10^{-16}$ at every recorded dictionary re-solve (1,842/6,457/6,739 columns) — the verdict is independent of the convergence level. Level trend: $P^{\text{T2}} = 0.8374$ ($N = 128$) vs 0.8332 ($N = 64$), both $\sim 5/6 - O(10^{-3})$, the pinned-baseline finite-size trend.

5.3 Absorption by slack: the structural finding

The near-CUE witnesses make the verdict structural. At $N = 64$: a 64-column law, marks $\{1, 2\}$ only, no pairs, no marks ≥ 3 , with $E|c_j|^2 = j + 1$ for every $j = 1, \dots, 63$ to 2×10^{-12} (maximum

deviation 1.84×10^{-12} , at $j = 62$) — all 63 upper pinning rows active at exactly the $+\tau_2$ allowance — and **the only active rows at the pinned optimum are the 63 pinning rows (and mass): F_1 , F' , C' , and every ladder row are strictly interior. The cubic block does not even bind.** The $+\tau_2$ allowance pays for the doubles; the cubic budget is met with room to spare. **Fraction** re-verification passes on the full row class (worst pinning margin -1.0×10^{-9} , inside the documented 10^{-6} grace). At $N = 128$ the same structure: 128 columns, marks $\{1, 2\}$, no pairs, upper pinning edge ridden at $\tau_2 - 10^{-5}$ (an interior shrink absorbing a solver-tolerance overshoot; the **Fraction** re-verification against the TRUE $\tau_2 = 1$ box passes with margin $+1.0 \times 10^{-5}$ strictly interior), $E[n_{\text{simple}}]/N = 0.6748$.

This is a stronger statement than Tier-1 absorption: inside the near-CUE class the cubic row is not absorbed by an adversary straining against it — it is slack at the optimum. There is no ceiling-class escape and no bite even inside the parent ceiling’s own data class, at both lattice sizes, at both budget centerings, subject to riders 1–3 of Section 5.1.

5.4 The $\tau_2 = 4$ annotation

At loose pinning ($\tau_2 = 4$) the cubic row is active and the residuals $\delta'_0 = 1.30 \times 10^{-3}/1.43 \times 10^{-3}$ ($\varepsilon = 0.05/0.02$) exceed their pricing bands ($[0.9 \times 10^{-3}, 1.5 \times 10^{-3}]$) and were still declining under enrichment ($1.28 \times 10^{-3} \rightarrow 1.17 \times 10^{-3}$ over five passes): they are **unconverged upper bounds at loose pinning, an order of magnitude below the M4 decision scale (the 10^{-2} – 3×10^{-2} δ_0 the proposal projected)**, obtained from deliberately shortened runs and recorded as such. They are diagnostic annotations, not certificate claims — and they sharpen the scoping: whatever residual cubic signal exists at this scale lives strictly between the near-CUE class ($\tau_2 = 1$: exact zero by slack) and the free Tier-1 class (exact zero by witness), i.e. in loosely pinned intermediate worlds, at sub-decision magnitude.

6 The simple-fraction benchmark

The parent certificate’s headline is the simple-zero fraction ($2/3$ at flat window; 0.6725 window-optimized; ceiling 0.6818287), so a no-go scoped only to N_d would be incomplete — this was the audit’s third major finding. The gate was therefore re-run with the objective $\min E_w[p_1]$, $p_1(c) = (\# \text{mark-1 atoms} + 2 \times \# \text{mult-1 pairs})/N$, on the follow-up-enriched dictionary (5,134 columns), with the literal pre-registered stop rule applied to both systems:

cell	$p_{1,\text{full}}$	$p_{1,\text{base}}$	δ_{p_1}	analytic corner $2 - (B_1/N)(1 + \varepsilon)$	converged
matched, $\varepsilon = 0.05$	0.6101913631697143	0.6101913631684138	1.3×10^{-12}	0.6101913631681821	yes: $S = 200$ clean, $\min_{\text{rc}} = 0.0$, both systems
asymptotic, $\varepsilon = 0.002$	0.6640000000012899	0.6640000000004871	8.0×10^{-13}	$0.6640000000000001 = 2 - (4/3)(1.002)$	yes: $S = 200$ clean, both systems

These were the campaign’s only fully clean-stop runs: zero improving columns at $S = 200$ for full AND base systems. Cross-checks on the final dictionary: coupled fuzz at the primary cell $\delta_{p_1} = 2.9 \times 10^{-13}$; W -scan over $\{2, 3, 4, 6, 8\}$ $\max |\delta_{p_1}| = 1.3 \times 10^{-12}$ (at $W = 8$; all $< 1.4 \times 10^{-12}$); the labeled on-line-only diagnostic variant 3.3×10^{-13} . The optimum is the exact marks- $\{1, 2\}$

doubles corner of Theorem 4.3(b) — pairs cannot help (a mult-1 pair buys 2 simple points at $\geq$ twice the two-simples Frobenius cost; a mult-2 pair is dominated by two doubles) — and the $\varepsilon \rightarrow 0$ value is $2 - 4/3 = \mathbf{2/3}$, the model's bandwidth-one analog of Montgomery's simple-zeros corner. The witness laws are the same ψ_1 -grid family as the Tier-1 witness, reweighted per cell, rational verification passing.

Two scoping notes. (i) The cited 0.6818287 ceiling belongs to the *window-optimized* certificate class (the Theorem-D effect), which the flat/flat gate deliberately does not consume; the flat-window model corner is $2/3$, and no statement in this paper re-decides the 0.6818287 ceiling — what is proved is that the cubic block does not move the flat/flat optimum for either objective. (ii) With this section, the no-go headline covers BOTH benchmarks: the distinct-zeros corner $5/6$ and the simple-fraction corner $2/3$, each with $\delta = 0$ at convergence.

7 Capacity and vacuity: no spectral escape

Two theorems close the spectral-escape routes from opposite ends, and together with the gate they complete the pricing: a scalar cubic tail row with any divergent cutoff is unconditionally vacuous (Theorem 7.4 — the “provable today, no LP needed” half of the no-go, mandated by the merge guidance); and under the repaired all- V ladder the same channel is capped at $o(N)$ with a sharp constant (Theorem 7.1). What remains is position/interference freedom — which is exactly the channel the witness of Section 4.4 uses.

7.1 The sharp capacity theorem

Setting (the continuum garnish LP). Heights normalized so the ladder's absolute floor is 1. Spectral escape is a positive measure μ on $[1, \infty)$ — $\mu([h, \infty))$ is the count per N of escaped eigenvalue mass at height $\geq h$ — subject to

$$\begin{aligned} \text{(Frobenius slack)} \quad & \int h^2 d\mu(h) \leq \varepsilon_g, \\ \text{(all-}V \text{ ladder)} \quad & \mu([h, \infty)) \leq Ch^{-4} \quad \text{for every } h \geq 1 \quad (C = C_{\text{led}}), \end{aligned}$$

and the objective is the absorbable cubic charge $\int h^3 d\mu(h)$. (Equivalently, in tail-function form $N(h) = \mu([h, \infty))$: maximize $N(1) + 3 \int_1^\infty h^2 N dh$ subject to $N(1) + 2 \int_1^\infty h N dh \leq \varepsilon_g$ and $N \leq Ch^{-4}$; the forms are identical by integration by parts.)

Theorem 7.1 (sharp capacity). *For every $C > 0$ and $0 < \varepsilon_g \leq 2C$:*

$$\max_{\mu} \int h^3 d\mu = 2\sqrt{2} \sqrt{C\varepsilon_g},$$

attained by the min-profile $N_(h) = C \min(a^{-4}, h^{-4})$, $a = \sqrt{2C/\varepsilon_g}$ (as a measure: no mass on $[1, a)$; density $4Ch^{-5} dh$ on (a, ∞)).*

Proof. Upper bound: for any feasible μ and ANY $a > 0$, the pointwise inequality $h^3 \leq ah^2 + (h^3 - ah^2)_+$ on $[1, \infty)$ gives

$$\int h^3 d\mu \leq a\varepsilon_g + \int (h^3 - ah^2)_+ d\mu.$$

Writing $g(h) = (h^3 - ah^2)_+ = \int_a^h g'(s) ds$ for $h \geq a$, with $g'(s) = 3s^2 - 2as > 0$ on (a, ∞) , Tonelli and the ladder give

$$\int g d\mu = \int_a^\infty g'(s) \mu([s, \infty)) ds \leq C \int_a^\infty (3s^2 - 2as) s^{-4} ds = C \left(\frac{3}{a} - \frac{a}{a^2} \right) = \frac{2C}{a}.$$

So $\int h^3 d\mu \leq a\varepsilon_g + 2C/a$ for every $a > 0$; minimizing over a (minimum at $a = \sqrt{2C/\varepsilon_g}$) gives $2\sqrt{2C\varepsilon_g}$. Attainment: $\varepsilon_g \leq 2C$ makes $a \geq 1$, so N_* is an admissible tail function; its Frobenius charge is $Ca^{-4} + Ca^{-4}(a^2 - 1) + Ca^{-2} = 2C/a^2 = \varepsilon_g$ exactly, and its cubic charge is $C/a + 3C/a = 4C/a = 2\sqrt{2}\sqrt{C\varepsilon_g}$, meeting the bound. $\square$

Remark (the squeeze, made exact). The optimal dual split is

$$[a \times \text{Frobenius}] + [\text{ladder integral above } a] :$$

bounded heights are blocked by the Frobenius cost (mass at height h buys h^3 of cubic but pays h^2 of Frobenius, ratio $h \leq a$), divergent heights by the ladder cap. The absolute floor C_{abs} (normalized to 1) appears NOWHERE in the upper bound — only in the attainability condition $a \geq 1$ — which is the precise form of the adjudication’s overruling of the “bite iff $2C_{\text{led}}/C_{\text{abs}} < 4/3$ ” claim: sharp constants at the floor are irrelevant against this adversary class.

Proposition 7.2 (the naive profile is infeasible; the corrected constant of record). *The point-mass-plus-tail profile (a ladder-saturating atom of size $n_0 = Cb^{-4}$ at height $b = \sqrt{3C/\varepsilon_g}$, plus the ladder-saturating tail above b) has Frobenius charge exactly ε_g and cubic charge $5C/b = (5/\sqrt{3})\sqrt{C\varepsilon_g} = 2.8868\sqrt{C\varepsilon_g}$ — but it violates the CUMULATIVE ladder on the open interval $(2^{-1/4}b, b)$, where its count $2Cb^{-4}$ exceeds the cap Ch^{-4} . The true optimum is Theorem 7.1’s $2\sqrt{2} = 2.8284271247\dots < 5/\sqrt{3}$; both earlier bookkeepings — the first-pass $5/\sqrt{3}$ and the adjudicator’s two-term dyadic $2\sqrt{C\varepsilon}$ — are superseded by the min-profile constant (same order; conclusions unchanged). The constant has three independent confirmations on record: the specification’s min-profile derivation, the implementer’s analytic re-derivation plus LP (2.8244 at $h_{\text{max}} = 400$, gap = the predicted truncation tail $3/h_{\text{max}}$), and the audit’s own optimum.*

Proof. The tail part has density $4Ch^{-5}$ on (b, ∞) . Frobenius: atom $Cb^{-2} + \text{tail } 2Cb^{-2} = 3Cb^{-2} = \varepsilon_g$ at the stated b . Cubic: atom $C/b + \text{tail } 4C/b$. Ladder violation: for $2^{-1/4}b < h < b$ the cumulative count is $n_0 + Cb^{-4} = 2Cb^{-4}$ while the cap is $Ch^{-4} < 2Cb^{-4}$ throughout the open interval. $\square$

Corollary 7.3 (capacity is $o(N)$: the positive complement of the no-go). *In the gate’s units, the maximal cubic shift purchasable by spectral escape under the repaired row system is*

$$|\Delta_{\text{cubic}}| \leq 2\sqrt{2}\sqrt{C_{\text{led}}\varepsilon_g}N = o(N) \quad \text{as } \varepsilon_g \rightarrow 0 \text{ at fixed } C_{\text{led}}$$

(e.g. $C_{\text{led}} = 1000$, $\varepsilon_g = 10^{-8}$: capacity $0.0089N$). A doubles-vs-pairs-scale swing ($\Theta(N)$, e.g. $(4/3)N$) can NEVER be produced by spectral escape once the all-V ladder is consumed: any absorption of the cubic block must run through position/interference freedom — the channel Theorem 3.9 exhibits and the witness uses. (In the gate’s LP this theorem enters as the coupled fuzz rows, tangent-cut on the concave $\sqrt{\cdot}$; the run found the fuzz row not even load-bearing for the absorption.)

Numerical confirmation (`verify_t5.py`, refinement runs re-executed and persisted to `verify_t5_out.json` this session): attainment profile exact to $4 \times 10^{-11}/9 \times 10^{-13}$; an independent discretized LP over point masses converges to 0.4000 from below under joint grid-and-truncation refinement — tail-corrected values 0.3950/0.3985/0.3991 at (600 points, $h_{\text{max}} = 2000$), (2000 points, $h_{\text{max}} = 2000$), (4000 points, $h_{\text{max}} = 8000$), $C = 1$, $\varepsilon_g = 0.02$, each the raw LP value plus the predicted truncation tail $3C/h_{\text{max}}$ — never exceeding the bound; the naive profile’s value 0.408248 and its ladder violation are reproduced; duality and scaling verified across (C, ε) over five orders of magnitude.

7.2 Divergent-cutoff vacuity: the garnish theorem

Setting (the as-written row system, family semantics). $N = N(T) \rightarrow \infty$; $V_0 = V_0(T) \rightarrow \infty$ any divergent cutoff with $V_0 = o(N^{1/3})$ (any polylog cutoff qualifies — in particular the proposal's $(\log \log T)^3$ and every $(\log T)^4$). The as-written system consumed its equalities “with $o(N)$ precision” — no fixed tolerance rate — so feasibility is a property of families (c_T) of model configurations (no mark cap was imposed as written), with spectral data read from isolated-block spectra, admissible iff

- (W1) mass: $\sum m = N$ (exact);
- (W2) Frobenius: $F = \kappa N + o(N)$, $\kappa = \kappa(1) = 4/3$ (the as-written system's bandwidth-one Frobenius row);
- (W3) signed cubic: $C = c_3 N + o(N)$;
- (W4) scalar tail row: $\sum_{|\lambda_i| \geq V_0} |\lambda_i|^3 = o(N)$;
- (W5) count/inertia rows imposed only at thresholds $V \geq V_0$, plus $n_- \leq p$;

objective $\liminf N_d/N$ (or $\liminf p_1$). (A fixed tolerance rate shrinking faster than $1/V_0$ would be a STRONGER system than the one written — and dishonest bookkeeping besides, since the consumed prime-side equalities carry no such rate.)

Theorem 7.4 (garnish absorption). *Let (c_T) satisfy (W1), (W2), (W4), (W5), contain at least $s_0 N$ simple on-line atoms for a fixed $s_0 > 0$ (a simple-atom fraction, unrelated to the marginal value δ_0), and fall short of the (W3) cubic equality by $\Delta_T N$ with $0 \leq \Delta_T \leq \Delta_{\max}$ fixed. Put $h_0 = \lfloor V_0/2 \rfloor$, $n_0 = \lceil \Delta_T N / h_0^3 \rceil$, and let c'_T be c_T with n_0 additional isolated on-line atoms of mark h_0 (at unoccupied, well-separated positions) and $n_0 h_0$ simple atoms deleted. Then for all large T , c'_T satisfies EVERY row of the as-written system, and*

$$N_d(c'_T) = N_d(c_T) - n_0(h_0 - 1) \leq N_d(c_T), \quad p_1(c'_T) \leq p_1(c_T).$$

Row-by-row (per N ; every shift $o(1)$): mass exact 0; Frobenius $+ \Delta_T/h_0 + O(1/h_0^2 + h_0^2/N) = o(1)$; cubic $+ \Delta_T + o(1)$ (landing the (W3) equality); tail row $+ 0$ (the garnish sits at $h_0 < V_0$); count rows at $V \geq V_0$: $+ 0$; inertia unchanged (only positive on-line eigenvalues added).

Proof. Feasibility: $n_0 h_0 \leq \Delta_{\max} N / h_0^2 + h_0 = o(N) < s_0 N$ for large T — enough simples exist to delete. Each garnish atom is an isolated on-line atom of mark h_0 : eigenvalue h_0 , Frobenius h_0^2 , cubic h_0^3 , mass h_0 . Mass: adds and deletes $n_0 h_0$ exactly. Cubic: adds $n_0 h_0^3$ in $[\Delta_T N, \Delta_T N + h_0^3]$, subtracts $n_0 h_0 \leq \Delta_{\max} N / h_0^2 + h_0$; net $\Delta_T N + O(h_0^3) + O(N/h_0^2) = \Delta_T N + o(N)$ since $h_0 \rightarrow \infty$ and $h_0^3 = o(N)$ — (W3) now holds. Frobenius: adds $n_0 h_0^2 \leq \Delta_{\max} N / h_0 + h_0^2 = o(N)$, inside (W2)'s window. Tail and count rows: every garnish eigenvalue equals $h_0 = \lfloor V_0/2 \rfloor < V_0$, and all thresholds sit at $V \geq V_0 > h_0$, so the added spectrum is invisible; deletions only decrease those rows. Objectives: N_d changes by $+n_0 - n_0 h_0 = -n_0(h_0 - 1) \leq 0$; the deleted atoms were simple and the added ones have mark $h_0 \geq 2$, so p_1 decreases. $\square$

Remark (the growth cap is removable). The hypothesis $V_0 = o(N^{1/3})$ is a convenience, not a constraint: for an arbitrary divergent cutoff, cap the garnish height at $h_0 = \min(\lfloor V_0/2 \rfloor, \lfloor N^{1/4} \rfloor)$. Then $h_0 \rightarrow \infty$, $h_0^3 \leq N^{3/4} = o(N)$, and $h_0 < V_0$ still holds, so every estimate in the proof goes through verbatim — the vacuity covers ANY divergent cutoff, as headlined.

Corollary 7.5 (vacuity of the scalar divergent-cutoff cubic row). *Suppose the two-moment system $\{(W1), (W2)\}$ admits optimizer families with a fixed positive simple fraction, spectra bounded by a fixed constant, and bounded cubic deficit ($C(c_T) \leq c_3N + o(N)$, deficit $\leq \Delta_{\max}N$). Then, for both benchmarks, the infimum over the as-written system equals the infimum over the two-moment system: the marginal value of the entire block $\{(W3) + (W4) + (W5)\}$ is ZERO — $\delta_0 = 0$ as written. The deficit hypothesis is the actual situation: the corner families of Theorem 4.3 have marks in $\{1, 2\}$ (spectra bounded by 2, simple fraction $2/3$ in the limit) and undershoot the cubic budget (Corollary 3.6: $G(1/2) = +1/2 > 0$ — the null budget sits strictly above the doubles plane, so doubles-saturated configurations are cubic-short; the shipped witness accordingly sits at the LOWER cubic edge). Whatever cubic demand the row was hoped to impose on the two-moment-degenerate corner — up to any fixed $\Delta_{\max}$, including the full “ $(4/3)N$ doubles-vs-pairs swing” of the proposal’s narrative — the garnish supplies it invisibly, at an objective DECREASE.*

Proof. One direction is trivial (more rows can only raise the infimum). Conversely, take the corner families: bounded spectra plus $V_0 \rightarrow \infty$ make (W4) and the $V \geq V_0$ count rows identically zero for large T , so they satisfy (W1), (W2), (W4), (W5); apply Theorem 7.4 to land (W3); the garnished family is admissible for the whole system at a weakly smaller objective. $\square$

Remark 7.6 (what this does and does not say — the repair structure). **1. It kills the scalar row, not the direction.** The vacuity is about the AS-WRITTEN system, whose count/tail information starts only at V_0 . The proposal’s own Chebyshev proof in fact yields the all- V ladder $n(V) \leq C_{\text{led}}NV^{-4}$ for every $V \geq C$ at $\theta < 1$ (the theorem underclaimed its proof — the adjudication’s central overruling); under THAT system the garnish channel is not free but capped, and the cap is exactly Theorem 7.1’s $2\sqrt{2}\sqrt{C_{\text{led}}\varepsilon_g}N = o(N)$. Repair R1 (“claim what the proof proves”) is precisely the difference between Corollary 7.5 and Theorem 7.1.

- 2. The garnish needs unbounded marks** ($h_0 = \lfloor V_0/2 \rfloor \rightarrow \infty$): it lives outside every fixed- W alphabet — which is why the repairs mandated the divergent- W adversary class and why “a break at marks $\{1, 2\}$ is not evidence.” (A pair-based garnish with slowly divergent pair multiplicity reaches the same conclusion through Proposition 3.2’s deep-pair blocks.)
- 3. The three-part structure of the no-go.** Theorem 7.4/Corollary 7.5 dispose of spectral escape as written; Theorem 7.1 caps it as repaired; and the gate’s decided content (Sections 4–6) is that the remaining channel — position/interference freedom — absorbs the cubic block too, at $\delta_0 = 0$ exactly. Together these ARE the sharpened no-go.

Numerical confirmation (`verify_t6.py`): the shift table at $\Delta = 4/3$, $V_0 = (\log \log T)^3$ over the adjudication’s $\log T$ grid extended to 10^6 — Frobenius shift $4/(3h_0) = 0.0992$ down to $0.0010 \rightarrow 0$, N_d shift negative throughout, cubic shift $+4/3$ exactly, tail and count rows 0 — reproducing the adjudication’s computation-(b) audit and the gate’s Tier-A regression; an abstract LP demonstration in which the cubic equality is INFEASIBLE without the garnish variable (the fake bite) and returns to the two-moment corner $5/6 - O(\text{tol})$ with it; and an exact-rational instance with all shifts exact.

8 Scope, residuals, and what survives

The credibility of an exact zero is its scoping. This section states precisely where $\delta_0 = 0$ is exact, where it is a bounded residual, what is finite-scale, and what is flagged — and closes with the one place a positive signal genuinely lives, labeled as outlook.

8.1 The residual table

The exact zero is **flat/flat-specific**. Theorem 3.9’s mechanism says exactly why: the flat bandwidth-one kernel’s zeros are equally spaced, so one site grid kills all $\lambda = 1$ cross-terms simultaneously; a cosine window’s kernel zeros are not equally spaced, so no grid does, and the degeneracy breaks by a microscopic but nonzero amount. The complete residual map:

configuration	residual δ_0	status
flat/flat, entire 940-record Tier-1 grid	0 ($< 10^{-12}$)	exact; witness-backed
flat at $\lambda = 1$ / $\cos(1.6s)$ at λ'	5.95×10^{-14} ($\varepsilon .05$), -1.14×10^{-13} ($\varepsilon .02$)	exact-zero class: changing the λ' window moves nothing — the absorber’s freedom is positional, not spectral
$\cos(\sqrt{2}s)$ [Montgomery–Taylor] at $\lambda = 1$ / flat	$+6.41 \times 10^{-5}$ ($\varepsilon .05$), $+2.28 \times 10^{-5}$ ($\varepsilon .02$)	bounded residual: the MT cosine breaks the exact grid degeneracy; unconverged upper bounds, three orders below the ε -slack scale
$\cos(\sqrt{2}s)$ [MT] / $\cos(1.6s)$	$+6.27 \times 10^{-5}$ ($\varepsilon .05$), $+1.91 \times 10^{-5}$ ($\varepsilon .02$)	same class
near-CUE $\tau_2 = 1$ (decision grade, $N = 64$ and 128, both centerings)	0 exact (reduced-cost bound $\leq 1.2 \times 10^{-5}$ at $N = 64$)	absorption by slack (Section 5.3)
near-CUE $\tau_2 = 4$ (loose box, diagnostic)	$1.30 \times 10^{-3}/1.43 \times 10^{-3}$, pricing bands $[0.9 \times 10^{-3}, 1.5 \times 10^{-3}]$	unconverged upper bounds, an order below the M4 decision scale (10^{-2} – 3×10^{-2}); the cubic row is active here
audit’s tight- ε Tier-2 probes (dictionary-limited, no column generation)	$\leq 5 \times 10^{-4}$, non-monotone in ε	upper bounds, same order as the documented $\sim 3 \times 10^{-4}$ pricing uncertainty — “where any residual signal concentrates”

Every nonzero entry is an unconverged UPPER bound on the converged marginal value, and every one sits at least an order of magnitude below the M4 decision scale. The audit’s binding wording bar is respected verbatim: $\delta_0 = 0$ exactly at the primary flat/flat configuration; $\leq O(10^{-4})$ across the window family and $\leq O(5 \times 10^{-4})$ under near-CUE pinning at that scale, within pricing-convergence uncertainty — now strengthened by the decision-grade $\tau_2 = 1$ exact zeros of Section 5, which post-date that wording.

8.2 Finite scale

The computational certification is at $N = 64$ and $N = 128$, verdict-stable across both (and across two budget centerings whose cubic centers differ by 5% — the F_1 centers by 0.7%, F' by 2.6%). This is not extrapolated silently: the continuum analog of the decoupling mechanism is exact (Proposition 3.11 — the integer lattice under the sinc kernel, with the finite- N grid its exact periodization), so the absorption has a structural reason to persist at every scale, and the model-level theorems of Sections 3, 4.1–4.3, and 7 are N -generic (the pair-channel capacity constants were certified at $N = 64$, the decision geometry; the mechanical $N = 128$ re-run, executed 2026-08-27, reproduces every identity and inequality — see O4 below). No claim about N beyond 128 rests on

computation.

8.3 Flags and honest residue (carried, not discharged)

- **F2 — the cubic budget center.** The identification $c_3(\lambda') = m_3(\lambda')$ ($= 5$ at $\lambda' = 1/2$) rests on Rudnick–Sarnak-range GUE 3-level correlations and the parent’s Section-5 diagonal method’s polylog-window uniformity; Theorem 3.5 proves the SINE value, not the identification. The absorption verdict is insensitive to this flag: $\delta_0 = 0$ across ε in 0.002–0.10 and Γ to 0.25, and across budget centers differing by 5% — an $O(1)$ -per-zero error in c_3 cannot flip it (the audit’s break attempt 7). [BGSTB24] licenses nothing at third order (rider 3).
- **F4 — the ladder’s provenance.** The all- V ladder consumed by Theorem 7.1’s system rests on Theorem 1(ii)-repaired, whose V -dependent bridge (repair R2) is unproven. Per the gate’s semantics the absorption branch is STRENGTHENED by assuming the ladder (absorption even with the ladder in force), and Theorem 7.4 needs no ladder at all; nothing in this paper depends on R2.
- **Heuristic pricing.** For classes without an exhibited witness (the nonzero rows of the residual table; the claim that no undiscovered configuration class bites), the evidence is converged column-generation pricing ($S = 200$ restart protocol), which is heuristic adversary-optimality — stated as such wherever used, and never load-bearing for the exact-zero claims, which are witness-backed LP equalities.
- **Pair-channel residue** (from the companion note): (O1) [updated 2026-08-27] The per-cell interval hardening was executed: the crowding cap’s ledger constant is interval-certified to $w \leq 0.98$ ($0.98465 < 1$); the earlier full-family ratio 0.9775 was a grid artifact — on $w \in (0.98, 1]$ the ledger fails (certified > 1) and coverage is the attacks (+17.46, re-run at the sliver) plus the 8/9 backstop; the cap itself on $w \in (0.82, 0.98]$ remains granted-not-proved (its $w \leq 0.82$ self-consistency is now interval-certified on both sides). (O2) Single-pair depths in $(0.45, 1.0)$ at unrestricted mass, and multi-pair beyond the family, are covered by numerics plus the 8/9 backstop only — not needed for any gate/LP statement. (O3) The sharp conjecture (Section 4.3) is open; near-tight configurations (+0.03) mean the constant-chasing is genuinely hard. (O4) The certified capacity constants are at $N = 64$; the mechanical $N = 128$ re-run (2026-08-27) reproduces every identity and inequality with $\leq \sim 2.5\%$ drift in the headline certified constants (deep-tail table values, $y \geq 1.5$, drift up to $\sim 8\%$), moves Theorem A(ii)’s deep threshold to $d \geq 1.1$, and shows the near-endpoint ledger failure at $N = 128$ as well (smaller sliver, better capped constant 0.9633). (O5) All pair-channel constants are for the flat $\lambda = 1$ window; none transfer to the MT-cosine window without recomputation — consistent with that window’s residual row above.
- **The formalized ceilings are not re-decided.** The 0.6818287 bandwidth-one ceiling belongs to the window-optimized certificate class; the 0.8453 trace/Frobenius cap stands on $\kappa \geq 2/\sqrt{3}$. The flat/flat gate consumes neither; this paper’s claim is that cubic augmentation moves nothing at the flat/flat operating point and produces only the bounded residuals above elsewhere.

8.4 Outlook: the exploratory λ' payoff curve (outside the proven ladder)

Labeled per repair R4 and merge-guidance item (4) (the MD(λ, δ) community-target quarantine): $\lambda' > 1/2 + o(1)$ is OUTSIDE the proven $\theta < 1$ ladder regime; these are M5 payoff-curve data for the

moderate-deviation community target $\text{MD}(\lambda, \delta)$, NOT certificate claims; column generation ran at reduced budget, so positive values are heuristic and upper-anchored.

λ'	ε	P_{base}	P_{full}	δ_0
0.55	0.05	0.8053310433508489	0.8053310433508484	0
0.55	0.02	0.8251787278264935	0.8251787278264899	0
0.60	0.05	0.8058514020011891	0.8062187975519491	$+3.67 \times 10^{-4}$
0.60	0.02	0.8256842190868753	0.8259625547887860	$+2.78 \times 10^{-4}$
0.65	0.05	0.8059503048717801	0.8086104956189072	$+2.66 \times 10^{-3}$
0.65	0.02	0.8263688470852664	0.8278795159840451	$+1.51 \times 10^{-3}$

The cubic row’s marginal value turns positive between $\lambda' = 0.55$ and 0.60 — precisely where the flat-window moment margin $\text{Marg}(\lambda) = 2m_2 - m_3$ changes sign (Corollary 3.6: $\lambda_0 = 0.610511$) — and grows toward the Rudnick–Sarnak endpoint $\lambda' = 2/3$. The first possible bite therefore sits at $\lambda' \gtrsim 0.6$; consuming it would require extending the proven large-value ladder into exactly the moderate-deviation regime $\text{MD}(\lambda, \delta)$ that the program has quarantined as a community target. At the proven operating point $\lambda' = 1/2 + o(1)$, the value is exactly zero — that is this paper’s theorem-grade content; the payoff curve is its honestly labeled boundary.

9 Consequences

1. **lemmaR_tight stands strengthened.** The two-moment degeneracy is not broken; it now carries an audited exact-zero certificate: two-bandwidth flat-window augmentation — λ' -Frobenius, signed cubic, all- V ladder, capacity fuzz, at $\lambda' = 1/2$ — adds exactly zero to the two-moment optimum at $N = 64/128$ scale, for both benchmarks, including inside the near-CUE class (by slack). The program’s map entry is updated from “exhausted” to “exhausted, and robust under RS-range cubic augmentation with capacity control.”
2. **The R2 bridge is not to be funded for the cubic payoff.** The gate was built to price the bridge investment; the payoff is $\delta_0 = 0$. Theorem 1(ii)-repaired retains its standalone value (the first spectral-tail statement for the Weil–Gabor matrix) and its other consumers.
3. **What remains live in A4:** (i) Theorem 1(ii)-repaired as the standalone analytic deliverable of record (all- V count ladder with explicit C_{led} , $\theta < 1$ strict, C^3 taper); separately, per repair R3, Theorem 1(i)’s fixed- λ divergent cutoff $V_0 = T^{2\lambda-1+\varepsilon}$ extends to all $\lambda \in (1/2, 2/3)$ via the MVT-4 branch alone — a divergent-cutoff statement, subject to Theorem 7.4’s vacuity, not an all- V ladder; (ii) partial orthonormalization off the $o(N)$ -dimensional exceptional subspace, where a divergent cutoff SUFFICES (unlike the cubic LP) — rank-perturbation bookkeeping already in the formalized linear-algebra library; (iii) the near-critical observation $X^2/T = (\log T)^{2\theta} = \text{polylog}$, shrinking the shift range needed by $\text{HL}^*(4)$ -type quartic input to logarithmic — a route into the existing 13/18 machinery via averaged Goldston–Montgomery/Montgomery–Soundararajan results; (iv) $\text{MD}(\lambda, \delta)$ as a community target, now carrying the quantified payoff curve of Section 8.4.
4. **Methodological consequence.** The gate architecture itself — explicit-configuration LP with realizability by construction, clustered null budgets, pre-registered decision semantics, witness-or-nothing absorption claims, adversarial audit with break attempts — is the reusable

instrument this paper certifies. In particular the correction of record (Theorem 3.4 superseding the “+8 vs 0” premise and the phase model) was produced by the gate’s own verification discipline before any LP ran.

10 Formalization: what is machine-checked, and what remains

The core of Sections 3 and 4 is **not a plan** — it is **formalized, compiled, and machine-checked** in Lean 4 against Mathlib, with no `sorry`, no `native_decide`, and no numeric certificates. Three files under `Zeta23/PairCeiling/` carry it (Lean 4.33.0-rc2; Mathlib rev 51e6992; last rebuilt 2026-08-27, “Build completed successfully”):

- `GridParseval.lean`. `flat_band_trace_sq` — **Lemma 3.8**, proved over any integral domain carrying a primitive M -th root of unity, for any band of M consecutive harmonics; and `trace_sq_grid` — **Theorem 3.9**, the literal $\text{tr } \tilde{G}^2 = \sum_k m_k^2$ collapse on the grid. Supporting: character orthogonality on $\mathbf{Z}/M$, DFT Parseval in the integer-mark pairing, the band reindexing, and the triangular-weight telescoping.
- `GridWitness.lean`. `vacancy_F1` — the $N = 64$ vacancy lattice has $F_1 = \sum m^2 = 64$ for *every* vacancy position; `vacancy_half_band_value` — its $\lambda' = 1/2$ half-band row is exactly $F' = 4128/33$, again for every vacancy position, the hand-checkable worked value of Section 3.4 and the shipped witness column, its integer core 136224 checked by `decide +kernel` (kernel-checked, not `native_decide`); and `half_band_alive` — $F' = 4128/33 \neq 64 = \sum m^2$, which is the “alive” half of **Proposition 3.10**, i.e. the two-bandwidth decoupling itself, in witness form. The finite/algebraic half of Proposition 3.10’s display is formalized as `sum_band_pair_tri / half_band_fold`, together with `sum_window_residues` — that the s -window is a complete residue system mod 65, so no residue-class telescoping can occur, unlike the $\lambda = 1$ band.
- `GridCorner.lean`. `mark_one_count_ge` — **Lemma 4.2** at the simple-fraction level; `grid_corner_pointwise` — **Theorem 4.3** pointwise on the grid class; `grid_corner_law` — the same in law form, i.e. the LP lower bound with no discretization; and `grid_corner_attained` — exact attainment at $N = 64$, $\varepsilon = 1/32$ by a marks- $\{1, 2\}$ column (12 doubles, 40 simples), meeting **both** corners with equality simultaneously.

Axiom footprint. `#print axioms` on each of the twelve theorems named above returns `[propext, Classical.choice, Quot.sound]` and nothing else — the three axioms of Lean’s classical foundation. In particular there is no `sorryAx`, which any admitted step would introduce, and no `Lean.ofReduceBool`, which `native_decide` would introduce and which would put the compiler’s evaluator into the trusted base. Full record, with the environment and the reproduction recipe: *supplementary material*, `formalization-status.md`.

The reason this much closes is structural: on the ψ_1 -zero grid every datum is algebraic, and Theorem 3.9 turns the first step of the corner chain from an inequality into an **equality** ($F_1 = \sum m^2$ exactly), after which the two integrality levels finish in finite arithmetic.

What is not formalized, and what each would take.

- **Proposition 3.10’s general position-sensitivity statement** — two grid configurations with the same mark multiset but different F' . Finite algebra at fixed N , but it needs a cyclotomic non-vanishing certificate; the precise gap is documented in `GridWitness.lean`’s header. The witness form is proved; the general form is not.

- **Theorem 4.3 off the grid** (atom-only, arbitrary positions) — needs the position-space form of Lemma 4.1, i.e. $F_1 \geq \sum m^2$ with the kernel $K = D^2$ psd. This is the one step the grid case gets for free.
- **Proposition 3.3 and Theorem 3.4** — polynomial identities; ring-adjacent, and the cheapest remaining items.
- **Theorem 3.5** — piecewise-polynomial integration against Mathlib’s integral.
- **Theorem 4.9** — a finite Hermitian matrix, inertia subadditivity, one scalar inequality; formalization-friendly, and the natural next unit from the pair channel.
- **Theorem 7.1 and Theorem 7.4** — one page of real analysis (Tonelli plus a one-variable optimization) and a bookkeeping argument respectively.
- **The pair-channel capacity results of Section 4.3** — Python-certified, with interval arithmetic. Formalizing the certified curves is a genuinely larger undertaking than the rest combined, and is not claimed here.
- **The full LP witness with its dual certificate** — the exact-rational route is available: the witness law is three columns with rational weights, integer site subsets and marks in $\{1, 2\}$, every row value rational ($F_1 = 64, 96, 128$; F' and C' rational combinations of the triangular weights, e.g. $4128/33$), so feasibility-plus-optimality is an exact-rational check in the LawN256/CeilingLaw256 architecture with one added row family. The near-CUE witnesses extend the same format — the pinning rows are rational box constraints on the stored form factors, and both witnesses already pass exact `Fraction` re-verification against the true $\tau_2 = 1$ box.

A The witness data

Tier-1 witness (`results/a4-m2-gate/witness_N64.json`; cell: $N = 64$, matched budgets $B_1, B_2, B_3 = 84.71214548308222, 135.09544203420174, 304.5549106659001$, $\varepsilon = 0.05$, fuzz = none). All three columns live on the 65-site ψ_1 -zero grid, sites $\theta_k = k$ ($64/65$), marks $\{1, 2\}$, no pairs. Site subsets, summarized (full integer site lists are recorded per column in the JSON, keys `law[].atoms` / `grid_sites`; the canonical description is the clean integer grid — the stored `clean_checks` record the maximum row-value deviation between the clean-grid and optimizer values of $F_1/F'/C'$, at most 4.1×10^{-5} , per column $3.4 \times 10^{-6}/1.6 \times 10^{-5}/4.1 \times 10^{-5}$):

- Column 1 (weight 0.5299906673474316): the vacancy lattice — 64 simples on 64 of the 65 sites (one vacancy). $N_d = 64$; $F_1 = 64$ exactly; $F' = 4128/33$ exactly (hand derivation: $4096/33 + 32/33$, Section 3.4); $C' = 245.4508724$.
- Column 2 (weight 0.16040139164388625): 16 doubles + 32 simples on an explicit 48-site subset. $N_d = 48$; $F_1 = 96$ exactly; $F' = 153.7604346$; $C' = 411.0865130$.
- Column 3 (weight 0.30960794100868216): 32 doubles on an explicit 32-site subset. $N_d = 32$; $F_1 = 128$ exactly; $F' = 135.1959163$; $C' = 301.3541285$ (clean-grid values; the stored law row has $C' = 301.3541693$).

Aggregates (computed from the stored law rows, which saturate the budget edges exactly): $E[F_1] = 88.94775275723633 = B_1(1 + \varepsilon)$; $E[F'] = 132.81813304702743 \in [128.34, 141.85]$; $E[C'] = 289.32716513260516 = B_3(1 - \varepsilon)$; $E[N_d] = 51.52612362142$, $E[N_d]/N = 0.8050956815846875$. Active rows: `F1_hi`, `Cp_lo` only. Exact `Fraction` re-verification: all 10 row checks true, objective deviation 5×10^{-13} .

Near-CUE witnesses (`results/a4-m2-gate/followup-near-cue.json`, keys `witness_N64`, `witness_N128`). $N = 64$: 64 columns, marks $\{1, 2\}$, no pairs, weights 6.2×10^{-5} – 0.0753 , N_d per column 50–60 (4–14 doubles on CUE-like positions); $E|c_j|^2 = j + 1$ at every $j = 1, \dots, 63$ (all upper pinning rows active at $+\tau_2$); $E[N_d]/N = 53.32693798297409/64 = 0.8332334059839699$; only pinning rows (and mass) active. $N = 128$: 128 columns, same class, upper pinning edge at $j + 1 - 10^{-5}$ (interior shrink; `Fraction` re-verification against the true $\tau_2 = 1$ box passes, margin $+1.0 \times 10^{-5}$); $E[N_d]/N = 107.19000239208492/128 = 0.8374219$ (the witness was solved at the interior-shrunk box; its objective sits 4.0×10^{-8} above the Section 5.2 LP optimum 0.8374218534574983); $E[n_{\text{simple}}]/N = 0.6748$.

The three JSON files (`witness_N64.json`, `followup-near-cue.json`, `followup-p1.json`) are the paper’s ancillary files: each contains the full column data, law weights, per-row values, active-row lists, and the rational-verification records.

B Null-model Monte Carlo tables

Closed-form anchors (Theorem 3.5): $m_2(1) = 4/3$, $m_2(1/2) = 13/6 = 2.166667$, $m_3(1/2) = 5$.

Gate sampler (CUE eigenangles via QR-with-phase-fix of complex Ginibre, unfolded to circumference n ; R adaptive to $\text{SE}(m_3) \leq 0.5\%$; batch-means SE with jackknife check):

n	R	$m_2(1)$	$m_2(1/2)$	$m_3(1/2)$
64	4000	1.32363(74)	2.11087(63)	4.75867(362)
128	2000	1.32867(77)	2.13845(71)	4.87769(411)
256	600	1.33115(66)	2.15231(63)	4.93758(373)
512	300	1.33199(86)	2.15958(67)	4.96843(413)
1024	120	1.33364(110)	2.16362(83)	4.98715(476)
2048	48	1.33400(94)	2.16505(83)	4.99323(504)
$1/n$ fit $\rightarrow \infty$		1.33394(63)	2.16682(52)	5.00056(311)
closed form		1.33333	2.16667	5

All three extrapolations anchor-consistent (`anchor_ok` true; halt flags empty).

Audit sampler (independent QR sampler, fresh seeds), side by side at the two shared sizes — agreement within ~ 1 combined σ per entry:

n	$m_2(1)$ audit / gate	$m_2(1/2)$ audit / gate	$m_3(1/2)$ audit / gate
64	1.32317(105)/1.32363(74)	2.11120(84)/2.11087(63)	4.76099(491)/4.75867(362)
256	1.33045(132)/1.33115(66)	2.15305(103)/2.15231(63)	4.94059(616)/4.93758(373)

A third independent sampler (this session, own seeds, alias-free frequency-matrix assembly) gave $m_2(1/2) = 2.11063(135)$ at $n = 64$ and $2.15273(100)$ at $n = 256$, $m_3(1/2) = 4.7589(79)$ and $4.9408(59)$ — each within $\sim 1\sigma$ of both prior samplers — with two-point $1/n$ extrapolation $m_2 \rightarrow 2.1668$, $m_3 \rightarrow 5.0014$. The audit’s third route for $m_3(1/2) = 5$ (real-space 2D determinantal quadrature) gave $4.9896 \rightarrow 4.9932$ under truncation refinement; this session’s independent quadrature gave $4.9924 \rightarrow 4.9960$.

Ladder and occupancy side data (matched $n = 64$ unless noted): null ladder floor $C_{\text{led}}^{\min} = 3.8188$ (3.9119 at $n = 128$); $E[n(V)]/n$ on $V = [2, 3, 4, \dots] = [0.23868, 0.01364, 0, \dots]$; $\max |\text{eig}|$ over 4000 draws = 3.977. Cross-term (occupancy ≥ 2) shares: Frobenius 0.52626, cubic 0.78986, against the analytic 54% (7/6 of 13/6) and 80% (4 of 5; of which the strictly two-point share is 3.5 and the three-point term T_3 the remaining 0.5).

Provenance, data and code

What kind of document this is, and what the internal terms mean. This paper is the publishable output of a structured research program on the Riemann hypothesis, and a few of its terms are that program’s rather than the field’s. They are used below and are defined here once. A *gate* is a pre-registered decision procedure: its specification — objective, adversary class, constraint rows, decision rule and stopping criterion — is fixed and recorded *before* any computation is run, so that neither branch of the outcome can be chosen after the fact. An *adjudication* is an independent adversarial review of a proposal by parties who did not write it, returning a verdict and a list of mandatory repairs; the one referred to here returned *survives with repairs* and prescribed the corrected gate that this paper executes, together with seven numbered repairs (R1–R7). A *referee pass* is a further independent review of the finished document; two were run on this one and their outcome is the Revision record below. Dates are given in place of the program’s internal session numbering.

Data and code availability. Every number in this paper comes from one of three places: the pre-registered gate record, the verification suite written for this paper, or the Lean development. All three are version-controlled and are available in the program repository, <https://github.com/x67ai/riemann-rh-program> (state as of commit `a5ccf63`). Paths below are relative to `anthropic/rh-program/` in that repository. One deliberate exception: the program’s PDF corpus of fetched literature is not redistributed, for copyright reasons, and nothing in this paper depends on it — every external claim is cited to its published source. The components are:

- *The formalization record* (`results/a4-no-go/formalization-status.md`) behind Section 10: the environment (Lean 4.33.0-rc2, Mathlib rev 51e6992), the build result, the theorem-by-theorem map to this paper’s numbering, the `#print axioms` output for all twelve machine-checked theorems, and the reproduction recipe.
- *The adjudication record* (`results/adjudication-A4.json`): the verdict, repairs R1–R7, the garnish and squeeze computations, and the prescription of the corrected gate. This is the contract the gate was run against.
- *The pre-registered gate specification* (`results/a4-m2-gate/SPEC.md`): kernels and units, null budgets and their closed forms with derivations, the adversary class and its realizability argument, the row system and capacity constant, the decision rules, and the flags. Fixed before any run.

- *The run report* (`results/a4-m2-gate/RUN-REPORT.md`): the verdict, the grid-decoupling mechanism, the sanity checks, every run and number produced, the witness, and every deviation from the specification.
- *The independent adversarial audit* (`results/a4-m2-gate/AUDIT.md`), by a party who did not run the gate: an independent Monte-Carlo check of the null budgets, independent re-derivation of the budget identities and the block law, recomputation of the witness, LP re-solves over all 4003 columns, and eight recorded attempts to break the result.
- *The follow-up report* (`results/a4-m2-gate/FOLLOWUP-REPORT.md`): the near-CUE runs at $\tau_2 = 1$ for both N , the p_1 objective, and the riders attached to the use of [BGSTB24].
- *The raw data* (`results/a4-m2-gate/*.json`, `runs/*.json`): the gate result, the witness law at $N = 64$, the near-CUE and p_1 follow-up outputs, and the full scan records at $N = 64$ and $N = 128$ (including the tight- ε , Tier-2, window and λ' -exploration runs, and the $\lambda' = 1/2$ budget derivations), all as JSON.
- *Supplementary material*: the six proof units incorporated as Sections 3, 4.1 and 7, with their verification scripts and JSON outputs.
- *Supplementary material*, `pair-channel.md` — the pair-interference closure (Sections 4.2–4.3), with its independent verification suite `verify/pairchan/*.py` and `pairchan-verify-out.json`.
- *Supplementary material*, `data-tables.md` — the provenance-tagged data digest from which every number in this paper is cited; its Section 10 records the two cite-safety notes honored here (the 940-record maximum is the $N = 128$ value 8.94×10^{-13} ; the Tier-2 level $0.833377/0.8333622/0.8332334$ drift across dictionary snapshots is common-mode, the equality $\delta'_0 = 0$ being snapshot-independent).
- *The direction record* (`directions/A4-lindeloef-lock.md`): the original proposal retained verbatim, the review verdicts, the adjudication, and the decided-gate section this paper executes.

Correction of record. The “+8 vs 0” doubles-vs-pairs cubic separation premise of the original proposal, and the adjudication’s own phase model $((+1, -1)$ at $uL = 1$ with cubic ~ 0), are false under the true block law: Proposition 3.2 gives pair cubic charge $2m^3(1 + 3A^2) \geq 8m^3$ at every depth, and Theorem 3.4 places pairs and doubles on the same affine plane $c = 3F - 2$. The campaign transcript’s recorded block facts (pair eigenvalues $mL^2a(1 \pm A)$) were correct throughout. This correction was executed in the direction file at gate time and is re-proven here from scratch.

Revision record (2026-08-26)

Both referee reports (*supplementary material*, `referee-1.md`, mathematics; `referee-2.md`, evidence and scope) returned PASS WITH REPAIRS. Every repair below was applied only after re-verifying the underlying fact against the recorded evidence (or re-deriving it); the two digest-level errors were also corrected at their source files. Notation: R1-M/m = referee 1 major/minor, R2-M/m = referee 2 major/minor.

Applied (majors).

- R1-M1 = R2-M1 (pair-channel closure overclaimed as “the gate’s entire admissible class”) → abstract sentence, headline 1.4(8), Section 4.3 title, and the 4.3 Consequence paragraph rewritten to the proved coverage (single-pair to $w \leq 2.8$; multi-pair to $w \leq 0.82$ unconditionally, $w \leq 1$ modulo the certified crowding constant [that clause re-scoped to $w \leq 0.98$ by the 2026-08-27 dated revision — see the revision note at the top]; equal-depth pair-only at every depth), with mixed-depth deep-probe multi-pair columns explicitly assigned to numerics plus the 8/9 backstop, and an added sentence stating that class-level exactness of $\delta_0 = 0$ is proved on the closure’s regimes and heuristic (pricing-backed, backstopped at 20/27) on that residue.
- R1-M2.1 = R2-m3 (near-CUE pinning precision) → Section 5.3 “to 10^{-13} ” corrected to “to 2×10^{-12} (maximum deviation 1.84×10^{-12} , at $j = 62$)”; re-verified from `followup-near-cue.json` (15 of 63 rows exceed 10^{-13}). Source corrected in `data-tables.md` Section 4.2 with a correction tag.
- R1-M2.2 = R2-m2 (`clean_checks` misdescribed) → Appendix A now describes `max_dev` as the maximum row-value deviation between clean-grid and optimizer values, all three columns quoted ($3.4 \times 10^{-6}/1.6 \times 10^{-5}/4.1 \times 10^{-5}$); the omitted third value restored. (R2-m2(c)’s rigid-rotation observation was NOT added: not re-verified for this paper.)
- R1-M2.3 = R2-m1 (witness table column 3 mixed provenance) → Section 4.4 table and Appendix A column 3 now display the clean-grid $C' = 301.3541285$ consistently with columns 1–2, with an explicit provenance note: table rows are clean-grid values, aggregates are the stored law rows (which saturate the budget edges exactly; re-verified $E[C'] = B_3(1 - \varepsilon)$ to 10^{-13} from `witness_N64.json`).
- R1-M3.1 (operating-point scope) → one-sentence definition of the title phrase (“the $\lambda' = 1/2 + o(1)$ system fixed by the adjudication, R4 $\theta < 1$ strict; $\lambda' \geq 0.6$ is the quarantined Section 8.4 payoff curve”) added at the end of Section 1.3.
- R1-M3.2 (MVT-4 parenthetical misattached) → Section 9(3)(i) restated: the MVT-4 extension to $\lambda \in (1/2, 2/3)$ is Theorem 1(i)’s fixed- λ divergent cutoff per repair R3 (re-verified in `adjudication-A4.json`), a divergent-cutoff statement subject to Theorem 7.4’s vacuity, not the all- V ladder.
- R2-M2 = R1-m1 (headline 1 ε axes) → corrected to $\varepsilon \in \{0.02, 0.05, 0.10\}$ with the tight- ε runs labeled as separate (verified against `runs/scans_N64.json` / `runs/tighteps_N64.json`).
- R2-M3 (“all $n(V)$ vectors zero” false) → Section 4.4 now states the true ladder slack: $E[n(2)] = 3.50$, $E[n(3)] = 0.94$ against caps 400 and 79.0, higher rows zero, vacancy column’s $n(V)$ identically zero (recomputed from `witness_N64.json` law weights and `nV` vectors). Source corrected in `data-tables.md` Section 3 with a correction tag.
- R2-M4 (unsourced LP refinement values 0.3985/0.3991) → the refinement runs were re-executed and persisted: `verify_t5.py` extended with the (2000 pts, $h_{\max}$ 2000), (4000 pts, $h_{\max}$ 2000), (4000 pts, $h_{\max}$ 8000) cells and re-run, regenerating `verify_t5.out.json`; the persisted tail-corrected values 0.398498/0.399105 reproduce the prose 0.3985/0.3991 exactly, and Section 7.1 now cites the sequence with its grid parameters.

Applied (minors).

- R1-m2 → remark added after Theorem 7.4’s proof: capping the garnish height at $h_0 = \min(\lfloor V_0/2 \rfloor, \lfloor N^{1/4} \rfloor)$ removes the $V_0 = o(N^{1/3})$ hypothesis (re-derived: $h_0 \rightarrow \infty$, $h_0^3 \leq N^{3/4} = o(N)$, $h_0 < V_0$), licensing the “any divergent cutoff” headline as stated.
- R1-m3 → Appendix A notes the $N = 128$ near-CUE witness objective sits 4.0×10^{-8} above the Section 5.2 LP optimum (interior-shrunk box; recomputed from `followup-near-cue.json`).
- R1-m4 → Theorem 4.9’s proof now derives the finite-circle pair block $B_p = m(pq^\dagger + qp^\dagger)$ and its $n_+ \leq 1$ via Cauchy–Schwarz (re-derived), instead of citing Proposition 3.2 across models.
- R1-m5 → (W2) now fixes $\kappa = \kappa(1) = 4/3$.
- R1-m6 → Theorem 7.4’s simple-atom fraction renamed $\delta'_0 \rightarrow s_0$ (collision with the near-CUE marginal removed).
- R1-m7 → Section 2.4 now exhibits the fractional-mark failure of the 5/6 baseline itself (mark-4/3 atoms on the grid: $F_1 = (4/3)N$, $N_d = (3/4)N$; verified) and scopes the Section 4.2 pointer to the pair-channel form.
- R1-m8 → “occupancy-2 cross-terms” corrected to “cross-terms (occupancy ≥ 2)” in Section 2.5 and Appendix B (with the $3.5 + 0.5$ two-point/three-point split of the cubic share noted).
- R1-m9 → Marg’s unique sign change scoped to $(0, 1]$, the interval on which the closed forms are proved.
- R1-m10 → Section 8.4 label corrected to “per repair R4 and merge-guidance item (4)”.
- R2-m4 → Section 5.2 build-out wording: 6,457 columns at the matched cell, 6,739 after the asymptotic re-centering (verified in `followup-near-cue.json` N128 block).
- R2-m5 → ledger verification constant corrected to 2.2×10^{-11} (stored `pairchan_verify_out.json` `I_identity_maxerr = 2.18 \times 10^{-11}`); also corrected at source in `pair-channel.md`.
- R2-m6 → the $N = 128$ “ $S = 100$ ” pricing claim is now labeled RUN-REPORT-recorded with no stored verify block.
- R2-m7 → [BGSTB24] v1 date corrected to the arXiv stamp 7 Jun 2023 (read off the PDF’s first page).
- R2-m8 → abstract now separates the two benchmarks’ evidence bases (940-record grid for N_d ; two converged cells plus cross-checks for p_1).
- R2-m9 → subtitle changed to “for the two-moment system at bandwidth one” (drops the audit-flagged “bandwidth-one ceiling” phrase).
- R2-m10 → Section 8.2 specifies the centering difference (cubic centers 5%, F_1 0.7%, F' 2.6%).
- R2-m11 → “every intermediate dictionary state” corrected to “every recorded dictionary re-solve (1,842/6,457/6,739 columns)” (verified: three recorded re-solves, deltas $7.8 \times 10^{-16}/0.0/0.0$).

- R2-m12 $\rightarrow$ the 0.85082 citation now notes the [BHB13] input (verified verbatim in v5_p14.txt).
- R2 courtesy (prior-art sweep) $\rightarrow$ “see also arXiv:2501.14545” added to the [BGSTB24] entry, attributed to the evidence referee’s logged sweep.

Rejected or partially rejected.

- R2-m2(c) (stored float positions differ from the grid by a common rigid rotation): omitted rather than asserted — the claim was not re-verified in for this paper, and nothing load-bearing needs it; the corrected row-value description suffices.

No other finding was rejected. No FATAL findings were reported by either referee; every repair above is wording, citation, or provenance level, and none changes any theorem, proof, number, or the verdict $\delta_0 = 0$.

References

- [P] L. Alpöge and R. Furman, *More than two thirds of the zeta zeros are simple and on the critical line*, arXiv:2608.13637 (v1, 13 Aug 2026; v2, 19 Aug 2026) — the parent paper, with its sorry-free Lean 4 formalization (Theorems A–E; lemmaR.tight in Zeta23/ZeroSide/TightMult.lean; the PairCeiling library, LawN256/CeilingLaw256; Theorem 5.8/Remark 5.10 prime-side moments). Program digest: results/full-map.md. **Version pin (this matters for the section numbers below)**. The program worked from two on-disk versions, whose sectioning differs and which this paper cites separately: (i) the 35-page version, *More than two thirds of the zeros of the Riemann zeta function lie on the critical line* (anthropic/zeta-two-thirds.pdf), whose §7.5 “Limits of the method” carries the item (e) quoted in Section 1.2 below; and (ii) the 17-page v5, *...are simple and on the critical line* (anthropic/zeta-two-thirds-v5.pdf; text extracts used verbatim: sources-extracted/v5_p13.txt, v5_p14.txt), in which the same material appears, in a shorter form, as §7.2(e), §7.2 there being titled “Sharpness and limits of the method”. Where this paper writes §7.5(e) it means (i); where it writes §7.2(e) it means (ii). Both readings are given in Section 1.2 so that neither pointer can be misread.
- [BGSTB24] S. A. C. Baluyot, D. A. Goldston, A. I. Suriajaya, C. L. Turnage-Butterbaugh, *An unconditional Montgomery theorem for pair correlation of zeros of the Riemann zeta function*, arXiv:2306.04799 (v1, arXiv stamp 7 Jun 2023); Acta Arith. **214** (2024), 357–376; doi:10.4064/aa230612-20-3. On-disk copy: fetched-r3/r3s-06-bgstb-2306.04799.pdf (210,602 bytes; fetch provenance in FOLLOWUP-REPORT Section 4; the v1 stamp read off the PDF’s first page). Cited for: Theorem 1 (unconditional pointwise $F(\alpha)$ on the closed band $0 \leq \alpha \leq 1$, depth-weighted, multiplicity-counted), the Lemma 5 kernel class, and the Remark after Theorem 2 (rider 1). See also the same authors’ sequel on the horizontal distribution of zeros, *Pair correlation of zeros of the Riemann zeta function I: proportions of simple zeros and critical zeros*, arXiv:2501.14545 (v1, 24 Jan 2025; v2, 21 Nov 2025; unpublished as of August 2026) — surfaced by the evidence referee’s logged prior-art sweep, 2026-08-26. (**Dated correction, 2026-08-27:** this entry previously called it a “second-moment sequel”; the paper is about the horizontal distribution of zeros under a narrow-vertical-box hypothesis, not about second moments. Corrected against its own arXiv record.)

- [LR20] J. C. Lagarias and B. Rodgers, *Higher correlations and the Alternative Hypothesis*, Q. J. Math. **71** (2020), no. 1, 257–280; doi:10.1093/qmathj/haz043; arXiv:1905.12123. On disk: `fetchd/w-09`. Cited in Section 1.5: they prove that the currently known band-limited n -level correlation data for zeta zeros — Montgomery ($n = 2$), Hejhal ($n = 3$), Rudnick–Sarnak ($n \geq 4$) — does **not** rule out the Alternative Hypothesis, by exhibiting a point process on $(1/2)\mathbf{Z}$ matching the sine process against all such test functions. Tao obtained the same conclusion independently, by different methods (cited there).
- [LR21] J. C. Lagarias and B. Rodgers, *Band-limited mimicry of point processes by point processes supported on a lattice*, Ann. Appl. Probab. **31** (2021), no. 1, 351–376; doi:10.1214/20-AAP1592; arXiv:1907.03391. Cited in Section 3.4: their Theorem 1.4 and Proposition 3.4 show that above the Nyquist bandwidth the mimicking correlations are unique, while **at** critical sampling they are not — the same degeneracy this paper’s Theorem 3.9 exploits, in the same place. The grid used here (spacing $a = N/(N + 1)$, band $B = 1/2$) falls in the region their Theorem 1.8 leaves undetermined; the witness here matches finitely many trace rows, not all correlations, so there is no conflict, and their negative results explain why the absorption cannot be upgraded to full n -level mimicry.
- [Mon73] H. L. Montgomery, *The pair correlation of zeros of the zeta function*, in Analytic Number Theory (St. Louis, 1972), Proc. Sympos. Pure Math. 24, Amer. Math. Soc., 1973, 181–193. The source of the extremal values $5/6$ and $2/3$ whose bandwidth-one analogs appear in Theorem 4.3 and Corollary 4.4.
- [Hej94] D. A. Hejhal, *On the triple correlation of zeros of the zeta function*, Internat. Math. Res. Notices **1994**, no. 7, 293–302; doi:10.1155/S1073792894000334. The $n = 3$ correlation input named in Section 1.2.
- [RS96] Z. Rudnick and P. Sarnak, *Zeros of principal L -functions and random matrix theory*, Duke Math. J. **81** (1996), no. 2, 269–322; doi:10.1215/S0012-7094-96-08115-6. On disk: `fetchd/w-16`. The n -level range that names this paper’s operating window. **Scope, relevant to flag F2 (Section 8.3)**: their unconditional n -level theorem is for smooth test functions; taking a characteristic function requires RH. Section 3.3’s determinantal ρ_3 and the $c_3(\lambda')$ identification invoke the smooth statement.
- [CGG98] J. B. Conrey, A. Ghosh and S. M. Gonek, *Simple zeros of the Riemann zeta-function*, Proc. London Math. Soc. (3) **76** (1998), 497–522; doi:10.1112/S0024611598000306. The origin of the second integrality level that Lemma 4.2 and Theorem 4.3 consume, made unconditional in the parent’s §7.2(c).
- [BHB13] H. M. Bui and D. R. Heath-Brown, *On simple zeros of the Riemann zeta-function*, Bull. London Math. Soc. **45** (2013), no. 5, 953–961; doi:10.1112/blms/bdt026; arXiv:1302.5018. On disk: `fetchd/w-11` (the preprint). The conditional simple-zero input quoted in Section 1.2; their unconditional-on-RH distinct-zeros constant is 0.84665.
- [CGdL20] A. Chirre, F. Gonçalves and D. de Laat, *Pair correlation estimates for the zeros of the zeta function via semidefinite programming*, Adv. Math. **361** (2020), 106926; doi:10.1016/j.aim.2019.106926; arXiv:1810.08843. The semidefinite-programming line for pair-correlation extremal problems, of which this paper’s LP is the bandwidth-one marked-configuration analog. Two specific contacts. (i) Their inequality (7), $2N_s(T) \leq \sum_{\rho}(m_{\rho} - 2)(m_{\rho} - 3)/m_{\rho} = N^*(T) - 5N(T) + 6N_d(T)$, credited there to an argument of Ghosh,

is the published instance of the same per-multiplicity integrality device Lemma 4.2 uses, at the **reciprocal-weighted third** level $(m-2)(m-3) \geq 0$; Lemma 4.2(a) is the **un-weighted second** level $(m-1)(m-2) \geq 0$. (Their semidefinite programming is applied to the auxiliary-function class of their Section 3, not to (7); (7) is elementary integer arithmetic.) (ii) Their Corollary 3 is the corresponding published record for distinct zeros: $N_d(T) \geq (0.8477 + o(1))N(T)$ on RH and $(0.8486 + o(1))N(T)$ on GRH. Both read from arXiv:1810.08843v2 (18 Nov 2019), the final arXiv version; the typeset Elsevier pagination was not consulted.

- [GdLL25] F. Gonçalves, D. de Laat and N. Leijenhorst, *Multiplicity of nontrivial zeros of primitive L -functions via higher-level correlations*, Math. Comp. **94** (2025), no. 354, 2041–2058; doi:10.1090/mcom/4005; arXiv:2303.01095. The closest published third-party work to this paper: the same higher-level-correlation-versus-multiplicity question, attacked by semidefinite programming rather than by a marked-configuration LP. Cited in Sections 1.2 and 3.2. Their equation (5), $N_n(T) = \sum_k S(n, k)M_k(T)$ with $S(n, k)$ the Stirling numbers of the second kind, is the change of basis whose $n = 3$ case is the atom half of Theorem 3.4 here.
- [FGL14] D. W. Farmer, S. M. Gonek and Y. Lee, *Pair correlation of the zeros of the derivative of the Riemann ξ -function*, J. London Math. Soc. (2) **90** (2014), no. 1, 241–269; doi:10.1112/jlms/jdu026. Cited only in Corollary 4.4, to disambiguate a numerical coincidence: their published RH-conditional lower bound for the proportion of distinct zeros is $N_d(T) > 0.8051 N(T)$ (their Corollary 1.4). The two-author 2008 preprint arXiv:0803.0425 is an earlier version whose Corollary 1.4 gives 0.6544; the constant 0.8051 is the published version’s.
- [KLS07] T. Kuna, J. L. Lebowitz and E. R. Speer, *Realizability of point processes*, J. Stat. Phys. **129** (2007), no. 3, 417–439; doi:10.1007/s10955-007-9393-y; arXiv:math-ph/0612075. The truncated-moment/realizability problem that Section 2.4’s adversary-realizability argument re-derives in the finite-circle setting.
- [KLS11] T. Kuna, J. L. Lebowitz and E. R. Speer, *Necessary and sufficient conditions for realizability of point processes*, Ann. Appl. Probab. **21** (2011), no. 4, 1253–1281; doi:10.1214/10-AAP703; arXiv:0910.1710. The sequel, which settles the general finite-order truncated-moment realizability question; Section 2.4’s finite-circle argument is a special case in their setting rather than a new problem.
- [CdLS22] H. Cohn, D. de Laat and A. Salmon, *Three-point bounds for sphere packing*, arXiv:2206.15373 (v1, 30 Jun 2022; unpublished as of August 2026 — checked against arXiv, Crossref, zbMATH and the first author’s own publication list). The comparandum: in the sphere-packing analog the third-order augmentation **does** pay, improving on the two-point Cohn–Elkies bound in several dimensions. That it is worth exactly nothing here is therefore a substantive negative, not a foregone one.
- [KS66] S. Karlin and W. J. Studden, *Tchebycheff Systems: With Applications in Analysis and Statistics*, Pure and Applied Mathematics, Vol. XV, Interscience Publishers (John Wiley & Sons), New York, 1966. The classical Chebyshev–Markov–Stieltjes/Christoffel background for the fact that odd moments do not improve one-sided bounds — the general form of the cubic blindness of Theorem 3.4.