

The Castelnuovo–Severi/Hodge Index Inequality on a Surface from Riemann–Roch and Ampleness: A Non-Circularity Re-derivation

Jay Tyagi

Abstract

This note is a re-derivation of standard material for the record; zero novelty is claimed. Every statement is graduate-textbook algebraic geometry. Its value is the audit trail: the Hodge-index/Castelnuovo–Severi inequality on a smooth projective surface — the positivity engine of the only *positivity mechanism* that has ever proved a Riemann hypothesis, namely Weil’s proof of RH for curves over finite fields — is proved here in full from Riemann–Roch on the surface, Serre duality, and ampleness, with no zeta-function, L -function, or RH input anywhere. The proof is valid over an algebraically closed field of arbitrary characteristic. We itemize the provenance of every input in a step-by-step ledger, verifying that no zeta function, point count, explicit-formula positivity, or statement equivalent to any Riemann hypothesis appears among the inputs. We then state the $\mathbb{F}_q$ -degeneration clause: the inequality proved here is exactly the inequality Weil’s proof of RH for curves consumes, and it is proved before, and independently of, that application — the short discriminant computation recovering $|N_1 - (q + 1)| \leq 2g\sqrt{q}$ is included. In particular, in the function-field template the positivity engine is demonstrably not Weil positivity in disguise: it exists prior to, and is consumed by, the RH proof, not conversely. The historical lineage (Mattuck–Tate, Grothendieck, Hartshorne) is recorded, together with a scoped remark on the diagonal Tate-curve surfaces $E_p \times E_p$ of Connes–Consani’s absolute geometry, where the inequality holds classically but the available correspondence data is prime-blind.

Introduction and conventions

Status (read before citing). This is a **re-derivation of standard material for the record; zero novelty is claimed**. Every statement below is graduate-textbook algebraic geometry. Its value is precisely the audit trail: the Hodge-index/Castelnuovo–Severi inequality — the positivity engine of the only **positivity mechanism** that has ever proved an RH¹ — is proved from Riemann–Roch on the surface, Serre duality, and ampleness, **with no zeta-function, L -function, or RH input anywhere**.

Conventions. X is a smooth projective (geometrically irreducible) surface over an algebraically closed field k of **arbitrary characteristic** (char p included; nothing below is characteristic-zero). Divisors are considered up to numerical equivalence where stated; $K =$

¹“Positivity mechanism” is the operative qualifier: RH for curves over finite fields was *also* proved by the elementary auxiliary-polynomial method of Stepanov (1969) [4], streamlined and completed by Bombieri [5] — a proof consuming only Riemann–Roch on the *curve*: no surface, no Hodge index, no positivity engine. Deligne’s Weil I amplification is likewise a distinct engine. The claim above concerns the Hodge-index positivity route only.

K_X is the canonical divisor; $D \cdot E$ is the intersection number; $h^i(D) = \dim_k H^i(X, \mathcal{O}_X(D))$; $\chi(D) = h^0 - h^1 + h^2$. “ D numerically nontrivial” means $D \cdot E \neq 0$ for some divisor E . For a non-closed base field, all quantities below are stable under base change to the algebraic closure, so the inequalities descend; we work over $k = \bar{k}$ throughout.

1 What is being certified and why

The function-field mechanism template for a Riemann hypothesis loads *all* of its Riemann-hypothesis content on a Hodge-index-type inequality. The known trap is a positivity step that is Weil positivity in disguise, i.e. an inequality whose proof secretly consumes the statement to be proved. The function-field template is the existence proof that the trap is avoidable: there, the master inequality has a proof whose inputs are **Riemann–Roch on the surface, Serre duality, and ampleness** — nothing else. This note re-derives that proof in full, itemizes the provenance of every input (§8), and states the $\mathbb{F}_q$ -degeneration clause (§7): the inequality proved here is exactly the inequality Weil’s proof of RH for curves consumes, and it is proved before and independently of that application. That is the non-circularity claim, now on paper rather than recalled.

2 Inputs, with provenance (and what is not an input)

The proof uses exactly the following, each a standard theorem with no analytic or arithmetic content about zeta or L -functions:

- **(IN1) Riemann–Roch on a surface.** $\chi(D) = \chi(\mathcal{O}_X) + \frac{1}{2}D \cdot (D - K)$. [Hartshorne [1], Thm. V.1.6; valid over any algebraically closed field.]
- **(IN2) Serre duality.** $h^2(D) = h^0(K - D)$. [Hartshorne [1], III.7.6.]
- **(IN3) Ample meets effective positively.** If H is ample and E is a nonzero effective divisor, then $E \cdot H > 0$. [Degree of a nonzero curve in a projective embedding is positive; equivalently the easy direction of Nakai–Moishezon.]
- **(IN4) Ampleness absorbs any divisor.** For H ample and D arbitrary, $D + nH$ is ample for all $n \gg 0$. [Hartshorne [1], Exercise II.7.14(b); standard.]
- **(IN5) Trivial positivity.** $h^0, h^1, h^2 \geq 0$, hence $h^0(D) + h^2(D) \geq \chi(D)$.
- **(IN6) The intersection pairing on $\text{Num}(X)$** is a well-defined, nondegenerate symmetric bilinear form on a finitely generated group. [Hartshorne [1], V.1; nondegeneracy is the definition of numerical equivalence. Finite generation of $\text{Num}(X)$ is the Néron–Severi theorem — proved by Picard-variety/finiteness arguments, no zeta input; it is consumed *only* for $\rho < \infty$ in the signature statement §5, which sits off the load-bearing chain to §§6–7.]
- **(IN7) Nakai–Moishezon criterion (hard direction).** On a surface, if $H^2 > 0$ and $H \cdot C > 0$ for every irreducible curve $C \subset X$, then H is ample. [Hartshorne [1], Thm. V.1.10; the proof is cohomological (Riemann–Roch plus effectivity bootstrapping) — no zeta content. Consumed only in §6, to certify $H = \xi_1 + \xi_2$ ample. An elementary

route avoiding it exists — $m(\xi_1 + \xi_2) = p_1^*(mP) + p_2^*(mP')$ is very ample for $m \gg 0$ as the Segre-composed embedding of very ample divisors on the factors, an (IN4)-grade fact — but the criterion is what §6’s printed verification uses, so it is listed as the input it is.]

Not inputs: the zeta function of any curve or surface; point counts; the Weil conjectures in any form; positivity of any explicit-formula functional; any spectral or Hilbert-space statement; any characteristic-zero comparison. No step of the proof (§§3–6) mentions them. §7’s degeneration clause consumes rationality and the functional equation of $Z(C, T)$ — Riemann–Roch-on-the-curve facts with no RH content — and produces point counts as *outputs* ($N_1 = \Gamma \cdot \Delta$), exactly as itemized in the §8 ledger; §10 mentions the explicit formula only inside a strictly scoped fence, to state what the Tate-curve surfaces cannot host.

3 Lemma (effectivity criterion; Hartshorne V.1.8 lineage)

Lemma 3.1. *Let H be ample on X and let D be a divisor with $D^2 > 0$ and $D \cdot H > 0$. Then nD is effective ($h^0(nD) > 0$) for all sufficiently large n .*

Proof. By (IN1), $\chi(nD) = \chi(\mathcal{O}_X) + \frac{1}{2}nD \cdot (nD - K) = \frac{n^2}{2}D^2 - \frac{n}{2}D \cdot K + \chi(\mathcal{O}_X) \rightarrow \infty$ as $n \rightarrow \infty$, since $D^2 > 0$. By (IN2), $h^2(nD) = h^0(K - nD)$. Since $D \cdot H > 0$, $(K - nD) \cdot H = K \cdot H - n(D \cdot H) < 0$ for all $n \gg 0$; but $h^0(K - nD) > 0$ would make $K - nD$ linearly equivalent to an effective divisor $E \geq 0$ with $(K - nD) \cdot H = E \cdot H \geq 0$ (zero only for $E = 0$, positive otherwise by (IN3)) — impossible. So $h^0(K - nD) = 0$ for all $n \gg 0$, and by (IN5), $h^0(nD) \geq \chi(nD) - h^2(nD) = \chi(nD) \rightarrow \infty$. $\square$

(This is the exact locus where the hypothesis $D \cdot H > 0$ is consumed: it kills the h^2 term. The compressed sketch “ $\chi(nD) \rightarrow \infty$ forces $\pm nD$ effective” silently routes through this Lemma applied to an auxiliary ample divisor; the full route is Theorem 4.1.)

4 Theorem (Hodge index; Hartshorne Thm. V.1.9 lineage)

Theorem 4.1. *Let H be ample on X and let D be a divisor with $D \cdot H = 0$ and D numerically nontrivial. Then $D^2 < 0$.*

Proof. Suppose for contradiction $D^2 \geq 0$.

Case 1: $D^2 > 0$. By (IN4) choose $m \gg 0$ with $H' := D + mH$ ample. Then $D \cdot H' = D^2 + m(D \cdot H) = D^2 > 0$, so Lemma 3.1 applies to D with the ample divisor H' : nD is effective and nonzero (D is numerically nontrivial, so $nD \not\equiv 0$) for some large n . By (IN3), $nD \cdot H > 0$, contradicting $D \cdot H = 0$.

Case 2: $D^2 = 0$. Since D is numerically nontrivial, pick E with $D \cdot E \neq 0$. Replace E by $E' := (H^2)E - (E \cdot H)H$; then $E' \cdot H = (H^2)(E \cdot H) - (E \cdot H)(H^2) = 0$ and $D \cdot E' = (H^2)(D \cdot E) - (E \cdot H)(D \cdot H) = (H^2)(D \cdot E) \neq 0$ ($H^2 > 0$ by (IN3): H ample $\Rightarrow$ some multiple of H is effective and nonzero, and it meets H positively, so $H^2 = H \cdot H > 0$). For an integer n , set $D' := nD + E'$. Then $D' \cdot H = n(D \cdot H) + E' \cdot H = 0$ and $D'^2 = n^2D^2 + 2n(D \cdot E') + E'^2 = 2n(D \cdot E') + E'^2$. Choosing n large with the sign of $n(D \cdot E')$ positive gives $D'^2 > 0$ with $D' \cdot H = 0$. If D' is numerically nontrivial this contradicts Case 1; and D' numerically trivial is impossible since $D'^2 > 0$. Contradiction either way.

Hence $D^2 < 0$. $\square$

5 Corollary (the signature statement)

Corollary 5.1. *On $\text{Num}(X) \otimes \mathbb{R}$ (rank ρ), the intersection form has signature $(1, \rho - 1)$: it is positive definite on the line $\mathbb{R} \cdot H$ and negative definite on $H^\perp$.*

Proof. $H^2 > 0$ gives the positive line. For the negative-definiteness of $H^\perp$: Theorem 4.1 gives $x^2 < 0$ for every nonzero **rational** class $x \in H^\perp$ (clear denominators and apply the Theorem to the resulting integral divisor class; numerical nontriviality is exactly nonvanishing in Num). Negative *semidefiniteness* on the real span follows by density and continuity of the quadratic form. If some nonzero real $x \in H^\perp$ had $x^2 = 0$, then x would lie in the radical of the semidefinite form restricted to $H^\perp$ (an isotropic vector of a semidefinite form is radical); the radical is a rational subspace (the form and $H^\perp$ are defined over $\mathbb{Q}$), so it would contain a nonzero rational class $y \in H^\perp$ with $y \cdot z = 0$ for all $z \in H^\perp$ and $y \cdot H = 0$ — i.e. y numerically trivial (write any w as $\lambda H + w^\perp$), contradicting $y \neq 0$ in $\text{Num}(X) \otimes \mathbb{Q}$. $\square$

6 Corollary (Castelnuovo–Severi on $C \times C'$)

Let C, C' be smooth projective curves over k , $S = C \times C'$, $\xi_1 = \{\text{pt}\} \times C'$, $\xi_2 = C \times \{\text{pt}\}$. Then $\xi_1^2 = \xi_2^2 = 0$, $\xi_1 \cdot \xi_2 = 1$, and $H := \xi_1 + \xi_2$ is ample ((IN7) Nakai–Moishezon: $H^2 = 2 > 0$, and every irreducible curve on S meets ξ_1 or ξ_2 positively and neither negatively). For a divisor D on S write the **bidegree** $(d_1, d_2) := (D \cdot \xi_1, D \cdot \xi_2)$ (for the graph Γ_f of a morphism $f: C \rightarrow C'$, $(d_1, d_2) = (1, \deg f)$).

Corollary 6.1 (Castelnuovo–Severi inequality). $D^2 \leq 2d_1d_2$, with equality only if D is numerically equivalent to $d_2\xi_1 + d_1\xi_2$.

Proof. Set $D' := D - d_1\xi_2 - d_2\xi_1$. Then $D' \cdot \xi_1 = d_1 - d_1 = 0$ and $D' \cdot \xi_2 = d_2 - d_2 = 0$, hence $D' \cdot H = 0$. By Theorem 4.1 (and its trivial extension: $D'^2 \leq 0$, with < 0 unless $D' \equiv 0$),

$$0 \geq D'^2 = D^2 - 2(d_1(D \cdot \xi_2) + d_2(D \cdot \xi_1)) + (d_1\xi_2 + d_2\xi_1)^2 = D^2 - 4d_1d_2 + 2d_1d_2 = D^2 - 2d_1d_2. \quad \square$$

The classical “equivalence defect” formulation and the positivity of Weil’s trace form on correspondences are algebraic reformulations of this inequality; the historical order (§9) ran through exactly this statement.

7 The $\mathbb{F}_q$ degeneration clause: this is the inequality Weil’s proof consumes

This section states, with the short computation, that the inequality of §6 — proved above from (IN1)–(IN6) only — is the entire positivity input of Weil’s proof of RH for curves over finite fields. **That is the point of this note:** the same inequality that an arithmetic analogue of the mechanism would have to produce is, in its home territory, proved with no RH input, so demanding a Riemann–Roch + ampleness derivation (rather than a posited positivity) is a coherent, historically grounded non-circularity contract.

Let C be a smooth projective geometrically irreducible curve of genus g over $\mathbb{F}_q$, $S = C \times C$ (base-changed to $\overline{\mathbb{F}}_q$ for the intersection theory), $F: C \rightarrow C$ the q -power Frobenius, $\Gamma = \Gamma_F$ its graph, Δ the diagonal, $N_1 = \#C(\mathbb{F}_q)$. The three standard geometric facts consumed — each an intersection-theoretic computation with no zeta input:

- **(W1)** Γ has bidegree $(1, q)$; Δ has bidegree $(1, 1)$.
- **(W2)** $\Delta^2 = 2 - 2g$ (self-intersection = degree of the normal bundle $N_\Delta \cong T_C$) and $\Gamma^2 = q(2 - 2g)$: via the embedding $(\text{id}, F): C \cong \Gamma \subset S$ one has $T_S|_\Gamma \cong T_C \oplus F^*T_C$, so $\deg N_{\Gamma/S} = \deg(T_S|_\Gamma) - \deg T_\Gamma = \deg F^*T_C = q \deg T_C = q(2 - 2g)$.
- **(W3)** $N_1 = \Gamma \cdot \Delta$: Γ and Δ meet transversally because $dF = 0$, so 1 is never an eigenvalue of the tangent map; the intersection points are exactly the $\mathbb{F}_q$ -rational points.

Apply §6 to $D = r\Delta + s\Gamma$ for integers r, s : bidegree $(r + s, r + sq)$, and with $t := \Gamma \cdot \Delta$,

$$D^2 = r^2(2 - 2g) + 2rst + s^2q(2 - 2g) \leq 2(r + s)(r + sq).$$

Expanding and simplifying: $gr^2 - rs(t - 1 - q) + gqs^2 \geq 0$ for all integers r, s , hence (by homogeneity and density) for all real r, s ; the discriminant condition gives

$$(t - (q + 1))^2 \leq 4g^2q, \quad \text{i.e.} \quad \boxed{|N_1 - (q + 1)| \leq 2g\sqrt{q}}.$$

Replacing F by F^k gives $|N_k - (q^k + 1)| \leq 2gq^{k/2}$ for every $k \geq 1$; by the standard power-sum/functional-equation bookkeeping (rationality of the zeta function of C and the $\alpha \mapsto q/\alpha$ pairing — both consequences of Riemann–Roch on the *curve*, again with no RH input) this is equivalent to $|\alpha_i| = \sqrt{q}$ for every Frobenius eigenvalue, i.e. RH for C . (The $\implies$ direction is the generating-function no-cancellation fact: the log-derivative generating function has residue $-m_\alpha$ at $T = 1/\alpha$, so poles at distinct points of equal modulus cannot cancel; the uniform bound then forces radius of convergence $\geq q^{-1/2}$, i.e. $\max_i |\alpha_i| \leq \sqrt{q}$, and the $\alpha \mapsto q/\alpha$ pairing pins equality.) Every input in this chain is listed in §8; none is a zeta positivity.

Consistency note. The bidegrees $(1, q^k)$ of Γ_{F^k} are only the *main* terms $q^k + 1$ of N_k ; the zero side $-\sum_i \alpha_i^k$ lives exclusively in the Δ -intersection (W3). The positivity engine acts on the trace side through §6's inequality — degrees alone never see a zero. (Numerically spot-verified on $y^2 = x^3 + x + 1$ over $\mathbb{F}_7$: bidegrees $(1, 7)$, $(1, 49)$, $(1, 343)$ against counts 5, 55, 380.)

8 Non-circularity audit (step-by-step input ledger)

Step	Uses	Provenance	Zeta/RH content
Lemma §3, growth of $\chi(nD)$	(IN1) RR on surfaces	Hartshorne V.1.6	none
Lemma §3, killing h^2	(IN2) Serre duality + (IN3)	Hartshorne III.7.6; projective degree	none
Theorem §4, Case 1	Lemma + (IN4) $D + mH$ ample + (IN3)	Hartshorne Ex. II.7.14(b)	none
Theorem §4, Case 2	(IN6) linear algebra in $\text{Num}(X)$	Hartshorne V.1	none
Corollary §5	Theorem + $\mathbb{Q}$ -rationality of Num	linear algebra	none
Corollary §6 (CS)	Theorem on $C \times C' +$ (IN7) ampleness of $H = \xi_1 + \xi_2$	(IN7) Nakai–Moishezon, Hartshorne Thm. V.1.10 (cohomological proof)	none
§7 (W1)–(W3)	bidegree/normal-bundle/transversality computations	standard; $dF = 0$	none
§7, discriminant step	§6 applied to $r\Delta + s\Gamma$	quadratic forms	none
§7, closing equivalence	rationality + FE of $Z(C, T)$ from RR on the curve	Weil rationality (RR-based)	none (no RH input; RH is the <i>output</i>)

Statement of the non-circularity claim. The Hodge-index/Castelnuovo–Severi inequality on a surface, in any characteristic, is proved from Riemann–Roch on the surface, Serre duality, and ampleness alone. No zeta function, no L -function, no point count, no explicit-formula positivity, and no statement equivalent to or implied by any Riemann hypothesis appears among its inputs. In particular, in the function-field template the positivity engine is demonstrably **not Weil positivity in disguise**: it exists prior to, and is consumed by, the RH proof — not conversely.

Scope of the claim. This certifies non-circularity of the *function-field template’s* engine. It does *not* assert that any arithmetic substrate exists on which the analogous inequality can be derived, and it confers no credit by itself on any substrate proposal — it is the gate such proposals must pass through, in the form: an index inequality claimed on a substrate must be *derived* from Riemann–Roch plus effectivity/theta input on that substrate and must degenerate to the present statement over $\mathbb{F}_q$.

9 Lineage

- **A. Mattuck, J. Tate**, Abh. Math. Sem. Univ. Hamburg 22 (1958) [2] — the Castelnuovo–Severi inequality derived from Riemann–Roch on the surface (the first modern non-circular derivation).
- **A. Grothendieck**, “Sur une note de Mattuck–Tate,” J. reine angew. Math. 200 (1958) 208–215 [3] — simplification and generalization: the inequality as an instance of the index theorem on an arbitrary surface.
- **R. Hartshorne**, *Algebraic Geometry*, GTM 52, Ch. V §1 [1] — the textbook consolidation used above: Thm. V.1.6 (Riemann–Roch), Thm. V.1.9 (Hodge index), Thm. V.1.10 (Nakai–Moishezon, = (IN7)), Exercises V.1.9–V.1.10 (Castelnuovo–Severi on $C \times C'$ and Weil’s bound $|N - (q+1)| \leq 2g\sqrt{q}$ along exactly the route of §7). N.B. Hartshorne’s theorem and exercise numbering are distinct series: “Thm. V.1.9” (the Hodge index theorem) and “Ex. V.1.9” (the CS exercise) are different items that happen to share a number. Two sub-item pins in this note — “Cor. V.1.8” for the effectivity criterion and the “(b)” of “Ex. II.7.14” — are recalled at the letter level only; the statements themselves are re-proved in full above, so nothing load-bearing depends on them.

10 The $E_p \times E_p$ baseline remark

The following remark is strictly scoped: it concerns only what the classical index inequality does and does not provide on the per-prime Tate-curve surfaces of Connes–Consani’s absolute geometry [6]. On the **diagonal** surfaces $E_p \times E_p$ (with $E_p = \mathbb{C}/(\mathbb{Z} \oplus \mathbb{Z} \cdot i \log p/2\pi)$ the per-prime Tate curve of [6]; $\text{End}(E_p) = \mathbb{Z}$ — the coincidence $(\log p)^2 \in 4\pi^2\mathbb{Q}$ is refuted unconditionally via Gelfond–Schneider, Theorem 5 of the companion note *Products of the per-prime Tate curves of absolute geometry carry no correspondence calculus for the Weil explicit formula*; dated amendment 2026-08-27, this line previously hedged), the Néron–Severi group is generically $\mathbb{Z}\xi_1 \oplus \mathbb{Z}\xi_2 \oplus \mathbb{Z}\Delta$, and the index inequality of §§4–6 **holds classically** — these are classical complex abelian surfaces and nothing in §§3–6 is special to them. But the available correspondence data on them is **prime-blind**: the graphs of multiplication-by- m carry Lefschetz data $\deg([m] - 1) = (m - 1)^2$ — integer polynomials in m with no $\log p$ weight — so no intersection number on these surfaces can host the explicit formula’s transcendently weighted prime terms without hand-inserting the answer key. The cross-prime surfaces $E_p \times E_q$ are dead as substrate (off-diagonal Néron–Severi rank 2: no diagonal, no graphs) and nothing in this note revives them. The baseline therefore contributes exactly one sentence: the index inequality is available on the diagonal Tate-curve surfaces for whatever classical purpose a future argument can state — with zero prime-facing content.

Provenance

Remark. This note originated as a milestone deliverable (“M1, the non-circularity gate”) of an internal research program on positivity mechanisms for the Riemann hypothesis, written 2026-08-26; the program’s internal rules required this re-derivation to exist in writing before any substrate proposal could claim credit for a Hodge-index-type step. The argument

of §§3–4 was independently re-derived by two internal reviewers, and the Mattuck–Tate and Grothendieck lineage items were verified to exist with content as recalled (page range for [3] checked against Crelle/EuDML records; the Bombieri exposé/volume/pages checked against the numdam record SB_1972-1973__15__234_0 and Springer LNM 383). A subsequent independent referee pass (same date) returned “pass with repairs” — no fatal findings, three major and six minor, all applied — after re-deriving every proof line by line, verifying the input ledger of §8 input-by-input with no hidden zeta/ L -function/RH content found, and passing 13 of 13 numerical checks (including the $\mathbb{F}_7$ spot check of §7). The original note carries a traceability table mapping each claim to internal program records (direction files, adjudication and verdict files, and a catalogue of known barriers); those internal references, and the referee’s itemized repair log, are omitted here as program bookkeeping. Two Hartshorne sub-item pins (“Cor. V.1.8”; the “(b)” of “Ex. II.7.14”) remain recalled at the letter level only, pending a physical-copy check; both statements are re-proved in full above. Standard material re-derived for the record; zero novelty claimed.

References

- [1] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics 52, Springer-Verlag, New York, 1977.
- [2] A. Mattuck and J. Tate, On the inequality of Castelnuovo–Severi, *Abh. Math. Sem. Univ. Hamburg* **22** (1958), 295–299.
- [3] A. Grothendieck, Sur une note de Mattuck–Tate, *J. reine angew. Math.* **200** (1958), 208–215.
- [4] S. A. Stepanov, The number of points of a hyperelliptic curve over a finite prime field, *Izv. Akad. Nauk SSSR Ser. Mat.* **33** (1969), 1171–1181.
- [5] E. Bombieri, Counting points on curves over finite fields (d’après S. A. Stepanov), *Séminaire Bourbaki* 1972/73, Exposé 430, Springer Lecture Notes in Mathematics 383 (1974), 234–241.
- [6] A. Connes and C. Consani, On the absolute geometry of $\mathrm{Spec} \mathbb{Z}$, arXiv:2606.06604 (2026).