

The polarized-Frobenius axiom class: an axiom statement with Davenport–Heilbronn and Epstein witnesses (not a theorem)

Jay Tyagi

August 26, 2026

Abstract

This note records an axiom class together with a witness record; it establishes no theorem. A *polarized Frobenius system* for a Dirichlet series $D(s)$ over a proper adelic curve (in the sense of Chen–Moriwaki, with nef adelic line bundles in the Yuan–Zhang calculus) consists of six axioms: a polarized square, a Frobenius family, a total composition law with primitivity exactly at the primes, effectivity, a transpose/Rosati-type functional-equation template, and a D -attachment clause. We show that the attachment clause cannot be formalized innocently: under a degree reading, an Euler product is forced by definition, and an explicit $\mathbb{P}^1 \times \mathbb{P}^1$ system satisfies axioms (PF1)–(PF4) while touching no coefficient of any D ; under a trace reading, counting data does not factor through composition, and an organizing Lefschetz-tie axiom must be added. On every branch the exclusion of RH-false examples is definitional. With effectivity as an axiom, a cheap monoid computation shows that admissible series have von Mangoldt data supported on prime powers and nonnegative. Exact witnesses are recorded: the Davenport–Heilbronn series and the Epstein zeta function of $x^2 + 5y^2$ fail admissibility, with closed-form von Mangoldt values; but so does the RH-true Dirichlet L -function $L(s, \chi_4)$. The class is therefore a positivity filter, not an Euler-product filter and not an RH filter, and Beurling-type generalized number systems satisfy all of its operative coefficient-level consequences, so the class as printed cannot decide the Riemann hypothesis. All numerical witnesses carry exact closed forms.

1 Introduction and status

This document is an **axiom statement plus a witness record**. It is NOT a theorem, it is NOT “provable today,” and it does NOT establish that the Davenport–Heilbronn and Epstein zeta functions admit no $\mathbb{F}_1$ -model (a claim that would require a formalized notion of $\mathbb{F}_1$ -model and remains open). An earlier “Lemma B” theorem framing of this material was refuted by two independent adversarial reviews (blind replication) and the refutation was upheld on independent re-derivation in an adjudication pass; the review record, and the meaning of the internal terms used below (“killers,” “adjudication,” “referee”), are summarized in the Provenance remark at the end of this note. What survives that process — per the adjudication’s partial overrule of one kill verdict, via a referee-verified minor finding — is exactly this: **with effectivity added as an axiom, the step-2 monoid computation under the degree reading is a correct, cheap formal computation**, so the axiom-format note is a legitimate, nonzero item rather than pure re-dressing. The content remains quasi-definitional: it is a known positivity-barrier witness in a new dialect, whose sharper formal home is the Blomer–Leung monoid converse [1].

Role. This note is the **axiom citation** for any argument that wants to consume “multiplicativity as geometry” — nothing more. It supersedes the earlier “Lemma B (target statement)” framing, which is retained in the program record as the immutable historical version.

Binding constraints. Every mathematical claim below was either (i) re-derived independently by both adversarial reviews and the adjudication, or (ii) computed and stored with exact closed forms (the Provenance remark traces every number to its source). Claims struck in adjudication (theorem status, “provable today,” discharge of the $\mathbb{F}_1$ -model question, any credit beyond the axiom-format role) do not appear here except in the negative.

2 The axiom class (repaired definition)

Let $S = (K, (\Omega, \nu), (|\cdot|_\omega))$ be a **proper adelic curve** in the Chen–Moriwaki sense [6] (“proper” = the product formula holds). A **polarized Frobenius system** for a Dirichlet series $D(s)$ over S consists of:

- **(PF1) Polarized square.** A fibered square X over S carrying a nef adelic $\mathbb{Q}$ -line bundle $\bar{L}$ in the Yuan–Zhang adelic-line-bundle calculus [12], with the two fiber classes ξ_1, ξ_2 and a correspondence composition $\circ$ and transpose t .
- **(PF2) Frobenius family.** A family of correspondences $\{\Psi_n\}_{n \geq 1}$ on X with $\Psi_1 = \Delta$ (the identity correspondence).
- **(PF3) Composition law and primitivity.** $\Psi_m \circ \Psi_n = \Psi_{mn}$ for all $m, n \geq 1$, and Ψ_p is primitive (not a composition of non-identity members) exactly at the primes p .
- **(PF4) Effectivity** (*axiom added by the adjudicated repair; the original definition omitted it and its step 3 is false without it — general correspondence classes are differences of effective ones and can have negative degrees*). Every Ψ_n is effective.
- **(PF5) FE/duality (transpose/Rosati-type)** (*axiom added by the adjudicated repair; the original definition’s “closed under transpose” was never tied to the Ψ_n by any axiom*). The family is stable under transpose, and a fixed Rosati-type identity ties Ψ_n^t to Ψ_n through the polarization $\bar{L}$, whose numerical shadow is the $s \mapsto 1 - s$ (eigenvalue $\alpha \mapsto q/\alpha$) symmetry of the functional equation of D . Template: in characteristic p the functional equation IS Rosati/Poincaré duality, relating Γ_F^t to the dual “ q/F ” ($\Gamma_F^t \circ \Gamma_F = q \cdot \Delta$ for the purely inseparable Frobenius F of degree q). Over an adelic curve this axiom is stated here at template level only: **no candidate substrate currently instantiates it for any D** , and this note makes no claim about how a sharpened transpose axiom would cut down the model class of §3.2 (unadjudicated).
- **(PF6) D -attachment.** The counting data of $D(s)$ is reproduced by the system. This clause is the definitional joint, and it cannot be formalized innocently: §3 shows that every reading makes the Davenport–Heilbronn/Epstein exclusion **definitional**, which is why this document is an axiom statement and not a theorem.

PF5 status (read together with §3.2 and §7). Two bookkeeping facts about (PF5), stated here so that no reader has to assemble them: **(a)** PF5’s transpose-stability clause already fails for the model system of §3.2 — $\Gamma_{z \mapsto z^n}^t$ has bidegree $(n, 1)$, which is not the bidegree $(1, m)$ of any Γ_m — so that exhibit certifies vacuity of the **(PF1)–(PF4) fragment**, and no transpose-stable instantiation of the full six-axiom class is currently known for any D . This is consistent with PF5’s uninstantiated-template status and changes nothing in the fork of §3: Horn A’s steps 1 and 3 do not use the exhibit. **(b)** In the §7 accounting, PF5 is excluded from “the axiom a

DMV world violates” because it is an uninstantiated template contributing no operative, checkable coefficient-level consequence. If its FE **shadow** were instead imposed directly on D as an analytic requirement, FE-free Beurling systems would fail it — but that would be an analytic condition inserted by hand, not an axiom any substrate enforces, and the class would still sit below the degree-one-rigidity bracket of §7 (no Ramanujan input, no degree-one-rigidity derivation). The positivity-filter classification and the relocation order of §7 are unchanged either way.

3 Why this is an axiom statement and not a theorem: the definitional fork

3.1 Preliminary: bidegrees multiply under composition; traces do not

For correspondences on a square, the bidegrees $(d_1, d_2) = (Z \cdot \xi_1, Z \cdot \xi_2)$ are multiplicative under composition: $d_i(\Psi' \circ \Psi) = d_i(\Psi') \cdot d_i(\Psi)$ (over a generic point the fibers compose; verified on graph models, four pairs, in the adjudication). Intersection numbers against the diagonal are NOT: $(\Psi' \circ \Psi \cdot \Delta)$ is not a function of $(\Psi' \cdot \Delta)$ and $(\Psi \cdot \Delta)$. Concretely, on $\mathbb{P}^1 \times \mathbb{P}^1$ with $\Gamma_n = \text{graph}(z \mapsto z^n)$: $(\Gamma_n \cdot \Delta) = n + 1$, verified two ways (numerical classes: $\Gamma_n \equiv n\xi_1 + \xi_2$, $\Delta \equiv \xi_1 + \xi_2$, product = $n + 1$; fixed points of $z^n = z$: the $n - 1$ roots of $z^{n-1} = 1$ plus 0 and ∞). And $n + 1$ is not multiplicative: $(\Gamma_6 \cdot \Delta) = 7 \neq 3 \cdot 4 = (\Gamma_2 \cdot \Delta)(\Gamma_3 \cdot \Delta)$.

3.2 Horn A — the degree reading

Read (PF6) as: *the degrees of the Ψ_n against the two fibers reproduce the counting data of D* (the reading the original commission’s own template sentence mandated). Then:

1. By §3.1 the attached degree data is **completely multiplicative by definition** — an Euler product is forced by (PF3) before any geometry is consulted.
2. **The class touches no coefficient of any D .** The system $X = \mathbb{P}^1 \times \mathbb{P}^1$ over the adelic curve of $\mathbb{Q}$ with the standard nef bundle $\mathcal{O}(1, 1)$ and $\Psi_n = \Gamma_{z \mapsto z^n}$ satisfies every axiom (PF1)–(PF4): $\Psi_m \circ \Psi_n = \Psi_{mn}$ exactly, Ψ_p primitive exactly at primes, every Ψ_n an effective irreducible curve, bidegrees $(1, n)$. Nothing in (PF1)–(PF4) references the coefficients of D , so the identical system would serve $D = \zeta$, $D = \text{DH}$, or $D = \text{Epstein}$ indifferently. (Both adversarial reviews constructed this counterexample independently; the adjudicator re-verified it. The printed (PF5) already excludes it — its transpose-stability clause fails, Γ_n^t having bidegree $(n, 1)$, not any Γ_m — so the exhibit certifies vacuity of the (PF1)–(PF4) fragment; see the PF5 status paragraph, §2. Whether some sharpened transpose axiom would cut down the model class further is not adjudicated and not claimed.)
3. Therefore, under Horn A, “reproduce the counting data of D ” must itself be an **axiom** — **an admissibility condition on D** , not a property the geometry proves. Under that admissibility axiom, prime-power support and $\Lambda_D \geq 0$ follow by the correct, cheap monoid computation of §5 — **quasi-definitionally**: the only geometric ingredient consumed is that degrees of effective correspondences against a nef class are ≥ 0 , which is a basic property of the Yuan–Zhang calculus [12], not a theorem being put to work.

3.3 Horn B — the trace reading

Read (PF6) as: *the counting data of D is reproduced by intersection against the diagonal, $(\Psi_n \cdot \Delta)$ -type numbers* — the only reading matching where counting data actually lives in the char- p template

(§3.5). Then the original step 2 (“composition law + primitivity $\implies$ prime-power support; formal monoid algebra; no analysis”) is **false as a formal claim**: by §3.1 trace data does not factor through composition ($(\Gamma_6 \cdot \Delta) = 7 \neq 12$ on $\mathbb{P}^1 \times \mathbb{P}^1$), and no formal argument forces $(\Psi_6 \cdot \Delta)$ -type data to vanish. What is needed is a **Lefschetz-tie axiom** (“the counting data of D is trace-organized by the monoid”) — in characteristic p that tie is a THEOREM about the existing cohomology; over $\mathbb{Q}$ it is exactly the missing substrate. Adding it as an axiom makes the D -attachment Euler-organized by definition, so the Davenport–Heilbronn/Epstein exclusion again follows from the definition plus the one-line coefficient witnesses of §6 — definitional once more.

3.4 Exhaustiveness of the fork

The primary dichotomy is formal, not empirical: the counting functional attached by (PF6) either **factors through composition** — through composition-multiplicative invariants (bidegrees and monomials in them; equivalently the induced action on H^0/H^2) — which is Horn A; or it **does not**. Any non-multiplicative reading whatsoever — the H^1 trace or any other — breaks step 2’s formal monoid algebra and therefore needs an organizing tie axiom, which makes the exclusion definitional: Horn B’s analysis applies verbatim to all of them. The H^0/H^2 -versus- H^1 trichotomy is the template interpretation of this split (in every known template the zero side IS the H^1 trace), not the load-bearing case division. Under Horn A the statement is an admissibility definition; under Horn B it needs a Lefschetz-type axiom that makes it an admissibility definition. Hence: axiom statement, on every branch. (Both adversarial reviews, independently; upheld in adjudication.)

3.5 The char- p accounting, corrected

The commission’s template sentence (“the char- p model — degrees of graph-of-Frobenius powers on $C \times C$ reproducing point counts — is the template”) was **factually wrong** and is corrected here. On $C \times C$ over $\mathbb{F}_q$, the bidegrees of Γ_{Frob^k} are $(1, q^k)$ — these are only the **main terms** of $\#C(\mathbb{F}_{q^k}) = q^k + 1 - \sum_i \alpha_i^k$. The zero side $-\sum_i \alpha_i^k$ lives **exclusively in the Δ -intersection** (Lefschetz: $N_k = (\Gamma_{F^k} \cdot \Delta)$); degrees never see it. A degree-based reading is therefore structurally blind to zeros. Numerically verified in the adjudication on $E: y^2 = x^3 + x + 1$ over $\mathbb{F}_7$: bidegrees $(1, 7)$, $(1, 49)$, $(1, 343)$ versus counts $N_1 = 5$, $N_2 = 55$, $N_3 = 380$ (Frobenius trace $a = 3$: $\alpha + \beta = 3$, $\alpha\beta = 7$).

4 The scope restriction: total multiplicativity pins degree-one Euler shape

(PF3) demands $\Psi_m \circ \Psi_n = \Psi_{mn}$ for **all** m, n — not merely coprime pairs. Two consequences, both stated here as adjudicated obligations:

- **(a) Degree-one Euler shape.** Matched at the level of the Dirichlet coefficients a_n of D , totally multiplicative coefficient data is exactly the Euler shape $D(s) = \prod_p (1 - a_p p^{-s})^{-1}$ — **degree one**. The axiom class as printed excludes every higher-degree Euler-intact L -function (e.g. ζ_K for a quadratic field K has degree-2 Euler factors and non-totally-multiplicative coefficients; it satisfies the coefficient-level consequence class of §5 but not the printed degree-one shape). A relaxed composition law (e.g. coprime-only multiplicativity plus per-prime tower data) is NOT provided by the printed axioms; any future widening is new design, outside this note.

- **(b) Relation to the one-generator function-field monoid.** The char- p template’s Frobenius monoid is generated by ONE element: $\{\Gamma_{F^k} : k \geq 0\} \cong (\mathbb{N}, +)$, indexed by the exponent k (the extension degree), with a single “prime.” The printed multi-prime monoid $\{\Psi_n\}$ over $(\mathbb{N}^\times, \cdot)$ — free commutative on all rational primes — is a different object: the commission’s template was a one-prime template misread as a multi-prime one. Total multiplicativity across distinct primes is exactly the new, restrictive content: at coefficient level it pins the degree-one Euler shape (a); if instead the degree data were matched against Λ -type (prime-power-supported) counting data directly, complete multiplicativity would contradict vanishing at products of distinct primes ($c_{pq} = c_p \cdot c_q \neq 0$ whenever two primes carry nonzero data) unless at most one prime carries data — i.e. exactly the one-generator char- p monoid again. (Both adversarial reviews; referee minor finding; upheld.)

5 The admissibility axiom and the exclusion computation (the correct cheap part)

Definition 5.1 (admissible D). $D(s) = \sum_n a_n n^{-s}$ ($a_1 = 1$) is *admissible* for the polarized-Frobenius axiom class if, writing $-D'/D(s) = \sum_n \Lambda_D(n) n^{-s}$, the counting data satisfies: Λ_D **is supported on prime powers and $\Lambda_D(n) \geq 0$ for all n** (with the FE/duality clause (PF5) named as a further requirement whose substrate-level content is currently uninstantiated). In the strict printed (degree-one) shape, $\Lambda_D(p^k) = a_p^k \cdot \log p$ with $a_p \geq 0$.

The computation (correct, cheap, and quasi-definitional). Under Horn A plus the admissibility axiom, the attachment functional is fixed explicitly: $a_n := d_2(\Psi_n) = (\Psi_n \cdot \xi_2)$, **the second-fiber degree** — the char- p analog of $\deg F^k = q^k$. (This is the multiplicative choice: by §3.1 the individual bidegree components d_i , and monomials in them, are monoid homomorphisms under composition, whereas the degree against the polarization itself — an $(\xi_1 + \xi_2)$ -type pairing — is NOT: on graphs, $(1+n)(1+m) \neq 1+nm$. Everything below runs verbatim for any declared multiplicative monomial in the bidegrees; nothing below may be read against the non-multiplicative polarization pairing.) (PF3) then makes $n \mapsto a_n = d_2(\Psi_n)$ a monoid homomorphism, i.e. totally multiplicative, forcing the degree-one Euler shape (§4(a)); primitivity places the generators exactly at the primes; hence $-D'/D$ has $\Lambda_D(p^k) = a_p^k \log p$ and $\Lambda_D(n) = 0$ off prime powers — formal monoid algebra, no analysis. (PF4) effectivity plus (PF1) nefness gives $a_p = d_2(\Psi_p) = (\Psi_p \cdot \xi_2) \geq 0$, because ξ_2 is nef and Ψ_p is effective, hence $\Lambda_D(n) \geq 0$. This is the step-2/step-3 chain the referee verified by writing out the monoid computation, valid **once effectivity is an axiom** — and it is an unpacking of the definition, not a theorem about D : the geometry contributes only “nef against effective is ≥ 0 .”

What would restore theorem status. Nothing currently available: a genuine exclusion theorem would require a substrate on which the Lefschetz tie of §3.3 is *proved* rather than posited — i.e. the missing substrate itself. Until then the correct citation form is: “ D is inadmissible for the polarized-Frobenius axiom class [this note, §§5–6],” never “ D admits no polarized Frobenius system [theorem].”

6 The witnesses (exact)

All values below are stored with exact closed forms; sources are traced in the Provenance remark. “Inadmissible” always means: fails the §5 admissibility condition, with the named witness.

6.1 Davenport–Heilbronn (RH-false, and inadmissible)

Coefficient system: a_n periodic mod 5, $(a_1, \dots, a_5) = (1, \kappa, -\kappa, -1, 0)$, with the Gauss-sum surd

$$\kappa = \frac{\sqrt{10 - 2\sqrt{5}} - 2}{\sqrt{5} - 1} = 0.28407904384041229602829183239312616909109$$

(41 digits; last digit rounded — the true continuation is ...9091088...).

DH is entire, satisfies a functional equation (verified to 5.6×10^{-40}), and has the verified off-line zero $\rho = 0.808517182456637 + 85.699348485377592i$ (Newton residual 7.6×10^{-41}). Its counting data violates BOTH halves of admissibility. The closed forms below follow from the von Mangoldt recursion $a_n \log n = \sum_{d|n} \Lambda_{\text{DH}}(d) a_{n/d}$ on the periodic coefficient array and match the stored record digit-for-digit at the printed precision:

n	exact form	value	reading
3	$-\kappa \cdot \log 3$	-0.3120927...	negativity at a prime (cheapest witness)
4	$-(2 + \kappa^2) \cdot \log 2$	-1.44223196...	negativity at a prime power
6	$+(1 + \kappa^2) \cdot \log 6$	+1.93635607662...	support off prime powers ($6 = 2 \cdot 3$)
12	$-\kappa \cdot [\kappa^2 \log 6 + (2 + \kappa^2) \log 2 + \log 3]$	-0.762877471988...	negativity at a composite (the original one-line witness)
$= -\kappa(1 + \kappa^2) \cdot \log 12$			

$\Lambda_{\text{DH}}(3) < 0$ and $\Lambda_{\text{DH}}(4) < 0$ are the adjudication-adopted “cheaper witnesses”: negativity is visible on prime powers alone, with no support-off-prime-powers preamble needed.

6.2 Epstein, $Q = x^2 + 5y^2$ ($D = -20$, $h = 2$); RH-false class, and inadmissible

“RH-false class” is classical: for binary quadratic forms of class number $h(D) > 1$, the Epstein zeta function has infinitely many zeros in $\text{Re } s > 1$ (Davenport–Heilbronn [7]) — cited for the label only; nothing in the exclusion computation below depends on it.

Normalization (as stored): $F(s) = \zeta_Q(s)/2$, $b_n = r_Q(n)/2$, $b_1 = 1$; Λ_Q from $-F'/F$ (independent of the constant). Exact rational-log arithmetic, not floating point; numeric rendering at 40 digits. The two witnesses:

- Support off prime powers ($6 = 2 \cdot 3$): the composition-law/primitivity conclusion fails —

$$\begin{aligned} \Lambda_Q(6) &= 2 \log 2 + 2 \log 3 = 2 \log 6 \\ &= +\mathbf{3.583518938456110001624954716761404545446}. \end{aligned}$$

- $\Lambda_Q < 0$: the effectivity/nef-positivity conclusion fails —

$$\begin{aligned} \Lambda_Q(36) &= -4 \log 2 - 4 \log 3 = -4 \log 6 \\ &= -\mathbf{7.167037876912220003249909433522809090892}. \end{aligned}$$

Support off prime powers for $n \leq 100$ at $\{6, 14, 21, 36, 46, 54, 69, 84, 86, 94\}$; negative values at $\{36, 54, 84\}$.

The two structural checks (both passed exactly for all $n \leq 200$):

1. **Genus identity.** $(r_{Q_1}(n) + r_{Q_2}(n))/2 = \sum_{d|n} \chi_{-20}(d)$ exactly, where $Q_1 = x^2 + 5y^2$ (principal) and $Q_2 = 2x^2 + 2xy + 3y^2$ (non-principal) — the coefficient machinery is verified against class-field theory, not merely enumerated.
2. **Contrast object.** $\zeta_K = \zeta \cdot L(\chi_{-20})$ (the full class-group average, Euler product intact) has Λ_K supported on prime powers and ≥ 0 over the same range. What is **proved** is the pair: (a) ζ_Q fails admissibility (the exact witnesses above); (b) the class-group average ζ_K — a theorem-level Euler product, being a Dedekind zeta — satisfies it. The violation therefore **tracks exactly the class-orbit-breaking**: the orbit average passes, the single-class evaluation fails; evaluating a single CM point of the Eisenstein torus period instead of the full class-group orbit is exactly what destroys Euler factorization. That mechanism attribution is the interpretation of (a) + (b) — the mechanism claim of the original Lemma B, surviving here at witness level.

6.3 The over-breadth witness: χ_4 (RH-true, and inadmissible)

$\Lambda_{\chi_4}(3) = a_3 \cdot \log 3 = -\log 3 = -1.0986 \dots < 0$ (computed in the adjudication to 50 digits). An RH-true, degree-one, Euler-intact Dirichlet L -function fails the admissibility condition. The filter is therefore **over-broad in a revealing way**: it is a *positivity* filter, not an Euler-product filter. Any $\mathrm{GL}(2)$ object with some $a_p < 0$ is excluded alongside DH.

7 The Beurling-instantiability and degree-one-rigidity clauses

Beurling/DMV instantiability. The axiom class as printed does **NOT** exclude Beurling/DMV worlds. Beurling generalized number systems (DMV: Diamond–Montgomery–Vorhauer [8]; the Broucke-school factories [4, 3, 5]) carry a full Euler product, multiplicativity at every generalized prime, and $\Lambda_P(n) \geq 0$ pointwise **automatically** — while violating RH maximally. Every **operative** coefficient-level consequence of the axiom class (prime-power support + $\Lambda \geq 0$) holds in these worlds by construction; (PF5), an uninstantiated template, contributes no operative coefficient-level consequence to check — its exclusion from this accounting, and why imposing its FE shadow by hand would change nothing, are stated in the PF5 status paragraph, §2. **The axiom a DMV world violates: currently NONE.** The candidate DMV-violated input, named per the adjudicated repair, is the substrate-level theta-effectivity $h_\theta^0(D) \geq \deg D + O(1)$ (Bost [2]; van der Geer–Schoof [9]) — a substrate-level input no current object supplies. Until such an axiom exists and is instantiated, the class cannot decide RH.

What the filter is. The polarized-Frobenius axiom class as printed is a **POSITIVITY filter separating** $\{\mathrm{DH}, \mathrm{Epstein}\}$ from $\{\zeta\} \cup \{\text{all Beurling worlds}\}$, and it additionally **excludes RH-true signed-coefficient L -functions** ($\Lambda_{\chi_4}(3) = -\log 3 < 0$, §6.3). At the coefficient level the substrate could only ever attach to nonnegative- Λ objects (ζ , ζ_K , Rankin–Selberg squares). It is not an Euler-product filter and not an RH filter.

Degree-one rigidity: the closing bracket. The two-sided calibration any de-novo axiom set must clear: below {FE + Euler product + Ramanujan} at degree one, counterexamples are mass-produced (the Beurling factory above); at them, rigidity — Selberg class degree $1 = \zeta$ + shifted Dirichlet L -functions only, no elements of degree $1 < d < 2$ (**Kaczorowski–Perelli** [10, 11]). The printed axioms do not derive this classification; it is hereby cited as the **closing-bracket target** the axiom class must eventually contain. The gap between Beurling-factory death and Kaczorowski–Perelli rigidity is where a sufficient axiom set must sit; the present class sits below it.

Relocation of RH content. Consequently, **all RH content must sit in the substrate + Hodge-index composite** (the substrate-and-positivity chain of the surrounding program; see the Provenance remark for its internal names). This note carries credit **ONLY** in the axiom-format role: it names the axioms a future substrate must instantiate and records which objects provably cannot satisfy their coefficient-level consequences. Arguments consuming “multiplicativity as geometry” should cite this note for the axiom statement and the **Blomer–Leung beyond-endoscopy monoid converse** [1] as the sharper formal home of the same exclusion — one axiom, two dialects.

Consistency rider. This section implements, in writing, the standing rider on the underlying barrier records: any future axiom-level filter of this kind must add FE/duality + degree-one-rigidity clauses and relocate RH content to the substrate + generator composite.

8 What this note may NOT be cited for (adjudicated exclusions)

1. **Not “provable today.”** Struck by the adjudication; the phrase may not be revived for any part of this material beyond the §5 monoid unpacking, which is quasi-definitional.
2. **Not a theorem.** “DH/Epstein admit no polarized Frobenius system” is not a theorem under either reading of the definition (§3); the citable statement is inadmissibility (§§5–6).
3. **Not a discharge of the $\mathbb{F}_1$ -model question.** The question asks for a proof that DH/Epstein have no *model* over $\mathbb{F}_1$; that requires a formalized notion of $\mathbb{F}_1$ -model and remains open.
4. **No credit beyond the axiom-format role.** In particular: no claim that the axiom class excludes Beurling/DMV worlds (§7 states the opposite); no vacuous “zeta instantiates the axioms” credit (the $\mathbb{P}^1 \times \mathbb{P}^1$ system of §3.2 shows axiom-level instantiation can be worthless); no full credit while no zeta instantiation of the axioms exists at any rank.
5. **Not a revival of per-prime-fiber assemblies.** Nothing here bears on per-prime-fiber assemblies (products of elliptic curves attached to distinct primes); those constructions are excluded on independent grounds recorded in the surrounding program and remain excluded.

Provenance. This note originated as deliverable M0 (axiom-format note) of the C3-r direction of an internal research program on the Riemann hypothesis, written in Session 6, 2026-08-26, per a binding adjudication of 2026-08-26. In the program’s terminology: the “killers” are two independent adversarial reviewers who refuted the original Lemma B theorem framing by blind replication (**results/verdicts-c3d1.json**, kill:C3 fatal 2, kill2:C3 fatal 1); the “adjudication” is the binding third pass that re-derived the findings and ordered the repairs

(`results/adjudication-C3.json`, the mandatory repair beginning “Reprice M0 ...”; adjudication entries are cited by quoted opening phrase, never by array index); the recommission order “C3-r” is in `directions/C3-geometric-substrate.md`. The consistency obligations engaged in the body correspond to the program’s barrier-catalog entries I.1 (the Blomer–Leung monoid converse as the sharper formal home), I.2 (the Beurling/DMV factory), I.7 (degree-one rigidity), III.21, and IV.10 (the rider implemented by §7; the same entry records the exclusion of the per-prime-fiber assemblies of §8, item 5) in `BARRIER-Z00.md`. The “missing substrate” of §3.3 and §5 is the program’s item M2; the “substrate + Hodge-index composite” of §7 is its M1 gate $\rightarrow$ M2c substrate $\rightarrow$ M3/M4 positivity chain, and the theta-effectivity axiom named there is its M4-level input.

Numeric traceability (paths relative to the program root `anthropic/rh-program/`): the fork, the $\mathbb{P}^1 \times \mathbb{P}^1$ counterexample, bidegree multiplicativity, $(\Gamma_n \cdot \Delta) = n + 1$, the non-multiplicativity $7 \neq 12$, the char- p correction, and the $\mathbb{F}_7$ verification are in `results/verdicts-c3d1.json` and `results/adjudication-C3.json` (computation b); κ (41 digits) and $\Lambda_{\text{DH}}(3), (4), (6), (12)$ in adjudication computation d and three independent computations, digit-for-digit, with the underlying DH machinery in `results/decisive-tests/ccm-dh-filter.json` and `results/ccm-dh-test/`; the Epstein exact pair, 40-digit values, support/negativity lists, genus identity, and ζ_K contrast in `results/c3-m0-epstein/n1_epstein_witness.json` (with its generating script); $\Lambda_{\chi_4}(3) = -\log 3$ in adjudication computation c. The exact expressions $-\kappa \log 3$, $-(2 + \kappa^2) \log 2$, $+(1 + \kappa^2) \log 6$, and $-\kappa[\kappa^2 \log 6 + (2 + \kappa^2) \log 2 + \log 3]$ were unpacked from the standard von Mangoldt recursion on the periodic DH coefficient array; each was checked against the stored decimal record and agrees to all printed digits. The Epstein closed forms are stored verbatim. An independent referee pass of 2026-08-26 (`results/c3-r/referee-m0.md`) returned PASS WITH REPAIRS — 0 fatal, 3 major, 7 minor; no mathematical error; all six witnesses, both closed-form families, the $\mathbb{F}_7$ accounting, and the $\mathbb{P}^1 \times \mathbb{P}^1$ counterexample were replicated from scratch, and the DH functional equation and off-line zero were independently re-verified to residuals $\leq 10^{-60}$; all repairs were applied the same day, none rejected or deferred.

References

- [1] V. Blomer and W. H. Leung, *A $GL(3)$ converse theorem via a “beyond endoscopy” approach*, Adv. Math. **485** (2026), Art. 110716; arXiv:2401.04037.
- [2] J.-B. Bost, *Theta invariants of Euclidean lattices and infinite-dimensional Hermitian vector bundles over arithmetic curves*, Progress in Mathematics 334, Birkhäuser, Cham, 2020; arXiv:1512.08946.
- [3] F. Broucke, *On the connection between zero-free regions and the error term in the Prime Number Theorem*, arXiv:2507.13780. (Cited for its construction of Beurling zeta functions with infinitely many zeros on a prescribed contour and none to the right of it.)
- [4] F. Broucke, G. Debruyne, and Sz. Gy. Révész, *Some examples of well-behaved Beurling number systems*, Trans. Amer. Math. Soc. **378** (2025), 477–501; arXiv:2309.01567. (Cited for its $[\alpha, \beta]$ -systems of Beurling generalized primes.)
- [5] F. Broucke and J. Vindas, *A new generalized prime random approximation procedure and some of its applications*, Math. Z. **307** (2024), Art. 62; arXiv:2102.08478. (Cited for its discretization of continuous Beurling systems preserving $\Lambda_P \geq 0$.)
- [6] H. Chen and A. Moriwaki, *Arakelov Geometry over Adelic Curves*, Lecture Notes in Mathematics 2258, Springer, 2020; arXiv:1903.10798.

- [7] H. Davenport and H. Heilbronn, *On the zeros of certain Dirichlet series (Second paper)*, J. London Math. Soc. **11** (1936), 307–312.
- [8] H. G. Diamond, H. L. Montgomery, and U. M. A. Vorhauer, *Beurling primes with large oscillation*, Math. Ann. **334** (2006), 1–36.
- [9] G. van der Geer and R. Schoof, *Effectivity of Arakelov divisors and the theta divisor of a number field*, Selecta Math. (N.S.) **6** (2000), 377–398; arXiv:math/9802121.
- [10] J. Kaczorowski and A. Perelli, *On the structure of the Selberg class, I: $0 \leq d \leq 1$* , Acta Math. **182** (1999), 207–241.
- [11] J. Kaczorowski and A. Perelli, *On the structure of the Selberg class, VII: $1 < d < 2$* , Ann. of Math. (2) **173** (2011), 1397–1441.
- [12] X. Yuan and S.-W. Zhang, *Adelic line bundles on quasi-projective varieties*, Annals of Mathematics Studies 221, Princeton University Press, 2026; arXiv:2105.13587.