

A counterexample to Haglund’s monotonic-zeros conjecture for the Riemann Ξ approximants

Kunal Tyagi*

October 1, 2026; revised October 3, 2026

Abstract

Haglund introduced the partial sums $\Xi_N = \sum_{n \leq N} \Phi_n$ of Riemann’s expansion of the Ξ -function in incomplete gamma functions and conjectured that for every N the zeros of Ξ_N in the closed first quadrant, listed by increasing real part, have nondecreasing imaginary parts. The conjecture implies the Riemann hypothesis; Haglund’s computations supported a weak form of it for $N \leq 10$. We show that the conjecture is false for $N = 27$: Ξ_{27} has a real zero in $(3144.8946, 3144.8947)$ and a non-real zero $z_0 = 3143.2206824215 \dots + 0.3152587993 \dots i$ in the first quadrant, whose real part is smaller. The weak form fails as well. Both facts are certified twice, independently, in interval arithmetic (Arb ball arithmetic; outward-rounded intervals with proved truncation bounds), and z_0 is enclosed in a square of half-width 10^{-48} . The same violation occurs at $N = 24$ for the companion approximants $\xi_N = \Xi_N + c_N$. The failure has a mechanism: on the critical line $\Xi_N < \Xi < \xi_N$, so the real zeros of Ξ_N lie in positive lobes of Ξ that rise above a positive level, and a low lobe followed by a higher one near the height $4(N+1)^2$, where the approximant departs from Ξ , can leave a non-real zero below real zeros. The same structure shows that the number of positive real zeros of Ξ_N , counted with multiplicity, is odd, and that $x^2 \Xi_N(x)$ tends to a negative limit on the real axis for every N .

1 Introduction

1.1 Haglund’s approximants and his conjecture

Write $s = \frac{1}{2} + iz$ and let

$$\Xi(z) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s) = \xi(s), \quad (1)$$

Riemann’s Ξ -function, which is even, entire, and real on the real axis; the Riemann hypothesis is the statement that all its zeros are real. Let $\mathcal{Q} = \{z : \operatorname{Re} z \geq 0, \operatorname{Im} z \geq 0\}$ be the closed first quadrant. Haglund [Hag09] starts from Riemann’s representation $\Xi(z) = \int_0^\infty \cos(zt)\varphi(t)dt$ with $\varphi = \sum_{n \geq 1} \varphi_n$, $\varphi_n(t) = \exp(-n^2\pi e^{2t})(8\pi^2 n^4 e^{4.5t} - 12\pi n^2 e^{2.5t})$, and asks what happens when φ is replaced by the finite sum $\sum_{n \leq N} \varphi_n$. With the “hyperbolic gamma function”

$$G(z; a, b) = 4 \int_0^\infty \cos(2zu) e^{2bu - ae^{2u}} du = \frac{\Gamma(b + iz, a)}{a^{b+iz}} + \frac{\Gamma(b - iz, a)}{a^{b-iz}} \quad (a > 0, b \in \mathbf{R}), \quad (2)$$

*hello.jay.tyagi@gmail.com. *Use of AI.* The mathematics in this paper — the derivations, the computations, the verification suite, and the text — was produced by Claude (Anthropic) working under the author’s direction; the author set the objectives, made the technical and editorial decisions, and is responsible for the content. Claude is a tool and is not an author. The complete working record is public: <https://github.com/x67ai/riemann-rh-program>.

where $\Gamma(w, a) = \int_a^\infty e^{-t} t^{w-1} dt$ is the upper incomplete gamma function, this gives Riemann's series $\Xi(z) = \sum_{n \geq 1} \Phi_n(z)$ and Haglund's Ξ -approximates

$$\Xi_N(z) := \sum_{n=1}^N \Phi_n(z), \quad \Phi_n(z) := 2\pi^2 n^4 G\left(\frac{z}{2}; n^2\pi, \frac{9}{4}\right) - 3\pi n^2 G\left(\frac{z}{2}; n^2\pi, \frac{5}{4}\right) \quad (3)$$

([Hag09, (1)–(3), (6), (10)–(14), pp. 1–3]; page and equation numbers refer to the arXiv version). Each Ξ_N is even, entire, and real on the real axis. Haglund defines: “We say that a given function $F(z)$ has monotonic zeros in a region D of the complex plane if, when we list the zeros of F in D by increasing real part, the imaginary parts of the zeros are monotone nondecreasing”, with the side assumption “(We assume F has at most one zero on the intersection of any vertical line with D).” [Hag09, p. 3]. He then states:

Conjecture 1 (Haglund [Hag09, p. 3]). “For $N \in \mathbf{N}$, $\Xi_N(z)$ has monotonic zeros in $\mathcal{Q}$.”

His Proposition 1 reads “Conjecture 1 implies the Riemann Hypothesis” [Hag09, pp. 3–4]; the proof is a Hurwitz-type argument, using that $\Xi_N \rightarrow \Xi$ uniformly on compacta and that Ξ has infinitely many real zeros. His Remark 1 records a weak form: “A weaker form of Conjecture 1, which still implies RH, is that there are no non-real zeros of $\Xi_N(z)$ in $\mathcal{Q}$ whose real part is less than the largest real zero of $\Xi_N(z)$ ”, and reports that “Computations along these lines indicate this weaker form of Conjecture 1 is true at least for $N \leq 10$ ” [Hag09, p. 4], the computations being floating-point ones for which “no attempt has been made to establish rigorous error bounds” [Hag09, p. 3]. The paper was published as [Hag11].

1.2 The result

Theorem 1.1. (a) Ξ_{27} has a real zero $x_1 \in (3144.8946, 3144.8947)$, and a second real zero $x_2 \in (3145.5998, 3145.5999)$.

(b) Ξ_{27} has exactly one zero z_0 , which is simple, in the open square of half-width $3.7 \cdot 10^{-11}$ about $c = 3143.2206824215 + 0.3152587994i$ (here $w + [-r, r]^2$ is the closed square $\{w + a + bi : |a|, |b| \leq r\}$). More precisely (certificate B), $z_0 \in z^* + [-10^{-48}, 10^{-48}]^2$ with

$$\begin{aligned} \operatorname{Re} z^* &= 3143.220682421536585287281295989417800119\dots, \\ \operatorname{Im} z^* &= 0.315258799378214845382286373009271765\dots, \end{aligned}$$

where the full explicit center z^* is given in Section 6.

(c) Consequently $z_0 \in \mathcal{Q}$ is non-real, with $\operatorname{Im} z_0 > 0.3152$ and $\operatorname{Re} z_0 < 3143.2207 < x_1$. Listed by increasing real part, the zeros of Ξ_{27} in $\mathcal{Q}$ do not have nondecreasing imaginary parts: Conjecture 1 is false for $N = 27$. Since z_0 is a non-real zero in $\mathcal{Q}$ whose real part is less than the real zero x_1 , and hence less than the largest real zero, the weak form of Remark 1 is false for $N = 27$ as well.

The pair (z_0, x_1) violates the conjecture under every enumeration of the zeros by increasing real part, so Haglund's side assumption (at most one zero on each vertical line) plays no role. Statements (a) and (b) are established by two independent computer-assisted proofs in rigorous interval arithmetic, described in Section 6: one in Arb ball arithmetic [Joh17, FLINT], one in outward-rounded interval arithmetic [mpm] with every truncation bounded by a proved remainder estimate. Each was first run on a ladder of known cases (two zeros from Haglund's table and a non-real zero of Ξ_1 from his appendix).

1.3 Scope

The paper does not decide Conjecture 1 for any $N \neq 27$: the scan that located the violation at $N = 27$ (Section 5) is numerical, covers a window near one height for each N , and is not a proof that any other N satisfies the conjecture.

1.4 The mechanism

The failure is not an accident of $N = 27$. On the critical line the approximants sit below Ξ by a positive amount: for every $N \geq 1$ and real t ,

$$\Xi_N(t) = \Xi(t) - Q_N(t), \quad Q_N(t) > 0$$

(Theorem 3.3). Hence $\Xi_N < 0$ wherever $\Xi \leq 0$, and the real zeros of Ξ_N lie in the *positive lobes* of Ξ (the intervals between consecutive zeros on which $\Xi > 0$): an even number in each positive lobe (γ_j, γ_{j+1}) and an odd number in the central lobe $(0, \gamma_1)$, where $0 < \gamma_1 < \gamma_2 < \dots$ are the positive real zeros of Ξ . A positive lobe carries real zeros of Ξ_N only if Ξ rises above the level Q_N there. Near the height $4(N+1)^2$ the level Q_N becomes comparable to the size of Ξ , and the heights of consecutive lobes fluctuate; a lobe that stays below the level, followed by a lobe that rises above it, is expected to leave a pair of non-real zeros below real zeros (it does in both cases computed below). At $N = 27$ the positive lobe of Ξ over $(3142.98, 3143.57)$ reaches only 0.411 of the level Q_{27} and the next one, over $(3144.49, 3146.28)$, reaches 1.325 of it: the pair $z_0, \bar{z}_0$ sits over the first, and the real zeros x_1, x_2 lie in the second (Section 5; these lobe figures are floating-point computations, not part of the proof).

The same sandwich gives two facts about the real zeros and the real-axis tail of Ξ_N .

- (i) The number of real zeros of Ξ_N on $(0, \infty)$, counted with multiplicity, is odd for every N (Corollary 3.5).
- (ii) On the real axis $x^2 \Xi_N(x)$ tends to a negative limit for every $N \geq 1$ (Proposition 3.4).

Both agree with the copy of Haglund's paper on the author's web page [Hag11w], dated February 9, 2011: its table lists 1, 7, 15, 31, 53, 79, 113, 155, 207, 263 real zeros for $N = 1, \dots, 10$ [Hag11w, p. 4], and it states that the coefficient of $1/x^2$ in $\Xi_N(x)$ "approaches zero from below as $N \rightarrow \infty$ " [Hag11w, p. 10]. The arXiv version [Hag09] and the journal version [Hag11] have 32 at $N = 4$ and, for the coefficient, "positive for $k \geq 3$ and negative for $1 \leq k < 3$ " [Hag09, p. 10] (his k is N); [Hag11w] corrects both.

1.5 The companion approximants, and why an ordering property

Haglund's Ξ_N differs by a constant from the more obvious truncation of Riemann's formula,

$$\xi_N(s) := \frac{1}{2} + \frac{1}{2}s(s-1) \sum_{n \leq N} g_n(s), \quad \Xi_N(z) = \xi_N\left(\frac{1}{2} + iz\right) - c_N, \quad c_N = \sum_{n > N} (4\pi n^2 - 1)e^{-\pi n^2} > 0$$

(Section 2). On the critical line ξ_N lies *above* Ξ (Theorem 3.1), so its real zeros lie in the negative lobes of Ξ , and the same ordering property fails for it at $N = 24$ (Theorem 7.1). Neither family is real-rooted: no member of the family $\xi_w = \frac{1}{2} + \frac{1}{2}s(s-1) \sum_n w_n g_n$ with $w \not\equiv 0$ and either $0 \leq w_n \leq 1$, $w \not\equiv 1$, or w finitely supported, has all its zeros on the critical line; it has finitely many zeros there and infinitely many off it (Theorem 4.1), and the same holds for every Ξ_N . With polynomially bounded weights, having all zeros on the line forces $w \equiv 1$ (Theorem 4.2). This is why a conjecture

about approximants of this kind has to be an ordering statement such as Conjecture 1, and not real-rootedness; and any approach through real-rootedness in a window that grows with N is a height filtration of the Riemann hypothesis, with nothing to prove beyond the Riemann hypothesis on the new window (Proposition 4.3).

1.6 Prior work

Haglund’s conjecture was stated in 2009 [Hag09, Hag11] and checked by him, in its weak form, for $N \leq 10$. Ahn’s thesis [Ahn12], written under Haglund’s supervision, computes zeros of the component functions of the approximants (incomplete gamma functions, hyperbolic gamma functions, and small linear combinations of the Φ_n) and restates the check: “Computations indicate this weaker form of Conjecture 1.6 [sic: his Conjecture 1.5] is true at least for $N \leq 10$ ” [Ahn12, pp. 10–11]; it tests no Ξ_N beyond $N = 10$. Haglund’s Conjecture 4 concerns the zeros of the pencil $\Xi_k + t\Phi_{k+1}$, $0 \leq t \leq 1$ [Hag09, p. 11]; its first case $k = 1$ is the subject of a 2026 preprint of Baccaro [Bac26], whose record states that “The result covers only the first interpolation.” We found no test of Conjecture 1 beyond $N = 10$, and no refutation of it, in the literature we could reach; a null search is not a proof of absence. Lagarias and Montague study a related Fourier-integral family whose members have only finitely many zeros on the critical line; they ask whether monotone increase of the imaginary parts of the zeros in the first quadrant would be a sufficient condition for the zeros of its derivative ξ to lie on the critical line, and point to Haglund’s paper as a related situation “involving the ξ -function, where such a monotonicity implies the RH” [LM11, p. 24]. Truncations that keep all but finitely many zeros *on* the line are known in another setting: Ki proved that for $y \geq 1$ all but finitely many complex zeros of any partial sum of the Chowla–Selberg formula are simple and on the line [Ki05, Corollary 1], and observed, restating it, that the Riemann hypothesis would follow if the words “but finitely many” could be removed for infinitely many N in the case $z = i$ [Ki06, p. 198]. The approximants here are the opposite case (Theorem 4.1). Finally, Liu, Matiyasevich, Oesterlé and Zaharescu prove, for a different family of approximants to ξ built from truncated Euler products, that “The Bounded Riemann Hypothesis is equivalent to the Riemann Hypothesis plus the Simplicity Conjecture” [LMOZ24, Theorem 1.7]; Proposition 4.3 is the analogue for the approximants of this paper.

1.7 Organization

Section 2 recalls Riemann’s identity and relates the two families. Section 3 proves the positivity and sandwich theorems and their consequences for the real zeros and the tail. Section 4 gives the structural theorems. Section 5 describes the departure height and the lobe scan. Section 6 is the certificate for Theorem 1.1, and Section 7 the companion violation at $N = 24$.

2 Riemann’s identity and the two families

Let $\psi(x) = \sum_{n \geq 1} e^{-\pi n^2 x}$. Splitting $\int_0^\infty \psi(x) x^{s/2-1} dx$ at $x = 1$ and using $\psi(1/x) = \sqrt{x} \psi(x) + (\sqrt{x} - 1)/2$ gives Riemann’s identity [Rie59, Edw74], valid for all $s \in \mathbf{C} \setminus \{0, 1\}$:

$$\Lambda(s) := \pi^{-s/2} \Gamma(s/2) \zeta(s) = -\frac{1}{s} - \frac{1}{1-s} + \sum_{n \geq 1} g_n(s), \quad (4)$$

where

$$g_n(s) = \frac{\Gamma(s/2, X)}{X^{s/2}} + \frac{\Gamma((1-s)/2, X)}{X^{(1-s)/2}}, \quad X = \pi n^2.$$

Hence $\xi(s) = \frac{1}{2}s(s-1)\Lambda(s) = \frac{1}{2} + \frac{1}{2}s(s-1)\sum_{n \geq 1} g_n(s)$. For weights $w: \mathbf{N} \rightarrow \mathbf{R}$ with $|w_n| \leq Cn^A$ for some constants C, A (so that the series converges absolutely and locally uniformly) put

$$\xi_w(s) := \frac{1}{2} + \frac{1}{2}s(s-1)\sum_{n \geq 1} w_n g_n(s), \quad (5)$$

and write ξ_N for $w = \mathbf{1}_{[1,N]}$. Every ξ_w satisfies $\xi_w(1-s) = \xi_w(s)$ and $\xi_w(\bar{s}) = \overline{\xi_w(s)}$, so $\xi_w(\frac{1}{2} + it)$ is real for real t ; and $|g_n(s)| \leq g_n(\operatorname{Re} s)$, since $|\Gamma(a, X)| \leq \Gamma(\operatorname{Re} a, X)$. Throughout, $s = \frac{1}{2} + iz$; on the critical line $z = t$ is real, and we write $\Xi(t) = \xi(\frac{1}{2} + it)$ and $P_w(t) := \xi_w(\frac{1}{2} + it) - \Xi(t)$.

Lemma 2.1. *For every $n \geq 1$, $\Phi_n(z) = \frac{1}{2}s(s-1)g_n(s) + (4\pi n^2 - 1)e^{-\pi n^2}$. Consequently*

$$\Xi_N(z) = \xi_N(s) - c_N, \quad c_N := \sum_{n > N} (4\pi n^2 - 1)e^{-\pi n^2} > 0.$$

Proof. Put $h(w) = X^{-w}\Gamma(w, X)$ with $X = \pi n^2$. Since $iz = s - \frac{1}{2}$, one has $\frac{9}{4} + \frac{iz}{2} = 2 + \frac{s}{2}$ and $\frac{9}{4} - \frac{iz}{2} = 2 + \frac{1-s}{2}$, and the same with 1 in place of 2 for $b = \frac{5}{4}$; so definition (3) reads $\Phi_n = 2X^2[h(\frac{s}{2} + 2) + h(\frac{1-s}{2} + 2)] - 3X[h(\frac{s}{2} + 1) + h(\frac{1-s}{2} + 1)]$. The recurrence $\Gamma(w+1, X) = w\Gamma(w, X) + X^w e^{-X}$ gives $h(w+1) = (wh(w) + e^{-X})/X$, and applying it twice, $2X^2h(a+2) - 3Xh(a+1) = (2a^2 - a)h(a) + (2a - 1 + 2X)e^{-X}$. For $a = s/2$ and for $a = (1-s)/2$ one has $2a^2 - a = \frac{1}{2}s(s-1)$, while $2a - 1$ equals $s - 1$ and $-s$ respectively; adding the two gives the first claim. Summing over $n \leq N$ and using $\sum_{n \geq 1} (4\pi n^2 - 1)e^{-\pi n^2} = \frac{1}{2}$ (differentiate $\theta(1/x) = \sqrt{x}\theta(x)$, $\theta = 1 + 2\psi$, at $x = 1$) gives the second. $\square$

Numerically $c_1 = 1.7180566258547875025 \cdot 10^{-4}$, $c_2 = 5.891258854396012012 \cdot 10^{-11}$, $c_3 = 2.9589851955455663928 \cdot 10^{-20}$, $c_4 = 2.4342009049056017 \cdot 10^{-32}$ and $c_5 = 3.443519624082649 \cdot 10^{-47}$. So Haglund's zero sets are the level sets $\{\xi_N = c_N\}$ of the truncations ξ_N , not their zero sets; the two families have different zeros, and both are studied below.

3 Positivity on the critical line

Theorem 3.1 (positivity). *For every $n \geq 1$ and real t , $g_n(\frac{1}{2} + it) > 0$. Consequently, for weights $0 \leq w_n \leq 1$,*

$$P_w(t) = \frac{1}{2}\left(\frac{1}{4} + t^2\right)\sum_{n \geq 1} (1 - w_n) g_n\left(\frac{1}{2} + it\right) \geq 0,$$

and $P_w(t) > 0$ for every real t unless $w \equiv 1$. In particular $\xi_1(\frac{1}{2} + it) > \xi_2(\frac{1}{2} + it) > \dots > \Xi(t)$ for every real t .

Proof. With $x = e^{2u}$,

$$g_n\left(\frac{1}{2} + it\right) = \int_1^\infty e^{-Xx} 2x^{1/4} \cos\left(\frac{t}{2} \log x\right) \frac{dx}{x} = 4 \int_0^\infty k(u) \cos(tu) du, \quad k(u) = \exp(u/2 - Xe^{2u}).$$

With $y = Xe^{2u} \geq X \geq \pi$ one has $k' = k(\frac{1}{2} - 2y) < 0$ and $k'' = k(4y^2 - 6y + \frac{1}{4}) > 0$, since the larger root of $4y^2 - 6y + \frac{1}{4}$ is $(3 + 2\sqrt{2})/4 = 1.457\dots < \pi$. So k is positive, strictly decreasing and strictly convex on $[0, \infty)$, and tends to 0. For $t \neq 0$, $\int_0^\infty k(u) \cos(tu) du = t^{-1} \int_0^\infty (-k'(u)) \sin(tu) du$; since $-k'$ is positive and strictly decreasing, the pieces of the sine integral over consecutive half-periods alternate in sign with strictly decreasing sizes, and the first is positive, so the integral is positive (this is the argument behind Pólya's criterion). At $t = 0$ the integral is $\int_0^\infty k > 0$. The formula for P_w follows by subtracting $\xi = \frac{1}{2} + \frac{1}{2}s(s-1)\sum_n g_n$ from (5), with $\frac{1}{2}s(s-1) = -\frac{1}{2}(\frac{1}{4} + t^2)$ on the line. $\square$

Corollary 3.2 (lobe law for ξ_w). *Let $0 \leq w \leq 1$, $w \neq 1$. Every zero of ξ_w on the critical line lies in an open negative lobe of Ξ (an interval between consecutive zeros of Ξ on which $\Xi < 0$), and each negative lobe contains an even number of them, counted with multiplicity.*

Proof. At the ends of a lobe $\xi_w = P_w > 0$, and on positive lobes $\xi_w \geq \Xi > 0$. $\square$

The real-zero counts of ξ_N observed numerically are 2, 6, 16, 32, 52, 80 for $N = 1, \dots, 6$, all even. Haglund's approximants lie on the other side of Ξ . For $n \geq 1$ put $k_n(u) = \exp(u/2 - \pi n^2 e^{2u})$ and

$$\tilde{\varphi}_n := k_n'' - \frac{1}{4}k_n = 2y(2y - 3)e^{u/2-y}, \quad y = \pi n^2 e^{2u},$$

which equals $(4\pi^2 n^4 e^{9u/2} - 6\pi n^2 e^{5u/2}) \exp(-\pi n^2 e^{2u})$, one half of the term $\varphi_n(u)$ of Riemann's kernel (Section 1.1).

Theorem 3.3 (sandwich). *For every $N \geq 1$ and real t ,*

$$\Xi_N(t) = \Xi(t) - Q_N(t) < \Xi(t) < \Xi(t) + P_N(t) = \xi_N(\tfrac{1}{2} + it),$$

where

$$Q_N(t) = c_N - P_N(t) = 2 \int_0^\infty \sum_{n>N} \tilde{\varphi}_n(u) \cos(tu) du > 0.$$

Proof. Integrating by parts twice,

$$\tfrac{1}{2}(\tfrac{1}{4} + t^2)g_n(\tfrac{1}{2} + it) = 2(\tfrac{1}{4} + t^2) \int_0^\infty k_n \cos(tu) du = -2k_n'(0) - 2 \int_0^\infty (k_n'' - \tfrac{1}{4}k_n) \cos(tu) du,$$

and $-2k_n'(0) = (4\pi n^2 - 1)e^{-\pi n^2}$; summing over $n > N$ gives $P_N = c_N - Q_N$, and $\Xi_N = \xi_N - c_N = \Xi + P_N - c_N = \Xi - Q_N$ by Lemma 2.1. For $n \geq 2$ one has $y \geq 4\pi$, and then $\tilde{\varphi}_n > 0$, $\tilde{\varphi}_n' = e^{u/2-y} y(-8y^2 + 30y - 15) < 0$ and $\tilde{\varphi}_n'' = e^{u/2-y} y(16y^3 - 112y^2 + 165y - 37.5) > 0$ (because $16y^2(y - 7) > 0$ and $165y > 37.5$). So $\sum_{n>N} \tilde{\varphi}_n$ is positive, strictly decreasing and strictly convex on $[0, \infty)$ for $N \geq 1$, and the argument of Theorem 3.1 gives $Q_N(t) > 0$. $\square$

For $n = 1$ the kernel is not monotone at $u = 0$: the factor $-8y^2 + 30y - 15$ is $+0.291$ at $y = \pi$, so $\tilde{\varphi}_1'(0) = e^{-\pi} \cdot \pi \cdot 0.291 = +0.0395 > 0$. This is why the theorem needs $N \geq 1$.

Proposition 3.4 (the $1/t^2$ tail). *For every $N \geq 1$, with $X = \pi n^2$,*

$$\lim_{t \rightarrow \infty} t^2 \Xi_N(t) = 2 \sum_{n>N} \tilde{\varphi}_n'(0) = -2 \sum_{n>N} (8X^3 - 30X^2 + 15X)e^{-X} < 0.$$

Proof. $\Xi(t)$ decays exponentially, and two integrations by parts in the integral for $Q_N(t)$ in Theorem 3.3 give $t^2 Q_N(t) \rightarrow -2 \sum_{n>N} \tilde{\varphi}_n'(0)$. At $u = 0$, $\tilde{\varphi}_n'(0) = -(8X^3 - 30X^2 + 15X)e^{-X}$, and each summand $8X^3 - 30X^2 + 15X$ is positive for $n \geq 2$. $\square$

Since the full kernel $\sum_{n \geq 1} \tilde{\varphi}_n$ is even (Haglund states it [Hag09, p. 2]; with $K(u) = e^{u/2}\psi(e^{2u})$ the theta relation gives $K(u) - K(-u) = -\sinh(u/2)$, so $K'' - K/4$ is even), $\sum_{n \geq 1} \tilde{\varphi}_n'(0) = 0$, and the limit is also the sum over $n \leq N$ of the coefficients of $1/t^2$ in the individual Φ_n , namely $-2\tilde{\varphi}_n'(0)$; this is how Haglund computes it. The limits are -0.0789975 , $-1.65306 \cdot 10^{-7}$, $-2.78345 \cdot 10^{-16}$, $-5.73947 \cdot 10^{-28}$ and $-1.70746 \cdot 10^{-42}$ for $N = 1, \dots, 5$ (at $t = 8000$, $t^2 \Xi_N(t)$ agrees with them to a relative 10^{-3}). Haglund lists the coefficients of $1/x^2$ in Φ_1 and Φ_2 as $-.01974938206$ and $.01974934121$ [Hag09, (52), p. 10], and to more digits in [Hag11w, (52)]; his values are one quarter

of ours (at $N = 1$, -0.01974938206 against -0.0789975), because his (51) applies his (50) at x although (3) evaluates G at $x/2$; the factor does not affect signs. The arXiv and journal versions conclude that the coefficient of $1/x^2$ in $\Xi_N(x)$ “is positive for $k \geq 3$ and negative for $1 \leq k < 3$ ” [Hag09, p. 10]; the author’s web copy states instead that it “approaches zero from below as $N \rightarrow \infty$ ” [Hag11w, p. 10], which is what Proposition 3.4 shows: the limit is negative for every N and tends to 0. The coefficients of Φ_1 , Φ_2 and Φ_3 nearly cancel: their sum at $N = 3$, of order 10^{-16} , is below the resolution of the printed values of (52) in either copy. Haglund’s use of these coefficients to exhibit non-monotone zeros of the pencil $t\Phi_1 + \Phi_2$ needs only that the coefficient of Φ_2 is positive and that of Ξ_2 negative, and both are.

Corollary 3.5 (real zeros of Ξ_N). *Let $N \geq 1$ and let $0 < \gamma_1 < \gamma_2 < \dots$ be the positive real zeros of Ξ . Every real zero of Ξ_N in $(0, \infty)$ lies in a positive lobe of Ξ . Counted with multiplicity, each positive lobe (γ_j, γ_{j+1}) contains an even number of them and the central lobe $(0, \gamma_1)$ an odd number. Consequently the number of real zeros of Ξ_N in $(0, \infty)$, counted with multiplicity, is finite and odd.*

Proof. Where $\Xi \leq 0$, $\Xi_N = \Xi - Q_N < 0$ (Theorem 3.3); in particular $\Xi_N(\gamma_j) = -Q_N(\gamma_j) < 0$ at both ends of every positive lobe. At the origin $\Xi_N(0) = \Xi(0) - Q_N(0) \geq 0.4971 - c_1 > 0$, since $Q_N = c_N - P_N \leq c_N \leq c_1$, while $\Xi_N(\gamma_1) < 0$. By Proposition 3.4, $\Xi_N < 0$ on $[T_0, \infty)$ for some T_0 , so only finitely many lobes carry zeros, and the zeros of the entire function Ξ_N in $[0, T_0]$ are finite in number. $\square$

The ten entries of Haglund’s table in [Hag11w, p. 4] (Section 1.4) are odd, in agreement with Corollary 3.5. A numerical census of Ξ_4 (real zeros by sign changes, the total by the argument principle, not interval-rigorous) finds 31 real zeros, the entry of [Hag11w], the largest being 103.3679880094135; the arXiv and journal versions print 32 there.

4 Structure: no member is real-rooted

Theorem 4.1. *Let $w \neq 0$, and suppose either (i) $0 \leq w_n \leq 1$ for all n , with $w_m < 1$ for some m , or (ii) w is finitely supported with arbitrary real values. Then ξ_w has only finitely many zeros on $\operatorname{Re} s = \frac{1}{2}$ and infinitely many zeros off it.*

Proof. (a) *Infinitely many zeros.* Since $|g_n(s)| \leq g_n(\operatorname{Re} s)$,

$$|\xi_w(s)| \leq \frac{1}{2} + \frac{1}{2}|s|^2 \sum_n |w_n| g_n(\operatorname{Re} s) = \exp(O(|s| \log |s|)).$$

With n_0 the least n such that $w_n \neq 0$, $\xi_w(\sigma) \sim \frac{1}{2}\sigma^2 w_{n_0} (\pi n_0^2)^{-\sigma/2} \Gamma(\sigma/2)$ as $\sigma \rightarrow +\infty$, since $\Gamma(\sigma/2, X)/\Gamma(\sigma/2) \rightarrow 1$ and the other terms are smaller by $(n_0/n)^\sigma$. So $\log |\xi_w(\sigma)| \geq (\sigma/2) \log \sigma (1 - o(1))$, ξ_w is not $e^{a+bs} P(s)$ with P a polynomial, and by Hadamard’s theorem it has infinitely many zeros.

(b) *Finitely many on the line.* Case (i): by Theorem 3.1, $\xi_w(\frac{1}{2} + it) \geq \Xi(t) + (1 - w_m)\frac{1}{2}(\frac{1}{4} + t^2)g_m(\frac{1}{2} + it)$. From $\gamma(a, X) = X^a e^{-X} \sum_{k \geq 0} X^k / (a)_{k+1}$ and $|(a)_{k+1}| \geq (|t|/2)^{k+1}$ on the line, $\frac{1}{2}(\frac{1}{4} + t^2)g_m(\frac{1}{2} + it) \rightarrow (4\pi m^2 - 1)e^{-\pi m^2} > 0$ as $|t| \rightarrow \infty$ (the parts involving the complete $\Gamma(a)$ are $O(|t|^{7/4} e^{-\pi|t|/4})$), while $\Xi(t) \rightarrow 0$; so $\liminf_{|t| \rightarrow \infty} \xi_w(\frac{1}{2} + it) \geq (1 - w_m)(4\pi m^2 - 1)e^{-\pi m^2} > 0$. Case (ii): integrating by parts twice as in Theorem 3.3,

$$\xi_w(\tfrac{1}{2} + it) = C_w + 2 \int_0^\infty \Phi_w(u) \cos(tu) du, \quad \Phi_w = K_w'' - \tfrac{1}{4}K_w = \sum_n w_n \tilde{\varphi}_n,$$

where $K_w(u) = e^{u/2} \sum_n w_n e^{-\pi n^2 e^{2u}}$ and $C_w = \frac{1}{2} + 2K'_w(0)$, and the integral has the asymptotic expansion $\sum_{j \geq 0} 2(-1)^{j+1} \Phi_w^{(2j+1)}(0) t^{-2j-2}$. If every coefficient of $\xi_w(\frac{1}{2} + it) \sim C_w + \sum_j a_j t^{-2j-2}$ vanished, the odd part of the real-analytic function Φ_w would have zero Taylor series at 0, so Φ_w would be even. But the expansion of $\Phi_w(u)$ as $u \rightarrow -\infty$ has as coefficient of $e^{(5/2+2k)u}$ the number $-2\pi(2k+3)(-\pi)^k/k! \cdot \sum_n w_n n^{2k+2}$, while an even Φ_w decays double-exponentially as $u \rightarrow -\infty$. So $\sum_n w_n n^{2k+2} = 0$ for all $k \geq 0$, and since only finitely many n occur, $w \equiv 0$ (Vandermonde), which is excluded. Hence some coefficient is nonzero, and $\xi_w(\frac{1}{2} + it)$ has constant sign for $|t|$ large. $\square$

For ξ_N the leading coefficient is $C_w = c_N$. The same proof applies to Haglund's $\Xi_N = \xi_N - c_N$, whose leading coefficient is 0 and whose t^{-2} coefficient is negative (Proposition 3.4): no Ξ_N is real-rooted either, and Haglund's own list of the zeros of Ξ_1 [Hag09, p. 16] shows one real zero followed by non-real ones.

Theorem 4.2. *Let $|w_n| \leq Cn^A$ for some constants C, A , and $w \neq 0$. If every zero of ξ_w lies on $\operatorname{Re} s = \frac{1}{2}$, then $w \equiv 1$.*

Proof. Part (a) of the proof of Theorem 4.1 holds for polynomially bounded w , so ξ_w has infinitely many zeros, all on the line by hypothesis. The expansion of part (b) also holds for such w (Φ_w is real-analytic on $\mathbf{R}$ and all its derivatives decay double-exponentially on $[0, \infty)$), so all its coefficients vanish, $C_w = 0$, and Φ_w is even. Then $D(u) := K_w(u) - K_w(-u)$ is odd with $D'' = D/4$, so $D = 2\alpha \sinh(u/2)$, that is, $\psi_w(1/x) = \sqrt{x} \psi_w(x) - \alpha(\sqrt{x} - 1)$ for $\psi_w(x) = \sum_n w_n e^{-\pi n^2 x}$. Differentiating at $x = 1$ gives $K'_w(0) = \alpha/2$, and $C_w = 0$ gives $\alpha = -\frac{1}{2}$: ψ_w satisfies Riemann's theta relation exactly. Then $\delta := \psi_w - \psi = \sum_n (w_n - 1) e^{-\pi n^2 x}$ satisfies $\delta(1/x) = \sqrt{x} \delta(x)$ and decays faster than every power of x at 0 and at ∞ (exponentially at ∞ , and like $x^{-1/2} \delta(1/x)$ at 0), so its Mellin transform $\pi^{-s/2} \Gamma(s/2) \sum_n (w_n - 1) n^{-s}$ is entire, symmetric under $s \mapsto 1 - s$, and bounded in vertical strips. By Hamburger's theorem [Ham21] the Dirichlet series $f(s) = \sum_n (w_n - 1) n^{-s}$ is a constant multiple of ζ ; since f is entire, the constant is 0 and $w \equiv 1$. $\square$

For bounded w , Hamburger's theorem is used in the form stated by Nakamura [Nak20, Theorem D, p. 4]: if $F(s) = \sum a(n) n^{-s}$ converges absolutely for $\sigma > 1$, $P(s)F(s)$ is entire of finite order for some polynomial P , and $\xi_F(1 - s) = \xi_F(s)$ with $\xi_F(s) := \pi^{-s/2} \Gamma(s/2) F(s)$, then $F = C\zeta$; here $F = f$ is entire of finite order. For polynomially bounded w , Burnol's extension [Bur12, pp. 2–3, Corollaire 2] needs convergence only for $\operatorname{Re} s$ large, together with finitely many poles, $f(s) = O(e^{\exp \epsilon |s|})$ in vertical strips, and, for $g(s) = \chi(s) f(1 - s)$, $g(s) = O(|s|^N y^{-s})$ with $y > \frac{1}{2}$ and $g(\sigma) = c + O(\sigma^{-k})$ for every k ; here $g = f = (w_1 - 1) + O(2^{-\sigma})$, so all of these hold. (Hamburger's theorem needs more than the functional equation: see [Nak20, p. 4] for Knopp's solutions when (H1) is weakened.)

Proposition 4.3 (windowed real-rootedness is a height filtration). *Let F_N be entire with $F_N(1 - s) = F_N(s)$ and $F_N(\bar{s}) = \overline{F_N(s)}$. Suppose that for each T there is $N_0(T)$ such that for $N \geq N_0(T)$: (i) $\sup_{R_T} |F_N - \xi| < \min_{\partial D} |\xi|$ for a finite family D of small discs about the zeros of ξ in $R_T = [-1, 2] \times [-T, T]$ and for ∂R_T ; and (ii) F_N has no zeros with $|\operatorname{Im} s| \leq T$ outside R_T . Then, by Rouché's theorem on each disc and on R_T minus the discs, the zeros of F_N in R_T are those of ξ , moved by less than the disc radii, with multiplicity, and a zero of multiplicity one near the line lies on it by the symmetry $s \mapsto 1 - \bar{s}$. Hence, if the zeros of ξ in R_T are simple, F_N is real-rooted on R_T for all large N if and only if the Riemann hypothesis holds on R_T .*

Remark 4.4. For $F_N = \xi_N$ one has $\sup_{R_T} |\xi_N - \xi| \leq \frac{1}{2}(T^2 + 4) \max_{-1 \leq \sigma \leq 2} \sum_{n > N} g_n(\sigma) \approx T^2 e^{-\pi(N+1)^2} / (\pi(N+1)^2)$, which is below the size of ξ on the line up to $T \approx 4(N+1)^2$; and (ii)

holds for all large N , by comparing ξ_N with ξ for $2 \leq \operatorname{Re} s \leq \pi(N+1)^2$, with the main term $\frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\sum_{n \leq N} n^{-s}$ for $\operatorname{Re} s \geq \pi(N+1)^2$, and using $s \mapsto 1-s$ on the left. So in any induction “members real-rooted up to T_N implies real-rooted up to T_{N+1} ”, the only content is the Riemann hypothesis on the new window $T_N < |\operatorname{Im} s| \leq T_{N+1}$; what the family proves by itself (Theorem 3.1: the set where $\xi_N(\frac{1}{2} + it) < 0$ grows with N , so a lobe that carries real zeros keeps carrying them) is the easy half. This is the frame in which Conjecture 1 lives: it asks for an ordering property to carry the induction, and Theorem 1.1 shows that the ordering property fails near the height at which the approximant leaves the window.

5 Departure height and the lobe scan

The figures in this section are floating-point computations in mpmath at 30 to 80 digits, cross-checked by a second formula or a second contour where zeros are reported; they explain where to look and are not part of the proof of Theorem 1.1.

Departure. On the line, $\Xi(t)$ is of size roughly $e^{-\pi t/4}$, and the corrections P_N and Q_N are of size roughly $e^{-\pi(N+1)^2}$. The two balance near $t = 4(N+1)^2$, the *departure height*: below it the approximant tracks Ξ , above it the approximant is dominated by its correction. For the truncations ξ_N (from a complete census of the zeros in $0 < \operatorname{Im} s \leq 200$ and a count of the lobes up to height 320) the first negative lobe of Ξ that carries no real zero of ξ_N is (25.01, 30.42), (40.92, 43.33), (69.55, 72.07), (107.17, 111.03), (150.05, 150.93), (202.49, 204.19), (265.56, 266.61) for $N = 1, \dots, 7$, with ratio of lobe depth to P_N equal to 0.0070, 0.20, 0.32, 0.24, 0.12, 0.27, 0.020: the departure height is roughly $4(N+1)^2 + 6$ to $4(N+1)^2 + 10$. Haglund’s table shows the same law: his largest real zeros 14.05, 39.53, 65.03, 103.37, $\dots$, 489.39 for $N = 1, 2, 3, 4, \dots, 10$ lie close to $4(N+1)^2$. Every off-line zero of ξ_N found below height 200 lies outside the critical strip, at $\operatorname{Re} s \geq 1.39$.

The ordering property fails near departure. By Theorem 3.3, apart from the central lobe the real zeros of Ξ_N come in pairs from the positive lobes of Ξ whose height exceeds the smooth level Q_N , and by Corollary 3.2 those of ξ_N come from the negative lobes deeper than P_N . A lobe that fails the level, followed by a lobe that passes it, is expected to leave (and in both cases below does leave) a pair of non-real zeros over the first lobe with real zeros above it, which violates the ordering. Lobe heights decrease on average by a factor $e^{-\pi^2/\log(t/2\pi)}$ per step between lobes of the same sign, but fluctuate by comparable factors, so such an inversion is to be expected near the departure height for some N . We scanned the windows $[4(N+1)^2 - 40, 4(N+1)^2 + 90]$ for $N = 6, \dots, 30$, counting zeros in every window and comparing each lobe’s height with the level. The only violations are at $N = 24$ for ξ_N and at $N = 27$ for Ξ_N . The closest other approach for Haglund’s family is at $N = 21$, where a lobe beyond the first failing one reaches 0.93 of the level (a violation needs more than 1); the next closest is $N = 20$, at 0.48. The scan is not a proof that Conjecture 1 holds for $N \leq 26$.

The violation at $N = 27$. The positive lobe (3142.97945673, 3143.56749918) of Ξ has height 0.411 times Q_{27} and carries no real zero of Ξ_{27} ; the next positive lobe, (3144.49048815, 3146.28352838), has height 1.325 times Q_{27} and carries the real zeros 3144.8946622186 and 3145.5998495765. For $\operatorname{Im} s \in [3142.95, 3144.0]$ and $\operatorname{Re} s \in [-\frac{1}{2}, \frac{3}{2}]$ the argument principle counts 2.0 zeros at 30 and at 45 digits, none of them on the line: $s = 0.8152587993782148453823 + 3143.220682421536585287i$, inside the critical strip, and its mirror $1 - \bar{s}$. In Haglund’s variable $z = -i(s - \frac{1}{2})$ the mirror is the zero z_0 of Theorem 1.1, whose real part is below both real zeros.

6 The certificate

Theorem 1.1 was certified twice, by two computations written independently: *certificate A* in Arb ball arithmetic [Joh17] through python-flint 0.6.0 (FLINT 3.0.1) [FLINT], and *certificate B* in the outward-rounded interval arithmetic of mpmath 1.3.0 [mpm], with no use of Arb. Both evaluate the objects (2)–(3) exactly as Haglund defines them. Each was first run on a ladder of known cases (Table 3).

6.1 Certificate A: Arb ball arithmetic

Certificate A evaluates Ξ_{27} by two routes. *Route L* is the literal sum (3), with $\Gamma(w, a)$ from FLINT's `acb_hypgeom_gamma_upper`. At $x = 3144.8946$ its terms fall from $|\Phi_1| \approx |\Phi_2| \approx 8.0 \cdot 10^{-9}$ to $|\Phi_{27}| \approx 1.5 \cdot 10^{-991}$ while the sum is about 10^{-1070} , so about 1062 digits cancel; route L is used only at exact points, at 4800 and 6400 bits, where the cancellation is fully resolved. *Route T* writes $\Xi_{27} = \Xi - \sum_{n=28}^{31} \Phi_n - R_{31}$, with Ξ from (1) (FLINT's ζ and Γ), the four terms as in route L, and the remainder bounded by Lemma 6.3; near z_0 all these terms are about 10^{-1066} or smaller ($\Phi_{28} \approx \Xi \approx 1.49 \cdot 10^{-1066}$, $\Phi_{29} \approx 3 \cdot 10^{-1144}$), so route T has no catastrophic cancellation and is sharp on ball inputs at 256 bits. Route T uses Riemann's series $\Xi = \sum_{n \geq 1} \Phi_n$ [Hag09, (12), p. 3], which also follows from (4) and Lemma 2.1; route L does not.

Lemma 6.1. *For $w \in \mathbf{C}$, $a > 0$, $b \in \mathbf{R}$: $|G(w; a, b)| \leq 2\Gamma(\beta, a)/a^\beta$ with $\beta = b + |\operatorname{Im} w|$.*

Proof. For $u \geq 0$, $|\cos(x + iy)|^2 = \cos^2 x + \sinh^2 y \leq \cosh^2 y$, so $|\cos(2wu)| \leq e^{2u|\operatorname{Im} w|}$ and $|G| \leq 4 \int_0^\infty e^{2\beta u - ae^{2u}} du$; the substitution $t = ae^{2u}$ turns this into $2a^{-\beta} \int_a^\infty t^{\beta-1} e^{-t} dt$. $\square$

Lemma 6.2. *For $\beta \geq 1$ and $a > \beta - 1$: $\Gamma(\beta, a)/a^\beta \leq e^{-a}/(a - \beta + 1)$.*

Proof. With $t = ax$, $\Gamma(\beta, a)/a^\beta = \int_1^\infty x^{\beta-1} e^{-ax} dx$, and $x^{\beta-1} \leq e^{(\beta-1)(x-1)}$ for $x \geq 1$ since $\log x \leq x - 1$ and $\beta \geq 1$; integrate. $\square$

Lemma 6.3. *Let $Y \geq |\operatorname{Im} z|$, $\beta_9 = \frac{9}{4} + Y/2$, $a_n = \pi n^2$, and $M \geq 1$ with $a_{M+1} - \beta_9 + 1 > 0$. Put $U_n = 2e^{-a_n}(2\pi^2 n^4 + 3\pi n^2)/(a_n - \beta_9 + 1)$. Then $|\sum_{n>M} \Phi_n(z)| \leq \sum_{n>M} U_n \leq 2U_{M+1}$.*

Proof. In (3), $|\operatorname{Im}(z/2)| \leq Y/2$, and Lemmas 6.1 and 6.2 (with $\beta = \frac{9}{4} + Y/2$ and $\frac{5}{4} + Y/2$, both ≥ 1) give $|\Phi_n| \leq U_n$ (since $\Gamma(\beta, a)/a^\beta = \int_1^\infty x^{\beta-1} e^{-ax} dx$ increases with β , $|\operatorname{Im}(z/2)|$ may be replaced by $Y/2$). For $n > M$, $U_{n+1}/U_n \leq e^{-3\pi} \cdot 16 < 1.3 \cdot 10^{-3}$, since the ratio of the polynomial factors is at most $((n+1)/n)^4 \leq 16$ and the ratio of the denominators is below 1; so $\sum_{n>M} U_n \leq 2U_{M+1}$. $\square$

At $N = 27$, $M = 31$ and $Y \leq 0.317$, and $2U_{32} \leq 2 \cdot 10^{-1393}$. The winding numbers use the elementary fact that if a path lies in a closed axis-parallel box that excludes 0, the box lies in an open half-plane through 0 and the continuous change of argument along the path equals the principal value $\operatorname{Arg}(\gamma(1)/\gamma(0)) \in (-\pi, \pi)$. The boundary of each square is cut into pieces, a piece is bisected until its ball enclosure excludes 0, and the increments are summed; by the argument principle a total in $(2\pi(k - \frac{1}{2}), 2\pi(k + \frac{1}{2}))$ certifies k zeros inside. Every arithmetic operation and special function is Arb's, whose contract is that the output ball contains the exact value for every input in the input balls; no floating-point value is used in any decision. The only other inputs are Lemmas 6.1–6.3, the half-plane fact, Riemann's series for route T, the intermediate value theorem and the argument principle.

6.2 Certificate B: interval arithmetic with proved remainders

Certificate B uses mpmath's real intervals with outward rounding, complex numbers as rectangular boxes built on them, and exact rational Bernoulli numbers; every truncation adds a disk whose radius is a proved bound, each targeting 2^{-190} relative to the size of the truncated quantity ($|\zeta| \approx 1$, and e^{-X} for h), at a working precision of 200 bits. It writes $\Phi_n = 2X^2[h(\frac{s}{2} + 2) + h(\frac{1-s}{2} + 2)] - 3X[h(\frac{s}{2} + 1) + h(\frac{1-s}{2} + 1)]$ with $h(w) = X^{-w}\Gamma(w, X) = \int_1^\infty e^{-Xu}u^{w-1}du$, as in the proof of Lemma 2.1, and evaluates $\Xi_{27} = \Xi - \Phi_{28} - \Phi_{29} - \Phi_{30} - T_{30}$ with the following bounds.

- Γ by Stirling's series with the remainder bound of [DLMF, 5.11.1, §5.11(ii)], shifting the argument when $|w| < 300$ or $\operatorname{Re} w < 0$.
- ζ by Euler–Maclaurin summation [DLMF, 2.10.1] with the remainder bound $|R_m| \leq (2 - 2^{1-2m})|B_{2m}| |(s)_{2m}| N^{1-\sigma-2m} / ((2m)! (\sigma + 2m - 1))$ from [DLMF, 24.9.2]; at this height $N = 1012$ and $m = 94$.
- h by repeated integration by parts, $h(w) = e^{-X} \sum_{k < K} \tau_k + \rho_K$ with $\tau_k = (w - 1) \cdots (w - k) / X^{k+1}$ and $|\rho_K| \leq e^{-X} |\tau_K| X / (X - \max(\beta, 0))$, $\beta = \operatorname{Re} w - 1 - K$.
- The tail: $|h(w)| \leq e^{-X} / (X - \beta')$ with $\beta' = \max(\operatorname{Re} w - 1, 0)$, so for $X \geq \max(2\beta', 12)$ (with β' maximized over the four arguments in Φ_n) one has $|\Phi_n| \leq 9Xe^{-X}$, whence $|\sum_{n > M} \Phi_n| \leq 18\pi(M + 1)^2 e^{-\pi(M+1)^2}$; at $N = 27$, $M = 30$ and the bound is $3.7 \cdot 10^{-1307}$.

At $x = 3144.8946$ one has $|\Xi| = 1.4923145335789 \cdot 10^{-1066}$ and $|\Phi_{28}| = 1.49249055304208 \cdot 10^{-1066}$, so this route resolves $\Xi_{27} \approx 10^{-1070}$ with about 50 digits to spare. As a second rigorous route for the real zeros, certificate B also evaluates the literal sum (3) at 4400 bits, with $\Phi_1, \dots, \Phi_{27}$ from the lower incomplete gamma series (remainder $\leq e^{-X} X^K / |(w)_K|$ once $\operatorname{Re} w + K \geq X + 1$), integration by parts where its remainder bound is small, and Stirling's series: no ζ , no Euler–Maclaurin summation, and no use of Riemann's series.

Winding numbers come from a rigorous Taylor model. Since $f = \Xi_{27}$ is entire, its Taylor coefficients about c satisfy $|a_k| \leq M_R / R^k$, with M_R an upper bound for $|f|$ on $|z - c| = R$ obtained from box enclosures covering the circle. The discrete Fourier transform of f on m points of the circle $|z - c| = \rho$ gives $D_k = \sum_{l \geq 0} a_{k+lm} \rho^{lm}$, so $|D_k - a_k| \leq (M_R / R^k) q^m / (1 - q^m)$ with $q = \rho / R$, and for $|z - c| \leq r' < R$, with $x = r' / R$,

$$\left| f(z) - \sum_{k < K} D_k (z - c)^k \right| \leq E := M_R \left[\frac{q^m}{1 - q^m} \sum_{k < K} x^k + \frac{x^K}{1 - x} \right].$$

A boundary piece is accepted when the model's box enclosure plus $[-E, E]^2$ excludes 0 and the ratio of the end values has positive real part; the change of argument along it is then the principal value of that ratio's argument. At $N = 27$: $R = 0.1$, $M_R = 2.09255 \cdot 10^{-1065}$, $\rho = 0.002$, $m = 24$, $K = 12$, giving, with $r' = \sqrt{2}r$ for a square of half-width r about c , $E = 3.51 \cdot 10^{-1106}$ for $r \leq 10^{-5}$ and $1.36 \cdot 10^{-1087}$ at $r = 10^{-3}$; the constant coefficient D_0 agrees with a direct enclosure of $\Xi_{27}(c)$. For the 10^{-48} enclosure a second model is built at the center

$$\operatorname{Re} z^* = 3143.220682421536585287281295989417800119257644961983569,$$

$$\operatorname{Im} z^* = 0.3152587993782148453822863730092717652956513976685290909,$$

with $\rho = 0.001$ and $m = 40$, giving $E = 2.09 \cdot 10^{-1145}$; at $r = 10^{-48}$ the smallest lower modulus over the boundary pieces, $5.9 \cdot 10^{-1114}$, still exceeds the radii of the point enclosures, about 10^{-1120} , by a factor 10^6 . The trust base of certificate B is mpmath's primitive interval operations and the cited DLMF remainder bounds; Riemann's series is used for the tail route and checked numerically, not re-derived.

6.3 The enclosures

Tables 1–3 give the certified values. Here $c = 3143.2206824215 + 0.3152587994i$, and a square of half-width r about a center w is $w + [-r, r]^2$.

x	certificate A (ball midpoint $\pm$ radius)	certificate B (interval)
3144.8946	$-1.76019463127752 \cdot 10^{-1070} \pm 4.1 \cdot 10^{-1085}$	$-1.760194631277516 \cdot 10^{-1070}$
3144.8947	$+1.06871649226171 \cdot 10^{-1070} \pm 3.3 \cdot 10^{-1085}$	$+1.068716492261707 \cdot 10^{-1070}$
3145.5998	$+1.12975694291664 \cdot 10^{-1070} \pm 1.9 \cdot 10^{-1085}$	$+1.129756942916642 \cdot 10^{-1070}$
3145.5999	$-1.14913915362778 \cdot 10^{-1070} \pm 2.9 \cdot 10^{-1085}$	$-1.149139153627783 \cdot 10^{-1070}$

Table 1: Enclosures of $\Xi_{27}(x)$ at the ends of the two real-zero intervals. Certificate A: balls as printed, display rounding included; the Arb radii are $3.3 \cdot 10^{-1435}$ (route L, 4800 bits), $6.0 \cdot 10^{-1915}$ (route L, 6400 bits) and $5.0 \cdot 10^{-1138}$ (route T, 256 bits), and all four evaluations (route T at 256 and 1024 bits, route L at 4800 and 6400 bits) give the same signs. Certificate B: tail route, box radii at most $6 \cdot 10^{-1121}$; its literal route at 4400 bits (radius about $1\text{--}2 \cdot 10^{-1171}$) agrees to all digits shown and overlaps.

cert.	center	half-width r	boundary pieces	k
A	c	10^{-3} (256 and 512 bits)	512, 512	1
A	c	$10^{-6}, 10^{-10}, 4 \cdot 10^{-11}$	512, 663, 807	1
A	$c + 2.5 \cdot 10^{-3}$	10^{-3}	269	0
A	$c + 1.5 \cdot 10^{-10}$	10^{-10}	714	0
B	c	$10^{-3}, 10^{-5}, 10^{-7}, 10^{-9}$	32 each	1
B	c	$10^{-10}, 5 \cdot 10^{-11}, 4 \cdot 10^{-11}$	32, 32, 33	1
B	c	$3.7 \cdot 10^{-11}$	36	1
B	c	$3.5 \cdot 10^{-11}$	33	0
B	$c + 0.01$	10^{-3}	32	0
B	z^*	$10^{-30}, 10^{-40}, 10^{-45}, 10^{-48}$	–	1

Table 2: Certified winding numbers k of Ξ_{27} along the boundaries of squares. In certificate A every run with $k = 1$ has total change of argument $6.28318530717959 \pm 4 \cdot 10^{-15}$. In certificate B at $r = 10^{-3}$ the smallest lower modulus over the pieces is $5.89 \cdot 10^{-1069}$ against the model error $E = 1.36 \cdot 10^{-1087}$; at $r = 3.7 \cdot 10^{-11}$ it is $1.97 \cdot 10^{-1078}$ against $E = 3.51 \cdot 10^{-1106}$. The run at $r = 3.5 \cdot 10^{-11}$ returns 0 because the zero lies just outside that square, and the rows with centers $c + 2.5 \cdot 10^{-3}$, $c + 1.5 \cdot 10^{-10}$ and $c + 0.01$ are zero-free controls.

Cross-checks. In certificate A, routes L and T overlap at all 18 points compared (the center, the reference zero, the corners and midpoints of the square of half-width 10^{-3} , the corners of the square of half-width $4 \cdot 10^{-11}$, and the four points of Table 1), with $|L - T| \leq 6.43 \cdot 10^{-1137}$ at 256 bits, and $|L - T| \leq 1.18 \cdot 10^{-1288}$ at 1024 bits at the fourteen points other than those of Table 1. In certificate B the literal and tail routes overlap at the four points of Table 1, its enclosures of Ξ contain mpmath's ordinary 400-bit values, and its two methods for h at $X = 784\pi$ overlap. The two certificates agree with each other to every digit printed. A third evaluation of the literal sum (3) in Arb at 4800 bits gives $\Xi_{27}(3144.8946) \in [-1.76019463128 \cdot 10^{-1070} \pm 2.49 \cdot 10^{-1082}]$ and $\Xi_{27}(3144.8947) \in [1.06871649226 \cdot 10^{-1070} \pm 1.71 \cdot 10^{-1082}]$, the same signs at 3145.5998 and 3145.5999, and Haglund's two ladder zeros; at the point $3143.220682421536585287 + 0.3152587993782148453823i$ it gives a value of modulus about $1.7 \cdot 10^{-1084}$, against about $6.1 \cdot 10^{-1072}$ at a point 10^{-6} to the right.

test	certificate A	certificate B
Ξ_1 at 14.04543957 and 14.04543959 (Haglund: 14.0454395788)	$+1.347060456147 \cdot 10^{-11},$ $-1.704102125315 \cdot 10^{-11}$	$+1.347060456146727 \cdot 10^{-11},$ $-1.704102125314551 \cdot 10^{-11}$
Ξ_2 at 39.5324810797 and 39.5324810799 (Haglund: 39.5324810798)	$+3.605517254360 \cdot 10^{-21},$ $-4.593418771015 \cdot 10^{-21}$	$+3.605517254360067 \cdot 10^{-21},$ $-4.5934187710153 \cdot 10^{-21}$
Ξ_1 about Haglund's zero 20.62534600592171760132974 + 2.697151842339519632505712 i	$k = 1$ at $r = 10^{-10}$ (1516 pieces)	$k = 1$ at $r = 10^{-3}$ and 10^{-8}
zero-free controls for Ξ_1	$k = 0$ (center shifted by 0.5, $r = 0.1$; shifted by $3 \cdot 10^{-10}$, $r = 10^{-10}$)	$k = 0$ ($17 + i$, $r = \frac{1}{4}$)

Table 3: The ladder, run before $N = 27$ with the same code: the largest real zeros of Ξ_1 and Ξ_2 from Haglund's table [Hag09, p. 4], the first non-real zero of Ξ_1 from his appendix [Hag09, p. 16], and zero-free squares.

6.4 Proof of Theorem 1.1

(a) By (2), $G(z; a, b)$ is real for real z , so Ξ_{27} is real and continuous on $\mathbf{R}$. The enclosures of Table 1 have strictly opposite signs at the ends of each interval, in both certificates; by the intermediate value theorem each interval contains a zero.

(b) Ξ_{27} is entire, being a finite sum of entire functions. Certificate B finds winding number 1 along the boundary of the square of half-width $3.7 \cdot 10^{-11}$ about c , with every boundary enclosure excluding 0; by the argument principle the open square contains exactly one zero z_0 , counted with multiplicity, so z_0 is simple. Certificate A independently finds winding number 1 at half-width $4 \cdot 10^{-11}$. The square $z^* + [-10^{-48}, 10^{-48}]^2$ lies inside the first square, since $z^* - c = 3.66 \dots \cdot 10^{-11} - 2.18 \dots \cdot 10^{-11} i$; winding number 1 there (certificate B) locates the same zero z_0 .

(c) From (b), $\text{Im } z_0 \geq 0.3152587994 - 3.7 \cdot 10^{-11} > 0.3152$ and $\text{Re } z_0 \leq 3143.2206824215 + 3.7 \cdot 10^{-11} < 3143.2207 < 3144.8946 < x_1$. So $z_0 \in \mathcal{Q}$ is non-real and precedes the real zero x_1 in any listing of the zeros of Ξ_{27} in $\mathcal{Q}$ by increasing real part, while $\text{Im } z_0 > 0 = \text{Im } x_1$. $\square$

7 The companion violation at $N = 24$

For the truncations ξ_N the ordering property reads: the zeros of $z \mapsto \xi_N(\frac{1}{2} + iz)$ in $\mathcal{Q}$, listed by increasing real part, have nondecreasing imaginary parts. Since $\xi_N \rightarrow \xi$ locally uniformly, the Hurwitz argument of Haglund's Proposition 1 shows that this property for every N would also imply the Riemann hypothesis. It fails at $N = 24$.

Theorem 7.1. (a) *The real function $t \mapsto \xi_{24}(\frac{1}{2} + it)$ changes sign on (2510.2026, 2510.2027) and on (2510.7086, 2510.7087), so ξ_{24} has zeros on the critical line at heights in these intervals.*

(b) *The function $z \mapsto \xi_{24}(\frac{1}{2} + iz)$ has exactly one zero in the square of half-width 10^{-10} about $2508.2839748053 + 0.3159896243 i$.*

(c) *Hence ξ_{24} has a zero ρ_{24} off the critical line, inside the critical strip, with $\text{Im } \rho_{24} < 2508.2839748054$, below zeros on the line: the ordering property fails for ξ_{24} .*

Both certificates prove this, with the same methods as in Section 6; here route T and certificate B use Riemann's identity (4), which rests on Jacobi's theta relation (cited, not re-proved), while

certificate A's route L, the literal sum, does not. The enclosures on the line are, in certificate A (both routes; route L is the literal sum at 4800 bits, cancelling about 855 digits)

$$\xi_{24}(\tfrac{1}{2} + it) \in \begin{cases} [+2.81317536164 \cdot 10^{-854} \pm 5 \cdot 10^{-866}], & t = 2510.2026, \\ [-1.65267129215 \cdot 10^{-854} \pm 3 \cdot 10^{-866}], & t = 2510.2027, \\ [-3.23281139505 \cdot 10^{-854} \pm 4 \cdot 10^{-866}], & t = 2510.7086, \\ [+6.14184061542 \cdot 10^{-855} \pm 4 \cdot 10^{-867}], & t = 2510.7087, \end{cases}$$

and in certificate B $+2.813175361644763 \cdot 10^{-854}$, $-1.65267129214774 \cdot 10^{-854}$, $-3.232811395053649 \cdot 10^{-854}$ and $+6.141840615423139 \cdot 10^{-855}$ at the same four points. Certificate A finds (in the s -plane; the symmetry $\xi_{24}(\frac{1}{2} + i\bar{z}) = \overline{\xi_{24}(\frac{1}{2} + iz)}$ carries its squares to those of (b)) winding number 1 along the s -plane squares of half-width 10^{-3} (151 pieces) and 10^{-10} (174 pieces) about $0.8159896243 + 2508.2839748053i$, and 0 for a control square shifted by $2.5 \cdot 10^{-3}i$; certificate B finds winding number 1 for $z \mapsto \xi_{24}(\frac{1}{2} + iz)$ on squares of half-width 10^{-3} , 10^{-6} , 10^{-9} and 10^{-10} about $2508.2839748053 + 0.3159896243i$ (Taylor model with $R = 0.1$, $\rho = 0.002$, $m = 24$, $K = 12$, $M_R = 4.07 \cdot 10^{-849}$; at $r = 10^{-3}$ the smallest lower modulus is $1.16 \cdot 10^{-852}$ against $E = 2.6 \cdot 10^{-871}$), and 0 for the control square shifted by 0.01. Numerically, at 50 digits, $\rho_{24} = 0.8159896243043424688941 + 2508.283974805324153202i$, and the zero in (b) is its mirror $1 - \bar{\rho}_{24}$.

The mechanism is the one of Section 5 with the signs reversed: the negative lobe (2507.87103958, 2508.7660461) of Ξ has depth 0.595 times P_{24} and carries no real zero of ξ_{24} , while the next negative lobe (2509.65299744, 2511.66018979) has depth 1.130 times P_{24} . The off-line zero lies inside the critical strip, at $\operatorname{Re} s = 0.8160$. As with Ξ_{27} , it is a zero of the approximant at a height where the approximant has departed from ξ .

Acknowledgment

I thank J. Haglund for pointing out the corrected copy [Hag11w] of his paper.

Data and code availability

The two certificates of Section 6 — their scripts and logs, and for each a document stating every bound used, with its proof or its source — are deposited with this paper on Zenodo as a self-contained archive, `haglund-counterexample-certificate.zip`, SHA-256 `4dfb1360042c3123b815e571a1916720081beca51fe9be8d6ed4760f41b1091c`, together with a third evaluation of the literal sum, second runs of both certificates, and a README giving the exact commands and their expected outputs. On one core of an Apple M4, certificate A runs in about 72 seconds (Python 3.9.6, python-flint 0.6.0) and certificate B in about 15 minutes (Python 3.9.6, mpmath 1.3.0), most of it the 4400-bit literal cross-check. The floating-point computations behind Sections 3 and 5 (the zero census, the tail coefficients, the lobe scan and the 50-digit zeros) are in the repository <https://github.com/x67ai/riemann-rh-program>.

References

- [Ahn12] S.-H. (Shirley) Ahn, *On the zeros of component functions of the Riemann Xi approximates*, M.A. thesis, University of Pennsylvania, 2012.

- [Bac26] M. L. Baccaro, *Haglund’s Zero-Trajectory Conjecture for the First Riemann Xi Approximant*, preprint, version v1.0.0, Zenodo, August 22, 2026, DOI 10.5281/zenodo.22059236.
- [Bur12] J.-F. Burnol, *Deux extensions de Théorèmes de Hamburger (portant sur l’équation fonctionnelle de la fonction zêta)*, Expo. Math. **30** (2012), 295–308; arXiv:1106.4749.
- [DLMF] *NIST Digital Library of Mathematical Functions*, <https://dlmf.nist.gov/>, release date 2026-09-15; equations 2.10.1, 5.11.1, 24.9.2 and §5.11(ii).
- [Edw74] H. M. Edwards, *Riemann’s Zeta Function*, Academic Press, New York, 1974; reprinted by Dover Publications, Mineola, NY, 2001.
- [FLINT] The FLINT team, *FLINT: Fast Library for Number Theory*, version 3.0.1, <https://flintlib.org>; used through python-flint 0.6.0, <https://github.com/flintlib/python-flint>.
- [Hag09] J. Haglund, *Some conjectures on the zeros of approximates to the Riemann Ξ -function and incomplete gamma functions*, arXiv:0910.5228v1, 27 October 2009. Page and equation numbers in this paper refer to this version, except where [Hag11w] is cited.
- [Hag11] J. Haglund, *Some conjectures on the zeros of approximates to the Riemann Ξ -function and incomplete gamma functions*, Cent. Eur. J. Math. **9** (2011), no. 2, 302–318, DOI 10.2478/s11533-010-0095-3.
- [Hag11w] J. Haglund, *Some conjectures on the zeros of approximates to the Riemann Ξ -function and incomplete gamma functions*, author’s copy dated February 9, 2011, <https://www2.math.upenn.edu/~jhaglund/preprints/rh8.pdf> (read October 3, 2026).
- [Ham21] H. Hamburger, *Über die Riemannsche Funktionalgleichung der ζ -Funktion (erste Mitteilung)*, Math. Z. **10** (1921), no. 3–4, 240–254.
- [Joh17] F. Johansson, *Arb: efficient arbitrary-precision midpoint-radius interval arithmetic*, IEEE Trans. Comput. **66** (2017), no. 8, 1281–1292, DOI 10.1109/TC.2017.2690633.
- [Ki05] H. Ki, *All but finitely many non-trivial zeros of the approximations of the Epstein zeta function are simple and on the critical line*, Proc. London Math. Soc. (3) **90** (2005), no. 2, 321–344, DOI 10.1112/S0024611504015060.
- [Ki06] H. Ki, *Zeros of the constant term in the Chowla–Selberg formula*, Acta Arith. **124** (2006), no. 3, 197–204.
- [LM11] J. C. Lagarias and D. Montague, *The integral of the Riemann ξ -function*, Comment. Math. Univ. St. Pauli **60** (2011), 143–169; arXiv:1106.4348.
- [LMOZ24] D. Liu, Y. Matiyasevich, J. Oesterlé and A. Zaharescu, *Euler product sieve*, arXiv:2406.00786, 2 June 2024.
- [mpm] The mpmath development team, *mpmath: a Python library for arbitrary-precision floating-point arithmetic*, version 1.3.0, <https://mpmath.org>.
- [Nak20] T. Nakamura, *Dirichlet series with periodic coefficients, Riemann’s functional equation and real zeros of Dirichlet L -functions*, arXiv:2008.02570v4, 10 September 2021.
- [Rie59] B. Riemann, *Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*, Monatsber. Königl. Preuss. Akad. Wiss. Berlin (1859), 671–680; English translation in [Edw74, Appendix].